

# Ideas and perspectives: Emerging contours of a dynamic exogenous kerogen cycle

Thomas M. Blattmann<sup>1</sup>

<sup>1</sup>Biogeochemistry Research Center, Research Institute for Marine Resources Utilization, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima-cho, 237-0061 Yokosuka, Japan

*Correspondence to:* Thomas M. Blattmann (blattmannt@jamstec.go.jp)

**Abstract.** Growing evidence points to the dynamic role that kerogen is playing on Earth's surface in controlling atmospheric chemistry over geologic time. Although quantitative constraints on the weathering of kerogen remain loose, its changing weathering behavior modulated by the activity of glaciers suggests that this largest pool of reduced carbon on Earth may have played a key part in atmospheric CO<sub>2</sub> variability across recent glacial-interglacial ~~cycles~~<sup>times</sup> and beyond. This work enunciates the possibility of kerogen oxidation as a major driver of atmospheric CO<sub>2</sub> increase in the wake of glacial episodes. This hypothesis of centennial and millennial-timescale-relevance for this chemical weathering pathway is substantiated by several lines of independent evidence synthesized in this contribution including the timing of CO<sub>2</sub> increase, CO<sub>2</sub> isotop~~ie~~<sup>ic</sup> composition (<sup>13</sup>C and <sup>14</sup>C), seawater osmium record, kerogen oxidation kinetics, observations of kerogen reburial, and modeling results. The author hypothesizes that the deglaciation of kerogen-rich lithologies in western Canada contributed in a major~~ly~~<sup>way</sup> to the characteristic deglacial increase in atmospheric CO<sub>2</sub>, which reached an inflection point ≤300 years after the Laurentide Ice Sheet retreated into the kerogen-poor Canadian Shield. To quantitatively constrain the contribution of kerogen oxidation to CO<sub>2</sub> rise at glacial terminations, systematic studies on CO<sub>2</sub> fluxes emanating from the weathering of different lithologies, oxidation kinetics of kerogen along glacial chronosequences, and high-resolution temporal changes in the aerial extent of glacially exposed lithological units are needed.

## 1 Introduction

Over geologic timescales, atmospheric CO<sub>2</sub> is controlled by the combined effects of chemical weathering of silicates and carbonates and the organic carbon cycle (Berner et al., 1983; Berner, 1990; Torres et al., 2014). Organic carbon in the form of kerogen<sup>1</sup> comprises around 15 million PgC, which encompasses over 99.9% of reduced carbon present on Earth. Tectonic uplift and denudation subjects 150 PgC/kyr of kerogen to weathering on Earth's surface (Hedges and Oades, 1997) thereby facilitating entry of this geologically ancient carbon into the atmosphere ~~and other surficial carbon pools standing in close communication (Hedges and Oades, 1997)~~. Upon oxidation of kerogen, O<sub>2</sub> is consumed and CO<sub>2</sub> is released to the atmosphere.

---

<sup>1</sup> In this work, kerogen is used as an umbrella term for all rock-derived forms of reduced carbon including soluble, insoluble, rock disseminated, rock forming, solid and liquid forms, as well as fossil palynomorphs, biogenic and abiogenic graphite.

In reverse, biospheric organic carbon burial in marine sediments removes carbon from ~~the~~ Earth's surface thereby drawing down atmospheric CO<sub>2</sub> and increasing O<sub>2</sub> over geological timescales (Galy et al., 2008). Therefore, kerogen weathering and sedimentary organic matter burial in the ocean play compensat~~ory~~~~ional~~ roles in governing atmospheric chemistry (Fig. 1) with kerogen oxidation considered important for atmospheric chemistry over million-year timescales (e.g., Petsch, 2014; Bolton et al., 2006).

However, the decay of kerogen on Earth's surface is incomplete (e.g., Hemingway et al., 2018; Leythaeuser, 1973), with physical erosion followed by riverine transport (Galy et al., 2015) and reburial in lacustrine and marine settings (e.g., Blattmann et al., 2018a; Blattmann et al., 2019b; Cui et al., 2016; Sackett et al., 1974; Sparkes et al., 2020). The operation of this "simple cycle besides the more complicated common circulation of carbon", enunciated by Sauramo (1938), begs the questions (i) what is the reburial efficiency of kerogen, (ii) what is the weathering efficiency of kerogen, (iii) what are their controlling factors, and (iv) ~~what are the implications of them changing for atmospheric chemistry~~how do reburial and weathering efficiency vary over geologic time and spacescales? The answers presented will suggest that the exogenous kerogen cycle behaves dynamically with ~~Here,~~ this contribution hypothesizinges a connection between kerogen oxidation and millennial/centennial-scale atmospheric CO<sub>2</sub> increases ~~during~~at glacial terminations.

## 2 Carbon isotopes and contradictions?

Driven mainly by orbital cyclicity, glaciers have rhythmically waxed and waned across Earth's surface together with changes in atmospheric chemistry, including the greenhouse gas CO<sub>2</sub> (Hays et al., 1976; Barnola et al., 1987). The amount and carbon isotope~~ie~~ composition of atmospheric CO<sub>2</sub> (Fig. 2) depend on (1) the terrestrial biospheric organic carbon pool, (2) the dissolved inorganic carbon pools residing in the surface and deep ocean, (3) exchanges between the atmosphere and the terrestrial biosphere and ocean, and (4) the export of organic matter and carbonate from the surface to deep waters and sediments of the oceans (Sigman and Boyle, 2000). ~~Aeross-During~~ deglaciation, an increasingly ~~voluminous-large~~ terrestrial biospher~~ic carbon poole~~ (Bird et al., 1994; Shackleton, 1977; Shackleton et al., 1983) is inferred to have controlled an increase in the stable carbon isotope~~ie~~ composition of dissolved inorganic carbon in ocean waters. However, this line of reasoning is complicated by various carbon pools changing in size at the same time as stable carbon isotope fractionation occurs, as carbon is exchanged between pools such as the terrestrial biosphere and pedosphere (see also Zeng, 2003, 2007)~~carbon pools changing in size coeval to carbon isotope fractionation as carbon rotates between these (e.g., terrestrial biosphere, pedosphere; see also discussions in Zeng (2003, 2007)).~~ Additionally~~In addition~~, during times of ~~most rapid~~highest CO<sub>2</sub> increase ~~during~~in transitions from glacial to interglacial periods, negative stable carbon isotope~~ie~~ shifts in atmospheric CO<sub>2</sub> ~~have~~ occurred (Fig. 3; Smith et al., 1999; Schmitt et al., 2012). ~~This, which~~ is a strong indicator ~~that~~of respired organic ~~matter~~carbon ~~acted~~ing as ~~source~~ea direct source of carbon~~ly~~ to the atmosphere (Bauska et al., 2016). Additionally, the constraints imposed by radiocarbon clearly indicate there must have been a source of carbon to the atmosphere and ocean depleted in or devoid of radiocarbon (Broecker and Clark 2010; Hain et al., 2014; Rafter et al., 2019; Zhao et al., 2018), thereby limiting the potential contributions from a

60 modern biospheric organic carbon source. Other studies have proposed that carbon sourced from dissolved inorganic carbon  
from the deep ocean was the predominant source for carbon ~~transferred to~~ the atmosphere ~~during~~ glacial terminations (e.g.,  
Hain et al., 2014). However, this hypothesis appears inconsistent with the negative fluctuation observed in the  $^{13}\text{C}$  fingerprint  
of atmospheric  $\text{CO}_2$  ~~during this time and requires a complex overlay of processes to reconcile (see discussion in~~ Broecker and  
McGee, 2013). Terrestrial organic carbon comprises the sum of biospheric and surface exposed lithogenic (i.e., kerogen) forms  
65 of organic carbon, with both exhibiting similar  $^{13}\text{C}$  signatures yet modern and dead  $^{14}\text{C}$  signatures, respectively (e.g., Lewan,  
1986; Meyers, 1994). Complementing the widely supported hypotheses of a strictly increasing terrestrial organic carbon pool  
and major changes in ocean-atmosphere  $\text{CO}_2$  exchange (e.g., Shackleton, 1977; Lindgren et al., 2018; Sigman and Boyle,  
2000). ~~As the author will argue,~~ the author will argue that the  $^{13}\text{C}$  and  $^{14}\text{C}$  lines of evidence suggest ~~major~~ the release of  $\text{CO}_2$  to  
the atmosphere via kerogen oxidation ~~to the atmosphere during deglaciation, contradicting (or complementing) the commonly~~  
70 ~~held notions of a strictly increasing terrestrial organic carbon pool and major changes in ocean-atmosphere  $\text{CO}_2$  exchange.~~

### 3 Kerogen and glaciers – Dynamic modulators of the global carbon cycle?

Chemical weathering of silicates and carbonates as well as the burial and oxidation of organic matter exert fundamental control  
over atmospheric chemistry on Earth over geologic timescales (Berner et al., 1983; Berner, 1990; Hilton and West, 2020;  
Petsch, 2014). On modern-day Earth, carbon release and sinks from mineral-based weathering (i.e., silicates and carbonates  
75 via carbonic and sulfuric acids) and biogeochemical fluxes (i.e., organic matter burial and kerogen oxidation) stand in fine  
balance to one another with contrasting concoctions of these under different tectonosedimentary settings (e.g., Bufer et al.,  
2021; Hilton and West, 2020; Blattmann et al., 2019a; Horan et al., 2019). Deglaciation likely modulated silicate and carbonate  
weathering resulting in enhanced (temporary) drawdown of atmospheric  $\text{CO}_2$  (e.g., Tranter, 1996; Gibbs and Kump, 1996;  
Munhoven and François 1996; c.f., Schachtman et al., 2019). Glaciers have been invoked as agents for accelerating chemical  
80 weathering of carbonate and silicate minerals by increasing sediment yield and creating reactive substrate with high surface  
area (Torres et al., 2017; Vance et al., 2009; Yu et al., 2021), with carbonate weathering constituting a source of  $\text{CO}_2$  to the  
atmosphere when sulfuric acid, stemming from oxidized sulfide minerals, is involved (c.f., Torres et al., 2014; Kölling et al.,  
2019). In the following, the idea of glaciers producing reactive substrate susceptible to chemical weathering is reviewed and  
explored for kerogen.

85 Evidence accumulated ~~from~~ over a century of scientific studies supports the idea that the reburial of kerogen ~~has been~~ was  
more extensive during cold interludes in Earth history where glacial erosion and ice rafting ~~was widespread~~ dominated  
(Blattmann et al., 2018b). This general pattern of kerogen reburial is also observed spatially today with more efficient reburial  
in high latitude, glaciated regions than in low latitude sedimentary systems (Cui et al., 2016). This enhanced reburial efficiency  
strengthens the short-circuiting of the exogenous kerogen cycle by keeping this ancient, reduced carbon locked away.  
90 However, under glaciofluvial conditions, the entry of kerogen-bound carbon into surficial carbon pools is promoted (Horan,  
2018), with, glacial meltwater releasing dissolved organic matter from kerogen, which is readily degraded and consumed by

microbes (Hood et al., 2009; Schillawski and Petsch, 2008; Sharp et al., 1999). Additionally, frost shattering, together with the retreat of glaciers, exposes finely ground, high surface area, reactive sediment, thereby accelerating oxidation and the release of kerogen-bound carbon-derived CO<sub>2</sub> to the atmosphere (Fischer and Gaupp, 2005; Horan et al., 2017). The initially strong input of kerogen-derived CO<sub>2</sub> would die down as the availability of glacially ground, reactive kerogen declines into the interglacial period. Enhanced degradation of kerogen in the wake of glacial episodes is consistent with observations from areas of ongoing deglaciation (Horan et al., 2017). Contemporaneous perturbations in CO<sub>2</sub> fluxes are expected to result for silicate and carbonate weathering via carbonic and sulfuric acids (Vance et al., 2009; Torres et al., 2014). Analogously, glaciers as agents for accelerating chemical weathering by increased sediment yield and creation of high surface area reactive substrate have also been invoked for silicate and carbonate minerals (Torres et al., 2017; Vance et al., 2009), with carbonate weathering constituting a source of CO<sub>2</sub> to the atmosphere when sulfuric acid is involved (Torres et al., 2014). However, kerogen oxidation, owing to its faster weathering kinetics and direct conversion to CO<sub>2</sub> leading to considerable losses in kerogen content over decadal timescales (Horan et al., 2017 and references therein; Fischer et al., 2007). This is a manifest process by which CO<sub>2</sub> can directly be injected directly into the atmosphere and intricately linked to impact glacial-interglacial cyclicity (Fig. 2).

In the following, quantitative arguments are presented: Of the approximately 150 PgC/kyr of kerogen reaching the Earth's surface ( $F_L$ ; Hedges and Oades, 1997),  $43^{+61}_{-25} \text{ PgC/kyr} \frac{\text{PgC}}{\text{km}^2 \text{yr}}$  is currently exported by rivers to oceans ( $F_{L-O}$ ; Galy et al., 2015; Copard et al., 2007), indicating that the modern-day reburial efficiency of this carbon lies in the ballpark of 30% (10-70%). Normalized to global land area ( $A_L = 149,000,000 \text{ km}^2$ ), the following average flux of carbon due to oxidation of kerogen to the atmosphere ( $F_{L-A}$ ) can be calculated from expected:

$$0.72^{+0.17}_{-0.41} \text{ MgC/km}^2 \text{yr} F_{L-A} = \frac{F_L - F_{L-O} 43^{+61}_{-25} \text{ PgC/kyr}}{A_L 149,000,000 \text{ km}^2} \quad (\text{Equation 1})$$

Here, equation 1 yields a modern-day global  $F_{L-A}$  of  $0.72^{+0.17}_{-0.41} \frac{\text{MgC}}{\text{km}^2 \text{yr}}$ . Reports of local and regional CO<sub>2</sub> fluxes from kerogen oxidation are based on disparate study sites and methods (Table 1). From this, reported CO<sub>2</sub> fluxes stemming from oxidation of rock disseminated forms of kerogen and across larger catchment areas under aerobic conditions span over two orders of magnitude ranging from 0.3 to  $64 \frac{\text{MgC}}{\text{km}^2 \text{yr}} \text{ MgC/km}^2 \text{yr}$ . The only quantities available for catchments of ongoing deglaciation stem from the Southern Alps of New Zealand, which are one to two orders of magnitude faster than those of the average Earth surface (ranging between  $9-50 \frac{\text{MgC}}{\text{km}^2 \text{yr}} \text{ MgC/km}^2 \text{yr}$ ; Horan et al., 2017). Taking this flux range as a working first-order estimate, how would kerogen oxidation translate into changes in atmospheric CO<sub>2</sub> concentration? In Zeng's (2003) modeling work postulating soil oxidation upon retreat of overriding glaciers, the effect of continuous inputs of organic matter-derived carbon directly to the atmosphere are calculated: 550 PgC oxidized lead to 60 PgC increase in atmospheric CO<sub>2</sub> (30 ppm increase). Similarly, modeling by Simmons et al. (2016) also propose glacially overridden the oxidation of overridden soil

organic carbon ~~oxidation upon subsequent to~~ deglaciation and calculate a 600 PgC release resulting in a 40-60 ppm increase  
 125 in atmospheric CO<sub>2</sub> (c.f., also Ciais et al. (2012) who calculate the deglacial oxidation of 700 PgC of an “inert” terrestrial  
 organic carbon pool). The idea of soil oxidation is mathematically analogous to kerogen oxidation in the hypothesis presented  
 here. How much area of deglaciaded terrain ( $A_D$ ) would therefore be needed to release 550-600 PgC of kerogen-derived carbon?  
 Using Zeng’s (2003) and Simmons et al.’s (2016) carbon release estimates ( $550 \text{ PgC} \leq C_R \leq 600 \text{ PgC}$ ) and Horan et al.’s  
 (2017) kerogen oxidation kinetics ( $9 \frac{\text{MgC}}{\text{km}^2 \text{yr}} \leq F_{KO} \leq 50 \frac{\text{MgC}}{\text{km}^2 \text{yr}}$ ) over a 6,000-year time window ( $T_{G-I}$ , timespan of greatest  
 130 pronounced CO<sub>2</sub> increase over the glacial-interglacial transition; see Fig. 3) and subtracting global average baseline kerogen  
 oxidation ( $F_{L-A} \approx 0.72^{+0.17}_{-0.41} \frac{\text{MgC}}{\text{km}^2 \text{yr}}$ ; see Eq. 1):

$$A_D = \frac{C_R}{T_{G-I} * (F_{KO} - F_{L-A})} \quad \text{(Equation 2)}$$

135 Between 2 to 12 million km<sup>2</sup> =  $\frac{\text{Between 550 to 600 PgC}}{6000 \text{ yr} * ([\text{Between 9 to 50}] - 0.72^{+0.17}_{-0.41}) \text{ MgC/km}^2 \text{ yr}}$  (Equation 2)

With calculated  $A_D$  yielding between 2 and 12 million km<sup>2</sup>, in the most conservative estimate, 12 million km<sup>2</sup> of deglaciaded  
 terrain containing sedimentary and metasedimentary rocks and their debris (glacial flour accumulated over millennia of  
 glaciation) would cover the source. In comparison, Canada covers approximately 10 million km<sup>2</sup>, which was mostly covered  
 140 by the Laurentide Ice Sheet during the Last Glacial Maximum. With this, a plausible scenario for releasing kerogen-derived  
 CO<sub>2</sub> to the atmosphere that could account for a 30-60 ppm rise during the glacial-interglacial transition encompassing an area  
 equal to or less than the terrestrial extent of the Laurentide Ice Sheet is identified. Similar to the oxidation fluxes reported from  
 deglaciating catchments (Horan et al., 2017) are those reported for the in-situ weathering of massive outcrop formations of  
 marls and shales (e.g., Soulet et al., 2018; Littke et al., 1991; see Table 1). Kerogen oxidation fluxes appear higher for  
 145 catchments showing larger glacial coverage (Horan et al., 2017) suggesting that the relative contribution of freshly exposed  
 and ground bedrock is highest for these areas and that kerogen oxidation fluxes progress over the course of deglaciation. The  
 CO<sub>2</sub> flux emanating from kerogen oxidation in soils formed from glacial till (Keller and Bacon, 1998) suggests that oxidation  
 fluxes greater by an order of magnitude greater than the global average can be sustained for millennia afterpast deglaciation.  
 However, a systematic chronosequence understanding is currently lacking that allows conclusions to be drawn as a variety of  
 150 factors (e.g., lithology, initial kerogen content, grain size) exert control over kerogen oxidation (c.f., Fischer and Gaupp, 2005;  
 Bolton et al., 2006; Martínez and Escobar, 1995). The bedrock and river sediments of the Southern Alps are relatively lean in  
 kerogen (Copard et al., 2007; Horan et al., 2017). Therefore, for deglaciating landscapes exposing catchments exhibiting  
 glacially ground kerogen-rich lithologies, larger oxidation fluxes than those reported from the Southern Alps can be expected.  
from such increasing and decreasing the plausible ranges for  $C_R$  and  $A_D$ , respectively deglaciaded landscapes (see next section).

155 Long-term accelerated decline in the radiocarbon concentration of atmospheric CO<sub>2</sub> parallel to an overall increase in CO<sub>2</sub>  
amount occurred since the Last Glacial Maximum (Reimer et al., 2013; Roth and Joos, 2013). Consistent with such a dilutional  
process, kerogen oxidation releases radiocarbon-dead CO<sub>2</sub> to the atmosphere. During the time period of greatest CO<sub>2</sub> increase  
from 17.5 until 11.5 kyr before present, the rate of decline in the concentration of <sup>14</sup>C in atmospheric CO<sub>2</sub> is greatest and  
averages about 35 ‰/kyr (Fig. 3; c.f., Broecker and Barker (2007)). With the release of 500 PgC (radiocarbon-dead) to the  
160 atmosphere over deglaciation, a drop in 100 to 200 ‰ Δ<sup>14</sup>C of atmospheric CO<sub>2</sub> can be expected (see calculations by Zeng,  
2007; c.f., Sarntheim et al. (2013) who propose a comparable release yet argue for an oceanic mechanism). Kerogen oxidation  
would also release isotopically light carbon to the atmosphere, which is also consistent with the atmospheric record (Fig. 3;  
Bauska et al., 2016). Finally, driven by orbital forcing (Hays et al., 1976), the response in atmospheric CO<sub>2</sub> faithfully echoing  
increasing global temperatures and diminishing glaciated terrain (Sigman and Boyle, 2000; Stips et al., 2016) goes hand in  
165 hand with continuous aerial exposure and enhanced oxidation of finely ground, reactive kerogen (Fig. 2).

#### 4 A Canadian tale. And what about microbes?

~~Controlled to a first order by lithology, k~~Kerogen exposed on Earth's surface is distributed unevenly, with kerogen-rich surface  
lithologies extending ~~across~~long much of western Canada (see Fig. 2 in Copard et al., 2007). These areas experienced dramatic  
deglaciation across the 17.5-11.5 ka timeframe (Fig. 4; Dyke, 2004; Dalton et al., 2020). In addition to the widespread  
170 occurrence and high abundance of rock disseminated forms of kerogen, there are also notable surficial occurrences of  
hydrocarbon-rich lithologies including coal and oil sands within the deglaciated terrains of western Canada. Within ~~the~~  
~~Province of~~ Alberta~~province~~, Andriashek and Pawlowicz (2002) report the widespread occurrence of reworked shale and  
bitumen in Quaternary till stemming from Cretaceous shales ~~and~~ the oil sands, with unoxidized and oxidized forms present,  
and the latter enhanced by aerial exposure ~~across~~long paleosurfaces. Today, bituminous erratic boulders are found strewn  
175 around the region ~~over~~across tens of thousands of square kilometers (Rutherford, 1928; Andriashek, 2018). As Andriashek  
(2018) points out, Rutherford (1928) states: "There are perhaps many more occurrences of bituminous sand within the glacial  
deposits of Alberta, but since they weather comparatively readily and become covered with soil, they are not likely to be  
detected unless by accident, ...". This observation of rapid weathering of bitumen indicates that today's occurrences in glacial  
tills represent the tip of the iceberg of what once was present. ~~Laboratory incubations simulating~~ CO<sub>2</sub> respiration from  
180 bituminous materials~~based on laboratory incubations~~ reveal fluxes that are markedly higher ~~than those associated with~~ ~~in~~  
~~comparison to~~ oxidation of rock disseminated forms of kerogen (see Table 1). Chang and Berner (1998, 1999) report subaquatic  
bituminous coal oxidation (calculated  $>300 \frac{\text{MgC}}{\text{km}^2 \text{yr}}$  ~~MgC/km<sup>2</sup>/yr~~) releasing CO<sub>2</sub> ~~at rates greater by~~ 1-2 orders of magnitude ~~higher~~  
than ~~those values~~ reported ~~from~~or oxidation of rock disseminated kerogen, ~~and~~ 3 orders of magnitude greater than ~~for~~ the  
average ~~for~~ Earth's surface. Microbes likely play key roles in assimilating and releasing this ancient organic carbon as CO<sub>2</sub> to  
185 the atmosphere (ZoBell, 1946; Hemingway et al., 2018) with microbial communities on the surface colonizing oil sand-derived  
bitumen under both summer and winter conditions (Wyndham and Costerton, 1981; Wong et al., 2015). Biodegradation is



accelerated during summer months when ~~the~~ temperatures of subaerially exposed outcrops of oil sands reach 60°C (Wong et al., 2015). Microbial degradation experiments ~~on~~ bitumen (Ait-Langomazino et al., 1991) reveal an even greater CO<sub>2</sub> release when extrapolated to natural systems (e.g., oil sands, see Table 1) exceeding 18 ~~MgC/km<sup>2</sup>/day~~ <sup>MgC</sup>/<sub>km<sup>2</sup> day</sub>, over one hundred times

greater than the highest CO<sub>2</sub> flux released by rock disseminated kerogen oxidation. Other laboratory-based studies (e.g., Uribe-Alvarez et al., 2011) that investigated the oxidative decay of hydrocarbon fractions also suggest similarly high fluxes when scaled to natural systems, even though these studies were conducted over periods of only a few weeks~~studying oxidative decay of hydrocarbon fractions also suggest similarly astronomical fluxes when scaled to natural systems, which however only monitor over time periods of weeks~~. Such observations likely overestimate long-term natural fluxes over multiple years and decades, while the fluxes reported by Chang and Berner (1998, 1999) likely represent an underestimation due to sample storage for several years prior to analysis and the absence of microbial activity under their experimental conditions. Additionally, comparable quantities of kerogen-derived CO<sub>2</sub> can also be released under anaerobic conditions, which may become relevant under warming glaciers (e.g., Bertassoli Jr. et al., 2016; Rogieri Pelissari et al., 2021; see also Sharp et al., 1999; Sharp and Tranter, 2017). Overall, it is conceivable that super carbon source terrains (hypothetically, areas laced or covered with coal, bituminous materials, etc.) across western Canada could supply an overproportionate quantity of radiocarbon dead CO<sub>2</sub> to the atmosphere during glacial-interglacial transitions. Therefore, it is unclear how much such potential super carbon source terrains supply a disproportionate quantity of radiocarbon dead CO<sub>2</sub> to the atmosphere over glacial-interglacial transitions.

By 11.5 ka before present, the pronounced rise in atmospheric CO<sub>2</sub> had subsided. By this time, the Laurentide Ice Sheet had retreated beyond the easternmost reaches of the Western Canadian Sedimentary Basin and was ~~now~~ exposing bedrock from the relatively kerogen-poor, highly metamorphosed Canadian Shield (Fig. 4; Dalton et al., 2020). The timing of the inflection point in the rise of CO<sub>2</sub> occurs  $\leq 300$  years after the entire western edge of the Laurentide Ice Sheet advanced~~moves fully onto~~ the Canadian Shield suggests ~~its~~ diminishing~~abating~~ decay of kerogen and relaxation of the landscape. By this time, the rate of soil organic carbon sequestration in these paraglacial landscapes begins to pick up substantially (Harden et al., 1992).

Lithologies of the Fennoscandian Shield and most of its adjacent landmasses (deglaciated eastern and southern sectors of the Fennoscandian Ice Sheet~~field~~; see Stroeve et al., 2016 and Copard et al., 2007) also contain relatively low kerogen contents, therefore limiting ~~its the~~ contributions to atmospheric CO<sub>2</sub> rise in the wake of Fennoscandian Ice Sheet retreat. Glacially ground graphite shed from these highly metamorphosed rocks is~~behaves~~ chemically recalcitrant and is mostly redeposited (Sauramo, 1938; Sackett et al., 1974; Sparkes et al., 2020). Geophysical arguments propose that the relatively narrow timing of the Laurentide Ice Sheet's isochrons retreating into the Canadian Shield is more than coincidence: the lithological base and its overlying glacial till, influence the dynamics of ice sheet retreat with the Western Canadian Sedimentary Basin representing a deformable base while the Canadian Shield behaves rigidly providing an explanation for why the geological and ice sheet isochron map show similarities (Clark, 1994 and see Fig. 3 therein; Licciardi et al., 1998). At the same time in the context of

the kerogen oxidation hypothesis, this offers an explanation for the well-defined inflection point in the rise of atmospheric CO<sub>2</sub> characteristic for glacial-interglacial transitions.

In the bigger picture, during Earth's recent glacial episodes, the Laurentide Ice Sheet ~~represents the aerially most~~was the most extensive ~~element~~part of the cryosphere that waxed and waned across the continents (Batchelor et al., 2019) and in conjunction with its lithological underpinning (Copard et al., 2007), likely played the largest role in releasing kerogen-derived CO<sub>2</sub> to the atmosphere upon glacial retreat. However, estimates ~~of~~in CO<sub>2</sub> fluxes emanating from bedrock show a wide range (Table 1) and there is considerable uncertainty ~~resides~~ in our current state of knowledge: Dedicated biogeochemical weathering studies ~~are lacking which that~~ provide estimates of CO<sub>2</sub>carbon fluxes ~~emanating~~ from bedrock-derived kerogen under relevant environmental conditions and over appropriate timescales are lacking. Chronosequence studies of kerogen oxidation rates in deglaciated terrains are needed to provide constraints on time-integrated CO<sub>2</sub> release to the atmosphere. In tandem with this, quantification is needed for (temporary) kerogen reburial in subaerial and subaquatic terrestrial systems (e.g., moraines, lakes) on global and regional scales (e.g., Meybeck, 1993; Vonk et al., 2016; Blattmann et al., 2019b; Fox et al., 2020). Such information in conjunction with high resolution reconstructions of changes in land ice extent and the lithologies of bedrock and glacial till (including data on kerogen content and its stable carbon isotope composition) being exposed by glacial retreat can, in theory, quantitatively disentangle the contribution of kerogen-derived CO<sub>2</sub> to the atmosphere during glacial-interglacial transitions.

~~high-resolution spatiotemporal changes in land ice extent together with lithology of bedrock and glacial till can theoretically quantitatively disentangle the contribution of kerogen-derived CO<sub>2</sub> to the atmosphere during glacial-interglacial transitions.~~

## 5 Tackling geologic deep time

While modern day kerogen weathering and reburial efficiencies are only loosely constrained, even less is known about how they varied back in geologic time. Overarching controls on these processes include the mode of erosion and transport ranging from glacial to glaciofluvial to fluvial, remineralization intensity, controlled by continental margin type and geomorphology, and the intrinsic reactivity of the kerogen present locally (Blair and Aller, 2012; Blattmann et al., 2018b). Also important ~~is~~are the bedrock lithology and regolith composition which have been hypothesized to exert feedback on temporal patterns of glacial-interglacial cyclicity (e.g., Roy et al., 2004; Zeng, 2007). For the Paleocene-Eocene Thermal Maximum, an extreme greenhouse episode, evidence increasingly ~~suggests that~~points towards enhanced remineralization and leaching of kerogen (Boucsein and Stein, 2009; Lyons et al., 2019), possibly enhanced by the activity of microbes (Hemingway et al., 2018; Petsch et al., 2001), ~~thereby~~increas~~ed~~ing the flux of carbon entering actively circulating pools on Earth's surface. ~~Over~~Across Earth's history ~~on~~over 10<sup>9</sup>-year timescales, ~~the kerogen~~reburial efficiency of kerogen presumably varied as a function of atmospheric O<sub>2</sub> content, with lower O<sub>2</sub> contents tied to higher reburial efficiency (Daines et al., 2017). Kerogen is surmised to have acted as a major source of carbon to the atmosphere and as an "antioxidant" during the early rise of atmospheric O<sub>2</sub> (Daines et al., 2017; Kump et al., 2011) and balancing O<sub>2</sub> levels over the Cenozoic (Galvez, 2020; see also Derry and France-Lanord (1996)



~~for discussion on Paleogene~~). Across multiple glacial-interglacial cycles, enhanced kerogen oxidation would also be consistent with declining atmospheric O<sub>2</sub> ~~on~~<sup>over</sup> 10<sup>6</sup> yr timescales (Stolper et al., 2016).

In order to understand ~~changes in~~ atmospheric chemistry through geologic time, in addition to comprehensively budgeting the effect of mineral chemical weathering (Blattmann et al., 2019a; Horan et al., 2019; Hilton et al., 2014), ~~the changing efficiency of the oxidation and~~ reburial ~~efficiency~~ of ~~sedimentary~~ kerogen needs to be evaluated. Direct quantification of kerogen found reburied in sediments is often associated with considerable uncertainty owing to uncertainties in organic matter source apportionment (e.g., Lin et al., 2020; Blattmann et al., 2019b) and ~~uneven spatial geospatial dispersal variability~~ (e.g., Blattmann et al., 2018a; Cui et al., 2016). While radiocarbon was paramount ~~for quantifying and establishing the importance of the reburial of kerogen in recent times (Blattmann et al., 2018b), its utility diminishes quickly for strata that pre-date the Last Glacial Maximum owing to its radioactive decay.~~ ~~for establishing the importance of and quantifying kerogen reburial in the Recent (Blattmann et al., 2018b), the utility of radiocarbon quickly diminishes for strata preceding the Last Glacial Maximum owing to its radioactive decay.~~ However, associated with kerogen are a promising suite of trace elements and their respective isotope signatures including, among others, rhenium (Hilton et al., 2014; Horan et al., 2017), osmium (Georg et al., 2013; Ravizza and Esser, 1993), and iodine (Moran et al., 1998), that can be exploited to trace sedimentary kerogen and its degradation. In the case of osmium, seawater records reveal isotopic shifts at the beginning of interglacial ~~periods~~ that are attributable to ~~the oxidation of~~ kerogen oxidation (Georg et al., 2013; compare discussions in Peucker-Ehrenbrink and Ravizza, 2020), consistent with the ~~presented~~ hypothesis presented here. More research constraining the exogenous kerogen cycle by quantification of reburied kerogen inputs (e.g., iodine isotopes, organic petrology) and kerogen oxidation recorded by chemical weathering proxies (e.g., osmium isotopes) is needed to put the presented hypothesis to the test.

## 6 Synthesis and outlook

In a nutshell, the ~~presented~~ hypothesis presented proposes the following: Less than 300 years after the Laurentide Ice Sheet retreated east past the ~~western~~<sup>eastern</sup> most edges of the kerogen-rich Western Canadian Sedimentary Basin, atmospheric CO<sub>2</sub> levels stabilized and ~~its~~<sup>the</sup> rate of decrease in <sup>14</sup>C ~~concentrations of CO<sub>2</sub> decline~~ subsided (Figs. 3 and 4). This inflection point is mirrored in the ~~surficial~~ lithologies of the Canadian Shield that were exposed during deglaciation, which contain relatively minor amounts of reactive kerogen ~~which exposed comparatively minor amounts of reactive kerogen to the elements upon further deglaciation~~. Within the context of the presented hypothesis, the coincidence in time of global trends in atmospheric chemistry with spatiotemporal patterns in the distribution of ~~of~~ freshly exposed deglaciated terrain impregnated with oxidizable and biodegradable kerogen and bituminous materials suggests that a burst (or bursts) of respired CO<sub>2</sub> contributed to the characteristic deglacial ~~increase~~<sup>rebound</sup> in atmospheric CO<sub>2</sub>. As soon as glacially ground shales and bituminous materials either exhausted their labile kerogen content and/or became buried by soil and vegetation taking hold ~~on~~<sup>of</sup> the deglaciated landscape, classically considered processes (land-ocean exchange, biospheric uptake, etc.) reassumed dominant control on fluctuations of atmospheric chemistry. As a corollary of this hypothesis: CO<sub>2</sub> rise within the envelope of glacial-interglacial

cyclicity would primarily have responded as a slave to orbitally controlled glacial retreat across kerogen impregnated landscapes as a result of global temperature increase. CO<sub>2</sub> rise may therefore lead or lag global temperature depending on the spatiotemporal patterns of glacial retreat ~~that~~ exposing glacially ground, kerogen-rich impregnated or even bituminous parent material.

Various mechanisms ~~for explaining CO<sub>2</sub> increases at glacial interglacial transitions~~ have been proposed to explain CO<sub>2</sub> increases at glacial interglacial transitions including for example the solubility pump hypothesis, iron fertilization hypothesis, ocean circulation hypotheses and many more (e.g. see hypotheses and reviews by Martin, 1990; Broecker and Peng, 1993; Kohfeld and Ridgwell, 2009; Rapp, 2019). However, as a cogwheel operating under manifold feedbacks in the greater Earth system (Sigman and Boyle, 2000), continuous glacial retreat, and the oxidation of finely ground kerogen, provides a hypothesis consistent with a CO<sub>2</sub> increase in the wake of glacial episodes. This dilution of radiocarbon-dead CO<sub>2</sub> in the atmosphere may well have been complemented by other terrestrial sources such as the oxidation of subglacial paleosols and permafrost-bound organic carbon oxidation (Zeng, 2007; Simmons et al., 2016), ~~permafrost-bound organic carbon oxidation~~ (Tesi et al., 2016; Winterfeld et al., 2018; Lindgren et al., 2018; Köhler et al., 2014; Ciais et al., 2012), and by ~~(time delayed)~~ volcanic emissions triggered by due to deglacial unloading of the lithosphere (Roth and Joos, 2012). In addition to studies of weathering in glacial forefields and source-to-sink tracing of sedimentary kerogen, several lines of geochemical evidence including atmospheric carbon isotope ~~ie~~ composition (<sup>13</sup>C and <sup>14</sup>C), which have thus far received contorted, partial explanations (Broecker and Clark, 2010; Schmitt et al., 2012; Broecker and McGee, 2013), glacial-interglacial changes in <sup>14</sup>C of dissolved inorganic carbon in seawater (Rafter et al., 2019, c.f., discussions therein), seawater osmium isotope changes, and long-term atmospheric O<sub>2</sub> content, conceptually go hand-in-hand with an opening of the exogenous kerogen cycle modulated by glacial activity. While geomagnetic variability and ocean ventilation together struggle to fully explain observed changes in atmospheric radiocarbon (Broecker and Barker, 2007; Cheng et al., 2018), the dilution of atmospheric CO<sub>2</sub> by accelerated oxidation of ancient terrestrial organic carbon ~~oxidation~~ at glacial terminations, in conjunction with other mechanisms including atmosphere-ocean gas exchange (e.g., Sigman et al., 2010; Marcott et al., 2014; Menviel et al., 2018; Martin, 1990; Sarntheim et al., 2013) appears as a simple and plausible explanation. In addition to kerogen oxidation at glacial terminations, the ~~presented~~ hypothesis presented carries other important implications, including the closure of the exogenous kerogen cycle during glacial periods (Fig. 2) potentially contributing to relatively high atmospheric <sup>14</sup>C signatures, a problem highlighted by Dincau et al. (2020), as a reduced <sup>14</sup>C-free CO<sub>2</sub> flux would reduce the dilution of the atmosphere's cosmogenic <sup>14</sup>C. Overall, increased reburial efficiency of kerogen can account for several tens of PgC over millennial timescales, entirely bypassing actively circulating carbon pools on Earth's surface (Fig. 1). In contrast, increased oxidation efficiency of kerogen in the wake of glacial episodes that have built up stores of finely ground reactive substrate and exposed fresh weathering profiles can account for several hundreds of PgC over millennial timescales released into actively circulating pools. Owing to increased bedrock exhumation over the Ice Ages (Herman et al., 2013; Herman et al., 2015) the dynamism of the exogenous kerogen cycle may have been intensified, with greater fluxes of detrital kerogen reburied in ocean sediments during glacial episodes and ~~increased enhanced~~ supplies of ground kerogen exposed to the elements in their wake.

While basic controls on kerogen reburial efficiency have emerged, its quantitative impact on atmospheric chemistry through geologic time remains conjectural. Mathematically analogous modeling results (Zeng, 2003; Simmons et al., 2016) suggest that kerogen oxidation could account for 30-60 ppm rise in atmospheric CO<sub>2</sub> over the course of the last deglaciation. However, there is a lack of data on kerogen weathering to provide sufficient quantitative constraint for testing this hypothesis. To work towards such a test, investigating kinetics of kerogen oxidation along glacial chronosequences with contrasting lithologies would provide numerical input for Earth system models. Such information in conjunction with spatiotemporal changes in land ice extent (e.g., Dalton et al., 2020; Stroeve et al., 2016) integrating over areal changes of glacially exposed lithological units (c.f., Copard et al., 2007) would constrain the quantitative impact of the exogenous kerogen cycle on atmospheric chemistry over glacial-interglacial cycles (e.g., see Fig. 2 in Kohfeld and Ridgwell, 2009). Additionally, the role of super carbon source terrains (e.g., surficial oil sands and coal) deserve special attention as biogeochemical weathering and associated CO<sub>2</sub> fluxes emanating from such areas conceivably contribute disproportionately, particularly upon deglaciation. Overall, the spatiotemporal deglaciation of contrasting source terrains across North America with the coeval progression of atmospheric chemistry provide a strong incentive to explore the potential role of the exogenous kerogen cycle in glacial-interglacial patterns. From a greenhouse perspective, further study of pivotal episodes such as the Paleocene-Eocene Thermal Maximum under this lens may provide an outlook for geological processes that is relevant today. In the context of our warming world, once critical thresholds are breached (Steffen et al., 2018), enhanced opening of the exogenous kerogen cycle may entrain the Earth system onto a new trajectory influencing the carbon cycle and climate for millennia to come.

## Acknowledgments

This work greatly benefitted from the input of ~~several~~<sup>two</sup> anonymous reviewers. The author is thankful for discussions with Timothy Eglinton, Dominik Letsch, Maarten Lupker, Jesper Suhrhoff, Valier Galy, [Naohiko Ohkouchi](#), and Robert Hilton. This work was supported by funding from JAMSTEC.

## Competing interests

The author declares that he has no conflict of interest.

## Author contributions

TMB conceived of and wrote this contribution.

## References

- Ait-Langomazino, N., Sellier, R., Jouquet, G. and Trescinski, M.: Microbial degradation of bitumen, *Experientia*, 47, 533-539, 1991.
- 345 Andriashek, L. D.: On the origin of oil sand and related bedrock erratics in glacial sediments of central Alberta; Alberta Energy Regulator / Alberta Geological Survey, AER/AGS Open File Report 2018-13, 42 p, 2018.
- Andriashek, L. D. and Pawlowicz, J. G.: Observations of naturally occurring hydrocarbons (bitumen) in Quaternary sediments, Athabasca Oil Sands area and areas west, Alberta; Alberta Energy and Utilities Board, EUB/AGS Geo-Note 2002-01, 27 p, 2002.
- 350 [Barnola, J. M., Raynaud, D., Korotkevich, Y. S., and Lorius, C.: Vostok ice core provides 160,000-year record of atmospheric CO<sub>2</sub>, \*Nature\*, 329, 408-414, 1987.](#)
- Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., Stokes, C. R., Murton, J. B. and Manica, A.: The configuration of Northern Hemisphere ice sheets through the Quaternary, *Nat. Commun.*, 10, 3713, 2019.
- Bauska T. K., Baggenstos D., Brook E. J., Mix A. C., Marcott S. A., Petrenko V. V., Schaefer H., Severinghaus J. P. and Lee
- 355 J. E.: Carbon isotopes characterize rapid changes in atmospheric carbon dioxide during the last deglaciation. *Proc. Natl. Acad. Sci. U.S.A.*, 113, 3465-3470, 2016.
- Berner R. A.: Atmospheric carbon dioxide levels over Phanerozoic time. *Science*, 249, 1382-1386, 1990.
- [Berner, R. A., Lasaga, A. C., and Garrels, R. M.: The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. \*Am. J. Sci.\*, 283, 641, 1983.](#)
- 360 Bertassoli Jr., D. J., Sawakuchi, H. O., Almeida, N. S., Castanheira, B., Alem, V. A. T., Camargo, M. G. P., Krusche, A. V., Brochsztain, S. and Sawakuchi, A. O.: Biogenic methane and carbon dioxide generation in organic-rich shales from southeastern Brazil, *Int. J. Coal Geol.*, 162, 1-13, 2016.
- Bird, M. I., Lloyd, J. and Farquhar, G. D.: Terrestrial carbon storage at the LGM. *Nature* 371, 566-566, 1994.
- Blair N. E. and Aller R. C.: The fate of terrestrial organic carbon in the marine environment. *Annu. Rev. Mar. Sci.*, 4, 401-
- 365 423, 2012.
- Blattmann T. M., Zhang Y., Zhao Y., Wen K., Lin S., Li J., Wacker L., Haghipour N., Plötze M., Liu Z. and Eglinton T. I.: Contrasting fates of petrogenic and biospheric carbon in the South China Sea. *Geophys. Res. Lett.*, 45, 9077-9086, 2018a.
- Blattmann T. M., Letsch D. and Eglinton T. I.: On the geological and scientific legacy of petrogenic organic carbon. *Am. J. Sci.*, 318, 861-881, 2018b.
- 370 Blattmann T. M., Wang S. L., Lupker M., Märki L., Haghipour N., Wacker L., Chung L. H., Bernasconi S. M., Plötze M. and Eglinton T. I.: Sulphuric acid-mediated weathering on Taiwan buffers geological atmospheric carbon sinks. *Sci. Rep.*, 9, 2945, 2019a.
- Blattmann T. M., Wessels M., McIntyre C. P. and Eglinton T. I.: Petrogenic organic carbon retention in terrestrial basins: A case study from perialpine Lake Constance. *Chem. Geol.*, 503, 52-60, 2019b.

- 375 Bolton E. W., Berner R. A. and Petsch S. T.: The weathering of sedimentary organic matter as a control on atmospheric O<sub>2</sub>: II. Theoretical modelling. *Am. J. Sci.*, 306, 575-615, 2006.
- Boucsein B. and Stein R.: Black shale formation in the late Paleocene/early Eocene Arctic Ocean and paleoenvironmental conditions: New results from a detailed organic petrological study. *Mar. Petroleum Geol.*, 26, 416-426, 2009.
- Broecker W. and Clark E.: Search for a glacial-age <sup>14</sup>C-depleted ocean reservoir. *Geophys. Res. Lett.*, 37, 2010.
- 380 Broecker, W. S. and McGee, D.: The <sup>13</sup>C record for atmospheric CO<sub>2</sub>: What is it trying to tell us? *Earth Planet. Sci. Lett.*, 368, 175-182, 2013.
- Broecker, W. and Barker, S.: A 190‰ drop in atmosphere's Δ<sup>14</sup>C during the “Mystery Interval” (17.5 to 14.5 kyr). *Earth Planet. Sci. Lett.*, 256, 90-99, 2017.
- Broecker, W. S. and Peng, T. H.: What caused the glacial to interglacial CO<sub>2</sub> change?, *in*: Heimann, M. *The Global Carbon Cycle*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 95-115, 1993.
- 385 [Bufe, A., Hovius, N., Emberson, R., Rugenstein, J. K. C., Galy, A., Hassenruck-Gudipati, H. J. and Chang, J.-M.: Co-variation of silicate, carbonate and sulfide weathering drives CO<sub>2</sub> release with erosion, \*Nat. Geosci.\*, 14, 211-216, 2021.](#)
- Chang, S. and Berner, R. A.: Humic substance formation via the oxidative weathering of coal, *Environ. Sci. Technol.*, 32, 2883-2886, 1998.
- 390 Chang, S. and Berner, R. A.: Coal weathering and the geochemical carbon cycle, *Geochim. Cosmochim. Acta*, 63, 3301-3310, 1999.
- Cheng, H., Edwards, R. L., Southon, J., Matsumoto, K., Feinberg, J. M., Sinha, A., Zhou, W., Li, H., Li, X., Xu, Y., Chen, S., Tan, M., Wang, Q., Wang, Y. and Ning, Y.: Atmospheric <sup>14</sup>C/<sup>12</sup>C changes during the last glacial period from Hulu Cave. *Science*, 362, 1293-1297, 2018.
- 395 Ciais, P., Tagliabue, A., Cuntz, M., Bopp, L., Scholze, M., Hoffmann, G., Lourantou, A., Harrison, S. P., Prentice, I. C., Kelley, D. I., Koven, C. and Piao, S. L.: Large inert carbon pool in the terrestrial biosphere during the Last Glacial Maximum, *Nat. Geosci.*, 5, 74-79, 2012.
- Clark, M. G., Drewitt, G. B. and Carey, S. K.: Energy and carbon fluxes from an oil sands pit lake, *Sci. Total Environ.*, 752, 141966, 2021.
- 400 [Clark, P. U.: Unstable behavior of the Laurentide Ice Sheet over deforming sediment and its implications for climate change, \*Quat. Res.\*, 41, 19-25, 1994.](#)
- Copard, Y., Amiotte-Suchet, P. and Di-Giovanni, C.: Storage and release of fossil organic carbon related to weathering of sedimentary rocks, *Earth Planet. Sci. Lett.*, 258, 345-357, 2007.
- Cui, X., Bianchi, T. S., Jaeger, J. M. and Smith, R. W.: Biospheric and petrogenic organic carbon flux along southeast Alaska, *Earth Planet. Sci. Lett.*, 452, 238-246, 2016.
- 405 Daines S. J., Mills B. J. W. and Lenton T. M.: Atmospheric oxygen regulation at low Proterozoic levels by incomplete oxidative weathering of sedimentary organic carbon. *Nat. Commun.*, 8, 14379, 2017.

- Dalton, A. S., Margold, M., Stokes, C. R., Tarasov, L., Dyke, A. S., Adams, R. S., Allard, S., Arends, H. E., Atkinson, N., Attig, J. W., Barnett, P. J., Barnett, R. L., Batterson, M., Bernatchez, P., Borns, H. W., Breckenridge, A., Briner, J. P., Brouard, E., Campbell, J. E., Carlson, A. E., Clague, J. J., Curry, B. B., Daigneault, R.-A., Dubé-Loubert, H., Easterbrook, D. J., Franzi, D. A., Friedrich, H. G., Funder, S., Gauthier, M. S., Gowan, A. S., Harris, K. L., Hétu, B., Hooyer, T. S., Jennings, C. E., Johnson, M. D., Kehew, A. E., Kelley, S. E., Kerr, D., King, E. L., Kjeldsen, K. K., Knaeble, A. R., Lajeunesse, P., Lakeman, T. R., Lamothe, M., Larson, P., Lavoie, M., Loope, H. M., Lowell, T. V., Lusardi, B. A., Manz, L., McMartin, I., Nixon, F. C., Occhietti, S., Parkhill, M. A., Piper, D. J. W., Pronk, A. G., Richard, P. J. H., Ridge, J. C., Ross, M., Roy, M., Seaman, A., Shaw, J., Stea, R. R., Teller, J. T., Thompson, W. B., Thorleifson, L. H., Utting, D. J., Veillette, J. J., Ward, B. C., Weddle, T. K. and Wright, H. E.: An updated radiocarbon-based ice margin chronology for the last deglaciation of the North American Ice Sheet Complex. *Quat. Sci. Rev.*, 234, 106223, 2020.
- [Derry, L. A. and France-Lanord, C.: Neogene growth of the sedimentary organic carbon reservoir, \*Paleoceanography\*, 11, 267-275, 1996.](#)
- Dinauer, A., Adolphi, F. and Joos, F.: Mysteriously high  $\Delta^{14}\text{C}$  of the glacial atmosphere: influence of  $^{14}\text{C}$  production and carbon cycle changes. *Clim. Past*, 16, 1159-1185, 2020.
- Dyke, A. S.: An outline of North American deglaciation with emphasis on central and northern Canada, *in*: Ehlers, J., Gibbard, P. L. *Developments in Quaternary Sciences*. Elsevier, pp. 373-424, 2004.
- Fischer, C. and Gaupp, R.: Change of black shale organic material surface area during oxidative weathering: Implications for rock-water surface evolution, *Geochim. Cosmochim. Acta*, 69, 1213-1224, 2005.
- Fischer, C., Karius, V. and Thiel, V.: Organic matter in black slate shows oxidative degradation within only a few decades, *J. Sediment. Res.*, 77, 355-365, 2007.
- [Fox, P. M., Bill, M., Heckman, K., Conrad, M., Anderson, C., Keiluweit, M. and Nico, P. S.: Shale as a source of organic carbon in floodplain sediments of a mountainous watershed, \*J. Geophys. Res. Biogeosci.\*, 125, 2020.](#)
- [Galvez, M. E.: Redox constraints on a Cenozoic imbalance in the organic carbon cycle, \*Am. J. Sci.\*, 320, 730, 2020.](#)
- Galy V., Beyssac O., France-Lanord C. and Eglinton T. I.: Recycling of graphite during Himalayan erosion: A geological stabilization of carbon in the crust. *Science*, 322, 943-945, 2008.
- Galy V., Peucker-Ehrenbrink B. and Eglinton T. I.: Global carbon export from the terrestrial biosphere controlled by erosion. *Nature*, 521, 204-207, 2015.
- Georg R. B., West A. J., Vance D., Newman K. and Halliday A. N.: Is the marine osmium isotope record a probe for  $\text{CO}_2$  release from sedimentary rocks? *Earth Planet. Sci. Lett.*, 367, 28-38, 2013.
- [Gibbs, M., and Kump, L.: Global chemical weathering during deglaciation, \*Fourth International Symposium on the Geochemistry of the Earth's Surface, University of Leeds\*, 733-737, 1996.](#)
- Hain, M. P., Sigman, D. M. and Haug, G. H.: Distinct roles of the Southern Ocean and North Atlantic in the deglacial atmospheric radiocarbon decline. *Earth Planet. Sci. Lett.*, 394, 198-208, 2014.



- Harden, J. W., Mark, R. K., Sundquist, E. T. and Stallard, R. F.: Dynamics of soil carbon during deglaciation of the Laurentide Ice Sheet, *Science*, 258, 1921, 1992.
- Hays J. D., Imbrie J. and Shackleton N. J.: Variations in the Earth's orbit: Pacemaker of the Ice Ages. *Science*, 194, 1121-1132, 1976.
- 445 Hedges J. I. and Oades J. M.: Comparative organic geochemistries of soils and marine sediments. *Org. Geochem.*, 27, 319-361, 1997.
- Herman F., Seward D., Valla P. G., Carter A., Kohn B., Willett S. D. and Ehlers T. A.: Worldwide acceleration of mountain erosion under a cooling climate. *Nature*, 504, 423-426, 2013.
- Herman F., Beyssac O., Brughelli M., Lane S. N., Leprince S., Adatte T., Lin J. Y. Y., Avouac J.-P. and Cox S. C.: Erosion  
450 by an Alpine glacier. *Science*, 350, 193-195, 2015.
- Hemingway J. D., Hilton R. G., Hovius N., Eglinton T. I., Haghpor N., Wacker L., Chen M.-C. and Galy V. V.: Microbial oxidation of lithospheric organic carbon in rapidly eroding tropical mountain soils. *Science*, 360, 209, 2018.
- [Hilton, R. G., and West, A. J.: Mountains, erosion and the carbon cycle, \*Nat. Rev. Earth Environ.\*, 1, 284-299, 2020.](#)
- Hilton R. G., Gaillardet J., Calmels D. and Birck J.-L.: Geological respiration of a mountain belt revealed by the trace element  
455 rhenium. *Earth Planet. Sci. Lett.*, 403, 27-36, 2014.
- Hood E., Fellman J., Spencer R. G. M., Hernes P. J., Edwards R., D'Amore D. and Scott D.: Glaciers as a source of ancient and labile organic matter to the marine environment. *Nature*, 462, 1044, 2009.
- Horan K., Hilton R. G., Dellinger M., Tipper E., Galy V., Calmels D., Selby D., Gaillardet J., Ottley C. J., Parsons D. R. and  
460 Burton K. W. Carbon dioxide emissions by rock organic carbon oxidation and the net geochemical carbon budget of the Mackenzie River Basin. *Am. J. Sci.*, 319, 473-499, 2019.
- Horan K., Hilton R. G., Selby D., Ottley C. J., Gröcke D. R., Hicks M. and Burton K. W.: Mountain glaciation drives rapid oxidation of rock-bound organic carbon. *Sci. Adv.*, 3, 2017.
- Horan K.: The oxidative weathering of organic matter and its carbon dioxide emissions: Insight from the trace elements rhenium and molybdenum, Ph.D. thesis, Durham University, United Kingdom, 223 pp., 2018.
- 465 [Kohfeld, K. E. and Ridgwell, A.: Glacial-interglacial variability in atmospheric CO<sub>2</sub>, in: \*Surface Ocean–Lower Atmosphere Processes\*, edited by: Le Quéré, C., and Saltzman, E. S., \*Geophys. Monogr. Ser.\*, \*American Geophysical Union\*, 251-286, 2009.](#)
- Köhler, P., Knorr, G. and Bard, E.: Permafrost thawing as a possible source of abrupt carbon release at the onset of the Bølling/Allerød. *Nat. Commun.* 5, 5520, 2014.
- [Kölling, M., Bouimtarhan, I., Bowles, M. W., Felis, T., Goldammer, T., Hinrichs, K.-U., Schulz, M. and Zabel, M.: Consistent CO<sub>2</sub> release by pyrite oxidation on continental shelves prior to glacial terminations, \*Nat. Geosci.\*, 12, 929-934, 2019.](#)  
470
- Kump L. R., Junium C., Arthur M. A., Brasier A., Fallick A., Melezhik V., Lepland A., Črne A. E. and Luo G.: Isotopic evidence for massive oxidation of organic matter following the Great Oxidation Event. *Science*, 334, 1694-1696, 2011.

- 475 [Lewan, M. D.: Stable carbon isotopes of amorphous kerogens from Phanerozoic sedimentary rocks, \*Geochim. Cosmochim. Acta\*, 50, 1583-1591, 1986.](#)
- [Leythaeuser, D.: Effects of weathering on organic matter in shales, \*Geochim. Cosmochim. Acta\*, 37, 113-120, 1973.](#)
- [Licciardi, J. M., Clark, P. U., Jenson, J. W. and Macayeal, D. R.: Deglaciation of a soft-bedded Laurentide Ice Sheet, \*Quat. Sci. Rev.\*, 17, 427-448, 1998.](#)
- 480 Lin, B., Liu, Z., Eglinton, T. I., Kandasamy, S., Blattmann, T. M., Haghipour, N., Huang, K.-F. and You, C.-F.: Island-wide variation in provenance of riverine sedimentary organic carbon: A case study from Taiwan, *Earth Planet. Sci. Lett.*, 539, 116238, 2020.
- [Lindgren, A., Hugelius, G. and Kuhry, P.: Extensive loss of past permafrost carbon but a net accumulation into present-day soils, \*Nature\*, 560, 219-222, 2018.](#)
- 485 Littke, R., Klusmann, U., Krooss, B. and Leythaeuser, D.: Quantification of loss of calcite, pyrite, and organic matter due to weathering of Toarcian black shales and effects on kerogen and bitumen characteristics, *Geochim. Cosmochim. Acta*, 55, 3369-3378, 1991.
- Lyons S. L., Baczynski A. A., Babila T. L., Bralower T. J., Hajek E. A., Kump L. R., Polites E. G., Self-Trail J. M., Trampush S. M., Vornlocher J. R., Zachos J. C. and Freeman K. H.: Palaeocene–Eocene Thermal Maximum prolonged by fossil carbon oxidation. *Nat. Geosci.*, 12, 54-60, 2019.
- 490 Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T. J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C., White, J. W. C. and Brook, E. J.: Centennial-scale changes in the global carbon cycle during the last deglaciation, *Nature*, 514, 2014.
- Martin, J. H.: Glacial-interglacial CO<sub>2</sub> change: The iron hypothesis, *Paleoceanography*, 5, 1-13, 1990.
- 495 [Martínez, M., and Escobar, M.: Effect of coal weathering on some geochemical parameters, \*Org. Geochem.\*, 23, 253-261, 1995.](#)
- Menviel, L., Spence, P., Yu, J., Chamberlain, M. A., Mearns, R. J., Meissner, K. J. and England, M. H.: Southern Hemisphere westerlies as a driver of the early deglacial atmospheric CO<sub>2</sub> rise, *Nat. Commun.*, 9, 2503, 2018.
- [Meybeck, M.: Riverine transport of atmospheric carbon: Sources, global typology and budget, \*Water Air Soil Pollut.\*, 70, 443-463, 1993.](#)
- 500 [Meyers, P. A.: Preservation of elemental and isotopic source identification of sedimentary organic matter, \*Chem. Geol.\*, 114, 289-302, 1994.](#)
- Miall, A. D. and Blakey, R. C.: The Phanerozoic tectonic and sedimentary evolution of North America, in: *The Sedimentary Basins of the United States and Canada (Second Edition)*, edited by: Miall, A. D., Elsevier, 1-38, 2019.
- Moran J. E., Fehn U. and Teng R. T. D.: Variations in <sup>129</sup>I/<sup>127</sup>I ratios in recent marine sediments: evidence for a fossil organic component. *Chem. Geol.*, 152, 193-203, 1998.
- 505 [Munhoven, G., and François, L. M.: Glacial-interglacial variability of atmospheric CO<sub>2</sub> due to changing continental silicate rock weathering: A model study, \*J. Geophys. Res. Atmos.\*, 101, 21423-21437, 1996.](#)

- Petsch S. T., Eglinton T. I. and Edwards K. J.:  $^{14}\text{C}$ -dead living biomass: Evidence for microbial assimilation of ancient organic carbon during shale weathering. *Science*, 292, 1127-1131, 2001.
- 510 Petsch S. T.: Weathering of organic carbon, *in*: Holland H.D., Turekian K.K., Treatise on Geochemistry. Elsevier, Oxford, pp. 217-238, 2014.
- Peucker-Ehrenbrink, B. and Ravizza, G. E.: Osmium isotope stratigraphy, *in*: Geologic Time Scale 2020, edited by: Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M., Elsevier, 239-257, 2020.
- Rafter, P. A., Carriquiry, J. D., Herguera, J.-C., Hain, M. P., Solomon, E. A. and Southon, J. R.: Anomalous > 2000-year-old  
515 surface ocean radiocarbon age as evidence for deglacial geologic carbon release. *Geophys. Res. Lett.*, 46, 13950-13960, 2019.
- Rapp, D.: Ice Ages and Interglacials: Measurements, Interpretation, and Models. 3<sup>rd</sup> edition, Springer Nature Switzerland, pp. 346, 2019.
- Ravizza G. and Esser B. K.: A possible link between the seawater osmium isotope record and weathering of ancient sedimentary organic matter. *Chem. Geol.*, 107, 255-258, 1993.
- 520 [Reed, J. C. Jr., Wheeler, J. O. and Tucholke, B. E.: Geologic map of North America: Decade of North American geology continental scale map 001, Boulder, Geological Society of America, scale 1:5,000,000, 2004.](#)
- Reimer P. J., Bard E., Bayliss A., Beck J. W., Blackwell P. G., Ramsey C. B., Buck C. E., Cheng H., Edwards R. L., Friedrich M., Grootes P. M., Guilderson T. P., Hafliðason H., Hajdas I., Hatté C., Heaton T. J., Hoffmann D. L., Hogg A. G., Hughen K. A., Kaiser K. F., Kromer B., Manning S. W., Niu M., Reimer R. W., Richards D. A., Scott E. M., Southon J. R., Staff R.  
525 A., Turney C. S. M. and van der Plicht J.: IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon*, 55, 1869-1887, 2013.
- Roffey, R. and Norqvist, A.: Biodegradation of bitumen used for nuclear waste disposal, *Experientia*, 47, 539-542, 1991.
- Rogieri Pelissari, M., Oliveira Sawakuchi, H., Bertassoli Junior, D. J., da Silva Almeida, N. and Oliveira Sawakuchi, A.: Water influence on  $\text{CH}_4$  and  $\text{CO}_2$  generation from tar sandstones: Insights from incubation experiments in the Pirambóia Formation,  
530 Paraná Basin, *J. S. Am. Earth Sci.*, 106, 103097, 2021.
- Roth R. and Joos F.: Model limits on the role of volcanic carbon emissions in regulating glacial–interglacial  $\text{CO}_2$  variations. *Earth Planet. Sci. Lett.*, 329-330, 141-149, 2012.
- Roth R. and Joos F.: A reconstruction of radiocarbon production and total solar irradiance from the Holocene  $^{14}\text{C}$  and  $\text{CO}_2$  records: implications of data and model uncertainties. *Clim. Past*, 9, 1879-1909, 2013.
- 535 Roy, M., Clark, P. U., Raisbeck, G. M. and Yiou, F.: Geochemical constraints on the regolith hypothesis for the middle Pleistocene transition, *Earth Planet. Sci. Lett.*, 227, 281-296, 2004.
- Rutherford R. L.; Two interesting boulders in the glacial deposits of Alberta. *J. Geol.*, 36, 558-563, 1928.
- [Sackett, W. M., Poag, C. W., and Eadie, B. J.: Kerogen recycling in the Ross Sea, Antarctica, \*Science\*, 185, 1045-1047, 1974.](#)
- Sarnthein, M., Schneider, B. and Grootes, P. M.: Peak glacial  $^{14}\text{C}$  ventilation ages suggest major draw-down of carbon into the  
540 abyssal ocean, *Clim. Past*, 9, 2595-2614, 2013.
- Sauramo M. R.: The mode of occurrence of carbon in Quaternary deposits. *Suomen Kemistilehti*, 3, 11-16, 1938.

Schachtman, N. S., Roering, J. J., Marshall, J. A., Gavin, D. G., and Granger, D. E.: The interplay between physical and chemical erosion over glacial-interglacial cycles, *Geology*, 47, 613-616, 2019.

- Schillawski S. and Petsch S.: Release of biodegradable dissolved organic matter from ancient sedimentary rocks. *Global Biogeochem. Cycles*, 22, 2008.
- Schmitt J., Schneider R., Elsig J., Leuenberger D., Laurantou A., Chappellaz J., Köhler P., Joos F., Stocker T. F., Leuenberger M. and Fischer H.: Carbon isotope constraints on the deglacial CO<sub>2</sub> rise from ice cores. *Science*, 336, 711-714, 2012.
- Shackleton N. J.: Carbon-13 in Uvigerina: Tropical rainforest history and the equatorial Pacific carbonate dissolution cycles, *in*: Andersen N. R. and Malahoff A. The Fate of Fossil Fuel CO<sub>2</sub> in the Oceans. Plenum Press, pp. 401-427, 1997.
- 550 Shackleton N. J., Hall M. A., Line J. and Shuxi C.: Carbon isotope data in core V19-30 confirm reduced carbon dioxide concentration in the ice age atmosphere. *Nature* 306, 319-322, 1983.
- Sharp M., Parkes J., Cragg B., Fairchild I. J., Lamb H. and Tranter M.: Widespread bacterial populations at glacier beds and their relationship to rock weathering and carbon cycling. *Geology* 27, 107-110, 1999.
- Sharp, M. and Tranter, M.: Glacier biogeochemistry, *Geochem. Perspect.*, 6, 173-174, 2017.
- 555 Sigman D. M., Hain M. P. and Haug G. H.: The polar ocean and glacial cycles in atmospheric CO<sub>2</sub> concentration. *Nature* 466, 47-55, 2010.
- Sigman D. M. and Boyle E. A.: Glacial/interglacial variations in atmospheric carbon dioxide. *Nature*, 407, 859-869, 2000.
- Simmons, C. T., Matthews, H. D. and Mysak, L. A.: Deglacial climate, carbon cycle and ocean chemistry changes in response to a terrestrial carbon release. *Clim. Dyn.*, 46, 1287-1299, 2016.
- 560 Smith H. J., Fischer H., Wahlen M., Mastroianni D. and Deck B.: Dual modes of the carbon cycle since the Last Glacial Maximum. *Nature*, 400, 248-250, 1999.
- Sparkes, R. B., Hovius, N., Galy, A. and Liu, J. T.: Survival of graphitized petrogenic organic carbon through multiple erosional cycles, *Earth Planet. Sci. Lett.*, 531, 115992, 2020.
- Steffen W., Rockström J., Richardson K., Lenton T. M., Folke C., Liverman D., Summerhayes C. P., Barnosky A. D., Cornell S. E., Crucifix M., Donges J. F., Fetzer I., Lade S. J., Scheffer M., Winkelmann R. and Schellnhuber H. J.: Trajectories of the Earth system in the Anthropocene. *Proc. Natl. Acad. Sci. U.S.A.*, 115, 8252, 2018.
- Stips A., Macias D., Coughlan C., Garcia-Gorriz E. and Liang X. S.: On the causal structure between CO<sub>2</sub> and global temperature. *Sci. Rep.*, 6, 21691, 2016.
- Stolper D. A., Bender M. L., Dreyfus G. B., Yan Y. and Higgins J. A.: A Pleistocene ice core record of atmospheric O<sub>2</sub>
- 570 concentrations. *Science*, 353, 1427-1430, 2016.
- Stroeven, A. P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B. W., Harbor, J. M., Jansen, J. D., Olsen, L., Caffee, M. W., Fink, D., Lundqvist, J., Rosqvist, G. C., Strömberg, B. and Jansson, K. N.: Deglaciation of Fennoscandia. *Quat. Sci. Rev.*, 147, 91-121, 2016.


- Tesi T., Muschitiello F., Smittenberg R. H., Jakobsson M., Vonk J. E., Hill P., Andersson A., Kirchner N., Noormets R.,  
575 Dudarev O., Semiletov I. and Gustafsson Ö.: Massive remobilization of permafrost carbon during post-glacial warming. *Nat. Commun.* 7, 13653, 2016.
- Torres M. A., West A. J. and Li G.: Sulphide oxidation and carbonate dissolution as a source of CO<sub>2</sub> over geological timescales. *Nature*, 507, 346-349, 2014.
- Torres M. A., Moosdorf N., Hartmann J., Adkins J. F. and West A. J.: Glacial weathering, sulfide oxidation, and global carbon  
580 cycle feedbacks. *Proc. Natl. Acad. Sci. U.S.A.*, 1-6, 2017.
- [Tranter, M.: Glacial runoff as a sink for atmospheric CO<sub>2</sub> during the last glacial-interglacial transition, Fourth International Symposium on the Geochemistry of the Earth's Surface, University of Leeds, 709-713, 1996.](#)
- Vance D., Teagle D. A. H. and Foster G. L.: Variable Quaternary chemical weathering fluxes and imbalances in marine  
geochemical budgets. *Nature*, 458, 493-496, 2009.
- 585 [Vonk, J. E., Dickens, A. F., Giosan, L., Hussain, Z. A., Kim, B., Zipper, S. C., Holmes, R. M., Montluçon, D. B., Galy, V. and Eglinton, T. I.: Arctic deltaic lake sediments as recorders of fluvial organic matter deposition, \*Front. Earth Sci.\*, 4, 2016.](#)
- Winterfeld, M., Mollenhauer, G., Dummann, W., Köhler, P., Lembke-Jene, L., Meyer, V. D., Hefter, J., McIntyre, C., Wacker, L., Kokfelt, U. and Tiedemann, R.: Deglacial mobilization of pre-aged terrestrial carbon from degrading permafrost. *Nat. Commun.*, 9, 3666, 2018.
- 590 Wong, M.-L., An, D., Caffrey, S. M., Soh, J., Dong, X., Sensen, C. W., Oldenburg, T. B. P., Larter, S. R. and Voordouw, G.: Roles of thermophiles and fungi in bitumen degradation in mostly cold oil sands outcrops, *Appl. Environ. Microbiol.*, 81, 6825, 2015.
- Wyndham, R. C. and Costerton, J. W.: In vitro microbial degradation of bituminous hydrocarbons and in situ colonization of bitumen surfaces within the Athabasca oil sands deposit, *Appl. Environ. Microbiol.*, 41, 791, 1981.
- 595 Uribe-Alvarez, C., Ayala, M., Perezgasga, L., Naranjo, L., Urbina, H. and Vazquez-Duhalt, R.: First evidence of mineralization of petroleum asphaltenes by a strain of *Neosartorya fischeri*, *Microb. Biotechnol.*, 4, 663-672, 2011.
- [Yu, Z., Wu, G., Li, F., Chen, M., Vi Tran, T., Liu, X., and Gao, S.: Glaciation enhanced chemical weathering in a cold glacial catchment, western Nyaingêntanglha Mountains, central Tibetan Plateau, \*J. Hydrol.\*, 126197, 2021.](#)
- Zeng N.: Glacial-interglacial atmospheric CO<sub>2</sub> change — The glacial burial hypothesis. *Adv. Atmospheric Sci.*, 20, 677-693,  
600 2003.
- Zeng N.: Quasi-100 ky glacial-interglacial cycles triggered by subglacial burial carbon release. *Clim. Past.*, 3, 135-153, 2007.
- Zhao N., Marchal O., Keigwin L., Amrhein D. and Gebbie G.: A synthesis of deglacial deep-sea radiocarbon records and their (in)consistency with modern ocean ventilation. *Paleoceanogr. Paleocl.*, 33, 128-151, 2018.
- ZoBell, C. E.: Action of microorganisms on hydrocarbons, *Bacteriol. Rev.*, 10, 1-49, 1946.

605 **Table 1: Fluxes of kerogen oxidation normalized to one year from onsite (soil and outcrops), catchment-wide studies (river), and laboratory-based incubation studies ordered approximately from small to large CO<sub>2</sub> release fluxes. For calculations, see online supplemental.**


| Area/ Material                | Oxidation flux [MgC/km <sup>2</sup> /yr] | Comments  | References                       |
|-------------------------------|--|---|----------------------------------|
| Andes                         | ≥0.3                                     | Kerogen oxidation in the Madeira floodplain based on radiocarbon isotope mass balancing of sedimentary organic carbon. Oxidation flux normalized to catchment area reported in Clark et al., (2017).  | Bouchez et al., (2010)           |
|                               | ~1.2                                     | Kerogen oxidation in the Madeira floodplain based on radiocarbon isotope mass balancing of sedimentary organic carbon refining the result by Bouchez et al. (2010).   | Clark et al., (2017)             |
| Mackenzie River Basin         | 0.45 <sup>+0.19</sup> <sub>-0.11</sub>   | Rhenium-based estimate integrating over catchment and riverine in-situ weathering.  | Horan et al., (2019)             |
| Global average                | 0.72 <sup>+0.17</sup> <sub>-0.41</sub>   | Calculated global average rate of kerogen oxidation on land. See Equation 1.  | See text.                        |
| Soil, glacial till (Canada)   | 4.2                                      | Holocene-averaged flux of onsite kerogen oxidation based on CO <sub>2</sub> evolution and carbon isotopes.  | Keller and Bacon (1998)          |
| Taiwan                        | 6.1-18.6                                 | Radiocarbon biogeochemistry reveals microbially-mediated weathering of kerogen onsite in soils.   | Hemingway et al., (2018)         |
|                               | 12±6                                     | Sedimentary organic carbon mass balancing with radiocarbon integrating over catchment and riverine in-situ weathering   | Hilton et al., (2011)            |
|                               | 7.4-13.0                                 | Rhenium-based estimate integrating over catchment and riverine in-situ weathering.  | Hilton et al., (2014)            |
| Shale (Posidonia, Germany)    | 11-16                                    | Onsite kerogen oxidation based on Holocene scenario described in reference using mass loss of kerogen and estimated erosion rates. Weathering depth of 5 m and organic carbon to organic matter conversion factor of 1.5 assumed.   | Littke et al., (1991)            |
| Southern Alps, New Zealand    | 9-50                                     | Rhenium-based estimate integrating over catchment and riverine in-situ weathering. Oxidation fluxes reported for 4 catchments with minimum and maximum reported here.   | Horan et al., (2017)             |
| Tar sandstone (Brazil)        | ~54                                      | Laboratory-based experiments over 1 year duration under water saturated and anoxic conditions at 20°C. Results scaled with density taken as 2000 kg/m <sup>3</sup> and 1 m weathering depth.  | Rogieri Pelissari et al., (2021) |
| Shale (Jurassic marl, France) | 61-64                                    | Field-based observations spanning over minutes to up to 101 days using zeolite traps.   | Soulet et al., (2018)            |
| Bituminous coal               | 320-530                                  | Laboratory-based subaquatic oxidation experiments at 24°C for air-saturated water over 179-442 days duration. Carbon release calculated assuming 30-50% of oxygen reacts to form CO <sub>2</sub> , an average specific surface area of coal of 1 m <sup>2</sup> /g, average density of 1400 kg/m <sup>3</sup> , and 1 m weathering depth. | Chang and Berner (1998, 1999)    |
| Shale (Brazil)                | 80-1050                                  | Laboratory-based experiments over 75 days duration under water saturated, dark, and anoxic conditions at 25°C. Results scaled with density taken as 2200 kg/m <sup>3</sup> and 1 m weathering depth.  | Bertassoli Jr. et al., (2016)    |
| Oil sands (Bitumen)           | 6800-13000                               | Laboratory-based experiments of 100 days duration for biodegradation of bitumen with different microbial cultures. Results scaled to oil sands with density taken as 2000 kg/m <sup>3</sup> with 10% bitumen content, and 1 m weathering depth.   | Ait-Langomazino et al., (1991)   |








610 Figure 1: Organic carbon cycle with the flow of kerogen (black solid lines) and the flow of biospheric carbon (green solid lines) showing **both** the fixation of atmospheric CO<sub>2</sub> by **both** terrestrial and marine primary productivity. The combined flux of reworked kerogen and biospheric carbon into ocean sediments constitutes **the** total organic carbon burial **entering into** the endogenous kerogen pool (Galy et al., 2015; Hedges and Oades, 1997).



620

Figure 2: Conceptual hypothesis overview: Changes in kerogen reburial efficiency and its effect on the reentry of ancient carbon into surficial carbon pools as a function of overall climate state. During glacial times, kerogen reburial is promoted by the activity of glaciers and ice sheets with relatively little oxidation of this carbon during its transit across Earth’s surface characteristic of a “closed” exogenous kerogen cycle. During glacial terminations and interglacials, the oxidation of kerogen is more efficient leading to the exhalation of this carbon to the atmosphere characteristic of an “open” exogenous kerogen cycle.







Figure 3: Evolution of atmospheric CO<sub>2</sub> and its carbon isotope composition starting from the Last Glacial Maximum with redrawn data from Schmitt et al. (2012) and <sup>14</sup>C data from Reimer et al. (2013). The area of the Laurentide Ice Sheet (including Greenland) is plotted after Dalton et al. (2020). The gray time envelope indicates the greatest rate of atmospheric CO<sub>2</sub> increase coinciding with a pronounced negative pulse in stable carbon isotope composition and accelerated decrease in radiocarbon concentration. In addition to the hypothesized role of kerogen oxidation, the superimposed effects of ocean-air gas exchange, a ~~growing~~ increasingly voluminous terrestrial biosphere, changes in radiocarbon production in the atmosphere, etc. all contribute to the evolution of these parameters.










Figure 4: Laurentide Ice sheet extents bracketing the time span of most pronounced glacial-interglacial CO<sub>2</sub> increase (redrawn after Dalton et al., 2020). ~~Sedimentary basins showing t~~The Western Canadian Sedimentary Basin (hatched area, with extension of related units into the United States) and the Cordilleran Basins ~~are to the west of and the~~ Canadian Shield ~~to the east and west, respectively~~ (simplified after Miall and Blakey, 2019). ~~The southern border of the Canadian shield is drawn with mainly Phanerozoic sedimentary units to the south (simplified after Reed et al., 2004).~~ The surficial kerogen content in the Canadian Shield is generally low, while high kerogen contents are present along the axes of the Cordilleran and Western Canadian Sedimentary Basins (see Fig. 2 in Copard et al., 2007). ≤300 years after glacial retreat extended into the Canadian Shield, the rise in atmospheric CO<sub>2</sub> subsided (compare with Fig. 3). Here, kerogen oxidation ~~mainly~~ in western Canada is hypothesized to have acted as a major source of CO<sub>2</sub> to the atmosphere in the direct wake of glacial retreat.