Nitrogen loss processes in response to upwelling in a Peruvian coastal setting dominated by denitrification

Kai G. Schulz¹, Eric P. Achterberg², Javier Arístegui³, Lennart T. Bach⁴, Isabel Baños³, Tim Boxhammer², Dirk Erler¹, Maricarmen Igarza⁵, Verena Kalter^{2,6}, Andrea Ludwig², Carolin Löscher⁷, Jana Meyer², Judith Meyer², Fabrizio Minutolo², Elisabeth von der Esch⁸, Bess B. Ward⁹, and Ulf Riebesell²

¹Centre for Coastal Biogeochemistry, School of Environment, Science and Engineering, Southern Cross University, Lismore, NSW, Australia

²GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

³Instituto de Oceanografía y Cambio Global, IOCAG, Universidad de Las Palmas de Gran Canaria ULPGC, Las Palmas, Spain

⁴Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

⁵Instituto del Mar del Perú (IMARPE), Dirección General de Investigaciones en Oceanografía y Cambio Climático, Callao, Peru

⁶Memorial University of Newfoundland, Department of Ocean Sciences, Logy Bay, Newfoundland, Canada

⁷Nordcee, DIAS, Department of Biology, University of Southern Denmark, Campusvej 55, Odense M-DK

⁸Institute of Hydrochemistry, Chair of Analytical Chemistry and Water Chemistry, Technical University of Munich, Munich, Germany

⁹Department of Geosciences, Princeton University, Princeton, New Jersey 08544

Correspondence: Kai G. Schulz (kai.schulz@scu.edu.au)

Abstract. Upwelling of nutrient-rich deep waters make Eastern Boundary upwelling systems (EBUS), such as the Humboldt Current System, hotspots of marine productivity. Associated settling of organic matter to depth and consecutive aerobic decomposition results in large sub-surface water volumes being oxygen-depleted. Under these circumstances organic matter remineralisation can continue via denitrification which represents a major loss pathway of bioavailable nitrogen. Another pro-

- 5 cess removing significant amounts of nitrogen in these areas is anaerobic ammonium oxidation. Here we assess the interplay of suboxic water upwelling and nitrogen cycling in a manipulative off-shore mesocosm experiment. Measured denitrification rates in the oxygen-depleted bottom layer of the mesocosms mostly ranged between 5.5–20 (interquartile range), reaching up to 80 nmol N₂ L⁻¹h⁻¹. However, realised actual in-situ rates did most likely not exceed 0.6–1.6 nmol N₂ L⁻¹h⁻¹ (interquartile range), due to substrate limitation in the mesocosms. This was in contrast to realised rates in In the surrounding Pacific
- 10 measured denitrification rates were similar, although no indications of substrate limitation were detected. Both in the mesocosms and the Pacific Ocean anammox made only a minor contribution to overall nitrogen loss when encountered, potentially related to organic matter C/N stoichiometry and/or process specific oxygen and hydrogen sulphide sensitivities. Over the first 38 days of the experiment, total nitrogen loss calculated from rates of denitrification and anammox rates was comparable to estimates from a full nitrogen budget in the mesocosms and ranged between $\sim 2-6 \,\mu$ mol N L⁻¹. This represents up to $\sim 20\%$
- 15 of the initially bioavailable inorganic and organic nitrogen standing stocks. Interestingly, this loss is comparable to the total amount of particulate organic nitrogen that was exported into the sediment traps at the bottom of the mesocosms in about 20

metres depth. Altogether, this suggests that a significant portion, if not the majority of nitrogen that could be exported to depth, is already lost, i.e. converted to N_2 in a relatively shallow layer of the surface ocean, provided oxygen-deficient conditions like during coastal upwelling in our study.

20 1 Introduction

25

30

Amongst the most productive marine ecosystems are Eastern Boundary upwelling systems (EBUS), which are mainly fuelled by wind-driven upwelling of dissolved inorganic nutrient-rich deep waters to the surface ocean, stimulating primary and associated higher trophic level productivity (Chavez and Messié, 2009; Kämpf and Chapman, 2016; FAO, 2018). This is particularly true for the Humboldt Current System off Peru (Montecino and Lange, 2009). High productivity and eventual export of organic matter to depth result in marked oxygen consumption by aerobic respiration, leading to so-called sub-surface oxygen-depleted zones as well as virtually anoxic oxygen minimum zones (ODZs and OMZs, respectively) at depth (e.g. Cline and Richards (1972); Paulmier and Ruiz-Pino (2009)). In the absence of oxygen, heterotrophic organic matter decomposition can continue with alternative electron acceptors such as nitrate (NO_3^-) or nitrite (NO_2^-) via nitric oxide (NO) and nitrous oxide (N_2O) to molecular nitrogen (N_2), in a series of separate steps carried out by a variety of bacteria (Zumft, 1997). In their entirety these processes are summarised as denitrification. A by-product of denitriciation denitrification is ammonium (NH_4^+) from the remineralised organic matter, but overall denitrification constitutes a net loss of bioavailable nitrogen (Paulmier et al., 2009). A second prevalent nitrogen loss pathway in ODZs and OMZs is autotrophic anaerobic ammonium oxidation (anammox) which

utilises NH₄⁺ and NO₂⁻ produced by heterotrophic processes to produce energy for carbon dioxide (CO₂) fixation and organic matter production (Thamdrup et al., 2006; Brandes et al., 2007). Again, the net balance in terms of bioavailable nitrogen is negative (Paulmier et al., 2009).

Observational and modeling studies have estimated the loss of bioavailable nitrogen via water column denitrification and anammox in suboxic ODZs and anoxic OMZs to amount to about 20-35% of the total nitrogen losses ocean-wide (for reviews

see Bianchi et al. (2012); Zhang et al. (2020)). In a context of climate/ocean change, which has been projected to enhance the temperature gradient between land and ocean and hence alongshore winds (Di Lorenzo, 2015), upwelling intensity and
frequency of deep waters would subsequently increase (Hauri et al., 2013; Wang et al., 2015). Such a scenario has been put forward for the Humboldt Current System off Peru but also other EBUS (Bakun and Weeks, 2008; Varela et al., 2015). Furthermore, due to increasing temperatures the ocean looses oxygen (O₂) and OMZs are expanding (e.g. Bopp et al. (2002); Bograd et al. (2008); Stramma et al. (2008); Oschlies et al. (2017)), changing biogeochemistry. Together with changes to

microbial activity, this modifies biogeochemical properties of upwelled waters . This is also the case for including, next to Q₂,
carbonate chemistry speciationas-, i.e. ongoing ocean acidification further decreases already low pH levels in these deep water pH levels (e.g. Feely et al. (2008); Franco et al. (2018); Schulz et al. (2019)). In essence, a better understanding of Changes in upwelling frequency/intensity, oxygen availability, temperature and pH can influence planktonic food web functioning in EBUS, with repercussions for nitrogen loss processes related to coastal upwelling is critical (Lam and Kuypers, 2011).

Here we make use of an off-shore mesocosm experiment to quantify the importance of nitrogen loss processes in overall nitrogen cycling following a-simulated deep-water upwelling event-in the Humboldt Current System.

2 Methods

55

The current experiment started on February 22, 2017 with the deployment of eight Kiel Off-Shore Mesocosms for Future Ocean Simulations (KOSMOS) about 4 nm-nautical miles (nm) off the Peruvian coast close to Isla San Lorenzo at 12.0555°S and 77.2348°W, and ended on April 16, 2017 after 50 days of sampling. Full details on the experimental setup, and the sampling, manipulation and maintenance schedule can be found in Bach et al. (2020).

2.1 Mesocosm setup and sampling

The KOSMOS system comprised 8 mesocosms, consisting of polyurethane bags of 2 m diameter extending to \sim 18.7 m depth and the last two metres being a funnel-shaped sediment trap. After two days of thorough flushing and soaking of the bags, which had been pulled under the surface and were open to water exchange through 3 mm mesh on both ends, the bags were se-

- 60 cured above the water line and the sediment traps attached, enclosing about 54 m^3 of a natural plankton assemblage. Sampling commenced on February 26, i.e. day 1. Sampling days typically started in the morning between 6:30–7:00 by removing the material that had accumulated in the sediment traps with a pump, followed by CTD casts. The CTD (Sea & Sun Technology 90M memory probe) was additionally equipped with an optical oxygen sensor and a calibrated AMT amperometric hydrogen sulphide (H₂S) sensor (for details on operation, the other sensors and calibrations/corrections see Schulz and Riebesell
- 65 (2013)). CTD casts were followed by sampling with an integrating water sampler (IWS, Hyrdobios). Because strong thermal stratification resulted in two distinct water masses at the surface (high oxygen and pH) and at the bottom (low oxygen and pH) of the mesocosms (Fig. 2A, B and C), two separate depth-integrated water samples were taken. According According to changes in stratification the depths were adjusted over time (days 1–2: surface 0-5 m and bottom 5-17 m, days 3–28: surface 0-10 m and bottom 10-17 m, and days 30–50: surface 0-12.5 m and bottom 12.5-17 m).
- Furthermore, to stabilise and maintain bottom water characteristics (as in the surrounding Pacific), a NaCl-brine solution was homogeneously injected below 10 m on day 13 and below 12.5 m on day 33, increasing salinity, and hence stratification, by about 0.7 and 0.5 psu, respectively (see Bach et al. (2020) for details).

While all seawater bulk parameters, such as particulate and dissolved organic matter or dissolved inorganic nutrients were measured on the two depth-integrated samples (see section 2.5 below for details), samples for N-loss process incubations (see

- restion 2.3 below for details) were taken with a Niskin sampler from 15.25 metres depth and treated as gas-sensitive, i.e. filled into two 100 ml glass-stoppered Duran bottles with at least one bottle volume of overflow, closed without headspace and kept cold and in the dark ($\sim 2-4$ hours) until processing. A direct comparison of oxygen concentrations in these samples by Winkler-titrations, following the recommendations of Carritt and Carpenter (1966); Bryan et al. (1976); Grasshoff et al. (1983), with CTD-oxygen-optode measurements revealed an offset by +13 μ mol L⁻¹ in the CTD data, probably related to the
- 80 2 second response time of the sensor and a strong gradient from surface to bottom (only down-cast data was used) although

response time hysteresis corrected ($\tau = 1$ s) as described in Fiedler et al. (2013). Hence, oxygen concentrations at depth as shown here from CTD casts, are likely to have been significantly lower.

2.2 Deep water collection and addition

To simulate upwelling with two distinct OMZ-signatures ('moderate' and 'extreme', in terms of N-deficit), deep water was

- 85 collected from 90 m at 12.044333°S, 77.377583°W, and 30 m at 12.028323°S, 77.223603°W, on days 5 and 10, respectively. Unfortunately, however, in terms of N/P ratios the waters targeted for the 'moderate' treatments had already quite a However, both waters had a quite strong N-deficit (N*), in comparison to a typical N/P of 16/1 required for phytoplankton growth (Redfield et al., 1963; Brzezinski, 1985), which was similar to the N-deficit in the deep water for the 'extreme' treatment and will be referred to as 'low N/P' and 'very low N/P' treatments in the following (compare Tab. 1). The deep waters were
- 90 collected into 100 m³ bags without headspace at the respective depths and sealed once brought back to the surface. Deep water was added by first removing about 20 m³ from each mesocosm and replacing it with the respective deep water that was injected into the bottom layer between 14-17 metres on day 11, and the surface layer between 1-9 metres on day 12. To minimise changes to deep water gas concentrations during injection, water was pumped from two metres depth out of the deep water bags. For further details on collection and addition see Bach et al. (2020).

95 2.3 N-loss processes incubations, measurements and calculations

The two main N-loss processes in oxygen-deficient waters off the Peruvian coast, i.e. denitrification and anammox (Fig. 1) were assayed with incubations of water from 15.25 m depth, using labelled ¹⁵NH₄Cl (≥ 98 atom %) or Na¹⁵NO₂ (98 atom %), i.e. by an addition of 3 µmol L⁻¹ each. During incubations with the former, anammox will be traced by the production of ²⁹N₂ (¹⁵N¹⁴N), and ³⁰N₂ (¹⁵N¹⁵N) if coupled to nitrification (although unlikely as being an oxic process – compare Fig. 1). During incubations with ¹⁵NO₂⁻, anammox would again produce ²⁹N₂, while denitrification would produce both,²⁹N₂ and ³⁰N₂ (Fig. 1). Here, a further complication for rate calculations would be coupled nitrate ammonification, aka DNRAdissimilatory nitrate or nitrite reduction to ammonium (DNRA), as also leading to ³⁰N₂ production via anammox (compare Fig. 1 and

- Holtappels et al. (2011)). Incubations were in 12 ml glass Exetainers ('Double Wadded', Labco Ltd.) in duplicates for each of the four time points, i.e. $0, \sim 2, \sim 6-7$ and ~ 20 hours after label addition (in order to avoid potential bottle effects), in
- 105 the dark and inverted at a failry fairly constant 17±0.2°C, close to respective in-situ conditions, and similar as described in Dalsgaard et al. (2003) and Ward et al. (2009). All sample handling, such as filling of the Exetainers with 8 ml of labelled sample water using an Eppendorf Multipette E3, were carried out in a glove box that had been evacuated three times with a vacuum pump followed by flushing with N₂ 4.5 gas (as well as the open Exetainers). After that To reduce large background ²⁸N₂ levels and facilitate detection of the small isotopic signal of labelled N₂ being produced during incubations, the Exetainers
- 110 were capped and sparged with Helium 5.0 at 3 psi for six minutes on a manifold that could hold all 16 Exetainers of a single mesocosm. Based on previous calculations and measurements, such setup will replace about 20 times the volume of each exetainer (unpublished data). This is lower than the 24 times, reported to ensure that the reduction in O₂ concentration is less than 20% compared to in-situ conditions (Holtappels et al., 2011). A similar reduction would also be observered for most other

gases (Wanninkhof, 1992), except those buffered by conjugate acid-base pairs, such as H₂S or CO₂, for which the reduction

115 would be even less. Incubations were stopped at each time point by the injection of 50 μ l of a ZnCl₂ solution (50% w/v), followd by thorough mixing.

Exetainer vials were stored upside down at room temperature in the dark. Prior to analysis, the headspace and water was equilibrated at room temperature by placing the Exetainers on a platform mixer at 100 rpm over night. For measurements, 100 μ l of headspace from each Exetainer was injected, using an autosampler (Carvalho and Murray, 2018), into a PLOT GC column

- 120 at 2 ml/min housed in a Trace GC oven and interfaced with a Thermo Delta V Plus mass spectrometer via a GC Combustion III unit, followed by a liquid nitrogen trap to minimise interference by CO (a constituent in air) and NO (a secondary product during ionisation of water/oxygen, followed by production of reactive oxygen species and recombination with nitrogen species) due to imperfect GC column peak separation. The mass spectrometer was calibrated for N_2 concentrations by injections of known amounts of air.
- Rate calculations were straight forward, as in most ¹⁵NH₄⁺ incubations no enrichment over time in ²⁹N₂ was detected, indicating the absence of anammox. Hence, denitrification in the ¹⁵NO₂⁻ incubations (and because ³⁰N₂ was quite noisy) was calculated from the increase in ²⁹N₂ and the known ratio of labelled to unlabelled NO₂⁻ (in the rare cases where anammox was detected in the ¹⁵NH₄⁺ incubations, denitrification was corrected for by subtraction). Rates were calculated from the linear regression slopes (Fig. A1) of the increase in the overall amount of N removed (as measured by areas ²⁸A and ²⁹A, and associated δ¹⁵N), which was determined by the following conversions (modified from Thamdrup et al. (2006); Holtappels et al.

(2011)):

$$N_{\rm removalt1-3} \left(\mu {\rm mol} \, N_2 \, {\rm L}^{-1}\right) = \left[\left({}^{15}{\rm r} \, \frac{\left({}^{28}{\rm A} + {}^{29} \, {\rm A}\right) \, {\rm cf}_{\rm MS} \, \frac{{\rm V}_{\rm h}}{{\rm V}_{\rm a}}}{{\rm f}_{\rm N_2 \, {\rm air}} \, {\rm cf}_{\rm 15} \, {\rm N}} \right)_{\rm t1-3} - \left({}^{15}{\rm r} \, \frac{\left({}^{28}{\rm A} + {}^{29} \, {\rm A}\right) \, {\rm cf}_{\rm MS} \, \frac{{\rm V}_{\rm h}}{{\rm V}_{\rm a}}}{{\rm f}_{\rm N_2 \, {\rm air}} \, {\rm cf}_{\rm 15} \, {\rm N}} \right)_{\rm t0} \right] \, \frac{1000}{{\rm V}_{\rm i}} \tag{1}$$

with

$${}^{5}r = \frac{{}^{15}R}{1 + {}^{15}R}$$
 (2)

135 and

140

1

$${}^{15}\mathrm{R} = \frac{\delta^{15}\mathrm{N}}{1000} \,{}^{15}\mathrm{R}_{\mathrm{air}} + 1 \tag{3}$$

where ¹⁵R and ¹⁵R_{air} denote the ratios of ¹⁵N/¹⁴N in a sample gas or air, respectively (the latter determined from measured ²⁸N₂ and ²⁹N₂ by Junk and Svec (1958), i.e. 0.00367647, and recommended by Coplen et al. (2002)), δ^{15} N the resulting isotopic composition (‰), V_h, V_i and V_a the volumes of headspace, sample incubated and analysed (ml), respectively, cf_{MS} the determined calibration factor for each measurement run to convert mass spectrometer peak area (²⁸A+²⁹A) into abundance (µmol), f_{N2air} the fraction of N₂ found in equilibrium in the 4 ml of headspace (V_h) in relation to the total amount, including the 8 ml of sample water (V_i), in an Exetainer, and t1-3 or t0 referring to the respective incubation sampling times. The conversion factor f_{N2air} was calculated from N₂ solubility (Hamme and Emerson, 2004) for a lab temperature of 21 °C and a salinity of 35. To extrapolate from measured removal of ¹⁵N to total N a conversion factor, cf_{15N} was calculated that takes into

145 account the availability of labelled and overall substrate(s), together with the likelihood of ${}^{29}N_2$ production, i.e. $2F_N(1-F_N)$ for denitrification incubations (binomial), and F_N for anammox, with F_N denoting the ratio of labelled to total $[NO_2^-]$ or $[NH_4^+]$, respectively.

2.4 In-situ substrate limitation of denitrification, total N-loss calculations and orni-eutrophication

In more than half of the denitrification incubations measured N-loss over a period of about 20 hours was higher than the combined concentrations of in-situ NO₃⁻ and NO₂⁻. This indicates that the 3 µmol L⁻¹ addition of Na¹⁵NO₂ alliviated alleviated substrate limitation and rates in the incubations were higher than theoretically sustainable under in-situ conditions. In these cases we report both rates, with the maximum in-situ denitrification rates estimated as the combined concentrations of in-situ NO₃⁻ and NO₂⁻ divided by the incubation period.

- Total N-loss by denitrification and anammox was calculated only for the first 38 days of the experiment. The reason was to 155 keep the estimates by rate measurements comparable to estimates from a full N-budget of the mesocoms second secon
- 160 from rate measurements by simply taking a mean of the individual rates in each mesocosm (in case also anammox was detected the measured rate was added to that of denitrification). This hourly mean N_2 loss was then multiplied by $24 \times 38 \times 2.24$ hrs per day×38 days×2 (conversion between N_2 to N) and divided by 3 (average contribution of bottom layer water to overall mesocosm volume), converting it to total N-loss over the first 38 days of the experiment. In cases measured rates exceeded those theoretically sustainable (see above for details), the latter was used.

165 2.5 Ancillary measurement parameters

Dissolved inorganic nutrient concentrations in seawater, i.e. $NO_3^- NO_2^- NH_4^+$ (together referred to as DIN), PO_4^{3-} and $Si(OH)_4$ were measured colorimetrically on a segmented flow analyser (QuAAtro, SEAL Analytical) on site (for details please see Bach et al. (2020)).

Dissolved organic nitrogen (DON) was calculated from a mass balance by subtracting measured DIN from total dissolved 170 nitrogen (TDN) concentrations. The latter was also determined by segmented flow analysis after an oxidising step with Oxisolv (Merk) for 30 minutes in an autoclave.

Particulate organic nitrogen and carbon (PON, POC) concentrations in the water column were determined by filtering known amounts of seawater over pre-combusted GF/F filters that were stored frozen until analysis on an elemental analyser (EuroVector EA 3000 - for details please see Bach et al. (2020)).

The removal of nitrogen from the water column via sedimentation (N_{sed}) was determined on material collected every other day from the sediment traps which was quantitatively precipitated, freeze-dried, weighed and an aliquot measured on an elemental analyser (EuroVector EA 3000 - for details please see again Bach et al. (2020) and Boxhammer et al. (2016)).

2.6 Statistical analyses

In order to assess what drives denitrification rates, stepwise multiple linear regressions with interaction terms (MLRs) were car-

- ried out. Prior to regressions, outliers in the measured/maximum sustained maximum-sustained denitrification (see section 2.4 and Tab. 2 for details) were identified in a box-whisker plot and removed, i.e. six values with rates higher than 4 nmol $L^{-1}h^{-1}$. In order to avoid over-fitting and find a balance between model complexity and explanatory power, we followed a backward elimination process, starting with the full seven potential measured predictors, i.e. PON, PON_{sed}, DON, NO₃⁻, NO₂⁻, O₂ and H₂S, all measured in the bottom layer of the mesocosms (with the exception of sedimenting PON_{sed}). Note that we have opted
- to not include POC, as being highly co-correlated with PON, as well as DOC which was not measured in the bottom layer (but probably would have a similar issue). In a next step, MLRs with all possible combinations of 6 potential predictors and their resulting R^2 were calculated, which was followed by futher-further MLRs, subsequently reducing the number of predictors each time by 1 (a total of 119 MLRs were fitted). Calculations were performed using the functions 'boxplot', 'stepwisefit' and 'plotEffects' in MATLAB.

190 3 Results

195

The experiment took place during the 2017 coastal El Niño, which was characterised by three significant surface ocean warming events throughout January to April (Garreaud, 2018) with the last two, end of February and mid March, clearly evident by water surface temperatures above 22°C at our mooring site (Fig. 2A). The El Niño was accompanied by torrential rains further inland which was reflected by periods of significant reductions in surface ocean salinity (Fig. 2B), coinciding with water discharge more than twice the typical rates of the nearby river Rimac (Fig. A3). During our experiments there were, however, also periods of deep water upwelling, as evidenced by colder surface ocean temperatures, as well as reduced oxygen saturation states and pH levels, reaching down to \sim 30% and 7.5 (total scale), respectively (Fig. 2C, D).

3.1 Temporal changes in oxygen, inorganic and organic nitrogen as well as hydrogen sulphide in the bottom layer of the mesocosms

- 200 Thermal stratification of Pacific Ocean and mesocosm waters during the initial phase of the experiment meant strict separation of well oxygenated surface (~0-10 metres) from oxygen-depleted bottom waters (Fig. 2C). And while in the surrounding Pacific bottom waters remained oxygen depleted throughout, corresponding oxygen levels in the mesocosms started to increase (Fig. 2 and 3E). This was caused by cooling of the surface waters due to upwelling in the surrounding Pacific and resulting mixing of surface and bottom waters in the mesocosms. Such artificial behaviour was mitigated by increasing bottom layer
- 205 salinity, and therefore stratification, on days 13 and 33, which brought oxygen concentrations at depth down to typical Pacific levels again (Fig. 3E).

All mesocosms started with significant amounts of dissolved inorganic nitrogen present as NO_3^- and NO_2^- , which however were quickly depleted within the first two weeks (Fig. 3C, D). The deep water addition significantly increased NO_2^- concentrations in the 'moderate low N/P' treatments, although it made only a minor contribution to the overall nitrogen budget (see

210 following sections for details).

Initial particulate and dissolved organic nitrogen concentrations (PON, DON) at depth were similar in all mesocosms (between 6 – 10 μ mol L⁻¹), and while there was no clear temporal trend for PON, DON saw a steady decline by about 30% until day 40 (Fig. 3A, B).

H₂S concentrations at depth were in the micro-molar range in all mesocosms as well as the surrounding Pacific and mostly
 oscillated between 3–10 μmol L⁻¹ (Fig. 3F), equivalent to 0.1–0.3 ppm. In contrast concentrations in surface waters were mostly in the high to low nano-molar range (data not shown).

3.2 Rates of denitrification and anammox, and overall nitrogen loss

Measured denitrification rates in the 24 hour incubations were similar in all mesocosms, ranging between less than 1 to up to ~80 nmol L⁻¹h⁻¹ (Tab. 2). More importantly however, towards the end, measured rates exceeded those theoretically sustainable by substrate availability, i.e. combined in-situ NO₃⁻ and NO₂⁻ concentrations. In comparison, measured denitrification rates in samples from the surrounding Pacific were comparable, although more variable between consecutive measurement days.

Anammox was only detected on day 12 in mesocosms 7 and 8, at rates significantly smaller than those typical for denitrification (Tab. 2). In contrast, anammox was occasionally detected in the surrounding Pacific Ocean, i.e. on days 8, 12 and 46.

When comparing potential nitrogen loss, calculated as the sum of average denitrification and anammox, over the first 38 days prior to the onset of orni-eutrophication (see section 2.4 and Tab. 2 for details), with estimates of total nitrogen losses in each mesocosm from a nitrogen budget approach (compare Tab. 2 and Fig. 4 and see next section for details), they were very similar, with a mean of $3.86 \,\mu\text{mol L}^{-1}$ for the former and $3.64 \,\mu\text{mol L}^{-1}$ for the latter. It is acknowledged, however, that there was no statistically significant correlation between the two approaches.

230

3.3 Nitrogen budget

Summing up all organic and inorganic nitrogen species (excluding dissolved gases) in the water column, i.e. PON, DON, NO₃⁻, NO₂⁻, NH₄⁺, N_{sed} (nitrogen exported through the sediment traps), and N_{DW} (overall deep water addition changes to the various nitrogen pools), revealed a first phase in all mesocosms for which the budget did not seem to be closed, as evidenced
by a constant decline in total N during the first two weeks (Fig. 4). This was followed by a steady increase in total N until day 40, which was still below starting levels in all mesocosms. This net loss reflects, as we will argue below, the combined nitrogen loss processes such as denitrification and anammox, exceeding nitrogen fixation, which was less than 0.1 μmol L⁻¹ over the entire water column in this period (Kittu pers. comm.). Finally the last ten days were characterised by a rapid increase in total N, mostly driven by rising surface PON concentrations, following continuous biomass production fuelled by external

240 orni-eutrophication, i.e. the introduction of nitrogen-rich bird faeces to the mesocosms (see Bach et al. (2020) for details).

3.4 Stepwise multiple linear regressions

In terms of \mathbb{R}^2 there was a turning point in the number of potential main variables to explain measured/maximum sustainable maximum-sustainable denitrification rates in stepwise multiple linear regressions (MLRs) with interactions. Above 4 indipendent independent variables, \mathbb{R}^2 only increased slightly, while below it decreased relatively quickly (Fig. 5A). Both models with the

245 highest R^2 for 4 main variables identified NO_2^- and PON as those with the highest positive main effect size, with the remaining two variables having relatively low effect sizes (Fig. 5B, C, Tab. 3). Interestingly, also the MLR with the highest R^2 for 3 main variables identified NO_2^- and PON as those with the largest positive main effect sizes, followed by NO_3^- (data not shown).

The resulting linear regression of measured/maximum sustainable maximum-sustainable (compare Tab. 2) versus predicted denitrification rates slightly deviated from an ideal 1:1 relationship, meaning that lower denitrification rates would be overwhile higher ones under-estimated (Fig. 5D).

4 Discussion

A discussion on the unusual situation of a coastal El Niño during our study period, i.e. significant warming of surface waters, increasing stratification and reducing upwelling intensity/frequency can be found elsewhere (Bach et al., 2020). Most importantly however, there were also multiple upwelling events in the surrounding Pacific during this time which were of similar

magnitude to our experimental upwelling (Chen et al., 2021), providing the natural context.

255 t

4.1 Denitrification rates and nitrogen budgets

Denitrification rates measured in the mesocosms and the surrounding Pacific waters of up to ~ 80 nmol N₂ L⁻¹ h⁻¹ (although mostly well below at a median of 12.4), were within the range of reported rates in Peruvian ODZ and OMZ waters (Farías et al., 2009; Dalsgaard et al., 2012). This was despite significantly higher oxygen concentrations here than typical for suboxic ODZs, i.e. less than ~6 µmol L⁻¹ (Tyson and Pearson, 1991; Yang et al., 2017). Although, denitrification can be relatively insensitive to oxygen concentrations of up to at least 30-40 µmol L⁻¹ (Farías et al., 2009), which was close to the is higher than levels in our study for most of the time (see Fig. 3 but also section 4.2.2).

Most interestingly, measured denitrification rates in incubations during the second half of the experiment exceeded those theoretically sustainable under conditions of in-situ substrate availability in the mesocosms (Tab. 2). The Nitrification, supplying additional nitrite for denitrification, can operate at low micromolar (and even nano-molar) levels such as those found in our study (Bristow et al., 2016). However, a significant contribution to measured denitrification rates is unlikely as there and elsewhere (Peng et al., 2016; Santoro et al., 2021) nitrification rates were usually at least an order of magnitude lower than our measured denitrification rates. Hence, the reason for higher measured rates is most likely found in the fact that both NO₃⁻

and NO_2^- concentrations had dropped below 0.1 or even 0.01 $\mu mol L^{-1}$ during this stage, indicating that the denitrifying community was actually substrate limited. Such observation was also made by Michiels et al. (2019), although in a temperate

Fjord system. In comparison, denitrification rates in the surrounding Pacific waters had a higher day to day variability and substrate limitation only occurred once. This is to be expected in a much more variable system, characterised by frequent upwelling events of oxygen depleted waters (Fig. 2) with relatively higher NO_3^- and NO_2^- concentrations (Fig. 3C, D). The

- 275 primary drivers of measured/maximum sustainable maximum-sustainable denitrification rates appeared to be the availability of NO_2^- , followed by particulate organic matter (nitrogen), as suggested indicated by multiple linear regression and effect size analyses analysis (Fig. 5). This finding is reasonable suggests that heterotrophic rather than chemolithotrophic denitrification was dominant (compare section 4.2.2), as both are substrates that might for the former process and eventually limit rates.
- Overall nitrogen loss, calculated from average denitrification rates over the first 38 days of the experiment prior to the onset of orni-eutrophication, was comparable to an alternative estimate that was based on a full nitrogen budget in each mesocosm 280 (Fig. 4, Tab. 2). The mean of both methods was indistinguishably close to each other, although there was no statistically significant correlation when all mesocosms were compared. However, given that the nitrogen budget calculations involved a mass balance of seven entities with individual measurement uncertainties, this probably does not come as a surprise. There were also no statistically significant differences between the two deep water addition treatments, which was most likely due
- 285 to relatively small differences in dissolved inorganic nitrogen in both waters in relation to the overall nitrogen pool in the mesocosms, as well as a similar N-deficit in both deep water batches (Tab. 1). It is noted, however, that overall nitrogen loss calculated by the two methods represent a lower boundary estimate. The reason are uncertainties associated with the onset of orni-eutrophication (which could have already have introduced small nitrogen perturbations prior to day 38) as well as the estimate of maximum sustainable maximum-sustainable denitrification rates when in-situ substrate limitation was encountered
- $(NO_3^-$ and NO_2^- standing stocks most likely underestimate availability as of hidden turn-over). 290

Finally, modeling exercises suggest that nitrogen loss in the water column is linked with mature El Niño/La Niña periods, with up to 70% reduced rates during the former, most likely linked to increased water column oxygen concentrations at reduced upwelling/enhanced stratification (Deutsch et al., 2011; Yang et al., 2017), as well as increasing mesoscale turbulence and associated offshore nutrient export by eddies (Espinoza-Morriberón et al., 2017). This suggests that the rates observed here could actually be significantly higher in the more frequent La Niña periods.

295

4.2 Rates of anammox and the lack thereof in the mesocosms

Anammox and denitrification were first reported in the Eastern Tropical South Pacific (ETSP) in 2006 (Thamdrup et al., 2006), and the ETSP remains the most frequently sampled and most thoroughly characterised open ocean OMZ region. Anammox was the dominant N loss process in several studies, where denitrification was either not (Hamersley et al., 2007) or only sporadically

300

(Kalvelage et al., 2013) detected. Subsequent studies in the ETSP detected both processes, and anammox was a substantial and mostly dominant part of the total N loss: average 78% anammox along a Chilean coastal transect (Dalsgaard et al., 2012), 82–90% at two high resolution stations (De Brabandere et al., 2013), and $\frac{48.849 \pm 20.320\%}{20.320\%}$ over four high resolution stations (Babbin et al., 2020).

Anammox requires a source of NH_4^+ and NO_2^- , which must either be produced in situ by remineralisation of organic matter in the absence of oxygen (as for instance by denitrification) or must be transported into the system from elsewhere, e.g., from 305

adjacent sediments (Ward, 2013). When both processes are supported by organic matter remineralisation, there is theoretical (Paulmier et al., 2009; Koeve and Kähler, 2010) and experimental evidence (Babbin et al., 2014) that the ratio of denitrification to anammox should be connected to the elemental composition of organic matter being decomposed, at least on larger spatial and temporal scales. The reasoning behind this connection is that complete anaerobic organic matter decomposition by denitri-

- fication produces NH⁺₄ and N₂ in quantities mostly dependent on the carbon to nitrogen ratio (C/N) of the organic matter being decomposed. Hence, this ratio dictates the denitrification (only utilising NO₃⁻ and or NO₂⁻) to anammox (utilising both NO₂⁻ and NH⁺₄) ratio in steady state (Babbin et al., 2014). However, dissimilatory nitrate or nitrite reduction to ammonium (DNRA)
 DNRA could also supply NH⁺₄ for anammox. Although there have been reports of DNRA in the ETSP (Lam et al., 2009), it is usually negligible in the water column (De Brabandere et al., 2013; Kalvelage et al., 2013) and rather restricted to shallow
- 315 coastal systems dominated by sediments (Jensen et al., 2011). In the absence of DNRA, complete anaerobic decomposition of average phytoplankton-derived organic matter, i.e. $C_{106}H_{175}O_{42}N_{16}P$ (Anderson, 1995), would require a ~ 28% contribution of anammox to overall N-loss via these two processes (Babbin et al., 2014).

The reason for an anammox dominance in several studies mentioned above, despite above outlined stoichiometric constraints could be partial denitrification of NO_3^- to NO_2^- without the following steps leading to N_2 production, supplying both $NO_2^$ and NH_4^+ for anammox (Lam et al., 2009; Jensen et al., 2011; Kalvelage et al., 2013; Peters et al., 2016). Furthermore, the ratio of NO_2^- to NH_4^+ produced by organic matter decomposition of a certain C/N ratio could also be influenced by anaerobic chemolithotrophic nitrite oxidation (compare Fig. 1), looping NO_2^- back to NO_3^- for further partial denitrification and associated NH_4^+ production (Babbin et al., 2020).

In summary, anammox in the absence of denitrification is difficult to explain (although partial denitrification is not easily 325 detected), while denitrification in the absences of anammox is rarely observed in the ETSP but presents no stoichiometric conflicts.

4.2.1 Organic matter C/N in mesocosms

For the first 38 days of the experiment, particulate organic matter in the bottom layer of the mesocosms, where the N-loss process incubation samples were taken from, had a C/N of \sim 7-8 (compare Bach et al. (2020)), which would not change the theoretical anammox contribution of $\sim 28\%$ by much. However, this ratio could be misleading as the particulate organic matter will be comprised by more and less labile fractions. A good proxy for that should be to just consider the freshly produced organic matter, i.e. in our case the particulate and dissolved matter in the sun-lit surface layer of the mesocosms. Interestingly, just looking at C/N of standing stocks for particulate organic matter, it was initially lower than in the bottom layer, indicative of preferential nitrogen remineralisation at depth, but increased to values of \sim 10 in all mesocosms but one (no bloom of

335 the dinoflagellate Akashiwo sanguinea - see Bach et al. (2020) for details) after the deep water addition (Fig. A4). Similarly, C/N in dissolved organic matter started already at about 10, increasing to up to 30 after the deep water addition, and when summed up to total organic matter C/N levels of up to 20 were reached (Fig. A4). Increasing C/N from 6.625 to 20 would reduce the theoretical anammox contribution to only about 10%. Furthermore, as mentioned already above, it should not be the standing stocks that are considered but the actual fresh production. Unfortunately, neither C nor N production were directly

- 340 measured, but gauging the change in total organic carbon (TOC) concentrations after deep water addition suggests that at least $\sim 100 \ \mu \text{mol L}^{-1}$ were produced (Fig. A4). In contrast, total organic nitrogen (TON) concentrations hardly changed at all, suggesting even higher C/N in freshly produced organic matter than in measured standing stocks. This could be the result of general dissolved inorganic nitrogen limitation and hence carbon overconsumption (Toggweiler, 1993), but also changes in phytoplankton community composition (see Bach et al. (2020) for details). It is noted though that changes in organic matter
- 345 standing stocks over time are not necessarily a good indicator of the quantity and quality of fresh production due to unknown and potentially different turnover times for each element, although probably better than using standing stocks themselves. In this sense the ODZ/OMZ signature of upwelled water, i.e. its nitrogen deficit, should influence the ratio of denitrification to anammox that is subsequently measured. Hence, regions with one or the other process dominating could be indicative of the upwelling source water history.

350 4.2.2 Oxygen and hydrogen sulphide in mesocosms

Varying ratios of anammox to denitrification could also be the result of different sensitivities of both processes to prevailing oxygen concentrations. However, while there are a few experiments that have directly addressed oxygen sensitivity of various N-cycling processes there appears to be no straight-forward answer. For instance, in manipulative experiments Jensen et al. (2008) found anammox at oxygen concentrations of up to $\sim 12.5 \,\mu \text{mol L}^{-1}$, with denitrification not encountered at all. This is

- similar to Kalvelage et al. (2011) who observed anammox to cease above $\sim 20 \ \mu \text{mol} \ \text{L}^{-1}$. And although no full denitrification was observed, the first step from NO_3^- to NO_2^- was found to occur even at the highest oxygen concentration of $\sim 25 \ \mu \text{mol} \ \text{L}^{-1}$. This is in turn consistent with nitrate reduction observed to at least N₂O at oxygen concentrations of up to $\sim 30\text{-}40 \ \mu \text{mol} \ \text{L}^{-1}$ (Frey et al., 2020). In contrast, both anammox and denitrification can already be significantly/completely inhibited by oxygen additions of 3 or 8 $\mu \text{mol} \ \text{L}^{-1}$ (Babbin et al., 2014) or even lower (Dalsgaard et al., 2014). In essence, there is no clear indication
- of one process being more sensitive to oxygen than the other, potentially also related to variability of oxygen concentrations on small scales, i.e. in mircoenvironments around particles which are not captured by bulk seawater oxygen concentration measurements. And indeed there is evidence that microbial diversity in the OMZ is critically linked to particles, as are rates of denitrification and anammox (Ganesh et al., 2014, 2015).
- The situation appears to be similar for H₂S, which furthermore has been observed to reach 10 μmol L⁻¹ (like in our study)
 and even higher in the coastal sub-surface ETSP close to our study location (Callbeck et al., 2018, 2019). H₂S has been reported to completely inhibit anammox at concentrations of 3 μmol L⁻¹also found. (Jensen et al., 2008), a concentration similar to that in the bottom layer of our mesocosms(Jensen et al., 2008), while Dalsgaard et al. (2014) found no effect on N₂ production by anammox, nor denitrification, at slightly lower levels of 1 μmol L⁻¹. Concerning denitrification, Dalsgaard et al. (2013) reported no effects of H₂S at even higher concentrations of up to 10 μmol L⁻¹, which is in the range of maximum concentration some instances, although most likely related to chemolithotrophic as opposed to heterotrophic denitrification (see Dalsgaard et al. (2013); Bonaglia et al. (2016) and references therein). Chemolithotrophic denitrification coupled to H₂S oxidation has also been hypothesised for coastal high H₂S stations off Peru by Kalvelage et al. (2013). Interestingly, by day 16 and onwards the anammox functional marker gene *hzo* (Schmid et al., 2008) was not anymore

detectable in any of the mesocosms (data not shown), which could be linked to relatively high H₂S concentrations, explaining

375 the lack of anammox, at least for most parts of the experiment.

4.3 ODZ/OMZ nitrogen budget implications

380

The loss of on average almost 4 μ mol L⁻¹ of nitrogen in our mesocosms fits well to zonal estimates by DeVries et al. (2012) for the ODZ in the Eastern South Pacific between 75–100 metres depth where oxygen levels were similar to those encountered here (Chang et al., 2010), although the latter study reported a 2-3 times higher nitrogen deficit. Furthermore, the overall amount of nitrogen loss measured in the Pacific was at the upper end of ranges encountered in the mesocosms, potentially connected to substrate limitation in the mesocosms during the second half of the experiment (Tab. 2).

Being closed systems, the mesocosms offer the unique opportunity to put the various nitrogen pathways, i.e. loss as opposed to initial bioavailable standing stocks and export, into a perspective context. Between $\sim 2-6 \ \mu mol L^{-1}$ of nitrogen were lost as N₂, representating up to 20% of the initially bioavailable inorganic and organic nitrogen until day 38 of the experiment 385 (compare Tab. 2 and Fig. 4). Interestingly, the amount of particulate organic nitrogen being exported below 20 metres, i.e. the approximate depth of the sediment traps at the bottom of the mesocosms (compare Fig. 1 in Bach et al. (2020)), was in the same range. This indicates that in the Peruvian EBUS about half of the nitrogen that could be exported to depth would already be lost, i.e. converted to N_2 , in a relatively shallow layer of the surface ocean, provided oxygen deficient conditions during coastal upwelling as in our study (Fig. 2). Furthermore, over the entire pelagic water column, nitrogen loss of exported 390 organic matter is likely to be even higher, suggesting that the majority of the dissolved inorganic nitrogen assimilated during

new production (equalling export production on larger scales) should actually be lost in EBUS.

Similar conclusions can also be reached by alternative means, i.e by starting with global export production. Recent estimates based on observations and models range between $\sim 5-15 \text{ Pg C v}^{-1}$, including next to the gravitation driven biological pump also those by particle injection (see Boyd et al. (2019) and refs. therein). Assuming a Redfieldian molar C/N of 6.625 (Redfield et al., 1963) would translate to $0.75-2.26 \text{ Pg N y}^{-1}$ being exported. Furthermore, considering that about 5% of 395 global primary production, and hence potential export production, is located in the surface ocean of the four major EBUS (Carr, 2002) above the ODZs and OMZs (ignoring the contributions by the Arabian Sea , Arabian Sea (Vijayaraghavan and Krishna Kumari, 1989) Bay of Bengal $\frac{1}{2}$ (Poornima et al., 2020) above the ODZs and OMZs, about 38–113 Tg N y⁻¹ could potentially be lost. In com-

parison, estimates of water column denitrification/anammox, i.e. 20-35% of total marine losses of about 260 ± 100 Tg N y⁻¹

400 (see Zhang et al. (2020) and references therein), then range between 52 ± 20 and 91 ± 35 Tg N y⁻¹, indicating that a significant portion, if not the majority, of the exported nitrogen is indeed lost in ODZs and OMZs. Finally

Nitrogen cycling in ODZs and OMZs currently plays a very important role in the overall marine nitrogen budget. However, the magnitude and direction of change in the actual nitrogen loss term in response to ongoing climate and ocean change (e.g. ocean stratification, acidification and/or changes in temperature and oxygen levels) is uncertain. This issue is further

complicated by uncertainties in future primary productivity and organic matter export estimates. For instance, depending on 405 the representative concentration pathway, future export production is projected to decrease by up to 16% at the end of this century, relative to the year 2000 could decrease as a result of changes to community structure (see Bindoff et al. (2019) for details and refs. therein), which. In summary, future changes in upwelling intensity and frequency, as well as the other potential biotic and abiotic factors mentioned above, could change the nitrogen (im)balance in ODZs and OMZs, having a significant impact on the overall marine nitrogen budget.

Data availability. The majority of data is available at https://doi.org/10.1594/PANGAEA.923395 (Bach et al., 2020), with some additional being presented here in tables.

Author contributions. KGS, LB, TB, AL, and UR designed the experiment. KGS, JA, LB, TB, MI, VK, AL, JaM, JuM, FM, EvdE and UR contributed to sampling and various measurements. DE helped with ¹⁵N data interpretation and BBW introduced KGS to denitrification and anammox assay incubations. All authors analysed the data and KGS wrote the manuscript with co-authors commenting.

Competing interests. The authors declare that they have no conflict of interests.

410

415

Acknowledgements. This project was supported by the Collaborative Research Centre SFB 754 Climate- Biogeochemistry Interactions in the Tropical Ocean financed by the German Research Foundation (DFG). Additional funding was provided by the EU project AQUACOSM and the Leibniz Award 2012 granted to UR. We thank all participants of the KOSMOS-Peru 2017 study for assisting in mesocosm sampling
and maintenance. We are particularly thankful to the staff of IMARPE for their support during the planning, preparation and execution of this study and to the captains and crews of BAP MORALES, IMARPE VI and BIC HUMBOLDT for support during deployment and recovery of the mesocosms and various operations during the course of this investigation. Special thanks go to the Marina de Guerra del Peru, in particular the submarine section of the Navy of Callao, and to the Dirección General de Capitanías y Guardacostas. We also acknowledge strong support for sampling and mesocosm maintenance by Jean-Pierre Bednar, Gabriela Chavez, Susanne Feiersinger, Peter Fritsche, Paul

- 425 Stange, Anna Schukat, Michael Krudewig. We want to thank Club Náutico Del Centro Naval for excellent hosting of our temporary filtration laboratory, office space and their great support and improvisation skills after two of our boats were lost. This work is a contribution in the framework of the Cooperation agreement between the IMARPE and GEOMAR through the German Ministry for Education and Research (BMBF) project ASLAEL 12-016 and the national project Integrated Study of the Upwelling System off Peru developed by the Direction of Oceanography and Climate Change of IMARPE, PPR 137 CONCYTEC. Furthermore, the SERVICIO NACIONAL DE METEOROLOGÍA
- 430 E HIDROLOGIA DEL PERU SENAMHI is thanked for providing the Rimac discharge data. Finally, Matheus Carvalho de Carvalho is thanked for N_2 isotope measurements.

Financial support. This research has been supported by the Collaborative Research Center SFB 754 Climate-Biogeochemistry Interactions in the Tropical Ocean financed by the German Research Foundation (DFG), the Leibniz Award 2012 (granted to Ulf Riebesell), the Helmholtz

International Fellow Award 2015 (granted to Javier Arístegui) and EU funding by AQUACOSM granted to a number of participants, including

435 Javier Arístegui and Kai G. Schulz.

References

Anderson, L. A.: On the hydrogen and oxygen content of marine phytoplankton, Deep Sea Res. Part I, 42, 1675–1680, 1995.

- Babbin, A. R., Keil, R. G., Devol, A. H., and Ward, B. B.: Organic Matter Stoichiometry, Flux, and Oxygen Control Nitrogen Loss in the Ocean, Science, 344, 406–408, 2014.
- 440 Babbin, A. R., Buchwald, C., Morel, F. M. M., Wankel, S. D., and Ward, B. B.: Nitrite oxidation exceeds reduction and fixed nitrogen loss in anoxic Pacific waters, Marine Chemistry, 224, 2020.
 - Bach, L. T., Paul, A. J., Boxhammer, T., von der Esch, E., Graco, M., Schulz, K. G., Achterberg, E., Aguayo, P., Aristegui, J., Ayon, P., Banos, I., Bernales, A., Boegeholz, A. S., Chavez, F., Chen, S.-M., Doering, K., Filella, A., Fischer, M., Grasse, P., Haunost, M., Hennke, J., Hernandez-Hernandez, N., Hopwood, M., Igarza, M., Kalter, V., Kittu, L., Kohnert, P., Ledesma, J., Lieberum, C., Lischka, S., Loescher,
- C., Ludwig, A., Mendoza, U., Meyer, J., Meyer, J., Minutolo, F., Ortiz Cortes, J., Piiparinen, J., Sforna, C., Spilling, K., Sanchez, S., Spisla,
 C., Sswat, M., Zavala Moreira, M., and Riebesell, U.: Factors controlling plankton community production, export flux, and particulate matter stoichiometry in the coastal upwelling system off Peru, Biogeosciences, 17, 4831–4853, 2020.
 - Bakun, A. and Weeks, S. J.: The marine ecosystem off Peru: What are the secrets of its fishery productivity and what might its future hold?, Progr. Oceanogr., 79, 290–299, 2008.
- 450 Bianchi, D., Dunne, J. P., Sarmiento, J. L., and Galbraith, E. D.: Data-based estimates of suboxia, denitrification, and N₂O production in the ocean and their sensitivities to dissolved O₂, Glob. Biogeochem. Cycl., 26, GB2009, 2012.
 - Bindoff, N. L., Cheung, W. W. L., Arístegui, J. G. K. J., Guinder, V. A., Hallberg, R., Hilmi, N., and ... Williamson, P.: Chapter 5: Changing ocean, marine ecosystems, and dependent communities. *IPCC special report oceans and cryospheres in changing climate*, Cambridge: University Press., 2019.
- 455 Bograd, S. J., Castro, C. G., Di Lorenzo, E., Palacios, D. M., Bailey, H., Gilly, W., and Chavez, F. P.: Oxygen declines and the shoaling of the hypoxic boundary in the California Current, Geophyss Ress Lett., 35, https://doi.org/10.1029/2008GL034185, 2008.
 - Bonaglia, S., Klawonn, I., De Brabandere, L., Deutsch, B., Thamdrup, B., and Brüchert, V.: Denitrification and DNRA at the Baltic Sea oxic–anoxic interface: Substrate spectrum and kinetics, Limnol. Oceanogr., 61, 1900–1915, 2016.

Bopp, L., Le Quéré, C., Heimann, M., Manning, A. C., and Monfray, P.: Climate-induced oceanic oxygen fluxes: Implications for the contemporary carbon budget, Glob. Biogeochem. Cycl., 16, 6–1–6–13, 2002.

- Bourbonnais, A., Letscher, R. T., Bange, H. W., Échevin, V., Larkum, J., Mohn, J., Yoshida, N., and Altabet, M. A.: N₂O production and consumption from stable isotopic and concentration data in the Peruvian coastal upwelling system, Biogeochm. Cycles, p. 678–698, 2017.
 - Boxhammer, T., Bach, L. T., Czerny, J., and Riebesell, U.: Technical Note: Sampling and processing of mesocosm sediment trap material for quantitative biogeochemical analysis, Biogeosci., 13, 2849–2858, 2016.
- 465 Boyd, P. W., Claustre, H., Levy, M., Siegel, D. A., and Weber, T.: Multi-faceted particle pumps drive carbon sequestration in the ocean, Nature, 568, 327–335, 2019.
 - Brandes, J. A., Devol, A. H., and Deutsch, C.: New Developments in the Marine Nitrogen Cycle, Chem. Rev., 107, 577-589, 2007.
 - Bristow, L. A., Dalsgaard, T., Tiano, L., Mills, D. B., Bertagnolli, A. D., Wright, J. J., Hallam, S. J., Ulloa, O., Canfield, D. E., Revsbech,N. P., and Thamdrup, B.: Ammonium and nitrite oxidation at nanomolar oxygen concentrations in oxygen minimum zone waters, Proc.
- 470 Natl. Acad. Sci. U.S.A., 113, 10 601–10 606, 2016.
 - Bryan, J. R., Riley, J. P., and Williams, P. J. L.: A Winkler procedure for making precise measurements of oxygen con-centration for productivity and related studies, J. Exp. Mar. Biol. Ecol., 21, 191–197, 1976.

Brzezinski, M. A.: The Si:C:N Ratio of Marine Diatoms: Interspecific Variability and the Effect of some Environmental Variables, J. Phycol., 21, 347–357, 1985.

- 475 Callbeck, C., Lavik, G., Ferdelman, T. G., Fuchs, B., Gruber-Vodicka, H. R., Hach, P. F., Littmann, S., Schoffelen, N. J., Kalvelage, T., Thomsen, S., Schunck, H., Löscher, C. R., Schmitz, R. A., and Kuypers, M. M. M.: Oxygen minimum zone cryptic sulfur cycling sustained by offshore transport of key sulfur oxidizing bacteria, Nature Comm., 9, 9:1729, 2018.
 - Callbeck, C., Pelzer, C., Lavik, G., Ferdelman, T. G., Graf, J. S., Vekeman, B., Schunck, H., Littmann, S., Fuchs, B. M., Hach, P. F., Kalvelage, T., Schmitz, R. A., and Kuypers, M. M. M.: *Arcobacter peruensis* sp. nov., a chemolithoheterotroph isolated from sulfide-and organic-rich coastal waters off Peru, Appl. Environ. Microbiol., 85, e01 344–19, 2019.
 - Carr, M. E.: Estimation of potential productivity in Eastern Boundary Currents using remote sensing., Deep-Sea Res., 49, 59–80, 2002.

Carritt, D. E. and Carpenter, J. H.: Comparison and Evaluation of Currently Employed Modifications of the Winkler Method for Determining Dissolved Oxygen in Seawater; A NASCO Report, J. Mar Res., 24, 286–318, 1966.

- Carvalho, M. C. and Murray, R. H.: Osmar, the open-source microsyringe autosampler, HardwareX, 3, 10–38, 2018.
- 485 Chang, B. X., Devol, A. H., and Emerson, S. R.: Denitrification and the nitrogen gas excess in the eastern tropical South Pacific oxygen deficient zone, Deep Sea Res., 57, 1092–1101, 2010.
 - Chavez, F. P. and Messié, M.: A comparison of Eastern Boundary Upwelling Ecosystems, Progr. Oceanogr., 83, 80–96, 2009.
 - Chen, S.-M., Riebesell, U., Schulz, K. G., von der Esch, E., and Bach, L. T.: Temporal dynamics of sea surface carbonate chemistry in response to natural and simulated upwelling events in the Peruvian oxygen minimum zone, Biogeosci. Discuss, 2021.
- 490 Cline, J. D. and Richards, F. A.: OXYGEN DEFICIENT CONDITIONS AND NITRATE REDUCTION IN THE EASTERN TROPICAL NORTH PACIFIC OCEAN, Limnol. Oceanogr., 17, 885–900, 1972.
 - Codispoti, L. A., Yoshinari, T., and Devol, A. H.: Respiration in Aquatic Ecosystems (Paul del Giorgio and Peter Williams (eds.)), chap. Suboxic respiration in the oceanic water column, Oxford University Press, 2005.
 - Coplen, T. B., Hopple, J. A., Bölke, J. K., Peiser, H. S., Riedler, S. E., Krouse, H. R., Rosman, K. J. R., Ding, T., Vocke Jr., R. D., Révész,
- 495 K. M., Lamberty, A., Tayler, P., and De Bièvre, P.: Compilation of Minimum and Maximum Isotope Ratios of Selected Elements in Naturally Occurring Terrestial Materials and Reagents, U.S. Geological Survey, Water-Resources Investigations Report 01-4222, 2002.
 - Dalsgaard, T., Canfield, D. E., Petersen, J., Thamdrup, B., and Acuña Gonsález, J.: N₂ production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica, Nature, 422, 606–608, 2003.
 - Dalsgaard, T., Thamdrup, B., Farías, L., and Revsbech, N. P.: Anammox and denitrification in the oxygen minimum zone of the eastern South
- 500 Pacific, Limnol. Oceanogr., 57, 1331–1346, 2012.

480

- Dalsgaard, T., De Brabandere, L., and Hall, P. O. J.: Denitrification in the water column of the central Baltic Sea, Geochim. Cosmochim. Acta, 106, 247–260, 2013.
 - Dalsgaard, T., Stewart, F. J., Thamdrup, B., De Brabandere, L., Revsbech, N. P., Ulloa, O., and Canfield, D. E. DeLong, E. F.: Oxygen at Nanomolar Levels Reversibly Suppresses Process Rates and Gene Expression in Anammox and Denitrification in the Oxygen Minimum
- 505 Zone off Northern Chile, mBio, 5, e01 966–14, 2014.
 - De Brabandere, L., Canfield, D. E., Dalsgaard, T., Friederich, G. E., Revsbech, N. P., Ulloa, O., and Thamdrup, B.: Vertical partitioning of nitrogen-loss processes across the oxic-anoxic interface of an oceanic oxygen minimum zone, Environm. Microbiol., 16, 3041–3054, 2013.
 - Deutsch, C., Brix, H., Ito, T., Frenzel, H., and Thompson, L. A.: Climate-Forced Variability of Ocean Hypoxia, Science, 333, 336–339, 2011.

510 DeVries, T., Deutsch, C., Primeau, F., Chang, B., and Devol, A.: Global rates of water-column denitrification derived from nitrogen gas measurements, Nature Geosci., 5, 547–550, 2012.

Di Lorenzo, E.: The future of coastal ocean upwelling, Nature, 518, 310-311, 2015.

- Espinoza-Morriberón, D., Echevin, V., Colas, F., Tam, J., Ledesma, J., Vásquez, L., and Graco, M.: Impacts of El Niño events on the Peruvian upwelling system productivity, J. Geophys. Res., 122, 5423–5444, 2017.
- 515 FAO: The State of World Fisheries and Aquaculture 2018 Meeting the sustainable development goals, Food and Agriculture Organization of the United Nations, Rome, 2018.
 - Farías, L., Castro-González, M., Cornjeo, M., Charpentier, J., Faúndez, J., Boontanon, N., and Yoshida, N.: Denitrification and nitrous oxide cycling within the upper oxycline of the eastern tropical South Pacific oxygen minimum zone, Limnol. Oceanogr., 54, 132–144, 2009.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for Upwelling of Corrosive 'Acidified' Water onto
 the Continental Shelf, Science, 320, 1490–1492, 2008.
 - Fiedler, B., Fietzek, P., Vieira, N., Silva, P., Bitting, H. C., and Körtzinger, A.: In Situ CO₂ and O₂ Measurements on a Profiling Float, J. Atmos. Ocean Technol., 30, 112 126, 2013.
 - Franco, A. C., Gruber, N., Frölicher, T., and Kropuenske Artman, L.: Contrasting Impact of Future CO₂ Emission Scenarios on the Extent of CaCO3 Mineral Undersaturation in the Humboldt Current System, J. Geophys. Res., 123, 2018–2036, 2018.
- 525 Frey, C., Bange, H. W., Achterberg, E. P., Jayakumar, A., Löscher, C. R., Arévalo-Martínez, Léon-Palmero, E., Sun, M., Sun, X., Xie, R. C., Oleynik, S., and Ward, B. B.: Regulation of nitrous oxide production in low-oxygen waters off the coast of Peru, Biogeosci., 17, 2263–2287, 2020.
 - Ganesh, S., Parris, D. J., Delong, E. F., and Stewart, F. J.: Metagenomic analysis of size-fractionated picoplankton in a marine oxygen minimum zone, ISME, 8, 187–211, 2014.
- 530 Ganesh, S., Bristow, L. A., Larsen, M., Sarode, N., Thamdrup, B., and Stewart, F. J.: Size-fraction partitioning of community gene transcription and nitrogen metabolism in a marine oxygen minimum zone, ISME, 9, 2682–2696, 2015.

Garreaud, R. D.: A plausible atmospheric trigger for the 2017 coastal El Niño, Int. J. Climatol., 38, e1296-e1302, 2018.

Grasshoff, K., Ehrhardt, M., and Kremling, K., eds.: Methods of Seawater Analysis. Second, Revised and Extended Edition, Verlag Chemie, Weinheim, Deerfield Beach, 1983.

- 535 Hamersley, M. R., Lavik, G., Woebken, D., Rattray, J. E., Lam, P., Hopmans, E. C., Sinninghe Damsté, J. S., Krüger, S., Graco, M., Gutiérrez, D., and Kuypers, M. M. M.: Anaerobic ammonium oxidation in the Peruvian oxygen minimum zone, Limnol. Oceanogr., 52, 923–933, 2007.
 - Hamme, R. and Emerson, S.: The solubility of neon, nitrogen and argon in distilled water and seawater, Deep-Sea Res. I, 51, 1517–1528, 2004.
- 540 Hauri, C., Gruber, N., McDonnell, A. M. P., and Vogt, M.: The intensity, duration, and severity of low aragonite saturation state events on the California continental shelf, Geophys. Res. Lett., 40, 3424–3428, 2013.
 - Holtappels, M., Lavik, G., Jensen, M. M., and Kuypers, M. M. M.: Methods in Enzymology, vol. 486, chap. ¹⁵N-Labeling Experiments to issect the Contributions of Heterotrophic Denitrification and Anammox to Nitrogen Removal in the OMS Waters of the Ocean, pp. 223–251, Elsevier, 2011.
- 545 Jensen, M. M., Kuypers, M. M. M., Lavik, G., and Thamdrup, B.: Rates and regulation of anaerobic ammonium oxidation and denitrification in the Black Sea, Limnol. Oceanogr., 53, 23–36, 2008.

Jensen, M. M., Lam, P., Revsbech, N. P., Nagel, B., Gaye, B., Jetten, M. S. M., and Kuypers, M. M. M.: Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium, ISME J., 5, 1660–1670, 2011.

Junk, G. and Svec, H. J .: The absolute abundance of the nitrogen isotopes in the atmosphere and compressed gas from various sources,

- 550 Geochim. Cosmochim. Acta, 14, 234–243, 1958.
 - Kalvelage, T., Jensen, M. M., Contreras, S., Revsbech, N. P., Lam, O., Günter, M., LaRoche, J., Lavik, G., and Kuypers, M. M. M.: Oxygen Sensitivity of Anammox and Coupled N-Cycle Processes in Oxygen Minimum Zones, PLoS One, 6, e29 299, 2011.
 - Kalvelage, T., Lavik, G., Lam, P., Contreras, S., Arteaga, L., Löscher, C. R., Oschlies, A., Paulmier, A., Stramma, L., and Kuypers, M. M. M.: Nitrogen cycling driven by organic matter export in the South Pacific oxygen minimum zone, Nature Geosci., 6, 228–234, 2013.
- Kämpf, J. and Chapman, P.: Upwelling Systems of the World, chap. The Functioning of Coastal Upwelling Systems, Springer, 2016.
 Koeve, W. and Kähler, P.: Heterotrophic denitrification vs. autotrophic anammox quantifying collateral effects on the oceanic carbon cycle, Biogeosci., 7, 2327–2337, 2010.
 - Lam, P. and Kuypers, M. M. M.: Microbial nitrogen cycling processes in oxygen minimum zones, Ann. Rev. Mar. Sci., 3, 317–345, 2011. Lam, P., Lavik, G., Jensen, M. M., Van Vossenberg, J. D., Schmid, M., Woebken, D., Gutiérrez, D., Amann, R., Jetten, M. S. M., and Kuypers,
- 560 M. M. M.: Revising the nitrogen cycle in the Peruvian oxygen minimum zone, Proc. Natl. Acad. Sci. U. S. A., 106, 4752–4757, 2009.
- Michiels, C. C., Huggins, J. A., Giesbrecht, K. E., Spence, J. S., Simister, R. L., Varela, D. E., Hallam, S. J., and Crowe, S. A.: Rates and Pathways of N₂ Production in a Persistently Anoxic Fjord: Saanich Inlet, British Columbia, Frontiers Mar. Sci., 6, https://doi.org/10.3389/fmars.2019.00027, 2019.
- Montecino, V. and Lange, C. B.: The Humboldt Current System: Ecosystem components and processes, fisheries, and sediment studies, Progr. Oceanogr., 83, 65–79, 2009.
 - Oschlies, A., Duteil, O., Getzlaff, J., Koeve, W., Landolfi, A., and Schmidtko, S.: Patterns of deoxygenation: Sensitivity to natural and anthropogenic drivers, Phil. Trans. Royal Soc. A, 375, 2017.
 - Paulmier, A. and Ruiz-Pino, D.: Oxygen minimum zones (OMZs) in the modern ocean, Progr. Oceanogr., 80, 113–128, 2009.

Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models,
Biogeosci., 6, 923–935, 2009.

- Peng, X., Fuchsman, C. A., Jayakumar, A., Warner, M. J., Devol, A. H., and Ward, B. B.: Revisiting nitrification in the Eastern Tropical South Pacific: A focus on controls, J. Geophys. Res., 121, 1667–1684, 2016.
- Peters, B. D., Babbin, A. R., Lettmann, K. A., Mordy, C. W., Ulloa, O., Ward, B. ., and Casciotti, K. L.: Vertical modeling of the nitrogen cycle in the eastern tropical South Pacific oxygen deficient zone using high-resolution concentration and isotope measurements, Glob.
- 575 Biogeochem. Cycl., 30, 1661–1681, 2016.
 - Poornima, D., Shanthi, R., Ranith, R., Srangi, R. K., Saravanakumar, A., and Thangaradjou, T.: Computational Methods for Long Term Monitoring of New Production and f-ratio Variability in Bay of Bengal, Int. J. of Oceans and Oceanogr., 14, 109–124, 2020.
 - Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of seawater, *In*: The Sea, 2nd edn., Hill, M. N. (Ed.), Wiley, New York, pp. 26–77, 1963.
- 580 Santoro, A., Buchwald, C., Knapp, A., Berelson, W., Capone, D., and Casciotti, K.: Nitrification and Nitrous Oxide Production in the Offshore Waters of the Eastern Tropical South Pacific, Glob. Biogeochem. Cycl., 35, 2021.
 - Schmid, M. C., Hooper, A. B., Klotz, M. G., Woebken, D., Lam, P., Kuypers, M. M. M., Pommerening-Roeser, A., Op Den Camp, H. J. M., and Jetten, M. S. M.: Environmental detection of octahaem cytochrome c hydroxylamine/hydrazine oxidoreductase genes of aerobic and anaerobic ammonium-oxidizing bacteria, Environ. Microbiol., 10, 3140–3149, 2008.

- 585 Schulz, K. G. and Riebesell, U.: Diurnal changes in seawater carbonate chemistry speciation in a high CO₂ world, Mar. Biol., 160, 1889– 1899, 2013.
 - Schulz, K. G., Hartley, S., and Eyre, B. D.: Upwelling Amplifies Ocean Acidification on the East Australia n Shelf: Implications for Marine Ecosystems, FMARS, 6, 636, https://doi.org/10.3389/fmars.2019.00636, 2019.
 - Stramma, L., Johnson, G. C., Sprintall, J., and Mohrholz, V.: Expanding Oxygen-Minimum Zones in the Tropical Oceans, Science, 320, 655–658, 2008.
 - Thamdrup, B., Dalsgaard, T., Jensen, M. M., Ulloa, O., Farías, L., and Escribano, R.: Anaerobic ammonium oxydation in the oxygen-deficient waters off northern Chile, Limnol. Oceanogr., 51, 2145–2156, 2006.

Toggweiler, J.: Carbon overconsumption, Nature, pp. 210-211, 1993.

590

Tyson, R. V. and Pearson, T. H.: Modern and Ancient Continental Shelf Anoxia, vol. 58, chap. Modern and ancient continental shelf anoxia:

an overview, pp. 1–24, Geological Society Special Publication, 1991.

- Varela, R., Álvarez, I., Santos, F., deCastro, M., and Gómez-Gesteira, M.: Has upwelling strengthened along worldwide coasts over 1982-2010?, Sci. Reports, 5, 10016, 2015.
- Vijayaraghavan, S. and Krishna Kumari, L.: Primary production in the southeastern Arabian Sea during southwest monsoon, Ind. J. of Mar. Sci., 18, 30–32, 1989.
- 600 Wang, D., Gouhier, T. C., Menge, B. A., and Ganguly, A. R.: Intensification and spatial homogenization of coastal upwelling under climate change, Nature, 518, 390–394, 2015.

Wanninkhof, R.: Relationship Between Wind Speed and Gas Exchange Over the Ocean, J. Geophys. Res., 97 (C5), 7373–7382, 1992.

Ward, B. B.: How nitrogen is lost, Science, 341, 352-353, 2013.

- Ward, B. B., Devol, A. H., Rich, J. J., Chang, B. X., Bulow, S. E., Naik, H., Pratihary, A., and Jayakumar, A.: Denitrification as the dominant
 nitrogen loss process in the Arabian Sea, Nature, 461, 78–81, 2009.
 - Yang, S., Gruber, N., Long, M. C., and Vogt, M.: ENSO driven variability of denitrification and suboxia in the Eastern Tropical Pacific Ocean, Glob. Biogeochem. Cycl., 31, 1470–1487, 2017.

Zhang, X., Ward, B. B., and Sigman, D. M.: Global Nitrogen Cycle: Critical Enzymes, Organisms, and Processes for Nitrogen Budgets and Dynamics, Chem. Rev., 120, 5308–5351, 2020.

610 Zumft, W.: Cell Biology and Molecular Basis of Denitrification, Microbiol. Mol. Biol. Rev., 61, 533-616, 1997.

Table 1. Concentrations of dissolved inorganic and organic nitrogen and phosphorus $(\mu \text{mol } L^{-1})$, as well as silicate in the two deep water batches and associated inorganic N/P/Si in comparison to a Redfieldian ratio of 16/1/16 (see text for details), as well as the inorganic N-deficit calculated as $N^* = \text{DIN} - 16 \times \text{PO}_4^{3-}$, with DIN denoting the sum of $\text{NO}_3^- \text{ NO}_2^-$ and NH_4^+ .

	NO_3^-	$\frac{NO_2^{2-}}{NO_2^{-}}$	NH_4^+	DON	PO_4^{3-}	DOP	$\mathbf{Si}(\mathbf{OH})_{4}$	N/P/Si	
Moderate DW-Low N/P DW	1.1	2.9	0.3	5.5	2.5	0.2	19.6	1.7 / 1 / 7.8	
Extreme DW-Very low N/P DW	0	0	0.3	5.2	2.6	0.2	17.4	0.1 / 1 / 6.7	

Table 2. Measured denitrification rates $(\text{nmol N}_2 \text{ L}^{-1} \text{h}^{-1})$ in the moderate (blue) low N/P and extreme (red) very low N/P deep water addition incubations and the surrounding Pacific, together with those for ananyox $\operatorname{ananyox}^*(\operatorname{nmol} \operatorname{N}_2 \operatorname{L}^{-1}\operatorname{h}^{-1})$ in green, when encountered, at various days. Numbers in brackets refer to the maximum in-situ denitrification estimates based on measured NO_3^- and NO_2^- concentrations. N-loss refers to the total amount of nitrogen (μ mol N L⁻¹) being lost through these processes for the first 38 days, for which an N-budget can be calculated (compare Fig. 4). N-loss was calculated from the rate mean (in case the incubation denitrification potential was higher than the maximum supply rate, the latter was used) for each mesocosm until day 38, assuming an average contribution of bottom water to the overall mesocosm volume of one third. For details on calculations see Methods.

Mesocosm				
Jow N/P	T8	T12	T16	T22
M1 6.68 18.65 13.8 32.34 30.62 24.79 29.32 0.89 3.9 7.39 4.45 - 3.44 (4.21) (1.30) (5.13) (11.26) M2 <u>M2</u>	14.99	7.77	1.96	28.92
				(3.37)
€W -ew	4.70	10.74	10.23	8.51
M4-M6	0.74- 3.86	43.33 - 20.46	11.46 3 .32	7.51- 7.08
	5.33 5 .94	82.23 3 .92	4.05 - <u>4.62</u>	25.34 -14.15
		$ \underbrace{0.44^*}_{0}$		$(\frac{2.30}{2.30}, \frac{30}{2.30})$
M6		3.86	Mean±SD	<mark>20.46−</mark> 2.85±1
very low N/P	7.08			
ĨŴ	16.74 6 . <u>68</u>	16.93- 18.65	14.37- 13.8	16.24- 32.34
			(2.83)	(1.664 .21)
M7-M4	5.94 0.74	3.92 4 3.33	4.62 - 11.46	14.15- 7 <u>.51</u>
		0.44		
Ĩ	<u>5.33</u>	82.23	4.05	25.34
				(2.79 2.30)
8W-8W-	10.81	7.85	4.67	13.23
		0.46 0.46*		(2.07)
PACIFIC	9.19	46.52	7.45	18.12

3.80

Mean

0.16*

7.48 (3.49)7.48*

0.16-(3.49)

Table 3. Multiple linear regression statistics, describing denitrification in response to various environmental variables, for the second-best best 4 variable model (in terms of \mathbb{R}^2) with interactions (compare Fig. 5C&D), as well as the best 3 variable model.

	Estimate	SE	t	р		Estimate	SE	
(Intercept)	2.772- 0.485	1.754- 1.050	1.580_ 0.462	0.120 -0.646	(Intercept)	-0.041	0.419	-0.098
PON DON	-0.101-0.040	0.129- 0.117	- 0.780- 0.341	0.438- 0.734	PON	0.113	0.054	2.10
$\frac{NO^{-}}{2} \underbrace{PON}{}$	33.28 0.114	5.606- 0.052	5.937-2.201	<u><0.001-0.032</u>	NO_3^-	1.060	0.256	4.13
$\Theta_2 NO_3$	-0.074- 1.438	0.040- 0.884	-1.852- 1.628	0.069- 0.109	NO_2^-	-6.037	3.271	-1.84
$H_2SNO_2^-$	-0.747_26.621	0.342- 10.761	-2.188 2.474	0.033 0.016	$PON:NO_3^-$	-0.086	0.028	-3.11
PON:NO ⁻ 2DON:NO ⁻ 3	-0.674 0.236	0.303- 0.086	-2.227- 2.749	0.030 <u>0.008</u>	$\mathrm{PON:NO}_2^-$	1.34	0.431	3.11
PON:H ₂ SDON:NO ₂	0.056 - <u>3.048</u>	0.027- 0.959	2.088_ - <u>3.179</u>	0.041 0.002	$\mathrm{NO}_3^-:\mathrm{NO}_2^-$	-1.412	0.2891	-4.88
$\frac{\mathrm{NO}_{2}^{-}:\mathrm{O}_{2}\mathrm{PON}:\mathrm{NO}_{3}^{-}}{\mathrm{PON}:\mathrm{NO}_{3}^{-}}$	-0.321 - <u>0.072</u>	0.054 0.027	-5.988 -2.637	<u><0.001-0.011</u>				
$\frac{\mathrm{NO}_{2}^{-}:\mathrm{H}_{2}\mathrm{S}\mathrm{PON}:\mathrm{NO}_{2}^{-}}{\mathrm{NO}_{2}^{-}}$	-1.950_1.018	0.524 0 .432	-3.721 2.355	< 0.001 _0.022				
$\Theta_2:H_2SNO_3:NO_2$	0.017- - <u>1.256</u>	0.007- 0.373	2.561 _ - <u>3.369</u>	0.013- 0.001				
			\mathbf{R}^2	0.5635 0.5710				R



Figure 1. Schematic reaction diagram of major marine nitrogen cycling processes, with (1) Hydroxylamine Oxidation, (2) Nitrifier Denitrification, (3) Denitrification, (4) Anammox, (5) Nitrification, (6) Nitrate Ammonification, aka DNRA, Anaerobic Nitrite Oxidation (7) and (8) Nitrogen Fixation. While processes (1–4) are considered nitrogen loss processes, (5–7) constitute neither a loss nor a gain, with (8) being the latter. Blue colours denote oxic and red suboxic/anoxic processes. Please note that the reactions have been chemically balanced and for electroneutrality, the exact amount of H^+ , e^- or water produced/consumed will depend on the actual organism/enzyme and reaction pathway. See Bourbonnais et al. (2017); Codispoti et al. (2005); Zumft (1997) and references therein for details.



Figure 2. Temporal development of upwelling and stratification in the surrounding Pacific waters at the mooring site (water depth between 25 and 30 metres) as evidenced by changes in CTD-derived (A) temperature (°C), (B) salinity (psu), (C) oxygen saturation (%) and (D) pH_T (total scale) depth profiles. Note that dips in salinity at the surface correspond with El Niño related torrential rain events further land inwards and increased discharge of freshwater by the nearby river Rimac.



Figure 3. Temporal evolution of depth-integrated bottom layer (A) PON, (B) DON, (C) nitrate (NO_3^-) , (D) nitrite (NO_2^-) , together with CTD-derived (E) oxygen (O_2) and (F) hydrogen sulphide (H_2S) concentrations at the Niskin sampling depth, in the bottom layer mesocosms (mostly between 10 or 12.5 and 17 mdepthM1-M8), with blue and the surrounding Pacific. Blue and red colours denote the extreme 'low N/P' and moderate 'very low N/P' deep water additions, respectively (see section 2.2 for details). The dashed green lines denote deep water additions on days 11 and 12, while the dotted black lines denote additions of a brine solution to the bottom layer to increase salinity, strengthen stratification and reduce mixing. See Methods for details.



Figure 4. Representative example of a total nitrogen budget from mesocosm M7, considering all relevant pools such as particulate organic nitrogen (PON), dissolved organic nitrogen (DON), dissolved inorganic nitrogen in the form of nitrate (NO_3^-) , nitrite (NO_2^-) and ammonium (NH_4^+) , cumulative particulate organic nitrogen exported to the sediment trap (N_{sed}) and the net change to all the above mentioned nitrogen species (with the exception of PON for which there was no deep water data) by deep water addition (N_{DW}) . Black horizontal markers denote the total amount of nitrogen in these pools, calculated as an average of three consecutive sampling days at the start of the experiment (days 1-3) and towards the end (days 36-40), prior to the onset of orni-eutrophication. The deficit in this N-budget comprises all nitrogen loss processes, dominated by denitrification (compare Tab. 2). Please note that the initial dip in nitrogen inventory is probably the result of a lag phase of nitrogen settling into the sediment trap.



Figure 5. Stepwise multiple linear regressions (MLRs) of measured/maximum sustained_maximum-sustained_denitrification rates in the mesocosms against up to 7 potential predictors and their interactions (PON, PON_{sed}, DON, NO₃⁻, NO₂⁻, O₂ and H₂S), with (**A**) reducing numbers of measured variables and resulting maximum R^2 . The red arrows denote the relatively small increase in maximum R^2 beyond 4 main variables and the relatively large decrease below, while the numbers in brackets denote the number of possible predictor combinations, i.e. MLRs fitted. (**B**) Main effect sizes of the stepwise MLR with four main variables and highest (0.571) and (**C**) second highest (0.564)- R^2 . (**P**C) Linear fit through measured and predicted denitrification rates (nmol N₂ L⁻¹h⁻¹) by the MLR shown in (**CB**).



Figure A1. Example of nitrogen removal during label incubations (Pacific on Day 8) as calculated according to Eq. 1. Denitrification or anammox rates were calculated from the slope of a linear fit to the data. It is acknowledged that in some samples a lag-phase was encountered.



Figure A2. Temporal evolution of depth-integrated (0-17 m) phosphate concentrations in the mesocosms (**A**), together with total particulate phosphorus (TPP) accumulating in the sediment traps (**B**). The black vertical line marks the onset of orni-eutrophication. <u>Blue and red colours</u> denote the 'low N/P' and 'very low N/P' deep water additions, respectively (see section 2.2 for details) in the various mesocosms (M1-M8).



Figure A3. Water discharge rates at station Chosica in the Rimac, \sim 45 km upstream the river mouth, and the latter being \sim 5 nm from the mesocosm mooring site. Red lines denote experimental days at which significant reductions in surface water salinities around the mesocosms were measured (compare Fig. 2). Note that the third measured surface freshening does not seem to coincide with such high discharge rates as for the first two. The freshwater could therefore stem from the River Chillón, \sim 9 nm to the North.



Figure A4. Particulate, dissolved and total organic carbon and nitrogen in the mesocosms' surface waters, and resulting ratios. Style and colour code follow those shown in Fig. $\frac{3}{2}A_2$, while black shows data for the surrounding Pacific