

Quantifying functional consequences of habitat degradation on a Caribbean coral reef

Alice E. Webb¹, Didier M. de Bakker^{2,3}, Karline Soetaert⁴, Tamara da Costa¹, Steven M. A. C. van Heuven⁵, Fleur C. van Duyl², Gert-Jan Reichart^{1,6}, Lennart J. de Nooijer¹

5 ¹ Department of Ocean Systems, NIOZ Royal Netherlands Institute for Sea Research, Den Hoorn, The Netherlands

² Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, Den Hoorn, The Netherlands

³ Wageningen Marine Research, Wageningen University and Research, Den Helder, The Netherlands

⁴ Department of Estuarine and Delta Systems, NIOZ Royal Netherlands Institute for Sea Research, Yerseke, The Netherlands

10 ⁵ Groningen University, Faculty of Science and Engineering, Groningen, The Netherlands

⁶ Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands

Correspondence to: Alice E. Webb (webbea4@gmail.com)

Abstract. Coral reefs are declining worldwide. The abundance of corals has decreased alongside a rise of filter feeders, turf, and algae in response to intensifying human pressures. This shift in prevalence of functional groups alters the biogeochemical 15 processes in tropical water ecosystems, thereby influencing reef biological functions. An urgent challenge is to understand the functional consequences of these shifts to develop suitable management strategies that aim at preserving the biological functions of reefs.

Here, we quantify biogeochemical processes supporting key reef functions (i.e., net community calcification (NCC) and production (NCP), and nutrient recycling) *in situ* for five different benthic assemblages currently dominating shallow degraded 20 Caribbean reef habitats. To this end, a transparent custom-made enclosure was placed over communities dominated by either one of five functional groups: coral, turf and macroalgae, bioeroding sponges, cyanobacterial mats, or sand, to determine chemical fluxes between these communities and the overlying water, during both day and night. To account for the simultaneous influence that distinct biogeochemical processes have on measured ~~fluxes~~variables, the rates ~~are~~were then derived by solving a model consisting of differential equations describing the contribution of each process to the measured 25 chemical fluxes.

Inferred rates were low compared to those known for reef flats worldwide. Reduced accretion potential was recorded, with negative or very modest net community calcification rates by all communities. Net production during the day was also low, suggesting limited accumulation of biomass through photosynthesis and remineralisation of organic matter at night was relatively high in comparison, resulting in net heterotrophy over the survey period by most communities. Estimated recycling 30 through nitrification and denitrification were high but denitrification did not fully counterbalance nutrient release from aerobic mineralisation, rendering all substrates sources of nitrogen. Results suggest similar directions and magnitudes of key biogeochemical processes of distinct communities on this shallow Curaçao reef. We infer that the amount and type of organic matter released by abundant algal turfs and cyanobacterial mats on this reef likely enhances heterotroph activity and stimulates

the proliferation of less diverse copiotrophic microbial populations, rendering the studied reef net heterotrophic and drawing
35 the biogeochemical ‘behaviour’ of distinct communities closer to each other.

1 Introduction

Community composition and biodiversity across all kinds of ecosystems are responding to escalating anthropogenic activities (McGill et al. 2015). In both terrestrial and aquatic systems, climate change, pollution and habitat fragmentation have promoted the expansion of opportunistic and tolerant species and the elimination of more sensitive yet key specialists (Clavel et al. 2011).
40 Communities within ecosystems and across spatial scales have become more biologically homogeneous (Burman et al. 2012; Cramer et al. 2021) which may lead to a decrease in functional diversity, therefore limiting services provided by biological communities (Matsuzaki et al. 2013; White et al. 2018). *Furthermore**Additionally*, this may cause synchronisation of the biological response to new or intensified anthropogenic pressures across local communities, thus reducing resilience of metacommunities (Tobias and Monika, 2012; Sonnier et al. 2014; Petsch et al. 2020).
45 Coral reefs support immense biodiversity and provide important ecosystem services to millions of people (Moberg and Folke, 1999). They are, however, in global decline as they are experiencing major loss in coral abundance and shifts in species composition in response to increasing human pressures and accelerating rates of environmental and climate change (Koop et al. 2001; Langdon and Atkinson, 2005; Andersson and Gledhill, 2013; De’ath et al. 2012; Chen et al. 2015). Returning degraded reefs to their original state is, in many cases, no longer an option (Hughes et al. 2017). Instead, today’s challenge is
50 to guide coral reefs through this transition while identifying and securing the ecosystem functions that underpin resilience and services of modern reef assemblages (Oliver et al. 2015).

This is particularly relevant for the depauperate reef systems in the Caribbean, which have, since the early 1970s, undergone considerable reorganisation with regards to community composition and structural appearance (Gardner et al. 2003; Jackson et al. 2014). The communities encountered on these reefs bear little resemblance to the systems once dominated by reef-
55 building *Acropora* spp. and *Orbicella* spp. (van Duyl, 1985; Alvarez-Filip et al. 2009). Major declines in the abundance of these species severely compromise reef function as they were the main drivers of critical processes including carbonate accretion, productivity, and structural complexity (Wolfe et al. 2020). This has led to low functional redundancy on Caribbean reefs, i.e., a reduced capacity of one or more species to functionally compensate for the loss of another, which makes them particularly vulnerable (Bellwood et al. 2003; McWilliam et al., 2018). On many reefs, areas covered by turf assemblages and
60 macroalgae, excavating sponges, cyanobacteria, rubble, and sand have increased alongside the decrease in stony corals (Aronson et al. 2005; Burman et al. 2012; Cramer et al. 2021). Although changes in community composition are well documented as they can be followed by monitoring the coverage of the various benthic taxa over time (Barott et al. 2012; de Bakker et al. 2016; de Bakker et al. 2017), assessing the impact of these shifts on the community ecophysiology in situ has proven more challenging.

65 The keystones of coral reef functioning include provision of a structural habitat through carbonate deposition, production and assimilation of biomass produced through photosynthesis and efficient cycling of nutrients within the ecosystem (Brandl et al. 2019). The biogeochemical processes that underlie these key functions are primary production, aerobic mineralisation, calcification, bioerosion, and nutrient release/uptake. Complementary to conventional monitoring efforts, quantification of the net budgets of these processes will provide insight into how reef degradation and community reorganisation affect reef
70 functioning (Brandl et al. 2019, Bellwood et al. 2019). However, obtaining accurate in situ measurements while accounting for the complexity of interactions between processes can render their quantification rather complicated.

In environments where the flow of water over the reef is relatively linear, the upstream/downstream method can be performed (Shaw et al., 2014; Kowek et al., 2015; Albright et al., 2016, 2018). For flow regimes that are not unidirectional, factors such as water residence time and biochemical and hydrological offshore conditions need to be considered (Courtney et al., 2016).

75 When conditions allow build-up of considerable chemical vertical gradients, net fluxes (of e.g., nutrients) can be measured (McGillis et al., 2011; Takeshita et al., 2016). On fully exposed reefs, where virtually no detectable accumulation occurs over the reef-flat - even within the boundary layer - incubating communities allow quantification of the fluxes into and out of the overlying water. Presently, efforts to quantify community processes have focused on individual functional groups (Brocke et al. 2015, 2018; den Haan et al. 2016; Webb et al. 2017; de Bakker et al. 2018) or on reconstructed communities ex situ (e.g.,
80 Dove et al. 2013, 2020). Moreover, the few studies that incubated whole communities in situ have so far not accounted for the complexity of interactions between biogeochemical processes (Yates and Halley, 2003, Kline et al. 2012, van Heuven et al. 2018, Roth et al. 2020, 2021).

Here, biogeochemical processes underlying key reef functions were quantified in situ across five different benthic assemblages found on the fringing reef of Curaçao, consisting of functional groups that currently characterise many degraded shallow reef
85 habitats throughout the wider Caribbean. To this end, a custom-made tent was placed over substrates dominated by either 1) coral, 2) turf and macroalgae, 3) bioeroding sponges, 4) benthic cyanobacteria mats or 5) sand. Chemical fluxes between water column and reef were then determined by monitoring nutrients, inorganic carbon chemistry and oxygen. This was done both during the day and at night to estimate overall net metabolism of these communities. To account for the simultaneous convoluted influence that various processes have on measured variables, the change in their concentrations is related to the
90 responsible metabolic processes by solving a model consisting of ordinary differential equations describing the contribution of each process to the measured chemical fluxes. With this approach, model parameters (i.e., rates of biogeochemical processes) are derived from concurrent changes in all measured variables. The aim being to provide accurate estimates of the rates of the biogeochemical processes that underlie functions of the newly configured shallow Caribbean reefs.

2 Materials and Methods

95 2.1 Study Site

Reef incubations were carried out on the leeward side of Curaçao (Piscadera Bay; $12^{\circ}07'16.3''N$ $68^{\circ}58'13.2''W$) between the 12th of February and 22nd of March 2018, at depths ranging from 5 to 7 meters. The water at the study site is characterised by episodes of high turbidity and is periodically eutrophied due to terrestrial runoff and ineffective waste-water treatment. Sediment plumes transporting high concentrations of nitrate, ammonium and phosphate into the shore's fringing reef are 100 commonly encountered after a period of heavy rainfall (den Haan et al. 2016). The shallow reef flat nearby the entrance of the bay in which we conducted our incubations is characterised by rubble and patchy distribution of small coral heads making this location particularly suitable for the deployment of tent incubations.

2.2 Tent Incubations

The incubation enclosure consists of a custom made, tetrahedron-shaped "tent" (Fig. 1). It has transparent, vinyl-and-butanyl 105 walls with rigid pole edges of 1 m in length, resembling the cBIT described by Haas et al. (2013). It also includes 0.5 m long flaps extending outward from each of the tent's three sides, allowing for better sealing of the tent to the substrate by placing weights (metal chains) on these flaps. The enclosure covers a 0.43 m^2 planar surface and encloses a 118 L volume. All three sides of the tent contained an opening to allow flushing of the enclosed volume between incubations: during incubations these 110 openings were sealed by zippers. Water enclosed in the incubation tent was homogenised during the experiment by means of a continuously running brushless submersible water pump (BLDC pump Co., Ltd.). This pump was attached to one of the tent poles, at half the height of the tent, generating a vertical circulating turbulence, while minimising the upsurge of sediment. Effectiveness of the stirring was demonstrated by rapid and even dispersal of a small dose of injected fluorescein prior to the 115 incubation. Surge movement was retained due to the non-rigid texture of the tent walls. Incubated communities included five different types of substrate dominated either by turf and macroalgae, sand, bioeroding sponges, benthic cyanobacteria mats or coral (Fig. 2), equalling a total of 15 studied communities (three of each type). Each community was incubated during the day (n=15) and due to practical reasons, only 2 of each type were incubated during the night (n=10) (i.e., for each type of 120 community, three daytime and 2 night-time incubations were carried out).

The incubations were carried out one at a time, over the study period and lasted four hours each. Prior to each incubation, the tent was placed with flaps open over the substrate and left for a minimum of 3 hours before the incubation was started. When 125 day incubations were terminated, the tent was left in place with flaps open so that night incubation could be carried out on the same substrate. Daytime incubations were started at 10:00 and night-time incubations started at 18:30.

2.3 Substrate Compositions

Substrates dominated by either coral, turf and macroalgae (TMA), bioeroding sponges (BES), cyanobacteria mats (BCM), or sand were incubated (Fig. 2). Three reef patches of each reef assemblage type were chosen depending on their cover of the

125 dominant benthic component (see Table S1 for detailed species composition and cover). In some cases, to fit adequate
incubation location and the tent capacity, pieces of rubble infested with sponge or covered in turf were added or retrieved from
the community. In these cases, the community was left to stabilise two or three days before starting incubations. Incubated
substrate included colonised hard substrate surrounded by bare hard substrate covered in a fine layer of sand for better enclosure
deployment (except for sand incubations). The incubated coral species are characteristic of degraded Caribbean reefs and
130 include some of the most prominent tolerant and opportunistic species found on modern reefs (Darling et al. 2012; de Bakker
et al. 2016; Cramer et al. 2021) (see Table S1). Turf here refers to the epilithic algal matrix, defined by Clements et al. (2016)
as ‘a conglomeration of short, turf-forming filamentous algae (< 1 cm high), macroalgae, microalgae, sediment, detritus
and associated fauna’. The benthic cyanobacterial mats in all three tent replicates were thick brown/reddish in colour and in
line with the description in Brocke et al. (2018) for mats found between 3 and 7 meters dominated by the species *Oscillatoria*
135 *bonnemaisonii*. Percentage cover was measured in situ after removal of the tent. For substrates dominated by coral, its cover
ranged from 34 to 36%. Turf and macroalgae cover ranged between 72 to 83%, bioeroding sponge cover varied from 38 to
40%, and cyanobacterial mats cover ranged from 83 to 91%, in their respective incubations (Fig. 2).

2.4 In Situ Measurements

140 Measurements of salinity (S), temperature (T), dissolved oxygen (O₂) and photosynthetically active radiation (PAR) within
the tent were recorded at one-minute intervals throughout the duration of the incubations. S and T were measured using a Star-
Oddi DST CTD, O₂ was recorded using a HOBO U26 dissolved oxygen sensor and data logger and PAR was assessed by an
Odyssey light logger (Dataflow Systems PTY Ltd., Christchurch, NZ), calibrated in air against a Walz instrument (Walz
ULM500, Walz GmbH, Effeltrich, Germany). In addition, S, T and PAR were measured for the duration of the incubations
outside the tent using the same sampling frequency. All instruments within the tent were attached to the ridges except the
145 Odyssey logger which was placed on the substrate facing upwards (covering approximately 150 cm² of the substrate).

2.5 Discrete Sampling

150 During each incubation, discrete samples were collected both inside and outside the tent at T₀, after two (T₂) and after four
hours (T₄) by SCUBA diving for the analyses of total alkalinity (A_T), total inorganic carbon (C_T) and nutrients (NO₂ + NO₃,
NO₂ and NH₄). pH was calculated from the former two parameters using the package Seacarb (Lavigne et al. 2009). Sampling
of the tent interior was carried out from the outside by drawing seawater through 150 ml plastic syringes connected to a 1.5 m
gas-impermeable tube (Tygon; Fig. 1). Syringes were flushed three times with the sampling water before collecting an actual
sample. The tubing was fixed around a rigid pole of the tent in such way that the seawater was sampled from the centre of the
tent incubation. The tube end located inside the tent was equipped with a Whatman ® filter (G/F 0.47 µm) which was replaced
daily to avoid the collection of particulate matter.
155 Analyses for A_T were performed within two hours upon sampling using spectrophotometrically guided single-step acid titration
(Liu et al. 2015) and samples for C_T were run on an autoanalyser Traacs 800 spectrophotometric system (Stoll et al. 2001) at

the NIOZ (Royal Netherlands institute for sea research). Accuracy of both instruments was set using certified reference material supplied by Scripps Institute of Oceanography (Dickson et al. 2007). Precision of replicates was $2.7 \mu\text{mol kg}^{-1}$ for C_T and $0.9 \mu\text{mol kg}^{-1}$ for A_T . Samples for dissolved inorganic macronutrients were prepared by dispensing sampled water through 160 $0.8/0.2 \mu\text{m}$ Acrodisk filters into 5 mL pony vials, and subsequently stored at -20°C until analysis at the NIOZ on a QuAAstro continuous flow analyser (SEAL Analytical, GmbH, Norderstedt, Germany) following GO-SHIP protocol (Hydes et al. 2010).

2.6 Rates of Water Exchange

After sampling water at T_0 for A_T , C_T , and nutrients in each incubation, 450ml of salt-saturated water was injected into the tent. The rate at which the elevated interior salinity equilibrated with ambient salinity during the incubation was used to 165 estimate the rate of water exchange with the surrounding sea water for each incubation.

The rate of change of salinity within the incubation can be solved by the differential equation below:

$$\frac{dS}{dt} = K (S_{\text{out}} - S)$$

Where $\frac{dS}{dt}$ is the rate at which salinity changes within the tent, S_{out} is the exterior salinity, S is the interior salinity and K is the water exchange rate.

170 The equation is solved using the function ‘ode’, within the package deSolve (Soetaert et al. 2010), the R routine that solves the differential equations. Function ‘modfit’ from the package FME (Soetaert and Petzoldt, 2010) was used to perform iterative minimisation (based on least squares) on residuals to find the best fit within lower and upper bounds.

2.7 Inverse Modelling and Model-Data Comparison

175 The use of inverse modelling is advantageous as it enables us to derive unknown parameters (here rates of biogeochemical processes) simultaneously from all measured data. The mathematical “state” of the incubation’s dynamic system can be described based on the mass balance between A_T , DIC , O_2 , NH_4 and NO_3 which is influenced by various biogeochemical processes. The rate of these processes are the unknown parameters that need to be quantified by fitting against an incomplete data set (only three time-points for A_T , DIC , NH_4 and NO_3).

180 The model consists of the five differential equations depicted below that relate the change in measured variable concentrations over time to the responsible processes, which are here assumed to have remained constant over time.

Since the involved processes affect the different chemical components simultaneously, the combination of these differential equations can be used to solve the contribution (in terms of rates) of the processes to the observed changes. The processes in 185 question include aerobic mineralisation (O_2 consumption related to mineralisation), primary production (PP), calcification, dissolution, nitrification and denitrification.

$$\begin{aligned}
\frac{dC_T}{dt} &= \text{mineralisation} - PP - \text{calcification} + \text{dissolution} + K (C_{Tout} - C_T) \\
\frac{dNH_4}{dt} &= \text{mineralisation} \times NC_{ratio} - PP \times NC_{ratio} \times Pnh4 - \text{nitrification} + K (NH_4out - NH_4) \\
\frac{dNO_3}{dt} &= -(mineralisation \times pDeni \times 0.8) - PP \times NC_{ratio} \times (1 - Pnh4) + \text{nitrification} + K (NO_3out - NO_3) \\
190 \quad \frac{dA_T}{dt} &= -2 \times \text{calcification} + 2 \times \text{dissolution} + \frac{dNH_4}{dt} - \frac{dNO_3}{dt} + K (A_{Tout} - A_T) \\
\frac{dO_2}{dt} &= -mineralisation \times OC_{ratio} \times (1 - pDeni) + PP \times OC_{ratio} - 2 \times \text{nitrification} + K (O_2meanout - O_2)
\end{aligned}$$

With mineralisation describing the degradation of an organic compound to its mineral components, i.e. carbon dioxide and inorganic nutrients. PP is the primary production and calcification is the deposition of calcium carbonate. Pnh4 is the part of N uptake as NH₄ for primary production. Dissolution results in an increase of calcium and carbonate ions by degradation of calcium carbonate shells and/or skeletons and K is the water exchange rate. Nitrification is the process by which ammonium (NH₄⁺) is converted into nitrate (NO₃⁻); two moles of oxygen are needed to oxidize one mole of ammonium during nitrification. pDeni is the fraction of mineralisation that respites nitrate (i.e., denitrification). The OC_{ratio} is the ratio between the concentrations of oxygen and C_T. The NC_{ratio} is the ratio between N and C_T. The 0.8 constant refers to the denitrification redox reaction (Soetaert et al. 2007).

We start by determining the parameters that can be fitted, based on parameter collinearity. After producing a best-fit set of the selected parameters, we quantify parameter uncertainty, and produce sensitivity ranges around the modelled variables.

The OC_{ratio}, NC_{ratio} and K parameters are always fixed and estimated from data prior to running the model. Others vary between fixed and free (to be fitted) depending on collinearity and light. For instance, primary production is fixed at 0 during night incubation, however during the day, only the dominant process can be estimated. Some parameters are highly correlated with each other such as primary production and remineralisation or calcification and dissolution and therefore cannot be estimated simultaneously. In general, when the collinearity index exceeds 20, the linear dependence is assumed to be critical (i.e., it will be impossible or difficult to estimate all the parameters in the combination together).

Collinearity of the parameter sets is measured using function 'collin' within the FME package (Soetaert and Petzold, 2010).

210 The model equations are specified in a function that calculates the rate of change of the state variables (dC_T, dNH₄, etc.). Inputs to the function are the model time (t), the values of the state variables (C_T, NH₄, NO₃, A_T and O₂) and the parameters (remineralisation, calcification, etc.). The differential equation model is solved using function 'ode', within the package deSolve, the R routine that solves the differential equations.

215 The discrepancy of the model solution with observed changes within the tents is calculated using function 'modCost' still in the FME package which estimates the residuals and the variable and model costs (sum of squared residuals).

Function ‘modfit’ was then used to perform iterative minimisation (based on least squares) on residuals to find the parameter giving the best fit within lower and upper bounds. Estimated parameters are the unknown fluxes (mineralisation, PP, calcification, etc.).

2.8 Conversion to fluxes

220 The best-fit parameters, i.e., the input rates R (in $\mu\text{mol kg}^{-1} \text{ min}^{-1}$), in the tent are converted to fluxes from the water-substrate interface ($\text{mmol m}^{-2} \text{ h}^{-1}$), assuming an enclosed mass of water of $108 \pm 10 \text{ kg}$ (tent encloses approximately 118 litres of volume; of which substrate volume is $\sim 10 \text{ L}$; seawater density $\sim 1022 \text{ kg m}^{-3}$) and an incubated planar surface of 0.43 m^2 . Net community calcification (NCC) fluxes were determined from the predicted calcification and dissolution. The model captures the dominant net flux and does not distinguish the relative contributions of gross calcification and dissolution to the 225 integrated NCC rate. Net community production (NCP) is the difference between remineralisation and primary production. Denitrification is estimated from the pDeni fraction and the mineralisation parameters.

2.9 Comparing biogeochemical signatures between incubations

230 To evaluate if water exchange rate had an impact on estimated processes, the non-parametric Kendall rank correlation test was performed. All inferred biogeochemical process rates (mineralisation, primary production, NCP, NCC, nitrification and denitrification) were tested against incubation water exchange rates.

Principal component analysis (PCA) was used to identify grouping among the 23 tent incubations (day $n = 13$, night: $n = 10$) in relation to their biogeochemical signature (i.e., NCC, NCP, nitrification and denitrification). The oxygen logger malfunctioned during two of the day incubations consisting of one BCM and one TMA- dominated community incubation. The model was therefore not run through these tents. PCA was conducted on a centred multivariate data set consisting of the 235 four main biogeochemical processes (i.e., NCC, NCP, nitrification and denitrification). Additionally, NCC was plotted against NCP to evaluate how the balance between both processes varied among distinct communities and by which process were communities dominated.

3. Results

3.1 Ambient conditions

240 In-tent light and temperature were only slightly impacted by the tent enclosure compared to the exterior (Fig. 3). Light was on average 17 % lower inside the tent and changes in temperature were damped within the tent. In-tent temperatures were on average $0.2 \text{ }^\circ\text{C}$ higher than those outside the tent. Average ambient A_T , DIC , pH , NH_4 and NO_3 was 2386.8 ± 13.9 , 2125.5 ± 20.0 , 7.9 ± 0.003 , 0.31 ± 0.15 and $0.32 \pm 0.14 \mu\text{mol kg}^{-1}$ respectively. Measured data for each incubation, inside and outside the tent for all three time-points, as well as the differences between T_0 and T_4 are presented in Fig. Table S2.

245

3.2 Water exchange quantification

Application of Equation (1) to salinity data collected during all incubations yielded dilution rate K ranging between 0.004 and 0.044 min^{-1} . This indicates that 2.1 to 4.8 kg of seawater (i.e., $K \times 108 \text{ kg}$) was exchanged every minute between the incubation enclosure and the environment. These rates correspond to the intensity of the water movement observed and recorded visually at the time of each incubation. Fig. 4 shows the data used to estimate the rate of water exchange of an incubation with relatively minor leakage (A) and one where leakage is more severe (B). In these examples, in-tent salinity returns to ambient concentrations after ~ 1 and ~ 2 hours respectively.

3.3 Model output

Figure 5 illustrates the output of our approach for all incubations. Using a minimisation routine, best fit parameters (mineralisation, PP, calcification, etc.) were predicted to best fit the model to the measured data. Individual graphic output for two incubations (including one carried out during the day on substrate dominated by BES and one performed at night-time on BCM-dominated substrate) with respective fixed and fitted parameters are presented in Fig. S1. Details for parameter prediction and best-fit can be found in Table S3. The model output shows a relatively good fit to the measured observations (Fig. 5), indicating that the interactions between processes and their effects on chemical fluxes were considered correctly. Overall fit is usually better on night data, which is mostly due to the inability of the model to predict irregular oxygen evolution caused by light variability during the daytime (Table S3).

As the process estimates are limited to net increase or decrease, fluxes for PP and mineralisation are presented as net community production (NCP) and calcification and dissolution are combined into net community calcification (NCC; Fig. 6).

265

3.4 Estimated biogeochemical processes

NCP showed a clear diurnal pattern (Fig. 6). While all NCP values were modestly skewed towards net autotrophy during the day (except for sand), the strongest signal was found for substrates dominated by BCM with an average daily NCP of 5.6 $\text{mmol m}^{-2} \text{ h}^{-1}$. Night values indicated net respiration ranging from an average of $-2.64 \text{ mmol m}^{-2} \text{ h}^{-1}$ on substrates dominated by TMA to $-16.28 \text{ mmol m}^{-2} \text{ h}^{-1}$ on substrates dominated by bioeroding sponges.

A clear diurnal signal also resided in NCC fluxes for all substrates involved (Fig. 6). Most NCC fluxes recorded during daytime (except for sand incubations) indicated net CaCO_3 precipitation. At night, most NCC fluxes indicated net CaCO_3 dissolution, especially on substrates dominated by BES and BCM. The absence of change in A_T for coral dominated substrates during the night indicated that dissolution equalled calcification during these incubations and hence, the average NCC at night was close to 0. Substrates dominated by coral generated the strongest decrease in A_T (net precipitation) during daytime, yielding an average NCC rates of $0.45 \text{ mmol CaCO}_3 \text{ m}^{-2} \text{ h}^{-1}$. Highest net dissolution was found at night-time for incubations of substrates

dominated by bioeroding sponges and cyanobacterial mats, with a comparable average of 0.56 and 0.63 mmol CaCO₃ m⁻² h⁻¹ respectively.

Nitrification was found to occur predominantly at night with higher fluxes in incubations of substrates dominated by bioeroding 280 sponges, cyanobacterial mats and corals. Denitrification also occurred mostly at night except on sand where daytime and night-time fluxes were small but relatively similar (Fig. 6).

The Kendall rank correlation test did not reveal significant correlation between water exchange rates and rates of mineralisation (p=0.79, tau=0.04), primary production (p=0.47, tau=0.12), NCP (p=0.75, tau=0.05), NCC (p=0.17, tau=0.21). Nitrification 285 (p=0.81, tau=0.04), and denitrification (p=0.27, tau=0.18). The Kendall correlation coefficient tau is closer to zero than to one in all cases, implying there is no significant association between the two tested variables.

3.5 Incubation comparison

The PCA based on the four main biogeochemical processes revealed different groups for night incubations and day incubations 290 (Figure 7A), except for sand incubations where night and day incubations grouped relatively close to each other. The first two principal component axes (PC1 and PC2) explained 88.68% of the total variability within the data. PC1 described a gradient in NCP and NCC from high (negative PCA scores) to low (positive PCA scores) and an opposite pattern for nitrification and denitrification. PC2 further explains the variability in NCC and nitrification, and to a lesser extent NCP and denitrification.

One of the communities dominated by bioeroding sponges (rep.1) is separate from other communities both during the day and 295 during the night due to considerably higher rates of nitrification and denitrification compared to other communities.

Figure 7B showed the position of the 15 communities in the different quadrants of the NCC vs. NCP diagram. A clear separation is observed between day incubations, characterised by net photosynthesis and calcification and night incubation indicating net respiration and dissolution. Day-time sand incubations are centred due to the low magnitude of the processes occurring on these substrates and are the only incubations exhibiting net dissolution during the daytime. For night-time 300 incubations, only coral-dominated communities displayed modest net calcification.

4. Discussion

The biogeochemical flux assessment has enabled us to identify and quantify the biological functions that are currently at play on this degraded Curaçao reef flat. Although the present study only investigated the shallow part of one single reef, it provides 305 insight into the ~~dire~~ effects that shifts in coral species and functional groups has had on the overall functioning of these communities. Comparison with coral reefs in a broader biogeographical context is needed to establish whether the rates obtained here are site-specific or representative of degrading Caribbean coral reefs in general.

The shallow reef communities investigated at this site barely support reef functions that are usually ascribed to a healthy coral reef. Overall, net community calcification and production on these substrates were low compared to reef flats worldwide (Atkinson, 2011). Very low or negative NCC rates were recorded on all substrates, suggesting reduced net accretion potential.

310 Net production was also low, likely indicating limited accumulation of biomass, while heterotrophic processes were prominent. Recycling processes, nitrification and denitrification, were high but did not prevent net nutrient release from aerobic mineralisation, rendering all substrates, sources of nitrogen. Although processes recorded on substrates dominated by coral, bioeroding sponges and cyanobacterial mats showed some variation between types of substrates, the overall behaviour of complementary processes for each of these assemblages was relatively comparable in terms of direction and magnitude.

315

4.1 Method considerations

Formatted: Font: Not Bold

~~Results indicate that the combination of in situ incubations and inverse modelling, which integrates the complexity of interactions between processes, is an effective tool to provide quantitative data on the functional state of coral reef patches.~~

320 ~~Quantification of the exchange of substances between reef communities and the overlaying water was achieved despite the presence of swell-induced seawater exchange, because this approach allowed for a volume exchange between the environment and the incubation. The incubations can thereby be replenished at some extent, keeping saturated O₂ levels within the tent, thus minimising unrepresentative reef community metabolism. Fluxes within our incubations can be treated as if acquired by an in situ flow through system.~~

325 ~~For the interpretation of the measured concentration differences, the multifaceted influence of metabolic processes on chemical fluxes was accounted for. The model shows a good fit of the observations around the fitted curve, indicating that the interactions between processes and their effects on chemical fluxes were considered correctly.~~

330 ~~Nonetheless, due the limited number of incubations that were carried out for this study, we interpret results with caution. Additionally, incubations were only deployed on the reef flat of one degraded reef site, future application of this or similar incubation methods should consider multiple sites. Lastly, methods would be further improved by continuous monitoring of the exchange rate, rather than assuming it to be constant throughout the incubation.~~

4.2 Net Dissolving Reef

Although net calcification was recorded during the day on all substrate types (except sand), it did not compensate for higher dissolution rates at night except on substrates dominated by coral. Diel shifts between net calcification and net dissolution are

335 ~~not uncommon for coral reefs and have been recorded on healthier reef systems than the one studied here (Yates and Halley, 2003; Albright et al. 2013; Albright et al. 2015; Kowek et al. 2015) with instances of net dissolution mainly taking place at night, coinciding with net respiration (and most likely with low gross calcification) (Cyronak et al. 2018). However, the community calcification budget over 24 hours resulting from this diurnal variability in the present study resulted in a modest average net calcification rate of 5.7 mmol CaCO₃ m⁻² day⁻¹ on coral-dominated substrates. This is low compared to~~

340 rates reported for reef flats worldwide (with an average around 130 mmol CaCO₃ m⁻² day⁻¹ and ranging from 20 to 250 mmol CaCO₃ m⁻² day⁻¹; Atkinson, 2011). Overall, the limited number of in situ flux-based experiments carried out in the wider Caribbean (Yates and Halley 2003; Muehllehner et al. 2016; van Heuven et al. 2018) suggest they are among the lowest NCC rates recorded worldwide (Albright et al. 2015; Shaw et al. 2015; Silverman, 2007) and particularly low compared to those in the Indo-Pacific region (Kowek et al. 2015; Takeshita et al. 2016). Surveys using a census-based approach (Perry et al. 2013; 345 de Bakker et al. 2019) also showed that some Caribbean reefs are net eroding. The recorded low rates of net carbonate production in the wider Caribbean may be expected simply due to the region-wide decrease in coral coverage since the 1970s. However, the decrease in net calcification relative to historical values is likely related to more than just coral cover loss. Indeed, the latter has subsequently left surfaces available for colonisation by turf, macroalgae (Hughes, 1994) and more recently, cyanobacterial mats (de Bakker et al. 2017). Shallow reefs around Curaçao (<10 m) are covered by filamentous algal turf 350 canopies that presently represent the most dominant benthic component on these reefs (Vermeij et al. 2010). Given their abundance and high release rates of dissolved organic carbon (Mueller et al. 2016), heterotrophic activity is likely to be stimulated. Furthermore, cyanobacterial mats release part of their photosynthetically fixed carbon as DOC into the water column at a higher rate than turf and macroalgae (Mueller et al. 2014; Brocke et al. 2015). They have been shown to be responsible for 79% of the total DOM release over a 24 h diel cycle at this same study site (Brocke et al. 2015). Considering 355 | their proliferation around the islands of Curaçao and Bonaire since ~2003 (Dede Bakker et al. 2017) and the prevalence of turf algae in the area, an accumulation of organic matter may have resulted in a reduction of pH due to oxidation of organic matter, i.e. stimulated heterotrophic activity, resulting in reduced calcification (Bates et al. 2010). Muehllehner et al. (2016) suggested that the seasonal character of reef dissolution they recorded on the Florida Reef Tract coincided with an accumulation of organic matter, following the die-off of annual sea grasses in the area.

360

4.32 Net heterotrophic reef

Low net community production rates in the current study indicate that autotrophic processes dominate modestly during the day. Integrating the NCP values over 24 hours (day + night) yielded rates skewed towards net respiration, indicating heterotrophy in all incubations. Although net community production of reef flats has been reported to vary notably over the 365 course of the day (Kowek et al. 2015), with values ranging from -220 to +310 mmol m⁻² day⁻¹ (Atkinson, 2011), large amplitude shifts between net autotrophy and net heterotrophy are usually recorded between day and night (Yates and Halley, 2003; Albright et al. 2013; Albright et al. 2015; Kowek et al. 2015). Here, the amplitude of this shift between day and night was modest. It should be noted however, that the reduction in light intensity ~~by 17~~ by 17 % on average may have resulted in a slight underestimation of NCP rates during the daytime. This would hold especially true for BCM incubations. Reductions in 370 light would intensify down the steep vertical physiochemical gradients present in these microbial mats and could interfere with light-controlled circadian regulation of photosynthesis and respiration in these cooperative communities (Hörlein et al. 2018), favouring respiration and decreasing net community productivity.

The reduction in the amplitude of the diel shift in net production and calcification recorded in the present study may have severe implications. For instance, metabolic fluctuations from reef biota cause strong temporal fluxes in compounds which
375 affect the oscillatory behaviour of reef seawater microbial communities (Kelly et al. 2019; Weber et al. 2020) leading to less distinct populations and more redundancy in microbial specialists' functions, i.e., a shift to a dominance in catabolic pathways. Organic material supplied to the ecosystem by benthic primary producers as exudates is also thought to play a pivotal role on microbial growth (Haas et al. 2011) and diversity (Nelson et al. 2013; Haas et al. 2016) depending on its origin. Studies on the effect of exudates of macroalgae and turf on microbial metabolism demonstrated that the composition of exudates stimulated
380 rapid growth of less diverse microbe communities compared to coral derived exudates. Consequently, reef microbial communities shift towards copiotrophic populations that have the potential to remineralise available organic nutrients at a high rate and encode greater numbers of potential virulence factor genes, ultimately harming corals and maintaining algal dominance (Nelson et al. 2013; Dinsdale and Rohwer, 2011). We infer that the amount and type of organic matter provided by abundant algal turfs mats on this reef, likely enhances heterotroph activity and stimulates the proliferation of less diverse
385 copiotrophic microbial populations, rendering the studied reef net heterotrophic regardless of substrate type.

4.43 Nitrogen cycling

Nitrogen pathways support high primary productivity in oligotrophic environments by supplying nutrients while simultaneously preventing the build-up of excess nutrients that may favour opportunistic primary producers such as algal turfs (e.g., O'Neil and Capone, 2008; Karcher et al. 2020). The abundance of non-coral primary producers on these reefs suggest
390 that nitrogen is not a limiting factor for growth. Results showed that all substrate types acted as NH_4^+ and NO_3^- sources during the day and the night, apart from sand and turf substrates which acted as sinks for NO_3^- . This is to be expected from overall net heterotrophic communities, however, even in instances of net autotrophy during the day, substrates still acted as DIN sources. This is comparable to recent results from in situ incubations carried out in the central Red Sea on net autotrophic coral and alga-dominated communities (Roth et al. 2020, Roth and 2021). It is likely that other community-wide processes, such
395 as the consumption and transformation of organic matter by microbial populations (e.g., Pfister and Altabet, 2019), masked the assimilation of DIN by primary producers.

Nitrification and denitrification rates measured in the present study generally fall within the published range of in situ measurements in tidal pools dominated by algae and corals (Webb and Wieber, 1975), in cavities covered in encrusting sponges (Scheffers et al. 2004), on cyanobacterial mats (Bonin and Michotey, 2006) and on carbonate sand (Capone et al. 1992; Eyre
400 et al. 2013b). However, there was no nitrification during the day (except for *theone* community dominated by sponges), which may be explained by light causing a reduced activity of nitrifiers (Kwon et al. 2020). Owing to the rather shallow depths of our experiment, nitrifiers may have been negatively affected by the light. As mentioned above, microbial communities are impacted by organic matter composition and temporal fluctuations in biochemicals. Shifts in diversity and abundance of the microbial communities inhabiting the reef substrate may also lead to diel shifts in nitrogen-cycling capacity (Rädecker et al.

405 2015). Further research investigating how alterations in diversity and abundance of these microbial functional groups relate to changes in the nitrogen-cycling capacity of reef assemblages is needed at this point.

4.54 Similar biogeochemical processes on coral-, bioeroding spongeby reef communities dominated by corals, sponges, and cyanobacteria-dominated reef communities.

410 Although processes recorded on distinct community assemblages showed some variation between substrate types, present results suggest that the various communities of this degraded reef of [Curaeao](#)[Curaçao](#) exhibit similar directions and magnitudes of key net biogeochemical processes. Results indicate that even on substrates with coral cover ranging from 34% to 36%, which is high for Curaçao reefs and relative to the wider Caribbean region (Jackson et al. 2014; de Bakker et al. 2016 and 2019), net community calcification is very low. In fact, daily rates are in a similar range to those for substrates dominated by bioeroding sponges where coral cover ranged from 1 to 9% and to substrates covered by cyanobacterial mats where no live 415 coral was recorded (coral: 0.45, BES: 0.45 and BCM: 0.33 mmol CaCO₃ m⁻² h⁻¹). Recent work by Romanó de Orte et al. (2021) found comparable results showing ~~that~~similar daytime calcification rates for live coral and dead coral substrate ~~were~~ comparable. Although hard coral is generally assumed to dominate the calcification signal on tropical reefs, these results suggest that coral might not be the sole key player in coral reef calcification dynamics on such impacted sites. Cementation/lithification processes carried out by coralline calcifying algae, micro-calcifiers (e.g., foraminifera and juvenile 420 shells) and benthic microbial communities, resulting in the trapping and binding of rubble and sediment in cryptic habitats and within/on the rubble, may play a comparably important role, counteracting some of the dissolution occurring in these communities.

425 The main differences between coral-dominated substrates and others, in terms of NCC, is that they were the only substrate able to balance out night-time dissolution. Primary production is barely compensating heterotrophic processes during the day on all substrates. Although substrates incubated in the present study are distinct in taxa dominance, they do share some similarities that may be drawing biogeochemical process differences closer to each other regardless of substrate type. For instance, turf covers any part of hard substrate available, the sand and rubble potentially harbour a variety of comparable cryptic organisms and the microbial community within and above each of these substrates may be shifting similarly towards generalist copiotrophic populations.

430 Shifts in community composition have resulted in the impairment of key reef processes and the present results may suggest that some degree of functional homogenisation (Clavel et al. 2011) exists ~~among~~among substrates with different community assemblages. It is noteworthy that seasonality may explain biogeochemical process similarity between major biogeochemical processes on reefs dominated by distinct functional groups. Roth et al. (2020) recently found that summer temperatures amplified functional differences between coral- and algae-dominated communities in the central Red Sea. Higher temperatures 435 benefit algae-dominated communities in terms of primary production and growth while coral-dominated communities shifted towards a more heterotrophic state with depressed net community calcification rates. The fact that coral-dominated substrates

Formatted: Space Before: 0 pt

studied here are already in a heterotrophic state with very low NCC values in winter temperatures attests to the differences in the studied systems and provides an opportunity for comparison between a relatively healthy system and a degraded one (Roff and Mumby, 2012). Additionally, average ambient pH at the current study site was 7.9, which is lower than average ‘summer’ 440 pH, usually between 8.1 and 8.2 (den Haan et al. 2016). This may suggest that depressed calcification rates in the Piscadera Bay are indeed linked to seasonality. However, further research and additional incubations are needed to better understand the effect of seasonal variation on the functional states of these degraded reefs.

4.5 Method considerations

Conclusions

← **Formatted:** Heading 2, Space Before: 0 pt

445 The combination of in situ incubations and inverse modelling, that takes into account which incorporates the complexity of interactions between processes, has proven to be an effective tool to provide quantitative data on the functional state of coral reef patches. Quantification of the exchange of substances between reef communities and the overlaying water was achieved despite the presence of swell-induced seawater exchange because this approach allowed for a volume exchange between the environment and the incubation. The incubations can thereby be replenished at some extent, keeping saturated O₂ levels within 450 the tent, thus minimising unrepresentative reef community metabolism. Fluxes within the incubations can be treated as if acquired by an in-situ flow through system.

For the interpretation of the measured concentration differences, the multifaceted influence of metabolic processes on chemical fluxes was accounted for. The model shows a good fit of the observations around the fitted curve, indicating that the interactions between processes and their effects on chemical fluxes were considered correctly.

455 Nonetheless, due the limited number of incubations that were carried out for this study, we interpret results with caution. Additionally, incubations were only deployed on the reef flat of one degraded reef site, future application of this or similar incubation methods should consider multiple sites. Lastly, methods would be further improved by continuous monitoring of the exchange rate, rather than assuming it to be constant throughout the incubation.

Conclusions

← **Formatted:** Heading 2, Space Before: 0 pt

460 Results acquired on this shallow Curaçaoan reef provide insight into the impact of habitat degradation and speciesbenthic composition shifts on reef functions. RemainingCurrently prevalent corals, although more resilient, calcify at a slower pace than more specialistpreviously abundant species (such as *Acropora* spp.) and cannot balance out heterotrophic processes from other functional groups. Coral presence does however contribute to counteracting dissolution processes at night, therefore acting as buffers, albeit marginal, to reef deconstruction. In the context of ongoing global change, the environmental resilience 465 of generalist species could be a determining factor of ecosystem stability (Clavel et al. 2011). For instance, on some reef terrace of the fringing reefs of Curaçao and Bonaire (southern Caribbean), certain stretches appear to harbour a considerable cover of steadily growing little boulder-constructing tolerant corals (including mainly (*Pseudo*)*Diploria* spp., *Porites astreoides* and

Siderastrea siderea) (de Bakker et al. 2019). Surprisingly, these are often found near areas which have locally suffered chronic stress from terrestrial sources (i.e., inflow, intense coastal development, factory outflow) but often limited to areas providing hard substrate and relatively little sand. Data on the processes underlying such developments however is virtually absent, but this could indicate that even the most severely degraded reefs could slowly regain essential functions when a critical adaptive capacity is reached.

While these sites may provide a spark of hope with regards to recovery potential, most of the reefs in the Caribbean presently reside in ecological states that closely resemble the reef site studied here (or are expected to do so in the near future). Ultimately, Caribbean reefs will benefit most notably from adequate mitigation strategies to give these systems a chance to adapt and restore key functions in the face of exacerbating environmental conditions.

Data and code availability Data and R code will be made available on request.

Author's contributions AW, DdB, SvH and LdN conceived the ideas and designed methodology; AW, DdB and TdC collected the data; AW and KS analysed the data; AW and LdN led the writing of the manuscript in consultation with DdB, SvH, FvD, KS and GR. All authors contributed critically to the drafts and gave final approval for publication.

Competing Interests The authors declare that they have no conflict of interest.

Acknowledgements The authors are particularly grateful to the Carmabi research station as a whole and especially Mark Vermeij for his support and facilitating the fieldwork on Curaçao. We would also like to thank Jasper de Goeij for lending us equipment. We extend our sincere gratitude to Karel Baker and Sharyn Ossebaar for analysing nutrient and DIC samples.

References

Atkinson, M. J.: Biogeochemistry of nutrients. In *Coral reefs: An ecosystem in transition*, 199-206, Springer, Dordrecht, 2011.

Albright, R., Langdon, C., and Anthony, K. R. N.: Dynamics of seawater carbonate chemistry, production, and calcification of a coral reef flat, central Great Barrier Reef, *Biogeosciences Discuss.*, 10(5), 7641-7676, 2013.

Albright, R., Benthuyzen, J., Cantin, N., Caldeira, K. and Anthony, K.: Coral reef metabolism and carbon chemistry dynamics of a coral reef flat, *Geophys. Res. Lett.*, 42(10), 3980-3988, doi: 10.1002/2015GL063488, 2015.

Albright, R., Caldeira, L., Hosfelt, J., Kwiatkowski, L., Maclare, J.K., Mason, B.M., Nebuchina, Y., Ninokawa, A., Pongratz, J., Ricke, K.L. and Rivlin, T.: Reversal of ocean acidification enhances net coral reef calcification, *Nature*, 531(7594), 362-365, 2016.

Albright, R., Takeshita, Y., Kowek, D.A., Ninokawa, A., Wolfe, K., Rivlin, T., Nebuchina, Y., Young, J. and Caldeira, K.: Carbon dioxide addition to coral reef waters suppresses net community calcification, *Nature*, 555(7697), 516-519, 2018

500 Alvarez-Filip, L., Dulvy, N. K., Gill, J. A., Côté, I. M. and Watkinson, A. R.: Flattening of Caribbean coral reefs: Region-wide declines in architectural complexity, *Proc. Royal Soc. B.*, 276(1669), 3019–3025, doi: 10.1098/rspb.2009.0339, 2009.

Andersson, A. J. and Gledhill, D.: Ocean Acidification and Coral Reefs: Effects on breakdown, dissolution, and net ecosystem calcification, *Ann Rev Mar Sci*, 5(1), 321–348, doi: 10.1146/annurev-marine-121211-172241, 2013.

Aronson, R. B., Macintyre, I. G., Lewis, S. A., and Hilbun, N. L.: Emergent zonation and geographic convergence of coral reefs, *Ecology*, 86(10), 2586-2600, 2005.

505 Bates, N. R., Amat, A. and Andersson, A. J.: Feedbacks and responses of coral calcification on the Bermuda reef system to seasonal changes in biological processes and ocean acidification, *Biogeosciences*, 2, 2509–2530, doi: 10.5194/bg-7-2509-2010, 2010.

Barott, K. L., Williams, G. J., Vermeij, M. J., Harris, J., Smith, J. E., Rohwer, F. L., and Sandin, S. A.: Natural history of 510 coral–algae competition across a gradient of human activity in the Line Islands, *Mar. Ecol. Prog. Ser.*, 460, 1-12, 2012.

Bonin, P. C., and Michotey, V. D.: Nitrogen budget in a microbial mat in the Camargue (southern France), *Mar. Ecol. Prog. Ser.*, 322, 75-84, 2006.

Bellwood, D. R., Hoey, A. S., and Choat, J. H.: Limited functional redundancy in high diversity systems: resilience and ecosystem function on coral reefs, *Ecology letters*, 6(4), 281-285, 2003.

515 Bellwood, D. R., Streit, R. P., Brandl, S. J., and Tebbett, S. B.: The meaning of the term ‘function’ in ecology: a coral reef perspective. *Functional Ecology*, 33(6), 948-961 - Wolfe, K., Anthony, K., Babcock, R., 2019.

Brandl, S. J., Rasher, D. B., Côté, I. M., Casey, J. M., Darling, E. S., Lefcheck, J. S., and Duffy, J. E.: Coral reef ecosystem functioning: eight core processes and the role of biodiversity, *Front. Ecol. Environ.*, 17(8), 445-454, 2019.

Brocke, H. J., Wenzhoefer, F., de Beer, D., Mueller, B., van Duyl, F. C. and Nugues, M. M.: High dissolved organic carbon 520 release by benthic cyanobacterial mats in a Caribbean reef ecosystem, *Sci. Rep.*, 5(1), 8852, doi: 10.1038/srep08852, 2015.

Brocke, H.J., Piltz, B., Herz, N., Abed, R.M., Palinska, K.A., John, U., den Haan, J., de Beer, D. and Nugues, M.M.: Nitrogen fixation and diversity of benthic cyanobacterial mats on coral reefs in Curaçao, *Coral Reefs*, 37(3), 861-874, 2018.

Burman, S. G., Aronson, R. B., & van Woesik, R.: Biotic homogenization of coral assemblages along the Florida reef tract, *Mar. Ecol. Prog. Ser.*, 467, 89-96, 2012.

525 Capone, D. G., Dunham, S. E., Horrigan, S. G., and Duguay, L. E.: Microbial nitrogen transformations in unconsolidated coral reef sediments, *Mar. Ecol. Prog. Ser.*, 75-88, 1992.

Chen, P. Y., Chen, C. C., Chu, L. F. and McCarl, B.: Evaluating the economic damage of climate change on global coral reefs, *Glob. Environ. Change*, 30, 12–20. doi: 10.1016/j.gloenvcha.2014.10.011, 2015.

Clavel, J., Julliard, R., and Devictor, V.: Worldwide decline of specialist species: toward a global functional homogenization?, 530 *Front. Ecol. Environ.*, 9(4), 222-228, 2011.

Clements, K. D., German, D. P., Piché, J., Tribollet, A., & Choat, J. H.: Integrating ecological roles and trophic diversification on coral reefs: multiple lines of evidence identify parrotfishes as microphages, *Biol. J. Linn. Soc.*, 120(4), 729-751, 2016.

Courtney, T.A., Andersson, A.J., Bates, N.R., Collins, A., Cyronak, T., de Putron, S.J., Eyre, B.D., Garley, R., Hochberg, E.J., Johnson, R. and Musielewicz, S.: Comparing chemistry and census-based estimates of net ecosystem calcification on a rim reef in Bermuda, *Front. Mar. Sci.*, 3, 181, 2016.

535 Cramer, K., Donovan, M., Jackson, J., Greenstein, B., Korpany, C., Cook, G., and Pandolfi, J.: The transformation of Caribbean coral communities since humans, *Authorea Preprints*, 2021.

Cyronak, T., Andersson, A. J., Langdon, C., Albright, R., Bates, R., Caldeira, K., Carlton, R., Corredor, J. E., Dunbar, R. B., Enochs, I., Erez, J., Eyre, B. D., Gattuso, J., Lantz, C., Lazar, B., Manzello, D., McMahon, A., Mele, M., Page, H. N., Santos, 540 I. R., Schulz, K. G., Shaw, E. and Silverman, J.: Taking the metabolic pulse of the world's coral reefs, *PLoS one*, 1–17, 2018.

Darling, E. S., Alvarez-Filip, L., Oliver, T. A., McClanahan, T. R., and Côté, I. M.: Evaluating life-history strategies of reef corals from species traits, *Ecol. Lett.*, 15(12), 1378–1386, 2012.

De'ath, G., Fabricius, K. E., Sweatman, H. and Puotinen, M.: The 27 – year decline of coral cover on the Great Barrier Reef and its causes, *Proc. Natl. Acad. Sci. U. S. A.*, doi: 10.1073/pnas.1208909109, 2012.

545 de Bakker, D. M., van Duyl, F. C., Perry, C. T., and Meesters, E. H.: Extreme spatial heterogeneity in carbonate accretion potential on a Caribbean fringing reef linked to local human disturbance gradients, *Glob. Change Biol.*, 25(12), 4092–4104, 2019.

de Bakker, D. M., Webb, A. E., van den Bogaart, L. A., van Heuven, S. M., Meesters, E. H., and van Duyl, F. C.: Quantification of chemical and mechanical bioerosion rates of six Caribbean excavating sponge species found on the coral reefs of Curaçao, 550 *PLoS one*, 13(5), e0197824, 2018.

de Bakker, D. M., van Duyl, F. C., Bak, R. P. M., Nugues, M. M., Nieuwland, G. and Meesters, E. H.: 40 Years of benthic community change on the Caribbean reefs of Curaçao and Bonaire: the rise of slimy cyanobacterial mats, *Coral Reefs*, Springer Berlin Heidelberg, 36(2), 355–367, doi: 10.1007/s00338-016-1534-9, 2017.

de Bakker, D. M., Meesters, E. H., Bak, R. P. M., Nieuwland, G. and Van Duyl, F. C.: Long-term Shifts in Coral Communities 555 On Shallow to Deep Reef Slopes of Curaçao and Bonaire: Are There Any Winners?, *Front. Mar. Sci.*, 1–14, doi: 10.3389/fmars.2016.00247, 2016.

den Haan, J., Huisman, J., Brocke, H.J., Goehlich, H., Latijnhouwers, K.R., Van Heeringen, S., Honcoop, S.A., Bleyenberg, T.E., Schouten, S., Cerli, C. and Hoitinga, L.: Nitrogen and phosphorus uptake rates of different species from a coral reef community after a nutrient pulse, *Sci. Rep.*, 6(1), 1–13, 2016.

560 Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO₂ measurements, North Pacific Marine Science Organization, 2007.

Dinsdale, E. A., and Rohwer, F.: Fish or germs? Microbial dynamics associated with changing trophic structures on coral reefs. In *Coral reefs: an ecosystem in transition*, 231–240, Springer, Dordrecht, 2011.

Dove, S.G., Kline, D.I., Pantos, O., Angly, F.E., Tyson, G.W. and Hoegh-Guldberg, O.: Future reef decalcification under a 565 business-as-usual CO₂ emission scenario, *PNAS*, 110(38), 15342–15347, 2013.

Dove, S.G., Brown, K.T., Van Den Heuvel, A., Chai, A. and Hoegh-Guldberg, O.: Ocean warming and acidification uncouple calcification from calcifier biomass which accelerates coral reef decline, *Comms earth*, 1(1), 1-9, 2020.

Eyre, B. D., Santos, I. R., and Maher, D. T.: Seasonal, daily and diel N2 effluxes in permeable carbonate sediments, *Biogeosciences*, 10(4), 2601-2615, 2013.

570 Gardner, T. A., Côté, I. M., Gill, J. A., Grant, A. and Watkinson, A. R.: Long-term Regional-wide declining in Caribbean corals, *Science*, 301(5635), 958-960, doi: 10.1126/science.1086050, 2003.

Haas, A.F., Nelson, C.E., Kelly, L.W., Carlson, C.A., Rohwer, F., Leichter, J.J., Wyatt, A. and Smith, J.E.: Effects of coral reef benthic primary producers on dissolved organic carbon and microbial activity, *PLoS one*, 6(11), e27973, 2011.

Haas, A. F., Nelson, C. E., Rohwer, F., Wegley-Kelly, L., Quistad, S. D., Carlson, C. A., Leichter, J. J., Hatay, M. and Smith, 575 J. E.: Influence of coral and algal exudates on microbially mediated reef metabolism, *PeerJ*, 1, e108, doi: 10.7717/peerj.108, 2013.

Haas, A.F., Fairoz, M.F., Kelly, L.W., Nelson, C.E., Dinsdale, E.A., Edwards, R.A., Giles, S., Hatay, M., Hisakawa, N., Knowles, B. and Lim, Y.W.: Global microbialization of coral reefs, *Nat. Microbiol.*, 1(6), 1-7, 2016.

Holm, S.: A simple sequentially rejective multiple test procedure, *Scand. J. Stat.*, 65-70, 1979.

580 Hörmlein, C., Confurius-Guns, V., Stal, L. J., and Bolhuis, H.: Daily rhythmicity in coastal microbial mats, *NPJ Biofilms Microbiomes*, 4(1), 1-11, 2018.

Hughes, T. P.: Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef, *Science*, 265(5178), 1547-1551, 1994.

Hughes, T.P., Barnes, M.L., Bellwood, D.R., Cinner, J.E., Cumming, G.S., Jackson, J.B., Kleypas, J., Van De Leemput, I.A., 585 Lough, J.M., Morrison, T.H. and Palumbi, S.R., Coral reefs in the Anthropocene, *Nature*, 546(7656), 82-90, 2017.

Hydes, D. J., Aoyama, M., Aminot, A., Bakker, K., Becker, S., Coverly, S., Daniel, A., Dickson, A. G., Gross, O., Kerouel, R., Van Ooijen, J., Sato, K., Tanhua, T., Woodward, M. and Zhang, J. Z.: Determination of dissolved nutrients (N, P, Si) in seawater with high precision and inter-comparability using gas-segmented continuous flow analysers, *The GO-SHIP Repeat Hydrography Manual: A Collection of Expert Reports and Guidelines*, 1-87, available at: <http://archimer.ifremer.fr/doc/00020/13141/>, 2010.

Jackson, J. B. C., Donovan, M. K., Cramer, K. L. and Lam, V.: Status and Trends of Caribbean Coral Reefs: 1970-2012, GCRMN, IUCN, Gland, Switzerland, 306, doi: 10.1242/jeb.061267, 2014.

Karcher, D.B., Roth, F., Carvalho, S., El-Khaled, Y.C., Tilstra, A., Kürten, B., Struck, U., Jones, B.H. and Wild, C.: Nitrogen eutrophication particularly promotes turf algae in coral reefs of the central Red Sea, *PeerJ*, 8, e8737, 2020.

595 Kelly, L.W., Nelson, C.E., Haas, A.F., Naliboff, D.S., Calhoun, S., Carlson, C.A., Edwards, R.A., Fox, M.D., Hatay, M., Johnson, M.D. and Kelly, E.L.: Diel population and functional synchrony of microbial communities on coral reefs, *Nat. Commun.*, 10(1), 1-9, 2019.

Kline, D.I., Teneva, L., Schneider, K., Miard, T., Chai, A., Marker, M., Headley, K., Opdyke, B., Nash, M., Valetich, M. and Caves, J.K.: A short-term in situ CO2 enrichment experiment on Heron Island (GBR), *Sci. Rep.*, 2(1), 1-9, 2012.

600 Koop, K., Booth, D., Broadbent, A., Brodie, J., Bucher, D., Capone, D., Coll, J., Dennison, W., Erdmann, M., Harrison, P.,
Hoegh-Guldberg, O., Hutchings, P., Jones, G. B., Larkum, A. W. D., O'Neil, J., Steven, A., Tentori, E., Ward, S., Williamson,
J. and Yellowlees, D.: ENCORE: The effect of nutrient enrichment on coral reefs. *Synthesis of results and conclusions*, Mar.
Pollut. Bull, 42(2), 91–120, doi: 10.1016/S0025-326X(00)00181-8, 2001.

605 Kwon, G., Le, L. T., Jeon, J., Noh, J., Jang, Y., Kang, D., and Jahng, D.: Effects of light and mass ratio of microalgae and
nitrifiers on the rates of ammonia oxidation and nitrate production, *Biochem. Eng. J.*, 107656, 2020.

Koeweek, D., Dunbar, R. B., Price, N., Mucciarone, D. and Teneva, L.: Environmental and ecological controls of coral
community metabolism on Palmyra Atoll, *Coral Reefs*, 339–351, doi: 10.1007/s00338-014-1217-3, 2015.

Langdon, C. and Atkinson, M. J.: Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with
seasonal change in temperature/ irradiance and nutrient enrichment, *J. Geophys. Res. Oceans*, 110(9), 1–16, doi:
610 10.1029/2004JC002576, 2005.

Lavigne, H., Proye, A., and Gattuso, J. P.: Package seacarb, Laboratoire d'Océanographie de Villefranche (LOV) France,
2009.

Liu, X., Byrne, R. H., Lindemuth, M., Easley, R. and Mathis, J. T.: An automated procedure for laboratory and shipboard
spectrophotometric measurements of seawater alkalinity: Continuously monitored single-step acid additions, *Mar. Chem.*, 174,
615 141–146, doi: 10.1016/j.marchem.2015.06.008, 2015.

Matsuzaki, S. I. S., Sasaki, T., & Akasaka, M.: Consequences of the introduction of exotic and translocated species and future
extirpations on the functional diversity of freshwater fish assemblages, *Glob. Ecol. Biogeogr.*, 22(9), 1071-1082, 2013.

McGill, B. J., Dornelas, M., Gotelli, N. J., & Magurran, A. E.: Fifteen forms of biodiversity trend in the Anthropocene, *Trends
Ecol. Evol.*, 30(2), 104-113, 2015.

620 McGillis WR, Langdon C, Loose B, Yates KK, Corredor J.: Productivity of a coral reef using boundary layer and enclosure
methods, *Geophys. Res. Lett.*, 38(3):1-5, 2011.

McWilliam, M., Hoogenboom, M. O., Baird, A. H., Kuo, C. Y., Madin, J. S., & Hughes, T. P.: Biogeographical disparity in
the functional diversity and redundancy of corals, *Proc. Natl. Acad. Sci. U. S. A.*, 115(12), 3084-3089, 2018.

Moberg, F., and Folke, C.: Ecological goods and services of coral reef ecosystems, *Ecol Econ.*, 29(2), 215-233, 1999.

625 Muehllehner, N., Langdon, C., Venti, A. and Kadko, D.: Dynamics of carbonate chemistry, production, and calcification of
the Florida Reef Tract (2009–2010): Evidence for seasonal dissolution, *Global Biogeochem Cycles*, 661–688, doi:
10.1002/2015GB005327, 2016.

Mueller, B., Goeij, J. M. De, Vermeij, M. J. A., Mulders, Y., Ent, E. Van Der, Ribes, M. and van Duyl, F. C.: Natural Diet of
Coral-Excavating Sponges Consists Mainly of Dissolved Organic Carbon (DOC), *PloS one*, 9(2), doi:
630 10.1371/journal.pone.0090152, 2014.

Mueller, B., Den Haan, J., Visser, P. M., Vermeij, M. J., and Van Duyl, F. C.: Effect of light and nutrient availability on the
release of dissolved organic carbon (DOC) by Caribbean turf algae, *Sci. Rep.*, 6(1), 1-9, 2016.

Nelson, C. E., Goldberg, S. J., Kelly, L. W., Haas, A. F., Smith, J. E., Rohwer, F., and Carlson, C. A.: Coral and macroalgal exudates vary in neutral sugar composition and differentially enrich reef bacterioplankton lineages, *ISME J*, 7(5), 962-979, 2013.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M. H. H., Oksanen, M. J., and Suggs, M. A. S. S.: The vegan package, *Community ecology package*, 10(631-637), 719, 2007.

Oliver, T.H., Heard, M.S., Isaac, N.J., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C.D.L., Petchey, O.L. and Proença, V.: Biodiversity and resilience of ecosystem functions , *Trends Ecol. Evol.*, 30(11), pp.673-684, 2015.

O'Neil, J. M., and Capone, D. G.: Nitrogen cycling in coral reef environments, *Nitrogen in the marine environment*, Eds.: Capone, D.G., Bronk, D.A., Mulholland, M.R., and Carpenter, E.J., 949-989, 2008.

Perry, C. T., Murphy, G. N., Kench, P. S., Smithers, S. G., Edinger, E. N., Steneck, R. S., and Mumby, P. J.: Caribbean-wide decline in carbonate production threatens coral reef growth, *Nat. Commun*, 4, 1402, 2013.

Petsch, D.K., Saito, V.S., Landeiro, V.L., Silva, T.S., Bini, L.M., Heino, J., Sozin, J., Tolonen, K.T., Jyrkäkallio-Mikkola, J., Pajunen, V. and Siqueira, T.: Beta diversity of stream insects differs between boreal and subtropical regions, but land use does not generally cause biotic homogenization, *Freshw. Sci.*, 40(1), 000-000, 2020.

Pfister, C. A., and Altabet, M. A.: Enhanced microbial nitrogen transformations in association with macrobiota from the rocky intertidal, *Biogeosciences*, 16(2), 193-206, 2019.

Rädecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., and Wild, C.: Nitrogen cycling in corals: the key to understanding holobiont functioning?, *Trends Microbiol.*, 23(8), 490-497, 2015.

Roff, G. and Mumby, P. J.: Global disparity in the resilience of coral reefs, *Trends Ecol. Evol.*, Elsevier Ltd, 27(7), 404-413, doi: 10.1016/j.tree.2012.04.007, 2012.

Romanó de Orte, M., Kowek, D.A., Cyronak, T., Takeshita, Y., Griffin, A., Wolfe, K., Szmant, A., Whitehead, R., Albright, R. and Caldeira, K.: Unexpected role of communities colonizing dead coral substrate in the calcification of coral reefs, *Limnol. Oceanogr.*, 66, pp.1793-1803, 2021.

Roth, F., Rädecker, N., Carvalho, S., Duarte, C.M., Saderne, V., Anton, A., Silva, L., Calleja, M.L., Morán, X.A.G., Voolstra, C.R. and Kürten, B.: High summer temperatures amplify functional differences between coral-and algae-dominated reef communities, *Ecology*, p.e03226, 2020.

Roth, F., El-Khaled, Y.C., Karcher, D.B., Rädecker, N., Carvalho, S., Duarte, C.M., Silva, L., Calleja, M.L., Morán, X.A.G., Jones, B.H. and Voolstra, C.R.: Nutrient pollution enhances productivity and framework dissolution in algae-but not in coral-dominated reef communities, *Mar. Pollut. Bull.*, 168, 112444, 2021.

Scheffers, S. R., Nieuwland, G., Bak, R. P. M., and Van Duyl, F. C.: Removal of bacteria and nutrient dynamics within the coral reef framework of Curaçao (Netherlands Antilles), *Coral Reefs*, 23(3), 413-422, 2004.

Shaw, E. C., Phinn, S. R., Tilbrook, B. and Steven, A.: Natural in situ relationships suggest coral reef calcium carbonate production will decline with ocean acidification. *Limnol. Oceanogr.*, 60(3), 777-788. doi: 10.1002/lno.10048, 2015.

Silverman, J., Lazar, B. and Erez, J.: Effect of aragonite saturation, temperature, and nutrients on the community calcification rate of a coral reef, *J. Geophys. Res. Oceans*, 112(5), 1–14, doi: 10.1029/2006JC003770, 2007.

Soetaert, K., Hofmann, A. F., Middelburg, J. J., Meysman, F. J., and Greenwood, J.: The effect of biogeochemical processes on pH, *Mar. Chem.*, 106(1-2), 380-401, 2007.

670 Soetaert, K. E., Petzoldt, T., and Setzer, R. W.: Solving differential equations in R: package deSolve, *J. Stat. Softw.*, 33, 2010.

Soetaert, K., and Petzoldt, T.: Inverse modelling, sensitivity and Monte Carlo analysis in R using package FME, *J. Stat. Softw.*, 33(3), 1-28, 2010.

Sonnier, G., Johnson, S. E., Amatangelo, K. L., Rogers, D. A., & Waller, D. M.: Is taxonomic homogenization linked to 675 functional homogenization in temperate forests?, *Glob. Ecol. Biogeogr.*, 23(8), 894-902, 2014.

Takeshita, Y., McGillis, W., Briggs, E.M., Carter, A.L., Donham, E.M., Martz, T.R., Price, N.N. and Smith, J.E.: Assessment of net community production and calcification of a coral reef using a boundary layer approach, *J. Geophys. Res. Oceans*, 121(8), 5655-5671, 2016.

Tobias, N., & Monika, W.: Does taxonomic homogenization imply functional homogenization in temperate forest herb layer 680 communities?, *Plant Ecol.*, 213(3), 431-443, 2012.

van Duyl, F. C.: Atlas of the living reefs of Curaçao and Bonaire (Netherlands Antilles), Utrecht, Netherlands Foundation for Scientific Research in Surinam and the Netherlands Antilles, 117, 1985.

van Heuven, S. M. A. C., Webb, A. E., Bakker, D. M. De, Meesters, E., Duyl, F. C. Van, Reichart, G. and Lennart, J.: In-situ 685 incubation of a coral patch for community-scale assessment of metabolic and chemical processes on a reef slope, *PeerJ*, 1–24, doi: 10.7717/peerj.5966, 2018.

Vermeij, M. J., Van Moorselaar, I., Engelhard, S., Hörlein, C., Vonk, S. M., and Visser, P. M.: The effects of nutrient enrichment and herbivore abundance on the ability of turf algae to overgrow coral in the Caribbean, *PloS one*, 5(12), e14312, 2010.

Webb, K. L., and Wiebe, W. J.: Nitrification on a coral reef, *Can. J. Microbiol.*, 21(9), 1427-1431, 1975.

690 Webb, A. E., van Heuven, S. M., de Bakker, D. M., van Duyl, F. C., Reichart, G. J., and de Nooijer, L. J.: Combined effects of experimental acidification and eutrophication on reef sponge bioerosion rates, *Front. Mar. Sci.*, 4, 311, 2017.

Weber, L., and Apprill, A.: Diel, daily, and spatial variation of coral reef seawater microbial communities, *PloS one*, 15(3), e0229442, 2020.

White, H. J., Montgomery, W. I., Storchová, L., Hořák, D., and Lennon, J. J.: Does functional homogenization accompany 695 taxonomic homogenization of British birds and how do biotic factors and climate affect these processes?, *Ecol. Evol.*, 8(15), 7365-7377, 2018.

Wolfe, K., Anthony, K., Babcock, R.C., Bay, L., Bourne, D.G., Burrows, D., Byrne, M., Deaker, D.J., Diaz-Pulido, G., Frade, P.R. and Gonzalez-Rivero, M.: Priority species to support the functional integrity of coral reefs. *Oceanogr. Mar. Biol.*, 2020

Yates, K. K. and Halley, R. B.: Measuring coral reef community metabolism using new benthic chamber technology, *Coral 700 Reefs*, 22(3), 247–255, doi: 10.1007/s00338-003-0314-5, 2003.






Figure 1: The tent incubation setup used. (a) Photograph depicting the tent incubation during the experiments. (b) Schematic cross section of the employed setup for enclosing a small patch of reef. A battery powered mixing propeller for maintaining water circulation, and analysers for salinity (S), temperature (T), oxygen (O_2), and light (PAR) are located inside the tent. Outside the enclosure another S/T and PAR analyser were placed, as well as the battery for the pump. Sampling of exterior and interior water (through sampling tube) was performed by divers using large volume syringes. Zippers allow for opening of tent windows for re-equilibrating the interior to the exterior conditions between incubations.



705

710

Figure 2: Benthic cover of the 15 incubated reef communities dominated either by coral, turf and macroalgae (TMA), bioeroding sponges (BES), cyanobacterial mats (BCM) or sand with exemplary photograph.



715 Figure 3: Exterior and interior measurements for three representative incubations performed throughout this study. (a) Difference in light between the inside of the tent and the ambient environment during an incubation. (b) Injection of salt within the tent at the start of the incubation and its gradual return to ambient salinity. (c) Temperature within the tent compared to exterior conditions, when the tent is opened at the end of the incubation, temperature immediately returns to ambient conditions.






Figure 4: (a, b) Best fit explaining in-tent salinity at any given time (equation 1, red line). In-tent salinity measurements are in black and ambient salinity measurements are in blue. (a) illustrates an incubation that leaked relatively slowly with $K = 0.0192$ ($0.0192 \times 108 = 2.1$ litres per minute), while (b) depicts a more rapidly leaking tent with $K = 0.044$ ($0.044 \times 108 = 4.8$ litres per minute).



725

Figure 5: Illustrative results of the model (coloured lines) employed to infer the process rates from measured data. The top graphs depict the model output of incubations performed during the day while the bottom ones show the output for night-time incubations. Points represent measured values of the state variables inside and outside the tent. The measured data was centred for graphic visualisation purposes (non-centred data for all incubations can be found in supplementary Table S2). The blue, green, orange, red and yellow colours represent communities dominated by coral, turf and macroalgae, bioeroding sponges, benthic cyanobacterial mats and sand respectively. Note that NH_4 and NO_3 measurements in BCM incubations at night-time were much higher than the rest. The y-axis of the graphs depicting NH_4 and NO_3 results were therefore truncated in order to better visualise model output of all other incubations (model output for the off chart BCM incubation can be found in Figure S1).

730



735

Figure 6: Process rates for all tent replicates (points) and respective average (line) estimated from observed concentration changes and model output in the tent enclosure on each substrate type. Processes occurring during the day are depicted in red, and processes at night are represented in purple.




Figure 7: A) Principal component analysis (PCA) diagram displaying the spatial variation of all incubations along the first two principal components. Processes are plotted as vectors and dots represent day-time incubations while triangles depict night-time incubations. Colour refers to the type of substrate incubated. B) NCP vs NCC rates for every incubation. Each dot/triangle represents an individual incubation and colour indicates community composition.