

1 **Reviews & Syntheses: Arctic Fire Regimes and Emissions in the 21st** 2 **Century**

3 Jessica L. McCarty¹, Juha Aalto^{2,3}, Ville-Veikko Paunu⁴, Steve R. Arnold⁵, Sabine Eckhardt⁶, Zbigniew
4 Klimont⁷, Justin J. Fain¹, Nikolaos Evangeliou⁶, Ari Venäläinen², Nadezhda M. Tchebakova⁸, Elena I.
5 Parfenova⁸, Kaarle Kupiainen⁹, Amber J. Soja^{10,11}, Lin Huang¹², and Simon Wilson¹³

6 ¹Department of Geography and Geospatial Analysis Center, Miami University, Oxford, OH, USA

7 ²Finnish Meteorological Institute, Helsinki, Finland

8 ³Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

9 ⁴Finnish Environment Institute (SYKE), Helsinki, Finland

10 ⁵Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, United
11 Kingdom

12 ⁶Norwegian Institute for Air Research, Kjeller, Norway

13 ⁷International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

14 ⁸V. N. Sukachev Institute of Forests, Siberian Branch, Russian Academy of Sciences, Krasnoyarsk, Russian Federation

15 ⁹Ministry of the Environment Finland, Aleksanterinkatu 7, Helsinki, P.O.Box 35, FI-00023 Government, Finland

16 ¹⁰National Institute of Aerospace, Hampton, VA, USA

17 ¹¹Langley Research Center, Hampton, VA, USA

18 ¹²Climate Research Division, ASTD/STB, Environment and Climate Change Canada, Toronto, Canada

19 ¹³Arctic Monitoring and Assessment Programme (AMAP) Secretariat, Tromsø, Norway

20
21 *Correspondence to:* Jessica L. McCarty (jmccarty@miamioh.edu)

22 **Abstract.** In recent years, the Pan-Arctic region has experienced increasingly extreme fire seasons. Fires in the northern high
23 latitudes are driven by current and future climate change, lightning, fuel conditions, and human activity. In this context,
24 conceptualizing and parameterizing current and future Arctic fire regimes will be important for fire and land management as
25 well as understanding current and predicting future fire emissions. The objectives of this review were driven by policy
26 questions identified by the Arctic Monitoring and Assessment Programme (AMAP) Working Group and posed to its Expert
27 Group on Short-Lived Climate Forcers. This review synthesises current understanding of the changing Arctic and boreal fire
28 regimes, particularly as fire activity and its response to future climate change in the Pan-Arctic has consequences for Arctic
29 Council states aiming to mitigate and adapt to climate change in the north. The conclusions from our synthesis are the
30 following: (1) Current and future Arctic fires, and the adjacent boreal region, are driven by natural (i.e., lightning) and human-
31 caused ignition sources, including fires caused by timber and energy extraction, prescribed burning for landscape management,
32 and tourism activities. Little is published in the scientific literature about cultural burning by Indigenous populations across
33 the Pan-Arctic and questions remain on the source of ignitions above 70°N in Arctic Russia. (2) Climate change is expected
34 to make Arctic fires more likely by increasing the likelihood of extreme fire weather, increased lightning activity, and drier

35 vegetative and ground fuel conditions. (3) To some extent, shifting agricultural land use, forest-steppe to steppe, tundra-to-
36 taiga, and coniferous-to-deciduous forest transitions in a warmer climate may increase and decrease open biomass burning,
37 [depending on land use in addition to climate-driven biome shifts](#). However, at the country- and landscape-scales, these
38 relationships are not well established. (4) Current black carbon and PM_{2.5} emissions from wildfires above 50°N and 65°N are
39 larger than emissions from the anthropogenic sectors of residential combustion, transportation, and flaring, respectively.
40 Wildfire emissions have increased from 2010 to 2020, particularly above 60°N, with 56% of black carbon emissions above
41 65°N in 2020 attributed to open biomass burning - indicating how extreme the 2020 wildfire season was and [how severe](#) future
42 Arctic wildfire seasons [can potentially be](#). (5) What works in the boreal zones to prevent and fight wildfires may not work in
43 the Arctic. Fire management will need to adapt to a changing climate, economic development, the Indigenous and local
44 communities, and fragile northern ecosystems, including permafrost and peatlands. (6) Factors contributing to the uncertainty
45 of predicting and quantifying future Arctic fire regimes include underestimation of Arctic fires by satellite systems, lack of
46 agreement between Earth observations and official statistics, and still needed refinements of location, conditions, and previous
47 fire return intervals on peat and permafrost landscapes. This review highlights that much research is needed in order to
48 understand the local and regional impacts of the changing Arctic fire regime on emissions and the global climate, ecosystems
49 and Pan-Arctic communities.

50 **1 Introduction**

51 For more than a decade, climate [modeling/modelling](#) studies have projected an “invasion” of fires to the Arctic regions
52 (Krawchuk et al., 2009). In this paper, we review the current understanding of the changing Arctic fire regime, and its impacts
53 on fires emissions, [providing a foundation for future systemic Pan-Arctic fire and fire emissions analyses and coordination in](#)
54 [the context of the Arctic Council Members, Permanent Participants, Observers, and Working Groups. This review paper is](#)
55 [also the first to link emissions with a changing fire regime for the Pan-Arctic](#). Previous published reviews on fires in the high
56 northern latitudes have linked increasing fire activity in the Arctic and the Boreal region to climate-driven warming and drying
57 (Hu et al., 2015; Walsh et al., 2020). While fires in the Arctic, defined as latitudes above 66°N by the Arctic Monitoring and
58 Assessment Programme (AMAP) definition (AMAP, 1998), are not new (Wein, 1976), a consensus of evidence suggests that
59 tundra fires are increasing (Hu et al., 2015; Masrur et al., 2018) with a potential for novel fire regimes (Young et al., 2016).
60 Fire regimes are often defined as the main characteristics of fire activity for a given location: frequency, typical sizes of fires,
61 annual burned area, severity, seasonality, type (surface, ground, or crown fires), and ignition cause (human or natural) (Hanes
62 et al., 2019).

63
64 Over the past four decades, fire activity has increased in Alaska and the Sakha Republic of Russia but decreased slightly in the
65 Northwest Territories of Canada, indicating large spatio-temporal variability of Pan-Arctic fire dynamics (York et al., 2020).
66 Further, in the past three years, there have been large fires in Fennoscandia in 2018, Alaska and Greenland in 2019, and the

67 Russian Federation in 2020, mainly in the Boreal zone, i.e., at and above 50°N, but with expanding fires into the Arctic region
68 (Walsh et al., 2020), even reaching as far north as the Arctic Ocean in eastern Siberia (Kharuk et al., 2021). Thus, quantifying
69 the impact of climate change, human ignition sources, and biophysical parameters, such as availability and/or distribution of
70 aboveground fuels, permafrost thaw, and drying of peat, on increased fire activity in the Arctic and Boreal are needed to
71 understand the emerging Arctic fire regime (Krawchuk and Moritz, 2011). Here we define an emerging fire regime in the
72 Arctic as documented increased frequency and lengthened seasonality (~~both earlier springtime fires and later fires later in fall~~)
73 of both natural and human-caused surface and ground fires (i.e., peat) increasing total fire emissions within the Arctic (see
74 Suppl. Table 1 in the Supplement for a list of all key terms).

75
76 For this review paper, the definition of open biomass burning in the Arctic will include wildland fires (i.e., sometimes referred
77 to as and encompassing of wildfires, forest fires, peat fires, as well as prescribed fires in natural areas) and fires in human-
78 dominated landscapes (i.e., agricultural open burning, prescribed burning in agroforestry, timber, rangelands, etc.), with
79 natural fires (lightning-caused ignitions) and human-caused fires differentiated where possible using reported statistics and
80 geospatial methods. Given the strong influence of boreal systems on the Arctic in terms of fire disturbance, emissions, and
81 shifting vegetation, we have included boreal fire regimes in this review, while specifically identifying each climatic zone as
82 needed. Open biomass burning is a known disturbance in the Arctic Council region¹ (AMAP, 2011; 2015). The 2015 Arctic
83 Monitoring and Assessment Programme (AMAP) assessment on black carbon (BC) and ozone as Arctic climate forcers noted
84 key characteristics of open biomass burning in the Arctic region, including human influence on both ignition and fuels
85 management, significant interannual variation of fire events and emissions, spatial and seasonal clustering of burning related
86 to active land management, and fuel conditions (AMAP, 2015). Since 2015, evidence of direct climate change influence on
87 large, early season fires has increased (Wang et al., 2017) as well as fueling extreme wildfires at the wildland-urban interface
88 (WUI) and not just remote Boreal forests and Arctic tundra (Abatzoglou and Williams, 2016; Kirchmeier-Young et al., 2019).
89 In terms of burned area, 2015 was the largest fire year for the Alaskan tundra ecoregion (Michaelides et al., 2019).

90
91 Under future climate change, an overall increase in fires is expected in the Arctic Council region, indicating that associated
92 emissions are also likely to increase. For instance, natural fires, defined as lightning-caused fires, may increase as lightning is
93 predicted to increase (Půček et al., 2017; Veraverbeke et al. 2017; Bieniek et al., 2020), under Representative Concentration
94 Pathways (RCPs) 4.5 (stabilising emissions) and 8.5 (high emissions) developed for the Intergovernmental Panel on Climate
95 Change (IPCC) Fifth Assessment Report (AR5). Likewise, using the same scenarios, wildfire emissions of BC, CO, NO_x,
96 PM_{2.5}, and SO₂ could exceed anthropogenic emissions in northeastern Europe, including Sweden and Finland, by 2090 (Knorr

¹ The Arctic Council membership comprises the eight member states: Canada, the Kingdom of Denmark, Finland, Iceland, Norway, the Russian Federation, Sweden, and the United States of America, as well six permanent participants representing Arctic Indigenous peoples, including the Aleut International Association, the Arctic Athabaskan Council, the Gwich'in Council International, the Inuit Circumpolar Council, the Russian Association of Indigenous Peoples of the North, and the Saami Council.

97 et al., 2016). There is a clear consensus that the emerging Arctic fire regime will be marked by shifts in fire seasons, i.e.,
98 likelihood of extreme fires later in the growing season, will occur in the Boreal forests of eastern Canada (Boulanger et al.,
99 2013); central and northwestern Canada (Boulanger et al., 2014); and European Russia, West Siberia, and the Far East
100 (Sherstyukov and Sherstyukov, 2014). By the end of the 21st century under RCP6.0 (stabilising emissions with higher CO₂
101 equivalency than RCP4.5), the annual chance of large tundra fire in Alaska will be almost one in four, i.e., a range of 13-23%
102 predicted increases (Hu et al., 2015). Moreover, Wang et al. (2017) noted that a recent lengthening in the fire season in Canada
103 has led to the increase in the total number of fire spread days, leading to large increases in total fire size and emissions for
104 early season fires like the Fort McMurray megafire in Alberta. Lengthening the fire season, a component of the emerging
105 Arctic fire regimes, means increased potential for more and larger fire emissions throughout the fire season, starting earlier in
106 spring and lasting later into autumn.

107
108 For the past two decades, it has been well established that understanding fire regimes improves emission estimates from fires
109 in high northern latitudes (Conard and Ivanova, 1997; Soja et al., 2004a) and may even be necessary for creating emission
110 models (van der Werf et al., 2010). Further, climate change is expected to alter fire regimes and likely increase emissions
111 (Sommers et al., 2014). For that reason, this review also includes emission estimates from adjacent boreal fires as well as
112 temperate fire sources known to impact the Arctic region via increased atmospheric abundance and deposition of black carbon
113 as well as greenhouse gas emissions. This review paper spawned from policy questions (Suppl. Table 2) that the Expert Group
114 on Short-Lived Climate Forcers (SLCF EG) of AMAP, a Working Group of the Arctic Council, was asked to answer for its
115 2021 Assessment Impacts of short-lived climate forcers on Arctic climate, air quality, and human health (AMAP, 2021). Our
116 specific objectives are to:

- 117 (1) identify and review the key drivers of the Arctic fires today and in the future to characterise an emerging Arctic fire regime,
118 with potential changes (paper sections 2-3 and policy question 1 in Suppl. Table 2);
- 119 (2) characterise fire emissions from ground- and satellite-based data sources in the Arctic, boreal, and temperate regions that
120 impact the Arctic (paper section 4 and policy questions 1,3-5 in Suppl. Table 2);
- 121 (3) contextualise emissions from the Arctic fire regime with other sectoral sources for the Pan-Arctic (paper section 5 and
122 policy questions 5-6 in Suppl. Table 2);
- 123 and (4) identify key challenges and research questions that could improve understanding, monitoring, and management of
124 Arctic fires in the 21st century (paper sections 6-8 and policy questions 2 and 6 in Suppl. Table 2). We
125
126 Our focus is on SLCF emissions but note that wildfires are also a source of CO₂ and other contaminants of environmental and
127 human health concern in the Arctic, including mercury and polycyclic aromatic hydrocarbons (PAH).

Formatted: Subscript

2 Drivers of Arctic fire regimes

For this review paper, the definition of open biomass burning in the Arctic will include wildland fires (i.e., wildfires, forest fires, peat fires) and fires in human-dominated landscapes (i.e., agricultural open burning, prescribed burning). Given the strong influence of boreal systems on the Arctic in terms of fire disturbance, emissions, and shifting vegetation, we have included boreal fire regimes in this review, while specifically identifying each climatic zone as needed. Broadly speaking, wildfires are driven by climate and weather conditions influencing flammability, fuels, and fuel conditions ~~Broadly speaking, wildfires are driven by climate and weather, fuels and fuel conditions, and ignition sources~~ (Silva and Harrison, 2010; de Groot et al., 2013). Ignition from lightning strikes, fire weather (i.e., temperature, humidity, precipitation, and wind), and fuel abundance (build-up) and conditions (moisture) are the typical controlling processes for 'natural' fires, i.e. fires not caused directly by human activity. Human-caused fires are driven by fuels management to reduce fire risk, land management in agricultural and timber landscapes, ~~and cultural practices~~, and accidental (Granström & Niklasson, 2008; Bowman et al., 2020). ~~Pollen-based reconstructions show prehistoric and early historic human settlements increased during wetter climates in Minusinsk Hollow in south-central Siberia, where grain and pasture yields increased twofold, rather than dry periods that favoured pastoralist (Blyakharchuk et al., 2014), highlighting the dependence of human-dominated landscapes and fire on climate.~~

Historically, both climate and humans have influenced fire activity the Pan-Arctic region. Paleofire meta analysis of boreal biomass burning during the Holocene (4,000 to 200 years BP) for the boreal zone of North America and Fennoscandia show general trends in boreal biomass burning were primarily controlled by climatic changes, mainly mean annual precipitation in Alaska, northern Quebec, and northern Fennoscandia and summer temperatures in central Canada and central Fennoscandia (Molinari et al., 2018). Boreal needleleaf evergreen fuel composition at the landscape-level across Alaska and central and southern Fennoscandia was secondary to climatic controls. These paleofire results align with recent findings by Walker et al. (2020), showing fine-scale drainage conditions, overstory tree species composition, and fuel accumulation rates across 417 sites in boreal and taiga ecoregions of northwestern Canada and Alaska were more important than incidental fire weather in terms of fire severity and subsequent carbon emissions. ~~Pollen-based reconstructions show prehistoric and early historic human settlements increased during wetter climates in Minusinsk Hollow in south-central Siberia, where grain and pasture yields increased twofold, rather than dry periods that favoured pastoralist (Blyakharchuk et al., 2014), highlighting the connections between fire, climate, and human-dominated landscapes.~~

~~Reported statistics and geospatial methods from Earth observations were used to quantify and differentiate open biomass burning as human-caused fires, i.e., agricultural open burning, timber and agroforestry, and natural fires, i.e., lightning-caused fires. Fire risk, or the likelihood of a fire occurring, in the Arctic region is often driven by climate and fire weather, fuel type, availability, and condition, and presence of people as ignition sources (more in Suppl. Table 1).~~

161 Ignition from lightning strikes, fire weather (i.e. temperature, humidity and wind), and fuel conditions (moisture) are the typical
162 controlling processes for 'natural' fires, i.e. fires not caused directly by human activity. End of century modeled fire climate
163 interactions under RCP6.0 for Alaska showed summer temperatures and annual precipitation are the most important climatic
164 factors driving the likelihood of new wildland fire regimes in tundra and the boreal forest-tundra boundary (Young et al.,
165 2016). Burned area is predicted to increase 40 to 50% in the high latitudes under climate forcing scenario 8.5 given modeled
166 changes in fuel loads, fuel moisture, and increased lightning frequency (Krause et al., 2014). Increased convective cloud
167 formation has been documented in the Russian Arctic (Chernokulsky and Esau, 2019) and the North American boreal forest
168 (Veraverbeke et al., 2017), with a 5% increase in convective storms in Northern Europe projected by the end of the 21st century
169 under RCPs 4.5 and 8.5 (Púčik et al., 2017). In general, lightning frequency is expected to increase over areas north of 50°N.
170 The strongest projected relative increase is approximately 100% across northern Europe under RCP 8.5 scenario by the end of
171 the century (Groenemeijer et al., 2016). Moreover, since summers are expected to become drier in the future (Venäläinen et
172 al., 2020), the role of lightning as an ignition source for wildfires may increase for northern Europe. These future models agree
173 with observations of past natural fires in the Arctic region. Paleofire meta-analysis of boreal biomass burning during the
174 Holocene (4,000 to 200 years BP) for the boreal zone of North America and Fennoscandia show general trends in boreal
175 biomass burning were primarily controlled by climatic changes, mainly mean annual precipitation in Alaska, northern Quebec,
176 and northern Fennoscandia and summer temperatures in central Canada and central Fennoscandia (Molinari et al., 2018).
177 Boreal needleleaf evergreen fuel composition at the landscape level across Alaska and central and southern Fennoscandia was
178 secondary to climatic controls. These paleofire results align with recent findings by Walker et al. (2020), showing fine-scale
179 drainage conditions, overstory tree species composition, and fuel accumulation rates across 417 sites in boreal and taiga
180 ecoregions of northwestern Canada and Alaska were more important than incidental fire weather in terms of fire severity and
181 subsequent carbon emissions.

182
183 Open biomass burning from anthropogenic activities like agriculture, timber, and energy extraction are expected to increase
184 in the Arctic as climate change expands human-dominated landscapes northward, increasing potential ignition sources (Fig.
185 1). The 2019 Greenland wildfire, which consumed surface vegetation and high carbon soils for nearly a month, was caused
186 when a campfire ignited dry ground near a public camping site of the world-renowned Arctic Circle Trail (McGwinn, 2019),
187 indicating that tourism will need to adapt to increased fire risk in tundra landscapes. Greenland wildfires in 2017 and 2019
188 occurred east of Sisimiut in tundra areas with low vegetative cover and degraded permafrost, but high carbon soils, during
189 warm, dry, and sunny summers (Evangelio et al., 2019). Timber extraction and site preparation, including operation of
190 machinery and vehicles on ground covered in dry wood residues, currently cause large wildfires in the Arctic Council region,
191 including the 2014 Västmanland fire in Sweden ignited by forestry vehicles during subsoiling activities (Lidskog et al., 2019),
192 which actively burned for 18 days creating a burn scar of over 14,000 ha (Pimentel and Arheimer, 2021). Northward
193 agricultural expansion will likely increase human-caused open burning as wheat and maize establish production is expected to
194 grow in previously permafrost areas of West Siberia (Parfenova et al., 2019), West Siberia is currently a minor source region

195 of agricultural burning (Hall and Loboda, 2017), with many farmers insisting that fire is necessary to clear fields under present-
196 day management and resource constraints despite bans on open agricultural burning (Theesfeld and Jelenik, 2017). This
197 northward agricultural land could expand into the cold regions of the boreal zone (Kicklighter et al., 2014; King et al.,
198 2018), and nearing the Arctic Circle for Central Siberia (Tchebakova et al., 2016). Of course, the northward agricultural
199 expansion-transitions will also be dependent on local and/or in-situ conditions limiting its expansion, such as inferior soils,
200 existing land uses not compatible with agricultural conversion, and topographic limitations (Ioffe and Nefedova, 2004; Dronin
201 and Kirilenko, 2011; Tchebakova et al., 2011). However, given the degraded conditions of most abandoned agricultural land
202 in the steppes of Siberia and high interest in northern agricultural development by neighbouring Asian countries, northward
203 development of grains and other commodity crops is expected (Prishchepov et al., 2020). Further, suppression of wildfire
204 in Canadian boreal communities has increased their likelihood of burning flammability, allowing fuels to build up in and near
205 populated places (Parisien et al., 2020), calling into question what other wildland-urban interfaces in the Arctic region may
206 have similar experience increased fire risk and fires due to long term aggressive fire suppression.

207 **3 Climate change and future Arctic fire activity**

208 **3.1 Climate change and future fires**

209 Many future fire modelling approaches use greenhouse gas emission scenarios to project the impact of climate change on
210 future temperature and precipitation - both influencing fuel ignition and subsequent burning (Veira et al., 2016). Increased fire
211 risk will not be uniform across the Pan-Arctic (Fig. 1). For instance, permafrost thaw will lead to a rewetting of soils (Wrona
212 et al., 2016), reducing above-ground and below-ground fire risk. Boike et al. (2016) showed that increasing areas of
213 thermokarst lakes were not coincident with areas of increasing fire in central Sakha Republic. Surface fires can cause
214 permafrost to thaw, producing thermokarst lakes (Jones et al., 2015), which previously have been considered to reduce fire
215 risk (Sofronov et al., 2000) but are not perfect fire breaks as wildfires can “jump” (Sofronov and Volokitina, 2010). Further,
216 changing precipitation regimes in the form of more rainfall in the Arctic for the months of March through December by end
217 of century using RCP8.5 projections (Bitanja and Andry, 2017) could both reduce fire risk through increased wetness but also
218 increase fire risk through more vegetation growth and/or shifting fuels regimes. End of century modelled fire-climate
219 interactions under RCP6.0 for Alaska showed summer temperatures and annual precipitation are the most important climatic
220 factors driving the likelihood of new wildland fire regimes in tundra and the boreal forest-tundra boundary (Young et al.,
221 2016). Burned area is predicted to increase 40 to 50% in the high latitudes under climate-forcing scenario RCP8.5 given
222 modelled changes in fuel loads, fuel moisture, and increased lightning frequency (Krause et al., 2014). Increased convective
223 cloud formation has been documented in the Russian Arctic (Chernokulsky and Esau, 2019) and the North American boreal
224 forest (Veraverbeke et al., 2017), with a 5% increase in convective storms in Northern Europe projected by the end of the 21st
225 century under RCPs 4.5 and 8.5 (Púčik et al., 2017). In general, lightning frequency is expected to increase over areas north
226 of 50°N. The strongest projected relative increase is approximately 100% across northern Europe under RCP 8.5 scenario by

Formatted: Heading 2

227 the end of the century (Groenemeijer et al., 2016). Moreover, since summers are expected to become drier in the future
228 (Venäläinen et al., 2020), the role of lightning as an ignition source for wildfires may increase for northern Europe.
229

230 Fig. 1 depicts transition themes and associated fire risks taken from the scientific literature, with general locations on the map
231 derived from the locations of these studies. These ecological and meteorological studies rely on gridded climate scenarios from
232 future greenhouse gas emission scenarios in order to predict fire risk for mid- (2050) and late-century (2100). First, as boreal
233 forests experience permafrost thaw, where initially wet soils (Wrona et al., 2016; O'Neill et al., 2020) are followed by
234 increasingly dry ground fuels (Turetsky et al., 2015; Box et al., 2019). Topography plays a crucial role in determining shifting
235 habitats, where drying will dominate on tilted surfaces and bogging will dominate on flat terrain (Tchebakova et al., 2009),
236 such that as ~~The~~ the Siberian Arctic tundra is dominated by relatively flat terrain, ~~consequently~~ bogging is predicted to prevail.
237 Second, anticipated transitions of boreal forest to deciduous forest stands would decrease fire risk in eastern Canada and small
238 regions of interior Alaska (Terrier et al., 2013; Foster et al., 2019; Mekonnen et al., 2019), as deciduous species are less
239 flammable than coniferous species (Päätao, 1998; Krawchuk et al., 2006).—Third, expansion of grassland ecosystems is
240 predicated in northwestern Canada and Alaska (Wang et al. 2019; Whitman et al., 2019) and Siberia (Tchebakova et al., 2009,
241 2016). Fourth, increased lightning strikes will ~~in turn~~ increase fire risk in Alaska (Veraverbeke et al. 2017) but also northern
242 Europe (Půček et al., 2017). Fifth, the interaction between climate-driven changes in fire regimes and permafrost will compel
243 a decrease in and a northern migration of Siberian taiga, which will result in the transition of tundra to taiga in northern Siberia
244 (Tchebakova et al., 2009, 2011; Sizov et al., 2021). Permafrost is not predicted to thaw deep enough to sustain dark-needled
245 taiga (*Pinus sibirica*, *Abies sibirica*, and *Picea obovata*), nonetheless light-needled coniferous Larix is predicted to continue
246 to dominate in eastern Siberia, maintaining a higher fire risk according to the ~~Russian~~ Russian fire hazard rankings (Melekhov, 1980).
247 The Russian fire hazard ranking systems shows a decrease in fire risk from light needle conifers (Scots pine, larch) to deciduous
248 broad-leaf tree species (birch, aspen, willow) that exist between the temperate and boreal zones, as well as along river valleys.
249 Fire risk is also lower in dark-leaf conifers (Melekhov, 1980). Fire return intervals (FRI) are consistent with Melekhov (1980),
250 with a mean FRI of 36 years (range 17-133) in light coniferous forest compared with a mean FRI of 196 years (range 75-725)
251 in dark-coniferous forest (Furyaev, 1996; Shvidenko and Nilsson, 2000; Soja et al., 2006). Larix are a fire-tolerant species,
252 and dark-coniferous species are shade-tolerant secondary-succession cohort (Shugart et al., 1992). Sixth, forest-steppe and
253 steppe is predicted to dominate over half of Siberia, largely forced by climate and increases in fire regimes (Tchebakova et
254 al., 2009). The forest-steppe that exists at the southernmost extent of the Siberian boreal forest is transitioning to steppe due
255 to: increases in extreme fires that burn the soil organic matter to mineral soil, and repeated fires and high temperatures that kill
256 regenerating seedlings. Seventh, northward agricultural expansion may increase human-caused agricultural burning as wheat
257 and maize (silage) establish in previously permafrost areas of East Siberia (Tchebakova et al., 2009; Parfenova et al., 2019),
258 expanding into the cold regions of the boreal zone (King et al., 2018) in North America as well. Finally, a threefold increase
259 of permafrost thaw in the boreal zone under RCP4.5 by 2100 is likely to increase the amount of peat fuels available for burning
260 (Nitzbon et al., 2020).

261
262 Previous work has identified the Arctic as a regional “hot spot” for interannual variability of key atmospheric constituents,
263 with wildfire being the major driver of this variability (Fisher et al., 2010; Monks et al., 2012; Voulgarakis et al., 2015). As
264 stated earlier, climate warming can cause more ignitions from lightning (Veraverbeke et al., 2017) and degraded permafrost
265 due to increasing dry ground fuels, including peat (Turetsky et al., 2015), that can and increased fire severity (Teufel and
266 Sushama, 2019). Using the RCP8.5 scenario, Teufel and Sushama (2019) estimate that a 2.0°C global threshold in temperature
267 increase, which could be reached around 2031, may cause 42% of pan-Arctic permafrost to abruptly degrade and increase fire
268 severity in Russia, Canada, and Alaska. Surface fires can cause permafrost to thaw, producing thermokarst lakes (Jones et al.,
269 2015), which previously have been considered to reduce fire risk (Sofronov et al., 2000) but are not perfect fire breaks as
270 wildfires can “jump” (Sofronov and Velokitina, 2010). By the end of the century, wildland fire risk is expected to increase,
271 with length of fire seasons - measured in terms of daily severe fire weather occurrence - predicted to expand by as much as 20
272 days for high northern latitudes using the A1B (roughly corresponding to RCP6.0), A2 (~ RCP8.5), and B1 (~RCP4.5)
273 scenarios (Flannigan et al., 2013). ~~By the end of the century, wildland fire risk is expected to increase, with length of fire~~
274 ~~seasons - measured in terms of daily severe fire weather occurrence - predicted to expand by as much as 20 days globally~~
275 ~~(Flannigan et al., 2013).~~ Similarly, Sherstyukov and Sherstyukov (2014) predict an increase of > 50 days of high fire risk days
276 by 2100 for Russia under RCP 8.5 scenario, with a potential to double annual forest fire burned area. Using CMIP5 model
277 intercomparisons, Lehtonen et al. (2016) found that large (> 0.1 km) boreal forest fires in Finland may double or even triple
278 by the end of century, using RCP4.5 and RCP 8.5 scenarios, but with large inter-model variability. ~~Using CMIP5 model~~
279 ~~intercomparisons, Lehtonen et al. (2016) estimate large ($\geq 0.1 \text{ km}^2$) boreal forest fires in Finland to increase by 1.9 times under~~
280 ~~RCP4.5 and 2.3 times under RCP8.5 by mid-century.~~ Robust predictions of future burned area in wildland and human-
281 dominated landscapes for the Arctic require an understanding and quantitative simulation of the major drivers of fire
282 (specifically climate and fire weather, ignition, fuels, and humans), including coupled dynamics between and among these
283 drivers (Riley et al., 2019).

284 285 **3.2.1 Biogeography of future fires** ~~Climate change will increase number of natural fires~~

286 The climate-induced vegetation shifts, which would also modify fire risk and related emissions, present a complex matrix for
287 the Arctic Council member states. Predictions of boreal forest transition to deciduous forest stands would decrease fire risk in
288 eastern Canada and interior Alaska (Terrier et al., 2013; Foster et al., 2019; Mekonnen et al., 2019). Wang et al. (2019) found
289 that these trends are already occurring in Alaska and Northwestern Canada using three decades of Landsat imagery with a 30
290 m resolution, as climate drives grass and shrub expansion in the Arctic and wildfires drive most of the evergreen forest
291 reduction and expansion of deciduous forests in the boreal. Further work in mature deciduous forests of Interior Alaska show
292 that current canopy “gaps” are related to ecological shifts to evergreen shrubs and lichens, grasses, and mosses, thus increasing
293 overall fire risk due to presence of these high flammability coniferous species in these small areas within low flammability

294 deciduous stands (Alexander and Mack, 2017). Further, moderate to high spatial and temporal resolution satellite mapping of
295 taiga-tundra vegetation show a northern expansion of trees, but with complex patterns of diffuse and abrupt transitions from
296 forests to non-forests (Montesano et al., 2020).

297
298 ~~The boreal and Arctic landscape is diverse, and thus so are natural fires, spanning from forests to grasslands and peatlands.~~
299 ~~Near-term warming means more ignitions from lightning (Veraverbeke et al., 2017) and degraded permafrost increasing dry~~
300 ~~ground fuels, including peat (Turetsky et al., 2015), and fire severity (Teufel and Sushama, 2019). By the end of the century,~~
301 ~~wildland fire risk is expected to increase, with length of fire seasons – measured in terms of daily severe fire weather occurrence~~
302 ~~– predicted to expand by as much as 20 days for high northern latitudes using the A1B (roughly corresponding to RCP6.0), A2~~
303 ~~(– RCP8.5), and B1 (– RCP4.5) scenarios (Flannigan et al., 2013). Similarly, Sherstyukov and Sherstyukov (2014) predict an~~
304 ~~increase of > 50 days of high fire risk days for Russia under RCP8.5 scenario, with a potential to double annual forest fire~~
305 ~~burned area by 2100. Using CMIP5 model intercomparisons, Lehtonen et al. (2016) found that large (≥ 0.1 km) boreal forest~~
306 ~~fires in Finland may double or even triple by the end of century, using RCP4.5 and RCP 8.5 scenarios, but with large inter-~~
307 ~~model variability. Robust predictions of future burned area in wildland and human-dominated landscapes for the boreal and~~
308 ~~Arctic require an understanding and quantitative simulation of the major drivers of fire (specifically climate and fire weather,~~
309 ~~ignition, fuels, and humans), including coupled dynamics between and among these drivers (Riley et al., 2019).~~

310
311 There is a consensus that prolonged fire seasons will become more common, increasing in the eastern boreal forests of Canada
312 (Boulanger et al., 2013); central and northwestern Canada (Boulanger et al., 2014); and European Russia (particularly the
313 Republic of Karelia and Leningradskaya oblast), West Siberia, and the Far East (Tchebakova et al., 2009; Sherstyukov and
314 Sherstyukov, 2014). Wang et al. (2017) note that recently the fire season in Canada is characterized by more total number of
315 fire spread days, leading to large increases in total fire size and emissions for early season fires like the Fort McMurray megafire
316 in Alberta, which burned both forests and peatlands and was caused by humans (Hanes et al., 2019). Lengthening the fire
317 season means increased potential for more and larger fire emissions throughout the fire season, starting earlier in spring and
318 lasting later into autumn. ~~Further, suppression of wildfire in Canadian boreal communities has increased their likelihood of~~
319 ~~flammability, allowing fuels to build up in and near populated places (Parisien et al., 2020), calling into question what other~~
320 ~~wildland-urban interfaces in the Arctic region may have similar risks due to long-term aggressive fire suppression. Ignition~~
321 ~~likelihood is often modeled modelled by considering the moisture conditions of ground fuels (i.e., litter) and the organic layer~~
322 ~~(i.e., forest canopy), whereby humans are the most likely source of fire on the ground and lightning the source for canopy fires~~
323 ~~(Wotton et al. 2003). Veraverbeke et al. (2017) introduced a positive feedback loop between climate, lightning, fires and~~
324 ~~northward forest expansion, whereby surface energy fluxes from forests appeared to be increasing the probability of lightning~~
325 ~~in Alaska.~~

327 ~~Climate change may have both positive and negative impacts on boreal forests and forestry (Reyer et al., 2017). Moreover,~~
328 ~~Boreal~~ fire regimes and related changes in spring albedo (relative reflectance) and the radiation balance are distinct in North
329 American (crown-fire dominated) and Northern Eurasian (surface-fire dominated, smaller negative shortwave forcing) systems
330 (Rogers et al., 2015). In the near future, these changes may be positive but become negative in the mid- and long-term. In
331 general, climate change accelerates forest growth at high northern latitudes due to a longer growing season. ~~Moreover,~~
332 ~~elevated CO₂ concentration decreases transpiration and increases photosynthetic rate and thus enhances forest growth (Peltola~~
333 ~~et al., 2002; Kellomäki et al., 2018). However, abiotic and biotic damages in particular may have opposite-negative effects on~~
334 ~~forest growth and dynamics (Seidl et al., 2014). For example, drought increases the risk of forest fires, but also negatively~~
335 ~~impacts the growth of Norway spruce (*Picea abies*) and exposes trees to biotic damages. Snow damages are estimated to~~
336 ~~increase in northeastern Europe but decrease elsewhere in Europe by end-of-century under RCP scenarios 4.5 and 8.5 scenarios~~
337 ~~(Groenemeijer et al., 2016). Wind damage risk is expected to increase due to the shortening of soil frost period (Venäläinen et~~
338 ~~al., 2020), as frozen soils anchor trees in the ground, thus making them less vulnerable to uprooting. Many forest insects~~
339 ~~responsible for bug kill of trees will benefit from climate change due to established linkage of increased habitat range and~~
340 ~~increased winter temperatures (Pureswaran et al., 2018). Climate-driven bug kill increases the amount of easily burnable~~
341 ~~material in forests and can influence fire risk. For example, a large-scale bark beetle invasion could increase the amount of~~
342 ~~fuels via dead wood, increasing ignition risk and crown fire risk as well as increasing the need, danger, and cost of fuels and~~
343 ~~fire management of insect attacked forests (Jenkins et al., 2014). According to Venäläinen et al. (2020), a warming climate is~~
344 ~~likely to increase the risk of bark beetle outbreaks and wood decay caused by *Heterobasidion spp.* root rot in Finland's~~
345 ~~coniferous forests. Siberian forests have already experienced a northern progression of the destructive Siberian moth~~
346 ~~(*Dendrolimus sibiricus* Tschetvericov) by a distance of ~ 0.5 degree and a decrease in its~~the~~ regeneration cycle from two to~~
347 ~~one year, prompted by drought and increasing temperatures (Baranchikov and Montgomery, 2014; Kharuk et al., 2017).~~
348 ~~Moreover, the probability of forest-damaging cascading and compounding events, i.e., large-scale wind damage followed by~~
349 ~~a widespread bark beetle outbreak, may increase remarkably in the future for the High Northern Latitudes. Future climate~~
350 ~~conditions are expected to become more favourable for forest fires in the boreal zone, even in highly managed regions.~~
351

Formatted: Font: Italic

352 3.2 Climate change will increase number of non-forest fires

353 Under RCP8.5, Stralberg et al. (2018) estimated that by 2100, grasslands will replace much of the upland conifer, mixed
354 forests, and deciduous forests for a large area of the boreal forest zone of northern Alberta. Shorter fire return intervals
355 combined with climate change-induced drought will reduce the resiliency of evergreen and broadleaf species to re-seed and/or
356 establish after wildfires, leading to expansion of grassland ecosystems in what is now Northern Canadian forests (Whitman et
357 al., 2019). Increased grass-dominated landscapes would create a new fire regime of frequent but low severity fires, with the
358 likelihood of SLCF transport to the Arctic most likely in the spring months of March through May (Hall and Loboda, 2018).
359 Grassland fires produce less energy, with smoke plumes more similar to crop residue burning, and are unlikely to breach the

Formatted: Swedish (Sweden)

360 tropopause for consistent, year-round transport of smoke to the Arctic (Hall and Loboda, 2017), unlike the current observed
361 deposition from boreal forest fires in the Arctic (Thomas et al., 2017). Further, Smirnov et al. (2015) found forest fires in
362 European Russia during 2008-2012 occurred mainly in June and August, with Siberia and the Russian Far East being the main
363 sources of BC emissions during a time when transport to the Arctic is unfavourable. In the Sakha Republic, Kirillina et al.
364 (2020) found that from 2011 onwards, fire seasons have been 13 days longer than previously, on average, and starting from
365 2009 onwards, fire seasons have started earlier in April, sooner than previous years. A peak fire occurrence across a three-
366 month period of May to July persists in Sakha. During the 2020 extreme fire season in Siberia, high resolution satellite data
367 from the European Space Agencies' Sentinel-2 detected fires around still-frozen thermokarst lakes above 70°N (McCarty et
368 al., 2020). This indicates that more BC from future early season burning in and near Arctic Siberia could be available for
369 transport, and thus deposition on snow and ice that accelerates melting, as well as associated climate feedback due to effect on
370 albedo. Given this, current and future early season fires are particularly relevant because Arctic snow and sea-ice coverage are
371 much more widespread in the early burning season than late season – meaning earlier BC deposition could accelerate
372 springtime melt to April, before the usual start of the melt season in May (Stroeve et al., 2014). Emission factors for biomass
373 burning in grassland and steppe ecosystems are generally smaller from those of boreal forests (Akagi et al., 2011; Andreae,
374 2019), which potentially implies different impacts on atmospheric chemistry and SLCFs. Therefore, while boreal forest fires
375 emit more SLCFs than grasslands and cropland fires, the springtime burning of northern grasslands, peatlands, and croplands
376 - often human-caused - means these emissions are more likely to be transported to the Arctic during favourable transport
377 conditions in March, April, and May than summertime forest fires.

378 4 Arctic fire emissions

379 In Section 4 and 5, we present new emissions work that builds on the 2015 AMAP assessment of BC and ozone (AMAP,
380 2015), which included 2005 biomass burning emissions from an the Global Fire Assimilation System (GFASv1.2; Kaiser et
381 al. 2012), Global Fire Emissions Database version 2 (GFEDv2; van der Werf et al. 2006), GFEDv3 (van der Werf et al. 2010),
382 the Global Inventory for Chemistry-Climate studies (GICC; Mieville et al. 2010), MACCity (Lamarque et al. 2010), and the
383 Fire Inventory from NCAR (FINNv1.5; Wiedinmyer et al., 2011) for above 60°N. For the 2021 AMAP assessment, we focused
384 on longitudinal biomass burning emission models for years 2005 through 2018 using the Global Fire Emissions Database with
385 small fires (GFEDv4s; van der Werf et al., 2017), FINNv1.5 (Wiedinmyer et al., 2011), GFASv1.2 (Kaiser et al., 2012), the
386 Quick Fire Emissions Dataset (QFEDv2.5r1; Koster et al., 2015), and the Fire Energetics and Emissions Research (FEER;
387 Ichoku and Ellison, 2014). These versions of GFAS, GFED, FINN, FEER, and QFED analysed rely on Moderate Resolution
388 Imaging Spectroradiometer (MODIS) thermal anomalies, with GFEDv4s integrating the MCD64A1 burned area product with
389 the MODIS active fire product to account for small fires. ~~It should be noted that the MCD64A1 algorithm used in GFEDv4s~~
390 ~~embeds the MODIS active fire data to seed burned area detection and growth~~ (Giglio et al., 2009). For each global fire
391 emissions model, the area of interest was defined roughly as 45° to 80° North (N) globally, split by latitude ranges of 45° to

392 50° N: Temperate, 50° to 60° N: boreal, 60° to 70° N: Low Arctic, and 70° to 80° N: High Arctic. Average annual emissions
393 from open biomass burning from all sources (agriculture, boreal forest, tundra, peat, etc.) were calculated for 2005-2018 for
394 BC, methane (CH₄), carbon monoxide (CO), and fine particulate matter (PM_{2.5}).

395

396 Since the Visible Infrared Imaging Radiometer Suite (VIIRS) provides daily, global observations of low-intensity fires
397 (Johnston et al., 2018), a custom AMAP open biomass burning emissions inventory was developed for the year 2018 to utilize
398 VIIRS's capabilities to detect smouldering fires which are common in peat landscapes. Suomi-NPP VIIRS active fire from
399 day and night detections (Oliva and Schroeder, 2015) were assumed to completely burn each 375 m² pixel. A 'best-guess' land
400 cover was created from three different land cover products, with a sample (n = 30 locations) validation of land cover type
401 performed for each country. Ultimately, the 750 m VIIRS Surface Type land cover product (Zhang et al., 2018) was used for
402 North America, Greenland, and the Russian Federation, augmented by the revised 1 km Circumpolar Arctic Vegetation Map
403 (Raster CAVM; Reynolds et al., 2019) for missing values in the high northern latitudes. For Norway, Sweden, and Finland,
404 the 10 m Land Cover Map of Europe 2017 from the Sentinel-2 Global Land Cover Project (Gromny et al., 2019) was used.
405 All land cover maps were reclassified into the International Geosphere-Biosphere Program (IGBP) classes for ease of emission
406 calculations. Fuel loadings and combustion completeness were taken from Van Leeuwen et al. (2014), with tundra values used
407 for Greenland. Emission factors were taken from Akagi et al. (2011), with updates from Andreae (2019).

408

409 Most fire activity and emissions occur between 50° and 60° N, with very few open biomass burning emissions between 70°
410 and 80° N and zero satellite observations of fire above 80° N (Fig. 2). The latitude band of 50° to 60° N corresponds to the
411 southern extents of the boreal region, an area experiencing increasing fires due to climate change (de Groot et al., 2013) and
412 includes the largest wildfires in British Columbia's history, burning 1,200 km² in summer 2017 (Kirchmeier-Young et al.,
413 2019). Note, ~~however, also~~ that fire activity detected by the 1km MODIS MCD14 Collection 6 active fire data (Giglio et al.,
414 2016), with confidence values > 50%, has a positive trend for fires occurring between 60° and 70° N, but not for the latitude
415 bands of 45° and 50° N or 50° and 60° N (Fig. 2).

416

417 In the 14-year emissions estimates from GFAS, GFED, and FINN, a clear shift has occurred in the zonal distribution of fire
418 since the mid-2000's. ~~More fire is now taking place~~ Fire emissions are increasing more north of 60° N ~~than in~~ compared to the
419 temperate zone of 45° to 50° N, where large amounts of human-caused burning and wildfires throughout North America,
420 Europe, and Eurasia occur (Fig. 2). This trend is pronounced in GFED and GFAS, with these two models showing a positive
421 trend (note the dotted line in Fig. 2), and FINN showing a slight decrease in later years even as total MODIS active fire
422 detections increased (bottom panels of Fig. 2). This trend is more pronounced in GFED and GFAS than in FINN, though all
423 models show a positive trend (note the dotted line in Fig. 2). The 2005 to 2018 multi-model annual average BC emissions from
424 all open biomass burning sources in the Arctic (60° to 80° N) and adjacent regions known to impact smoke transport into the
425 Arctic (45° to 60° N) is 0.34 Tg. The years with the highest multi-model average are 2012, 2008, and 2015 with BC emissions

426 of 0.45 Tg, 0.44 Tg, and 0.41 Tg, respectively. The lowest annual average BC emission from the five global fire emissions
427 models are 2007⁸ and 2013, [both](#) with 0.27 Tg. The fire emissions model with the consistently highest BC emissions is QFED,
428 with an annual average of 0.68 Tg (Fig. 3). FEER, GFAS, and GFED have more agreement, with annual BC emission averages
429 of 0.32 (± 0.07) Tg, 0.30 (± 0.07) Tg, and 0.25 (± 0.06) Tg, respectively. FINN has the lowest annual average BC emissions
430 of 0.130 Tg, with higher emissions in 2012 (0.20 Tg) and 2008 (0.19 Tg). The AMAP model designed specifically for the Pan-
431 Arctic, which was based on VIIRS active fire data and region-specific land cover types, produced slightly higher emission
432 estimates than FINN (Fig. 3) for year 2018. The AMAP model predicts BC emissions of 0.13 Tg and CH₄ emissions of 1.39
433 Tg, compared to FINN's 0.11 Tg of BC and 1.19 Tg of CH₄. Compared for 2018 only, GFED has marginally higher BC
434 emissions than GFAS, while methane emission estimates from GFAS are substantially higher than GFED.

435
436 Ground-based official statistics vary greatly by country or sub-region (i.e., Alaska and Greenland) for circa 2019 (Table [12](#)).
437 Suppl. Table [32](#) provides the emission variables used to calculate emissions for each country or sub-region of the Pan-Arctic
438 reporting official burned area statistics. The Russian Federation has the highest burned area, with over 100,000 km² burned.
439 In 2019, open biomass burning in European Russia - comprising [the](#) Northwestern, Central, Southern, North Caucasus, and
440 Volga Federal Districts - accounted for only 190 km² of burned area (ФБУ "АВИАЛЕСООХРАНА", 2019). Approximately
441 98.2% of burned area in Russia occurred in the Urals, Siberia, and Far East Federal Districts. In general, Greenland,
442 Fennoscandia, and European Russia are the regions with the lowest burned area and open biomass burning emissions, with all
443 regions experiencing the most burning in 50° to 60°N and the second most burning in the latitudinal band of 60° to 70°N.
444 Alaska and Canada account for approximately 29,000 km² of total pan-Arctic biomass burning and 17% of the BC emissions,
445 while the contiguous United States (CONUS) accounted for 24% of BC emissions. It should be noted that while Canada and
446 CONUS reported similar official statistics for burned area, fires in temperate zones of the CONUS tend to emit double the
447 emissions of boreal ecosystems (Table [21](#)) [due to higher fuel loadings, emission factors, and combustion completeness \(Suppl.](#)
448 [Table 3\)](#). Greenland is a novel fire regime in the Arctic, with two relatively substantial wildfires in 2017 ([Evangelidou et al.,](#)
449 [2019](#)) and 2019 ([Table 1](#)), that accounted for more burned area and emissions than Norway or Finland. In 2019, the majority
450 of open biomass burning and related emissions for the Arctic Council member states originated in Siberia and the Russian Far
451 East, followed by the CONUS, Canada, and Alaska.

452
453 Focusing on a potentially novel Arctic fire regime in Greenland allows us to localise the impact of fires on [BC](#) deposition and
454 ice, and what that may hold for the future. Unusual fires were observed in western Greenland by pilots and also confirmed by
455 satellites between 31 July and 21 August 2017, after a period of warm, dry and sunny weather. The largest wildfire grew to
456 approximately 22 km² in size, eventually extinguished by rain (Cartier, 2017)-. The fires burned > 20 km² of high carbon soils
457 - potentially peat due to smouldering and fire spread behaviour - that became vulnerable due to permafrost degradation (Daanen
458 et al., 2011). Work by Evangelidou et al. (2019) estimated the 2017 wildfire consumed a fuel amount of about 0.12 Tg of Carbon
459 (C) and emitted about 0.00002 Tg ([20 Mg](#)) of BC and 0.0007 Tg ([700 Mg](#)) of Organic Carbon (OC), including 0.00014 Tg

Formatted: Subscript

460 (140 Mg) of Brown Carbon (BrC - the portion of OC that absorbs towards shorter wavelengths). Although these fires were
461 small compared to fires burning at the same time in North America and Eurasia, a large fraction of the BC, OC, and BrC
462 emissions (30%) was deposited on the Greenland ice sheet. Measurements of aerosol optical depth in western Greenland
463 showed that the air was strongly influenced by the Canadian forest fires. Even so, the Greenland fires had an observable impact,
464 doubling the column concentrations of BC. The spatiotemporal evolution and, in particular, the top height of the plume was
465 also confirmed using the vertical cross section of total attenuated backscatter (at 532 nm) from Cloud-Aerosol Lidar and
466 Infrared Pathfinder Satellite Observations (CALIOP) Lidar. The maximum albedo change due to BC and BrC deposition from
467 the Greenland fires was -0.007 at maximum, while the average instantaneous BOA (Bottom Of the Atmosphere) radiative
468 forcing over Greenland at noon on 31 August 2017 (post-fire) was between 0.03 and 0.04 Wm⁻², with locally occurring maxima
469 up to 0.77 Wm⁻². Here, the BOA included both the aerosol effects of BC and BrC in the atmosphere and deposited on the snow.
470 The albedo effect (a decrease) was very low (0.007), practically unmeasurable. The summer 2017 fires in Greenland had a
471 small impact on the Greenland ice sheet, causing almost negligible extra radiative forcing. This was due to the comparably
472 small size of the fires in Greenland, in a global and Pan-Arctic context. However, with 30% of the emissions deposited on the
473 Greenland ice sheet, the 2017 Greenland wildfires were very efficient climate forcers on a per unit emission basis and adding
474 to current BC deposition from North American boreal forest fires (Thomas et al., 2017). Thus, while the fires in 2017 were
475 small in size on a global scale, if the expected future warming of the Arctic (IPCC, 2013) produces more and larger fires in
476 Greenland (Keegan et al., 2014), this could indeed cause substantial albedo changes and, in turn, contribute to accelerated
477 melting of the Greenland ice sheet.

478 **5 Relevance of fire sources in global and Arctic emissions: Non-fire anthropogenic versus fire emissions**

479 To place current Arctic fire emissions into context, GFASv1.2 emissions (Kaiser et al., 2012) were compared to total
480 anthropogenic emissions of BC, PM_{2.5}, and CH₄ estimated with the integrated assessment model GAINS (Greenhouse gas –
481 Air pollution Interactions and Synergies) (Amann et al., 2011; Klimont et al., 2017). The GAINS model explicitly considers
482 environmental policies and assesses their impact on current and future emissions (Amann et al., 2011; Klimont et al., 2017;
483 Amann et al, 2020) and projects emissions from various anthropogenic sectors until 2050; here we compare emissions
484 estimated for 2010, 2015, and 2020. To place current Arctic fire emissions into context, GFASv1.2 emissions (Kaiser et al.,
485 2012) were compared to non-fire anthropogenic emissions from the integrated assessment Greenhouse gas – Air pollution
486 Interactions and Synergies, or GAINS model (Amann et al., 2011, Klimont et al., 2017), with a focus on BC and PM_{2.5}. The
487 GAINS model was chosen because it considers explicitly environmental policies and assess their impact on current and future
488 emissions (Amann et al., 2011; Klimont et al., 2017; Amann et al, 2020) and projects emissions from various anthropogenic
489 sectors on five year time steps. For this comparison, we use 2010, 2015, and 2020. Global GFAS data was downloaded from
490 the European Centre for Medium-Range Weather Forecasts (ECMWF, <https://apps.ecmwf.int/datasets/data/cams-gfas/>).
491 GFAS was chosen for this comparison because it was produced in near real-time on the global scale, unlike GFED which is a

492 historical product and at the time of this writing had not completed the 2020 emission estimates. GFAS also did not show
493 consistently low emissions for the Pan-Arctic region, like FINN (Fig. 2). Further, GFAS is currently used as an operational
494 product for global and regional forecasting (Inness et al., 2019), thus likely to be integrated into policy-making decisions on
495 fire management. The GFAS ~~“wildfire”~~ wildfire and biomass burning emissions include all open biomass burning activity,
496 with no differentiation between human-caused ignitions and natural sources, like lightning, but attempt to remove spurious
497 fire emissions from industrial, volcanic, and geothermal sources (Rémy et al., 2017). Data was clipped to Pan-Arctic extents
498 at 50°N, 60°N, and 65°N. ~~The GFAS wildfire emissions data, referred to as wildfire emissions in this review due to inability~~
499 ~~to differentiate fire types in the emissions data,~~ has a spatial resolution of 0.1°, so it was aggregated to 0.5° for comparison
500 with GAINS. ~~Since the 2020 wildland fire season in the Arctic was unprecedented (Witze, 2020), with approximately 27% of~~
501 ~~fires in Siberia burning above 65°N (Conard and Ponomarev, 2020), the 2020 GFAS emissions can be used to represent what~~
502 ~~potential future fire regimes by mid-century, i.e., 2050, may be like, with climate change-driven expansion of fire seasons and~~
503 ~~likelihood for extreme fire weather and risk (see Sect. 3). Since the 2020 wildland fire season in the Arctic was unprecedented~~
504 ~~(Witze, 2020), with approximately 27% of fires in Siberia burning above 65°N (Conard and Ponomarev, 2020), we used the~~
505 ~~2020 GFAS emissions to replicate potential future fire regimes by mid-century, i.e., 2050, with climate change-driven~~
506 ~~expansion of fire seasons and likelihood for extreme fire weather and risk (see Sect. 3).~~

507
508 Fig. 4, 5, and 6 present 2010, 2015, and 2020 annual BC, PM_{2.5}, and CH₄ emissions, respectively, from four main source
509 sectors of GAINS ECLIPSEv6b (<https://iiasa.ac.at/web/home/research/researchPrograms/air/ECLIPSEv6.html>; Höglund-
510 Isaksson et al., 2020) and biomass burning from GFAS at the global-scale (left) and above 50° N and 60° N. Globally,
511 residential combustion, i.e., oil, coal, wood, etc. used for heating, is the main anthropogenic source of BC emissions for these
512 years and is the largest overall when compared with GFAS wildfire emissions (Fig. 4, left). Bond et al. (2004) estimated BC
513 emissions from open biomass burning from wildlands and agricultural fires to be higher than other sources, but we did not find
514 that when comparing GAINS emissions with GFAS fire emissions at the global scale. However, in the northern latitudes,
515 wildfires surpass the four anthropogenic sources: residential, transportation, gas flaring during oil and gas exploration and
516 production, and the sum of all other sources, i.e. ‘Others’. North of 60° N, gas flaring is the main anthropogenic source, with
517 comparable but still smaller emissions than GFAS wildfire emissions estimates. As Fig. 4 shows, 2020 was an extreme year
518 for Arctic wildfires (York et al., 2020), with BC emissions above 60° N twice as high as in 2010 and 2015. For PM_{2.5}, wildfires
519 have higher emissions than the anthropogenic sectors globally, and the difference increases in the northern latitudes (Fig. 5).
520 Globally, the agriculture sector is the main source of CH₄, with fossil fuel industry (coal, oil, gas exploration, distribution, and
521 use)energy, flaring, and waste sectors all emitting more than wildfires (Fig. 6). Above 50° N, the same anthropogenic sectors
522 are the main CH₄ sources, though in 2020 wildfires emitted more methane than the others sector. A similar phenomenon
523 occurred above 60° N, where across all years, wildfire emissions are higher than the other anthropogenic sectors except for
524 energy sector. Above 50° N, the same anthropogenic sectors are the main CH₄ sources, though in 2020 wildfires emitted more

methane than the energy sector. A occurred above 60° N, where depending on the year, wildfire emissions are comparable to the energy sector, while flaring has higher emissions than all the other sectors combined, including agriculture.

Arctic shipping is often brought up as a potentially important source of BC within the Arctic in the future. According to GAINS, in 2015 shipping comprised only 0.6 % of anthropogenic BC emissions north of 60° N. However, according to a white paper by the International Council on Clean Transportation (ICCT; Comer et al., 2020), BC emissions from Arctic shipping increased by 85% between 2015 and 2019. Their definition of Arctic is as described in the International Maritime Organization (IMO) Polar Code, i.e., they assessed shipping in much of the High Arctic above the Barents and Kara Seas but inclusive of waters between Alaska and Russia as far south as 60°N. In our comparison, shipping is included in the transport sector of GAINS emissions.

Fig. 7 shows the monthly BC emissions averaged from 2010, 2015, and 2020 for the globe and the three northern latitude breaks of 50°N, 60°N, and 65°N for the two leading sectors - wildfires and residential combustion. As with global annual emissions (Fig. 4), residential combustion is the main source sector in most months. However, in July and September the wildfire emissions are similar to residential combustion, and in August they are higher at the global-scale. These two sectors show opposite temporal profiles during the year (Suppl. Fig. 1). Residential combustion is highest in the winter months, but wildfires grow during the spring and reach their maximum in the summer, generally falling off in September with the exception of 2020 fire emissions. In the northern latitudes, strong spring emissions in April correspond with the global signal (upper panels of Fig. 7), while the summer months comprise an even larger share of the annual emissions than in the global average.

Consistently, wildfire emissions account for more than half of all black carbon emissions north of 60°N and 65°N (Fig. 8), representing up to 74% and 82% of 2020 BC emissions, respectively (Suppl. Table 34). At these northern latitudes, wildfires and flaring are the main sources of black carbon, especially north of 65°N with these two sectors accounting for 93% of black carbon emissions, compared to 88% for 60°N. North of 50°N, residential, transport, and flaring are proportionally larger than north of 60°N and 65°N, but still less than wildfire emissions (Suppl. Fig. 2). North of 60°N, wildfire emissions have increased from 2010 to 2020, particularly above 65°N. Of those wildfire emissions from GFAS that were above 60°N, 21% in 2010 and 27% in 2015 occurred above 65°N (Suppl. Table 34). However, in 2020 the percentage was 56% (Fig. 8), indicating how extreme the 2020 wildfire year was in the Arctic.

Given the large portion of black carbon emissions from fires in comparison to anthropogenic sources as modeled by GAINS, understanding the local climate and air pollution impacts for the Arctic Council region is key. For example, the timing of fires in agricultural landscapes, boreal forest fires, and the Arctic tundra occur during the early spring to early summer months (i.e., March through May for 50° N and May and June for 60° N and 65° as seen in Suppl. Fig. 1) when BC transport and deposition to the Arctic is possible and critical for the cryosphere (Hall and Loboda, 2018) and air pollution (Law and

559 Stohl, 2007), both from long-range (Thomas et al., 2017) and local sources of BC deposition (Evangelidou et al., 2019). For
560 example, BC transport is possible as early as March into mid-May for agricultural landscapes of eastern Europe (Hall and
561 Loboda, 2017) and peatlands, grasslands, and forests in North America (Qi and Wang, 2019), ~~and with~~ fires in grasslands,
562 forests, and agricultural lands most common in southern Siberia (Kukavskaya et al., 2016) and the Russian Far East (Hayasaka
563 et al., 2020) during the spring months of March, April, and May. The boreal forest fire season starts in April and May in
564 Canada (Tymstra et al., 2020) and Siberia (Soja et al., 2004b; Conard and Ponomarev, 2020), moving north into Alaska by
565 early June (Partain et al., 2015). Fires and associated transport of black carbon to the Arctic in the spring months of March to
566 June tend to be climatically important when deposition on the cryosphere can accelerate surface melting (Bond et al., 2013).
567 In spring and summer of 2020, fires in the Arctic landscape of northern Sakha Republic were burning as early as the beginning
568 of May (McCarty et al., 2020), indicating a local source of black carbon. Likewise, wildfires in Greenland in July 2017 and
569 July 2019 confirm that a local source of BC deposition on the Greenland Ice Sheet is possible (Evangelidou et al., 2019).
570 Wildfire PM_{2.5} emissions are local sources of air pollution for urban and rural communities across the Arctic (Mölders and
571 Kramm, 2018; Schmale et al., 2018), often peaking in summer months.

572 **6 Fire management in the Arctic**

573 ~~Active-f~~Fuels management, like prescribed fires and even allowing wildfires to burn under non-severe fire weather conditions,
574 may be more effective than fire suppression and/or efforts to eliminate all fire from northern landscapes (McWethy et al.,
575 2019), including in novel landscapes caused by warming in the Arctic. Fuel treatments in the boreal zones of Alaska were
576 ~~modeled~~modelled to be effective for at least 14 years post-treatment, especially in shaded fuel breaks that reduce canopy cover
577 and ladder fuels (Little et al., 2018). However, in dried and degraded peatlands of the Arctic region, fuels management will be
578 more complicated outside the boreal forest and forest-tundra gradient, where mulching treatments that convert canopy and
579 surface fuels to a masticated fuel bed can limit peat burn depth in Black Spruce (*Picea mariana*) stands (Wilkinson et al., 2018).
580 Privately-owned grassy tussock tundra and dwarf shrub tundra vegetation types are more likely to burn than low shrub tundra
581 in Alaska (Hu et al., 2015), ~~showing with~~ relatively rapid vegetation re-greening within a decade after burning for shrub and
582 tussock tundra (Rocha et al., 2012) - ~~potentially a re-establishing the shrub and tussock tundra fuelbed for repeat burns~~. While
583 prescribed burning could be effective in fuel management for tussock and dwarf shrub landscapes of the tundra, prescribed
584 burning effectiveness for peatlands is less clear. Peat fire risk and burn depth, however, is less influenced by canopy and ground
585 vegetation and more by soil bulk density (impacting air availability in soils), the water table depth, and precipitation (Kieft et
586 al., 2016). After the devastating 2010 fires in the Moscow region, the regional government undertook an ambitious 70,000 ha
587 peatland rewetting project to reduce fire risk (Sirin et al., 2014), a landscape-scale process that can be monitored using existing
588 Earth observation sensors at the moderate resolution (30 m Landsat to 10 m Sentinel-2; Sirin et al., 2018). To date, the
589 effectiveness of this campaign is unclear, but in ~~practical-term~~theory it should reduce fire risk. In the larger context of CH₄,
590 Günther et al. (2020) used a radiative forcing model to determine that methane emissions from peatland rewetting are less

591 significant in the short-term when compared to the CO₂ emissions from degraded or drained peatlands increasing long-term
592 warming when rewetting is postponed. Adaptive management strategies of the timber industry in Fennoscandia could also
593 reduce fire risk. Intensive management via ditch network maintenance and fertilization of drained peatlands will increase
594 timber values while also rewetting the peat (Ahtikoski and Hökkä, 2019). Prescribed burning for silvicultural retention and
595 maintaining and regenerating pure stands can also reduce fuel loadings while increasing biodiversity (Lindberg et al., 2020).

596
597 Human ignition sources, ~~including and determining~~ predicting future how demographic, migration, and/or development
598 patterns in these changing northern landscapes, will impact fire activity and related emissions (Robinne et al. 2016; Reilly et
599 al. 2019). For example, consider agricultural landscapes as one source of fire. Expanding climate-driven agricultural frontiers
600 in the high northern latitudes under RCP8.5 scenario for 2060-2080 could add 8.5 million km² of new croplands in Canada
601 and Russia alone, expanding wheat and maize production into areas with carbon-rich or peat soils (Hannah et al., 2020).
602 Further, Parfenova et al. (2019) found crop growing conditions would be established in some of the permafrost zones of Siberia
603 under RCPs 2.6 and 8.5 by 2080, ~~favorable~~ favorable for wheat and maize (silage) production. These crops are commonly
604 managed via open burning practices in the U.S., eastern Europe, Russia, and Canada (Kutcher and Malhi, 2010; McCarty et
605 al., 2017; Theesfeld and Jelinek, 2017; Shiwakoti et al., 2019; Thompson and Morrison, 2020). Thus, ~~seasonality of burns and~~
606 ~~management of burning of~~ croplands, grasslands, and deciduous forests ~~often may~~ occur at times when transport of fire
607 emissions to the Arctic is likely, i.e., late winter/early spring for Russia (Hall and Loboda, 2018; Qi and Wang, 2019) as well
608 ~~as and~~ Canada and north central U.S. (Viatte et al., 2015), respectively.

609
610 While open biomass burning emissions are episodic in nature when considering emissions from single extreme wildland fire
611 events and even wildfire seasons, the spring to early summer human-caused fires are a consistent source of BC and PM_{2.5} that
612 can be managed and potentially reduced. From the policy perspective, and how these events will contribute to Pan-Arctic
613 pollution, ~~fires are is~~ important to consider for future Arctic Council collaboration and coordination among Member States,
614 Arctic Indigenous Permanent Participants, and non-Arctic Observer States. For future Arctic fires, policy controls are
615 effectively limited to fuels management, reduction of human ignitions, and wildland firefighting in the Arctic and the boreal
616 zone (Flannigan et al., 2013). Further, wildland firefighting techniques in the boreal forest will not be appropriate for the more
617 fragile permafrost- and peat-dominated Arctic tundra and will need to be specifically tailored, for example, to the tundra
618 (French et al., 2015). Collaboration, cooperation, and innovation are needed for future Arctic wildland firefighting techniques,
619 practices, and implementation, particularly in the context of potential emissions ~~reductions~~ mitigation.

620 **7 Knowledge gaps and associated uncertainties**

621 Here we highlight the key problems summarized from the review of scientific literature in an attempt to focus future research
622 efforts. It is important to reduce the uncertainties below to understand Arctic fire regimes and emissions, especially given that

623 climate change potentially introduces a new fast-moving uncertainty. Improving the understanding of the current and future
624 Arctic and boreal fire regimes will be important for Arctic policymakers as well, given a rapidly changing Arctic and the
625 influence of these fire regimes on climate systems, fragile Arctic ecosystems, and society (Rogers et al., 2020). Overall, a
626 major uncertainty exists in conceptualising and documenting what constitutes a shift in fire regimes of a certain region or even
627 Pan-Arctic (i.e., current fire climatology versus fuel types) and what happens when a new regime is about to emerge (i.e.,
628 future projections of climatic and ecological conditions). Specific recommendations are made in each subsection to propose
629 next steps.

630 **7.1 Spatial and temporal modelling of future fire landscapes and regimes** ~~Future fire landscapes and regimes~~

631 Modelling future fire landscapes and regimes, in terms of coupled fire-climate-land use-ecological models, remains uncertain.

632 Future Arctic fire regimes will be ~~influenced~~impacted by ~~changing-shifting~~ vegetation on the landscape types (Tchebakova et
633 al., 2009; Sizov et al., 2021), with both climate change and subsequent fire seasons, i.e., fire disturbance, ~~impacting-determining~~
634 the species and locations of future vegetation on Arctic and boreal landscapes (Foster et al., 2019). For example, fire and the
635 thawing of permafrost are considered to be the principal mechanisms that will shape new vegetation physiognomies for Siberia
636 (Polikarpov et al., 1998; Tchebakova et al., 2010). It is important to note that moisture from summertime thaw of the active
637 layer of permafrost provides necessary moisture for forest growth in the dry environment of interior Siberia, otherwise only
638 steppe could exist without this additional moisture (Shumilova, 1962). In the dry climate in interior Siberia, frequent fires
639 eliminate any of the dark conifer undergrowth that may have become established in suitable sites within the permafrost zone.
640 The fire return interval in the light conifer (larch, *Larix spp.*, and Scots pine, *Pinus sylvestris*) middle taiga in central Siberia
641 is 20–30 years (Furyaev et al., 2001), compared to 200–300 years in dark conifer (Siberian pine, *Pinus sibirica*, and fir, *Abies*
642 *sibirica*) forests in southern Siberia, including mountain taiga. Slowly growing dark conifers are not adapted to frequent fires
643 and typically die; additionally, they are not light-tolerant, so they are not likely to be the first species to succeed following fire
644 events. On the other hand, *Larix dahurica* is evolutionarily adapted to fire and successfully regenerates when cones open
645 following fire events. For East Siberia, Polikarpov et al. (1998) speculated that post-fire succession would mean that dark
646 conifers would be replaced by Scots pine in southern dry climates and by larch on cold soils in a warmer climate. ~~Zonal-d~~Dark
647 conifers, which survive in specific climatic zones, would shift northwards and eastwards following permafrost retreat, and
648 light-needled tree species (e.g., *Pinus sylvestris* and *Larix sibirica*) would follow them, expanding from the south. In the
649 transition zone between dark-needled and light-needled tree species, birch and mixed light conifer-hardwoods subtaiga and
650 forest-steppe would dominate, likely reducing fire risk. In the southern tundra of Yamalo-Nenets Autonomous Okrug in
651 northwest Siberia, a transition from dry dwarf shrub to woodlands (< 50% of area is covered by trees) has been documented
652 in previously burned areas (Sizov et al., 2021).

653
654 Total area of Siberian forests are predicted to decrease and shift northwards, with-and forest-steppe and steppe ecosystems ~~are~~
655 predicted to dominate 50% of Siberia by 2080 under RCP 8.5 (Parfenova et al., 2019), meaning agriculture in Siberia would

656 likely benefit from climate warming. About 50-85% of central Siberia was predicted to be climatically suitable for agriculture
657 (Tchebakova et al., 2011), although potential croplands would be limited by availability of suitable soils. Crop production may
658 increase by twofold. The introduction of new agricultural crops could likely be less costly than afforestation with new tree
659 species climatotypes. Farming may be a preferred land use choice in the future where forests would fail due to climate change,
660 with regional business and economy authorities determining what specific measures may be undertaken to support forestry,
661 agriculture, or mixed agriculture and forestry practices in order to optimize economic loss or gain effects of climate change.
662 Therefore, understanding how climate change and ongoing fire disturbance in the boreal and Arctic will impact species
663 distribution, and thus fuel availability, remains complex (Shuman et al., 2017) and more work in coupled fire-climate-
664 ecological models, with considerations for permafrost and human-driven land use and ignition in emerging agricultural
665 systems, for the Arctic and boreal is needed.

666 7.2 Peatlands

667 Peat smouldering can emit large quantities of smoke, contributing to hazardous air quality (Hu et al., 2018). Current global
668 fire emissions inventories underestimate peat fires, as forest fuel types currently drive fuels maps and profiles (Liu et al., 2020).
669 Boreal zone peatland fires are not well quantified in terms of fuel loadings (Van Leuwen et al., 2014). High uncertainty in
670 emission factors for boreal peat fires (Hu et al., 2018) has led to improved laboratory-derived emission factors from sampled
671 peat from Russia and Alaska (Watson et al., 2019). [Recent laboratory work on fire mechanisms of organic soils and how peat
672 fires spread improves the understanding of these processes \(for example, Huang et al., 2017; Huang et al., 2015; Prat-Guitart,
673 2016; Huang et al., 2019; Christensen et al., 2020; Santoso et al., 2021; Yuan et al., 2021\), though a need for Pan-Arctic field
674 observations persists.](#) Burn depth is also not well captured outside of localized spatial scales, like sampling plots, given lack
675 of Earth observation sensing capabilities and pre-fire and post-fire soil surveys (Rogers et al., 2014), which can lead to
676 emissions underestimations.

677
678 With a warming climate, there is a risk of increasing peatland and “legacy carbon” fires (Ingram et al., 2019) in boreal forests,
679 particularly in stands younger than 60 years where drying limits the resilience of the carbon rich soils (Walker et al., 2019)
680 and in drying fen watersheds near large settlements, like the costliest wildfire in Canada’s history - the May 2016 Horse
681 River/Fort McMurray fire (Elmes et al., 2018). Future emission estimates from peat fires will need to be informed by where
682 and in what condition these carbon-rich soils reside, particularly as predicted moderate and severe drought in boreal peatlands
683 western Canada are expected to increase fire size by over 500% (Thompson et al., 2019). [Current Earth system models do not
684 typically characterise well or include peat fires and related feedbacks \(Lasslop et al., 2019; Loisel et al., 2020\), further limiting
685 our ability to predict future emissions from peatland burning.](#) Mapping pan-Arctic peatlands has proved challenging (Yu et al.,
686 2010; Xu et al., 2018), with recent improvements linking permafrost to peat storage (Hugelius et al., 2020). Further, difficulties
687 in estimating and/or accounting for water table depth and moisture content of peat when modeling depth of burn and associated
688 emissions during smouldering is a key observational uncertainty (Kiely et al, 2019). Future fuels data will need to account for

689 how the complexities of the boreal and Arctic peat topography will impact rate of post-fire peat soil accumulation (Ingram et
690 al., 2019), with some landscapes remaining resilient with other marginal peat areas with severe smouldering and fewer
691 sediment inputs becoming sources of legacy carbon emissions, thus driving future fuels availability. Current Earth system
692 models underestimate evaporative water loss and overestimate current and future water availability for boreal peatland systems
693 under RCP 4.5 and 8.5 warming scenarios [when compared to current climatic conditions](#), perhaps underestimating fire risk,
694 activity, and emissions in peat systems (Helbig et al., 2020).

695
696 Climate mitigation efforts, like restoration or rewetting of peatlands, do not eliminate the role of fire as a management tool
697 (Davies et al., 2016) nor the risk of wildland fire in peat landscapes. Thus, estimates of future fire emissions will need to
698 assimilate complexities associated with peat fuel conditions and loadings. For example, restoration of peat is not a linear
699 process, with previous results in Canada showing one to two decades needed for restoration and rewetting of degraded
700 peatlands that have residual peat and vegetation to ‘seed’ the sites (Nugent et al., 2019). Until these restored peatlands have
701 sufficient moisture and vegetation cover, they are still susceptible to fire risk. Burn depth in peat can be limited in naturally
702 wet and rewetted peatlands if the surface maintains a high moisture content via hydrological and vegetation processes (Granath
703 et al., 2016). Maintaining these needed hydrological processes is difficult for degraded, unmanaged peatlands. In Alberta,
704 wildland peat sites lacking constant sources of water and depositional inputs experienced severe burning on margins (Ingram
705 et al., 2019), while Wilkinson et al. (2019) found forested peatland margins were extremely vulnerable to peat smouldering
706 combustion, especially in previously burned areas with > 60 years since fire. Ronkainen et al. (2013) expect a warmer climate
707 to lower water tables via evapotranspiration for unmanaged peatlands in Finland, thus increasing wildfire risk. Producing more
708 complete estimates of fuel loadings for peatlands across the Arctic region can follow methodologies set by Johnston et al.
709 (2015) to augment the dynamic boreal, taiga, and tundra fuel loadings, e.g. Innes (2013) and Ivanova et al. (2019).

710 7.3 Permafrost

711 Approximately half of all peatlands in the Northern Hemisphere are coincidental with permafrost (Hugelius et al., 2020), with
712 many discontinuous permafrost sites dominated by peatlands in Canada (Estop-Aragonés et al., 2018; Gibson et al., 2018),
713 Russia (Hugelius et al., 2014), and Sweden (Chang et al., 2019). In the flat West Siberian terrain, Kotlyakov and Khromova
714 (2002) and Malevsky-Malevich et al. (2001) show no continuous or discontinuous permafrost below 65°N, which influences
715 the viable vegetation for the tundra and sparse *Larix sibirica* taiga. Current climate models may be missing the link between
716 melting ground ice, sometimes referred to as thermokarst processes, and potential permafrost degradation of the currently
717 stable and carbon-rich northeast Siberian Arctic lowlands (NESAL). Nitzbon et al. (2020) indicate that we can expect a
718 threefold increase of permafrost thaw in the NESAL region under RCP4.5 (a stabilization scenario) by 2100 when thermokarst
719 processes are combined with increased temperature projections in numerical modelling, potentially increasing the amount of
720 peat fuels in an already high fire activity region. Combining current peatland distribution maps with newer [modeledmodelled](#)
721 datasets of predicted mid-century and late-century permafrost extent and geohazard indices under climate-forcing scenarios

722 (Karjalainen et al., 2019) can reduce uncertainties to determine: 1) increased peat fire risk and locations due to permafrost
723 thaw and 2) decreased capability to deploy ground-level wildland firefighting, thus limiting ability to control future peat fires
724 and fire emissions in the Pan-Arctic. Further, permafrost thawing changes hydrology (e.g. greater river discharge or
725 disappearing lakes) and geomorphology (solifluction and thermokarst processes) across broad expanses of the contemporary
726 permafrost zone. In a warmer and drier climate, many locations in the Arctic may be affected by solifluction, with thermokarst
727 modified by frequent catastrophic fires, and deeper active layer thaw. As a whole, retreating permafrost should cause a
728 reduction in the area of forests and their replacement by steppe on well-drained, tilted geomorphology (Lawrence and Slater,
729 2005) or by bogs on poorly-drained plains (Tchebakova et al., 2009).

730

731 Permafrost areas, especially at their southern distributions, are being disturbed by wildfires (Holloway et al., 2020). In Alaska
732 and northwestern Canada, the impacts of wildfire disturbances on permafrost have been well quantified. For instance, post-
733 fire permafrost change in Alaska showed surface warming greater in boreal sites than tundra, with surface temperatures higher
734 for previously burned sites than at unburned sites, even after vegetation recovered for one to four decades (Jiang et al., 2017).
735 In the North Slope of Alaska, recent evidence suggests that a transition from grasses to shrubbier conditions is occurring post-
736 tundra fires (Jones et al., 2013). Though the vast majority of fires in the continuous and discontinuous permafrost zones occur
737 in deciduous needleleaf forests (Lorant et al., 2016), knowledge gaps on post-fire permafrost resiliency exist for larch-
738 dominated forests (*Larix spp.*) in Siberia. For instance, recent work in Sakha Republic found that a 36 km² wildfire in an open
739 larch with shrub and moss lichen landscape northwest of the Batagaika megaslump resulted in approximately 3.5 million cubic
740 meters of thawed permafrost five years later (Yanagiya & Furuya, 2020). Likewise, uncertainties persist for post-fire
741 permafrost resiliency in the boreal forests of eastern Canadian, like Quebec and Labrador (Holloway et al., 2020). As with
742 peatlands, improved geospatial products advance our understanding of the potential for impacts of wildfires across large spatial
743 scales (Hugelius et al., 2020).

744

745 **7.4 Satellite-based fire emissions**

746 Fire regimes for the boreal are often described by impacts to and from fire emissions (Rogers et al., 2020), with many modeling
747 emissions in the high northern latitudes using Earth observations. Uncertainties in emission models are driven by availability
748 and quality of fire activity data from satellite- and ground-based sources, as well as incomplete knowledge of fuels and
749 emission factors. Current global fire emission inventories rely on satellite-derived fire activity from active fire detections,
750 burned area mapping, and fire radiative power (Liu et al., 2020). A comparison of four satellite-based global fire emissions
751 databases over North America - GFED, FINN, GFAS, QFED - found that assumed portions of dry matter in fuels and not
752 emission factors were creating biomass burning aerosol estimates that differ by factors of four to seven, essentially limiting
753 the ability to accurately quantify the impact of smoke on climate and air quality (Carter et al., 2020). Given the international

754 scientific community's reliance on two main fire emissions factor sources (Akagi et al., 2011; Andraea, 2019 as an update to
755 Andraea & Merlet, 2001), information available for a robust uncertainty analysis for this variable is limited (Pan et al., 2020).

756

757 Satellite-based observations of fire in the Arctic and boreal regions underestimate open burning in agricultural landscapes,
758 surface fires in boreal forests, and smouldering peat fires. For example, current emission inventories based on satellite-derived
759 products of burned area, like GFEDv4, underestimate human-caused burning in agricultural landscapes and mixed forests in
760 Eurasia between 50° to 65° N by approximately 2,100 km² annually (Zhu et al., 2017), indicating that actual burned area from
761 anthropogenic ignitions in the Eurasian boreal zone is currently underestimated by as much as 16%. Surface fires under forest
762 canopies dominate fire regimes in much of Northern Eurasia, but these fires are not well quantified in current satellite-based
763 burned area products (Rogers et al., 2015; Duncan et al., 2020) and thus emission inventories. Smouldering fires in carbon-
764 rich humus and peat landscapes will be difficult to detect, as smouldering combustion occurs at much lower temperatures than
765 flaming combustion; 500°C to 700°C versus 1500°C to 1800°C, respectively (Rein et al. 2008). As previously mentioned,
766 daily, global observations of low-intensity fire from existing satellite systems are limited currently to VIIRS (Johnston et al.,
767 2018), as it was designed to detect smaller and cooler fires than MODIS. For this review, the versions of GFAS, GFED, FINN,
768 FEER, and QFED analysed rely on MODIS thermal anomalies, unlike the custom AMAP fire emissions which used VIIRS
769 only. Smouldering fires in the Arctic can be mapped via regionally-tuned algorithms designed to ingest daily active fire
770 detections from multispectral VIIRS (Waigl et al., 2017) and hyperspectral Hyperion (Waigl et al., 2019) sensors. In general,
771 satellite and drone detections (Burke et al., 2019) of smouldering peat fires are difficult because ground fires are low
772 temperature and can burn underground and re-emerge in new locations (Rein, 2016), with additional existing detection
773 constraints from coarse resolution (> 1 km) global satellite sensors, canopy cover, and cloud cover (Johnston et al., 2018).

774

775 A further complication is that peat fires can smoulder for months, years, and even decades (Hu et al., 2018), burning laterally
776 and vertically below surface, appearing to be extinguished, but releasing smoke at the surface in a different location from the
777 original ignition site. This phenomenon is referred to as holdover, overwintered, and/or zombie fires, and makes it difficult to
778 allocate as a single - but complex - fire event from cumulative satellite active fire and burned area pixels. For example, in April
779 2020, the Alaska Division of Forestry was monitoring several active smouldering peat fires from the ~ 5 km² Deshka Landing
780 Fire of August 2019 that had overwintered near Willow, Alaska despite heavy snow melt (Alaska Wildland Fire Information,
781 2020). Preliminary results by Scholten and Veraverbeke (2020), indicate that overwintering fires are more likely to be
782 holdovers from high severity fires, emerging more frequently in lowland black spruce-dominated boreal forests. McCarty et
783 al. (2020) hypothesize that some of the earliest fires along still-frozen thermokarst lakes of Sahka Republic in May 2020 may
784 be holdover fires, as the drivers and extent of early season human-caused ignitions are still not well-documented in the
785 scientific literature for much of the Arctic.

786

7.5 Lack of agreement between official statistics and satellite observations

Earth observations from satellite products are powerful tools for forecasting (Pickell et al., 2017), improving rapid response post-fire modelling (Miller et al., 2017), and quantifying fire in the boreal and Arctic regions (Hislop et al., 2020). Consistently, however, there has been little correlation between satellite-derived and official estimates of burned area (Fusco et al., 2019). Loepfe et al. (2012) found that multiple satellite fire products had high correlation with official reports of burned areas for Sweden, but little to no correlation with official statistics for Finland. Agreement of burned area within Siberian forests between official Russian burned area-statistics and four satellite-based burned area products from fires in Siberian forests was less than 10% ~~when compared to four satellite-based burned area products~~ (Kukavskaya et al., 2013). Average official satellite-derived Russian burned area estimates differ by a mean of 48% for 2002 to 2015 in comparison to the Loboda et al. (2017) regionally-tuned product, which only differs by a mean of 18% in comparison to official burned area statistics for Alaska and Canada. One reason for these differences could be regional-to-global scale algorithms may not have the sensitivity necessary to define surface fire, which is the dominant fire type in Siberia in normal fire years. Also, North American and Nordic countries have long-term ground-based boreal burned area records that span 50 years or greater, which aids in calibrating current satellite data records and analysing relationships between fire regimes, vegetation, weather, and climate. Long-term accurate fire records do not exist for much of Russia, primarily because fire was not historically recorded in the remote ‘unprotected territories’ (Sofronov et al., 1998; Soja et al., 2004). Consequently, understanding of the balance between surface-to-crown fire and the ecosystem-dependent areas that burn in Siberia is limited, which adversely affects fire emissions estimates. The Global Wildfire Information System (GWIS; <https://gwis.jrc.ec.europa.eu/>), a joint program between the Group on Earth Observations (GEO; https://www.earthobservations.org/geoss_wp.php), Copernicus (<https://www.copernicus.eu/en/services/emergency>), and NASA (<https://www.nasa.gov/>), uses the MODIS MOD64A1 Collection 6 Burned Area product (Giglio et al., 2018) to create country-level burned area statistics. GWIS satellite-derived burned area overestimates open biomass burning in both Norway and Finland by 199% and 129%, respectively, when compared to official statistics (Table 3). Though, GWIS underestimates open biomass burning in Sweden by 48%. The work of the SLCF EG was unable to determine exact reasons for why this mismatch occurs, though previous work has shown that satellite-based fire observations are more likely to align with official records as fire sizes increase (Fusco et al., 2019). Both Norway and Finland reported the lowest fire activity and burned area (Table 1). Future open biomass burning emissions will need improved satellite fire detection methodologies for the Arctic and boreal regions ~~and also~~ shorter latency in ground reports and statistics from official agencies. Further, verifying and relating satellite detections of fires ~~via to~~ ground-level verification will require a concerted effort and likely lead to a better understanding of how and why these two fire data sources do not presently align.

817 **8 People and future Arctic fire regimes**

818 Prevention and management of Pan-Arctic fires are limited to reduction of human-caused ignitions and management of
819 landscape fuels (Flannigan et al., 2013). The impact of humans on fire risk is dependent on local- to national-scale actions that
820 may increase fire and emissions via deforestation, transportation networks, energy extraction, and agricultural open burning
821 as well as decrease fire and fire emissions via active suppression. On a practical level, people are the main ignition sources for
822 fires in the Arctic region, while lightning ignitions tend to lead to larger fires. In interior Alaska, where lightning-caused fires
823 account for 95% of total burned area (Veraverbeke et al., 2017), 52% of total ignitions were human in origin but occurred in
824 areas of high fire suppression resulting in only 5% of total burned area from 1990 to 2016 (Calef et al., 2017). Archard et al.
825 (2008) estimated 65% of all forest fires in the Russian Federation were caused by human ignition, and a more recent study
826 found approximately half of all fires in Sakha Republic are caused by anthropogenic activities (Kirillina et al., 2020).
827 Throughout boreal Canada, anthropogenic factors increase fire probability (Parisien et al., 2016), with humans igniting most
828 fires close to roads while lightning-caused fires are responsible for the majority of burned area in the more remote locations
829 (Gralewicz et al., 2012). Blouin et al. (2016) found that 45% of wildfires in Alberta were started by lightning, but responsible
830 for 71% of burned area. In Finland, lightning-caused fires account for less than 15% of forest fires (Larjavaara et al., 2005).
831 Machines used for forestry operations in stony areas of Sweden account for 7-10% of total annual ignitions and 40% of total
832 burned area (Sjöström et al., 2019). For the 19 European countries reporting fires and ignition sources to the European Forest
833 Fire Information System (EFFIS; <https://effis.jrc.ec.europa.eu/>), de Rigo et al. (2017) determined only 4% of fires were from
834 natural sources, with half of the fire records lacking a verified cause.

835
836 Indigenous Fire Management (IFM) and understanding Indigenous use of fire, as well as fire risk and response to fire events
837 (Mottershead et al., 2020), are needed in a changing Arctic environment. IFM is more frequently being deployed in fire-prone
838 and/or fire-adapted areas (Nikolakis et al., 2020), which accounts for much of the boreal but not necessarily Arctic ecosystems.
839 Cogos et al. (2019) documented historical place names in northern Sweden (e.g., *roavve* and *roavvi*) related to historical Saami
840 practices of burning pine heath landscapes to improve long term foraging of reindeer. Approximately one out of every ten
841 people in the Arctic are Indigenous (Nordregio, 2019), comprising an estimated 15% of the population of Alaska, 53% of
842 the northern territories of Canada, and 98% of Greenland, for a total of 1.13 million Indigenous peoples in the Pan-Arctic
843 (Young and Bjerregaard, 2019). Arctic communities are demanding more leadership roles in climate research and applications
844 (Stone, 2020). Research- and experiential-driven recommendations on how to incorporate traditional, Indigenous knowledge
845 into Arctic Council working groups efforts, including (1) Use of participatory methodology; (2) Use of Indigenous
846 methodologies; (3) Recognition that traditional ecological knowledge is local; (4) Application to policy; and (5) Cross-cultural
847 understanding (Sidorova, 2020), align well with community- and landscape-driven fire science methodologies needed to
848 predict future fire risk (Bowman et al., 2020; Johnston et al., 2020) and to answer many of the fire regime and emission,

849 including ignition and fuel type, uncertainties raised in this review. Who better to ask - and to lead - than the people who live
850 there?

851 **9 Conclusions**

852 Since the mid-2000s, emissions from open biomass burning have increased above 60°N, with fires above 66°N occurring
853 earlier in the year and burning later into the growing season, indicative of changing Arctic fire regime. Compared to
854 anthropogenic sources in the GAINS model, biomass burning already accounts for more BC and PM_{2.5} emissions than
855 anthropogenic sources north of 60°N, including flaring from associated gas from oil and natural gas extraction. Increased
856 length in fire seasons is coupled with prediction of increased fire severity, with predictions of essentially physically
857 unmanageable crown fires in the boreal as soon as 2050 (Wootton et al., 2017). Future emissions from fires are difficult to
858 predict and here more work is needed. For example, emissions from functionally uncontrollable fires in boreal forests are not
859 well quantified due to uncertainties in combustion efficiency observations and estimates (Xu et al., 2020). [Improving our](#)
860 [understanding of the future of Arctic fires and fire emissions will also allow us to better predict future Earth system processes](#)
861 [- both at high latitudes and globally.](#)
862

863 In contributing to the AMAP 2021 assessment of SLCFs, this review was driven by policy questions identified by member
864 states of the Arctic Council (Table 1), and builds on the 2011 (AMAP, 2011) and 2015 (AMAP, 2015) reports, which included
865 some analysis and discussion of natural, ‘semi-natural’ (i.e., human-caused ignitions in wildland landscapes), and agricultural
866 field burning. We did not perform a systematic review of the fire research literature (Robinne et al., 2020), and the existing
867 literature cited was not assessed for limitations or errors. Further, while the authors attempted to cite published literature and
868 official fire statistics for the seven Arctic Council states experiencing open biomass burning (excluding Iceland), we know that
869 bias may still be present in the over 200 peer-reviewed sources of literature and data chosen for this review (Johnston et al.,
870 2020). This review is a starting point, a foundation for future Pan-Arctic research agendas for fire monitoring and needed
871 systematic reviews (Haddaway et al., 2020) of future fire risk, fire emissions, and fire prevention and management in the Arctic
872 - all needed to accurately describe future Arctic fire regimes.
873

874 Future Arctic fire regimes will likely be driven by climate change impacts on fuels, including the interactions between peat
875 and permafrost, fire weather, and ignition sources as well as the complexities of climate and fire disturbance changing
876 vegetation types (Tchebakova et al., 2009; Shuman et al., 2017). The consensus of current literature is that climate change and
877 human activity will increase fire risk in the Arctic, via increased lightning strikes, thawing of permafrost, transitions to grasses,
878 taiga, and dry peat, and more human-caused ignitions. In eastern Canada, the northward expansion of deciduous forests will
879 likely decrease fire risk, which may also be true for portions of southern Siberia and Fennoscandia. Human- and lightning-
880 caused fires are likely to increase given expansion of energy extraction, transportation networks, tourism, and climate change.

881 Further, Arctic landscapes are complex, with high levels of localized heterogeneity due to polygonal tundra landforms (Lara
882 et al., 2020), complex and endemic vegetation types and communities (Raynolds et al., 2019), and topography (Morin et al.,
883 2016). Future fire emissions studies will need to integrate multiple datasets to accurately quantify Arctic fire regimes (Masrur
884 et al., 2018), including climate, permafrost conditions, aboveground, surface, and peat fuels, topography, land use, Indigenous
885 and local fire management, seasonality of burns, and ignition sources.

886
887 Human activity and communities in the Arctic will need to adapt to increasing fire risk. To prepare for these 21st century
888 changes to the Arctic fire regime, evidence-based fire monitoring and management - including prevention strategies - must
889 incorporate Indigenous and local knowledge in the Arctic. This will require increasing transdisciplinary research (Sidorova,
890 2020) to understand and predict fire in the North, how humans are and must adapt to a new fire prone landscape in the
891 Anthropocene (Bowman et al., 2020), and Pan-Arctic collaboration and cooperation. Understanding ecological landscape
892 changes, predicted to substantially increase across Asian Russia, is crucial information for developing viable strategies for
893 long-term economic and social development in preparation for climate migration and strategic adaptation planning (Parfenova
894 et al., 2019).

895
896 The Arctic Council's role as an agent of change in the region is promising, as it has moved its role from policy informing to
897 policy making (Barry et al., 2020). Given the extreme fire season of 2020, an Arctic Council-led initiative for Pan-Arctic fire
898 monitoring, prevention, and management is strongly needed for a rapidly changing Arctic (McCarty et al., 2020). Such efforts
899 have started, including the Arctic Wildland Fire Ecology Mapping and Monitoring Project (Arctic FIRE;
900 <https://www.caff.is/arcticfire>) led by the Gwich'in Council International, an Indigenous Permanent Participant, via the
901 Conservation of Arctic Flora and Fauna (CAFF) working group of the Arctic Council, as well as other Arctic Council activities.
902 Potentially expanding existing efforts or coordinating with new initiatives to incorporate the five other Indigenous permanent
903 participants, as well as more efforts from the science and disaster response agencies of the eight member states and the expertise
904 of other Arctic Council working groups, could create the type of community- and Arctic-centric science needed for Pan-Arctic
905 fire policies and to increase the capacity for the Indigenous peoples of the Arctic to monitor and protect their Arctic homelands
906 (Wilson, 2020) from fire risk and to adapt to the changing Arctic fire regime.

908 **References**

909 Abatzoglou, J. T., and Williams, A. P.: Impact of anthropogenic climate change on wildfire across western US forests, *Proc.*
910 *Natl. Acad. Sci.*, 113, 11770-11775. <https://doi.org/10.1073/pnas.1607171113>, 2016.

911 Ahtikoski, A., and Hökkä, H: Intensive forest management—does it pay off financially on drained peatlands?, *Can. J. For.*
912 *Res.*, 49, 1101-1113, <https://doi.org/10.1139/cjfr-2019-0007>, 2019.

913 Akagi, S. K., Yokelson, R.J., Wiedinmyer, C., Alvarado, M.J., Reid, J.S., Karl, T., Crouse, J.D. and Wennberg, P.O.:
914 Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039,
915 <https://doi.org/10.5194/acp-11-4039-2011>, 2011.

916 Alaska Division of Forestry: 2019 EOY handout, available at:
917 <http://forestry.alaska.gov/Assets/pdfs/firestats/2019%20Alaska%20Fire%20Statistics.pdf>, 2020.

918 Alaska Wildland Fire Information: Despite heavy snow melt, Deshka Landing hot spots still smoldering, available at:
919 <https://akfireinfo.com/2020/04/24/despite-heavy-snow-melt-deshka-landing-hot-spots-still-smoldering/>, 2020.

920 Alexander, H. D., and Mack, M. C.: Gap regeneration within mature deciduous forests of Interior Alaska: Implications for
921 future forest change, *For. Ecol. Manage.*, 396, 35-43, <https://doi.org/10.1016/j.foreco.2017.04.005>, 2017.

922 Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M.,
923 Rafaj, P., and Sandler, R.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy
924 applications, *Environ. Model Softw.*, 26, 1489-1501, <https://doi.org/10.1016/j.envsoft.2011.07.012>, 2011.

925 Amann, M., Kiesewetter, G., Schöpp, W., Klimont, Z., Winiwarter, W., Cofala, J., Rafaj, P., Höglund-Isaksson, L., Gomez-
926 Sabriana, A., Heyes, C. and Purohit, P.: Reducing global air pollution: the scope for further policy interventions, *Philos T R*
927 *Soc A.*, 378, 20190331, <https://doi.org/10.1098/rsta.2019.0331>, 2020.

928 AMAP: AMAP Assessment Report: Arctic Pollution Issues, Arctic Monitoring and Assessment Programme (AMAP), Oslo,
929 Norway, xii+859 pp, available at: <https://www.amap.no/documents/doc/amap-assessment-report-arctic-pollution-issues/68>,
930 1998.

931 AMAP Assessment 2011: The Impact of Black Carbon on Arctic Climate. Arctic Monitoring and Assessment Programme
932 (AMAP), Oslo, Norway, Technical Report no. 4, available at: <https://www.amap.no/documents/download/977/inline>, 2011.

933 AMAP Assessment 2015: Black carbon and ozone as Arctic climate forcers. Arctic Monitoring and Assessment Programme
934 (AMAP), Oslo, Norway, available at: <http://hdl.handle.net/11374/1607>, 2015.

935 AMAP Assessment 2021: Impacts of short-lived climate forcers on Arctic climate, air quality, and human health, Arctic
936 Monitoring and Assessment Programme (AMAP), Tromsø, Norway (in prep).

- 937 Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15,
938 955-966, <https://doi.org/10.1029/2000GB001382>, 2001.
- 939 Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*,
940 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- 941 Baranchikov, Y.N. and Montgomery, M.E.: Chapter XXXVI - Siberian Moth, in *The use of classical biological control to*
942 *preserve forests in North America*, edited by Van Driesche, R. and Reardon, R.C., United States Department of Agriculture,
943 Forest Service, Forest Health Technology Enterprise Team, Morgantown, WV, USA,383-391, 2014.
- 944 Barry, T., Davíðsdóttir, B., Einarsson, N., and Young, O. R.: The Arctic Council: an agent of change?, *Glob Environ Change*,
945 63, 102099, <https://doi.org/10.1016/j.gloenvcha.2020.102099>, 2020.
- 946 [Betänkande av 2018 års skogsbrandsutredning: Skogsbränderna sommaren 2018 \[Forest fires in summer 2018, in Swedish\].](#)
947 [Statens offentliga utredningar \(SOU\) 2019:7, Stockholm. 1-334, 2019.](#)
- 948 Bieniek, P.A., Bhatt, U.S., York, A., Walsh, J.E., Lader, R., Strader, H., Ziel, R., Jandt, R.R., and Thoman, R.L.: Lightning
949 variability in dynamically downscaled simulations of Alaska’s present and future summer climate, *J Appl Meteorol Climatol*,
950 59, 1139-1152, <https://doi.org/10.1175/JAMC-D-19-0209.1>, 2020.
- 951 Blyakharchuk, T. A., Tchebakova, N.M., Parfenova, E.I., and Soja, A.J.: Potential influence of the late Holocene climate on
952 settled farming versus nomadic cattle herding in the Minusinsk Hollow, south-central Siberia, *Environ. Res. Lett.*, 9, 065004,
953 <https://doi.org/10.1088/1748-9326/9/6/065004>, 2014.
- 954 Boike, J., Grau, T., Heim, B., Günther, F., Langer, M., Muster, S., Gouttevin, I. and Lange, S.: Satellite-derived changes in the
955 permafrost landscape of central Yakutia, 2000–2011: Wetting, drying, and fires, *Glob Planet Change*, 139, 116,
956 <https://doi.org/10.1016/j.gloplacha.2016.01.001>, 2016.
- 957 Blouin, K. D., Flannigan, M. D., Wang, X., and Kochtubajda, B.: Ensemble lightning prediction models for the province of
958 Alberta, Canada, *Int J Wildland Fire*, 25, 421-432, <https://doi.org/10.1071/WF15111>, 2016.
- 959 Bond, T.C., Streets, D.G., Yarber, K.F., Nelson, S.M., Woo, J.H. and Klimont, Z.: A technology-based global inventory of
960 black and organic carbon emissions from combustion, *J. Geophys. Res. Atmos*, 109, 203,
961 <https://doi.org/10.1029/2003JD003697>, 2004.

Formatted: Swedish (Sweden)

- 962 Bond, T.C., Doherty, S.J., Fahey, D.W., Forster, P.M., Bernsten, T., DeAngelo, B.J., Flanner, M.G., Ghan, S., Kärcher, B.,
963 Koch, D., and Kinne, S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.*
964 *Atmos.*, 118, 5380-5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- 965 Boulanger, Y., Gauthier, S., Gray, D. R., Le Goff, H., Lefort, P., and Morissette, J.: Fire regime zonation under current and
966 future climate over eastern Canada, *Ecol Appl.*, 23, 904-923, <https://doi.org/10.1890/12-0698.1>, 2013.
- 967 Boulanger, Y., Gauthier, S., & Burton, P. J.: A refinement of models projecting future Canadian fire regimes using
968 homogeneous fire regime zones, *Can. J. For. Res.*, 44, 365-376, <https://doi.org/10.1139/cjfr-2013-0372>, 2014.
- 969 Bowman, D. M., Kolden, C. A., Abatzoglou, J. T., Johnston, F. H., van der Werf, G. R., and Flannigan, M.: Vegetation fires
970 in the Anthropocene, *Nat. Rev. Earth Environ.*, 1, 500-515, <https://doi.org/10.1038/s43017-020-0085-3>, 2020.
- 971 Burke, C., Wich, S., Kusin, K., McAree, O., Harrison, M.E., Ripoll, B., Ermiasi, Y., Mulero-Pázmány, M., and Longmore, S.:
972 Thermal-Drones as a Safe and Reliable Method for Detecting Subterranean Peat Fires, *Drones*, 3, 23,
973 <https://doi.org/10.3390/drones3010023>, 2019.
- 974 Calef, M. P., Varvak, A., and McGuire, A. D.: Differences in human versus lightning fires between urban and rural areas of
975 the boreal forest in interior Alaska, *Forests*, 8, 422, <https://doi.org/10.3390/f8110422>, 2017.
- 976 Carter, T.S., Heald, C.L., Jimenez, J.L., Campuzano-Jost, P., Kondo, Y., Moteki, N., Schwarz, J.P., Wiedinmyer, C.,
977 Darmenov, A.S., da Silva, A.M. and Kaiser, J.W.: 2020. How emissions uncertainty influences the distribution and radiative
978 impacts of smoke from fires in North America, *Atmos. Chem. Physics*, 20: 2073–2097, [https://doi.org/10.5194/acp-20-2073-](https://doi.org/10.5194/acp-20-2073-2020)
979 2020, 2020.
- 980 Cartier, K. M. S.: Southern Greenland wildfire extinguished. *Eos*, 98, <https://doi.org/10.1029/2017EO080905>, 2017.
- 981 Chang, K. Y., Riley, W. J., Crill, P. M., Grant, R. F., Rich, V. I., and Saleska, S. R.: Large carbon cycle sensitivities to climate
982 across a permafrost thaw gradient in subarctic Sweden, *The Cryosphere*, 13, 647–663, <https://doi.org/10.5194/tc-13-647-2019>,
983 2019.
- 984 Chernokulsky, A., and Esau, I: Cloud cover and cloud types in the Eurasian Arctic in 1936–2012. *Int J Climatol*, 39, 5771-
985 5790, <https://doi.org/10.1002/joc.6187>, 2019.
- 986 [Christensen, E.G., Fernandez-Anez, N. and Rein, G.: Influence of soil conditions on the multidimensional spread of](https://doi.org/10.1016/j.combustflame.2019.11.001)
987 [smouldering combustion in shallow layers, *Combust Flame*, 214, 361-370,](https://doi.org/10.1016/j.combustflame.2019.11.001)
988 <https://doi.org/10.1016/j.combustflame.2019.11.001>, 2020.

Formatted: Finnish

989 CIFFC: Canadian Interagency Forest Fire Centre: Fire Hectares by Year, available at: <https://ciffc.net/en/ext/hectares-by-year>,
990 2020.

991 Comer, B., Osipova, L., Georgeff, E., and Mao, X.: The International Maritime Organization's proposed Arctic heavy fuel oil
992 ban: Likely implications and opportunities for improvement, International Council on Clean Transportation, available at:
993 <https://theicct.org/sites/default/files/publications/Arctic-HFO-ban-sept2020.pdf> , 2020.

994 Conard, S.G. and Ivanova, G.A.: Wildfire in Russian boreal forests—Potential impacts of fire regime characteristics on
995 emissions and global carbon balance estimates, *Environ. Pollut.*, 98, 305, [https://doi.org/10.1016/S0269-7491\(97\)00140-1](https://doi.org/10.1016/S0269-7491(97)00140-1),
996 1997.

997 Conard, S.G., and Ponomarev, E.: Fire in the North, *Wildfire Magazine*, available at: [https://www.iawfonline.org/article/fire-](https://www.iawfonline.org/article/fire-in-the-north-the-2020-siberian-fire-season/)
998 [in-the-north-the-2020-siberian-fire-season/](https://www.iawfonline.org/article/fire-in-the-north-the-2020-siberian-fire-season/), 2020.

999 Cogos, S., Östlund, L. and Roturier, S.: Forest fire and indigenous Sami land use: place names, fire dynamics, and ecosystem
1000 change in Northern Scandinavia, *Human Ecology*, 47, 51-64, <https://doi.org/10.1007/s10745-019-0056-9>, 2019.

1001 Daanen, R. P., Ingeman-Nielsen, T., Marchenko, S. S., Romanovsky, V. E., Foged, N., Stendel, M., Christensen, J. H., and
1002 Hornbech Svendsen, K.: Permafrost degradation risk zone assessment using simulation models, *The Cryosphere*, 5, 1043–
1003 1056, <https://doi.org/10.5194/tc-5-1043-2011>, 2011.

1004 Davies, G. M., Kettridge, N., Stoof, C.R., Gray, A., Ascoli, D., Fernandes, P.M., Marrs, R., Allen, K.A., Doerr, S.H., Clay,
1005 G.D. and Mc Morrow, J.: The role of fire in UK peatland and moorland management: the need for informed, unbiased debate,
1006 *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, 371, 20150342, <https://doi.org/10.1098/rstb.2015.0342>, 2016.

1007 De Groot, W. J., Flannigan, M. D., and Stocks, B. J.: Climate change and wildfires, González-Cabán, Armando, tech. coord.
1008 Proceedings of the fourth international symposium on fire economics, planning, and policy: climate change and wildfires,
1009 available at: https://www.fs.fed.us/psw/publications/documents/psw_gtr245/psw_gtr245_001.pdf, 2013.

1010 de Rigo, D., Libertà, G., Houston Durrant, T., Artés Vivancos, T., and San-Miguel-Ayanz, J.: Forest fire danger extremes in
1011 Europe under climate change: variability and uncertainty, Publication Office of the European Union, Luxembourg,
1012 <https://doi.org/10.2760/13180>, 2017.

1013 Dronin, N., and Kirilenko, A.: Climate change, food stress, and security in Russia, *Reg Environ Change*, 11(1), 167-178,
1014 <https://doi.org/10.1007/s10113-010-0165-x>, 2011.

1015 DSB: Direktoratet for samfunnsikkerhet og beredskap, Personal communication, March 2020, Homepage:
1016 <https://www.dsb.no/>, 2020.

1017 Duncan, B. N., Ott, L. E., Abshire, J. B., Brucker, L., Carroll, M. L., Carton, J., et al.: Space-Based Observations for
1018 Understanding Changes in the Arctic-Boreal Zone. *Rev. Geophys.*, 58, e2019RG000652,
1019 <https://doi.org/10.1029/2019RG000652>, 2020.

1020 Elmes, M. C., Thompson, D. K., Sherwood, J. H., and Price, J. S.: Hydrometeorological conditions preceding wildfire, and the
1021 subsequent burning of a fen watershed in Fort McMurray, Alberta, Canada, *Nat. Hazards Earth Syst. Sci.*, 18, 157–170,
1022 <https://doi.org/10.5194/nhess-18-157-2018>, 2018.

1023 Estop-Aragonés, C., Czimczik, C. I., Heffernan, L., Gibson, C., Walker, J. C., Xu, X., and Olefeldt, D.: Respiration of aged
1024 soil carbon during fall in permafrost peatlands enhanced by active layer deepening following wildfire but limited following
1025 thermokarst, *Environ. Res. Lett.*, 13, 085002, <https://doi.org/10.1088/1748-9326/aad5f0>, 2018.

1026 Evangeliou, N., Kylling, A., Eckhardt, S., Myrioniuk, V., Stebel, K., Paugam, R., Zibtsev, S., and Stohl, A.: Open fires in
1027 Greenland in summer 2017: transport, deposition and radiative effects of BC, OC and BrC emissions, *Atmos. Chem. Phys.*,
1028 19, 1393–1411, <https://doi.org/10.5194/acp-19-1393-2019>, 2019.

1029 [Fisher, J. A., Jacob, D. J., Purdy, M. T., Kopacz, M., Le Sager, P., Carouge, C., Holmes, C. D., Yantosca, R. M., Batchelor,](#)
1030 [R. L., Strong, K., Diskin, G. S., Fuelberg, H. E., Holloway, J. S., Hyer, E. J., McMillan, W. W., Warner, J., Streets, D. G.,](#)
1031 [Zhang, Q., Wang, Y., and Wu, S.: Source attribution and interannual variability of Arctic pollution in spring constrained by](#)
1032 [aircraft \(ARCTAS, ARCPAC\) and satellite \(AIRS\) observations of carbon monoxide, *Atmos. Chem. Phys.*, 10, 977–996,](#)
1033 <https://doi.org/10.5194/acp-10-977-2010>, 2010.

1034 Flannigan, M., Cantin, A. S., De Groot, W. J., Wotton, M., Newbery, A., and Gowman, L. M.: Global wildland fire season
1035 severity in the 21st century, *Forest Ecol Manag.*, 294, 54–61, <https://doi.org/10.1016/j.foreco.2012.10.022>, 2013.

1036 Foster, A.C., Armstrong, A.H., Shuman, J.K., Shugart, H.H., Rogers, B.M., Mack, M.C., Goetz, S.J., and Ranson, K.J.:
1037 Importance of tree-and species-level interactions with wildfire, climate, and soils in interior Alaska: Implications for forest
1038 change under a warming climate, *Ecol Modell.*, 409, 108765, <https://doi.org/10.1016/j.ecolmodel.2019.108765>, 2019.

1039 French, N. H., Jenkins, L. K., Loboda, T. V., Flannigan, M., Jandt, R., Bourgeau-Chavez, L. L., and Whitley, M.: Fire in arctic
1040 tundra of Alaska: past fire activity, future fire potential, and significance for land management and ecology, *Int J Wildland*
1041 *Fire*, 24, 1045–1061, <https://doi.org/10.1071/wf14167>, 2015.

1042 [Furyaev, V.V.: Pyrological regimes and dynamics of the southern taiga forests in Siberia, in: Fire in ecosystems of boreal](#)
1043 [Eurasia, edited by: Goldammer J.G., and Furyaev V.V., Springer, Dordrecht, Netherlands, 168-185,](#)
1044 https://doi.org/10.1007/978-94-015-8737-2_12, 1996.

1045 [Furyaev, V.V., Vaganov, E.A., Tchebakova, N.M., and Valendik, E.N.: Effects of fire and climate on successions and structural](#)
1046 [changes in the Siberian boreal forest, Eurasian J. For. Res. 2, 1-15, 2001.](#)

1047 Fusco, E. J., Finn, J. T., Abatzoglou, J. T., Balch, J. K., Dadashi, S., and Bradley, B. A.: Detection rates and biases of fire
1048 observations from MODIS and agency reports in the conterminous United States, *Remote Sens. Environ.*, 220, 30-40,
1049 <https://doi.org/10.1016/j.rse.2018.10.028>, 2019.

1050 Gibson, C. M., Chasmer, L. E., Thompson, D. K., Quinton, W. L., Flannigan, M. D., and Olefeldt, D.: Wildfire as a major
1051 driver of recent permafrost thaw in boreal peatlands, *Nat. Commun.*, 9(1), 1-9, <https://doi.org/10.1038/s41467-018-05457-1> ,
1052 2018.

1053 Giglio, L., Loboda, T., Roy, D. P., Quayle, B., and Justice, C. O.: An active-fire based burned area mapping algorithm for the
1054 MODIS sensor, *Remote Sens. Environ.*, 113, 408-420, <https://doi.org/10.1016/j.rse.2008.10.006>, 2009.

1055 Giglio, L., Schroeder, W. and Justice, C.O.: The collection 6 MODIS active fire detection algorithm and fire products, *Remote*
1056 *Sens. Environ.*, 178, 31, <https://doi.org/10.1016/j.rse.2016.02.054>, 2016.

1057 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L., and Justice, C. O.: The Collection 6 MODIS burned area mapping
1058 algorithm and product, *Remote Sens. Environ.*, 217, 72-85, <https://doi.org/10.1016/j.rse.2018.08.005>, 2018.

1059 Girardin, M. P., and Terrier, A.: Mitigating risks of future wildfires by management of the forest composition: an analysis of
1060 the offsetting potential through boreal Canada, *Clim. Change*, 130(4), 587-601, <https://doi.org/10.1007/s10584-015-1373-7>,
1061 2015.

1062 Gralewicz, N. J., Nelson, T. A., and Wulder, M. A.: Factors influencing national scale wildfire susceptibility in Canada, *Forest*
1063 *Ecol Manage.*, 265, 20-29, <https://doi.org/10.1016/j.foreco.2011.10.031>, 2012.

1064 Granath, G., Moore, P. A., Lukenbach, M. C., and Waddington, J. M.: Mitigating wildfire carbon loss in managed northern
1065 peatlands through restoration, *Sci. Rep.*, 6(1), 1-9, <https://doi.org/10.1038/srep28498>, 2016.

1066 Granström, A., and Niklasson, M.: Potentials and limitations for human control over historic fire regimes in the boreal forest,
1067 *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, 363, 2351-2356, <https://doi.org/10.1098/rstb.2007.2205>, 2008.

1068 Groenemeijer, P., Vajda, A., Lehtonen, I., Kämäräinen, M., Venäläinen, A., Gregow, H., Becker, N., Nissen, K., Ulbrich, U.,
1069 Paprotny, D., & Morales Napoles, O.: Present and future probability of meteorological and hydrological hazards in Europe,
1070 Final report of Deliverable 2.5 for the Risk Analysis of Infrastructure Networks in response to extreme weather (RAIN) project,
1071 available at: http://rain-project.eu/wp-content/uploads/2016/09/D2.5_REPORT_final.pdf, 2016.

1072 Gromny, E., Lewiński, S., Rybicki, M., Malinowski, R., Krupiński, M., Nowakowski, A., and Jenerowicz, M.: Creation of
1073 training dataset for Sentinel-2 land cover classification, In *Photonics Applications in Astronomy, Communications, Industry,
1074 and High-Energy Physics Experiments 2019* (Vol. 11176, p. 111763D), International Society for Optics and Photonics,
1075 available at: <http://s2glc.cbk.waw.pl/>, 2019.

1076 Günther, A., Barthelmes, A., Huth, V., Joosten, H., Jurasinski, G., Koebsch, F., and Couwenberg, J.: Prompt rewetting of
1077 drained peatlands reduces climate warming despite methane emissions, *Nat. Commun.*, 11, 1644.
1078 <https://doi.org/10.1038/s41467-020-15499-z>, 2020.

1079 Haddaway, N.R., Bethel, A., Dicks, L.V., Koricheva, J., Macura, B., Petrokofsky, G., Pullin, A.S., Savilaakso, S. and Stewart,
1080 G.B.: Eight problems with literature reviews and how to fix them, *Nat. Ecol. Evol.*, 4, 1582, <https://doi.org/10.1038/s41559-020-01295-x>, 2020.

1082 Hall, J. V., and Loboda, T. V.: Quantifying the Potential for Low-Level Transport of Black Carbon Emissions from Cropland
1083 Burning in Russia to the Snow-Covered Arctic. *Front. Earth Sci.*, 5, 109, <https://doi.org/10.3389/feart.2017.00109>, 2017.

1084 Hall, J., and Loboda, T.: Quantifying the variability of potential black carbon transport from cropland burning in Russia driven
1085 by atmospheric blocking events, *Environ. Res. Lett.*, 13, 055010, <https://doi.org/10.1088/1748-9326/aabf65>, 2018.

1086 Hanes, C.C., Wang, X., Jain, P., Parisien, M.A., Little, J.M. and Flannigan, M.D.: Fire-regime changes in Canada over the last
1087 half century, *Can. J. For. Res.*, 49, 256, <https://doi.org/10.1139/cjfr-2018-0293>, 2019.

1088 Hannah, L., Roehrdanz, P.R., KC, K.B., Fraser, E.D., Donatti, C.I., Saenz, L., Wright, T.M., Hijmans, R.J., Mulligan, M.,
1089 Berg, A., and van Soesbergen, A.: The environmental consequences of climate-driven agricultural frontiers, *PLoS One*, 15,
1090 e0228305, <https://doi.org/10.1371/journal.pone.0228305>, 2020.

1091 Hayasaka, H., Sokolova, G. V., Ostroukhov, A., and Naito, D: Classification of Active Fires and Weather Conditions in the
1092 Lower Amur River Basin, *Rem. Sens.*, 12, 3204, <https://doi.org/10.3390/rs12193204>, 2020.

1093 Helbig, M., Waddington, J.M., Alekseychik, P., Amiro, B.D., Aurela, M., Barr, A.G., Black, T.A., Blanken, P.D., Carey, S.K.,
1094 Chen, J. and Chi, J.: Increasing contribution of peatlands to boreal evapotranspiration in a warming climate, *Nat. Clim. Chang.*,
1095 10, 555, <https://doi.org/10.1038/s41558-020-0763-7>, 2020.

1096 Hlásny, T., Krokene, P., Liebhold, A., Montagné-Huck, C., Müller, J., Qin, H., Raffa, K., Schelhaas, M., Seidl, R., Svoboda,
1097 M. and Viiri, H.: Living with bark beetles: impacts, outlook and management options (No. 8), European Forest Institute,
1098 available at: https://efi.int/sites/default/files/files/publication-bank/2019/efi_fstp_8_2019.pdf, 2019.

1099 Hislop, S., Haywood, A., Jones, S., Soto-Berelov, M., Skidmore, A., and Nguyen, T. H.: A satellite data driven approach to
1100 monitoring and reporting fire disturbance and recovery across boreal and temperate forests, *Int. J Appl. Earth Obs*, 87, 102034,
1101 <https://doi.org/10.1016/j.jag.2019.102034>, 2020.

1102 Höglund-Isaksson, L., Gómez-Sanabria, A., Klimont, Z., Rafaj, P., Schöpp, W.: Technical potentials and costs for reducing
1103 global anthropogenic methane emissions in the 2050 timeframe –results from the GAINS model, *Environmental Research*
1104 *Communications*, 2, 025004, <https://doi.org/10.1088/2515-7620/ab7457>, 2020.

1105 Holloway, J. E., Lewkowicz, A. G., Douglas, T. A., Li, X., Turetsky, M. R., Baltzer, J. L., and Jin, H.: Impact of wildfire on
1106 permafrost landscapes: A review of recent advances and future prospects, *Permafr. Periglac. Process.*, 31, 371,
1107 <https://doi.org/10.1002/ppp.2048>, 2020.

1108 Hu, F.S., Higuera, P.E., Duffy, P., Chipman, M.L., Rocha, A.V., Young, A.M., Kelly, R. and Dietze, M.C.: Arctic tundra fires:
1109 natural variability and responses to climate change, *Front. Ecol. Environ.*, 13, 369, <https://doi.org/10.1890/150063.2015>.

1110 Hu, Y., Fernandez-Anez, N., Smith, T. E., and Rein, G.: Review of emissions from smouldering peat fires and their contribution
1111 to regional haze episodes, *Int J Wildland Fire*, 27, 293, <https://doi.org/10.1071/wf17084>, 2018.

1112 [Huang, X., and Rein, G.: Computational study of critical moisture and depth of burn in peat fires, *Int J Wildland Fire*, 24,](#)
1113 [798-808, <https://doi.org/10.1071/WF14178>, 2015.](#)

1114 [Huang, X., and Rein, G.: Downward spread of smouldering peat fire: the role of moisture, density and oxygen supply, *Int J*](#)
1115 [Wildland Fire., 26, 907-918, <https://doi.org/10.1071/WF16198>, 2017.](#)

1116 [Huang, X. and Rein, G.: Upward-and-downward spread of smoldering peat fire, *Proc Combust Inst*, 37, 4025-4033,](#)
1117 [https://doi.org/10.1016/j.proci.2018.05.125, 2019.](#)

1118 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirrmeister, L., Grosse, G., Michaelson,
1119 G.J., Koven, C.D., and O'Donnell, J. A.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges
1120 and identified data gaps, *Biogeosciences*, 11(23), 6573, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.

1121 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R.B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M.,
1122 Siewert, M.B., and Treat, C.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw, *Proc Natl Acad*
1123 *Sci*, 117, 20438, <https://doi.org/10.1073/pnas.1916387117>, 2020.

1124 Ichoku, C., and Ellison, L.: Global top-down smoke-aerosol emissions estimation using satellite fire radiative power
1125 measurements, *Atmos. Chem. Phys.*, 14, 6643, <https://doi.org/10.5194/acp-14-6643-2014>, 2014.

1126 Ingram, R. C., Moore, P. A., Wilkinson, S., Petrone, R. M., and Waddington, J. M.: Postfire soil carbon accumulation does
1127 not recover boreal peatland combustion loss in some hydrogeological settings, *J. Geophys. Res. Biogeosci.*, 124, 775,
1128 <https://doi.org/10.1029/2018jg004716>, 2019.

1129 Innes, R.J.: Fire regimes of Alaskan tundra communities, U.S. Department of Agriculture, Forest Service, Rocky Mountain
1130 Research Station, Fire Sciences Laboratory (Producer), available at:
1131 www.fs.fed.us/database/feis/fire_regimes/AK_tundra/all.html, 2013.

1132 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R.,
1133 Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy,
1134 S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19, 3515–3556,
1135 <https://doi.org/10.5194/acp-19-3515-2019>, 2019.

1136 Ioffe, G., and Nefedova, T.: Marginal farmland in European Russia, *Eurasian Geogr. Econ.*, 45(1), 45,
1137 <https://doi.org/10.2747/1538-7216.45.1.45>, 2004.

1138 IPCC Climate Change 2013: The Physical Science Basis. Contribution to the Fifth Assessment Report of the Intergovernmental
1139 Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels,
1140 A., Xia, Y., Bex, V. and Midgley, P.M., available at: <https://www.ipcc.ch/report/ar5/wg1/>, 2013.

1141 Ivanova, G. A., Kukavskaya, E. A., Ivanov, V. A., Conard, S. G., and McRae, D. J.: Fuel characteristics, loads and consumption
1142 in Scots pine forests of central Siberia, *J For Res*, 31, 2507, <https://doi.org/10.1007/s11676-019-01038-0>, 2019

1143 Jain, P., Tye, M. R., Paimazumder, D., and Flannigan, M.: Downscaling fire weather extremes from historical and projected
1144 climate models, *Clim Change*, 1-28, <https://doi.org/10.1007/s10584-020-02865-5>, 2020.

1145 Jenkins, M. J., Runyon, J. B., Fettig, C. J., Page, W. G., and Bentz, B. J.: Interactions among the mountain pine beetle, fires,
1146 and fuels, *For Sci*, 60, 489-501, <https://doi.org/10.5849/forsci.13-017>, 2014.

1147 Jiang, Y., Rocha, A. V., O'Donnell, J. A., Drysdale, J. A., Rastetter, E. B., Shaver, G. R., and Zhuang, Q.: Contrasting soil
1148 thermal responses to fire in Alaskan tundra and boreal forest, *J. Geophys. Res. Earth. Surf.*, 120, 363,
1149 <https://doi.org/10.1002/2014jf003180>, 2015.

1150 Johnston, D. C., Turetsky, M. R., Benscoter, B. W., and Wotton, B. M.: Fuel load, structure, and potential fire behaviour in
1151 black spruce bogs, *Can J Forest Res*, 45, 888, <https://doi.org/10.1139/cjfr-2014-0334>, 2015.

1152 Johnston, J. M., Johnston, L. M., Wooster, M. J., Brookes, A., McFayden, C., and Cantin, A. S.: Satellite detection limitations
1153 of sub-canopy smouldering wildfires in the North American Boreal Forest, *Fire*, 1, 28, <https://doi.org/10.3390/fire1020028>,
1154 2018.

1155 Johnston, L.M., Wang, X., Erni, S., Taylor, S.W., McFayden, C.B., Oliver, J.A., Stockdale, C., Christianson, A., Boulanger,
1156 Y., Gauthier, S., and Arseneault, D.: Wildland fire risk research in Canada, *Environ. Rev.*, 28, 164,
1157 <https://dx.doi.org/10.1139/er-2019-0046>, 2020.

1158 Jones, B.M., Breen, A.L., Gaglioti, B.V., Mann, D.H., Rocha, A.V., Grosse, G., Arp, C.D., Kunz, M.L. and Walker, D.A.:
1159 Identification of unrecognized tundra fire events on the north slope of Alaska, *J. Geophys. Res. Biogeosci.*, 118, 1334,
1160 <https://doi.org/10.1002/jgrg.20113>, 2013.

1161 Jones, B. M., Grosse, G., Arp, C. D., Miller, E., Liu, L., Hayes, D. J., and Larsen, C. F.: Recent Arctic tundra fire initiates
1162 widespread thermokarst development, *Sci. Rep.*, 5, 15865, <https://doi.org/10.1038/srep15865>, 2015.

1163 Kaiser, J.W., Heil, A., Andreae, M.O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.J., Razinger, M., Schultz, M.G.,
1164 Suttie, M., and Van Der Werf, G.R.: Biomass burning emissions estimated with a global fire assimilation system based on
1165 observed fire radiative power, *Biogeosciences*, 9, 527-554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.

1166 Karjalainen, O., Aalto, J., Luoto, M., Westermann, S., Romanovsky, V.E., Nelson, F.E., Etzelmüller, B. and Hjort, J.:
1167 Circumpolar permafrost maps and geohazard indices for near-future infrastructure risk assessments, *Scientific Data*, 6, 190037,
1168 <https://doi.org/10.1038/sdata.2019.37>, 2019.

1169 Keegan, K. M., Albert, M. R., McConnell, J. R., and Baker, I.: Climate change and forest fires synergistically drive widespread
1170 melt events of the Greenland Ice Sheet. *Proc. Natl. Acad. Sci.*, 111, 7964, <https://doi.org/10.1073/pnas.1405397111>, 2014.

1171 Kellomäki, S., Strandman, H., Heinonen, T., Asikainen, A., Venäläinen, A., and Peltola, H.: Temporal and spatial change in
1172 diameter growth of boreal Scots pine, Norway spruce, and birch under recent-generation (CMIP5) global climate model
1173 projections for the 21st century. *Forests*, 9, 118, <https://doi.org/10.3390/f9030118>, 2018.

1174 Ketola, J.: Forest fire activity and burned area for Finland, Emergency Services Academy, Personal communication to Henrik
1175 Lindberg, based on rescue service database PRONTO, available at: <https://prontonet.fi/Pronto3/online3/OnlineTilastot.htm>,
1176 2020.

1177 [Kicklighter, D.W., Cai, Y., Zhuang, Q., Parfenova, E.I., Paltsev, S., Sokolov, A.P., Melillo, J.M., Reilly, J.M., Tchepakova,](#)
1178 [N.M., and Lu, X.: Potential influence of climate-induced vegetation shifts on future land use and associated land carbon fluxes](#)
1179 [in Northern Eurasia, *Environ. Res. Lett.* 9, 035004, <https://doi.org/10.1088/1748-9326/9/3/035004>, 2014.](#)

1180 Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borcen-Kleefeld, J., and Schöpp, W.: Global
1181 anthropogenic emissions of particulate matter including black carbon. *Atmos. Chem. Phys.*, 17, 8681,
1182 <https://doi.org/10.5194/acp-17-8681-2017>, 2017.

1183 Kharuk, V. I., Im, S.T., Ranson, K.J., and Yagunov, M.N.: Climate-Induced Northerly Expansion of Siberian Silkmoth Range,
1184 *Forests*, 8, 301, <https://doi.org/10.3390/f8080301>, 2017.

1185 Kharuk, V.I., Ponomarev, E.I., Ivanova, G.A., Dvinskaya, M.L., Coogan, S.C. and Flannigan, M.D.: Wildfires in the Siberian
1186 taiga, *Ambio*, 1, 1-22, <https://doi.org/10.1007/s13280-020-01490-x>, 2021.

1187 Kieft, J., Smith, T., Someshwar, S., and Boer, R.: Towards Anticipatory Management of Peat Fires to Enhance Local Resilience
1188 and Reduce Natural Capital Depletion, *Ecosystem-Based Disaster Risk Reduction and Adaptation in Practice*, Springer, 2016.

1189 Kiely, L., Spracklen, D. V., Wiedinmyer, C., Conibear, L., Reddington, C. L., Archer-Nicholls, S., Lowe, D., Arnold, S. R.,
1190 Knote, C., Khan, M. F., Latif, M. T., Kuwata, M., Budisulistiorini, S. H., and Syaufina, L.: New estimate of particulate
1191 emissions from Indonesian peat fires in 2015, *Atmos. Chem. Phys.*, 19, 11105–11121, [https://doi.org/10.5194/acp-19-11105-](https://doi.org/10.5194/acp-19-11105-2019f)
1192 [2019f](https://doi.org/10.5194/acp-19-11105-2019f), 2019.

1193 King, M., Altdorff, D., Li, P., Galagedara, L., Holden, J., and Unc, A.: Northward shift of the agricultural climate zone under
1194 21 st-century global climate change, *Sci. Rep.*, 8, 7904, <https://doi.org/10.1038/s41598-018-26321-8>, 2018.

1195 Kim, J. S., Kug, J. S., Jeong, S. J., Park, H., and Schaepman-Strub, G.: Extensive fires in southeastern Siberian permafrost
1196 linked to preceding Arctic Oscillation, *Sci. Adv.*, 6, eaax3308, <https://doi.org/10.1126/sciadv.aax3308>, 2020.

1197 Kirchmeier-Young, M. C., Gillett, N. P., Zwiers, F. W., Cannon, A. J., and Anslow, F. S.: Attribution of the Influence of
1198 Human-Induced Climate Change on an Extreme Fire Season, *Earths Future*, 7, 2, <https://doi.org/10.1029/2018ef001050>, 2019.

1199 Kirillina, K., Shvetsov, E. G., Protopopova, V. V., Thiesmeyer, L., and Yan, W.: Consideration of anthropogenic factors in
1200 boreal forest fire regime changes during rapid socio-economic development: case study of forestry districts with increasing
1201 burnt area in the Sakha Republic, Russia. *Environ. Res. Lett.*, 15, 035009, <https://doi.org/10.1088/1748-9326/ab6c6e>, 2020.

1202 Knorr, W., Dentener, F., Hantson, S., Jiang, L., Klimont, Z., and Arneth, A.: Air quality impacts of European wildfire emissions
1203 in a changing climate, *Atmos. Chem. Phys.*, 16, 5685, <https://doi.org/10.5194/acp-16-5685-2016>, 2016.

1204 Koster, R. D., Darmenov, A. S., and da Silva, A. M.: The Quick Fire Emissions Dataset (QFED): Documentation of Versions
1205 2.1, 2.2 and 2.4, Technical Report Series on Global Modeling and Data Assimilation, available at:
1206 <https://ntrs.nasa.gov/search.jsp?R=20180005253>, 2015.

1207 Kotlyakov, V. and Khromova, T.: Land Resources of Russia – Maps of Permafrost and Ground Ice, Boulder, Colorado USA:
1208 National Snow and Ice Data Center, available at: <https://nsidc.org/data/GGD600/versions/1>, 2002.

1209 Krause, A., Kloster, S., Wilkenskjeld, S., and Paeth, H.: The sensitivity of global wildfires to simulated past, present, and
1210 future lightning frequency, *J. Geophys. Res. Biogeosci.*, 119, 312, <https://doi.org/10.1002/2013jg002502>, 2014.

1211 [Krawchuk, M. A., Cumming, S.G., Flannigan, M.D., and Wein, R.W.: Biotic and abiotic regulation of lightning fire initiation](#)
1212 [in the mixedwood boreal forest, *Ecology*, 87, 458-468, <https://doi.org/10.1890/05-1021>, 2006.](#)

1213 Krawchuk, M. A., Moritz, M. A., Parisien, M. A., Van Dorn, J., and Hayhoe, K.: Global pyrogeography: the current and future
1214 distribution of wildfire, *PLOS ONE*, 4, e5102, <https://doi.org/10.1371/journal.pone.0005102>, 2009.

1215 Krawchuk, M.A., and Moritz, M.A.: Constraints on global fire activity vary across a resource gradient, *Ecology*, 92, 121,
1216 <https://doi.org/10.1890/09-1843>, 2011.

1217 Krylov, A., McCarty, J. L., Potapov, P., Loboda, T., Tyukavina, A., Turubanova, S., and Hansen, M. C.: Remote sensing
1218 estimates of stand-replacement fires in Russia, 2002–2011, *Environ. Res. Lett.*, 9, 105007, <https://doi.org/10.1088/1748-9326/9/10/105007>, 2014.

1220 Kukavskaya, E. A., Soja, A. J., Petkov, A. P., Ponomarev, E. I., Ivanova, G. A., and Conard, S. G.: Fire emissions estimates
1221 in Siberia: evaluation of uncertainties in area burned, land cover, and fuel consumption, *Can. J. Forest Res.*, 43, 493,
1222 <https://doi.org/10.1139/cjfr-2012-0367>, 2013.

- 1223 Kukavskaya, E.A., Buryak, L.V., Shvetsov, E.G., Conard, S.G. and Kalenskaya, O.P.: The impact of increasing fire frequency
1224 on forest transformations in southern Siberia, *Forest Ecol. Manag.*, 382, 225, <https://doi.org/10.1016/j.foreco.2016.10.015>,
1225 2016.
- 1226 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C., Mieville, A., Owen, B.,
1227 Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N.,
1228 McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass
1229 burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039,
1230 <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- 1231 Lara, M.J., McGuire, A.D., Euskirchen, E.S., Genet, H., Yi, S., Rutter, R., Iversen, C., Sloan, V. and Wullschleger, S.D.:
1232 Local-scale Arctic tundra heterogeneity affects regional-scale carbon dynamics, *Nat Commun*, 11, 4925,
1233 <https://doi.org/10.1038/s41467-020-18768-z>, 2020.
- 1234 Larjavaara, M., Kuuluvainen, T., and Rita, H.: Spatial distribution of lightning-ignited forest fires in Finland, *Forest Ecol.*
1235 *Manag.*, 208, 177, <https://doi.org/10.1016/j.foreco.2004.12.005>, 2005.
- 1236 [Lasslop, G., Coppola, A.I., Voulgarakis, A., Yue, C. and Veraverbeke, S.: Influence of Fire on the Carbon Cycle and Climate,](#)
1237 [Curr Clim Change Rep. 5, 112–123, <https://doi.org/10.1007/s40641-019-00128-9>, 2019.](#)
- 1238 Law, K. S., and Stohl, A.: Arctic air pollution: Origins and impacts, *Science*, 315, 1537,
1239 <https://doi.org/10.1126/science.1137695>, 2007.
- 1240 Lawrence, D. M., and Slater, A. G.: A projection of severe near-surface permafrost degradation during the 21st century,
1241 *Geophys. Res. Lett.*, 32,L24401, <https://doi.org/10.1029/2005GL025080>, 2005.
- 1242 Lee, S. H., Lawrence, Z. D., Butler, A. H., and Karpechko, A. Y.: Seasonal Forecasts of the Exceptional Northern Hemisphere
1243 Winter of 2020, *Geophys. Res. Lett.*, e2020GL090328, <https://doi.org/10.1029/2020GL090328>, 2020.
- 1244 Lehtonen, I., Venäläinen, A., Kämäräinen, M., Peltola, H., and Gregow, H.: Risk of large-scale fires in boreal forests of Finland
1245 under changing climate, *Natural Hazards Earth Syst. Sci.*, 16, 239, <https://doi.org/10.5194/nhess-16-239-2016>, 2016.
- 1246 Lidskog, R., Johansson, J., and Sjödin, D.: Wildfires, responsibility and trust: public understanding of Sweden's largest
1247 wildfire, *Scand. J. For. Res.*, 34, 319, <https://doi.org/10.1080/02827581.2019.1598483>, 2019.
- 1248 Lindberg, H., Punttila, P., and Vanha-Majamaa, I.: The challenge of combining variable retention and prescribed burning in
1249 Finland, *Ecological Processes*, 9, 4, <https://doi.org/10.1186/s13717-019-0207-3>, 2020.

1250 Little, J. M., Jandt, R. R., Drury, S., Molina, A., and Lane, B.: Evaluating the effectiveness of fuel treatments in Alaska-Final
1251 Report to the Joint Fire Science Program, JFSP Project No. 14-5-01-27, University of Alaska-Fairbanks, available at:
1252 <https://www.fs.fed.us/psw/pubs/58856>, 2018.

1253 Liu, T., Mickley, L. J., Marlier, M. E., DeFries, R. S., Khan, M. F., Latif, M. T., and Karambelas, A.: Diagnosing spatial biases
1254 and uncertainties in global fire emissions inventories: Indonesia as regional case study, *Remote Sens. Environ.*, 237, 111557,
1255 <https://doi.org/10.31223/osf.io/nh57j>, 2020.

1256 Loboda, T. V., Hall, J.V., Hall, A.H., and Shevade, V.S.: ABoVE: Cumulative Annual Burned Area, Circumpolar High
1257 Northern Latitudes, 2001-2015, available at: https://daac.ornl.gov/cgi-bin/dsvviewer.pl?ds_id=1526, ORNL DAAC, Oak
1258 Ridge, TN, USA, <https://doi.org/10.3334/ORNLDAAC/1526>, 2017.

1259 Loepfe, L., Lloret, F., and Román-Cuesta, R. M.: Comparison of burnt area estimates derived from satellite products and
1260 national statistics in Europe, *Int J Remote Sens.*, 33, 3653, <https://doi.org/10.1080/01431161.2011.631950>, 2012.

1261 [Loisel, J., Gallego-Sala, A.V., Amesbury, M.J., Magnan, G., Anshari, G., Beilman, D.W., Benavides, J.C., Blewett, J., Camill,](#)
1262 [P., Charman, D.J. and Chawchai, S.: Expert assessment of future vulnerability of the global peatland carbon sink, *Nat. Clim.*](#)
1263 [Chang., 11, 70–77, <https://doi.org/10.1038/s41558-020-00944-0>, 2021.](#)

1264 Loranty, M. M., Lieberman-Cribbin, W., Berner, L. T., Natali, S. M., Goetz, S. J., Alexander, H. D., and Kholodov, A. L.:
1265 Spatial variation in vegetation productivity trends, fire disturbance, and soil carbon across arctic-boreal permafrost ecosystems,
1266 *Environ. Res. Lett.*, 11, 095008, <https://doi.org/10.1088/1748-9326/11/9/095008>, 2016.

1267 Malevsky-Malevich, S.P., Molkentin, E.K., Nadyozhina, E.D. and Shklyarevich, O.B.: Numerical simulation of permafrost
1268 parameters distribution in Russia, *Cold Reg Sci Technol*, 32, 1-11, [https://doi.org/10.1016/s0165-232x\(01\)00018-0](https://doi.org/10.1016/s0165-232x(01)00018-0), 2001.

1269 Markuse, P.: Before/After Comparison of the July/August 2019 Greenland Wildfire: Analysis from Sentinel-2, available at:
1270 <https://pierre-markuse.net/2019/08/19/before-after-comparison-of-the-july-august-2019-greenland-wildfire/>, 2019.

1271 McCarty, J. L., Krylov, A., Prishchepov, A. V., Banach, D. M., Tyukavina, A., Potapov, P., and Turubanova, S.: Agricultural
1272 fires in European Russia, Belarus, and Lithuania and their impact on air quality, 2002–2012, In *Land-Cover and Land-Use*
1273 *Changes in Eastern Europe after the Collapse of the Soviet Union in 1991*, Springer, 2017.

1274 McCarty, J. L., Smith, T. E., and Turetsky, M. R.: Arctic fires re-emerging, *Nat. Geosci.*, 13, 658,
1275 <https://doi.org/10.1038/s41561-020-00645-5>, 2020.

1276 McGwinn, K.: Hikers warned as Greenland wildfire burns out of control, Arctic Today, available at:
1277 <https://www.arctictoday.com/hikers-warned-as-greenland-wildfire-burns-out-of-control/>, 2019.

1278 McWethy, D. B., Schoennagel, T., Higuera, P. E., Krawchuk, M., Harvey, B. J., Metcalf, E. C., Schultz, C., Miller, C., Metcalf,
1279 A.L., Buma, B. and Virapongse, A.: Rethinking resilience to wildfire, Nat. Sustain, 2, 797, [https://doi.org/10.1038/s41893-](https://doi.org/10.1038/s41893-019-0353-8)
1280 019-0353-8, 2019.

1281 Mekonnen, Z.A., Riley, W.J., Randerson, J.T., Grant, R.F., and Rogers, B.M.: Expansion of high-latitude deciduous forests
1282 driven by interactions between climate warming and fire, Nat. Plants, 5, 952, <https://doi.org/10.1038/s41477-019-0495-8>,
1283 2019.

1284 [Melekhov, I.S.: Forest Science, Moscow \(in Russian\), 1980.](#)

1285 Michaelides, R. J., Schaefer, K., Zebker, H. A., Parsekian, A., Liu, L., Chen, J., Natali, S., Ludwig, S. and Schaefer, S.R.:
1286 Inference of the impact of wildfire on permafrost and active layer thickness in a discontinuous permafrost region using the
1287 remotely sensed active layer thickness (ReSALT) algorithm, Environ. Res. Lett., 14, 035007, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/aaf932)
1288 9326/aaf932, 2019.

1289 Mieville, A., Granier, C., Lioussé, C., Guillaume, B., Mouillot, F., Lamarque, J.F., Grégoire, J.M. and Pétron, G.: Emissions
1290 of gases and particles from biomass burning during the 20th century using satellite data and an historical reconstruction, Atmos.
1291 Environ., 44, 1469, <https://doi.org/10.1016/j.atmosenv.2010.01.011>, 2010.

1292 Miller, M. E., Billmire, M., Bourgeau-Chavez, L., Elliot, W. J., Robichaud, P. R., and MacDonald, L.: Rapid response tools
1293 and datasets for post-fire modeling in Boreal and Arctic Environments, Spring 2017 AFSC Remote Sensing Workshop:
1294 Opportunities to Apply Remote Sensing in Boreal/Arctic Wildfire Management and Science, available at:
1295 https://digitalcommons.mtu.edu/mtri_p/290, 2017.

1296 Mölders, N., and Kramm, G.: Climatology of Air Quality in Arctic Cities—Inventory and Assessment, Open Journal of Air
1297 Pollution, 7(1), 48-93, <https://doi.org/10.4236/ojap.2018.71004>, 2018.

1298 Molinari, C., Lehsten, V., Blarquez, O., Carcaillet, C., Davis, B.A., Kaplan, J.O., Clear, J. and Bradshaw, R.H.: The climate,
1299 the fuel and the land use: Long-term regional variability of biomass burning in boreal forests, Glob Chang Biol., 24, 4929-
1300 4945, <https://doi.org/10.1111/gcb.14380>, 2018.

1301 [Monks, S. A., Arnold, S.R., and Chipperfield, M.P.: Evidence for El Niño-Southern Oscillation \(ENSO\) influence on Arctic](#)
1302 [CO interannual variability through biomass burning emissions, Geophys. Res. Lett., 39, L14804, doi:10.1029/2012GL052512,](#)
1303 [2012.](#)

1304 Montesano, P.M., Neigh, C.S., Macander, M., Feng, M. and Noojipady, P.: The bioclimatic extent and pattern of the cold edge
1305 of the boreal forest: the circumpolar taiga-tundra ecotone, Environ. Res. Lett., 15, 105019, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/abb2c7)
1306 [9326/abb2c7](https://doi.org/10.1088/1748-9326/abb2c7), 2020.

1307 Morin, P., Porter, C., Cloutier, M., Howat, I., Noh, M.J., Willis, M., Bates, B., Willamson, C. and Peterman, K.: ArcticDEM:
1308 a [publicly](#) available, high resolution elevation model of the Arctic, available at:
1309 <https://livingatlas2.arcgis.com/arcticdemexplorer/>, 2016.

1310 Mottershead, K.D., McGee, T.K. and Christianson, A.: Evacuating a First Nation Due to Wildfire Smoke: The Case of Dene
1311 Tha' First Nation, Int J Disaster Risk Sci 11, 274, <https://doi.org/10.1007/s13753-020-00281-y>, 2020.

1312 NIFC: National Interagency Fire Center: Total Wildland Fires and Acres (1926-2019), available at:
1313 https://www.nifc.gov/fireInfo/fireInfo_stats_totalFires.html, 2019.

1314 Nikolakis, W., Roberts, E., Hotte, N. and Ross, R.M.: Goal setting and Indigenous fire management: a holistic perspective, Int
1315 J Wildland Fire, 29, 974, <https://doi.org/10.1071/WF20007>, 2020.

1316 Nitzbon, J., Westermann, S., Langer, M., Martin, L. C., Strauss, J., Laboor, S., and Boike, J.: Fast response of cold ice-rich
1317 permafrost in northeast Siberia to a warming climate, Nat Commun, 11, 2201, <https://doi.org/10.1038/s41467-020-15725-8>,
1318 2020.

1319 Nordregio.: Indigenous population in the Arctic, available at: <https://nordregio.org/maps/indigenous-population-in-the-arctic/>,
1320 2019.

1321 Nugent, K. A., Strachan, I. B., Roulet, N. T., Strack, M., Frohling, S., and Helbig, M.: Prompt active restoration of peatlands
1322 substantially reduces climate impact, Environ. Res. Lett., 14, 124030, <https://doi.org/10.1088/1748-9326/ab56e6>, 2019.

1323 Oliva, P. and Schroeder, W.: Assessment of VIIRS 375 m active fire detection product for direct burned area mapping, Remote
1324 Sens. Environ., 160, 144, <https://doi.org/10.1016/j.rse.2015.01.010>, 2015.

1325 Overland, J. E., and Wang, M.: The 2020 Siberian Heat Wave, Int J Climatol, <https://doi.org/10.1002/joc.6850>, 2020.

1326 [Päätaalo, M.-L.: Factors influencing occurrence and impacts of fires in northern European forests, *Silva Fenn.*, 32, 185-202,](#)
1327 <https://doi.org/10.14214/sf.695>, 1998.

1328 Pan, X., Ichoku, C., Chin, M., Bian, H., Darmenov, A., Colarco, P., Ellison, L., Kucsera, T., da Silva, A., Wang, J., Oda, T.,
1329 and Cui, G.: Six global biomass burning emission datasets: intercomparison and application in one global aerosol model,
1330 *Atmos. Chem. Phys.*, 20, 969–994, <https://doi.org/10.5194/acp-20-969-2020>, 2020.

1331 Parfenova, E., Tchebakova, N., and Soja, A.: Assessing landscape potential for human sustainability and ‘attractiveness’ across
1332 Asian Russia in a warmer 21st century, *Environ. Res. Lett.*, 14, 065004, <https://doi.org/10.1088/1748-9326/ab10a8>, 2019.

1333 Parisien, M. A., Miller, C., Parks, S. A., DeLancey, E. R., Robinne, F. N., and Flannigan, M. D.: The spatially varying influence
1334 of humans on fire probability in North America, *Environ. Res. Lett.*, 11, 075005, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/11/7/075005)
1335 [9326/11/7/075005](https://doi.org/10.1088/1748-9326/11/7/075005), 2016.

1336 Parisien, M.A., Barber, Q.E., Hirsch, K.G., Stockdale, C.A., Erni, S., Wang, X., Arseneault, D. and Parks, S.A.: Fire deficit
1337 increases wildfire risk for many communities in the Canadian boreal forest, *Nat Commun*, 11, 2121,
1338 <https://doi.org/10.1038/s41467-020-15961-y>, 2020.

1339 Partain Jr, J.L., Alden, S., Strader, H., Bhatt, U.S., Bieniek, P.A., Bretschneider, B.R., Walsh, J.E., Lader, R.T., Olsson, P.Q.,
1340 Rupp, T.S., and Thoman Jr, R.L.: An assessment of the role of anthropogenic climate change in the Alaska fire season of
1341 2015, *Bull Am Meteorol Soc*, 97, S14, <https://doi.org/10.1175/bams-d-16-0149.1>, 2016.

1342 Peltola, H., Kilpeläinen, A., and Kellomäki, S.: Diameter growth of Scots pine (*Pinus sylvestris*) trees grown at elevated
1343 temperature and carbon dioxide concentration under boreal conditions, *Tree Physiol*, 22, 963-972,
1344 <https://doi.org/10.1093/treephys/22.14.963>, 2002.

1345 ФБУ «Авиалесоохрана»: Сведения о лесопожарной обстановке на территории субъектов РФ на 31.12, available at:
1346 https://aviales.ru/files/documents/2019/fds_svedeniya/%D1%81%D0%B2%D0%B5%D0%B4%D0%B5%D0%BD%D0%B8%D1%8F%20%D0%BE%20%D0%BB%D0%B5%D1%81%D0%BE%D0%BF%D0%BE%D0%B6%D0%B0%D1%80%D0%BD%D0%BE%D0%B9%20%D0%BE%D0%B1%D1%81%D1%82%D0%B0%D0%BD%D0%BE%D0%B2%D0%BA%D0%B5%20%D0%BD%D0%B0%20%D1%82%D0%B5%D1%80%D1%80%D0%B8%D1%82%D0%BE%D1%80%D0%B8%D0%B8%20%D1%81%D1%83%D0%B1%D1%8A%D0%B5%D0%BA%D1%82%D0%BE%D0%B2%20%D1%80%D1%84%20%D0%BD%D0%B0%2031.12.2019.pdf, 2019.

- 1352 Pickell, P. D., Coops, N. C., Ferster, C. J., Bater, C. W., Blouin, K. D., Flannigan, M. D., and Zhang, J.: An early warning
1353 system to forecast the close of the spring burning window from satellite-observed greenness, *Scientific Rep*, 7, 1-10,
1354 <https://doi.org/10.1038/s41598-017-14730-0>, 2017.
- 1355 Pimentel, R., and Arheimer, B.: Hydrological impacts of a wildfire in a Boreal region: The Västmanland fire 2014 (Sweden),
1356 *Sci. Total Environ.* 756, 143519, <https://doi.org/10.1016/j.scitotenv.2020.143519>, 2021.
- 1357 Polikarpov, N.P., Andreeva, N.M., Nazimova, D.I., Sirotinina, A.V. and Sofronov, M.A.: Formation composition of the forest
1358 zones in Siberia as a reflection of forest-forming tree species interrelations, *Russ J For Sci*, 5, 3-11, 1998.
- 1359 [Prat-Guitart, N., Rein, G., Hadden, R.M., Belcher, C.M. and Yearsley, J.M.: Propagation probability and spread rates of self-](#)
1360 [sustained smouldering fires under controlled moisture content and bulk density conditions, *Int J Wildland Fire*, 25, 456-465,](#)
1361 <https://doi.org/10.1071/WF15103>, 2016.
- 1362 [Prishchepov, A.V., Schierhorn, F., Dronin, N., Ponkina, E.V., and Müller, D.: 800 Years of Agricultural Land-use Change in](#)
1363 [Asian \(Eastern\) Russia, in: KULUNDA: Climate Smart Agriculture, Innovations in Landscape Research, edited by: Frühauf,](#)
1364 [M., Guggenberger, G., Meinel, T., Theesfeld, I., Lentz, S., Springer, Cham, Switzerland, 67-87, \[https://doi.org/10.1007/978-\]\(https://doi.org/10.1007/978-3-030-15927-6_6\)](#)
1365 [3-030-15927-6_6](#), 2020.
- 1366 Púčik, T., Groenemeijer, P., Rädler, A.T., Tijssen, L., Nikulin, G., Prein, A.F., van Meijgaard, E., Fealy, R., Jacob, D., and
1367 Teichmann, C.: Future changes in European severe convection environments in a regional climate model ensemble, *J Clim*,
1368 30, 6771, <https://doi.org/10.1175/jcli-d-16-0777.1>, 2017.
- 1369 Pureswaran, D. S., Roques, A., and Battisti, A.: Forest insects and climate change, *Curr. For. Rep.*, 4,
1370 35, <https://doi.org/10.1007/s40725-018-0075-6>, 2018.
- 1371 Qi, L., and Wang, S.: Sources of black carbon in the atmosphere and in snow in the Arctic, *Sci Total Environ*, 691, 442-454,
1372 <https://doi.org/10.1016/j.scitotenv.2019.07.073>, 2019.
- 1373 Raynolds, M.K., Walker, D.A., Balsler, A., Bay, C., Campbell, M., Cherosov, M.M., Daniëls, F.J.A., Eidesen, P.B., Ermokhina,
1374 K.A., Frost, G.V., Jedrzejek, B., Jorgenson, M.T., Kennedy, B.E., Kholod, S.S., Lavrinenko, I.A., Lavrinenko, O.V.,
1375 Magnússon, B., Matveyeva, N.V., Metúsalemsson, S., Nilsen, L., Olthof, I., Pospelov, I.N., Pospelova, E.B., Pouliot, D.,
1376 Razzhivin, V., Schaepman-Strub, G., Šibík, J., Telyatnikov, M.Y., Troeva, E.: A raster version of the Circumpolar Arctic
1377 Vegetation Map (CAVM), *Remote Sens Environ*, 232, 111297, <https://doi.org/10.1016/j.rse.2019.111297>, 2019.

- 1378 Rein, G., Cleaver, N., Ashton, C., Pironi, P., and Torero, J. L.: The severity of smouldering peat fires and damage to the forest
1379 soil. *Catena*, 74(3), 304-309, <https://doi.org/10.1016/j.catena.2008.05.008>, 2008.
- 1380 Rein, G.: Smoldering combustion, in: *SFPE Handbook of Fire Protection Engineering*, edited by: Hurley, M.J., Gottuk, D.,
1381 Hall, J.R., Harada, K., Kuligowski, E., Puchovsky, M., Torero, J., Watts, J.M., and C. Wieczoreks, C., Springer New York,
1382 New York, New York, 581-603, https://doi.org/10.1007/978-1-4939-2565-0_19, 2016.
- 1383 Rémy, S., Veira, A., Paugam, R., Sofiev, M., Kaiser, J. W., Marengo, F., Burton, S. P., Benedetti, A., Engelen, R. J., Ferrare,
1384 R., and Hair, J. W.: Two global data sets of daily fire emission injection heights since 2003, *Atmos. Chem. Phys.*, 17, 2921-
1385 2942, <https://doi.org/10.5194/acp-17-2921-2017>, 2017.
- 1386 ~~Reyer, C.P., Bathgate, S., Blennow, K., Borges, J.G., Bugmann, H., Delzon, S., Faias, S.P., Garcia-Gonzalo, J., Gardiner, B.,~~
1387 ~~Gonzalez-Olabarria, J.R. and Gracia, C.: Are forest disturbances amplifying or canceling out climate change induced~~
1388 ~~productivity changes in European forests?, *Environ. Res. Lett.*, 12, 034027, <https://doi.org/10.1088/1748-9326/aa5ef1>, 2017.~~
- 1389 Riley, K. L., Williams, A. P., Urbanski, S. P., Calkin, D. E., Short, K. C., and O'Connor, C. D.: Will Landscape Fire Increase
1390 in the Future? A Systems Approach to Climate, Fire, Fuel, and Human Drivers. *Curr. Pollut. Rep.*, 5, 9,
1391 <https://doi.org/10.1007/s40726-019-0103-6>, 2019.
- 1392 Robinne, F. N., Parisien, M. A., and Flannigan, M.: Anthropogenic influence on wildfire activity in Alberta, Canada, *Int J*
1393 *Wildland Fire*, 25(11), 1131-1143, <https://doi.org/10.1071/wf16058>, 2016.
- 1394 Robinne, F. N., Hallema, D. W., Bladon, K. D., and Buttle, J. M.: Wildfire impacts on hydrologic ecosystem services in North
1395 American high-latitude forests: A scoping review, *J Hydrol.*, 581, 124360, <https://doi.org/10.1016/j.jhydrol.2019.124360>,
1396 2020.
- 1397 Rocha, A.V., Loranty, M.M., Higuera, P.E., Mack, M.C., Hu, F.S., Jones, B.M., Breen, A.L., Rastetter, E.B., Goetz, S.J. and
1398 Shaver, G.R.: The footprint of Alaskan tundra fires during the past half-century: implications for surface properties and
1399 radiative forcing, *Environ. Res. Lett.*, 7, 044039, <https://doi.org/10.1088/1748-9326/7/4/044039>, 2012.
- 1400 Rogers, B.M., Veraverbeke, S., Azzari, G., Czimczik, C.I., Holden, S.R., Mouteva, G.O., Sedano, F., Treseder, K.K. and
1401 Randerson, J.T.: Quantifying fire-wide carbon emissions in interior Alaska using field measurements and Landsat imagery, *J.*
1402 *of Geophys. Res. Biogeosci.*, 119, 1608, <https://doi.org/10.1002/2014JG002657>, 2014.
- 1403 Rogers, B. M., Soja, A.J., Goulden, M.L., and Randerson, J.T.: Influence of tree species on continental differences in boreal
1404 fires and climate feedbacks, *Nature Geosci.*, 8, 228, <https://doi.org/10.1038/ngeo2352>, 2015.

1405 Rogers, B.M., Balch, J.K., Goetz, S.J., Lehmann, C.E. and Turetsky, M.: Focus on changing fire regimes: interactions with
1406 climate, ecosystems, and society, *Environ. Res. Lett.*, 15, p.030201, <https://doi.org/10.1088/1748-9326/ab6d3a>, 2020.

1407 Ronkainen, T., Välranta, M., and Tuittila, E.-S.: Fire pattern in a drainage-affected boreal bog, *Boreal Environ Res*, 18, 309,
1408 2013.

1409 [Santoso, M.A., Cui, W., Amin, H.M., Christensen, E.G., Nugroho, Y.S. and Rein, G.: Laboratory study on the suppression of](#)
1410 [smouldering peat wildfires: effects of flow rate and wetting agent, *Int J Wildland Fire*, 30, 378-390,](#)
1411 <https://doi.org/10.1071/WF20117>, 2021.

1412 Schmale, J., Arnold, S.R., Law, K.S., Thorp, T., Anenberg, S., Simpson, W.R., Mao, J., and Pratt, K.A.: Local Arctic air
1413 pollution: A neglected but serious problem, *Earth's Future*, 6, 1385, <https://doi.org/10.1029/2018EF000952>, 2018.

1414 Scholten, R., and Veraverbeke, S.: Alaska Fire Science Consortium: Spatiotemporal patterns of overwintering fire in Alaska,
1415 available at: https://akfireconsortium.files.wordpress.com/2020/03/fsh_2020mar25_holdoverfires-1.pdf, 2020.

1416 Seidl, R., Schelhaas, M. J., Rammer, W., and Verkerk, P. J.: Increasing forest disturbances in Europe and their impact on
1417 carbon storage, *Nature Clim. Chang.*, 4, 806-810, <https://doi.org/10.1038/nclimate2318>, 2014.

1418 Sherstyukov, B. G., and Sherstyukov, A. B.: Assessment of increase in forest fire risk in Russia till the late 21st century based
1419 on scenario experiments with fifth-generation climate models, *Russ Meteorol Hydro+*, 39, 292,
1420 <https://doi.org/10.3103/s1068373914050021>, 2014.

1421 Shiwakoti, S., Zheljazkov, V. D., Gollany, H. T., Kleber, M., Xing, B., and Astatkie, T. (2019). Micronutrients in the Soil and
1422 Wheat: Impact of 84 Years of Organic or Synthetic Fertilization and Crop Residue Management, *Agronomy*, 9, 464,
1423 <https://doi.org/10.3390/agronomy9080464>, 2019.

1424 [Shugart, H.H., Leemans, R. and Bonan, G.B.: A Systems Analysis of the Global Boreal Forest, Cambridge University Press,](#)
1425 [New York, USA, 1-565, https://doi.org/10.1017/CBO9780511565489.022, 1992.](#)

1426 Shuman, J.K., Foster, A.C., Shugart, H.H., Hoffman-Hall, A., Krylov, A., Loboda, T., Ershov, D. and Sochilova, E.: Fire
1427 disturbance and climate change: implications for Russian forests, *Environ. Res. Lett.*, 12, 035003,
1428 <https://doi.org/10.1088/1748-9326/aa5eed>, 2017.

1429 Shumilova, L.V.: Botanical Geography of Siberia, Tomsk University Press, Tomsk, USSR, 1962.

- 1430 Shvetsov, E. G., Kukavskaya, E. A., Buryak, L. V., and Barrett, K.: Assessment of post-fire vegetation recovery in Southern
1431 Siberia using remote sensing observations, *Environ. Res. Lett.*, 14, 055001, <https://doi.org/10.1088/1748-9326/ab083d>, 2019.
- 1432 [Shvidenko, A.Z., and Nilsson, S.: Extent, distribution, and ecological role of fire in Russian forests, in: Fire, climate change,
1433 and carbon cycling in the boreal forest, edited by Kasischke E.S., and Stocks B.J., Springer, New York, NY, 132-150,
1434 \[https://doi.org/10.1007/978-0-387-21629-4_16\]\(https://doi.org/10.1007/978-0-387-21629-4_16\), 2000.](#)
- 1435 Sidorova, E. J.: The incorporation of Traditional Ecological Knowledge in the Arctic Council: Lip service?, *Polar Rec (Gr
1436 Brit)*, 56, e28, <https://doi.org/10.1017/S0032247420000273>, 2020.
- 1437 Silva, J. S., and Harrison, S. P.: Humans, Climate and Land Cover as Controls on European Fire Regimes, in *Towards
1438 integrated fire management-Outcomes of the European Project Fire Paradox*, edited by: Silva, J.S., Rego, F.C., Fernandes, P.,
1439 and Rigolot, E. European Forest Institute, Joensuu, Finland, 49-59, 2010.
- 1440 Sirin, A., Maslov, A., Medvedeva, M., Vozbrannaya, A., Valyaeva, N., Tsyganova, O., Glukhova, T., and Makarov, D.:
1441 Multispectral Remote Sensing Data as a Tool for Assessing the Need and the Effectiveness for Peatland Restoration, In
1442 *Proceedings of the 9th European Conference on Ecological Restoration*, edited by: Tolvanen, A., Hekkala, A.M., Finnish
1443 Forest Research Institute, Oulu, Finland, 133, 2014.
- 1444 Sirin, A., Medvedeva, M., Maslov, A., and Vozbrannaya, A.: Assessing the Land and Vegetation Cover of Abandoned Fire
1445 Hazardous and Rewetted Peatlands: Comparing Different Multispectral Satellite Data, *Land*, 7, 71,
1446 <https://doi.org/10.3390/land7020071>, 2018.
- 1447 Sizov, O., Ezhova, E., Tsymbarevich, P., Soromotin, A., Prihod'ko, N., Petäjä, T., Zilitinkevich, S., Kulmala, M., Bäck, J., and
1448 Köster, K.: Fire and vegetation dynamics in northwest Siberia during the last 60 years based on high-resolution remote sensing,
1449 *Biogeosciences*, 18, 207–228, <https://doi.org/10.5194/bg-18-207-2021>, 2021.
- 1450 Sjöström, J., Plathner, F. V., and Granström, A.: Wildfire ignition from forestry machines in boreal Sweden, *Int J Wildland
1451 Fire*, 28, 666, <https://doi.org/10.1071/wf18229>, 2019.
- 1452 Smirnov, N. S., Korotkov, V. N., and Romanovskaya, A. A.: Black carbon emissions from wildfires on forest lands of the
1453 Russian Federation in 2007–2012, *Russ Meteorol Hydro+*, 40, 435, <https://doi.org/10.3103/s1068373915070018>, 2015.
- 1454 Sofronov, M.A., Volokitina, A.V. and Shvidenko, A.Z.: Wildland fires in the north of Central Siberia, *Commonw. For. Rev.*,
1455 77, 124, 1998.

1456 Sofronov, M. A., Volokitina, A., Kajimoto, T., Matsuura, Y., and Uemura, S.: Zonal peculiarities of forest vegetation
1457 controlled by fires in northern Siberia, *Eurasian Journal of Forest Research*, 1, 51, 2000.

1458 Sofronov, M.A., and Volokitina A.V.: Wildfire Ecology in Continuous Permafrost Zone. In *Permafrost Ecosystems, Ecological
1459 Studies (Analysis and Synthesis)*, vol 209, edited by: Osawa, A., Zyryanova, O., Matsuura, Y., Kajimoto, T., and Wein, R.,
1460 Springer, Dordrecht, Netherlands, 59, https://doi.org/10.1007/978-1-4020-9693-8_4, 2010.

1461 Soja, A.J., Cofer, W.R., Shugart, H.H., Sukhinin, A.I., Stackhouse, P.W., McRae, D.J. and Conard, S.G.: Estimating fire
1462 emissions and disparities in boreal Siberia (1998–2002), *J Geophys Res - Atmos*, 109, <https://doi.org/10.1029/2004JD004570>,
1463 2004a.

1464 Soja, A. J., Sukhinin, A.I., Cahoon Jr., D.R., Shugart, H.H., and Stackhouse Jr, P.W.: AVHRR-derived fire frequency,
1465 distribution and area burned in Siberia, *Int. Journal of Remote Sens.*, 25, 1939, <https://doi:10.1080/01431160310001609725>,
1466 2004b.

1467 [Soja, A.J., Shugart, H.H., Sukhinin, A., Conard, S., and Stackhouse Jr., P.W.: Satellite-Derived Mean Fire Return Intervals As
1468 Indicators Of Change In Siberia \(1995–2002\), *Mitig Adapt Strat Glob Change*, 11, 75–96, \[https://doi.org/10.1007/s11027-
1469 006-1009-3\]\(https://doi.org/10.1007/s11027-006-1009-3\), 2006.](#)

1470 Sokolik, I. N., Soja, A. J., DeMott, P. J., and Winker, D.: Progress and Challenges in Quantifying Wildfire Smoke Emissions,
1471 Their Properties, Transport, and Atmospheric Impacts, *J. Geophys. Res. Atmos.*, 124, 13005,
1472 <https://doi.org/10.1029/2018JD029878>, 2019.

1473 Sommers, W.T., Loehman, R.A. and Hardy, C.C.: Wildland fire emissions, carbon, and climate: Science overview and
1474 knowledge needs, *For. Ecol. Manage.*, 317, 1, <https://doi.org/10.1016/j.foreco.2013.12.014>, 2014.

1475 Stralberg, D., Wang, X., Parisien, M.A., Robinne, F.N., Sólomos, P., Mahon, C.L., Nielsen, S.E., and Bayne, E.M.: Wildfire-
1476 mediated vegetation change in boreal forests of Alberta, Canada. *Ecosphere*, 9, e02156, <https://doi.org/10.1002/ecs2.2156>,
1477 2018.

1478 Stone, R.: As the Arctic thaws, Indigenous Alaskans demand a voice in climate change research, *Science Magazine*, available
1479 at: <https://www.sciencemag.org/news/2020/09/arctic-thaws-indigenous-alaskans-demand-voice-climate-change-research#>,
1480 <https://doi.org/10.1126/science.abe7149>, 2020.

1481 [Stroeve, J.C., Markus, T., Boisvert, L., Miller, J. and Barrett, A.: Changes in Arctic melt season and implications for sea ice
1482 loss, *Geophys. Res. Lett.*, 41, 1216-1225, <https://doi.org/10.1002/2013GL058951>, 2014.](#)

1483 Tchebakova, N.M., Parfenova, E. and Soja, A.J.: The effects of climate, permafrost and fire on vegetation change in Siberia
1484 in a changing climate, *Environ. Res. Lett.*, 4, 045013, <https://doi.org/10.1088/1748-9326/4/4/045013>, 2009.

1485 Tchebakova, N.M., Rehfeldt, G.E., and Parfenova, E.: From Vegetation Zones to Climates: Effects of Climate Warming on
1486 Siberian Ecosystems, in: *Permafrost Ecosystems, Ecological Studies (Analysis and Synthesis)*, edited by Osawa, A.,
1487 Zyryanova, O., Matsuura, Y., Kajimoto, T., and Wein, R., Springer, Dordrecht, Germany, 427-446,
1488 https://doi.org/10.1007/978-1-4020-9693-8_22, 2010.

1489 Tchebakova, N.M., Parfenova, E.I., Lysanova, G.I., and Soja, A.J.: Agroclimatic potential across central Siberia in an altered
1490 twenty-first century, *Environ. Res. Lett.*, 6, 045207, <https://doi.org/10.1088/1748-9326/6/4/045207>, 2011.

1491 Tchebakova, N. M., Chuprova, V. V., Parfenova, E. I., Soja, A. J., and Lysanova, G. I.: Evaluating the agroclimatic potential
1492 of Central Siberia. In: *Novel Methods for Monitoring and Managing Land and Water Resources in Siberia*, edited by: Mueller
1493 L., Sheudshen A., and Eulenstein F. Springer, Cham, https://doi.org/10.1007/978-3-319-24409-9_10, 2016.

1494 Terrier, A., Girardin, M. P., Périé, C., Legendre, P., and Bergeron, Y.: Potential changes in forest composition could reduce
1495 impacts of climate change on boreal wildfires, *Ecol Appl*, 23, 21, <https://doi.org/10.1890/12-0425.1>, 2013.

1496 Teufel, B., and Sushama, L.: Abrupt changes across the Arctic permafrost region endanger northern development, *Nat. Clim.*
1497 *Chang.*, 9, 858, <https://doi.org/10.1038/s41558-019-0614-6>, 2019.

1498 Theesfeld, I., and Jelinek, L.: A misfit in policy to protect Russia's black soil region. An institutional analytical lens applied
1499 to the ban on burning of crop residues. *Land use policy*, 67, 517, <https://doi.org/10.1016/j.landusepol.2017.06.018>, 2017.

1500 Thomas, J.L., Polashenski, C.M., Soja, A.J., Marelle, L., Casey, K.A., Choi, H.D., Raut, J.C., Wiedinmyer, C., Emmons, L.K.,
1501 Fast, J.D. and Pelon, J.: Quantifying black carbon deposition over the Greenland ice sheet from forest fires in Canada, *Geophys.*
1502 *Res. Lett.*, 44, 7965, <https://doi.org/10.1002/2017gl073701>, 2017.

1503 Thompson, D. K. and Morrison, K.: A classification scheme to determine wildfires from the satellite record in the cool
1504 grasslands of southern Canada: considerations for fire occurrence modelling and warning criteria, *Nat. Hazards Earth Syst.*
1505 *Sci.*, 20, 3439–3454, <https://doi.org/10.5194/nhess-20-3439-2020>, 2020.

1506 Thompson, D. K., Simpson, B. N., Whitman, E., Barber, Q. E., and Parisien, M. A.: Peatland hydrological dynamics as a driver
1507 of landscape connectivity and fire activity in the boreal plain of Canada, *Forests*, 10, 534, <https://doi.org/10.3390/f10070534>,
1508 2019.

1509 Tymstra, C., Stocks, B. J., Cai, X., and Flannigan, M. D.: Wildfire management in Canada: Review, challenges and
1510 opportunities. *Progress in Disaster Science*, 5, 100045, <https://doi.org/10.1016/j.pdisas.2019.100045>, 2020.

1511 Val Martin, M., Kahn, R. A., and Tosca, M. G.: A Global Analysis of Wildfire Smoke Injection Heights Derived from Space-
1512 Based Multi-Angle Imaging, *Remote Sens.*, 10, 1609, <https://doi.org/10.3390/rs10101609>, 2018.

1513 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability
1514 in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, [https://doi.org/10.5194/acp-6-](https://doi.org/10.5194/acp-6-3423-2006)
1515 [3423-2006](https://doi.org/10.5194/acp-6-3423-2006), 2006.

1516 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin,
1517 Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat
1518 fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.

1519 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E.,
1520 Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth*
1521 *Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.

1522 van Leeuwen, T. T., van der Werf, G. R., Hoffmann, A. A., Detmers, R. G., Rücker, G., French, N. H. F., Archibald, S.,
1523 Carvalho Jr., J. A., Cook, G. D., de Groot, W. J., Hély, C., Kasischke, E. S., Kloster, S., McCarty, J. L., Pettinari, M. L.,
1524 Savadogo, P., Alvarado, E. C., Boschetti, L., Manuri, S., Meyer, C. P., Siegert, F., Trollope, L. A., and Trollope, W. S. W.:
1525 Biomass burning fuel consumption rates: a field measurement database, *Biogeosciences*, 11, 7305–7329,
1526 <https://doi.org/10.5194/bg-11-7305-2014>, 2014.

1527 Veira, A., Lasslop, G., and Kloster, S.: Wildfires in a warmer climate: emission fluxes, emission heights, and black carbon
1528 concentrations in 2090–2099, *J. Geophys. Res. Atmos.*, 121, 3195, <https://doi.org/10.1002/2015jd024142>, 2016.

1529 Venäläinen, A., Lehtonen, I., Laapas, M., Ruosteenoja, K., Tikkanen, O.P., Viiri, H., Ikonen, V.P. and Peltola, H.: Climate
1530 change induces multiple risks to boreal forests and forestry in Finland: A literature review, *Glob Chang Biol.*, 26, 4178,
1531 <https://doi.org/10.1111/gcb.15183>, 2020.

1532 Veraverbeke, S., Rogers, B. M., Goulden, M. L., Jandt, R. R., Miller, C. E., Wiggins, E. B., and Randerson, J. T.: Lightning
1533 as a major driver of recent large fire years in North American boreal forests, *Nature Clim Change*, 7, 529,
1534 <https://doi.org/10.1038/nclimate3329>, 2017.

1535 [Voulgarakis, A., Marlier, M. E., Faluvegi, G., Shindell, D. T., Tsigaridis, K., and Mangeon, S.: Interannual variability of](#)
1536 [tropospheric trace gases and aerosols: the role of biomass burning emissions, *J. Geophys. Res.-Atmos.*, 120, 7157–7173,](#)
1537 <https://doi.org/10.1002/2014JD022926>, 2015.

1538 Waigl, C. F., Stuefer, M., Prakash, A., and Ichoku, C.: Detecting high and low-intensity fires in Alaska using VIIRS I-band
1539 data: An improved operational approach for high latitudes, *Remote Sens. Environ.*, 199, 389,
1540 <https://doi.org/10.1016/j.rse.2017.07.003>, 2017.

1541 Waigl, C. F., Prakash, A., Stuefer, M., Verbyla, D., and Dennison, P.: Fire detection and temperature retrieval using EO-1
1542 Hyperion data over selected Alaskan boreal forest fires, *Int J Appl Earth Obs.*, 81, 72, <https://doi.org/10.1016/j.jag.2019.03.004>,
1543 2019.

1544 Walker, X.J., Rogers, B.M., Baltzer, J.L., Cumming, S.G., Day, N.J., Goetz, S.J., Johnstone, J.F., Schuur, E.A., Turetsky, M.R.
1545 and Mack, M.C.: Cross-scale controls on carbon emissions from boreal forest megafires, *Glob Chang Biol.*, 24,
1546 4251, <https://doi.org/10.1111/gcb.14287>, 2018.

1547 Walker, X.J., Baltzer, J.L., Cumming, S.G., Day, N.J., Ebert, C., Goetz, S., Johnstone, J.F., Potter, S., Rogers, B.M., Schuur,
1548 E.A. and Turetsky, M.R.: Increasing wildfires threaten historic carbon sink of boreal forest soils, *Nature*, 572, 520,
1549 <https://doi.org/10.1038/s41586-019-1474-y>, 2019.

1550 Walker, X.J., Rogers, B.M., Veraverbeke, S., Johnstone, J.F., Baltzer, J.L., Barrett, K., Bourgeau-Chavez, L., Day, N.J., de
1551 Groot, W.J., Dieleman, C.M. and Goetz, S.: Fuel availability not fire weather controls boreal wildfire severity and carbon
1552 emissions, *Nat. Clim. Chang.*, 10, 1130, <https://doi.org/10.1038/s41558-020-00920-8>, 2020.

1553 Walsh, J.E., Ballinger, T.J., Euskirchen, E.S., Hanna, E., Mård, J., Overland, J.E., Tangen, H. and Vihma, T.: Extreme weather
1554 and climate events in northern areas: A review. *Earth-Sci. Rev.*, 209, 103324, <https://doi.org/10.1016/j.earscirev.2020.103324>,
1555 2020.

1556 Wang, X., Parisien, M.A., Taylor, S.W., Candau, J.N., Stralberg, D., Marshall, G.A., Little, J.M. and Flannigan, M.D.:
1557 Projected changes in daily fire spread across Canada over the next century, *Environ. Res. Lett.*, 12, 025005,
1558 <https://doi.org/10.1088/1748-9326/aa5835>, 2017.

1559 Wang, J. A., Sulla-Menashe, D., Woodcock, C. E., Sonnentag, O., Keeling, R. F., and Friedl, M. A.: Extensive land cover
1560 change across Arctic–Boreal Northwestern North America from disturbance and climate forcing, *Glob Chang Biol.*, 26, 807,
1561 <https://doi.org/10.1111/gcb.14804>, 2019.

1562 Watson, J. G., Cao, J., Chen, L.-W. A., Wang, Q., Tian, J., Wang, X., Gronstal, S., Ho, S. S. H., Watts, A. C., and Chow, J.
1563 C.: Gaseous, PM_{2.5} mass, and speciated emission factors from laboratory chamber peat combustion, *Atmos. Chem. Phys.*, 19,
1564 14173–14193, <https://doi.org/10.5194/acp-19-14173-2019>, 2019.

1565 Wein, R. W.: Frequency and characteristics of arctic tundra fires, *Arctic*, 29, 213, <https://doi.org/10.14430/arctic2806>, 1978.

1566 Whitman, E., Parisien, M. A., Thompson, D. K., and Flannigan, M. D.: Short-interval wildfire and drought overwhelm boreal
1567 forest resilience, *Sci Rep*, 9, 18796, <https://doi.org/10.1038/s41598-019-55036-7>, 2019.

1568 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire
1569 INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model
1570 Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

1571 Wilkinson, S. L., Moore, P. A., Thompson, D. K., Wotton, B. M., Hvenegaard, S., Schroeder, D., and Waddington, J. M.: The
1572 effects of black spruce fuel management on surface fuel condition and peat burn severity in an experimental fire, *Can. J. For.
1573 Res.*, 48, 1433, <https://doi.org/10.1139/cjfr-2018-0217>, 2018.

1574 Wilkinson, S. L., Moore, P. A., and Waddington, J. M.: Assessing Drivers of Cross-Scale Variability in Peat Smoldering
1575 Combustion Vulnerability in Forested Boreal Peatlands. *Front. For. Glob. Change*, 2, 84,
1576 <https://doi.org/10.3389/ffgc.2019.00084>, 2019.

1577 Wilson, G. N.: Indigenous Internationalism in the Arctic, in: *The Palgrave Handbook of Arctic Policy and Politics*, edited by:
1578 Coates, K.S., and Holroyd, C., Palgrave Macmillan, Cham, Switzerland, 27-40, https://doi.org/10.1007/978-3-030-20557-7_3,
1579 2020.

1580 Witze, A.: The Arctic is burning like never before-and that's bad news for climate change. *Nature*, 585, 336,
1581 <https://doi.org/10.1038/d41586-020-02568-y>, 2020.

1582 Wotton, B. M., Martell, D. L., and Logan, K. A.: Climate change and people-caused forest fire occurrence in Ontario, *Clim.
1583 Change*, 60, 275, <https://doi.org/10.1023/A:1026075919710>, 2003.

1584 Wotton, B. M., Flannigan, M. D., and Marshall, G. A.: Potential climate change impacts on fire intensity and key wildfire
1585 suppression thresholds in Canada, *Environ. Res. Lett.*, 12, 095003, <https://doi.org/10.1088/1748-9326/aa7e6e>, 2017.

1586 Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-
1587 analysis, *Catena*, 160, 134, <https://doi.org/10.1016/j.catena.2017.09.010>, 2018.

- 1588 Xu, W., He, H. S., Hawbaker, T. J., Zhu, Z., and Henne, P. D.: Estimating burn severity and carbon emissions from a historic
1589 megafire in boreal forests of China, *Sci Total Environ*, 136534, <https://doi.org/10.1016/j.scitotenv.2020.136534>, 2020.
- 1590 Yanagiya, K., and Furuya, M.: Post-wildfire surface deformation near Batagay, Eastern Siberia, detected by L-band and C-
1591 band InSAR, *J. Geophys. Res*, 125, e2019JF005473, <https://doi.org/10.1029/2019JF005473>, 2020.
- 1592 York, A., Bhatt, U.S., Gargulinski, E., Garbinski, Z., Jain, P., Soja, A., Thoman, R.L., and Ziel, R.: Wildland Fire in High
1593 Northern Latitudes, in: *Arctic Report Card 2020*, edited by: Thoman, R.L., Richter-Menge, J., and Druckenmiller, M.L.,
1594 available at: [https://arctic.noaa.gov/Report-Card/Report-Card-2020/ArtMID/7975/ArticleID/903/Wildland-Fire-in-High-](https://arctic.noaa.gov/Report-Card/Report-Card-2020/ArtMID/7975/ArticleID/903/Wildland-Fire-in-High-Northern-Latitudes)
1595 *Northern-Latitudes*, <https://doi.org/10.25923/2gef-3964>, 2020.
- 1596 Young, A. M., Higuera, P. E., Duffy, P. A., and Hu, F. S.: Climatic thresholds shape northern high-latitude fire regimes and
1597 imply vulnerability to future climate change, *Ecogeg*, 40, 606, <https://doi.org/10.1111/ecog.02205>, 2016.
- 1598 Young, T. K., and Bjerregaard, P.: Towards estimating the indigenous population in circumpolar regions. *Int. J. Circumpolar*
1599 *Health*, 78, 1653749, <https://doi.org/10.1080/22423982.2019.1653749>, 2019.
- 1600 Yu, P., Toon, O.B., Bardeen, C.G., Zhu, Y., Rosenlof, K.H., Portmann, R.W., Thornberry, T.D., Gao, R.S., Davis, S.M., Wolf,
1601 E.T. and de Gouw, J.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365,
1602 587, <https://doi.org/10.1126/science.aax1748>, 2019.
- 1603 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum,
1604 *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010gl043584>, 2010.
- 1605 [Yuan, H., Richter, F., and Rein, G.: A multi-step reaction scheme to simulate self-heating ignition of coal: Effects of oxygen](#)
1606 [adsorption and smouldering combustion, *Proc Combust Inst*, 38, 4717-4725, <https://doi.org/10.1016/j.proci.2020.07.016>,](#)
1607 [2021.](#)
- 1608 Zhang, R., Huang, C., Zhan, X., Jin, H., and Song, X. P.: Development of S-NPP VIIRS global surface type classification map
1609 using support vector machines, *Int. J. Digit. Earth*, 11, 212, <https://doi.org/10.1080/17538947.2017.1315462>, 2018.
- 1610 Zhu, C., Kobayashi, H., Kanaya, Y., & Saito, M. (2017). Size-dependent validation of MODIS MCD64A1 burned area over
1611 six vegetation types in boreal Eurasia: Large underestimation in croplands, *Sci Rep*, 7, 4181, [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-017-03739-0)
1612 [017-03739-0](#), 2017.

Formatted: Finnish

1613 Zielinski, T., Bolzacchini, E., Cataldi, M., Ferrero, L., Grafl, S., Hansen, G., Mateos, D., Mazzola, M., Neuber, R., Pakszys,
1614 P. and Posyniak, M., 2020. Study of Chemical and Optical Properties of Biomass Burning Aerosols during Long-Range
1615 Transport Events toward the Arctic in Summer 2017, Atmosphere, 11, 84, <https://doi.org/10.3390/atmos11010084>, 2020.

1616 **Tables and Figures**

1617 **Table 1: Policy driven questions tasked to be answered by the SLCF EG of AMAP, a Working Group of the Arctic Council, for the**
1618 **AMAP Assessment 2021: Arctic climate, air quality, and health impacts from short-lived climate forcers (SLCFs).**

1. What are the impacts of climate change on fire risk and fire emissions?
2. What are current and future fire management strategies in the Arctic?
3. What are the long term emissions from open biomass burning?
4. What are the emissions from 'natural' fires?
5. What are the emissions from human caused open biomass burning?
6. What are the uncertainties in future natural and open biomass burning emissions?

1619
1620






Figure 1: A sample of peer-reviewed future Arctic fire risk variables due to expected ecological and meteorological transitions by mid- and late 21st century climate change for Arctic Council member states. 'Up arrows' indicate increase in fire risk and 'down arrows' indicate a decrease in fire risk, with the location of the arrows approximate to the location of fire risk from the literature

1626
1627
1628
1629

and not projections for a given country; the dashed line indicates the boundary between European Russia, and Siberia and the Russian Far East. Note that taiga is used in northern forest zones completely contained in Russia while boreal is used for the rest of the Pan-Arctic northern forests.



1630
1631
1632
1633
1634
1635

Figure 2: Annual black carbon (BC) emissions in Tg from three commonly used global fire emissions models and annual fire activity from the MODIS Collection 6 active fire product (Terra and Aqua) split by latitude ranges for the Arctic Council Region, 2005 - 2018; note the y-axis has been standardized for each model for ease of comparison; dotted line is the positive trend for BC emissions from open biomass burning and 1 km MODIS active fire detections (Terra and Aqua) for 60° to 70° N.




Figure 3: Annual 2018 BC and CH₄ emissions in Tg from five global fire emissions models and a custom AMAP fire emissions model for north of 45°N.

Table 21: Summary table of BC, PM_{2.5}, and CH₄ emissions in teragrams (Tg) from reported statistics on burned area from the Arctic Council members; sources for burned area include Norway (DSB, 2020), Greenland (Markuse, 2019), Finland (Ketola, 2020), Sweden (Betänkande av 2018 års skogsbrandsutredning, 2019), Canada (CIFFC, 2020), Alaska (Alaska Division of Forestry, 2020), the contiguous United States (NIFC, 2019), and the Russian Federation (ФБУ "АВИАЛЕСООХРАНА", 2019); fuel loadings and combustion completeness from Van Leeuwen et al. (2014) for boreal forests, with tundra values used for Greenland and temperate forests for the USA/CONUS; emission factors taken from GFED4.

Country/ Region	Year	Official Burned Area (km ²)	BC (Tg)	PM _{2.5} (Tg)	CH ₄ (Tg)
Norway	2019	0.03	7.61E-12	2.33E-10	9.08E-11
Denmark/ Greenland	2019	8	1.27E-10	2.88E-08	6.59E-08
Finland	2019	6	2.00E-09	5.00E-08	2.00E-08
Sweden	2018	250	6.30E-08	1.94E-06	7.60E-07
Canada	2019	18,389	4.67E-06	1.43E-04	5.56E-05
USA/ Alaska	2019	10,481	2.66E-06	8.14E-05	3.17E-05

USA/ CONUS	2019	18,876	1.02E-05	3.43E-04	9.64E-05
Russia	2019	100,785	2.56E-05	7.83E-04	3.05E-04
Total		148,795	4.30E-05	1.35E-03	4.90E-04


1646
1647



1648

1649 **Figure 4: Annual black carbon emissions for 2010, 2015, and 2020 from four anthropogenic source sectors (residential, transport,**
1650 **flaring, others) from GAINS and wildfires from GFAS, presented globally (left) and (right) 50°- 60°N (lighter colours of the**
1651 **cumulative bar) and north of 60°N latitude (darker colours of the cumulative bar).**





1652



1653


1654 **Figure 5: Annual PM_{2.5} emissions for 2010, 2015, and 2020 from four anthropogenic source sectors (residential, transport, flaring,**
 1655 **others) from GAINS and wildfires from GFAS, presented globally (left) and (right) 50°- 60°N (lighter colours of the cumulative bar)**
 1656 **and north of 60°N latitude (darker colours of the cumulative bar).**

1657



1660
1661
1662
1663
1664

Figure 6: Annual CH₄ emissions for 2010, 2015, and 2020 from four anthropogenic source sectors (agriculture, energy (including flaring, residential, transport, waste flaring, others) from GAINS and wildfires from GFAS, presented globally (left) and (right) 50°-60°N (lighter colours of the cumulative bar) and north of 60°N latitude (darker colours of the cumulative bar).







Figure 7: Monthly black carbon emissions from the leading anthropogenic sector, residential heating, in GAINS and wildfires from GFAS by latitudinal ranges (left) and based on global estimates (left/right) and by latitudinal ranges (right); emissions are averaged from the given years of 2010, 2015 and 2020 to align with the GAINS data availability.



1671

1672 **Figure 8: Sectoral black carbon emissions above 60° N (lighter colours) and 65° N (darker colours) for 2010, 2015, and 2020;**
 1673 **anthropogenic emissions from GAINS and wildfire emissions from GFAS.**

1674

1675 **Code and Data Availability**

1676 The GFEDv4s, FINNv1.5, GFASv1.2, QFEDv2.5r1, FEERv1.0-G1.2 fire emissions data for 2005 through 2018 were
 1677 downloaded from <https://globalfires.earthengine.app/view/firecam>. The AMAP SLCF EG 2018 Pan-Arctic fire emissions
 1678 database can be downloaded at <https://zenodo.org/record/4648723#.YGTq469KhPY> (embargoed access until review is
 1679 complete, can be provided to editors to share with reviewers) and R code used to compute it can be downloaded at
 1680 <https://github.com/fainjj>. 2020 global GFAS emissions data was downloaded from: [https://apps.ecmwf.int/datasets/data/cams-](https://apps.ecmwf.int/datasets/data/cams-gfas/)
 1681 [gfas/](https://apps.ecmwf.int/datasets/data/cams-gfas/). GAINS global emission data can be accessed at
 1682 <https://iiasa.ac.at/web/home/research/researchPrograms/air/ECLIPSEv6.html>.

1683 **Supplement**

1684 The supplement related to this article is available online at:

1685 **Author Contributions**

1686 JLM coordinated the review, designed the fire emission models comparison, co-led the creation of the custom AMAP fire
1687 emissions model with JJF, archived the literature for the review, wrote the manuscript, and led the revision. JA, VVP, ZK, SE,
1688 AV, SRA, NE, NMT, EIP, AJS, and KK provided major efforts in manuscript design, organization, and revision. VVP, JJF,
1689 ZK, and JLM led the GAINS and GFAS analysis. JJF, VVP, SE, and JLM created the figures and supplemental materials, with
1690 input from the remaining authors. All authors contributed to interpretation and verification of the review, as well as contributing
1691 to the writing of the manuscript.

1692 **Competing Interests**

1693 The authors declare that they have no conflict of interest.

1694 **Special Issue Statement**

1695 This article is part of the special issue “Arctic climate, air quality, and health impacts from short-lived climate forcers (SLCFs):
1696 contributions from the AMAP Expert Group”.

1697 **Acknowledgements**

1698 This paper was developed as part of the Arctic Monitoring Assessment Programme (AMAP), AMAP 2021 Assessment:
1699 Arctic climate, air quality, and health impacts from short-lived climate forcers (SLCFs). The authors would like to thank the
1700 Arctic Monitoring and Assessment Programme Secretariat for providing additional input and review. C.J. Mescher is
1701 acknowledged for his contributions in designing and creating original scientific artwork used in this manuscript. The authors
1702 would also like to thank the anonymous reviewers, whose reviews improved this manuscript.-

1703 **Financial support**

1704 This research has been supported by: JLM and JJF acknowledge internal research support from Miami University; JA and AV
1705 acknowledges funding from the Ministry for Foreign Affairs of Finland (IBA Forest Fires, decision PC0TQ4BT-53); VVP
1706 acknowledges support from Business Finland (BC Footprint; grant no. 1462/31/2019); SRA acknowledges support from the
1707 ACROBEAR project, funded by the Belmont Forum Climate, Environment and Health (CEH) Collaborative Research Action
1708 and the UK Natural Environment Research Council (grant no. NE/T013672/1), and support from the Arctic Monitoring and
1709 Assessment Programme (AMAP); N.E. and S.E. acknowledge AMAP funding; N.M.T and E.I.P. acknowledge funding from
1710 the Russian Foundation for Basic Research (RFBR grant no. 19-45-240004) and a joint project of the Government of

1711 Krasnoyarsk Territory and Russian Foundation for Basic Research (GKT KRFS & RFBR grant no. 20-05-00540); A.J.S.
1712 acknowledges support from NASA’s Weather and Data Analysis program; and L.H. acknowledges the Climate Adaptation
1713 Research Fund from Environment & Climate Change Canada. This publication was produced with the financial support of the
1714 European Union via the EU-funded Action on Black Carbon in the Arctic. Its contents are the sole responsibility of J.L.M.,
1715 V.V.P., Z.K., and J.J.F., and do not necessarily reflect the views of the European Union.

