

Subsurface iron accumulation and rapid aluminium removal in the Mediterranean following African dust deposition

Matthieu Bressac^{1,2}, Thibaut Wagener³, Nathalie Leblond⁴, Antonio Tovar-Sánchez⁵, Céline Ridame⁶, Vincent Taillandier¹, Samuel Albani^{7,8}, Sophie Guasco³, Aurélie Dufour³, Stéphanie H. M. Jacquet³,
5 François Dulac⁸, Karine Desboeufs⁹, and Cécile Guieu¹

¹ Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, Villefranche-sur-mer, 06230, France

² Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

³ Aix Marseille Univ., CNRS, IRD, Université de Toulon, MIO UMR 110, Marseille, 13288, France

⁴ Sorbonne Université, CNRS, Institut de la Mer de Villefranche, IMEV, 06230 Villefranche-sur-Mer, France

10 ⁵ Department of Ecology and Coastal Management, Institute of Marine Sciences of Andalusia (ICMAN-CSIC), 07190 Puerto Real, Spain

⁶ Sorbonne Université, LOCEAN, 4 Place Jussieu – 75252 Paris Cedex 05, France

⁷ Department of Environmental and Earth Sciences, University of Milano–Bicocca, Milan, Italy

15 ⁸ Laboratoire des Sciences du Climat et de l'Environnement (LSCE), UMR 8212 CEA-CNRS-UVSQ, Institut Pierre-Simon Laplace, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

⁹ Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR7583 CNRS, Université de Paris, Université Paris-Est Créteil, Institut Pierre-Simon Laplace, France

Correspondence to: Matthieu Bressac (matthieu.bressac@imev-mer.fr)

Abstract. Mineral dust deposition is an important supply mechanism for trace elements in the low-latitude ocean. Our
20 understanding of the controls of such inputs has been mostly built onto laboratory and surface ocean studies. The lack of direct observations and the tendency to focus on near surface waters prevent a comprehensive evaluation of the role of dust in oceanic biogeochemical cycles. In the frame of the PEACETIME project (ProcEss studies at the Air-sEa Interface after dust deposition in the MEditerranean sea), the responses of the aluminium (Al) and iron (Fe) cycles to two dust wet deposition events over the central and western Mediterranean Sea were investigated at a timescale of hours to days using a comprehensive dataset
25 gathering dissolved and suspended particulate concentrations, along with sinking fluxes.

Dissolved Al (dAl) removal was dominant over dAl released from dust. Fe/Al ratio of suspended and sinking particles revealed that biogenic particles, and in particular diatoms, were key in accumulating and exporting Al relative to Fe. By combining these observations with published Al/Si ratios of diatoms, we show that adsorption onto biogenic particles, rather than active uptake, represents the main sink for dAl in Mediterranean waters. In contrast, systematic dissolved Fe (dFe) accumulation
30 occurred in subsurface waters (~100-1000 m), while dFe input from dust was only transient in the surface mixed-layer. The rapid transfer of dust to depth, the Fe-binding ligand pool in excess to dFe in subsurface (while nearly-saturated in surface), and low scavenging rates in this particle-poor depth horizon are all important drivers of this subsurface dFe enrichment.

At the annual scale, this previously overlooked mechanism may represent an additional pathway of dFe supply for the surface ocean through diapycnal diffusion and vertical mixing. However, low subsurface dFe concentrations observed at the basin
35 scale (<0.5 nmol kg⁻¹) questions the residence time for this dust-derived subsurface reservoir, and hence its role as a supply

mechanism for the surface ocean, stressing the need for further studies. Finally, these contrasting responses indicate that dAl is a poor tracer of dFe input in the Mediterranean Sea.

1 Introduction

Globally, iron (Fe) supply to the surface ocean sets the productivity of major phytoplankton groups (Moore et al., 2013).
40 Among the multiple sources of Fe, atmospheric deposition of mineral dust represents an important supply mechanism in the low-latitude ocean (Duce et al., 1991; Jickells et al., 2005; Conway and John, 2014), and plays a key role in the functioning of Low Nutrient Low Chlorophyll (LNLC) systems (e.g., Guieu et al., 2014a). Despite widespread attention over the last three decades, large uncertainties remain in the factors controlling aerosol Fe solubility (Meskhidze et al., 2019). Consequently, poorly constrained controls of Fe solubility partly explain the substantial inter-model difference in the atmospheric dFe input
45 flux to the global ocean ($\sim 1\text{-}30 \text{ Gmol yr}^{-1}$; Tagliabue et al., 2016), and hinder accurate predictions of the impact of dust on ocean productivity.

African dust deposition events have long been known to impact trace element concentrations and fluxes in the upper water column of the Mediterranean (e.g., Buat-Ménard et al., 1989; Davies and Buat-Ménard, 1990; Quézel et al., 1993; Guerzoni et al., 1999; Heimbürger et al., 2011). Our understanding of the role of dust in marine biogeochemical cycles remains limited,
50 however, partly resulting from the difficulty in quantifying atmospheric dust fluxes to the surface ocean at short timescales. In the absence of direct assessments of atmospheric inputs, marine concentrations of tracers such as aluminium (Al) have been widely used to constrain these fluxes (e.g., Measures and Brown, 1996; Han et al., 2008; Anderson et al., 2016; Menzel Barraqueta et al., 2019). Al is predominantly of crustal origin and is characterized by a similar fractional solubility as Fe with a longer residence time in seawater. Al could thus be used to constrain the integrated input of dust Fe over seasonal timescales
55 (Dammshäuser et al., 2011). However, the fact that the distribution of Al can itself be controlled by biological activity (e.g., Mackenzie et al., 1978; Moran and Moore, 1988a; Li et al., 2013; Middag et al., 2015; Rolison et al., 2015) questions its quality as a tracer.


In addition, dust deposition being highly episodic in time and spatially patchy (Donaghay et al., 1991; Guieu et al., 2014a; Vincent et al., 2016), direct observations at sea are extremely challenging, and hence sparse (e.g., Croot et al., 2004; Rijkenberg
60 et al., 2008). To overcome this limitation, a variety of small-volume enclosed systems have been used to quantify Fe solubility from dust. Although yielding important insights into atmospheric trace element solubilities (Baker and Croot, 2010 and references therein), these systems do not fully simulate *in situ* conditions (de Leeuw et al., 2014), motivating the development of larger volume experiments ($>100 \text{ L}$) where dust particles are free to sink and interact with dissolved and particulate organic matter while sinking (Bressac and Guieu, 2013; Guieu et al., 2014b; Herut et al., 2016; Gazeau et al., 2021). Two key findings
65 emerged from these large volume experiments. First, they demonstrated the pivotal role played by the *in situ* biogeochemical conditions at the time of deposition in controlling post-depositional processes (i.e., dissolution, organic complexation, adsorption, colloidal aggregation), and their interplay (Wagener et al., 2010; Bressac and Guieu, 2013; Wuttig et al., 2013).

The most striking and unexpected consequence is that upon deposition, dust can act as a net sink of dFe through scavenging (Wagener et al., 2010; Ye et al., 2011). Second, the large range in Fe solubility observed in seawater depending on the season reveals that oceanic rather than atmospheric conditions, determine the flux of ‘truly’ bioavailable Fe to the surface ocean (Bressac and Guieu, 2013). However, these findings are valid in the first meters of the water column and direct observations of the whole water column are needed if we are to fully understand the role of dust in the oceanic iron cycle. For this purpose, the Mediterranean Sea is a particularly relevant region. This semi-enclosed basin, characterized by a west-to-east gradient in oligotrophy, receives some of the largest dust inputs of the ocean (Guerzoni et al., 1999), mostly in the form of wet deposition in the central and western part of the Basin, and a few intense events may account for the bulk of the annual deposition (Loÿe-Pilot and Martin, 1996; Vincent et al., 2016). The PEACETIME project (ProcEss studies at the Air-sEa Interface after dust deposition in the MEditerranean sea) and oceanographic campaign on board the R/V *Pourquoi Pas?* provided a unique opportunity to directly observe the biogeochemical effects of two mineral dust wet deposition events of contrasting intensity that occurred during late spring 2017 in the central and western open Mediterranean Sea (Guieu et al., 2020). The presence of the R/V before, during, and/or few days after deposition allowed investigating (1) the parameters and processes shaping the contrasting distributions of dAl and dFe, (2) the importance of the timescale considered when assessing the flux of bioavailable Fe to the surface ocean, and (3) the relevance of using dAl to constrain dFe input from dust.

2 Materials and Methods

2.1 Oceanographic cruise

The PEACETIME cruise (doi.org/10.17600/15000900) was conducted during late spring conditions in May and June 2017 aboard the R/V *Pourquoi Pas?* in the central and western Mediterranean Sea. In total, 10 short stations (~8 hours) and 3 long stations located in the Tyrrhenian Sea (TYR; occupation = 4 days), the Ionian Sea (ION; 4 days), and in the western Algerian basin (FAST; 5 days) were occupied (Fig. 1). FAST was an opportunistic station dedicated to investigating the biogeochemical effects of a dust deposition event by combining atmospheric and oceanographic in situ measurements before, during and after deposition (Guieu et al., 2020). At all stations, a standard CTD rosette with 24 Niskin bottles (12 L) equipped with a Sea-Bird SBE9 CTD, transmissometer, chlorophyll fluorescence (Chelsea Acquatracka), photosynthetically active radiation and oxygen (SBE43) sensors, and a trace metal-clean (TMC) titanium rosette were deployed to sample the water column for biological and chemical parameters. Samples for aluminium and iron analyses were collected using the TMC titanium rosette mounted with GO-FLO bottles deployed on a Kevlar cable with a dedicated clean winch, while samples for particulate Al (pAl) determination were also collected at all the stations from the standard rosette (see section 2.3 and Supp. Table 1).



100 **Figure 1: Sampling locations during the PEACETIME cruise (filled circles). The cruise track consisted of 10 short stations and 3 long stations (TYR, ION, and FAST). Open circles correspond to the stations 64PE370_12 (orange), 64PE374_13 (red), and 64PE374_12 (green) from the GEOTRACES GA04 section (Rolison et al., 2015; Gerringa et al., 2017) used for comparison in Fig. 3. Note that the same color code is used in figures 3, 6 and 7.**

2.2 Dissolved Al and Fe concentrations

105 Immediately after recovery, the GO-FLO bottles were transferred inside a class-100 clean laboratory container. Seawater samples were directly filtered from the GO-FLO bottles through acid-cleaned 0.2 μm capsule filters (Sartorius Sartobran-P-capsule 0.45/0.2 μm). Dissolved Fe and Al samples were stored in acid-washed low-density polyethylene bottles and immediately acidified to pH 1.8 (quartz-distilled HCl) under a laminar flow hood.

110 Dissolved Al analyses were conducted on board using the fluorometric method described by Hydes and Liss (1976). Briefly, the samples were buffered to pH 5 with ammonium-acetate and the reagent lumogallion was added. The samples were then heated to 80°C for 1.5 h to accelerate the complex formation. The fluorescence of the sample was measured with a Jasco FP-2020 Plus spectrofluorometer (excitation wavelength 495 nm, emission wavelength 565 nm). Calibration was realized with additions of Al standard solution in seawater. The detection limit (DL; 3 times the standard deviation (SD) of the concentrations measured from the dAl-poor seawater used for calibration) varied between 0.2 and 0.5 nmol kg^{-1} . The reagent blank determined by measuring acidified ultrapure water varied between 0.9 and 1.7 nmol kg^{-1} .

115 Dissolved Fe concentrations were measured (mostly on board in the class-100 clean laboratory) using an automated Flow Injection Analysis (FIA) with online preconcentration and chemiluminescence detection (Bonnet and Guieu, 2006). The stability of the analysis was assessed by analyzing daily an internal acidified seawater standard. On average, the DL was 15 pmol kg^{-1} (3 times the SD of the concentration measured 5 times from the same dFe-poor seawater) and the accuracy of the

method was controlled by analysing on a regular basis the GEOTRACES seawater standards SAFe D1 ($0.64 \pm 0.13 \text{ nmol kg}^{-1}$ ($n = 19$), consensus value $0.67 \pm 0.04 \text{ nmol kg}^{-1}$), GD ($1.04 \pm 0.10 \text{ nmol kg}^{-1}$ ($n = 10$), consensus value $1.00 \pm 0.10 \text{ nmol kg}^{-1}$) and GSC ($1.37 \pm 0.16 \text{ nmol kg}^{-1}$ ($n = 4$), consensus value $1.50 \pm 0.11 \text{ nmol kg}^{-1}$).

2.3 Suspended particulate trace elements

Just prior to sampling for particulate trace elements (pTM), GO-FLO bottles were gently mixed and pTM were sampled directly on-line from the pressurized ($0.2 \mu\text{m}$ filtered N_2) GO-FLO bottles onto acid-cleaned 25 mm diameter Supor $0.45 \mu\text{m}$ polyethersulfone filters mounted on Swinnex polypropylene filter holders (Millipore), following GEOTRACES recommendations. Filtration was stopped when the filter clogged or the bottle was empty. On average, each particulate concentration was obtained by filtering 4.8 L (range 1.1–10.2 L). When the filtration was complete, filter holders were transferred under a laminar flow hood and residual seawater was removed using a polypropylene syringe. Filters were stored in acid-cleaned petri-slides, left open under the laminar flow hood for ~ 24 h to allow the filters to dry. Particulate samples were digested (10% HF / 50% HNO_3 (v/v)) following the protocol described in the ‘GEOTRACES Cookbook’ (Cutter et al. 2010) and Planquette and Sherrell (2012). Procedural blanks consisted of unused acid-cleaned filters. Analyses were performed on a HR-ICP-MS (High Resolution Inductively Coupled Plasma Mass Spectrometry; Element XR, Thermo-Fisher Scientific). The accuracy of the measurements was established using the certified reference materials (CRM) MESS-4 and PACS-3 (marine sediments, National Research Council Canada) (Supp. Table 2).

In addition, pAl concentrations were also obtained at all the stations from the standard rosette (Supp. Table 1). This additional pAl dataset already published by Jacquet et al. (in revision) was obtained according to the sampling, processing and analysis methods described in Jacquet et al. (2015). Briefly, 4 to 6 L of seawater collected with the Niskin bottles were filtered onto acid-cleaned 47 mm polycarbonate filters ($0.4 \mu\text{m}$ porosity). Filters were rinsed with Milli-Q grade water and dried at 50°C . A total digestion of the membranes was performed using a tri-acid mixture (0.5 mL HF / 1.5 mL HNO_3 / 1 mL HCl), and analyses were performed on the same HR-ICP-MS. A good agreement was obtained when comparing pAl concentrations obtained with the TMC and standard rosettes at ION and FAST (difference in sampling time at TYR prevents quantitative comparison; see Sect. 4.1) (Supp. Fig. 1), demonstrating the absence of contamination for pAl when using the standard rosette.

2.4 Export fluxes and composition

Sinking particles were collected at ~ 200 , 500, and 1000 m depth using PPS5 sediment traps (Technicap, France; 1 m^2 collection area) deployed on a free-drifting mooring for 4 (TYR and ION) and 5 days (FAST). Cups were filled with filtered seawater and buffered formaldehyde (2% final concentration) as a biocide. Once recovered, each cup representing 24 hours of collection was stored in the dark at 4°C until processed. Samples were treated following the standard protocol followed at the national service ‘‘Cellule Piège’’ of the French INSU-CNRS (Guieu et al., 2005) following the JGOFS’ protocol. After removing the swimmers, the remaining sample was rinsed three times with ultrapure water in order to remove salt, and then freeze-dried.

150 The total amount of material collected was weighed to quantify the total exported flux. Several aliquots were then used to
measure the following components: total and organic carbon, particulate Al and Fe, lithogenic and biogenic silica (LSi and
BSi, respectively). Total carbon, particulate organic carbon (POC) (after removing inorganic carbon by acidification with HCl
2N), and particulate organic nitrogen (PON) were measured on an elemental analyzer CHN (2400 Series II CHNS/O Elemental
Analyzer Perkin Elmer). For one sample (TYR 1000 m), 5 aliquots were analysed, yielding a coefficient of variation (CV) of
6%. Particulate inorganic carbon (PIC) was quantified by subtracting POC from total particulate carbon. Particulate Fe and Al
155 concentrations were determined by ICP-AES (Inductively Coupled Plasma Atomic Emission Spectrometry, Spectro ARCOS
Ametek) after acid-digestion following the protocol described in Ternon et al. (2010). Blanks were negligible (<0.8% of the
lowest Al and Fe concentrations of the digested aliquots) and the efficiency of the acid digestion was established using the
CRM GBW-07313 (marine sediment, National Research Center for CRMs, China) (Supp. Table 2). Samples for BSi and LSi
(2 or 3 aliquots) were digested (NaOH at 95°C and HF at ambient temperature, respectively) and concentrations analysed by
160 colorimetry (Analytikjena Specor 250 plus spectrophotometer) (Nelson et al. 1989; Mullin and Riley, 1955). Mean export
fluxes and composition of exported material are presented in Table 1.

165 **Table 1: Mean flux values for total mass (dry weight), particulate organic carbon (POC), particulate inorganic carbon (PIC),
particulate organic nitrogen (PON), biogenic silica (BSi), lithogenic silica (LSi), particulate aluminium and iron collected at TYR,
ION and FAST (data not available at 500 m depth at TYR due to technical problems with the sediment trap). Values in parentheses
correspond to the standard deviation of the arithmetic mean of the 4 (TYR and ION) and 5-day (FAST) deployment period.**

	Latitude	Longitude	Depth	Mass	POC	PIC	PON	BSi	LSi	Al	Fe
	°N	°E	[m]	[mg m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]	[μmol m ⁻² d ⁻¹]
			222	64.5 (54.3)	340.7 (215.1)	157.1 (109.1)	40.2 (13.4)	88.2 (78.7)	339.6 (308.9)	100.6 (101.9)	25.3 (25.6)
TYR	39.34	12.59	500	-	-	-	-	-	-	-	-
			1056	76.0 (21.4)	185.2 (49.1)	225.2 (51.9)	16.8 (4.5)	86.0 (27.4)	458.2 (192.1)	135.6 (40.0)	34.9 (11.6)
			265	34.1 (19.1)	315.6 (146.5)	56.5 (29.9)	31.3 (14.4)	64.4 (53.1)	126.8 (94.3)	34.0 (28.3)	8.5 (6.7)
ION	35.49	19.80	560	30.1 (10.2)	144.2 (22.9)	108.3 (37.2)	12.6 (1.2)	48.3 (15.6)	125.7 (39.5)	46.2 (18.3)	11.9 (4.4)
			1097	30.1 (9.3)	115.7 (16.5)	98.7 (56.6)	7.6 (2.0)	36.3 (12.1)	126.8 (36.5)	45.7 (15.7)	12.3 (3.9)
			259	29.1 (14.1)	368.2 (139.5)	67.6 (62.0)	45.8 (20.6)	90.8 (58.9)	72.8 (48.6)	26.5 (19.9)	6.1 (4.5)
FAST	37.95	2.91	498	33.9 (10.5)	403.8 (150.3)	116.6 (85.0)	56.7 (26.0)	60.7 (42.3)	68.6 (64.3)	27.1 (22.2)	6.3 (5.2)
			1004	36.4 (17.7)	379.0 (318.5)	74.1 (44.2)	53.0 (50.7)	59.3 (29.8)	82.4 (57.0)	38.0 (19.6)	9.1 (4.8)

2.5 Inventories and Kd

170 Discrete measurements at different depths were used to calculate the water column integrated Al and Fe inventories ($\mu\text{mol m}^{-2}$) by trapezoidal integration. The concentration measured nearest to the surface was assumed to be constant up to 0 m. At FAST, six replicate measurements of dAl and dFe were performed at 5 and 400 m depth from 2 sets of 6 x GO-FLO bottles. The CV obtained at 5 and 400 m depths were used to determine the uncertainties in the 0-20 m and 0-200 m inventories, respectively. Variability among replicates was higher for dFe (CV = 11.3 and 6.9% at 5 and 400 m depth, respectively) than for dAl (CV = 5.3 and 1.1% at 5 and 400 m depth, respectively), potentially reflecting a small-scale variability in the dFe distribution.

175 At the FAST station, the partitioning coefficient between the particulate and dissolved phases (K_d ; $[\text{particulate}]/[\text{dissolved}]$) was used to investigate exchanges between dissolved and particulate pools of Al and Fe. Following the relative change over time of this metric allowed excluding potential artefacts related to change in water masses driven by lateral advection (Guieu et al., 2020).

180 3 Results

3.1 Biogeochemical conditions

The PEACETIME cruise took place in late spring when the stratification of the upper water column was well established with the mixed-layer depth ranging between ~10 and 20 m along the cruise track (Van Wambeke et al., 2020). Surface chlorophyll *a* concentrations ($<0.1 \mu\text{g L}^{-1}$) were typical of oligotrophic conditions (Guieu et al., 2020). A diatom-dominated deep chlorophyll maximum (DCM) that coincided with a maximum in biomass and primary production was well developed and observed all along the cruise track (Marañón et al., 2021). POC downward fluxes measured at 200 m depth were similar at the 3 long stations, while downward fluxes of Al and LSi, two proxies for dust, were maximum at TYR (Table 1). At the surface, dAl distribution was characterized by a marked west-to-east increasing gradient (Supp. Fig. 2b) driven by advective mixing between (dAl poor) Atlantic and Mediterranean waters and by the accumulation of dust, and reflected by a strong relationship between surface dAl concentrations and salinity (Guerzoni et al., 1999; Rolison et al., 2015). All along the transect, dFe concentrations were high in the upper 100 m (up to 2.7 nmol kg^{-1}), and decreasing to levels $<0.5 \text{ nmol kg}^{-1}$ below the euphotic layer (Supp. Fig. 2c). Subsurface patches of high dFe concentrations previously observed in the eastern Mediterranean Basin, and attributed to hydrothermal activity and mud-volcanoes (Gerringa et al., 2017), were not observed along our cruise track.


3.2 Dust deposition over the central and western Mediterranean Sea

195 The impact in the water column of two dust deposition events of contrasting magnitudes were studied during the cruise. They occurred in the area of the TYR and FAST stations (Fig. 1), on the 11-12 May and 3-5 June, respectively. The first deposition

event in the southern Tyrrhenian Sea was not directly observed but hypothesized based on satellite observations of intense dust plume transport and water-column Al inventory presented in the following. The combined analysis of time-series of quick-looks of operational aerosol products from MSG/SEVIRI (Fig. 2a) and from meteorological and dust transport models available during the campaign (Guieu et al., 2020; Supp. Fig. 3) allowed us to suspect that a red rain event likely occurred over the southern Tyrrhenian Sea on the 11th of May and possibly on the early 12th. The daytime daily mean aerosol optical depth (AOD) product over oceanic areas (Thieuleux et al., 2005) shows that a large dust plume was exported from the Tunisian and Libyan coasts towards southern Italy and Greece from the 10th of May (Fig. 2a). Up to 1.5-1.6 AOD at 550 nm (AOD₅₅₀) (i.e., about 2 g m⁻² of dust in the column assuming a specific extinction cross-section of dust of 0.77 m² g⁻¹ following Dulac et al., 1992) was observed on the 12th north of Sicily in the area of the Tyrrhenian stations, a rather high value relatively unusual in this area (Gkikas et al., 2016). On the 11th, clouds developed over most of the area affected by dust, whereas clear (turbid) sky was again present during daytime on the 12th. The extension and dynamics of this dust transport event were reasonably well forecasted by the various existing regional dust transport models available during the campaign (Supp. Fig. 3), including the set of models of the World Meteorological Organization Sand and Dust Storm Warning Advisory and Assessment System operated by the Barcelona Supercomputing Center (BSC; Huneeus et al., 2016), the SKIRON/Dust model operated by the Atmospheric Modeling and Weather Forecasting Group of the National University of Athens (Spyrou et al., 2010), the CAMS model operated by the European Centre for Medium-range Weather Forecast (ECMWF; Flemming et al., 2015), the NCEP/GFS model operated by the U.S. National Weather Service (Han et al., 2017), the NAAPS model modified from that of Christensen (1997) and operated by the U.S. Naval Research Laboratory, and the TAU/DREAM8 model operated by the Weather Research Centre of the Tel Aviv University (TAU; Kishcha et al., 2008). The dust plume extension in the cloudy area on the 11th is illustrated by Supp. Fig. 3a and b. Most meteorological models predicted significant precipitation over the Tyrrhenian Sea on the 11th (Supp. Fig. 3c). Dust transport models producing dust deposition fluxes generally forecasted dust wet deposition on 11th of May between Tunisia and Italy, but with significant variability on the location, extent and schedule. The NMMB/BSC and SKIRON models predicted a significant wet deposition flux of dust, with at least 1.5 g m⁻² over 6 h in the area of our stations 5, TYR, and 6 in the afternoon of the 11th of May (Supp. Fig. 3d and e). The DREAM model versions operated by the BSC and TAU, however, forecasted much lower values or even no dust wet deposition in the Tyrrhenian stations area (Supp. Fig. 3f). However, model reanalyses now available do not show any precipitation in the area between the 10th and 13th of May (Fig. 2b). However, the half-hourly, 0.1°-resolution Global Precipitation Measurement (GPM) mission Integrated Multi-satellitE Retrievals for GPM (IMERG) final run images of the rainfall rate (product GPM_3IMERGHH_v06; Huffman et al., 2015) report the occurrence of light rains in the Tyrrhenian Sea in the early morning of the 11th of May, especially around 12°E and between 39°N and 40°N (<1 mm of accumulated precipitation, with a large error of up to several mm; Fig. 2b). The study of the TRMM-3B42-v6 product, a former multi-satellite, 3-hourly precipitation product in the western Mediterranean region has shown that the detection of light rainfall is difficult when compared to rain gauge observations, with many occurrences missed by the satellite product (Sarrand et al., 2012). Indeed, the minimum corresponding random error image is 0.238 mm hr⁻¹ for pixels without detected rain, which can be assimilated to a lower detection limit. During the whole

day, some light rain cells remain visible near Sardinia, Sicily or southern Italy. It is therefore well possible that a much larger area of the Tyrrhenian Sea than that reported by the GPM IMERG images has been affected by light precipitation on that day. For simplicity, the 11th of May 2017, 18:00 UTC will be considered as the time of deposition, that is approximately 3 to 10 days before our sampling of the area.

235 During the early June deposition event in the western Algerian basin, precipitation was directly observed in the area of the R/V and even sampled onboard (Desboeufs et al., in revision), associated with a dust transport event of moderate extent and intensity over the southwestern Mediterranean basin. The AOD₅₅₀ peaked at about 0.40 in the area of the FAST station (Desboeufs et al., in revision), corresponding to a maximum columnar dust load <0.4 g m⁻², assuming a non-dust background AOD₅₅₀ in the boundary layer of 0.10-0.15 as observed north of the plume or the day before the plume arrived. This dust plume
240 encountered a massive rain front covering ~80,000 km² and moving eastward from Spain and North Africa regions (Desboeufs et al., in revision). Direct atmospheric and oceanographic observations of this event were possible thanks to a dedicated ‘fast action’ strategy (see Guieu et al. (2020) for details). Two rain periods concomitant with the dust plume transported in altitude (1 to 4 km) allowed below-cloud deposition of dust in the FAST station area, as confirmed by on-board Lidar records (Desboeufs et al., in revision). The first rain period occurred on the 3rd of June in the neighbouring area of the R/, and the
245 second one occurred from the 4th (22:00 UTC) to the 5th of June (9:00 UTC), and was sampled on board the R/V on the 5th of June from 00:36 to 01:04 UTC (Desboeufs et al., in revision). This second rain event was characterized by a dust signature clearly revealed by the chemical composition of the rain. From the total (dissolved + particulate) Al concentration measured in this rainwater sample, a dust flux of 65 ± 18 mg m⁻² was derived (Desboeufs et al., in revision). This sampled flux, considered as relatively modest compared to the multi-year record in this area (Vincent et al., 2016), was likely in the lower range of the
250 total dust deposition flux that affected the whole area between the 3rd and 5th of June.




255 **Figure 2: (a) MSG/SEVIRI-derived daily (daytime) mean aerosol optical depth; the white ellipse on the 11th of May image includes the location of the 5 Tyrrhenian stations reported in Fig. 1; (b) Left: Time averaged map of the GPM mission multi-satellite precipitation final run estimate with gauge calibration (mm hr⁻¹) over 04:00-05:30 UTC on the 11th of May 2017 in the western Mediterranean region, from the GPM_3IMERGHH v06, half-hourly, 0.1°-resolution (Huffmann et al., 2015). Right: Corresponding random error (mm hr⁻¹). After images produced by the Giovanni online data system (Acker and Lepkouth, 2007).**

3.3 Reconstruction of the dust deposition fluxes

260 The absence of direct measurement of the dust deposition flux over the Tyrrhenian Sea and the limited spatial coverage of collection of atmospheric dust and rain at the FAST station call for an alternative approach to estimate dust deposition fluxes. For this purpose, we used the water-column AI inventory. We acknowledge that this approach involves uncertainties, as do all the observational approaches employed so far to quantify deposition (Anderson et al., 2016). Caveats include (1) other sources of pAI, and (2) some uncertainties into the derived dust fluxes that could come from the sampling method (Twining et al., 2015a), the time lag between deposition and sampling favouring dispersion of dust by lateral mixing, and to a lesser extent, 265 the limited vertical resolution below 500 m depth (Fig. 3a-d).

3.3.1 Central Mediterranean Sea

Dust deposition flux over the Tyrrhenian Sea was estimated from the Al_{excess} inventories corresponding to the difference between the measured 0-1000 m pAl inventories and a background 0-1000 m pAl inventory. In the absence of pre-depositional observations and historic pAl data (to the best of our knowledge), the median pAl vertical profile obtained during the cruise at the other stations unimpacted by this event (grey bold line on Fig. 3a-d), similar or slightly higher than pAl data available for the open Mediterranean Sea (e.g., Sarthou and Jeandel, 2001), was used as a background level. The comparison between the measured pAl vertical profiles and this background level revealed a marked excess in pAl south of Sardinia (ST04) and in the southern Tyrrhenian (ST05, TYR and ST06; Fig. 3a-d and Table 2). This spatial extent is in good agreement with the maps of precipitation and dust wet deposition provided for the 11th of May by the ARPEGE, SKIRON, and NMMB/BSC models (Supp. Fig. 3). The obtained Al_{excess} inventories were further corrected for the loss of pAl associated with the sinking flux using the pAl downward flux measured at 1000 m depth at the TYR station (assuming a constant flux over the 3 to 10-day period after deposition). Assuming that Al represents 7.1% of the dust in mass (Guieu et al., 2002), and further assuming that Al_{excess} resulted from a single dust event, a dust deposition flux ranging between 1.7 (ST06) and 9.2 g m⁻² (ST04) was derived from these Al_{excess} inventories (Table 2). Large uncertainties are associated with these dust flux estimates, partly due to potential additional sources of pAl that are suspected for fluxes derived from ST04 and ST05 but unlikely for TYR and ST06 (see Section 4.1). Nevertheless, the approach remains valuable to estimate the magnitude of this dust event.



285 **Figure 3: Vertical distribution of pAl (upper panels), dAl (middle panels), and dFe concentrations (lower panels) obtained during the PEACETIME cruise (filled circles) at ST04 (a, e, i), ST05 (b, f, j), TYR (c, g, k), and ST06 (d, h, l). Particulate Al concentrations in (c) correspond to the vertical profiles TYR_2 and TYR_3 (Fig. 5). Previously published vertical profiles of dAl (Rolison et al., 2015) and dFe (Gerringa et al., 2017) obtained at similar locations (Fig. 1) are superimposed (open circles). The median pAl vertical profile (bold lines) used as background level (see Sect. 3.3.1 for details), and the 1st and 3rd quantiles (grey-shaded areas) are represented in a-d.**

290

This range of dust deposition flux is of similar magnitude to the annual flux observed during former periods in the west central ($7.4 \text{ g m}^{-2} \text{ yr}^{-1}$; Vincent et al., 2016) and northwestern Mediterranean Sea ($11.4 \text{ g m}^{-2} \text{ yr}^{-1}$; Ternon et al., 2010), highlighting the remarkable magnitude of this event. Furthermore, this comparison with annual fluxes confirms that the annual deposition of African dust in the Mediterranean region is generally driven by only a few intense events (Loÿe-Pilot and Martin, 1996; Guerzoni et al., 1999; Kubilay et al., 2000; Desboeufs et al., 2018). The strong spatial variability of these dust flux estimates,

295

with a marked west-to-east gradient, might result from the varying time lag between deposition and sampling of the water column at these different stations (Table 2), but also from the patchiness of the rainfalls associated with the rain front (Supp. Fig. 3). Indeed, Vincent et al. (2016) showed that high deposition events in the western Mediterranean are often limited spatially although the associated dust plumes may affect a large part of the basin. By assuming that the deposition was spatially homogeneous over the southern Tyrrhenian, an Al export flux of more than 4000 $\mu\text{mol m}^{-2} \text{d}^{-1}$ is needed to explain the difference in the $\text{Al}_{\text{excess}}$ inventory observed between ST04, ST05 and TYR (i.e., ~ 3.6 to 8.4 days after deposition). This is an order-of-magnitude difference compared to the Al export flux measured at TYR ~ 5 to 8 days after deposition ($136 \pm 40 \mu\text{mol m}^{-2} \text{d}^{-1}$; Table 1), indicating that the observed spatial variability was primarily driven by the precipitation patchiness rather than related to the time lag between deposition and observations.

305

Table 2: Estimates of the input of pAl, dAl and dust south of Sardinia (ST04) and over the Tyrrhenian Sea (ST05, TYR and ST06) based on the 0-1000 m $\text{Al}_{\text{excess}}$ inventories. The relatively low vertical resolution at TYR_1 precludes accurate estimates.

	time since deposition ¹	$\text{Al}_{\text{excess}}$ ²	Al loss ³	pAl input ⁴	dAl input ⁵	dust flux ⁶
	[d]	$[\mu\text{mol m}^{-2}]$	$[\mu\text{mol m}^{-2}]$	$[\mu\text{mol m}^{-2}]$	$[\mu\text{mol m}^{-2}]$	$[\text{g m}^{-2}]$
ST04	3.6	23684	485	24169	363	9.2
ST05	4.5	17773	610	18382	276	6.9
TYR_1	5.6	-	-	-	-	-
TYR_2	6.4	3591	874	4465	67	1.7
TYR_3	8.4	4837	1143	5980	90	2.3
ST06	10.5	3126	1423	4549	68	1.7

¹ 11/05/17 18:00 UTC is considered as the time of deposition

² Difference between the measured and median 0-1000 m pAl inventories (see section 3.3.1 for details)

³ Estimates based on the downward Al flux (1000 m depth; TYR) and assuming a constant flux over time

⁴ Corresponds to $\text{Al}_{\text{excess}}$ corrected for Al loss

⁵ Estimates based on an Al fractional solubility of 1.5% (Wuttig et al. 2013)


⁶ Estimates based on an Al composition of the dust of 7.1% (Guieu et al. 2002)

310 3.3.2 Western Mediterranean Sea

At the FAST station, dissolved and particulate Al and Fe concentrations were measured at high temporal and vertical resolutions before, during, and after the wet deposition of dust (Supp. Fig. 4). About 6 h after deposition, the total (dissolved + particulate) Al and Fe inventories within the upper 20 m increased by ~ 145 and $48 \mu\text{mol m}^{-2}$, respectively (Fig. 4a and c). This increase in the 0-20 m inventories was consistent but higher than the atmospheric Al and Fe fluxes collected on the R/V

315 (~98 and 25 $\mu\text{mol m}^{-2}$, respectively; Desboeufs et al., in revision). Based on the increase in the 0-20 m total Al inventory and assuming 7.1% Al in the dust (Guieu et al., 2002), a total dust input of 55 mg m^{-2} was derived. Although direct collection of atmospheric dust aerosols represents the most straightforward approach for quantifying the dust flux, it has only a limited spatial coverage. In contrast, the upper water-column inventory integrated most of the patchy rainfalls associated with this large rain front. This difference in time and space integrations is best illustrated by the ~70% increase in the 0-20 m pAl and pFe inventories observed the 4th of June (Fig. 4c), i.e., several hours before the rainfall collected onboard the R/V and probably associated with surrounding precipitations. It must be noted that the water-column approach is also subject to uncertainties and we cannot exclude an under-estimation of the deposition flux due to the rapid sinking of the largest dust particles (e.g., Bressac et al., 2012). However, no evidence of these fast-sinking particles was found deeper in the water column (Fig. 4d), nor within the sediment traps (not shown).

325



330

Figure 4: Temporal evolution of the dissolved (a and b) and particulate (c and d) Al (black) and Fe (orange) inventories within the 0-20 m (left) and 0-200 m (right) depth ranges at the FAST station. Only dissolved inventories were measured the 3rd of June. The relative change in the coefficient of partitioning between the particulate and dissolved fractions (Kd) for Al and Fe is presented in e and f. Note that 0-200 m inventories measured the 4th of June (instead of the 2nd of June) were used for the initial Kd in f (see Sect. 4.1 for details). Grey vertical bars indicate the two dusty rain events that occurred in the FAST station area. The grey-dotted vertical line corresponds to the time of the dusty rainfall sampled on board the R/V (Desboeufs et al., in revision).

4 Discussion

335 4.1 Advective transport from continental margins in the central Mediterranean Sea


In the absence of direct atmospheric measurements, large uncertainties are associated with the estimates of the dust deposition flux over the Tyrrhenian Sea. These uncertainties are partly driven by potential additional sources of pAl, and in particular the resuspension of sediments and their advective transport from continental margins (e.g., Misic et al., 2008). The Strait of Sicily, characterized by high turbidity values (Gdaniec et al., 2018), represents a zone of formation for nepheloid layers. Levantine
340 intermediate waters can then act as a conveyor belt that accumulates and transports particles from the eastern to western basin of the Mediterranean Sea (Taillandier et al., 2020). Stations ST04 and ST05, located in the southwestern sector of the Tyrrhenian Sea (i.e., the branch of circulation between the Strait of Sicily and the Sardinian Channel), could be potentially impacted by particles driven by this mechanism, contributing to the excess in pAl observed at those stations. At the opposite, the central part of the Tyrrhenian Sea is characterized by low turbidity values relative to the rest of the Mediterranean Sea
345 (Gdaniec et al., 2018). During PEACETIME, lateral advection was negligible (A. Doglioli, pers. comm., 2020) at stations TYR and ST06 precluding any contribution of lithogenic particles other than from dust atmospheric deposition. Evidences of a recent dust deposition event over the Tyrrhenian Sea, traced by the excess in pAl, are discussed in the next section.

4.2 Dust dynamic in the water column

In the Tyrrhenian Sea, deposition of dust was evidenced by the >3 times higher Al and LSi downward fluxes measured at 200
350 and 1000 m depth ~5 to 9 days after deposition relative to those measured at ION and FAST at the same depths (Table 1). At TYR, Al and LSi fluxes both increased by 35% between 200 and 1000 m depth, suggesting that a significant fraction of the dust particles were rapidly transferred to depth. This trend is consistent with the pAl vertical profiles at the 4 stations likely impacted by this event, as a subsurface maximum was depicted between ~200 and 500 m depth (Fig. 3a-d). In addition, three pAl vertical profiles performed at TYR over ~72 h showed a continuous decrease in surface pAl concentration of $20 \mu\text{mol m}^{-2} \text{d}^{-1}$ that was accompanied by subsequent increases within the ~150-500 m depth layer (Fig. 5). It is worth noting that ~5.6 days after the event, a remarkably high pAl concentration was observed at 1000 m depth (~260 nmol kg⁻¹ (TYR_1); not shown), which could indicate that dust particles were sinking at a rate of ~180 m d⁻¹. This finding confirms that dust particles can be rapidly transferred to depth either alone (Bressac et al., 2012) or incorporated into biogenic aggregates (e.g., Hamm, 2002; Bressac et al., 2014; Laurenceau-Cornec et al., 2020; van der Jagt et al., 2018). Together, these observations demonstrate the
360 atmospheric origin of pAl observed in the southern Tyrrhenian (rather than sediment resuspension or advective inputs), and confirm that a significant fraction of the dust particles (coarse fraction) can rapidly leave the surface mixed-layer when the stratification is strong (Croot et al., 2004; TERNON et al., 2010; Nowald et al., 2015), while the remaining fraction (small-sized particles) likely accumulates along the thermocline until the disruption of the stratification (Migon et al., 2002).

At the FAST station, a two-orders of magnitude lower dust deposition flux ($\sim 55 \text{ mg m}^{-2}$) led to an increase by 78% of the 0-20 m pAl inventory (Fig. 4c). About 24 h after deposition, only $\sim 40\%$ of this signal was still present in the mixed-layer. This is consistent with a short residence time in surface water for a significant fraction of the dust, although we cannot exclude the effect of lateral advection (Guieu et al., 2020). Deeper in the water column, the trend is more complicated to interpret with a 40% decrease ($\sim 2000 \text{ } \mu\text{mol m}^{-2}$) of the 0-200 m pAl inventory that occurred before/during deposition (Fig. 4d). This unexpected decrease cannot be explained by the vertical transfer of pAl, as only $\sim 130 \text{ } \mu\text{mol m}^{-2}$ of pAl were exported out of the upper 200 m over 5 days (data not shown). On the other hand, a southwestward flow disrupted the water column in the ~ 25 -100 m depth range from the 3rd of June bringing water masses of distinct properties (Guieu et al., 2020). Therefore, it is likely that the water mass sampled before deposition (2nd of June) was different from the one sampled during the rest of the time-series. For this reason, inventories obtained the 4th of June (instead of the 2nd) were used as background level to investigate the temporal evolution of $K_d(\text{Al})$ and $K_d(\text{Fe})$ in the 0-200 m depth range (Fig. 4f).

375




380 **Figure 5: Time evolution of the pAl vertical distribution measured at three different times over ~ 72 h at the TYR station. Light-shaded areas correspond to the uncertainty in pAl measurement. The orange (TYR_1), red (TYR_2), and blue (TYR_3) vertical profiles were measured approximately 5.6, 6.4, and 8.4 days after the dust deposition event, respectively.**

4.3 Impact on the dAl inventory

4.3.1 Absence of dAl anomaly

A relatively large range in Al fractional solubility (1-15%; defined as the fraction of dust-derived Al that dissolves in rainwater or seawater) has been reported (e.g., Orians and Bruland, 1986; Baker et al., 2006; Measures et al., 2010; Han et al., 2012; Shelley et al., 2018). Assuming a conservative Al fractional solubility of 1.5% in seawater (Wuttig et al., 2013), dust deposition over the Tyrrhenian Sea led to a dAl input ranging between 68 and 363 $\mu\text{mol m}^{-2}$ (Table 2). Further assuming an homogeneous distribution within the 0-20 m mixed-layer, this dust event represented a dAl input of 3.3-17.7 nmol kg^{-1} . However, the absence of noticeable anomaly in the long-recognized relationship between surface dAl concentrations and salinity, when compared with published data (Rolison et al., 2015), reveals a limited or transient impact of this event on surface dAl concentrations (Fig. 6). Several mechanisms can be invoked here to explain the absence of dAl signal in the upper water column following the deposition. First, high surface dAl concentrations ($>20 \text{ nmol kg}^{-1}$) might mask any additional input. At ST06, the putative dAl input of 3.3 nmol kg^{-1} (Table 2) is within the range of variability of both published and observed surface concentrations (30-43 nmol kg^{-1} ; Fig. 3h). At the other end (ST04), the situation is different with a dAl input of 17.7 nmol kg^{-1} that would represent more than 55% of pre-depositional surface dAl concentrations (31-32 nmol kg^{-1} ; Rolison et al., 2015). Considering the short time lag between deposition and observations at ST04 (~ 3.6 days), and the low horizontal advection at the TYR station area (A. Doglioli, pers. comm., 2020), it is unlikely that advective mixing diluted any elevated dAl signal from this event. Deeper in the water column, no clear trend was obtained with subsurface dAl concentrations lower (ST04; Fig. 3e) or slightly higher than background levels (TYR; Fig. 3g). Similarly, no noticeable increase in dAl could be observed at the FAST station in the mixed-layer (Fig. 4a and 6). In contrast to $K_d(\text{Fe})$, $K_d(\text{Al})$ was still higher than pre-depositional value 4 days after deposition (Fig. 4e), potentially reflecting a lower fractional solubility for dust-derived Al relative to Fe, and/or higher removal rate for dAl. Below the mixed-layer, $K_d(\text{Al})$ remained relatively constant and similar to initial value (Fig. 4f). Together, these observations indicate that wet deposition of dust over the FAST station area had a limited impact on the dAl inventory.



405

Figure 6: Dissolved Al concentrations versus salinity measured in the upper 30 m of the Mediterranean Sea during the PEACETIME (filled circles), 64PE370 and 64PE374 cruises (open circles; Rolison et al., 2015). Yellow, orange, blue, red, and green circles correspond to data obtained at FAST, ST04, ST05, TYR and ST06, respectively (Fig. 1). Grey circles correspond to data obtained during the PEACETIME cruise at the other stations.

410

4.3.2 Drivers of the rapid removal of dAl

415

An Al fractional solubility of 5% was measured in rainwater for dust aerosols collected at the FAST station (Desboeufs et al., in revision), i.e., well above the conservative value of 1.5% obtained for a dust flux of 10 g m^{-2} (Wuttig et al., 2013) and used to estimate dAl inputs over the Tyrrhenian Sea. This confirms that dust loading partly controls the Al fractional solubility (e.g., Shelley et al., 2018). In addition, a biological control on dAl distribution in the Mediterranean Sea has been proposed to explain the strong coupling between dAl and orthosilicic acid (Si(OH)_4) in subsurface waters (Chou and Wollast, 1997; Rolison et al., 2015). Several laboratory and field studies have demonstrated that marine phytoplankton, in particular diatoms (mainly incorporated into the frustules; Gehlen et al., 2002), can take up and/or scavenge dAl (Mackenzie et al., 1978; Orians and Bruland, 1986; Moran and Moore, 1988b; Loucaides et al., 2010; Twining et al., 2015b; Wuttig et al., 2013; Liu et al., 2019). To investigate the respective role of particle adsorption and biological uptake in removing dAl, Al was compared to Fe – a particle-reactive and bioactive element (Tagliabue et al., 2017) predominantly of crustal origin in the Mediterranean Sea – through the Fe/Al content of suspended and sinking particles collected at different depth horizons (Fig. 7).

420

425

For suspended particles, the median Fe/Al ratio was maximum within the surface mixed-layer, and minimum at the DCM (60–100 m; Fig. 7a), highlighting a strong contrast in Fe/Al between the diatom-dominated particle assemblage at the DCM (Marañón et al., 2021) and detrital/lithogenic particles in the rest of the water column. This contrast supports the significant role played by phytoplankton, and in particular diatoms, in accumulating Al via active uptake (Gehlen et al., 2002; Liu et al., 2019) and/or adsorption onto cell membranes (Dammshäuser and Croot, 2012; Twining et al., 2015b). Note that we cannot exclude atmospheric input of BSi associated with dust particles (e.g., Folger et al., 1967). Regarding sinking particles, the

relationship between the Fe/Al ratio and the relative proportion of LSi and BSi was relatively well represented by a two-component model with BSi having a fixed Fe/Al ratio of 0.21 mol mol⁻¹ (value observed in the diatom-dominated DCM; Fig. 7a), and LSi having a Fe/Al ratio ranging between 0.26 (dusty rainwater sampled at FAST; Desboeufs et al., in revision) and 0.30 mol mol⁻¹ (Saharan dust end-member; Guieu et al., 2002) (Fig. 7b). Sparse Al/Si ratios available for natural diatom communities range between ~1 and 10 μmol mol⁻¹ (van Bennekom et al., 1989; Gehlen et al., 2002; Koning et al., 2007). Using this range of values, the Al downward flux at 200 m depth driven by (and incorporated into) BSi would represent only 0.1-1.1 μmol m⁻² d⁻¹, i.e., ~0.2-3.3% of the total Al flux. This two-orders of magnitude difference with our conservative estimates of dAl inputs over the Tyrrhenian Sea (68-371 μmol m⁻²; Table 2) indicates that adsorption onto biogenic particles (including BSi), rather than active uptake by diatoms, was likely the main sink for dAl in that region.




Figure 7: (a) Box whisker plot of the Fe/Al molar ratio (mol mol⁻¹) for suspended particles collected at TYR, ION, and FAST. The Fe/Al median values are 0.27 ($n = 37$; 0-20 m), 0.24 ($n = 13$; 20-60 m), 0.21 ($n = 17$; 60-100 m), 0.24 ($n = 12$; 100-200 m), 0.24 ($n = 9$; 200-500 m), and 0.24 ($n = 10$; 500-3000 m). For clarity, two outliers (Fe/Al = 0.50 and 1.02) observed in the 0-20 m depth range are not represented. (b) Fe/Al versus LSi/BSi molar ratios (mol mol⁻¹) of sinking particles collected at ~200 m (circles), 500 m (squares), and 1000 m depth (triangles) at the stations TYR (red), ION (purple), and FAST (yellow). The two black-dotted curves correspond to a two-component model with BSi having a fixed Fe/Al molar ratio of 0.21 mol mol⁻¹ (i.e., value observed in the DCM in (a)), and LSi having a fixed Fe/Al molar ratio of 0.26 (lower curve; Desboeufs et al., in revision) or 0.30 mol mol⁻¹ (upper curve; Guieu et al., 2002). Yellow-shaded areas in (a) and (b) represent the range in Fe/Al molar ratio proposed for Saharan dust, with the lower limit corresponding to dust aerosols collected at FAST (0.26 mol mol⁻¹; Desboeufs et al., in revision) and the upper limit to Saharan dust end-member (0.30 mol mol⁻¹; Guieu et al., 2002).

450 4.4 Vertical pattern in dFe input

4.4.1 Transient dFe increase in the surface mixed-layer

The absence of pre-depositional observations in the Tyrrhenian Sea is more problematic for Fe compared to Al, as no clear longitudinal trend has been reported in the Mediterranean Sea for that element. Dissolved Fe vertical profiles were thus compared to previously published data that were obtained at similar locations (Fig. 1), and at the same period of the year for
455 ST04 (mid-April) but about 2 months later at TYR and ST06 (early August) (Gerringa et al., 2017). Consequently, this approach ignores interannual and seasonal variabilities in dFe, and cannot be used to strictly quantify dFe input but remains valuable to investigate its magnitude and vertical distribution.

Assuming a Fe content of 4.45% in dust (Guieu et al., 2002), this dust event over the Tyrrhenian Sea represented a Fe input of
~1300-7000 $\mu\text{mol m}^{-2}$ (with a short retention time within the sea surface microlayer (Tovar-Sánchez et al., 2020)). Yet, dFe
460 concentrations within the surface mixed-layer were at background levels (ST04 and ST06) or slightly below (TYR) (Fig. 3i-l). These observations made ~3 to 10 days after deposition indicate that this event had no impact on dFe in the surface mixed-layer at a timescale of days. At a shorter timescale, sampling performed at a high temporal resolution at the FAST station revealed two distinct increases of the 0-20 m dFe inventory that occurred during (+13 $\mu\text{mol m}^{-2}$) and about 6 h after deposition (+15 $\mu\text{mol m}^{-2}$; Fig. 4a). These ~50% increases were only transient and the pre-depositional level was rapidly recovered.
465 Considering that Fe cycling in this LNLC system is dominated by physico-chemical rather than biological processes, our findings are consistent with rapid scavenging of dFe in surface Mediterranean waters following dust deposition, as already reported in some mesocosm and minicosm dust addition experiments (Wagener et al., 2010; Wuttig et al., 2013; Bressac and Guieu, 2013). Overall, the Fe-binding ligand pool is nearly saturated in surface Mediterranean waters (Gerringa et al., 2017). As a consequence, any new input of dFe will tend to precipitate, pointing to the importance of the initial dFe and Fe-binding
470 ligand concentrations in setting the net effect of dust input on dFe in the surface mixed-layer (Ye et al., 2011; Wagener et al., 2010; Wuttig et al., 2013).

4.4.2 Enrichment in dFe below the surface mixed-layer

A key feature in the southern Tyrrhenian was the systematic subsurface excess in dFe observed from ~40 m (ST04) and 200 m depth (TYR and ST06) (Fig. 3i-l), and mirroring the vertical distribution of $\text{Al}_{\text{excess}}$ (Fig. 3a-d). Similarly, wet dust deposition
475 over the FAST station area led to a net input of dFe mainly below the mixed-layer, as revealed by the opposite trends in $\text{Kd}(\text{Fe})$ observed in the 0-20 m and 0-200 m depth ranges (Fig. 4e-f). This increase in dFe relative to pFe was persistent on a timescale of days (Fig. 4f), and was primarily driven by dust dissolution (Fig. 4b) rather than ballasting of pre-existing pFe (Fig. 4d), as evidenced by the low export Fe flux collected at 200 m depth (1.7-12.3 $\mu\text{mol m}^{-2} \text{d}^{-1}$). This systematic excess in dFe observed below the mixed-layer and extending to 1000 m suggests that the mechanisms involved are independent of the dust flux – that
480 differed by two-orders of magnitude – and timescale considered (hours to weeks). Such dust-related subsurface enrichment in dFe (without enhanced surface dFe concentrations) has already been observed in the subarctic Pacific and tropical North

Atlantic. This feature was attributed either to low oxygen levels allowing Fe(II) to stay in solution (Schallenberg et al., 2017), or to remineralization of organic matter formed in the dust-laden surface ocean (Measures et al., 2008; Fitzsimmons et al., 2013); two mechanisms that cannot be invoked here considering the oxygen levels in subsurface (170-200 μM), the short
485 timescale considered, and the low mesopelagic Fe regeneration efficiency (Bressac et al., 2019).

To account for this dFe excess below the surface mixed-layer, dust-bearing Fe must continue to dissolve as dust particles settle through the mixed-layer and reach the mesopelagic. The short residence time for dust in surface water (Sect. 4.1), and the presence of a ‘refractory’ Fe pool within dust particles that dissolves over several days (Wagener et al., 2008) confirm that dust dissolution can occur in subsurface. It is also likely that the low particle concentration encountered at these depths relative
490 to the particle-rich surface waters at the time of deposition prevented rapid removal of dFe (e.g., Spokes et al., 1994; Bonnet and Guieu, 2004). Furthermore, the Fe-binding ligand pool is pivotal in setting the Fe fractional solubility (Rijkenberg et al., 2008; Wagener et al., 2008, 2010; Ye et al., 2011; Fishwick et al., 2014), and its magnitude, composition, and distribution likely shape patterns of dFe supply. While nearly saturated in surface, the Fe-binding ligand pool is in relatively large excess to dFe in subsurface Mediterranean waters (Gerringa et al., 2017), and hence available to stabilize new dFe. Importantly, this
495 subsurface pool is constantly replenished by bacterial degradation of sinking biogenic particles (Boyd et al., 2010; Velasquez et al., 2016; Bressac et al., 2019; Whitby et al., 2020). Thus, there is a permanent resetting of the ligand pool while dust particles settle (Bressac et al., 2019), and conceptually, we can imagine that the binding equilibrium between available ligands and Fe is rarely reached at these depths and timescale. This fundamental difference in the surface waters (and batch experiments) could explain the high Fe fractional solubility of 4.6-13.5% derived in the southern Tyrrhenian from the increase
500 in the 0-1000 m dFe inventories (relative to published data; Fig. 3i-l), and assuming 4.45% Fe in the dust (Guieu et al., 2002). By feeding the subsurface dFe reservoir, dust deposition could represent an indirect supply route for the surface ocean through vertical mixing and diapycnal diffusion (e.g., Tagliabue et al., 2014). However, the residence time of this dust-derived reservoir remains an open question. Relatively low subsurface dFe concentrations observed at the basin-scale ($<0.5 \text{ nmol kg}^{-1}$; Supp. Fig. 2c), compared to Atlantic waters for instance (Gerringa et al., 2017), argue in favour of a short residence time. Scavenging
505 by sinking (dust) particles (e.g., Wagener et al., 2010; Bressac et al., 2019), and bacterial removal of humic-like ligands (Dulaquais et al., 2018; Whitby et al., 2020) represent two potential sinks for this subsurface dFe reservoir that need to be explored.

5 Conclusions

During the PEACETIME cruise performed in May-June 2017 in the western and central Mediterranean, we observed two
510 atmospheric wet deposition events while measuring Al and Fe water-column distributions, providing important insights into the timescale and pattern of dAl and dFe inputs from African dust in the remote Mediterranean Sea. The water-column Al inventories were successfully utilised to assess dust deposition fluxes, complementing atmospheric measurements and the ‘fast-action’ strategy used during the campaign to directly sample dusty rain events. Our observations show that dAl removal

through adsorption onto biogenic particles was dominant over dAl released from dust at a timescale of hours to days. While
515 surface dAl concentrations reflect seasonal changes and large-scale patterns in dust deposition, this finding indicates that this
tracer may not be appropriate to trace the imprint of a single dust deposition event in highly dust-impacted areas. Furthermore,
dust deposition represented a significant input of dFe in the surface mixed-layer only on a timescale of hours. On a longer
timescale (days/weeks), dFe inputs occurred primarily below the surface mixed-layer extending to 1000 m depth where the
Fe-binding ligand pool likely in excess of dFe allows stabilizing any additional input of dFe. This mechanism may represent
520 an additional pathway of dFe resupply for the surface ocean (through vertical mixing and diapycnal diffusion), although the
residence time of this dust-derived dFe reservoir still needs to be investigated.

Data availability

Underlying research data are being used by researcher participants of the PEACETIME campaign to prepare other papers, and
therefore data are not publicly accessible at the time of publication. Data will be accessible once the special issue is completed
525 (June 2021) (<http://www.obs-vlfr.fr/proof/php/PEACETIME/peacetime.php>; last access 25/10/2021). The policy of the
database is detailed here: <http://www.obs-vlfr.fr/proof/dataconvention.php> and <https://www.seanoe.org/data/00645/75747/>
(last access 25/10/2021).

Author contribution

M.B., T.W., and C.G. designed the study. M.B. wrote the manuscript. M.B., T.W., N.L., A.T.-S., C.R., V.T., S.A., S.G., and
530 A.D. collected and/or analysed the samples. F.D. and K.D. interpreted the atmospheric data. All the authors commented on
and contributed to the improvement of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

535 We thank the captain and crew of the R/V *Pourquoi Pas?*, the DT INSU for the design and preparation of the mooring line,
N. Bhairy, G. De Liège, and G. Rougier for their help with the mooring. Hélène Ferré and the AERIS/SEDOO service are
acknowledged for real-time collection during the cruise of maps produced from operational satellites and models used in this
study, with appreciated contributions of EUMETSAT and ICARE for MSG/SEVIRI products, Météo-France for ARPEGE
model outputs, the WMO SDS-WAS operated by the Barcelona Supercomputing Center (BSC) for DREAM and NMMB
540 model outputs, and the AM&WFG of the University of Athens for SKIRON model outputs. This study is a contribution to the

PEACETIME project (<http://peacetime-project.org>; last access 25/10/2021), a joint initiative of the MERMEX and ChArMEX components supported by CNRS-INSU, IFREMER, CEA and Météo-France as part of the decadal programme MISTRALS coordinated by INSU. PEACETIME was endorsed as a process study by GEOTRACES and is also a contribution to IMBER and SOLAS international programs. M.B. was funded by a Marie Skłodowska-Curie Postdoctoral European Fellowship (European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. PIOF-GA-2012-626734 (IRON-IC project)). S.A. acknowledges funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement 708119, for the project “DUSSt, Climate, and the Carbon Cycle” (DUSC3).

References

- Acker, J. G., and Leptoukh, G.: Online analysis enhances use of NASA Earth science data, *Eos Trans. AGU*, 88(2), 14–17, <https://doi.org/10.1029/2007EO020003>, 2007.
- Anderson, R. F., Cheng, H., Edwards, R. L., Fleisher, M. Q., Hayes, C. T., Huang, K.-F., Kadko, D., Lam, P. J., Landing, W. M., Lao, Y., Lu, Y., Measures, C. I., Moran, S. B., Morton, P. L., Ohnemus, D. C., Robinson, L. F., and Shelley, R. U.: How well can we quantify dust deposition to the ocean?, *Philos. Trans. R. Soc. A*, 374(2081), 20150285, <https://doi.org/10.1098/rsta.2015.0285>, 2016.
- Baker, A. R., and Croot P. L.: Atmospheric and marine controls on aerosol iron solubility in seawater, *Mar. Chem.*, 120(1–4), 4–13, <https://doi.org/10.1016/j.marchem.2008.09.003>, 2010.
- Baker, A. R., Jickells, T. D., Witt, M., and Linge, K. L.: Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean, *Mar. Chem.*, 98(1), 43–58, <https://doi.org/10.1016/j.marchem.2005.06.004>, 2006.
- Bonnet, S. and Guieu, C.: Atmospheric forcing on the annual iron cycle in the Mediterranean Sea. A one-year survey, *J. Geophys. Res.*, 111(C9), C09010, <https://doi.org/10.1029/2005JC003213>, 2006.
- Bonnet, S., and Guieu, C.: Dissolution of atmospheric iron in seawater, *Geophys. Res. Lett.*, 31(3), L03303, <https://doi.org/10.1029/2003GL018423>, 2004.
- Boyd, P. W., Ibsanmi, E., Sander, S. G., Hunter, K. A., and Jackson, G. A.: Remineralization of upper ocean particles: Implications for iron biogeochemistry, *Limnol. Oceanogr.*, 55(3), 1271–1288, <https://doi.org/10.4319/lo.2010.55.3.1271>, 2010.
- Bressac, M., and Guieu, C.: Post-depositional processes: What really happens to new atmospheric iron in the ocean surface?, *Global Biogeochem. Cycles*, 27, 859–870, <https://doi.org/10.1002/gbc.20076>, 2013.

- 570 Bressac, M., Guieu, C., Doxaran, D., Bourrin, F., Desboeufs, K., Leblond, N., and Ridame, C.: Quantification of the lithogenic carbon pump following a simulated dust-deposition event in large mesocosms, *Biogeosciences*, 11, 1007–1020, <https://doi.org/10.5194/bg-11-1007-2014>, 2014.
- Bressac, M., Guieu, C., Doxaran, D., Bourrin, F., Obolensky, G., and Grisoni, J. M.: A mesocosm experiment coupled with optical measurements to assess the fate and sinking of atmospheric particles in clear oligotrophic waters, *Geo-Mar. Lett.*, 32, 153–164, <https://doi.org/10.1007/s00367-011-0269-4>, 2012.
- 575 Bressac, M., Guieu, C., Ellwood, M. J., Tagliabue, A., Wagener, T., Laurenceau Cornec, E. C., Whitby, H., Sarthou, G., and Boyd, P. W.: Resupply of mesopelagic dissolved iron controlled by particulate iron composition, *Nat. Geosci.*, 12, 995–1000, <https://doi.org/10.1038/s41561-019-0476-6>, 2019.
- Buat-Ménard, P., Davies, J., Remoudaki, E., Miquel, J. C., Bergametti, G., Lambert, C. E., Ezat, U., Quétel, C., La Rosa, J., and Fowler, S. W.: Non-steady-state biological removal of atmospheric particles from Mediterranean surface waters. *Nature* 340, 131–134, <https://doi.org/10.1038/340131a0>, 1989.
- 580 Chou, L., and Wollast, R.: Biogeochemical behavior and mass balance of dissolved aluminum in the western Mediterranean Sea, *Deep-Sea Res. II*, 44(3), 741–768, [https://doi.org/10.1016/S0967-0645\(96\)00092-6](https://doi.org/10.1016/S0967-0645(96)00092-6), 1997.
- Christensen, J. H.: The Danish eulerian hemispheric model - A three-dimensional air pollution model used for the Arctic, *Atmos. Environ.*, 31(24), 4169–4191, [https://doi.org/10.1016/S1352-2310\(97\)00264-1](https://doi.org/10.1016/S1352-2310(97)00264-1), 1997.
- 585 Conway, T. M., and John, S. G.: Quantification of dissolved iron sources to the North Atlantic Ocean, *Nature*, 511(7508), 212–215, <https://doi.org/10.1038/nature13482>, 2014.
- Croot, P. L., Streu, P., and Baker, A. R.: Short residence time for iron in surface seawater impacted by atmospheric dry deposition from Saharan dust events, *Geophys. Res. Lett.*, 31(23), L23S08, <https://doi.org/10.1029/2004GL020153>, 2004.
- Cutter, G. A., Andersson, P., Codispoti, L., Croot, P., Francois, R., Lohan, M., Obata, H., and Rutgers van der Loeff, M.: Sampling and Sample-handling Protocols for GEOTRACES Cruises. 2010 GEOTRACES Standards and Intercalibration
- 590 Committee, 2010.
- Dammshäuser, A., and Croot, P. L.: Low colloidal associations of aluminium and titanium in surface waters of the tropical Atlantic, *Geochim. Cosmochim. Acta*, 96, 304–318, <https://doi.org/10.1016/j.gca.2012.07.032>, 2012.
- Dammshäuser, A., Wagener, T., and Croot, P. L.: Surface water dissolved aluminum and titanium: Tracers for specific time scales of dust deposition to the Atlantic?, *Geophys. Res. Lett.*, 38(24), L24601, <https://doi.org/10.1029/2011GL049847>, 2011.
- 595 Davies, J. E. and Buat-Ménard, P.: Impact of atmospheric deposition on particulate manganese and aluminium distribution in northwestern Mediterranean surface water, *Global Planet. Change*, 3(1-2), 35–45, [https://doi.org/10.1016/0921-8181\(90\)90054-G](https://doi.org/10.1016/0921-8181(90)90054-G), 1990.

- de Leeuw, G., Guieu, C., Arneth, A., Bellouin, N., Bopp, L., Boyd, P. W., Denier van der Gon, H. A. C., Desboeufs, K. V., Dulac, F., Cristina Facchini, M. C., Gantt, B., Langmann, B., Mahowald, N. M., Marañón, E., O'Dowd, C., Olgun, N., Pulido-Villena, E., Rinaldi, M., Stephanou, E.G., and Wagener, T.: Ocean-Atmosphere interactions of particles, in: Ocean-Atmosphere Interactions of Gases and Particles, edited by: Liss, P., and Johnson, M., , Springer Earth System Sciences, Springer Berlin Heidelberg, 171–246, https://doi.org/10.1007/978-3-642-25643-1_4, 2014.
- Desboeufs, K., Bon Nguyen, E., Chevaillier, S., Triquet, S., and Dulac, F.: Fluxes and sources of nutrients and trace metal atmospheric deposition in the northwestern Mediterranean, *Atmos. Chem. Phys.*, 18, 14477–14492, <https://doi.org/10.5194/acp-18-14477-2018>, 2018.
- Desboeufs, K., Fu, F., Bressac, M., Tovar-Sánchez, A., Triquet, S., Doussin, J. F., Giorio, C., Chazette, P., Dinasquet, J., Feron, A., Formenti, P., Maisonneuve, F., Rodriguez-Romero, A., Zapf, P., Dulac, F., and Guieu, C. : Wet deposition in the remote western and central Mediterranean as a source of trace metals to surface seawater, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2021-624>, 2021.
- Donaghay, P. L., Liss, P.S., Duce, R.A., Kester, D.R., Hanson, A.K., Villareal, T., Tindale, N.W., and Gifford, D.J.: The role of episodic atmospheric nutrient inputs in the chemical and biological dynamics of oceanic ecosystems, *Oceanography*, 4, 62–70, <https://doi.org/10.5670/oceanog.1991.04>, 1991.
- Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Menard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., Hansen, L., Jickells, T. D., Knap, A. H., Reinhardt, K. H., Schneider, B. Soudine, A., Tokos, J. J., Tsunogai, S., Wollast, R., and Zhou, M.: The atmospheric input of trace species to the world ocean, *Global Biogeochem. Cycles*, 5, 193–259, <https://doi.org/10.1029/91GB01778>, 1991.
- Dulac, F., Tanré, D., Bergametti, G., Buat-Ménard, P., Desbois, M., and Sutton, D.: Assessment of the African airborne dust mass over the western Mediterranean Sea using Meteosat data, *J. Geophys. Res.: Atmospheres*, 97(D2), 2489–2506, <https://doi.org/10.1029/91JD02427>, 1992.
- Dulaquais, G., Waeles, M., Gerringa, L. J., Middag, R., Rijkenberg, M. J., and Riso, R.: The biogeochemistry of electroactive humic substances and its connection to iron chemistry in the North East Atlantic and the Western Mediterranean Sea, *J. Geophys. Res.: Oceans*, 123(8), 5481–5499, <https://doi.org/10.1029/2018JC014211>, 2018.
- Fishwick, M. P., Sedwick, P. N., Lohan, M. C., Worsfold, P. J., Buck, K. N., Church, T. M., and Ussher, S. J.: The impact of changing surface ocean conditions on the dissolution of aerosol iron, *Global Biogeochem. Cycles*, 28(11), 1235–1250, <https://doi.org/10.1002/2014GB004921>, 2014.
- Fitzsimmons, J.N., Zhang, R., and Boyle, E.A.: Dissolved iron in the tropical North Atlantic Ocean, *Mar. Chem.*, 154, 87–99, <https://doi.org/10.1016/j.marchem.2013.05.009>, 2013.

- Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E., Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and
630 Tsikerdekis, A.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, *Geosci. Model Dev.*, 8(4), 975–1003, <https://doi.org/10.5194/gmd-8-975-2015>, 2015.
- Folger, D. W., Burekle, L. H., and Heezen, B. C.: Opal phytoliths in a North Atlantic dust fall, *Science*, 155(3767), 1243–1244, <https://doi.org/10.1126/science.155.3767.1243>, 1967.
- Gazeau, F., Ridame, C., VanWambeke, F., Alliouane, S., Stolpe, C., Irisson, J.-O., Marro, S., Grisoni, J.-M., De Liège, G.,
635 Nunige, S., Djaoudi, K., Pulido-Villena, E., Dinasquet, J., Obernosterer, I., Catala, P., and Guieu, C.: Impact of dust addition on Mediterranean plankton communities under present and future conditions of pH and temperature: an experimental overview, *Biogeosciences*, 18, 5011–5034, <https://doi.org/10.5194/bg-18-5011-2021>, 2021.
- Gdaniec, S., Roy-Barman, M., Foliot, L., Thil, F., Dapoigny, A., Burckel, P., Garcia-Orellana, J., Masqué, P., Mörth, C.-M., and Andersson, P. S.: Thorium and protactinium isotopes as tracers of marine particle fluxes and deep water circulation in the
640 Mediterranean Sea, *Mar. Chem.*, 199, 12–23, <https://doi.org/10.1016/j.marchem.2017.12.002>, 2018.
- Gehlen, M., Beck, L., Calas, G., Flank, A.-M., Van Bennekom, A. J., and Van Beusekom, J. E. E. : Unraveling the atomic structure of biogenic silica: evidence of the structural association of Al and Si in diatom frustules, *Geochim. Cosmochim. Acta*, 66(9), 1601–1609, [https://doi.org/10.1016/S0016-7037\(01\)00877-8](https://doi.org/10.1016/S0016-7037(01)00877-8), 2002.
- Gerringa, L. J. A., Slagter, H. A., Bown, J., van Haren, H., Laan, P., De Baar, H. J. W., and Rijkenberg, M. J. A.: Dissolved
645 Fe and Fe-binding organic ligands in the Mediterranean Sea–GEOTRACES G04, *Mar. Chem.*, 194, 100–113, <https://doi.org/10.1016/j.marchem.2017.05.012>, 2017.
- Gkikas, A., Basart, S., Hatzianastassiou, N., Marinou, E., Amiridis, V., Kazadzis, S., Pey, J., Querol, X., Jorba, O., Gassó, S., and Baldasano, J. M.: Mediterranean intense desert dust outbreaks and their vertical structure based on remote sensing data, *Atmos. Chem. Phys.*, 16, 8609–8642, <https://doi.org/10.5194/acp-16-8609-2016>, 2016.
- 650 Guertzoni, S., Chester, R., Dulac, F., Herut, B., Loÿe-Pilot, M. D., Measures, C., Migon, C., Molinaroli, E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, *Prog. Oceanogr.*, 44, 147–190, [https://doi.org/10.1016/S0079-6611\(99\)00024-5](https://doi.org/10.1016/S0079-6611(99)00024-5), 1999.
- Guieu, C., Loÿe-Pilot, M.D., Ridame, C., and Thomas, C.: Chemical characterization of the Saharan dust end-member; some biological implications for the western Mediterranean, *J. Geophys. Res.*, 107 (D15), 4258,
655 <https://dx.doi.org/10.1029/2001JD000582>, 2002.
- Guieu, C., Roy-Barman, M., Leblond, N., Jeandel, C., Souhaut, M., Le Cann, B., Dufour, A., and Bournot, C.: Vertical particle flux in the northeast Atlantic Ocean (POMME experiment), *J. Geophys. Res. Oceans*, 110(C7), CO7S18, <https://doi.org/10.1029/2004JC002672>, 2005.

- 660 Guieu, C., Aumont, O., Paytan, A., Bopp, L., Law, C. S., Mahowald, N., Achterberg, E. P., Marañón, E., Salihoglu, B., Crise, A., Wagener, T., Herut, B., Desboeufs, K., Kanakidou, M., Olgun, N., Peters, F., Pulido-Villena, E., Tovar-Sanchez, A., and Völker, C.: The significance of episodicity in atmospheric deposition to Low Nutrient Low Chlorophyll regions, *Global Biogeochem. Cycles.*, 28, 1179–1198, <https://doi.org/10.1002/2014GB004852>, 2014a.
- 665 Guieu, C., Ridame, C., Pulido-Villena, E., Bressac, M., Desboeufs, K., and Dulac, F.: Impact of dust deposition on carbon budget: a tentative assessment from a mesocosm approach, *Biogeosciences*, 11, 5621–5635, <https://doi.org/10.5194/bg-11-5621-2014>, 2014b.
- Guieu, C., D'Ortenzio, F., Dulac, F., Taillandier, V., Doglioli, A., Petrenko, A., Barrillon, S., Mallet, M., Nabat, P., and Desboeufs, K.: Introduction: Process studies at the air–sea interface after atmospheric deposition in the Mediterranean Sea—objectives and strategy of the PEACETIME oceanographic campaign (May–June 2017), *Biogeosciences*, 17(22), 5563–5585, <https://doi.org/10.5194/bg-17-5563-2020>, 2020.
- 670 Hamm, C. E.: Interactive aggregation and sedimentation of diatoms and clay-sized lithogenic material, *Limnol. Oceanogr.*, 47(6), 1790–1795, <https://doi.org/10.4319/lo.2002.47.6.1790>, 2002.
- Han, Q., Moore, J. K., Zender, C., Measures, C., and Hydes, D.: Constraining oceanic dust deposition using surface ocean dissolved Al, *Global Biogeochem. Cycles*, 22 (2), <https://doi.org/10.1029/2007GB002975>, 2008.
- 675 Han, Q., Zender, C., Moore, J. K., Buck, C. S., Chen, Y., Johansen, A., and Measures, C.: Global estimates of mineral dust aerosol iron and aluminum solubility that account for particle size using diffusion-controlled and surface-area controlled approximations, *Global Biogeochem. Cycles*, 26, GB2038, <https://doi.org/10.1029/2011GB004186>, 2012.
- Han, J., Wang, W., Kwon, Y.C., Hong, S.Y., Tallapragada, V. and Yang, F., 2017. Updates in the NCEP GFS cumulus convection schemes with scale and aerosol awareness, *Weather Forecast.*, 32(5), 2005–2017, <https://doi.org/10.1175/WAF-D-17-0046.1>, 2017.
- 680 Heimbürger, L.E., Migon, C. and Cossa, D. Impact of atmospheric deposition of anthropogenic and natural trace metals on Northwestern Mediterranean surface waters: A box model assessment, *Environ. Pollut.*, 159(6), 1629–1634, <https://doi.org/10.1016/j.envpol.2011.02.046>, 2011.
- 685 Herut, B., Rahav, E., Tsagaraki, T. M., Giannakourou, A., Tsiola, A., Psarra, S., Lagaria, A., Papageorgiou, N., Mihalopoulos, N., Theodosi, C. N., Violaki, K., Stathopoulou, E., Scoullou, M., Krom, M. D., Stockdale, A., Shi, Z., Berman-Frank, I., Meador, T. B., Tanaka, T., and Paraskevi, P.: The potential impact of Saharan dust and polluted aerosols on microbial populations in the East Mediterranean Sea, an overview of a mesocosm experimental approach, *Frontiers Mar. Sci.*, 3, 226, <https://doi.org/10.3389/fmars.2016.00226>, 2016.
- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., and Tan, J. : Integrated Multi-satellitE Retrievals for GPM (IMERG) technical documentation, NASA/GSFC Code, 612(47), 2015.

- 690 Huneus, N., Basart, S., Fiedler, S., Morcrette, J.-J., Benedetti, A., Mulcahy, J., Terradellas, E., Pérez García-Pando, C., Pejanovic, G., Nickovic, S., Arsenovic, P., Schulz, M., Cuevas, E., Baldasano, J. M., Pey, J., Remy, S., and Cvetkovic, B.: Forecasting the northern African dust outbreak towards Europe in April 2011: a model intercomparison, *Atmos. Chem. Phys.*, 16, 4967–4986, <https://doi.org/10.5194/acp-16-4967-2016>, 2016.
- Hydes, D. and Liss, P.: Fluorimetric method for the determination of low concentrations of dissolved aluminium in natural
695 waters, *Analyst*, 101, 922–931, <https://doi.org/10.1039/AN9760100922>, 1976.
- Jacquet, S. H. M., Dehairs, F., Lefèvre, D., Cavagna, A. J., Planchon, F., Christaki, U., Monin, L., André, L., Closset, I., and Cardinal D.: Early Spring Mesopelagic Carbon Remineralization and Transfer Efficiency in the Naturally Iron-Fertilized Kerguelen Area, *Biogeosciences*, 12(6), 1713–1731, <https://doi.org/10.5194/bg-12-1713-375>, 2015.
- Jacquet, S. H. M., Tamburini, C., Garel, M., Dufour, A., Van-Vambeke, F., Le Moigne, F. A. C., Bhairy, N., and Guasco, S.:
700 Particulate biogenic barium tracer of mesopelagic carbon remineralization in the Mediterranean Sea (PEACETIME project), *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2020-271>, 2020.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Science*, 308(5718), 67–71,
705 <https://doi.org/10.1126/science.1105959>, 2005.
- Kishcha P., Nickovic S., Ganor E., Kordova L., and Alpert P.: Saharan Dust over the Eastern Mediterranean: Model Sensitivity, in: *Air Pollution Modeling and Its Application XIX*, edited by: Borrego C., and Miranda A.I., NATO Science for Peace and Security Series Series C: Environmental Security, Springer, Dordrecht, https://doi.org/10.1007/978-1-4020-8453-9_39, 2008.
- Koning, E., Gehlen, M., Flank, A.-M., Calas, G., and Epping, E.: Rapid post-mortem incorporation of aluminum in diatom
710 frustules: evidence from chemical and structural analyses, *Mar. Chem.*, 106(1), 208–222, <https://doi.org/10.1016/j.marchem.2006.06.009>, 2007.
- Kubilay, N., Nickovic, S., Moulin, C., and Dulac, F.: An illustration of the transport and deposition of mineral dust onto the eastern Mediterranean, *Atmos. Environ.*, 34(8), 1293–1303, [https://doi.org/10.1016/S1352-2310\(99\)00179-X](https://doi.org/10.1016/S1352-2310(99)00179-X), 2000.
- Laurenceau-Cornec, Le Moigne, F. A. C., Gallinari, M., Moriceau, B., Toullec, J., Iversen, M. H., Engel, A., and De La Rocha,
715 C. L.: New guidelines for the application of Stokes' models to the sinking velocity of marine aggregates, *Limnol. Oceanogr.*, 65(6), 1264–1285, <https://doi.org/10.1002/lno.11388>, 2020.
- Li, F., Ren, J., Yan, L., Liu, S., Liu, C., Zhou, F., and Zhang, J.: The biogeochemical behavior of dissolved aluminum in the southern Yellow Sea: Influence of the spring phytoplankton bloom, *Chin. Sci. Bull.*, 58, 238–248, <https://doi.org/10.1007/s11434-012-5512-5>, 2013.

- 720 Liu, Q., Zhou, L., Liu, F., Fortin, C., Tan, Y., Huang, L., and Campbell, P. G.: Uptake and subcellular distribution of aluminum in a marine diatom, *Ecotoxicology and environmental safety*, 169, 85–92, <https://doi.org/10.1016/j.ecoenv.2018.10.095>, 2019.
- Loucaides, S., Michalopoulos, P., Presti, M., Koning, E., Behrends, T., and Van Cappellen, P.: Seawater-mediated interactions between diatomaceous silica and terrigenous sediments: results from long-term incubation experiments, *Chem. Geol.*, 270(1-4), 68-79, <https://doi.org/10.1016/j.chemgeo.2009.11.006>, 2010.
- 725 Loÿe-Pilot, M.-D., and Martin, J.-M.: Saharan dust input to the western Mediterranean: an eleven years record in Corsica, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S., and Chester, R., Environmental Science and Technology Library, 11, 191–199, Springer, Dordrecht, https://doi.org/10.1007/978-94-017-3354-0_18, 1996.
- Mackenzie, F. T., Stoffyn, M., and Wollast, R.: Aluminum in seawater: control by biological activity, *Science*, 199(4329), 680–682, <https://doi.org/10.1126/science.199.4329.680>, 1978.
- 730 Marañón, E., Van Wambeke, F., Uitz, J., Boss, E. S., Dimier, C., Dinasquet, J., Engel, A., Haëntjens, N., Pérez-Lorenzo, M., Taillandier, V., and Zäncker, B.: Deep maxima of phytoplankton biomass, primary production and bacterial production in the Mediterranean Sea, *Biogeosciences*, 18(5), 1749–1767, <https://doi.org/10.5194/bg-18-1749-2021>, 2021.
- Measures, C. I., and Brown, E. T.: Estimating Dust Input to the Atlantic Ocean Using Surface Water Aluminium Concentrations, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S., and Chester, R., Environmental Science and Technology Library, 11, 301–311, Springer, Dordrecht, https://doi.org/10.1007/978-94-017-3354-0_30, 1996.
- 735 Measures, C. I., Landing, W. M., Brown, M. T., Buck, C. S.: High-resolution Al and Fe data from the Atlantic Ocean CLIVAR-CO2 repeat hydrography A16N transect: extensive linkages between atmospheric dust and upper ocean geochemistry, *Global Biogeochem. Cycles*, 22(1), GB1005, <https://doi.org/10.1029/2007GB003042>, 2008.
- 740 Measures, C., Sato, T., Vink, S., Howell, S., and Li, Y.: The fractional solubility of aluminium from mineral aerosols collected in Hawaii and implications for atmospheric deposition of biogeochemically important trace elements, *Mar. Chem.*, 120(1–4), 144–153, <https://doi.org/10.1016/j.marchem.2009.01.014>, 2010.
- Menzel Barraqueta, J. L., Klar, J. K., Gledhill, M., Schlosser, C., Shelley, R., Planquette, H. F., Wenzel, B., Sarthou, G., and Achterberg, E. P.: Atmospheric deposition fluxes over the Atlantic Ocean: a GEOTRACES case study, *Biogeosciences*, 16(7), 1525–1542, <https://doi.org/10.5194/bg-16-1525-2019>, 2019.
- 745 Meskhidze, N., Völker, C., Al-Abadleh, H. A., Barbeau, K., Bressac, M., Buck, C., Bundy, R. M., Croot, P., Feng, Y., Ito, A., Johansen, A. M., Landing, W. M., Mao, J., Myriokefalitakis, S., Ohnemus, D., Pasquier, B., and Ye, Y.: Perspective on identifying and characterizing the processes controlling iron speciation and residence time at the atmosphere-ocean interface, *Mar. Chem.*, 217, 103704, <https://doi.org/10.1016/j.marchem.2019.103704>, 2019.

- 750 Middag, R., van Hulten, M. M. P., van Aken, H. M., Rijkenberg, M. J. A., Gerringa, L. J. A., Laan, P., and de Baar, H. J. W.: Dissolved aluminium in the ocean conveyor of the West Atlantic Ocean: mirror image of the biological cycle?, *Mar. Chem.*, 177, 69–86, <https://doi.org/10.1016/j.marchem.2015.02.015>, 2015.
- Migon, C., Sandroni, V., Marty, J. C., Gasser, B., and Miquel, J. C.: Transfer of atmospheric matter through the euphotic layer in the northwestern Mediterranean: seasonal pattern and driving forces, *Deep-Sea Res. II*, 49(11), 2125–2141, [https://doi.org/10.1016/S0967-0645\(02\)00031-0](https://doi.org/10.1016/S0967-0645(02)00031-0), 2002.
- 755 Misic, C., Castellano, M., Ruggieri, N., and Harriague, A. C.: Variations in ectoenzymatic hydrolytic activity in an oligotrophic environment (Southern Tyrrhenian Sea, W Mediterranean), *J. Mar. Syst.*, 73(1-2), 123–137, <https://doi.org/10.1016/j.jmarsys.2007.10.003>, 2008.
- Moore, C. M., Mills, M., Arrigo, K., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6(9), 701–710, <https://doi.org/10.1038/ngeo1765>, 2013.
- 760 Moran, S. B., Moore, R. M.: Temporal variations in dissolved and particulate aluminium during a spring bloom, *Estuar. Coast. Shelf Sci.*, 27, 205–215, [https://doi.org/10.1016/0272-7714\(88\)90090-X](https://doi.org/10.1016/0272-7714(88)90090-X), 1988a.
- 765 Moran, S. B., and Moore, R. M.: Evidence from mesocosm studies for biological removal of dissolved aluminium from sea water, *Nature*, 335(6192), 706–708, <https://doi.org/10.1038/335706a0>, 1988b.
- Mullin, J., and Riley, J. P.: The colorimetric determination of silicate with special reference to sea and natural waters, *Anal. Chim. Acta*, 12, 162-176, [https://doi.org/10.1016/S0003-2670\(00\)87825-3](https://doi.org/10.1016/S0003-2670(00)87825-3), 1955.
- Nelson, D. M., Smith Jr, W. O., Muench, R. D., Gordon, L. I., Sullivan, C. W., and Husby, D. M.: Particulate matter and nutrient distributions in the ice-edge zone of the Weddell Sea: relationship to hydrography during late summer, *Deep Sea Res. I*, 36(2), 191–209, [https://doi.org/10.1016/0198-0149\(89\)90133-7](https://doi.org/10.1016/0198-0149(89)90133-7), 1989.
- 770 Nowald, N., Iversen, M. H., Fischer, G., Ratmeyer, V., and Wefer, G.: Time series of in-situ particle properties and sediment trap fluxes in the coastal upwelling filament off Cape Blanc, Mauritania, *Progr. Oceanogr.*, 137A, 1–11, <https://doi.org/10.1016/j.pocean.2014.12.015>, 2015.
- 775 Orians, K. J., and Bruland, K. W.: The biogeochemistry of aluminum in the Pacific Ocean, *Earth Planet. Sci. Lett.*, 78(4), 397–410, [https://doi.org/10.1016/0012-821X\(86\)90006-3](https://doi.org/10.1016/0012-821X(86)90006-3), 1986.
- Planquette, H., and Sherrell, R. M.: Sampling for particulate trace element determination using water sampling bottles: methodology and comparison to in situ pumps, *Limnol. Oceanogr.: Methods*, 10(5), 367–388, <https://doi.org/10.4319/lom.2012.10.367>, 2012.

- 780 Quétel, C. R., Remoudaki, E., Davies, J. E., Miquel, J. C., Fowler, S. W., Lambert, C. E., Bergametti, G. and Buat-Ménard, P.: Impact of atmospheric deposition on particulate iron flux and distribution in northwestern Mediterranean waters, *Deep Sea Res. I*, 40(5), 989–1002, [https://doi.org/10.1016/0967-0637\(93\)90085-H](https://doi.org/10.1016/0967-0637(93)90085-H), 1993.
- Rijkenberg, M., Powell, C., Dall'Osto, M., Nielsdottir, M., Patey, M., Hill, P., Baker, A., Jickells, T., Harrison, R., and Achterberg, E.: Changes in iron speciation following a Saharan dust event in the tropical North Atlantic Ocean, *Mar. Chem.*, 110, 56–67, <https://doi.org/10.1016/j.marchem.2008.02.006>, 2008.
- 785 Rolison, J. M., Middag, R., Stirling, C. H., Rijkenberg, M. J. A., and De Baar, H. J. W.: Zonal distribution of dissolved aluminium in the Mediterranean Sea, *Mar. Chem.*, 177, 87–100, <https://doi.org/10.1016/j.marchem.2015.05.001>, 2015.
- Sarrand, B., Dulac, F., Baldi, M., Bargaoui, Z., Cindrić, K., Ducrocq, V., Labiadh, M., Schiavone, J., de Silvestri, L., Somot, S., and Tovar-Sánchez, A.: Precipitation in the Mediterranean basin as seen from the 2000-2010 TRMM-3B42-v6 database, *EGU General Assembly Conference Abstracts*, p.11965, 2012.
- 790 Sarthou, G., and Jeandel, C.: Seasonal variations of iron concentrations in the Ligurian Sea and iron budget in the Western Mediterranean Sea, *Mar. Chem.*, 74(2-3), 115–129, [https://doi.org/10.1016/S0304-4203\(00\)00119-5](https://doi.org/10.1016/S0304-4203(00)00119-5), 2001.
- Schallenberg, C., Ross, A. R., Davidson, A. B., Stewart, G. M., and Cullen, J. T.: Temporal variability of dissolved iron species in the mesopelagic zone at Ocean Station PAPA, *J. Mar. Syst.*, 172, 128–136, <https://doi.org/10.1016/j.jmarsys.2017.03.006>, 2017.
- 795 Shelley, R. U., Landing, W. M., Ussher, S. J., Planquette, H., and Sarthou, G.: Regional trends in the fractional solubility of Fe and other metals from North Atlantic aerosols (GEOTRACES cruises GA01 and GA03) following a two-stage leach, *Biogeosciences*, 15(8), 2271–2288, <https://doi.org/10.5194/bg-15-2271-2018>, 2018.
- Spokes, L. J., Jickells, T. D., and Lim, B.: Solubilisation of aerosol trace metals by cloud processing: A laboratory study, *Geochim. Cosmochim. Acta*, 58, 3281–3287, [https://doi.org/10.1016/0016-7037\(94\)90056-6](https://doi.org/10.1016/0016-7037(94)90056-6), 1994.
- 800 Spyrou, C., Mitsakou, C., Kallos, G., Louka, P., and Vlastou, G.: An improved limited area model for describing the dust cycle in the atmosphere, *J. Geophys. Res.*, 115(D17), D17211, <https://doi.org/10.1029/2009JD013682>, 2010
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J. P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J. K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., and Yool, A.: How well do global ocean biogeochemistry models simulate dissolved iron distributions?, *Global Biogeochem. Cycles*, 30(2), 149–174, <https://doi.org/10.1002/2015GB005289>, 2016.
- 805 Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron in ocean biogeochemistry, *Nature*, 543(7643), 51-59, <https://doi.org/10.1038/nature21058>, 2017.
- Tagliabue, A., Sallée, J. B., Bowie, A. R., Lévy, M., Swart, S., and Boyd, P. W.: Surface-water iron supplies in the Southern Ocean sustained by deep winter mixing, *Nat. Geosci.*, 7(4), 314–320, <https://doi.org/10.1038/ngeo2101>, 2014.

- 810 Taillandier, V., Prieur, L., d'Ortenzio, F., Ribera d'Alcalà, M., and Pulido-Villena, E.: Profiling float observation of thermohaline staircases in the western Mediterranean Sea and impact on nutrient fluxes, *Biogeosciences*, 17(13), 3343–3366, <https://doi.org/10.5194/bg-17-3343-2020>, 2020.
- Ternon, E., Guieu, C., Loÿe-Pilot, M.-D., Leblond, N., Bosc, E., Gasser, B., Miquel, J.-C., and Martín, J.: The impact of Saharan dust on the particulate export in the water column of the North Western Mediterranean Sea, *Biogeosciences*, 7, 809–
815 826, <https://doi.org/10.5194/bg-7-809-2010>, 2010.
- Thieuleux, F., Moulin, C., Bréon, F. M., Maignan, F., Poitou, J., and Tanré, D.: Remote sensing of aerosols over the oceans using MSG/SEVIRI imagery, *Ann. Geophys.*, 23(12), 3561–3568, <https://doi.org/10.5194/angeo-23-3561-2005>, 2005.
- Tovar-Sánchez, A., Rodríguez-Romero, A., Engel, A., Zäncker, B., Fu, F., Marañón, E., Pérez-Lorenzo, M., Bressac, M., Wagener, T., Triquet, S., Siour, G., Desboeufs, K., and Guieu, C.: Characterizing the surface microlayer in the Mediterranean
820 Sea: trace metal concentrations and microbial plankton abundance, *Biogeosciences*, 17(8), 2349–2364, <https://doi.org/10.5194/bg-17-2349-2020>, 2020.
- Twining, B. S., Rauschenberg, S., Morton, P. L., and Vogt, S.: Metal contents of phytoplankton and labile particulate material in the North Atlantic Ocean, *Progr. Oceanogr.*, 137, 261–283, <https://doi.org/10.1016/j.pocean.2015.07.001>, 2015b.
- Twining, B. S., Rauschenberg, S., Morton, P. L., Ohnemus, D. C., and Lam, P. J.: Comparison of particulate trace element
825 concentrations in the North Atlantic Ocean as determined with discrete bottle sampling and in situ pumping, *Deep-Sea Res. Pt II*, 116, 273–282, <https://doi.org/10.1016/j.dsr2.2014.11.005>, 2015a.
- Van Bennekom A. J., Jansen F., Van Der Gaast S., Van Iperen J., and Pieters J.: Aluminum-rich opal: An intermediate in the preservation of biogenic silica in the Zaire (Congo) deep-sea fan, *Deep-Sea Res.*, 36, 173–190, [https://doi.org/10.1016/0198-0149\(89\)90132-5](https://doi.org/10.1016/0198-0149(89)90132-5), 1989.
- 830 Van der Jagt, H., Friese, C., Stuut, J. B. W., Fischer, G., and Iversen, M. H.: The ballasting effect of Saharan dust deposition on aggregate dynamics and carbon export: Aggregation, settling, and scavenging potential of marine snow, *Limnol. Oceanogr.*, 63(3), 1386–1394, <https://doi.org/10.1002/lno.10779>, 2018.
- Van Wambeke, F., Taillandier, V., Deboeufs, K., Pulido-Villena, E., Dinasquet, J., Engel, A., Marañón, E., Ridame, C., and Guieu, C.: Influence of atmospheric deposition on biogeochemical cycles in an oligotrophic ocean system, *Biogeosciences*,
835 18, 5699–5717, <https://doi.org/10.5194/bg-18-5699-2021>, 2021.
- Velasquez, I. B., Ibisani, E., Maas, E. W., Boyd, P. W., Nodder, S., and Sander, S. G.: Ferrioxamine siderophores detected amongst iron binding ligands produced during the remineralization of marine particles, *Frontiers in Marine Science*, 3, 172, <https://doi.org/10.3389/fmars.2016.00172>, 2016.

- Vincent, J., Laurent, B., Losno, R., Bon Nguyen, E., Roullet, P., Sauvage, S., Chevaillier, S., Coddeville, P., Ouboulmane, N.,
840 di Sarra, A. G., Tovar-Sánchez, A., Sferlazzo, D., Massanet, A., Triquet, S., Morales Baquero, R., Fournier, M., Coursier, C.,
Desboeufs, K., Dulac, F., and Bergametti, G.: Variability of mineral dust deposition in the western Mediterranean basin and
south-east of France, *Atmos. Chem. Phys.*, 16, 8749–8766, <https://doi.org/10.5194/acp-16-8749-2016>, 2016.
- Wagener, T., Pulido-Villena, E., and Guieu, C.: Dust iron dissolution in seawater: Results from a one-year time-series in the
Mediterranean Sea, *Geophys. Res. Lett.*, 35(16), L16601, <https://doi.org/10.1029/2008GL034581>, 2008.
- 845 Wagener, T., Guieu, C., and Leblond, N.: Effects of dust deposition on iron cycle in the surface Mediterranean Sea: results
from a mesocosm seeding experiment, *Biogeosciences*, 7, 3769–3781, <https://doi.org/10.5194/bg-7-3769-2010>, 2010.
- Whitby, H., Bressac, M., Sarthou, G., Ellwood, M. J., Guieu, C., and Boyd, P. W.: Contribution of electroactive humic
substances to the iron-binding ligands released during microbial remineralization of sinking particles, *Geophys. Res. Lett.*, 47,
e2019GL086685, <https://doi.org/10.1029/2019GL086685>, 2020.
- 850 Wuttig, K., Wagener, T., Bressac, M., Dammshäuser, A., Streu, P., Guieu, C., and Croot, P. L.: Impacts of dust deposition on
dissolved trace metal concentrations (Mn, Al and Fe) during a mesocosm experiment, *Biogeosciences*, 10, 2583–2600,
<https://doi.org/10.5194/bg-10-2583-2013>, 2013.
- Ye, Y., Wagener, T., Völker, C., Guieu, C., and Wolf-Gladrow, D. A.: Dust deposition: iron source or sink? A case study,
Biogeosciences, 8, 2107–2124, <https://doi.org/10.5194/bg-8-2107-2011>, 2011.