

1 **Supplementary Material**

2

3 **Trace gas fluxes from tidal salt marsh soils: implications for carbon-**
4 **sulfur biogeochemistry**

5

6 Margaret Capooci¹ and Rodrigo Vargas ¹

7 ¹ Department of Plant and Soil Science

8 University of Delaware

9 152 Townsend Hall

10 531 South College Ave.

11 Newark, DE, USA 19716

12

13

14 *Correspondence to:*

15 Rodrigo Vargas

16 152 Townsend Hall

17 531 South College Ave.,

18 Newark, DE 19716

19 rvargas@udel.edu

20 Phone: 302-831-1386 Phone: 302-831-1386

21 Supplementary Table 1 – Overview of CS₂ and DMS fluxes from various *Spartina alterniflora*
 22 marshes.


Location	Time Frame/ Location	CS ₂ (nmol m ⁻² s ⁻¹)	DMS (nmol m ⁻² s ⁻¹)	References
Wallops Island	August/September Annual	0.006 0.68	NA 1.85	Carroll et al., 1986; Adams et al., 1981
North Carolina (Cedar Island Wildlife Refuge, Cox's Landing)	Summer <i>Spartina</i>	0.48	1.75	Aneja et al., 1981, 1979a, 1979b; Lamb et al., 1987; Goldan et al., 1987; Adams et al., 1981b
	Mud Flat	< 0.02	< 0.04	
	Single Plant	0.10	NA	
St. Marks National Wildlife Refuge, FL	January <i>Spartina</i> (W) Sand	< 0.0009 < 0.0009–0.004	0.09–1.63 0.004–0.01	De Mello et al., 1987; Cooper et al., 1987; Adams et al., 1981b
	May <i>Spartina</i> (WD)	< 0.0009–0.003	0.002–4.82	
	October <i>Spartina</i> (WD)	< 0.0009–0.08	0.14–1.27	
	Sand	< 0.0009–0.04	<0.001–0.15	
	Annual	0.52	1.22	
Chapman's Marsh, NH	August	NA	1.1–1.67 Max: 5.3	Morrison and Hines, 1990
Long Island, NY	Fall	NA	Max: 3.80	Hill et al., 1978
Louisiana	Annual	0.002–0.02	0.01–1.25	DeLaune et al., 2002
Canary Marsh, DE	August Intertidal	0.006	0.04	Adams et al., 1981a, 1981b
	Infrequently Flooded	0.06	0.90	
	Unknown	0.07	0.48	
	Annual	0.08	2.84	
Great Sippewissett Marsh, MA	Annual	0.08	2.84	Steudler and Peterson, 1985, 1984

23

24 Notes:

- 25 - For locations with multiple studies the highest mean or widest range is reported here
- 26 - For two locations, the maximum flux was reported in addition to other data (Chapman's
- 27 Marsh and Long Island)


28



29

30 Supplementary Figure 1. Full range of CH₄ fluxes from each chamber during S2. Each color

31 designates a different chamber.




32

33 Supplementary Figure 2. Density plots comparing the distribution of fluxes throughout all
 34 campaigns (continuous) to those measured during daytime low tide (discrete) for (a) CH₄, (b)

35 N₂O, (c) CS₂, and (d) DMS. Note: The scales on the x- and y-axes are different. This figure

36 shows the full distributions of the density plots shown in Figure 4.



37

38 Supplementary Figure 3. Plots comparing the daily average of continuous to discrete

39 measurements for CH₄. Red dashed line is the 1:1 line, while the black solid line is the trend line.

40 This figure shows the full error bars for Figure 5b.

Supplementary References

- 41
42
43 Adams, D. F., Farwell, S. O., Pack, M. R., and Robinson, E.: Biogenic sulfur gas emissions from
44 soils in eastern and southeastern United States, *J. Air Pollut. Control Assoc.*, 31, 1083–
45 1089, <https://doi.org/10.1080/00022470.1981.10465330>, 1981a.
46 Adams, D. F., Farwell, S. O., Robinson, E., Pack, M. R., and Bamesberger, W. L.: Biogenic
47 sulfur source strengths, *Environ. Sci. Technol.*, 15, 1493–1498,
48 <https://doi.org/10.1021/es00094a012>, 1981b.
49 Aneja, V. P., Overton, J. H., Cupitt, L. T., Durham, J. L., and Wilson, W. E.: Carbon disulphide
50 and carbonyl sulphide from biogenic sources and their contributions to the global sulphur
51 cycle, *Nature*, 282, 493–496, <https://doi.org/10.1038/282493a0>, 1979a.
52 Aneja, V. P., Overton, J. H., Cupitt, L. T., Durham, J. L., and Wilson, W. E.: Direct
53 measurements of emission rates of some atmospheric biogenic sulfur compounds, 31,
54 174–178, <https://doi.org/10.1111/j.2153-3490.1979.tb00895.x>, 1979b.
55 Aneja, V. P., Overton, J. H., and Aneja, A. P.: Emission survey of biogenic sulfur flux from
56 terrestrial surfaces, *J. Air Pollut. Control Assoc.*, 31, 256–258,
57 <https://doi.org/10.1080/00022470.1981.10465218>, 1981.
58 Carroll, M. A., Heidt, L. E., Cicerone, R. J., and Prinn, R. G.: OCS, H₂S, and CS₂ fluxes from a
59 salt water marsh, *J. Atmos. Chem.*, 4, 375–395, <https://doi.org/10.1007/BF00053811>,
60 1986.
61 Cooper, D. J., De Mello, W. Z., Cooper, W. J., Zika, R. G., Saltzman, E. S., Prospero, J. M., and
62 Savoie, D. L.: Short-term variability in biogenic sulphur emissions from a Florida
63 *Spartina alterniflora* marsh, *Atmos. Environ.*, 21, 7–12, 1987.
64 DeLaune, R. D., Devai, I., and Lindau, C. W.: Flux of reduced sulfur gases along a salinity
65 gradient in Louisiana coastal marshes, *Estuar. Coast. Shelf Sci.*, 54, 1003–1011,
66 <https://doi.org/10.1006/ecss.2001.0871>, 2002.
67 Goldan, P. D., Kuster, W. C., Albritton, D. L., and Fehsenfeld, F. C.: The measurement of
68 natural sulfur emissions from soils and vegetation: three sites in the Eastern United States
69 revisited, *J. Atmos. Chem.*, 5, 439–467, <https://doi.org/10.1007/BF00113905>, 1987.
70 Hill, F. B., Aneja, V. P., and Felder, R. M.: A technique for measurement of biogenic sulfur
71 emission fluxes, *J. Environ. Sci. Heal. Part A Environ. Sci. Eng.*, 13, 199–225,
72 <https://doi.org/10.1080/10934527809374804>, 1978.
73 Lamb, B., Westberg, H., Allwine, G., Bamesberger, L., and Guenther, A.: Measurement of
74 biogenic sulfur emissions from soils and vegetation: application of dynamic enclosure
75 methods with Natusch filter and GC/FPD analysis, *J. Atmos. Chem.*, 5, 417–437,
76 <https://doi.org/10.1007/BF00113904>, 1987.
77 De Mello, W. Z., Cooper, D. J., Cooper, W. J., Saltzman, E. S., Zika, R. G., Savoie, D. L., and
78 Prospero, J. M.: Spatial and diel variability in the emissions of some biogenic sulfur
79 compounds from a Florida *Spartina alterniflora* coastal zone, *Atmos. Environ.*, 21, 987–
80 990, [https://doi.org/10.1016/0004-6981\(87\)90095-3](https://doi.org/10.1016/0004-6981(87)90095-3), 1987.
81 Morrison, M. C. and Hines, M. E.: The variability of biogenic sulfur flux from a temperate salt
82 marsh on short time and space scales, *Atmos. Environ. Part A. Gen. Top.*, 24A, 1771–
83 1779, [https://doi.org/10.1016/0960-1686\(90\)90509-L](https://doi.org/10.1016/0960-1686(90)90509-L), 1990.
84 Steudler, P. A. and Peterson, B. J.: Contribution of gaseous sulphur from salt marshes to the
85 global sulphur cycle, *Nature*, 311, 455–457, <https://doi.org/10.1038/311455a0>, 1984.
86 Steudler, P. A. and Peterson, B. J.: Annual cycle of gaseous sulfur emissions from a New

87 England *Spartina alterniflora* marsh, Atmos. Environ., 19, 1411–1416,
88 [https://doi.org/10.1016/0004-6981\(85\)90278-1](https://doi.org/10.1016/0004-6981(85)90278-1), 1985.
89