

Nutrient release and flux dynamics of CO₂, CH₄, and N₂O in a coastal peatland driven by actively induced rewetting with brackish water from the Baltic Sea

Daniel Lars Pönisch^{1*}, Anne Breznikar^{2*}, Cordula Nina Gutekunst³, Gerald Jurasinski³, Maren Voss²,

5 Gregor Rehder¹

** these authors contributed equally to this work and share first authorship*

¹ Department of Marine Chemistry, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Rostock, Germany

² Department of Biological Oceanography, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Rostock, Germany

³ Department of Landscape Ecology, Faculty for Agriculture and Environmental Sciences, University of Rostock, Germany

10 Correspondence to: Daniel L. Pönisch (daniel.poenisch@io-warnemuende.de) and Anne Breznikar (anne.breznikar@io-warnemuende.de)

Abstract. The rewetting of drained peatlands supports long-term nutrient removal in addition to reducing emissions of carbon dioxide (CO₂) and nitrous oxide (N₂O). However, rewetting may lead to short-term nutrient leaching into adjacent water and high methane (CH₄) emissions. The consequences of rewetting with brackish water on nutrient and greenhouse gas (GHG) 15 fluxes remain unclear, although beneficial effects such as lower CH₄ emissions seem likely. Therefore, we studied the actively induced rewetting of a coastal peatland with brackish water, by comparing pre- and post-rewetting data from the peatland and the adjacent bay.

Both the potential transport of nutrients into adjacent coastal water and the shift of GHG fluxes (CO₂, CH₄, N₂O) accompanying the change from drained to inundated conditions were analyzed based on measurements of the surface water 20 concentrations of nutrients (dissolved inorganic nitrogen (DIN), phosphate (PO₄³⁻)), oxygen (O₂), components of the CO₂ system, CH₄, and N₂O together with manual closed-chamber measurements of GHG fluxes.

Our results revealed higher nutrient concentrations in the rewetted peatland than in the adjacent bay, indicating that nutrients leached out of the peat and were exported to the bay. A comparison of DIN concentrations of the bay with those of an unaffected reference station showed a significant increase after rewetting. The maximum estimated nutrient export (mean 25 ± 95 % confidence level) out of the peatland was calculated to be 33.8 ± 9.6 t yr⁻¹ for DIN-N and 0.24 ± 0.29 t yr⁻¹ for PO₄-P, depending on the endmember (bay vs. reference station).

The peatland was also a source of GHG in the first year after rewetting. However, the spatial and temporal variability decreased and high CH₄ emissions, as reported for freshwater rewetting, did not occur. CO₂ fluxes (mean ± SD) decreased slightly from 0.29 ± 0.82 g m⁻² h⁻¹ (pre-rewetting) to 0.26 ± 0.29 g m⁻² h⁻¹ (post-rewetting). The availability of organic matter 30 (OM) and dissolved nutrients were likely the most important drivers of continued CO₂ production. Pre-rewetting CH₄ fluxes ranged from 0.13 ± 1.01 mg m⁻² h⁻¹ (drained land site) to 11.4 ± 37.5 mg m⁻² h⁻¹ (ditch). After rewetting, CH₄ fluxes on the

formerly dry land increased by one order of magnitude ($1.74 \pm 7.59 \text{ mg m}^{-2} \text{ h}^{-1}$), whereas fluxes from the former ditch decreased to $8.5 \pm 26.9 \text{ mg m}^{-2} \text{ h}^{-1}$. These comparatively low CH_4 fluxes can likely be attributed to the suppression of methanogenesis and oxidation of CH_4 by the available O_2 and sulfate in the rewetted peatland, which serve as alternative electron acceptors. The post-rewetting N_2O flux was low, with an annual mean of $0.02 \pm 0.07 \text{ mg m}^{-2} \text{ h}^{-1}$.

Our results suggest that rewetted coastal peatlands could account for high, currently unmonitored nutrient inputs into adjacent coastal water, at least on a short time scale such as a few years. However, rewetting with brackish water may decrease GHG emissions and might be favored over freshwater rewetting in order to reduce CH_4 emissions.

1. Introduction

Pristine peatlands are natural sinks for nutrients, in particular nitrate (NO_3^-), and greenhouse gases (GHG) such as mostly carbon dioxide (CO_2) and occasionally nitrous oxide (N_2O) (Martikainen et al., 1993; Regina et al., 1996; Strack, 2008; Kaat and Joosten, 2009). Globally, peatlands store up to 550 Gt of carbon (C), which is twice the C stock of total forest biomass (Moore et al., 1998; Joosten and Clarke, 2002; Kaat and Joosten, 2009).

The drainage of peatlands leads to the mineralization of the topmost peat layer and the accumulation of nutrients (Smolders et al., 2006; Geurts et al., 2010). After rewetting, peatlands can therefore be sources of nutrients, especially ammonium (NH_4^+) and phosphate (PO_4^{3-}) (Lamers et al., 2002; Duhamel et al., 2017). Conversely, due to the anoxic conditions in the water-saturated peat, rewetted peatlands can also act as nutrient sinks, mainly for NO_3^- (Fisher and Acreman, 2004). Whether rewetting leads to nutrient release or uptake is, besides other factors, controlled by the degree of peat decomposition (Zak and Gelbrecht, 2007; Cabezas et al., 2012), the water level (Duhamel et al., 2017) and the salinity (Liu and Lennartz, 2019). Nutrient release is highest in strongly degraded peat in formerly drained peatlands (Zak and Gelbrecht, 2007; Cabezas et al., 2012). Therefore, removal of the topsoil before rewetting has been recommended as a measure to greatly reduce the release of PO_4^{3-} and nitrogen (N) (Harpenslager et al., 2015; Zak et al., 2017). However, nutrient release from peat after rewetting has mostly been assessed in laboratory and incubation studies. To our knowledge, field data on nutrient leaching and potential exports to adjacent waters are lacking.

The GHG exchange of peatlands is strongly influenced by the prevailing biogeochemical and physical conditions, which in turn are largely determined by vegetation and the water level and thus the ratio of oxic and anoxic conditions (Kaat and Joosten, 2009). In drained peatlands, the low water table enables the aerobic decomposition of peat, which is accompanied by increased CO_2 emissions (Joosten and Clarke, 2002). In rewetted peatlands, CO_2 emissions are regulated by photosynthesis, decomposition, and temperature within the upper oxygen-rich soil layer and the overlying water column (Parish, 2008; Oertel et al., 2016). In the anoxic water-saturated zones, the formerly oxygen-induced decomposition of organic matter (OM) is slowed down and relies on alternative terminal electron acceptors (TEAs) such as NO_3^- , manganese (Mn^{4+}), iron (Fe^{3+}), and sulfate (SO_4^{2-}), leading to lowered CO_2 emissions (Strack, 2008; Dean et al., 2018). However, methanogenesis, as the last step in the mineralization of OM and a depletion of TEAs, may become more important in anoxic zones.

Methane (CH₄) emissions in drained peatlands are virtually negligible at water levels < 20 cm below the surface
65 (Jurasinski et al., 2016). Although CH₄ is formed in anoxic zones via methanogenesis, most of it is oxidized as it passes through
the oxic soil layer (Kaat and Joosten, 2009; Dean et al., 2018). Consequently, drained peatlands are a minor source of
atmospheric CH₄. In rewetted peatlands, CH₄ is microbially produced in water-saturated, anoxic soil layers mainly by archaea,
when all other TEAs are depleted (Schönheit et al., 1982; Oremland, 1988; Segers and Kengen, 1998), so that rewetted
peatlands are often significant sources of CH₄ (Hahn et al., 2015). However, in coastal peatlands that receive marine water and
70 therefore SO₄²⁻, the contribution of methanogenesis might be reduced, as methanogenic archaea are outcompeted by sulfate-
reducing bacteria (SRB) (Bartlett et al., 1987; Capone and Kiene, 1988; Oremland, 1988; Jørgensen, 2006). Additionally, any
CH₄ produced may be oxidized by anaerobic methane oxidation coupled to SO₄²⁻ reduction (e.g. Boetius et al., 2000).

N₂O is an intermediate in microbial processes, mostly nitrification, denitrification and nitrifier-denitrification (Kool
et al., 2011). In degraded peatlands, all of these processes are fueled by the accumulated nutrients. Drained peatlands can be
75 weak (Martikainen et al., 1993) or strong sources of N₂O (Liu et al., 2019), depending mainly on the climate zone and land
use (Petersen et al., 2012; Leppelt et al., 2014). Rewetted, and thus water-saturated, peat usually acts as N₂O sink over long-
term scales, due to the formation of anoxic zones where N₂O is consumed (Strack, 2008). However, rewetting can at least
temporarily increase the N₂O production and thus its release into the atmosphere due to the high nutrient availability in strongly
degraded peat, which enables higher rates of nitrification and denitrification (Moseman-Valtierra et al., 2011; Chmura et al.,
80 2016; Roughan et al., 2018).

In temperate latitudes, coastal peatlands are widespread at the interface between marine and terrestrial ecosystems.
However, for many coastal peatlands, the sinking of their ground level due to degradation and peat shrinkage over decades has
made them vulnerable to rising sea level and sinking coasts (Jurasinski et al., 2018). In Mecklenburg-Vorpommern
(northeastern Germany), currently drained coastal peatlands along the low-lying coastline cover an area of ~360–400 km²
85 (Bockholt, 1985; Holz et al., 1996). Nowadays, peatlands are rewetted to restore their habitat function and biodiversity, thereby
preventing CO₂ and N₂O emissions and, in the long-term, reestablishing their C- and N-storage capacity (Strack, 2008;
Zielinski et al., 2018).

Coastal drained peatlands may be rewetted in different ways depending on the available water source. The rewetting
can consist of permanent flooding with freshwater (from groundwater or rivers), episodic inundations with brackish water
90 and permanent brackish water flooding. While the effects of freshwater rewetting (Richert et al., 2000; Hogan et al., 2004; Zak
and Gelbrecht, 2007) and episodic inundations with brackish water on nutrient dynamics and GHG have been investigated
(Chmura et al., 2011; Neubauer et al., 2013; Hahn et al., 2015; Koebisch et al., 2019), less is known about the impact of
permanent brackish water flooding.

In this study we examined the immediate effects of rewetting with brackish water on nutrient (NO₃⁻, nitrite (NO₂⁻),
95 NH₄⁺ and PO₄³⁻) and GHG fluxes (CO₂, CH₄, N₂O) in a low-lying, highly degraded coastal peatland at the German Baltic Sea
coast, by comparing pre- and post-rewetting conditions. Due to the unique formation of a permanent brackish water column
above formerly drained peat, this is the first study to combine marine shallow-water and terrestrial peatland research. We

investigated how the rewetting with brackish water affects (1) nutrient leaching and the potential transport from a nutrient-enriched, flooded peatland to the adjacent bay driven by frequent water exchange, (2) the GHG dynamics in the surface water within the first year after rewetting and (3) the GHG fluxes along the transition from drained to inundated conditions.

2. Material and methods

2.1 Study area

The study area is a low-lying, highly degraded coastal peatland that had been transformed from a drained, agriculturally used polder to a brackish wetland. The “Polder Drammendorf” (referred to in the following as “peatland”) is located at the northeastern German Baltic Sea coast, on the western part of the island of Rügen (Mecklenburg-Vorpommern, Germany), bordering on the Kubitzer Bodden (Figure 1). The climate is oceanic, with a mean annual air temperature of 9.1 °C and a mean annual precipitation height of 626 mm (Deutscher Wetterdienst (DWD), 1991–2020). The central Kubitzer Bodden has a mean surface water temperature of 11.4 ± 6.6 °C and a mean surface salinity of 8.5 ± 1.4 (referred to in the following as “central bay”, data retrieved from a monitoring station of the Landesamt für Umwelt, Naturschutz und Geologie Mecklenburg Vorpommern (LUNG MV), 2006–2020, 54.40° N, 13.11° E, Figure 1b). For comparison, the Arkona Basin, the near-by open Baltic Sea basin to the north of the island of Rügen that influences the water in the Kubitzer Bodden, has a mean surface water temperature of 10.2 ± 5.6 °C and a mean surface salinity of 8.0 ± 0.5 (MARNET, data originator: Leibniz Institute for Baltic Sea Research Warnemünde, Germany, 2006–2020, 54.88° N, 13.86° E).





Figure 1. (a) Overview of the study area located in the southern Baltic Sea. (b) Coastline of northeast Germany in Mecklenburg-Vorpommern and study area location (“Polder Drammendorf”, red) on the island of Rügen, bordering on the Kubitzer Bodden, where a monitoring station served as reference (“central bay”, purple). The Kubitzer Bodden is connected with the Arkona Basin in the north. Bathymetry refers to Seifert et al., (2001), and borders were retrieved from National Oceanic And Atmospheric Administration (NOAA) - National Centers For Environmental Information (NCEI).

Like most peatlands in northern Germany, Drammendorf was artificially drained for agricultural use (pasture and grassland) in the 1960s, by establishing a sandy dike and an extensive ditch system that affected an area of ~ 2.2 km². The northwestern

part (mostly mineral soil, higher elevation) served as grassland while the northeastern part was used for agriculture with seasonal fertilizer application only until the 1990s ($100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). The southern compartment (organic soil) provided an area for cattle grazing (~ 30 cows). The topsoil of the central part consists of up to 50–70 cm highly degraded peat (Brisch, 2015), classified as H7 according to the von Post humification scale (Wang et al., 2021). This highly degraded topsoil layer was not removed prior to rewetting. Underneath the degraded topsoil follows a well-preserved peat layer with a thickness of ~ 100 cm. Peat deposits of up to 220 cm thickness are largest in the western part, near the former dike. The long-lasting drainage and ongoing peat degradation have led to the formation of a local land depression with an average soil elevation of around -0.5 meters above sea level (masl). To control the water expansion after rewetting, a new dike was built in the southern part before flooding (Figure 2a). Additionally, a drainage ditch that receives water from the catchment was rebuilt and a new pumping station was installed. A significant input of nutrients from this additional water supply can be excluded due to the low pumping activity and the absence of a permanent hydrological connection to the study area (Wasser- und Bodenverband Rügen (WBV), pers. comm., 2020).



135 **Figure 2.** Topography of the study area and overview of the stations in the inner bay (purple), the flooded peatland (black)
and along the transect of the GHG flux measurements (red). (a) Water coverage at mean sea level. The new dike is shown in
dark red. (b) Transect stations that were sampled for atmospheric chamber-based GHG flux measurements (before and after
rewetting) and for surface water GHG concentration measurements (after rewetting). Data from station BTD7 were used for a
comparison of the chamber-based measurements with the calculated air-sea fluxes after rewetting. Topography data retrieved
140 from the Landesamt für innere Verwaltung MV, Amt für Geoinformation, Vermessungs- und Katasterwesen, Fachbereich
Geodatenbereitstellung.

The area was rewetted by the targeted removal of a 20 m wide dike section in November 2019 that caused an immediate
flooding of the low-lying area behind the dike. The newly built channel represents the only permanent hydrological connection
145 between the peatland and the Kubitzer Bodden that allows major surface water exchange. The remaining section of the dike
(~650 m) was removed down to the surface elevation level and is hence only flooded at very high water levels.

The restored area covers ~0.8 km² in total and is characterized by a permanently water-covered area of ~0.5 km², with
a mean water depth of ~0.5 m, compared to 1.0–1.5 m in the Kubitzer Bodden. The extent of the inundated area depends
directly on the water level of the Baltic Sea, which is highly dynamic despite the absence of regular tides (Figure A1).
150 Therefore, minor changes in the water level lead to major changes in the water-covered area. For instance, if the water level
rises from –0.5 to + 0.5 masl, the water-covered area increases from 0.08 to 0.7 km² (Figure A2, Figure A3). The ditch system
was only partly removed and hence, some deeper areas with water depths of up to 4 m remained. It is noteworthy that in the
first months after rewetting, former grassland and ditch vegetation (*Elymus repens* L. (Gould) (Couch grass), *Phragmites*
australis (Cav.) Trin. ex Steud. (Common reed)) died almost completely and the cover of emergent macrophytes was then
155 negligible. However, *Phragmites* was able to grow back during the growing season and expanded especially around the ditches.

2.2 Sampling

2.2.1 Surface water sampling

Before rewetting, surface water samples for nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻) and chlorophyll a were collected from the
inner Kubitzer Bodden (referred to in the following as “inner bay”) at station D1 (Figure 2a) and irregularly at a second station
160 right in front of the now removed dike section, which was abandoned after rewetting and therefore merged with station D1.
Environmental variables (water temperature, dissolved oxygen (O₂), salinity) were measured on-site. Both stations were
reached from land and sampling was conducted monthly from June to November 2019, except in August.

After rewetting, surface water samples were collected with a small boat and the sampled variables were extended for
the concentrations of GHGs (CO₂, CH₄, N₂O) and dissolved organic carbon (DOC). The first sampling took place one week
165 after the dike removal. Sampling was continued over one year (25 sampling dates until December 2020) at weekly (December
2019 to January 2020) or biweekly (February 2020 to September 2020, except for August) intervals. From October 2020 to
December 2020, sampling was conducted monthly. In the inner bay, three stations (D1, D3, D14), and in the flooded peatland
six stations (D4, D5, D11, D12, D15, BTD8) were sampled (Figure 2a). The inner bay station “D14” was sampled from March

2020 onwards. DOC sampling started in April 2020. For the air-sea gas exchange calculation, data from station D10_1, located
170 in the channel, were also included.

Moreover, surface water samples for the analysis of GHG concentrations (CO_2 , CH_4 , N_2O) were sampled at eight stations along a transect (Figure 2b). This sampling was carried out simultaneously with the sampling described in Sect. 2.2.2 to link GHG air-sea exchange calculations based on surface water samples with chamber-based flux measurements.

175 Surface water temperature, O_2 , and salinity were measured directly in the field using a HACH HQ40D multimeter (HACH Lange GmbH, Germany) equipped with two outdoor electrodes (LDO10105, CDC40105). Depending on the prevailing water depth, additional measurements were conducted in the peatland 15 cm above the soil surface (excluding the ditches) on 22 of the 25 sampling dates. The precision of the electrodes was ± 0.3 °C, ± 0.8 %, and ± 0.1 for temperature, O_2 saturation, and salinity, respectively.

180 Surface water samples were taken using a horizontal 7 L Niskin bottle to sample the upper 20 cm of the water column. These included 250 mL subsamples for $\text{CH}_4/\text{N}_2\text{O}$ analysis (bottles capped with butyl rubber stoppers and crimp-sealed), analysis of the CO_2 system (one bottle each for total CO_2 (C_T), total alkalinity (A_T), and pH) and 15 mL subsamples for the analysis of nutrients and DOC. Water for chlorophyll a determination was taken using 3 L canisters.

185 In the laboratory, $\text{CH}_4/\text{N}_2\text{O}$ and CO_2 samples were poisoned with 500 μL and 200 μL of saturated HgCl_2 , respectively, and stored in the dark at 4°C until analysis. Subsamples for nutrients and DOC were filtered in the field with pre-combusted (450 °C for 4 h) 0.7 μm glass-fiber filters (GF/F, Whatman®) and stored at -20 °C. Samples for chlorophyll a were filtered in the laboratory with non-combusted 0.7 μm glass-fiber filters (GF/F, Whatman®) and likewise stored at -20 °C.

2.2.2 Chamber-based atmospheric GHG flux sampling for CO_2 and CH_4

Starting in June 2019, nearly 6 months before rewetting, GHG exchange was regularly measured using dynamic closed chambers (Livingston and Hutchinson, 1995) along a transect representing a soil humidity gradient (Figure 2b). The
190 measurements were conducted twice a month, for a total of eleven sampling days at six peatland stations and two additional stations in the north-south-oriented main ditch. Each station was sampled up to eight times per sampling day, resulting in overall 418 CO_2 and 184 CH_4 pre-rewetting flux measurements.

195 For each measurement, the chambers were placed on permanently installed collars and connected through an air-tight seal, with a closure period between 180 and 300 s. Between the measurements, chambers were lifted to vent them until atmospheric GHG concentrations were reached. To ensure coverage of photosynthetic and respiration activity, CO_2 measurements on the terrestrial peatland were conducted using opaque and transparent chambers for NEE (net ecosystem exchange) and Reco (ecosystem respiration) determination, respectively. To cover a broad spectrum of solar radiation, two additional measurements were conducted with cloth-covered transparent chambers, resulting in a reduced photosynthetically active photon flux density (PPFD). Ditch stations' GHG concentrations were determined in three consecutive measurements
200 with floating opaque chambers placed on the water surface. Changes in GHG concentrations in the chamber headspace were measured using a portable laser-based analyzer (Picarro G4301, GasScouter, Santa Clara, USA; LI-820, LI-COR Biosciences,

Lincoln, USA and an Ultraportable Greenhouse Gas Analyzer (UGGA), Los Gatos Research Inc., Mountain View, Calif., USA).

205 After rewetting, the stations along the transect covered a gradient of ground elevations, including stations that fell dry at low water levels and stations that remained permanently flooded. Atmospheric GHG fluxes were measured twice a month using floating opaque chambers positioned above the same sampling locations of the flooded peatland. Since the flooding caused most plants to die, and almost all measurement locations were covered by water during the study period, we reduced the number of NEE measurements with transparent chambers to stations and days with low water table. Approximately six measurements per station were made during 23 sampling days between December 2019 and December 2020, with a total of 210 698 CO₂ and 482 CH₄ fluxes determined during the post-rewetting year.

2.3 Data processing, statistics, and definition of seasons and means

Data analysis and visualization were performed using R (R Core Team, 2020) and the packages *tidyverse* (Wickham et al., 2019), *lubridate* (Grolemund and Wickham, 2011), *patchwork* (Pedersen, 2020) *car* (Fox and Weisberg, 2019), and *flux* (Jurasinski et al., 2014). The relationships between environmental variables, nutrient concentrations, and GHG 215 concentrations/fluxes were investigated in linear regression analyses. The significance level was set to $p < 0.05$.

To describe temporal patterns during the entire sampling period, we defined two pre- and four post-rewetting periods, roughly corresponding to seasons (Table 1). For a direct comparison between the pre- and post-rewetting periods, we compared nutrient and GHG flux data from summer and autumn 2019 with those from summer and autumn 2020 (Table 3) by using the Mann-Whitney-U test.

220 **Table 1.** Defined seasons of the investigation period

	Pre-rewetting		Post-rewetting			
season	summer 2019	autumn 2019	winter 2019/2020	spring 2020	summer 2020	autumn 2020
months	June–August	September–November	December–February	March–May	June–August	September–December

We analyzed the data among the respective peatland and inner bay stations in order to verify the use of means for each sampling site (peatland and inner bay separately) and date. The difference between spatial (sampling stations of either sites) and temporal (sampling seasons) data variability was tested by using a Two-Way ANOVA and showed a higher temporal variability 225 ($p < 0.05$). Therefore, we decided to combine the stations within the peatland and within the inner bay, respectively, to report mean values and standard deviations (single values can be found in the published data set). The Two-Way ANOVA was also used to identify seasonal differences between the peatland and the inner bay (Table 2).

At station D3, in the inner bay, the pH, CH₄, and pCO₂ values differed significantly from those of the remaining stations of the inner bay during the year after rewetting (ANOVA, Kruskal-Wallis test). Since the differences in water 230 temperature, salinity, and O₂ were not significant, we decided to include the data from D3 for these variables to obtain a larger data pool for the inner bay and to exclude D3 for all other variables. The exclusion was conducted because variables such as

pH, CH₄, and pCO₂ are related to biological activity which can vary, while the more physically influenced variables (temperature, salinity, and O₂) are rather constant.

2.4 Nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻), chlorophyll a and DOC

235 2.4.1 Analysis

Nutrient analyses were carried out according to standard photometric methods (Grasshoff et al., 2009) by using a continuous segmented flow analyzer (SEAL Analytical QuAAtro, SEAL Analytical GmbH, Norderstedt, Germany). Detection limits were 0.2 μM for NO₃⁻, 0.05 μM for NO₂⁻, 0.5 μM for NH₄⁺ and 0.1 μM for PO₄³⁻. Measurements of the nutrient concentrations were partly below the detection limit for the peatland, the inner bay and the central bay (flagged in the published dataset). For 240 such measurements below detection limit, it is recommended to use the actual values of these measurements (e.g. Fiedler et al., 2022) to achieve a robust statistical analysis. Since these data were not available, we decided to use randomly generated values between 0 and the respective detection limit with a uniform distribution for these measurements.

Chlorophyll a was extracted from glass-fiber filters (GF/F, Whatman®) by incubation with 96 % ethanol for 3 h and afterwards analyzed by using a fluorometer (TURNER 10-AU-005, Turner Designs, San José, USA) at 670 nm according to 245 Edler (1979). DOC was analyzed after high-temperature combustion using a Multi 2100S instrument (Analytik Jena GmbH, Jena, Germany) and detected by non-dispersive infrared spectrometry according to ISO 20236, ISO 8245 I, and EN 1484.

2.4.2 Use of reference data from a monitoring station

Coastal nutrient data (NO₃⁻, NO₂⁻, NH₄⁺ and PO₄³⁻ concentrations) from a monitoring station in the Kubitzer Bodden (“central bay”, Figure 1b) ~15 km away from the study area were obtained as reference. Monitoring data from 2016 to 2020 were 250 included. These data were used (1) to compare them with nutrient concentrations from the inner bay before and after rewetting to detect potentially higher concentrations, resulting from nutrient leaching within the peatland and a subsequent export into the inner bay and (2) to calculate the total possible export out of the peatland (Sect. 2.4.3) by using the monitoring station as a second, unaffected endmember besides the inner bay, which is by contrast potentially affected by the rewetting. Due to transformations and potential losses along the way to the monitoring station, especially of the nitrogen species, the calculated 255 total possible exports have to be considered as maximum estimates.

2.4.3 Nutrient transport calculation (DIN-N and PO₄-P)

To calculate the bulk exchanges of dissolved inorganic nitrogen (DIN-N) and PO₄-P between the flooded peatland and the inner bay/central bay, the water level was transformed to water volume by creating a hypsographic curve with increments of 0.1 m and a resolution of 1x1 m (Figure A3). Water level data from a nearby monitoring station (“Barhöft”, 54.43° N, 13.03° E) 260 and topography data with a resolution of 1x1 m were obtained from the Wasserstraßen- und Schifffahrtsamt Ostsee (WSA) and the Landesamt für innere Verwaltung MV, respectively. To ensure that the water level data of the monitoring station were

valid for the peatland, the water level data of the latter, measured between August and December 2020, were compared with the data from the monitoring station, which showed a strong correlation ($r_s = 0.95$, $p < 0.001$, 15-min intervals, data not shown).

265 A water level of -1.6 masl, as the lowest recorded water level within the last 25 years, was used as the starting point to derive the cumulative water volumes of the peatland. The water volumes were then assigned to the corresponding water levels to finally calculate the water volume changes (Q , in $\text{m}^3 \text{s}^{-1}$) according to Eq. (1):

$$Q(t) = \frac{dV}{dt} \quad (1)$$

where V is the water volume and t the time. Positive volume changes ($Q > 0$) indicate an inflow of water into the peatland and *vice versa*. For each season, the mean inflow (Q_{in}) and outflow (Q_{out}) volumes were calculated according to Eqs. (2) and (3):

$$Q_{in} = \frac{1}{\Delta T} \int_t^{t+\Delta T} Q^{positive} dt \quad \text{for } Q > 0 \quad (2)$$

$$Q_{out} = \frac{1}{\Delta T} \int_t^{t+\Delta T} Q^{negative} dt \quad \text{for } Q < 0 \quad (3)$$

270 where ΔT denotes the season length. Note that Q_{out} is negative. Seasonal mean values of nutrient concentrations (DIN and PO_4^{3-}) were calculated and converted from μM to kg m^{-3} by using the molecular masses of the basic elements N and P to derive DIN-N and $\text{PO}_4\text{-P}$. After the conversion, nutrient masses of the peatland ($c_{peatland}$) and the inner bay (c_{IB}) vs. peatland and central bay (c_{CB}), respectively, were multiplied by Q_{out} and Q_{in} and integrated to calculate the net nutrient transport (NNT, in tonnes, equivalent to megagrams) according to Eqs. (4) and (5):

$$NNT = \int_t^{t+\Delta T} Q_{in} c_{IB} dt + \int_t^{t+\Delta T} Q_{out} c_{peatland} dt \quad (4)$$

$$NNT = \int_t^{t+\Delta T} Q_{in} c_{CB} dt + \int_t^{t+\Delta T} Q_{out} c_{peatland} dt \quad (5)$$

275 Negative values indicate a net nutrient export from the peatland into the inner/central bay, and positive values display a net nutrient import into the peatland. Uncertainty ranges for the seasonal NNTs (u_{NNT} , as 95 % confidence level) were calculated by using an error propagation according to Eq. (6):

$$u_{NNT} = \sqrt{(c_{bay} dt u_{Q_{in}})^2 + (c_{peat} dt u_{Q_{out}})^2 + (Q_{out} dt u_{c_{peat}})^2 + (Q_{in} dt u_{c_{bay}})^2} \quad (6)$$

where terms with “ u ” denote the respective 95 % confidence level. To derive the annual uncertainty range of the NNT, all seasonal errors were added up.

280 2.5 GHG concentrations and fluxes

2.5.1 Inorganic carbon system analysis

Directly measured variables (C_T , A_T , pH)

The inorganic carbon system was determined by analyzing the total CO₂ (C_T), total alkalinity (A_T), and pH of the water samples. C_T was measured with an automated infrared inorganic carbon analyzer (AIRICA, S/N #027, Marianda, Kiel, Germany). The system acidifies a discrete sample volume (phosphoric acid, 10 %), whereby the inorganic carbon species of C_T are shifted to CO_{2(g)}. A carrier gas stream (99.999 % N₂) transfers the gaseous components to a Peltier device and a Nafion® drying tube (Perma Pure Nafion®, Ansyco GmbH, Karlsruhe, Germany) to remove water residues. The produced CO_{2(g)} is detected by an infrared detector (LICOR 7000; LI-COR Environmental GmbH, Bad Homburg, Germany). Certified reference materials (CRM; Scripps Institution of Oceanography, University of California, San Diego, USA) were used for calibration. Triplicate measurements were conducted for each sample, and a precision of ± 5 μmol kg⁻¹ was achieved.

A_T was measured by potentiometric titration (glass electrode type LL Electrode plus 6.0262.100, Metrohm, Filderstadt, Germany) in the open-cell configuration, after Dickson et al. (2007). The system was calibrated with the same CRM as used for C_T and resulted in the same precision.

The pH was analyzed spectrophotometrically using the pH-sensitive indicator dye m-cresol purple (mCP, 2 mM, Contros System and Solution GmbH, Kiel, Germany). The measurement principle and instrumental setup are described elsewhere (Dickson et al., 2007; Carter et al., 2013). In brief, absorption was measured using the Agilent 8453 UV-visible spectroscopy system (Agilent Technology, Waldbronn, Germany); pH parameterization for brackish water was calculated following Müller and Rehder (2018). Quality control was performed by measuring buffer solutions (salinity of 20) prepared according to Müller et al. (2018). An external buffer solution with a salinity of 35 (Scripps Institution of Oceanography, University of California, San Diego, USA) was additionally used. All pH values are reported given on the total scale (pH_T).

Calculated variables

The CO₂ partial pressure in the water phase (pCO₂), the value of which was required for the CO₂ air-water flux calculations (Sect. 2.5.3), was calculated from C_T and pH using the R packages *seacarb* (Gattuso et al., 2019), with K₁ and K₂ from Millero (2010), K_s from Dickson (1990), and K_f from Dickson and Riley (1979). C_T and pH were preferred because non-oceanic components, in particularly organic acid-base systems, contribute significantly to A_T (Kuliński et al., 2014). A_T was also calculated from C_T and pH and the values compared with measured values, thus revealing the magnitude of the contributions of those components to A_T.

2.5.2 Dissolved CH₄ and N₂O concentration analysis

Dissolved CH₄ and N₂O concentrations were determined by gas chromatography on an Agilent 7890B instrument (Agilent Technologies, Santa Clara, USA) coupled to a flame ionization detector (FID) and an electron capture detector (ECD). A purge and trap technique, explained in detail in Sabbaghzadeh et al. (2021) was used. In brief, a helium gas stream was used to purge 10 mL of seawater to extract volatile compounds. The gas stream passed through a purifier (VICI Valco Instruments Co. Inc., Houston, USA) and was dried using a Nafion® tube (Perma Pure Nafion®, Ansyco GmbH, Karlsruhe, Germany) and a SICAPENT® tube (Merck KGaA, Darmstadt, Germany). The relevant compounds were enriched by cryofocusing on a trap filled with HayeSep D® (CS Chromatographie Service GmbH, Langerwehe, Germany) maintained at -120 °C using an

ethanol/nitrogen cooling bath. After 10 minutes of heating in a 95 °C water bath, the compounds were desorbed and separated by two capillary columns linked to the detectors by a Deans Switch (Pönisch, 2018).

For quality control, a calibration standard (gas composition: 9.9379 ± 0.0159 ppm CH₄ and 1982.07 ± 3.77 ppb N₂O) was measured daily before and after the sample measurements; the standard deviation was < 1 %. The calibration range was adjusted using multi-loop injection of the calibration gas to ensure that the samples were within the limits of the calibration. The standard was recalibrated according to high-precision standards (ICOS-CAL laboratory, Max Planck Institute, Jena, Germany).

2.5.3 GHG flux calculations

Atmospheric fluxes based on closed-chamber measurements

CO₂ and CH₄ fluxes were calculated using the ideal gas law (Livingston and Hutchinson, 1995), as formulated in Eq. (7):

$$F = \frac{MpV}{RTA} * \frac{dc}{dt} \quad (7)$$

where F is the GHG flux ($\text{g m}^{-2} \text{h}^{-1}$), M is the molar mass of the gas (g mol^{-1}), p is the standard air pressure (101,300 Pa), V is the chamber volume (m^3), R is the gas constant ($\text{m}^3 \text{Pa K}^{-1} \text{mol}^{-1}$), T is the temperature in the chamber (K), A is the surface area of the measurement collar (m^2), and dc/dt is the change in concentration over time. The latter was derived from the slope of a linear regression based on the medians of the gas concentrations. The atmospheric sign convention was applied; thus, positive fluxes indicated a release of GHG by the soil and negative fluxes GHG uptake by the soil. The fluxes were estimated using the function *fluxx()* of the R package *flux* (Jurasinski et al., 2014) and the SLP method. Outlier identification (using a histogram) resulted in the exclusion of CO₂ fluxes which were smaller than -2.5 and larger than $4 \text{ g m}^{-2} \text{h}^{-1}$. Likewise, CH₄ fluxes larger than $200 \text{ mg m}^{-2} \text{h}^{-1}$ were discarded due to a high risk of capturing ebullition-based CH₄ emissions instead of diffusive fluxes.

335

Atmospheric fluxes based on air-sea gas exchange parameterization (velocity k model)

The air-sea gas exchange (F , $\text{g m}^{-2} \text{h}^{-1}$) is a function of the gas transfer velocity (k) and the concentration difference between the bulk liquid (C_w) and the top of the liquid boundary layer adjacent to the atmosphere (C_a). It was calculated as reported in Wanninkhof (2014) and as shown in Eq. (8):

$$F = k (C_w - C_a) \quad (8)$$

where k was derived from an empirical relationship between a coefficient of gas transfer (0.251) and the wind speed $\langle U^2 \rangle$ (Wanninkhof, 2014) and Schmidt number (Sc), as expressed by Eq. (9):

$$k = 0.251 \langle U^2 \rangle (Sc/660)^{(-0.5)} \quad (9)$$

Wind speeds originated from the nearby (~15 km away) monitoring station Putbus and were measured at 10 m height (DWD; 54.3643° N, 13.4771° E, WMO-ID 10093). The average wind speed was defined in this study ± 3 h from midday, because the wind speed over 24 h was lowest at night and highest at midday and because sampling was usually conducted within the

345 selected time interval. The Schmidt number was approximated by a linear interpolation between the freshwater and seawater values. Atmospheric-equilibrium conditions (C_a) were calculated using the atmospheric data for CO_2 and CH_4 obtained from the ICOS station “Utö” (Finnish Meteorological Institute, Helsinki). Due to the seasonal changes in the atmospheric dry molar fraction of CO_2 and CH_4 , mean values for each season were computed. For N_2O , the atmospheric dry mole fraction from station Mace Head was selected (National University of Ireland, Galway; data from the NOAA GML carbon cycle cooperative global
350 air sampling network (Dlugokencky et al., 2019a, 2019b)). A mean value of the atmospheric N_2O concentration during the investigation period was calculated due to its minor seasonality. Equilibrium concentrations were then calculated using the solubility coefficient (K_0) from Weiss and Price (1980). We acknowledge that the air-sea exchange model we used (Wanninkhof, 2014) was developed for open ocean waters and is a doubtful approach for deriving fluxes in small enclosed areas such as our working area. However, the lack of an appropriate parameterization, and the convincing result of the
355 comparison of our two approaches (see below and Appendix C: Comparability of two independent approaches to atmospheric flux determination) justify our approach.

Comparability of two independent approaches to atmospheric flux determination

We evaluated the comparability of the two previously described methods by comparing the results of a representative station (BTD7) for each post-rewetting season. The comparison showed no significant differences between the fluxes of CO_2 and CH_4
360 derived with the different methods and therefore, it seems appropriate to combine the fluxes for each GHG into one pooled post-rewetting data set. The pooled post-rewetting flux values were compared with the pre-rewetting values to investigate the effect of rewetting on CH_4 and CO_2 fluxes (Table 3). For more details concerning the comparability assessment, see Appendix C: Comparability of two independent approaches to atmospheric flux determination. Due to the large variability and the pooling of chamber-based measurements with k model data, the GHG fluxes after rewetting are hardly suitable for upscaling
365 and thus, the single values in the published data should be used.

3. Results

3.1 Surface water properties (temperature, salinity, O_2 , DOC, chlorophyll a)

In the first year after rewetting, there were no significant differences between the peatland and the inner bay with respect to surface water temperature, salinity and O_2 saturation (Figure 3a–c, Table 2), suggesting a pronounced water exchange between
370 the peatland and the inner bay that was driven by frequent changes in the water level (Figure A1). Additionally, no significant differences between summer and autumn 2019 and summer and autumn 2020 were found in the inner bay.

Temperature and salinity measurements near the peat surface showed no significant differences between the surface and bottom water over the year ($n_{\text{surface}} = 140$, $n_{\text{bottom}} = 86$, data not shown), which suggested that vertical exchange processes and mixing were highly pronounced. However, a significant difference in O_2 saturation between the surface and bottom water
375 in summer ($p < 0.01$) indicated that local and temporary gradients are possible.




Figure 3. Time series of the mean (a) temperature, (b) salinity, (c) O₂ saturation, (d) DOC concentration and (e) chlorophyll a concentration (\pm standard deviations) in the surface water from June 2019 to December 2020. Data from the flooded peatland ($n = 6$) are shown in blue and data from the inner bay ($n = 2$ or 3 , as explained in Sect. 2.3) in black. The vertical black line indicates the rewetting event.

380 DOC concentrations were significantly higher in the peatland than in the inner bay in spring and summer, with the highest concentration ($\sim 30 \text{ mg L}^{-1}$) measured in the peatland (Figure 3d, Table 2). Chlorophyll a concentrations after rewetting showed clear seasonal and spatial differences, with significantly higher concentrations in the peatland in spring and summer (max. $\sim 125 \text{ } \mu\text{g L}^{-1}$, Figure 3e, Table 2). A comparison of pre- and post-rewetting chlorophyll a concentrations in the inner bay in summer and autumn showed higher concentrations after rewetting (pre-rewetting: $2.5 \pm 0.9 \text{ } \mu\text{g L}^{-1}$, post-rewetting: 385 $15.4 \pm 11.5 \text{ } \mu\text{g L}^{-1}$).

Table 2. Seasonal comparison of the surface water means (\pm standard deviation) in the peatland (“peat”) as opposed to the inner bay (“bay”) for all in situ variables. The number of observations is shown in parentheses, and significant seasonal differences ($p < 0.05$) between the inner bay and the peatland are indicated in bold.

		Pre-rewetting		Post-rewetting			
		summer 2019	autumn 2019	winter 2019	spring 2020	summer 2020	autumn 2020
temperature (°C)	peat	N/A	N/A	3.73 \pm 1.25 (45)	12.03 \pm 4.17 (35)	19.85 \pm 2.44 (30)	12.94 \pm 6.61 (30)
	bay	25.17 \pm 3.27 (3)	13.95 \pm 3.59 (6)	3.86 \pm 0.99 (17)	12.17 \pm 4.09 (17)	19.36 \pm 2.68 (15)	12.52 \pm 6.58 (15)
salinity	peat	N/A	N/A	6.67 \pm 0.68 (45)	8.23 \pm 0.66 (35)	8.96 \pm 0.50 (30)	8.22 \pm 0.33 (30)
	bay	9.21 \pm 0.69 (4)	8.39 \pm 0.38 (6)	6.99 \pm 0.65 (17)	8.27 \pm 0.56 (17)	8.86 \pm 0.63 (15)	8.13 \pm 0.32 (15)
O ₂ (mg L ⁻¹)	peat	N/A	N/A	11.19 \pm 0.74 (45)	11.72 \pm 1.93 (35)	8.60 \pm 1.86 (30)	9.34 \pm 1.35 (30)
	bay	7.66 \pm 1.70 (3)	7.48 \pm 3.87 (6)	11.18 \pm 0.67 (17)	10.03 \pm 3.48 (17)	8.26 \pm 2.26 (15)	8.86 \pm 1.80 (15)
chlorophyll a (μ g L ⁻¹)	peat	N/A	N/A	8.55 \pm 10.80 (24)	40.03 \pm 26.39 (12)	74.03 \pm 29.01 (10)	30.57 \pm 37.50 (10)
	bay	2.66 \pm N/A (1)	2.42 \pm 1.09 (3)	4.76 \pm 2.31 (8)	13.52 \pm 8.90 (8)	21.91 \pm 11.04 (10)	8.83 \pm 7.76 (10)
DOC (mg L ⁻¹)	peat	N/A	N/A	N/A	14.82 \pm 2.13 (18)	16.95 \pm 6.09 (27)	12.07 \pm 3.47 (29)
	bay	N/A	N/A	N/A	11.78 \pm 2.12 (6)	10.72 \pm 2.73 (10)	11.09 \pm 2.54 (10)
NO ₃ ⁻ (μ M)	peat	N/A	N/A	100.03 \pm 57.66 (45)	25.22 \pm 46.03 (35)	0.14 \pm 0.10 (29)	3.69 \pm 3.99 (30)
	bay	0.36 \pm 0.30 (4)	2.33 \pm 2.80 (6)	68.50 \pm 40.67 (9)	15.38 \pm 30.68 (11)	0.16 \pm 0.12 (10)	3.38 \pm 3.56 (10)
NO ₂ ⁻ (μ M)	peat	N/A	N/A	1.49 \pm 0.62 (45)	0.43 \pm 0.44 (35)	0.23 \pm 0.12 (29)	0.99 \pm 1.03 (30)
	bay	0.11 \pm 0.07 (4)	0.19 \pm 0.11 (6)	1.04 \pm 0.49 (9)	0.29 \pm 0.33 (11)	0.16 \pm 0.12 (10)	1.11 \pm 1.20 (10)
NH ₄ ⁺ (μ M)	peat	N/A	N/A	30.02 \pm 26.13 (45)	2.27 \pm 1.56 (35)	5.54 \pm 6.48 (29)	18.78 \pm 19.50 (30)
	bay	1.67 \pm 1.33 (3)	3.00 \pm 1.70 (6)	21.47 \pm 23.42 (9)	1.71 \pm 1.13 (11)	2.82 \pm 3.87 (10)	17.03 \pm 21.78 (10)
PO ₄ ³⁻ (μ M)	peat	N/A	N/A	0.37 \pm 0.41 (45)	0.26 \pm 0.28 (35)	0.49 \pm 0.26 (29)	0.35 \pm 0.33 (30)
	bay	1.30 \pm 1.90 (4)	0.12 \pm 0.08 (6)	0.21 \pm 0.21 (9)	0.09 \pm 0.13 (11)	0.22 \pm 0.21 (10)	0.26 \pm 0.28 (10)
CH ₄ (nmol L ⁻¹)	peat	N/A	N/A	47.96 \pm 49.52 (46)	300.49 \pm 414.29 (35)	1502.36 \pm 693.36 (30)	733.74 \pm 699.17 (30)
	bay	N/A	N/A	81.37 \pm 106.93 (7)	130.12 \pm 190.54 (11)	502.47 \pm 479.31 (10)	194.70 \pm 186.49 (20)
N ₂ O (nmol L ⁻¹)	peat	N/A	N/A	85.53 \pm 152.45 (46)	15.42 \pm 4.97 (35)	6.95 \pm 1.35 (30)	14.34 \pm 4.04 (30)
	bay	N/A	N/A	26.74 \pm 9.69 (7)	13.13 \pm 4.13 (11)	8.76 \pm 1.26 (10)	16.68 \pm 5.27 (10)
pCO ₂ (μ atm)	peat	N/A	N/A	1403.89 \pm 674.79 (46)	925.64 \pm 868.56 (35)	4016.69 \pm 2120.03 (30)	2197.11 \pm 1771.41 (30)
	bay	N/A	N/A	1050.00 \pm 552.68 (7)	297.81 \pm 93.57 (11)	1161.74 \pm 1275.46 (10)	1151.68 \pm 968.31 (10)
pH	peat	N/A	N/A	7.66 \pm 0.21 (46)	8.01 \pm 0.33 (35)	7.35 \pm 0.34 (30)	7.60 \pm 0.32 (30)
	bay	N/A	N/A	7.78 \pm 0.20 (7)	8.32 \pm 0.13 (11)	7.95 \pm 0.48 (10)	7.86 \pm 0.36 (10)
C _T (μ mol kg ⁻¹)	peat	N/A	N/A	2153.61 \pm 121.07 (46)	2471.11 \pm 223.74 (35)	2539.09 \pm 225.34 (30)	2273.41 \pm 312.95 (30)
	bay	N/A	N/A	2113.87 \pm 73.73 (7)	2201.63 \pm 98.45 (11)	2094.51 \pm 208.11 (10)	2106.76 \pm 282.17 (10)
A _T (μ mol kg ⁻¹)	peat	N/A	N/A	2154.43 \pm 155.12 (46)	2614.86 \pm 209.57 (35)	2546.03 \pm 239.96 (30)	2290.59 \pm 272.70 (30)
	bay	N/A	N/A	2144.41 \pm 94.49 (7)	2414.45 \pm 123.87 (11)	2270.25 \pm 125.07 (10)	2187.83 \pm 213.75 (10)

Table 3. Statistical comparison of pre- and post-rewetting nutrient concentrations and GHG fluxes. For pre- and post-rewetting phases, summer and autumn seasons were used (June to November 2019 and 2020, respectively). Nutrient concentrations are compared for the inner bay and GHG fluxes for the peatland site. *** and "n.s." indicate $p < 0.001$ and not significant, respectively.

	location	Pre-rewetting		Post-rewetting		<i>p</i>
		mean \pm SD	n	mean \pm SD	n	
NH ₄ ⁺ (μ M)	inner bay	2.6 \pm 1.6	9	9.9 \pm 16.9	20	n.s.
NO ₃ ⁻ (μ M)	inner bay	1.5 \pm 2.3	10	1.8 \pm 2.9	20	n.s.
NO ₂ ⁻ (μ M)	inner bay	0.2 \pm 0.1	10	0.6 \pm 1.0	20	n.s.
PO ₄ ³⁻ (μ M)	inner bay	0.6 \pm 1.3	10	0.2 \pm 0.2	20	n.s.
CO ₂ flux (g m ⁻² h ⁻¹)	transect + area	0.3 \pm 0.8	330	0.3 \pm 0.3	450	n.s.
CO ₂ flux (g m ⁻² h ⁻¹)	ditch	0.3 \pm 0.1	87	0.3 \pm 0.3	92	n.s.
CH ₄ flux (mg m ⁻² h ⁻¹)	transect + area	0.1 \pm 1.0	97	1.7 \pm 7.6	320	***
CH ₄ flux (mg m ⁻² h ⁻¹)	ditch	11.4 \pm 37.5	85	8.5 \pm 26.9	92	***

3.2 Nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻)

395 3.2.1 Pre- and post-rewetting spatio-temporal dynamics and comparison with a nearby monitoring station

In the inner bay, all N-nutrient concentrations were substantially higher at the first sampling after rewetting than prior to rewetting, while PO₄³⁻ concentrations were only slightly higher post-rewetting (Figure 4). This increase of N-nutrients led to a drastic increase of the N:P ratio from ~73 in autumn 2019 before rewetting to ~1600 shortly after rewetting in winter 2019. A comparison of the same pre- and post-rewetting seasons (summer and autumn 2019/2020) showed generally higher N-
400 nutrient concentrations in the inner bay after rewetting which could not be confirmed statistically (Mann-Whitney-U-test, Table 3).

During winter, all N-nutrients were high in the peatland and inner bay. After a rapid decrease in spring, N-nutrient concentrations reached their lowest values during summer, with NH₄⁺ and NO₂⁻ then increasing in autumn again. PO₄³⁻ concentrations followed a different pattern, with the highest concentrations determined in summer and fewer fluctuations over
405 the year.

The spatial differences in nutrient concentrations between the inner bay and the peatland after rewetting varied greatly between the nutrient species. From the N-nutrients, only NO₂⁻ concentrations were significantly higher once in winter, shortly after rewetting, whereas NH₄⁺ and NO₃⁻ concentrations showed no significant differences in any season (Table 2). Significantly higher PO₄³⁻ concentrations in the peatland occurred during spring and summer ($p < 0.05$). Some significant correlations
410 between nutrient species were found (Figure D1), especially between NO₂⁻/NH₄⁺ and NO₃⁻/NO₂⁻ both in the peatland and the inner bay.

Nutrient concentrations of the monitoring station (“central bay”) showed a low inter-annual variability during the years 2016–2020 and often lower concentrations than the inner bay (Figure 5). A detailed comparison of nutrient data from the monitoring station with those from the inner bay showed that before rewetting, only the NH₄⁺ concentrations were

415 significantly higher in the inner bay. After rewetting, NO_3^- and NO_2^- concentrations in the inner bay increased and were significantly higher than in the central bay ($p < 0.001$ and $p < 0.05$, respectively). In spring, N-nutrient concentrations were similar at the two locations whereas in summer, all N-nutrients were significantly higher in the inner bay ($p < 0.01$). In autumn, NO_2^- and NH_4^+ concentrations increased again and thus, showed significantly higher concentrations in the inner bay. PO_4^{3-} again followed a pattern different from that of the N-nutrients. Shortly before rewetting, its concentrations in the inner bay were significantly lower than those in the central bay ($p < 0.05$). After rewetting, PO_4^{3-} concentrations showed no significant differences in any season.

420




Figure 4. Time series of the mean (a) PO_4^{3-} , (b) NO_3^- , (c) NO_2^- , and (d) NH_4^+ concentrations (\pm standard deviations) in the surface water from June 2019 to December 2020. Data from the flooded peatland ($n = 6$) are shown in blue and data from the inner bay (until 11 March 2020: $n = 1$, thereafter: $n = 2$) in black. The vertical black line indicates the rewetting event.

425




Figure 5. Seasonal nutrient concentrations of (a) NO_3^- , (b) NH_4^+ , (c) NO_2^- , and (d) PO_4^{3-} at the nearby monitoring station (central bay, red) and in the inner bay (inner bay, blue) from pre- to post-rewetting. The vertical black line indicates the rewetting event. Note that 5-year-data (2016–2020) are shown for the central bay (see Sect. 2.4.2). ns = not significant, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.

3.2.2 Nutrient export from the rewetted peatland into the inner bay


The rewetted peatland was a net source of DIN-N and $\text{PO}_4\text{-P}$ for the inner bay (Table B1). During the first year after rewetting, $10.8 \pm 17.4 \text{ t yr}^{-1}$ DIN-N and $0.24 \pm 0.29 \text{ t yr}^{-1}$ $\text{PO}_4\text{-P}$ were exported into the inner bay (given as mean \pm 95 % confidence level, equivalent to megagrams yr^{-1}). DIN-N export was highest during the winter directly after rewetting ($8.6 \pm 9.9 \text{ t}$) and lowest during summer ($0.3 \pm 0.5 \text{ t}$). DIN-N and $\text{PO}_4\text{-P}$ were only exported from the peatland into the inner bay in all seasons.

N-nutrient concentrations showed a gradient from the peatland through the inner bay to the central bay. Therefore, nutrient data from the central bay were also taken into account to estimate the total possible export from the peatland to the sea. This resulted in an estimated total net export of $33.8 \pm 9.6 \text{ t yr}^{-1}$ DIN-N. In contrast to the comparison of the peatland and the inner bay, $\text{PO}_4\text{-P}$ was once imported from the central bay into the peatland in autumn ($0.03 \pm 0.10 \text{ t}$). Additionally, it was noticeable that the $\text{PO}_4\text{-P}$ concentrations in the central bay were permanently higher than in the inner bay, leading to a lower annual export of $0.09 \pm 0.32 \text{ t yr}^{-1}$ $\text{PO}_4\text{-P}$.

3.3 GHG in the surface water after rewetting

3.3.1 Inorganic C system


During the winter after rewetting, the differences in the CO₂ system (C_T, A_T, pH, pCO₂) between the inner bay and the peatland were not significant (Figure 6, Figure 7a). All variables increased slightly until spring, coinciding with a slight increase in salinity over the same period. From spring onwards, however, the components of the CO₂ system followed contrasting patterns, with C_T and A_T remaining relatively constant in the inner bay but reaching significantly higher values in the peatland ($p < 0.05$), including maximum values in summer (Table 2). The pH also showed significant seasonal differences, with lower values and a minimum in summer in the peatland ($p < 0.05$). C_T and A_T values in the inner bay and in the peatland aligned in autumn whereas the pH remained significantly different ($p < 0.05$). The mean pCO₂ (calculated from C_T and pH) of the surface water in winter was $1050.0 \pm 55.7 \mu\text{atm}$ in the inner bay and $1403.9 \pm 674.8 \mu\text{atm}$ in the peatland (Figure 7a). The pCO₂ values were highest during the first few weeks after inundation and then steadily decreased, with the lowest mean values occurring in spring (peatland) and summer (inner bay). The summer was characterized by high pCO₂ values in general, including earlier and stronger increases in the peatland than in the inner bay that resulted in significant differences in spring and summer ($p < 0.05$ for both seasons). pCO₂ values were highest in summer with $4016.7 \pm 2120.0 \mu\text{atm}$ (peatland) and $1161.7 \pm 1275.5 \mu\text{atm}$ (inner bay) (Table 2). In October, all of the examined CO₂ quantities had a short-term inversion of the prevailing pattern.



460 **Figure 6.** Time series of the mean (a) total CO₂ (C_T), (b) total alkalinity (A_T), and (c) pH (± standard deviations) in the surface water after rewetting, as measured from December 2019 to December 2020. Data from the flooded peatland (n = 6) are shown in blue and data from the inner bay (until 11 March 2020: n = 1, thereafter: n = 2) in black. The vertical black line indicates the rewetting event.

3.3.2 CH₄

During the first few months after flooding (in winter), the CH₄ concentrations in both the inner bay and the peatland were low and did not differ significantly (Figure 7b, Table 2): 48.0 ± 49.5 nmol L⁻¹ (peatland) and 81.4 ± 107.0 nmol L⁻¹ (inner bay). From mid-spring onwards, CH₄ concentrations in the inner bay and the peatland increased such that during summer and autumn 2020, the differences at the two areas were significant (*p* < 0.05). Mean CH₄ values were highest in summer and amounted 1502.4 ± 693.4 nmol L⁻¹ in the peatland and 502.5 ± 479.3 nmol L⁻¹ in the inner bay. Further, the peatland was characterized by a considerable short-term variability in spring and summer, expressed in four peaks representing elevated concentrations. A positive significant correlation (*r*_S = 0.73, n = 72, *p* < 0.001) was found in the peatland between the surface water CH₄ concentrations and a water temperature > 10 °C, but not < 10 °C.



470

Figure 7. Time series of the mean (a) pCO₂, (b) CH₄ concentration (cCH₄), and (c) N₂O concentration (cN₂O) (± standard deviations) after rewetting in the surface water from December 2019 to December 2020. Data from the flooded peatland (n = 6) are shown in blue and data from the inner bay (until 11 March 2020: n = 1, thereafter: n = 2) in black. The vertical black line indicates the rewetting event.

3.3.3 N₂O

475 The highest N₂O concentration of 486.3 nmol L⁻¹ was measured in the peatland one week after rewetting (Figure 7c), followed by 4–5 weeks of still elevated N₂O concentrations between 19.9 and 91.8 nmol L⁻¹. During winter, significant positive correlations were determined in the peatland between N₂O and NH₄⁺ ($r_s = 0.61$, $n = 45$, $p < 0.001$) and between N₂O and NO₂⁻ ($r_s = 0.46$, $n = 45$, $p < 0.01$). From spring onwards, N₂O decreased rapidly, both in the peatland and the inner bay, with the lowest values of 4.7 to 7.9 nmol L⁻¹ reached in summer. Other positive correlations of N₂O with N-nutrients in the peatland
480 included NO₃⁻ ($r_s = 0.74$, $n = 35$, $p < 0.001$) and NO₂⁻ ($r_s = 0.70$, $n = 35$, $p < 0.001$) in spring and all N species in autumn (NO₃⁻: $r_s = 0.85$, $n = 30$, $p < 0.001$; NO₂⁻: $r_s = 0.70$, $n = 30$, $p < 0.001$; NH₄⁺: $r_s = 0.80$, $n = 30$, $p < 0.001$).

Spatial differences in N₂O concentrations between the inner bay and the peatland were low and not significant in winter, spring or autumn, whereas significantly lower concentrations were measured in the peatland during summer (Table 2).


3.4 Pre- and post-rewetting GHG fluxes (CO₂, CH₄, N₂O)

485 Terrestrial CO₂ fluxes before rewetting, during summer and autumn 2019, were highly variable ranging from -3.3 to 3.0 g m⁻² h⁻¹ with a mean ± SD of 0.29 ± 0.82 g m⁻² h⁻¹ (Figure 8a). Within the ditch, pre-rewetting CO₂ fluxes ranged from -0.008 to 0.6 g m⁻² h⁻¹, but on average were comparable to the fluxes determined at the terrestrial (dry) surface.

After rewetting, formerly terrestrial CO₂ fluxes decreased in amplitude (-0.5 to 1.4 g m⁻² h⁻¹), while the summer and autumn averages were unchanged compared to the pre-rewetting fluxes (Table 3). In the ditch, the mean and minimum post-
490 rewetting CO₂ fluxes were within the range of those determined pre-rewetting (mean: 0.26 ± 0.29 g m⁻² h⁻¹, min: -0.02 g m⁻² h⁻¹) but the maximum flux (1.1 g m⁻² h⁻¹) was almost twice as high as the pre-rewetting ditch flux (max: 0.6 g m⁻² h⁻¹).

Pre-rewetting CH₄ fluxes (mean ± SD) in summer and autumn 2019 varied between -0.9 to 8.4 mg m⁻² h⁻¹ (terrestrial) and -1.1 to 193.6 mg m⁻² h⁻¹ (drainage ditch; Figure 8b). While mean terrestrial CH₄ fluxes were 0.13 ± 1.01 mg m⁻² h⁻¹, the
495 mean ditch fluxes were 11.4 ± 37.5 mg m⁻² h⁻¹. In summer and autumn 2020, after rewetting, average CH₄ fluxes on formerly terrestrial land increased slightly but significantly (1.74 ± 7.59 mg m⁻² h⁻¹), whereas in the ditch they decreased considerably (8.5 ± 26.9 mg m⁻² h⁻¹). Flux amplitudes at the ditch station before and after rewetting were comparable.

Data on N₂O fluxes are available only for the post-rewetting period. The rewetted peatland was a small source of N₂O, with an annual mean (± SD) flux of 0.02 ± 0.07 mg m⁻² h⁻¹ in the first year after rewetting (Figure 8c). The highest N₂O
500 flux of 0.4 mg m⁻² h⁻¹ occurred one week after rewetting, followed by lower N₂O fluxes between 0.007 and 0.2 mg m⁻² h⁻¹ within the following 4–5 weeks. Afterwards, N₂O fluxes remained constantly close to zero. Negative fluxes, indicating N₂O uptake, were measured only in summer.



505 **Figure 8.** Time series of the mean (a) CO₂, (b) CH₄, and (c) N₂O fluxes (\pm standard deviations) from June 2019 to December 2020. Fluxes of the permanently wet drainage ditch are shown in purple and those derived from the two methods employed in this study in green. The vertical black line indicates the rewetting event.

4. Discussion

4.1 Nutrient dynamics and export

The seasonal dynamics of the nutrients followed a typical course over the year. Thus, after rewetting, NH₄⁺, NO₃⁻, and NO₂⁻ concentrations in the water column were high in winter and autumn, which is typically due to the mineralization of OM followed by nitrification (Voss et al., 2010). By contrast, the low DIN concentrations during spring and summer reflected the consumption of nutrients by plants and phytoplankton. The very high chlorophyll a concentration (up to 125 μ g L⁻¹) in the peatland indicated the presence of a highly phototrophic community, likely driven by the higher availability of nutrients compared to the inner bay. Due to these distinct seasonal differences with the lowest nutrient concentrations in spring and summer, a rewetting within these seasons would probably be more beneficial to reduce a potential nutrient export into the inner bay, at least during the first few months after rewetting.

510

515

To assess whether the flooded peatland served as a nutrient source for the inner bay, nutrient concentrations of the peatland were compared with those of the inner bay and of an unaffected monitoring station (“central bay”) and showed generally higher mean concentrations. Due to drainage, the mineralization of upper peat layers can lead to an accumulation of

520 nutrients within the soil (Zak and Gelbrecht, 2007; Cabezas et al., 2012). After rewetting, nutrient concentrations in the
porewater and ultimately in the overlying water increase (van de Riet et al., 2013; Harpenslager et al., 2015; Zak et al., 2017).
The leaching of nutrients is driven by concentration differences across the soil-water interface, but it is also dependent on
factors such as salinity (Rysgaard et al., 1999; Steinmuller and Chambers, 2018), the oxygen availability in the soil (Lennartz
and Liu, 2019), and the effects of the latter on microbial processes (Burgin and Groffman, 2012), as well as on the degree of
525 peat decomposition (Cabezas et al., 2012). For instance, highly degraded peat, such as at our study area, can store and release
more nutrients than less degraded peat (Cabezas et al., 2012), meaning that the highly degraded peat of our study area was
prone to leach high amounts of nutrients. Occasional measurements of porewater nutrient concentrations in the peat of our
study area revealed DIN and PO_4^{3-} concentrations up to one order of magnitude higher than those in the surface water (Anne
Breznikar, unpublished data), providing further support for the leaching of nutrients out of the peatland and into the inner bay.

530 The estimated annual nutrient exports (mean \pm 95 % confidence level) from the peatland of $10.8 \pm 17.4 \text{ t yr}^{-1}$ DIN-N
and $0.24 \pm 0.29 \text{ t yr}^{-1}$ $\text{PO}_4\text{-P}$ (peatland/inner bay) and $33.8 \pm 9.6 \text{ t yr}^{-1}$ DIN-N and $0.09 \pm 0.32 \text{ t yr}^{-1}$ $\text{PO}_4\text{-P}$ (peatland/central
bay) were high, given the small size of the flooded peatland ($\sim 0.5 \text{ km}^2$ at 0 masl). For comparison, the Warnow, a small river
that flows into the Baltic Sea near the city of Rostock, Mecklenburg-Vorpommern, draining an area of $\sim 3300 \text{ km}^2$, had a mean
annual DIN-N and $\text{PO}_4\text{-P}$ export of $1200 \pm 500 \text{ t yr}^{-1}$ and $19.9 \pm 7.6 \text{ t yr}^{-1}$, respectively, over the last 25 years (HELCOM,
535 2019). Therefore, the total nutrient export from the flooded peatland to the inner bay and to the central bay in the first year
after rewetting accounted for ~ 1 and ~ 3 %, respectively, of the annual DIN-N and $\text{PO}_4\text{-P}$ loads of the Warnow. When
normalized to the same dimensions, our study area exported $21.6\text{--}67.6 \text{ t DIN-N km}^{-2} \text{ yr}^{-1}$ and $0.18\text{--}0.48 \text{ t PO}_4\text{-P km}^{-2} \text{ yr}^{-1}$,
whereas the Warnow exported only $0.36 \text{ t DIN-N km}^{-2} \text{ yr}^{-1}$ and $0.01 \text{ t PO}_4\text{-P km}^{-2} \text{ yr}^{-1}$.

However, we also want to shortly address the reasons for the high uncertainty range of our calculated nutrient exports.
540 Firstly, they derive from high fluctuating nutrient concentrations in the surface water within the seasons. This is also visible in
the high standard deviations (Table 2). Therefore, the 95 % confidence level of the nutrient exports is high and reflects the
natural dynamic. Secondly, we conducted default error propagation during the export calculation which leads to even higher
ranges on top of the high natural dynamic.

Compared to the Warnow river, it is noticeable that the range of uncertainties is highly different for the two sources.
545 While our uncertainties are mostly higher and in the same order of magnitude compared to the means, the uncertainties of the
river data are one order of magnitude lower. This is likely due to the different time scales of the two data sets. Our export data
were generated by taking only the first post-rewetting year into account in which the system was still in a transition state and
thus, showed very dynamic nutrient concentrations. The uncertainties of the river exports were generated by using 25 years of
data, leading to lower uncertainties than using data from only one year and they were calculated as standard deviation and not
550 as 95 % confidence level, as was done for the exports of our study site. Therefore, this has to be considered when their
uncertainty ranges are compared directly. Nevertheless, our results highlight the importance of currently still unmonitored and
small, independent draining areas along the coastline of the Baltic Sea, in particular those that become intentionally flooded
(HELCOM, 2019).

4.2 Assessment of the GHG dynamics

555 4.2.1 CO₂

The carbon system in our study area is governed by a variety of processes (e.g. Wolf-Gladrow et al., 2007; Kuliński et al., 2017; Schneider and Müller, 2018). C_T and A_T were transported with the brackish water from the central bay and ultimately from the Arkona Basin. Additional alkalinity can be added either by a supply of freshwater, which in the southwestern Baltic Sea is characterized by higher alkalinities than the brackish or even saltwater endmember (Beldowski et al., 2010; Müller et al., 2016), or can be introduced by mineralization processes from the seafloor in the inner bay and the flooded peatland. Primary production (i.e., carbon fixation) will decrease C_T, lower the pCO₂ and increase pH during the formation of organic matter. The mineralization of OM from various sources (new primary production, mineralization of the inundated former vegetation and from the underlying peat) will enhance C_T and A_T concentrations, increase pCO₂ and decrease pH. Air-sea exchange during our study is fostered by a pCO₂ that is above atmospheric levels throughout the year, except for a short period in spring in the inner bay and the peatland, where outgassing of CO₂ occurred, resulting in lower pCO₂ and a decrease in C_T.

We observed three main developments in the surface water CO₂ system and air-sea flux pattern: (i) in winter 2019/2020, the CO₂ system hardly differed between the peatland and the inner bay; (ii) from spring to autumn, there were significant differences in the CO₂ system between the peatland and the inner bay, with higher pCO₂, C_T and A_T values and lower pH in the peatland coinciding with an enrichment in chlorophyll a; (iii) overall, the first post-rewetting year showed sustained high, but less variable CO₂ fluxes compared to pre-rewetting conditions. In the following, we will discuss these three observations and set them into context.

4.2.1.1 Initial post-rewetting CO₂ dynamics

The first weeks after the rewetting were characterized by high nutrient concentrations, a continuous increase in A_T, C_T and pH and a decrease in pCO₂ (Figure 4, Figure 6, Figure 7). The increase in C_T and A_T coincided with a steady increase in salinity (Figure 3), which is in line with a general increase of A_T with increasing salinity known for the western Baltic Sea (e.g. Kuliński et al., 2022).

Still, the A_T values at the given salinity were higher in the inner bay and the peatland than would be expected from a linear A_T/salinity relation found for surface waters in the open Baltic Sea from the central Gotland Sea to the Kattegat (Beldowski et al., 2010; Müller et al., 2016). Thus, the high A_T in the inner bay and peatland were likely associated with local carbonate (CaCO₃) weathering from terrestrial sources and/or a transport by groundwater (Schneider and Müller, 2018). C_T and A_T values during this period were consistently higher by ~70–80 μmol kg⁻¹ in the peatland than in the inner bay, consistent with enhanced leaching from the recently inundated peat. Besides, local CaCO₃ weathering as well as local anoxic processes, such as SO₄²⁻ reduction may have increased the A_T in the submerged soil and finally contributed to higher A_T values compared to the inner bay.

The oversaturation in pCO₂ and potentially the excess leaching of alkalinity from the soil might have contributed to the decrease in pCO₂ and increase in pH in the peatland in winter 2019/2020. This was apparently reinforced by a short episode

of primary production mid/end January, indicated by a steeper decline of the pCO₂ and a steeper pH increase. This coincided with a short increase in chlorophyll a (~30 µg L⁻¹) and a slight intermittent increase of the surface water temperatures (Figure 3). This short, unusually early productive period might have resulted from the high nutrient availability induced by the rewetting of the peatland (Sect. 3.2.1), in particular the high NH₄⁺ levels, which simultaneously showed a sharp intermittent minimum.

4.2.1.2 The predominance of production and mineralization shaped the productive period (spring to autumn)

In late winter and the first half of spring, pCO₂ continuously decreased in the peatland as well as in the inner bay. Lowest pCO₂ was measured between March and May and coincided with enhanced chlorophyll a concentrations and a high availability of nutrients in the peatland and in the inner bay, which decreased until mid spring. This resulted in a slight CO₂ uptake in the peatland of -0.005 g m⁻² h⁻¹ for a short period of time, so that spring was the only season when pCO₂ was on average below atmospheric concentrations in the inner bay (Figure 8). This finding can be attributed to the onset of the productive period, at still moderate surface water temperatures below 10 °C until mid April. During this period, productivity clearly exceeded mineralization, as suggested by the decreasing pCO₂ and increasing pH, despite rising temperatures, as well as increasing O₂ oversaturation in the surface waters. These trends were slightly more pronounced in the peatland than in the inner bay, in accordance with higher nutrient concentrations available for production.

From mid spring until late summer, the peatland was characterized by increased pCO₂ and a variable CO₂ system together with high mean chlorophyll a concentrations of up to 106.0 µg L⁻¹. N-nutrients were very low and the system was clearly nitrogen-limited, with only slightly elevated NH₄⁺ concentrations in late summer (Figure 3, Figure 4). Furthermore, the O₂ saturation shifted from over- to undersaturated conditions. These observations suggest that the peatland and the inner bay were characterized by simultaneous production and mineralization processes from mid spring until autumn that kept the N-nutrients (except PO₄³⁻) low. Mineralization of OM in the water column, sediment and soil dominated over production, leading to the observed high pCO₂, lowered pH, and enhanced A_T and C_T concentrations. Mineralization during this period was more pronounced in the peatland than in the inner bay, leading to the higher pCO₂, A_T, and C_T values in the peatland, and a stronger and more pronounced reduction of the pH. This stronger mineralization, in particular in the warm summer months, also led to higher DOC concentrations in summer, with a maximum in June/July coinciding with maximum surface water temperatures. The enhanced mineralization in the peatland was likely fueled by higher OM availability from high decomposition rates of fresh plant substrate from inundated plant residuals (Glatzel et al., 2008; Hahn-Schöfl et al., 2011). In addition, aerobic and anaerobic oxidation of CH₄, that was produced in anoxic zones, might have led to increased CO₂ production, especially during increased water temperatures (e.g. Treude et al., 2005; Dean et al., 2018), due to the availability of SO₄²⁻ and O₂.

The calculated A_T (from C_T and pH) in the peatland was consistently lower than the measured A_T with a difference in the range of 55–122 µmol kg⁻¹ and thus of 2.7–4.7 % (data not shown). This difference was higher than in the Baltic Sea, where the contribution of organic A_T is estimated to be 1.5–3.5% (Kuliński et al., 2014). Due to closer vicinity to the coast and the high amount of degradable OM, this higher contribution of organic A_T was to be expected. The highest discrepancy between measured A_T values and those calculated from pH and C_T occur in early summer, simultaneously to the highest values in DOC,

in particular in the peatland (Figure 3). This suggests that the organic A_T related to the occurrence of DOC (and thus dissolved organic matter (DOM)), contributed to the excess of A_T . The higher DOM formation in summer in the peatland might partly explain the difference in A_T between the inner bay and the peatland.

4.2.1.3 Brackish water flooding caused sustained high but less variable CO_2 fluxes

625 The amplitude of the CO_2 fluxes from formerly drained parts of the study area decreased after rewetting with brackish water, while the amplitude of CO_2 fluxes from the ditch (inundated after flooding but with deeper, probably incompletely exchanged water) did not differ strongly before and after rewetting (Figure 8). An increased water table is the main driver for the reduction of CO_2 emissions on formerly drained locations. A similar scenario has been reported for terrestrial sites (Bubier et al., 2003; Strack, 2008). In a nearby coastal peatland, both photosynthesis and ecosystem respiration were strongly reduced after
630 rewetting (Koebsch et al., 2013). The rewetting of our study area probably caused a die-back of the highly productive grassland vegetation, which most likely lead to a reduction of the CO_2 flux amplitude.

Average summer/autumn CO_2 fluxes after rewetting had a mean of $0.26 \pm 0.29 \text{ g m}^{-2} \text{ h}^{-1}$ and remained thus relatively high compared with those fluxes from 2019. They were also higher than the fluxes determined in studies of shallow coastal or near-shore waters in the northwestern Bornholm Sea of up to $0.01 \text{ g m}^{-2} \text{ h}^{-1}$ (Thomas and Schneider, 1999) or the Bothnian
635 Bay of around $\sim 0.0007 \text{ g m}^{-2} \text{ h}^{-1}$ (Löffler et al., 2012). In a nearby coastal fen recently influenced by brackish water inflow, ecosystem respiration was two orders of magnitude lower (Koebsch et al., 2020) compared to our study site, where the ongoing decomposition of submerged substrate from plant residuals and the fresh soil may have fueled the continuously high CO_2 fluxes in the first year after rewetting (Hahn-Schöfl et al., 2011). The mineralization of OM from primary production driven by the high initial nutrient availability, as well as aerobic and anaerobic oxidation (Dean et al., 2018) of easily degradable
640 substrates or CH_4 , might have additionally contributed to these CO_2 fluxes. We expect that CO_2 emissions will further decrease, likely because substrates become exhausted and a novel ecosystem will be established (Kreyling et al., 2021), with developing algae fostering CO_2 fixation.

4.2.2 CH_4

We observed three main developments in surface water methane concentrations and flux patterns: (i) a short-term, very
645 moderate increase in CH_4 concentrations directly after rewetting in winter 2019/2020; (ii) an increase in the CH_4 concentrations mainly from spring to autumn, that was significantly higher and more variable in the peatland than in the inner bay and correlated with water temperature; (iii) in the first year after rewetting, much lower CH_4 fluxes than reported for nearby peatlands rewetted by freshwater. These three observations are discussed and set into context in the following.

4.2.2.1 Short-term, moderate increase in the CH_4 concentrations in the winter after rewetting

650 The measurements in winter, immediately after rewetting, showed a short-term but moderate increase in the CH_4 concentrations (Figure 7). The rewetting resulted in inundation of the degraded peat and the remaining vegetation. Therefore, it is assumed that methanogenesis was not limited by the availability of high-quality OM, which is often a major controlling factor (Heyer and Berger, 2000; Parish, 2008), and likely originated from decomposition of the residual plant material.

655 However, since CH₄ concentration remained low, a temperature control is assumed, which has been frequently described in the literature. A major control of temperature has been reported, for example, for a nearby shallow coastal area of the Baltic Sea, between the islands of Rügen and Hiddensee, where low CH₄ emission rates and variability were found together with low temperatures (Heyer and Berger, 2000).

660 The rewetting with brackish water transported water with a salinity of 6–7.4 into the peatland such that there were no significant differences in salinity compared to the inner bay in winter (same as for temperature; Table 2). Thus, sulfate reached the peatland immediately after rewetting. As a terminal electron acceptor (TEA), SO₄²⁻ promotes the activity of sulfate-reducing bacteria (SRB), which outcompete methane-producing microorganisms (methanogens) for substrates (Segers and Kengen, 1998; Jørgensen, 2006; Segarra et al., 2013). This process was shown to play an important role in flat brackish water systems (e.g. Heyer and Berger, 2000). The availability of other TEAs, such as NO₃⁻ that had high concentrations of ~100 ± 58 μM in our study, could have further suppressed methanogenesis (Table 2) (Jørgensen, 2006). Beside competitive 665 mineralization, aerobic and anaerobic CH₄ oxidation may have reduced the CH₄ concentrations (Heyer and Berger, 2000; Reeburgh, 2007; Knittel and Boetius, 2009; Steinle et al., 2017), supported by the effective exchange of water masses. Overall, the rewetting with brackish water during the cold winter season apparently inhibited methanogenesis and/or facilitated effective CH₄ oxidation, resulting in low CH₄ concentrations and a small CH₄ flux into the atmosphere.

4.2.2.2 Increased and variable CH₄ concentrations during the vegetation period

670 The temperature increase from spring to autumn was accompanied by elevated, albeit variable, CH₄ concentrations. Temperature is of crucial importance in controlling the CH₄ cycle in shallow coastal brackish water (Bange et al., 1998; Heyer and Berger, 2000) and in the North Sea (e.g. Borges et al., 2018). Similar relationships have been described for wetlands, e.g. for permanently inundated wetlands (e.g. Koebsch et al., 2015) and in a peatland close to our study site during the first year after rewetting (Hahn et al., 2015). Accordingly, CH₄ concentrations in the peatland ($r_s = 0.75$, $n = 74$, $p < 0.05$) and the inner 675 bay ($r_s = 0.55$, $n = 29$, $p < 0.05$) also correlated significantly and positively with temperature. In the study of Heyer and Berger (2000) the temperature range influenced the temporal variability in CH₄ emissions, which were highest in late spring. Since the temperature range in the peatland of our study was variable (e.g. maximum difference of ~6 °C between samplings), with the highest values between spring and autumn (7.4–23.1 °C), this variability may have strongly contributed to the observed CH₄ dynamics.

680 The peatland and the inner bay were clearly influenced by the same hydrographic conditions, evidenced by their very similar salinities and temperatures. However, the peatland showed higher CH₄ concentrations from spring to late autumn, likely due to the high availability of OM as described by Heyer and Berger (2000) and Bange et al. (1998). Incubation experiments of a degraded fen grassland demonstrated the accumulation of fresh plant litter in a new sediment layer after flooding that resulted in high rates of CH₄ and CO₂ production (Hahn-Schöfl et al., 2011). A further potential driver of OM 685 availability is the sedimentation of freshly produced OM originating from primary production, as described for shallow areas in the Baltic Sea (Bange et al., 1998) and for a shallow bight in the North Sea, which in the latter led to a yearly peak in the seasonal CH₄ cycle (Borges et al., 2018). Although our observations were not made in OM-poor sediments, an impact of

freshly produced OM on enhanced CH₄ concentrations in the OM-rich Drammendorf peatland is possible, given the significant positive correlation of the surface CH₄ concentrations and the chlorophyll a concentration ($r_s = 0.41$, $n = 56$, $p < 0.05$).

690 4.2.2.3 Brackish water rewetting and low CH₄ emissions

Despite high surface water CH₄ concentrations in the peatland and their inter-seasonal and spatial variability, rewetting with brackish water resulted in CH₄ emissions considerably lower than those from temperate fens rewetted with freshwater, where CH₄ emissions strongly increased (Augustin and Chojnicki, 2008; Couwenberg et al., 2011; Hahn et al., 2015; Franz et al., 2016; Jurasinski et al., 2016).

695 At our study site, although average CH₄ fluxes on formerly terrestrial locations increased significantly by one order of magnitude after rewetting, the overall increase from 0.13 ± 1.01 to 1.74 ± 7.59 mg m⁻² h⁻¹ (Figure 8) was lower than that reported for freshwater rewetted fens under similar climatological boundary conditions (e.g. Hahn et al., 2015; Franz et al., 2016). Even several years after rewetting, the annual CH₄ budgets of a shallow lake on a formerly drained fen varied between 13.2 and 52.6 g m⁻² yr⁻¹ (Franz et al., 2016), which corresponds to approximately 1.5 to 6.0 mg m⁻² h⁻¹. Our CH₄ fluxes were
700 also lower than the emissions reported from coastal-near shallow waters of the Baltic Sea, where fluxes of 39.9–104.2 mg m⁻² h⁻¹ were measured in June/July (Heyer and Berger, 2000). For the same months, mean CH₄ fluxes at the formerly dry stations in our study site were 0.5–4.9 mg m⁻² h⁻¹. However, compared to CH₄ fluxes from continental shelves (0.015–0.024 mg m⁻² h⁻¹; adapted from Bange et al., (1994)), the fluxes of our study site were two orders of magnitude higher. Despite low average fluxes, emission peaks could be distinguished with the highest flux from the now inundated ditch of
705 149.2 mg m⁻² h⁻¹ in September 2020 and 108.3 mg m⁻² h⁻¹ in October 2020. While these values were still lower than the maximum value of 243.0 mg m⁻² h⁻¹ reported by Heyer and Berger (2000), it is important to stress that our study site was a source of CH₄ already in its drained state, especially within the drainage ditch, where CH₄ fluxes were comparable to the ~0.2 mg m⁻² h⁻¹ reported from undrained fens (Danevčič et al., 2010).

The lower CH₄ emissions of the brackish rewetted Drammendorf peatland might be attributed to the availability of
710 TEAs, especially SO₄²⁻, which (1) may have contributed to a suppression in methanogenesis by competitive inhibition (Segers and Kengen, 1998; Jørgensen, 2006; Segarra et al., 2013) or (2) fostered the anaerobic oxidation of methane (AOM) as an effective pathway to reduce CH₄ emissions, and by (3) fast aerobic CH₄ oxidation mediated by oxygen-rich water. The high variability of CH₄ concentrations may also be related to changing rates of AOM, as the process is sensitive to the introduction of O₂ mediated by sporadic wind-driven resuspension (Treude et al., 2005). Since our study area was shallow and likely
715 experiences regular wind-driven resuspension, spatially and temporally dynamic AOM can be assumed. The low CH₄ fluxes suggested that effective aerobic and anaerobic oxidation of CH₄ likely occurred. Moreover, higher CH₄ concentrations in the peatland compared to the inner bay in combination with the high lateral water exchange due to frequent changes in the water level (Figure A3) should have driven a net advective export of CH₄-enriched water to the inner bay. This would have further contributed to the low peatland CH₄ emissions and the observed high variability.

720 While CH₄ production and emission were likely prevented by rewetting with oxygen-rich, sulfate-containing brackish water, the possibility remains that the total CH₄ release was underestimated by insufficient accounting for ebullition. In the

marine environment, bubble-mediated transport is attributed to gassy sediments and an effective mechanism of vertical CH₄ migration (e.g. Borges et al., 2016). Although neither of the methods used to determine CH₄ fluxes specifically account for ebullition, we estimated that 6.9 % of all analyzed chamber-based fluxes were partly bubble-influenced. We estimated this percentage by counting the measurements which showed irregular data points in graphical depiction, but did not influence the linear slope. We observed further that in another 9.6 % of the chamber-based flux measurements the CH₄ concentration patterns indicated ebullition (flux measurements with exponential slope, which was clearly steeper than the linear regression of the majority of data points), but these were not accounted for in the final calculations of diffusive flux. Thus, given that only 16.5 % of the chamber-based flux measurements indicated bubble-mediated CH₄ transport and in almost half of those cases, the resulting perturbation was small and was included in the flux amplitude, the magnitude of the ebullition-driven underestimation of our flux estimates is considered to be small.

In summary, the increase in CH₄ concentrations after rewetting in winter was small, short-lived and associated with the die-back of plants. CH₄ fluxes in the first year after rewetting remained relatively low and were lower than typical for post-rewetting conditions. They also followed a seasonal pattern common for shallow organic-rich systems, with a strong correlation with temperature in spring and summer. We anticipate that continuing reduction in OM availability after the initial die-back of vegetation will likely lead to a further decrease of CH₄ emissions in subsequent years.

4.2.3 N₂O

The rewetted peatland was a source of N₂O in the first year after rewetting, although the mean annual N₂O flux of $0.02 \pm 0.07 \text{ mg m}^{-2} \text{ h}^{-1}$ was very low (Figure 8). This was expected since permanent inundation leads to anoxic conditions in the peat, preventing the production of N₂O by nitrification, as well as by denitrification due to the lack of NO₃⁻ (e.g. Succow and Joosten, 2001; Strack, 2008). However, the range of post-rewetting N₂O fluxes in the first three months (winter) was clearly much larger than during the rest of the year, which indicated that N₂O was strongly and immediately affected by the rewetting, as shown elsewhere (Goldberg et al., 2010; Jørgensen and Elberling, 2012). The highest N₂O flux ($0.4 \text{ mg m}^{-2} \text{ h}^{-1}$) and the highest NH₄⁺ concentration (78.0 μM) was measured one week after rewetting and a significant positive correlation between these two variables was found in winter ($r_s = 0.61$, $n = 45$, $p < 0.001$). Additionally, N₂O correlated significantly positive with NO₂⁻ in winter ($r_s = 0.46$, $n = 45$, $p < 0.01$), whereas no correlation with NO₃⁻ was found. The accumulation of N₂O, but also of NO₂⁻ and NO₃⁻ can generally be interpreted as a result of shifting O₂ conditions in the freshly inundated ecosystem, such that incomplete process chains of e.g. nitrification and denitrification were favored (Rassamee et al., 2011). However, it seems likely that the high N₂O concentrations in winter originated from nitrification due to the correlations of N₂O with its substrate (NH₄⁺) and its main accumulating intermediate product (NO₂⁻), as well as a trend of increasing NO₃⁻ concentrations throughout the winter.

During late spring and early summer, undersaturation of the surface water with N₂O, compared to the atmosphere, pointed to consumption within suboxic/anoxic zones of the peat. Consumption in the surface water was unlikely because anoxic conditions were never found near the peat surface. The undersaturation of N₂O a few months after rewetting evidenced

755 the change in O₂ conditions in the peat, from oxic to hypoxic/anoxic, turning the rewetted peatland into an N₂O sink, at least temporarily. This change was likely driven by the higher availability of fresh OM (measured as chlorophyll a) in the peatland compared to the inner bay, finally leading to significantly lower N₂O concentrations in the peatland in summer ($p < 0.001$, Table 2).

760 Previously reported N₂O fluxes in drained peatlands range from 0.002 to 0.45 mg m⁻² h⁻¹, with a clear trend towards higher fluxes in fertilized or naturally N-rich areas (Flessa et al., 1998; Glatzel and Stahr, 2001; Augustin, 2003; Strack, 2008; Minkkinen et al., 2020). Augustin et al. (1998) examined multiple degraded fens in Mecklenburg-Vorpommern and Brandenburg (Germany) and calculated N₂O fluxes of 0.04 to 0.10 mg m⁻² h⁻¹ in extensively and intensively used fen grasslands, respectively (Augustin et al., 1998). N₂O fluxes in drained peatlands result from a low water level which allows the permanent penetration of atmospheric oxygen into the peat to fuel N₂O producing processes that are dependent on oxygen
765 (Martikainen et al., 1993; Regina et al., 1999). As the water level in our study site was permanently below the soil surface before rewetting, it is likely that the drained peat was a source of N₂O. The mean post-rewetting N₂O flux determined in our study area (0.02 ± 0.07 mg m⁻² h⁻¹) is in the lower range of reported fluxes from drained peatlands. Therefore, as shown in other studies (Succow and Joosten, 2001; Minkkinen et al., 2020), the rewetting probably led to a reduction of N₂O fluxes, since they were likely high before the rewetting.

770 In general, the N₂O fluxes in rewetted peatlands are in the same range as fluxes from pristine ones (Minkkinen et al., 2020), indicating that rewetting is a very effective measure to reduce N₂O emissions to natural levels. Literature values range from up to 0.01 and 0.02 mg m⁻² h⁻¹ for rewetted and undrained boreal peatlands (Minkkinen et al., 2020), respectively, to 0.08 mg m⁻² h⁻¹ for a rewetted riparian wetland near a freshwater meadow (Kandel et al., 2019). Although it is difficult to compare the N₂O fluxes determined in this study with those from other sites with different salinity, hydrology and history of
775 use, our mean annual post-rewetting value is in the lower range of N₂O fluxes previously reported for rewetted and pristine peatlands.

5. Conclusions and Outlook

The effects of rewetting a drained coastal peatland with brackish water in winter and the subsequent formation of a permanently inundated area were studied over one year.

780 We found a strong pulse of DIN leaching out of the peat followed by the transport of DIN into the inner bay, leading to a high export especially in winter compared to the Warnow, a nearby river. However, due to a rapid decrease of nutrient concentrations in spring, the nutrient export after a rewetting in spring or summer would likely be lower compared to rewetting in winter, at least during the first few months thereafter.


785 Further, CO₂ concentrations and emissions seem to remain relatively high after the rewetting with brackish water compared to the dry conditions before rewetting. This was likely driven by the high OM availability from the residual vegetation but also by the high rate of primary production in the water column. However, the flux amplitude decreased after

rewetting and thus, peak emissions during the vegetation period were prevented. The lack of a strong increase of CH₄ emissions in the first year after rewetting with brackish water, in contrast to nearby areas rewetted with freshwater, suggests that especially during the colder months, rewetting with brackish water or seawater would minimize CH₄ emissions and thus maximize the effect on integrated GHG emission reduction. Moreover, a rapid elevation of the water level, as occurred at our study site, will promote the oxidation of peat-derived CH₄ in the water column. Future CH₄ emissions will depend on processes, such as the development of vegetation and will likely decrease. According to literature, dry peatlands were found to be rather large sources of N₂O due to their drainage for agricultural use. However, the permanent inundation of our study site led to a rapid decrease of N₂O emissions and converted the peatland into a N₂O sink during summer, with fluxes similar to pristine peatlands.


With the ongoing formation of salt grass meadows, livestock farming at our study area can and will continue. However, the area's use has not hindered its positive development towards an ecosystem with the potential to eventually become a carbon and nutrient sink in the future. We expect that both the nutrient export and GHG emissions will slowly decrease due to a shrinking reservoir of substrates. Nonetheless, because degraded peat is both nutrient- and OM-enriched, this decrease will occur slowly, given that the topsoil was not removed prior to flooding to diminish nutrients and OM, as was demonstrated by other studies. Whether or not the area will act as a C sink in the future depends on the success and speed of the establishment of vascular vegetation and its burial in the anoxic parts of the sediment.

Nutrient export from peatlands and the re-establishment of the nutrient and C-sequestration functions of highly degraded coastal peatlands after rewetting are complex processes whose elucidation requires long-term investigations. The pronounced seasonal dynamics highlight the need for approaches that include a high temporal resolution, such as achieved with sensor-based or eddy-supported measurements.

Appendix A: Study area



810 **Figure A1.** Water level data from the monitoring station “Barhöft” (Wasserstraßen- und Schifffahrtsamt Ostsee), representing the Kubitzer Bodden, from the beginning of rewetting (26 November 2019) until the end of the investigation period. The red dots indicate the sampling days. The dashed horizontal line represents 0 masl. The minimum and maximum water levels of the investigation period are shown by the blue horizontal lines (−0.7 masl and 1.1 masl, respectively). See also Figure A2 and Figure A3.



815 **Figure A2.** The changing water level and its effect on the water coverage of the study area, shown for (a) −0.5 masl and (b) 0.5 masl. Topography data retrieved from the Landesamt für innere Verwaltung Mecklenburg-Vorpommern, Amt für Geoinformation, Vermessungs- und Katasterwesen, Fachbereich Geodatenbereitstellung.




Figure A3: Hypsographic curve of the study area, in increments of 0.1 m. The red dots represent the observed range of the water level during the study. For a water level time series during the sampling period, see Figure A1.

820 Appendix B: Nutrient export calculation

Table B1. Seasonal water volume exchanges (Q_{in}/Q_{out} , $m^3 s^{-1}$) and nutrient masses ($kg m^{-3}$) \pm standard error in the inner bay (c_{IB}), the central bay (c_{CB}), the peatland ($c_{peatland}$), and the resulting net nutrient transport (NNT, in tonnes) for DIN-N and PO₄-P. Negative values of NNT indicate an export from the peatland into the inner bay/central bay and *vice versa*.

season	Q_{in} ($m^3 s^{-1}$)	Q_{out} ($m^3 s^{-1}$)	c_{IB}		NNT DIN-N (t)	$c_{peatland}$		NNT PO ₄ -P (t)
			DIN-N ($kg m^{-3}$)	DIN-N ($kg m^{-3}$)		PO ₄ -P ($kg m^{-3}$)	PO ₄ -P ($kg m^{-3}$)	
winter	1.9 ± 0.1	-1.9 ± 0.1	1270 x 10 ⁻⁶ ± 506 x 10 ⁻⁶	1840 x 10 ⁻⁶ ± 267 x 10 ⁻⁶	-8.6 \pm 9.9	6.5 x 10 ⁻⁶ ± 5.0 x 10 ⁻⁶	11.5 x 10 ⁻⁶ ± 3.7 x 10 ⁻⁶	-0.08 \pm 0.10
spring	1.3 ± 0.1	-1.3 ± 0.1	243 x 10 ⁻⁶ ± 289 x 10 ⁻⁶	391 x 10 ⁻⁶ ± 220 x 10 ⁻⁶	-1.5 \pm 3.8	2.8 x 10 ⁻⁶ ± 2.8 x 10 ⁻⁶	8.1 x 10 ⁻⁶ ± 3.1 x 10 ⁻⁶	-0.05 \pm 0.04
summer	1.1 ± 0.1	-1.1 ± 0.1	44.0 x 10 ⁻⁶ ± 38.2 x 10 ⁻⁶	82.7 x 10 ⁻⁶ ± 34.6 x 10 ⁻⁶	-0.3 \pm 0.5	6.8 x 10 ⁻⁶ ± 4.7 x 10 ⁻⁶	15.2 x 10 ⁻⁶ ± 3.1 x 10 ⁻⁶	-0.07 \pm 0.05
autumn	1.2 ± 0.1	-1.2 ± 0.1	301 x 10 ⁻⁶ ± 218 x 10 ⁻⁶	328 x 10 ⁻⁶ ± 104 x 10 ⁻⁶	-0.4 \pm 3.2	8.1 x 10 ⁻⁶ ± 6.2 x 10 ⁻⁶	10.9 x 10 ⁻⁶ ± 3.7 x 10 ⁻⁶	-0.04 \pm 0.10
total (peatland / inner bay)					-10.8 \pm 17.4			-0.24 \pm 0.29
season	Q_{in} ($m^3 s^{-1}$)	Q_{out} ($m^3 s^{-1}$)	c_{CB}		NNT DIN-N (t)	$c_{peatland}$		NNT PO ₄ -P (t)
			DIN-N ($kg m^{-3}$)	DIN-N ($kg m^{-3}$)		PO ₄ -P ($kg m^{-3}$)	PO ₄ -P ($kg m^{-3}$)	
winter	1.9 ± 0.1	-1.9 ± 0.1	169 x 10 ⁻⁶ ± 63.1 x 10 ⁻⁶	1840 x 10 ⁻⁶ ± 267 x 10 ⁻⁶	-26.2 \pm 5.4	9.9 x 10 ⁻⁶ ± 5.9 x 10 ⁻⁶	11.5 x 10 ⁻⁶ ± 3.7 x 10 ⁻⁶	-0.02 \pm 0.11
spring	1.3 ± 0.1	-1.3 ± 0.1	85.1 x 10 ⁻⁶ ± 42.1 x 10 ⁻⁶	391 x 10 ⁻⁶ ± 220 x 10 ⁻⁶	-3.1 \pm 2.4	4.3 x 10 ⁻⁶ ± 4.7 x 10 ⁻⁶	8.1 x 10 ⁻⁶ ± 3.1 x 10 ⁻⁶	-0.04 \pm 0.06
summer	1.1 ± 0.1	-1.1 ± 0.1	20.2 x 10 ⁻⁶ ± 9.5 x 10 ⁻⁶	82.7 x 10 ⁻⁶ ± 34.6 x 10 ⁻⁶	-0.5 \pm 0.3	8.4 x 10 ⁻⁶ ± 3.4 x 10 ⁻⁶	15.2 x 10 ⁻⁶ ± 3.1 x 10 ⁻⁶	-0.06 \pm 0.04
autumn	1.2 ± 0.1	-1.2 ± 0.1	26.5 x 10 ⁻⁶ ± 9.1 x 10 ⁻⁶	328 x 10 ⁻⁶ ± 104 x 10 ⁻⁶	-3.9 \pm 1.5	13.0 x 10 ⁻⁶ ± 6.5 x 10 ⁻⁶	10.9 x 10 ⁻⁶ ± 3.7 x 10 ⁻⁶	0.03 \pm 0.10
total (peatland / central bay)					-33.8 \pm 9.6			-0.09 \pm 0.32

Appendix C: Comparability of two independent approaches to atmospheric flux determination

825 Since the gas transfer velocity k model (Sect. 2.5.3) requires a water-air interface and thus cannot be applied to dry conditions, atmospheric flux measurements obtained by manual closed-chambers along a representative transect (Figure 2b) were available to determine pre-rewetting GHG fluxes (CO_2 and CH_4). After rewetting, data from manual closed-chambers (transect) and from surface water sampling for the k model (transect and peatland stations) were used. The two methodologies were applied at the same locations along the transect only after rewetting (Table C1).

830 **Table C1.** Overview of the methods used to determine the atmospheric GHG fluxes

Pre-rewetting	Post-rewetting	
transect (Figure 2b)	transect (Figure 2b)	peatland area (Figure 2a)
chamber-based	chamber-based ^{1,2}	k model ²
	k model ^{1,2}	

¹ inter-methodological comparison at station BTD7
² formed the data representing post-rewetting fluxes


To evaluate the inter-comparability of the flux estimates obtained with the two methods, the results from station BTD7 were compared for each post-rewetting season (Figure C1). Data from this station were chosen because it was permanently flooded after rewetting and thus assured a valid baseline for comparison. The dynamics of the CO_2 fluxes determined by the two methods were the same and thus did not differ significantly in any of the seasons (Kruskal-Wallis test, $p > 0.05$).

CH_4 fluxes also did not differ significantly, except in autumn (Kruskal-Wallis test, $p < 0.001$), when the average flux calculated according to the two methods differed by a factor of 2.7. However, the data of the k model had less impact, due to the smaller number of measurements ($n = 6$). Given the smaller data set compared to that of the closed chambers ($n = 17$), the same statistical analysis was conducted without a seasonal division. The results showed no significant differences in the two methods for CH_4 fluxes (Kruskal-Wallis test, CO_2 and CH_4). Therefore, it was deemed appropriate to combine the flux-estimation methods for each GHG into one post-rewetting data set, as this allowed the consideration of a broader range of possible flux amplitudes. In addition, the post-rewetting data acquired along the transect were pooled with data distributed throughout the peatland area. Although the area covered by the transect was smaller than the covered by the k -model data from the peatland, such that pooling of the post-rewetting-data risked spatial bias, two positive effects of pooling were identified:

845 (1) The transect stations were representative of the entire area after flooding, because they covered a water-level-gradient (several cm to > 2 m in the ditch) that coincided with the conditions of the peatland stations. (2) The transect stations represented a large heterogeneity in the peatland before rewetting that decreased post-rewetting. This was also evident from the CO_2 flux measurements, which showed a high variability (data not shown) at each station before rewetting. After rewetting, there was less variability such that the stations became more similar in their atmospheric C-exchange patterns, likely due to

850 the mixing patterns triggered by lateral exchange with the Baltic Sea (Sect. 3.1). Largely similar conditions were therefore assumed at all stations within the peatland.

The pooled post-rewetting flux values were compared with the pre-rewetting values to investigate the direct effect of rewetting on CH₄ and CO₂ fluxes.



855 **Figure C1.** Seasonal post-rewetting fluxes of (a) CO₂ and (b) CH₄ at station BT7 which is part of the GHG flux transect. Chamber-based atmospheric GHG fluxes are shown in blue and air-sea GHG fluxes from the k model in red. The methodological comparisons within seasons are based on a significance level of $p < 0.05$. ns: not significant; *** $p < 0.001$.

Appendix D: Nutrient cross plots

860 Cross plots with linear regression analyses were generated for nutrients (NH₄⁺, NO₃⁻, NO₂⁻, PO₄³⁻) and DOC concentrations across all seasons to investigate potential correlations (Figure D1). Significant correlations are shown with red asterisks ($p < 0.05$).




Figure D1. Cross plots of the measured nutrient (NH_4^+ , NO_3^- , NO_2^- , PO_4^{3-}) and DOC concentrations in (a) the inner bay and (b) the peatland across all seasons. Significant correlations are indicated by asterisks.

865

Data availability. The raw data used in this study are archived at <http://doi.io-warnemuende.de/10.12754/data-2022-0003>. The calculated GHG emission data used in this study are archived at <http://doi.io-warnemuende.de/10.12754/data-2022-0004>.

Author contributions. All authors designed the concept of the study. DLP, AB and CNG conducted the field work, data analysis, and interpretation. DLP and AB wrote the first draft of the manuscript. DLP created the figures and organized the data. AB conducted the statistical analysis. CNG wrote sections of the manuscript. All authors contributed to manuscript revision and approved the submitted version.

875

Competing interests. The authors declare that they have no conflict of interest.

880

Acknowledgements. The authors would like to thank Cindy Hoppe and Henning Sack for their great support during the field work; Lara Prella, Petra Mutinova and the Biologische Station Zingst (all University of Rostock) for providing and measuring some additional nutrient data; Christian Burmeister, Dr. Stefan Otto (both IOW) and Dr. Stefan Köhler (University of Rostock) for their technical laboratory assistance; Dr. Joachim Dippner, Dr. Marvin Lorenz and Dr. Christiane Hassenrück (all IOW) for their help on the nutrient export calculation and statistical analyses, respectively. Dr. Bitu Sabbaghzadeh and Dr. Oliver Schmale (both IOW) provided valuable feedback on the manuscript. We are grateful to the Ostseestiftung and especially to Rasmus Klöpffer, who guided the cooperation required for the project and provided valuable data on the study area. We thank Sascha Klatt for information on the study area and especially for technical support during field work. We also thank the

Wasserstraßen- und Schifffahrtsamt Ostsee (WSA Ostsee) for water level data, the Landesamt für innere Verwaltung
885 Mecklenburg-Vorpommern (LAI V MV), Fachbereich Geodatenbereitstellung for topography data, the Landesamt für Umwelt,
Naturschutz und Geologie Mecklenburg-Vorpommern (LUNG MV), especially Mario von Weber, for nutrient monitoring data
and the DWD for meteorological data. Finally, we would like to thank our editor Carolin Löscher and the reviewers for a fair
and pleasant review process.

890 *Financial support.* This study was conducted within the framework of the Research Training Group 'Baltic TRANSCOAST'
funded by the DFG (Deutsche Forschungsgemeinschaft) under grant number GRK 2000 (www.baltic-transcoast.uni-rostock.de). This is Baltic TRANSCOAST publication no. GRK2000/0062. A. Breznikar was funded by a doctoral scholarship
from the Deutsche Bundesstiftung Umwelt (DBU).

References

895 Augustin, J. (Ed.): Gaseous emissions from constructed wetlands and (re)flooded meadows, in: International Conference:
Constructed and Riverine Wetlands for Optimal Control of Wastewater at Catchment Scale, edited by: Mander, Ü., Vohla, C.
and Poom, A., Tartu Univ. Press, 2003.

Augustin, J. and Chojnicki, B.: Austausch von klimarelevanten Spurengasen, Klimawirkung und Kohlenstoffdynamik in den
ersten Jahren nach Wiedervernässung von degradiertem Niedermoorgrünland, Berichte des Leibniz-Institut für
900 Gewässerökologie und Binnenfischerei, 50–61, 2008.

Augustin, J., Merbach, W., Steffens, L., and Snelinski, B.: Nitrous Oxide Fluxes Of Disturbed Minerotrophic Peatlands,
Agribiological research (Germany), 51, 47–57, 1998.

Bange, H. W., Bartell, U. H., Rapsomanikis, S., and Andreae, M. O.: Methane in the Baltic and North Seas and a reassessment
of the marine emissions of methane, *Global Biogeochem. Cycles*, 8, 465–480, <https://doi.org/10.1029/94GB02181>, 1994.

905 Bange, H. W., Dahlke, S., Ramesh, R., Meyer-Reil, L.-A., Rapsomanikis, S., and Andreae, M. O.: Seasonal Study of Methane
and Nitrous Oxide in the Coastal Waters of the Southern Baltic Sea, *Estuarine, Coastal and Shelf Science*, 47, 807–817,
<https://doi.org/10.1006/ecss.1998.0397>, 1998.

Bartlett, K. B., Bartlett, D. S., Harriss, R. C., and Sebacher, D. I.: Methane emissions along a salt marsh salinity gradient,
Biogeochemistry, 4, 183–202, <https://doi.org/10.1007/BF02187365>, 1987.

910 Beldowski, J., Löffler, A., Schneider, B., and Joensuu, L.: Distribution and biogeochemical control of total CO₂ and total
alkalinity in the Baltic Sea, *Journal of Marine Systems*, 81, 252–259, <https://doi.org/10.1016/j.jmarsys.2009.12.020>, 2010.

Bockholt, R.: Flächen-, Ertrags- und Problemanalyse des Überschwemmungsgrünlandes der Ostsee-, Bodden- und
Haffgewässer, Forschungsbericht Universität Rostock, 17, 1985.

- 915 Boetius, A., Ravensschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R., Jørgensen, B. B., Witte, U.,
and Pfannkuche, O.: A marine microbial consortium apparently mediating anaerobic oxidation of methane, *Nature*, 407, 623–
626, <https://doi.org/10.1038/35036572>, 2000.
- Borges, A. V., Champenois, W., Gypens, N., Delille, B., and Harlay, J.: Massive marine methane emissions from near-shore
shallow coastal areas, *Scientific reports*, 6, 27908, <https://doi.org/10.1038/srep27908>, 2016.
- 920 Borges, A. V., Speeckaert, G., Champenois, W., Scranton, M. I., and Gypens, N.: Productivity and Temperature as Drivers of
Seasonal and Spatial Variations of Dissolved Methane in the Southern Bight of the North Sea, *Ecosystems*, 21, 583–599,
<https://doi.org/10.1007/s10021-017-0171-7>, 2018.
- Brisch, A.: Erkundung von Torfmächtigkeit und Vegetation in zwei potenziellen Wiedervernässungsgebieten bei Ramin und
Grosow (Rügen), Unpublished expert opinion by Naturschutzstiftung Deutsche Ostsee, 2015.
- 925 Bubier, J., Crill, P., Mosedale, A., Frolking, S., and Linder, E.: Peatland responses to varying interannual moisture conditions
as measured by automatic CO₂ chambers, *Global Biogeochem. Cycles*, 17, 1–35, <https://doi.org/10.1029/2002GB001946>,
2003.
- Burgin, A. J. and Groffman, P. M.: Soil O₂ controls denitrification rates and N₂O yield in a riparian wetland, *J. Geophys. Res.*,
117, 1–15, <https://doi.org/10.1029/2011JG001799>, 2012.
- 930 Cabezas, A., Gelbrecht, J., Zwirnmann, E., Barth, M., and Zak, D.: Effects of degree of peat decomposition, loading rate and
temperature on dissolved nitrogen turnover in rewetted fens, *Soil Biology and Biochemistry*, 48, 182–191,
<https://doi.org/10.1016/j.soilbio.2012.01.027>, 2012.
- Capone, D. G. and Kiene, R. P.: Comparison of microbial dynamics in marine and freshwater sediments: Contrasts in anaerobic
carbon catabolism, *Limnol. Oceanogr.*, 33, 725–749, <https://doi.org/10.4319/lo.1988.33.4part2.0725>, 1988.
- 935 Carter, B. R., Radich, J. A., Doyle, H. L., and Dickson, A. G.: An automated system for spectrophotometric seawater pH
measurements, *Limnol. Oceanogr. Methods*, 11, 16–27, <https://doi.org/10.4319/lom.2013.11.16>, 2013.
- Chmura, G. L., Kellman, L., and Guntenspergen, G. R.: The greenhouse gas flux and potential global warming feedbacks of a
northern macrotidal and microtidal salt marsh, *Environ. Res. Lett.*, 6, 1–6, <https://doi.org/10.1088/1748-9326/6/4/044016>,
2011.
- 940 Chmura, G. L., Kellman, L., van Ardenne, L., and Guntenspergen, G. R.: Greenhouse Gas Fluxes from Salt Marshes Exposed
to Chronic Nutrient Enrichment, *PloS one*, 11, 1–13, <https://doi.org/10.1371/journal.pone.0149937>, 2016.
- Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Bärtsch, S., Dubovik, D., Liashchynskaya, N., Michaelis, D., Minke,
M., Skuratovich, A., and Joosten, H.: Assessing greenhouse gas emissions from peatlands using vegetation as a proxy,
Hydrobiologia, 674, 67–89, <https://doi.org/10.1007/s10750-011-0729-x>, 2011.

- 945 Danevčič, T., Mandic-Mulec, I., Stres, B., Stopar, D., and Hacin, J.: Emissions of CO₂, CH₄ and N₂O from Southern European peatlands, *Soil Biology and Biochemistry*, 42, 1437–1446, <https://doi.org/10.1016/j.soilbio.2010.05.004>, 2010.
- Dean, J. F., Middelburg, J. J., Röckmann, T., Aerts, R., Blauw, L. G., Egger, M., Jetten, M. S. M., Jong, A. E. E. de, Meisel, O. H., Rasigraf, O., Slomp, C. P., in't Zandt, M. H., and Dolman, A. J.: Methane Feedbacks to the Global Climate System in a Warmer World, *Rev. Geophys.*, 56, 207–250, <https://doi.org/10.1002/2017RG000559>, 2018.
- 950 Dickson, A. and Riley, J.: The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water — Kw, *Marine Chemistry*, 7, 89–99, [https://doi.org/10.1016/0304-4203\(79\)90001-X](https://doi.org/10.1016/0304-4203(79)90001-X), 1979.
- Dickson, A. G.: Standard potential of the reaction: AgCl(s) + 1/2H₂(g) = Ag(s) + HCl(aq), and the standard acidity constant of the ion HSO₄⁻ in synthetic sea water from 273.15 to 318.15 K, *The Journal of Chemical Thermodynamics*, 22, 113–127, [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.
- 955 Dickson, A. G., Sabine, C. L. and Christian, J. R. (Eds.): Guide to best practices for ocean CO₂ measurements, North Pacific Marine Science Organization, 2007.
- Dlugokencky, E., Crotwell, A., Mund, J., Crotwell, M., and Thoning, K.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983-2018, <https://doi.org/10.15138/wkgj-f215>, 2019b.
- 960 Dlugokencky, E., Crotwell, A., Mund, J., Crotwell, M., and Thoning, K.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983-2018, <https://doi.org/10.15138/VNCZ-M766>, 2019a.
- Duhamel, S., Nogaro, G., and Steinman, A. D.: Effects of water level fluctuation and sediment–water nutrient exchange on phosphorus biogeochemistry in two coastal wetlands, *Aquat Sci*, 79, 57–72, <https://doi.org/10.1007/s00027-016-0479-y>, 2017.
- 965 Edler, L.: Recommendations on Methods for Marine Biological Studies in the Baltic Sea: Phytoplankton and chlorophyll, *Baltic Marine Biologists*, 1979.
- Fiedler, J., Fuß, R., Glatzel, S., Hagemann, U., Huth, V., Jordan, S., Jurasinski, G., Kutzbach, L., Maier, M., Schäfer, K., Weber, T., and Weymann, D.: Best Practice Guideline Measurement of carbon dioxide, methane and nitrous oxide fluxes between soil-vegetation-systems and the atmosphere using non-steady state chambers, <https://doi.org/10.23689/figeo-5422>,
- 970 2022.
- Fisher, J. and Acreman, M. C.: Wetland nutrient removal: a review of the evidence, *Hydrol. Earth Syst. Sci.*, 8, 673–685, <https://doi.org/10.5194/hess-8-673-2004>, 2004.

- Flessa, H., Wild, U., Klemisch, M., and Pfadenhauer, J.: Nitrous oxide and methane fluxes from organic soils under agriculture, 49, 1998.
- 975 Fox, J. and Weisberg, S.: An {R} Companion to Applied Regression, Third Edition. Thousand Oaks CA: Sage, <https://socialsciences.mcmaster.ca/jfox/Books/Companion/> (last access: 03 April 2022), 2019.
- Franz, D., Koebsch, F., Larmanou, E., Augustin, J., and Sachs, T.: High net CO₂ and CH₄ release at a eutrophic shallow lake on a formerly drained fen, *Biogeosciences*, 13, 3051–3070, <https://doi.org/10.5194/bg-13-3051-2016>, 2016.
- Gattuso, J.-P., Epitalon, J.-M., Lavigne, H., and Orr, J.: seacarb: Seawater Carbonate Chemistry, <https://CRAN.R-project.org/>
 980 <https://CRAN.R-project.org/package=seacarb> (last access: 06 February 2022), R package version 3.2.15, 2019.
- Geurts, J. J. M., Smolders, A. J. P., Banach, A. M., van de Graaf, J. P. M., Roelofs, J. G. M., and Lamers, L. P. M.: The interaction between decomposition, net N and P mineralization and their mobilization to the surface water in fens, *Water research*, 44, 3487–3495, <https://doi.org/10.1016/j.watres.2010.03.030>, 2010.
- Glatzel, S., Forbrich, I., Krüger, C., Lemke, S., and Gerold, G.: Environmental controls of greenhouse gas release in a restoring
 985 peat bog in NW Germany, *Biogeosciences Discussions, European Geosciences*, 213–242, <https://doi.org/10.5194/bgd-5-213-2008>, 2008.
- Glatzel, S. and Stahr, K.: Methane and nitrous oxide exchange in differently fertilised grassland in southern Germany, *Plant and Soil*, 231, 21–35, 2001.
- Goldberg, S. D., Knorr, K.-H., Blodau, C., Lischeid, G., and Gebauer, G.: Impact of altering the water table height of an acidic
 990 fen on N₂O and NO fluxes and soil concentrations, *Global change biology*, 16, 220–233, <https://doi.org/10.1111/j.1365-2486.2009.02015.x>, 2010.
- Grasshoff, K., Kremling, K., and Ehrhardt, M.: *Methods of Seawater Analysis*, 2009.
- Grolemund, G. and Wickham, H.: Dates and Times Made Easy with lubridate. *Journal of Statistical Software*, <https://www.jstatsoft.org/v40/i03/> (last access: 03 April 2022), 2011.
- 995 Hahn, J., Köhler, S., Glatzel, S., and Jurasinski, G.: Methane Exchange in a Coastal Fen in the First Year after Flooding-A Systems Shift, *PloS one*, 10, 1-25, <https://doi.org/10.1371/journal.pone.0140657>, 2015.
- Hahn-Schöfl, M., Zak, D., Minke, M., Gelbrecht, J., Augustin, J., and Freibauer, A.: Organic sediment formed during inundation of a degraded fen grassland emits large fluxes of CH₄ and CO₂, *Biogeosciences*, 8, 1539–1550, <https://doi.org/10.5194/bg-8-1539-2011>, 2011.
- 1000 Harpenslager, S. F., van den Elzen, E., Kox, M. A., Smolders, A. J., Ettwig, K. F., and Lamers, L. P.: Rewetting former agricultural peatlands: Topsoil removal as a prerequisite to avoid strong nutrient and greenhouse gas emissions, *Ecological Engineering*, 84, 159–168, <https://doi.org/10.1016/j.ecoleng.2015.08.002>, 2015.

HELCOM: HELCOM Guidelines for the annual and periodical compilation and reporting of waterborne pollution inputs to the Baltic Sea (PLC-Water), http://nest.su.se/helcom_plc/ (last access 17 December 2021), HELCOM, 2019.

1005 Heyer, J. and Berger, U.: Methane Emission from the Coastal Area in the Southern Baltic Sea, *Estuarine, Coastal and Shelf Science*, 51, 13–30, <https://doi.org/10.1006/ecss.2000.0616>, 2000.

Hogan, D. M., Jordan, T. E., and Walbridge, M. R.: Phosphorus retention and soil organic carbon in restored and natural freshwater wetlands, *Wetlands*, 24, 573–585, [https://doi.org/10.1672/0277-5212\(2004\)024\[0573:PRASOC\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2004)024[0573:PRASOC]2.0.CO;2), 2004.

1010 Holz, R., Herrmann, C., and Müller-Motzfeld, G.: Vom Polder zum Ausdeichungsgebiet: Das Projekt Karrendorfer Wiesen und die Zukunft der Küstenüberflutungsgebiete in Mecklenburg-Vorpommern, *Natur und Naturschutz in MV*, Schriftenreihe des Institutes für Landschaftsökologie und Naturschutz Greifswald, Band 32, 1996.

Joosten, H. and Clarke, D.: *Wise use of mires and peatlands, Background and principles including a framework for decision-making*, 2002.

1015 Jørgensen, B. B. (Ed.): *Bacteria and Marine Biogeochemistry*, in: *Marine Geochemistry*, https://doi.org/10.1007/3-540-32144-6_5, 2006.

Jørgensen, C. J. and Elberling, B.: Effects of flooding-induced N₂O production, consumption and emission dynamics on the annual N₂O emission budget in wetland soil, *Soil Biology and Biochemistry*, 53, 9–17, <https://doi.org/10.1016/j.soilbio.2012.05.005>, 2012.

1020 Jurasinski, G., Günther, A. B., Huth, V., Couwenberg, J., and Glatzel, S.: Paludiculture – productive use of wet peatlands., *Ecosystem services provided by paludiculture – Greenhouse gas emissions*, 79–94, 2016.

Jurasinski, G., Janssen, M., Voss, M., Böttcher, M. E., Brede, M., Burchard, H., Forster, S., Gosch, L., Gräwe, U., Gründling-Pfaff, S., Haider, F., Ibenthal, M., Karow, N., Karsten, U., Kreuzburg, M., Lange, X., Leinweber, P., Massmann, G., Ptak, T., Rezanezhad, F., Rehder, G., Romoth, K., Schade, H., Schubert, H., Schulz-Vogt, H., Sokolova, I. M., Strehse, R., Unger, V., Westphal, J., and Lennartz, B.: Understanding the Coastal Ecocline: Assessing Sea–Land Interactions at Non-tidal, Low-Lying
1025 Coasts Through Interdisciplinary Research, *Front. Mar. Sci.*, 5, <https://doi.org/10.3389/fmars.2018.00342>, 2018.

Jurasinski, G., Koebsch, F., Guenther, A., and Beetz, S.: Flux rate calculation from dynamic closed chamber measurements., <https://CRAN.R-project.org/package=flux> (last access: 06 February 2022), R package version 0.3-0, 2014.

Kaat, A. and Joosten, H.: *Factbook for UNFCCC policies on peat carbon emissions*, 2009.

1030 Kandel, T. P., Lærke, P. E., Hoffmann, C. C., and Elsgaard, L.: Complete annual CO₂, CH₄, and N₂O balance of a temperate riparian wetland 12 years after rewetting, *Ecological Engineering*, 127, 527–535, <https://doi.org/10.1016/j.ecoleng.2017.12.019>, 2019.

- Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process, *Annual review of microbiology*, 63, 311–334, <https://doi.org/10.1146/annurev.micro.61.080706.093130>, 2009.
- 1035 Koebsch, F., Glatzel, S., Hofmann, J., Forbrich, I., and Jurasinski, G.: CO₂ exchange of a temperate fen during the conversion from moderately rewetting to flooding, *J. Geophys. Res. Biogeosci.*, 118, 940–950, <https://doi.org/10.1002/jgrg.20069>, 2013.
- Koebsch, F., Gottschalk, P., Beyer, F., Wille, C., Jurasinski, G., and Sachs, T.: The impact of occasional drought periods on vegetation spread and greenhouse gas exchange in rewetted fens, *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 375, 20190685, <https://doi.org/10.1098/rstb.2019.0685>, 2020.
- 1040 Koebsch, F., Jurasinski, G., Koch, M., Hofmann, J., and Glatzel, S.: Controls for multi-scale temporal variation in ecosystem methane exchange during the growing season of a permanently inundated fen, *Agricultural and Forest Meteorology*, 204, 94–105, <https://doi.org/10.1016/j.agrformet.2015.02.002>, 2015.
- Koebsch, F., Winkel, M., Liebner, S., Liu, B., Westphal, J., Schmiedinger, I., Spitzzy, A., Gehre, M., Jurasinski, G., Köhler, S., Unger, V., Koch, M., Sachs, T., and Böttcher, M. E.: Sulfate deprivation triggers high methane production in a disturbed and rewetted coastal peatland, *Biogeosciences*, 16, 1937–1953, <https://doi.org/10.5194/bg-16-1937-2019>, 2019.
- 1045 Kool, D. M., Dolfing, J., Wrage, N., and van Groenigen, J. W.: Nitrifier denitrification as a distinct and significant source of nitrous oxide from soil, *Soil Biology and Biochemistry*, 43, 174–178, <https://doi.org/10.1016/j.soilbio.2010.09.030>, 2011.
- Kreyling, J., Tanneberger, F., Jansen, F., van der Linden, S., Aggenbach, C., Blüml, V., Couwenberg, J., Emsens, W.-J., Joosten, H., Klimkowska, A., Kotowski, W., Kozub, L., Lennartz, B., Liczner, Y., Liu, H., Michaelis, D., Oehmke, C., Parakenings, K., Pleyl, E., Poyda, A., Raabe, S., Röhl, M., Rücker, K., Schneider, A., Schrautzer, J., Schröder, C., Schug, F., 1050 Seeber, E., Thiel, F., Thiele, S., Tiemeyer, B., Timmermann, T., Urich, T., van Diggelen, R., Vegelin, K., Verbruggen, E., Wilming, M., Wrage-Mönnig, N., Wołajko, L., Zak, D., and Jurasinski, G.: Rewetting does not return drained fen peatlands to their old selves, *Nature communications*, 12, 5693, <https://doi.org/10.1038/s41467-021-25619-y>, 2021.
- Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C., Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm, A., Slomp, C. P., Sofiev, 1055 M., Sobek, A., Szymczycha, B., and Undeman, E.: Biogeochemical functioning of the Baltic Sea, *Earth Syst. Dynam.*, 13, 633–685, <https://doi.org/10.5194/esd-13-633-2022>, 2022.
- Kuliński, K., Schneider, B., Hammer, K., Machulik, U., and Schulz-Bull, D.: The influence of dissolved organic matter on the acid–base system of the Baltic Sea, *Journal of Marine Systems*, 132, 106–115, <https://doi.org/10.1016/j.jmarsys.2014.01.011>, 2014.
- 1060 Kuliński, K., Schneider, B., Szymczycha, B., and Stokowski, M.: Structure and functioning of the acid–base system in the Baltic Sea, *Earth Syst. Dynam.*, 8, 1107–1120, <https://doi.org/10.5194/esd-8-1107-2017>, 2017.

- Lamers, L. P., Smolders, A. J., and Roelofs, J. G.: The restoration of fens in the Netherlands, *Hydrobiologia*, 478, 107–130, <https://doi.org/10.1023/A:1021022529475>, 2002.
- 1065 Lennartz, B. and Liu, H.: Hydraulic Functions of Peat Soils and Ecosystem Service, *Front. Environ. Sci.*, 7, <https://doi.org/10.3389/fenvs.2019.00092>, 2019.
- Leppelt, T., Dechow, R., Gebbert, S., Freibauer, A., Lohila, A., Augustin, J., Drösler, M., Fiedler, S., Glatzel, S., Höper, H., Järveoja, J., Lærke, P. E., Maljanen, M., Mander, Ü., Mäkiranta, P., Minkkinen, K., Ojanen, P., Regina, K., and Strömgren, M.: Nitrous oxide emission budgets and land-use-driven hotspots for organic soils in Europe, *Biogeosciences*, 11, 6595–6612, <https://doi.org/10.5194/bg-11-6595-2014>, 2014.
- 1070 Liu, H. and Lennartz, B.: Short Term Effects of Salinization on Compound Release from Drained and Restored Coastal Wetlands, *Water*, 11, 1549, <https://doi.org/10.3390/w11081549>, 2019.
- Liu, H., Zak, D., Rezanezhad, F., and Lennartz, B.: Soil degradation determines release of nitrous oxide and dissolved organic carbon from peatlands, *Environ. Res. Lett.*, 14, 94009, <https://doi.org/10.1088/1748-9326/ab3947>, 2019.
- Livingston, G. P. and Hutchinson, G.: Enclosure-based measurement of trace gas exchange: applications and sources of error., 1075 In: Matson, P.A. and Harris, R.C., Eds., *Biogenic trace gases: measuring emissions from soil and water.*, Blackwell Science Ltd., Oxford, UK., 14–51, 1995.
- Löffler, A., Schneider, B., Perttilä, M., and Rehder, G.: Air–sea CO₂ exchange in the Gulf of Bothnia, Baltic Sea, *Continental Shelf Research*, 37, 46–56, <https://doi.org/10.1016/j.csr.2012.02.002>, 2012.
- Martikainen, P. J., Nykänen, H., Crill, P., and Silvola, J.: Effect of a lowered water table on nitrous oxide fluxes from northern 1080 peatlands, *Nature*, 366, 51–53, <https://doi.org/10.1038/366051a0>, 1993.
- Millero, F. J.: Carbonate constants for estuarine waters, *Mar. Freshwater Res.*, 61, 139, <https://doi.org/10.1071/MF09254>, 2010.
- Minkkinen, K., Ojanen, P., Koskinen, M., and Penttilä, T.: Nitrous oxide emissions of undrained, forestry-drained, and rewetted boreal peatlands, *Forest Ecology and Management*, 478, 118494, <https://doi.org/10.1016/j.foreco.2020.118494>, 2020.
- 1085 Moore, T. R., Roulet, N. T., and Waddington, J. M.: Uncertainty in Predicting the Effect of Climatic Change on the Carbon Cycling of Canadian Peatlands, *Climatic Change*, 40, 229–245, <https://doi.org/10.1023/A:1005408719297>, 1998.
- Moseman-Valtierra, S., Gonzalez, R., Kroeger, K. D., Tang, J., Chao, W. C., Crusius, J., Bratton, J., Green, A., and Shelton, J.: Short-term nitrogen additions can shift a coastal wetland from a sink to a source of N₂O, *Atmospheric Environment*, 45, 4390–4397, <https://doi.org/10.1016/j.atmosenv.2011.05.046>, 2011.

- 1090 Müller, J. D., Bastkowski, F., Sander, B., Seitz, S., Turner, D. R., Dickson, A. G., and Rehder, G.: Metrology for pH Measurements in Brackish Waters—Part 1: Extending Electrochemical pH Measurements of TRIS Buffers to Salinities 5–20, *Front. Mar. Sci.*, 5, <https://doi.org/10.3389/fmars.2018.00176>, 2018.
- Müller, J. D. and Rehder, G.: Metrology of pH Measurements in Brackish Waters—Part 2: Experimental Characterization of Purified meta-Cresol Purple for Spectrophotometric pH Measurements, *Front. Mar. Sci.*, 5, 1095 <https://doi.org/10.3389/fmars.2018.00177>, 2018.
- Müller, J. D., Schneider, B., and Rehder, G.: Long-term alkalinity trends in the Baltic Sea and their implications for CO₂-induced acidification, *Limnol. Oceanogr.*, 61, 1984–2002, <https://doi.org/10.1002/lno.10349>, 2016.
- Neubauer, S. C., Franklin, R. B., and Berrier, D. J.: Saltwater intrusion into tidal freshwater marshes alters the biogeochemical processing of organic carbon, *Biogeosciences*, 10, 8171–8183, <https://doi.org/10.5194/bg-10-8171-2013>, 2013.
- 1100 Oertel, C., Matschullat, J., Zurba, K., Zimmermann, F., and Erasmi, S.: Greenhouse gas emissions from soils—A review, *Geochemistry*, 76, 327–352, <https://doi.org/10.1016/j.chemer.2016.04.002>, 2016.
- Oremland, R. S. (Ed.): The biogeochemistry of methanogenic bacteria, in: The biology of microorganisms. Available online at <http://pubs.er.usgs.gov/publication/70198767>, 1988.
- Parish, F.: Assessment on peatlands, biodiversity and climate change, Main report, 2008.
- 1105 Pedersen, T. L.: patchwork: The Composer of Plots. R package version 1.1.1, <https://CRAN.R-project.org/package=patchwork>, 2020.
- Petersen, S. O., Hoffmann, C. C., Schäfer, C.-M., Blicher-Mathiesen, G., Elsgaard, L., Kristensen, K., Larsen, S. E., Torp, S. B., and Greve, M. H.: Annual emissions of CH₄ and N₂O, and ecosystem respiration, from eight organic soils in Western Denmark managed by agriculture, *Biogeosciences*, 9, 403–422, <https://doi.org/10.5194/bg-9-403-2012>, 2012.
- 1110 Pönisch, D. L.: Methodenentwicklung und –anwendung zur Analytik von Methan und Lachgas in Seewasser, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Master thesis, 2018.
- Rassamee, V., Sattayatewa, C., Pagilla, K., and Chandran, K.: Effect of oxic and anoxic conditions on nitrous oxide emissions from nitrification and denitrification processes, *Biotechnology and bioengineering*, 108, 2036–2045, <https://doi.org/10.1002/bit.23147>, 2011.
- 1115 Reeburgh, W. S.: Oceanic methane biogeochemistry, *Chemical reviews*, 107, 486–513, <https://doi.org/10.1021/cr050362v>, 2007.
- Regina, K., Nykänen, H., Silvola, J., and Martikainen, P. J.: Fluxes of nitrous oxide from boreal peatlands as affected by peatland type, water table level and nitrification capacity, *Biogeochemistry*, 35, 401–418, <https://doi.org/10.1007/BF02183033>, 1996.

- 1120 Regina, K., Silvola, J., and Martikainen, P. J.: Short-term effects of changing water table on N₂O fluxes from peat monoliths from natural and drained boreal peatlands, *Global change biology*, 5, 183–189, <https://doi.org/10.1046/j.1365-2486.1999.00217.x>, 1999.
- Richert, M., Dietrich, O., Koppisch, D., and Roth, S.: The Influence of Rewetting on Vegetation Development and Decomposition in a Degraded Fen, *Restor Ecology*, 8, 186–195, <https://doi.org/10.1046/j.1526-100x.2000.80026.x>, 2000.
- 1125 Roughan, B. L., Kellman, L., Smith, E., and Chmura, G. L.: Nitrous oxide emissions could reduce the blue carbon value of marshes on eutrophic estuaries, *Environ. Res. Lett.*, 13, 44034, <https://doi.org/10.1088/1748-9326/aab63c>, 2018.
- Rysgaard, S., Thastum, P., Dalsgaard, T., Christensen, P. B., Sloth, N. P., and Rysgaard, S.: Effects of Salinity on NH₄⁺ Adsorption Capacity, Nitrification, and Denitrification in Danish Estuarine Sediments, *Estuaries*, 22, 21, <https://doi.org/10.2307/1352923>, 1999.
- 1130 Sabbaghzadeh, B., Arévalo-Martínez, D. L., Glockzin, M., Otto, S., and Rehder, G.: Meridional and Cross-Shelf Variability of N₂O and CH₄ in the Eastern-South Atlantic, *J. Geophys. Res. Oceans*, 126, <https://doi.org/10.1029/2020JC016878>, 2021.
- Schneider, B. and Müller, J. D.: Biogeochemical Transformations in the Baltic Sea, <https://doi.org/10.1007/978-3-319-61699-5>, 2018.
- Schönheit, P., Kristjansson, J. K., and Thauer, R. K.: Kinetic mechanism for the ability of sulfate reducers to out-compete
1135 methanogens for acetate, *Archives of microbiology*, 132, 285–288, 1982.
- Segarra, K. E., Comerford, C., Slaughter, J., and Joye, S. B.: Impact of electron acceptor availability on the anaerobic oxidation of methane in coastal freshwater and brackish wetland sediments, *Geochimica et Cosmochimica Acta*, 115, 15–30, <https://doi.org/10.1016/j.gca.2013.03.029>, 2013.
- Segers, R. and Kengen, S.: Methane production as a function of anaerobic carbon mineralization: A process model, *Soil
1140 Biology and Biochemistry*, 30, 1107–1117, [https://doi.org/10.1016/S0038-0717\(97\)00198-3](https://doi.org/10.1016/S0038-0717(97)00198-3), 1998.
- Seifert, T., Tauber, F., and Kayser, B.: A high resolution spherical grid topography of the Baltic Sea – 2nd edition, *Baltic Sea Science Congress, Stockholm 25-29. November 2001, Poster #147*, 2001.
- Smolders, A. J. P., Lamers, L. P. M., Lucassen, E. C. H. E. T., van der Velde, G., and Roelofs, J. G. M.: Internal eutrophication: How it works and what to do about it—a review, *Chemistry and Ecology*, 22, 93–111,
1145 <https://doi.org/10.1080/02757540600579730>, 2006.
- Steinle, L., Maltby, J., Treude, T., Kock, A., Bange, H. W., Engbersen, N., Zopfi, J., Lehmann, M. F., and Niemann, H.: Effects of low oxygen concentrations on aerobic methane oxidation in seasonally hypoxic coastal waters, *Biogeosciences*, 14, 1631–1645, <https://doi.org/10.5194/bg-14-1631-2017>, 2017.

- Steinmuller, H. E. and Chambers, L. G.: Can Saltwater Intrusion Accelerate Nutrient Export from Freshwater Wetland Soils? An Experimental Approach, *Soil Sci. Soc. Am. j.*, 82, 283–292, <https://doi.org/10.2136/sssaj2017.05.0162>, 2018.
- Strack, M. (Ed.): *Peatlands and climate change*, Internat. Peat Soc., 2008.
- Succow, M. and Joosten, H. (Eds.): *Landschaftsökologische Moorkunde*, E. Schweizerbart'sche Verlagsbuchhandlung (Nägele u. Obermiller), 2001.
- Thomas, H. and Schneider, B.: The seasonal cycle of carbon dioxide in Baltic Sea surface waters, *Journal of Marine Systems*, 22, 53–67, [https://doi.org/10.1016/S0924-7963\(99\)00030-5](https://doi.org/10.1016/S0924-7963(99)00030-5), 1999.
- Treude, T., Krüger, M., Boetius, A., and Jørgensen, B. B.: Environmental control on anaerobic oxidation of methane in the gassy sediments of Eckernförde Bay (German Baltic), *Limnol. Oceanogr.*, 50, 1771–1786, <https://doi.org/10.4319/lo.2005.50.6.1771>, 2005.
- van de Riet, B. P., Hefting, M. M., and Verhoeven, J. T. A.: Rewetting Drained Peat Meadows: Risks and Benefits in Terms of Nutrient Release and Greenhouse Gas Exchange, *Water Air Soil Pollut. (Water, Air, & Soil Pollution)*, 224, <https://doi.org/10.1007/s11270-013-1440-5>, 2013.
- Voss, M., Deutsch, B., Liskow, I., Pastuszak, M., Schulte, U., and Sitek, S.: Nitrogen retention in the Szczecin Lagoon, Baltic Sea, *Isotopes in environmental and health studies*, 46, 355–369, <https://doi.org/10.1080/10256016.2010.503895>, 2010.
- Wang, M., Liu, H., and Lennartz, B.: Microtopography effects on carbon accumulation and nutrient release from rewetted coastal wetlands., AGU2021 Fall Meeting, 2021.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr. Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- Weiss, R. F. and Price, B. A.: Nitrous oxide solubility in water and seawater, *Marine Chemistry*, 8, 347–359, [https://doi.org/10.1016/0304-4203\(80\)90024-9](https://doi.org/10.1016/0304-4203(80)90024-9), 1980.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H.: Welcome to the tidyverse. *Journal of Open Source Software*, <https://doi.org/10.21105/joss.01686>, 2019.
- Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes, *Marine Chemistry*, 106, 287–300, <https://doi.org/10.1016/j.marchem.2007.01.006>, 2007.
- Zak, D. and Gelbrecht, J.: The mobilisation of phosphorus, organic carbon and ammonium in the initial stage of fen rewetting (a case study from NE Germany), *Biogeochemistry*, 85, 141–151, <https://doi.org/10.1007/s10533-007-9122-2>, 2007.

1180 Zak, D., Meyer, N., Cabezas, A., Gelbrecht, J., Mauersberger, R., Tiemeyer, B., Wagner, C., and McInnes, R.: Topsoil removal to minimize internal eutrophication in rewetted peatlands and to protect downstream systems against phosphorus pollution: A case study from NE Germany, *Ecological Engineering*, 103, 488–496, <https://doi.org/10.1016/j.ecoleng.2015.12.030>, 2017.

Zielinski, T., Sagan, I. and Surosz, W. (Eds.): *Interdisciplinary Approaches for Sustainable Development Goals*, Springer International Publishing, <https://doi.org/10.1007/978-3-319-71788-3>, 2018.