

# Ideas and perspectives: Land-ocean connectivity through groundwater

Damian L. Arévalo-Martínez <sup>1,2,3\*</sup> Amir Haroon <sup>1,\*</sup>, Hermann W. Bange <sup>1</sup>, Ercan Erkul <sup>2</sup>, Marion Jegen <sup>1</sup>,  
5 Nils Moosdorf <sup>2,4</sup>, Jens Schneider von Deimling <sup>2</sup>, Christian Berndt <sup>1</sup>, Michael Ernst Böttcher <sup>5,6,7</sup>, Jasper  
Hoffmann <sup>8</sup>, Volker Liebetrau <sup>1,†</sup>, Ulf Mallast <sup>9</sup>, Gudrun Massmann <sup>10</sup>, Aaron Micallef <sup>1,11</sup>, Holly A.  
Michael <sup>12</sup>, Hendrik Paasche <sup>9</sup>, Wolfgang Rabbel <sup>2</sup>, Isaac Santos <sup>13</sup>, Jan Scholten <sup>2</sup>, Katrin Schwalenberg  
14, Beata Szymczyscha <sup>15</sup>, Ariel T. Thomas <sup>11</sup>, Joonas J. Virtasalo <sup>16</sup>, Hannelore Waska <sup>10</sup>, Bradley Weymer  
17.

10 <sup>1</sup> GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, 24105, Germany

<sup>2</sup> Kiel University, Kiel, 24118, Germany

<sup>3</sup> Now at: Radboud University, Nijmegen, 6525 AJ, The Netherlands

<sup>4</sup> Leibniz Centre for Tropical Marine Research (ZMT), Bremen, 28359, Germany

<sup>5</sup> Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Rostock, 18119, Germany

<sup>6</sup> Marine Geochemistry, University of Greifswald, Greifswald, 17489, Germany

<sup>7</sup> Interdisciplinary Faculty, University of Rostock, Rostock, 18051, Germany

<sup>8</sup> Alfred-Wegener-Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, 27515, Germany

<sup>9</sup> Helmholtz Centre for Environmental Research, Leipzig, 04318, Germany

<sup>10</sup> Carl von Ossietzky University of Oldenburg, Oldenburg, 26129, Germany

<sup>11</sup> University of Malta, Msida, MSD 2080, Malta

<sup>12</sup> University of Delaware, Newark, DE 19716, USA

<sup>13</sup> Department of Marine Science, University of Gothenburg, Gothenburg, 40539, Sweden

<sup>14</sup> Federal Institute for Geosciences and Natural Resources, Hannover, 30655, Germany

<sup>15</sup> Institute of Gdańsk Polish Academy of Sciences, Sopot, 81-712, Poland

<sup>16</sup> Marine Geology, Geological Survey of Finland (GTK), Espoo, 02150, Finland

<sup>17</sup> School of Oceanography, Shanghai Jiao Tong University, Shanghai, China

† deceased

30 \*Correspondence to:

Damian L. Arévalo-Martínez ([darevalo@geomar.de](mailto:darevalo@geomar.de)) or Amir Haroon ([aharoon@geomar.de](mailto:aharoon@geomar.de))

**Abstract.** For millennia humans have gravitated towards coastlines for their resource potential and as geopolitical centres for global trade. A basic requirement ensuring water security for coastal communities relies on a delicate balance between the supply and demand of potable water. The interaction between 35 freshwater and saltwater in coastal settings is, therefore, complicated by both natural and human-driven environmental changes at the land-sea interface. In particular, ongoing sea level rise, warming and deoxygenation might exacerbate such perturbations. In this context, an improved understanding of the

40 nature and variability of groundwater fluxes across the land-sea continuum is timely, yet remains out of reach. The flow of terrestrial groundwater across the coastal transition zone as well as the extent of  
freshened groundwater below the present-day seafloor are receiving increased attention in marine and  
coastal sciences because they likely represent a significant, yet highly uncertain component of  
(bio)geochemical budgets, and because of the emerging interest in the potential use of offshore freshened  
groundwater as a resource. At the same time, “reverse” groundwater flux from offshore to onshore is of  
45 prevalent socio-economic interest as terrestrial groundwater resources are continuously pressured by  
overpumping and seawater intrusion in many coastal regions worldwide. An accurate assessment of the  
land-ocean connectivity through groundwater and its potential responses to future anthropogenic  
activities and climate change will require a multidisciplinary approach combining the expertise of  
geophysicists, hydrogeologists, (bio)geochemists and modellers. Such joint activities will lay the  
50 scientific basis for better understanding the role of groundwater in societal-relevant issues such as climate  
change, pollution and the environmental status of the coastal oceans within the framework of the United  
Nations Sustainable Development Goals. Here, we present our perspectives on future research directions  
to better understand land-ocean connectivity through groundwater, including the spatial distributions of  
the essential hydrogeological parameters, highlighting technical and scientific developments, and briefly  
discussing its societal relevance in rapidly changing coastal oceans.

## 55 1 Background

The exchange of groundwater between land and ocean is a wide-spread phenomenon, which has significant impacts on the biogeochemical cycles of the coastal ocean (e.g. Church, 1996; Moore, 2010; Santos et al., 2021). Coastal margins play a disproportionately important role for marine ecosystems compared to the open ocean due to their greater biological productivity, sediment-water interactions and  
60 air-sea transfer of climate-relevant trace gases (Liu et al., 2010). Increasing anthropogenic activities result in high nutrient fluxes into the coastal ocean, leading to eutrophication, deoxygenation and release of greenhouse gases, which in turn could exacerbate the current global warming trend and significantly affect the livelihood of nations that rely on coastal ecosystem services (e.g. Van Meter et al., 2018; Oehler et al., 2021; Rocha et al., 2021). In addition, accelerating global sea level rise (GSLR) can negatively  
65 influence terrestrial coastal aquifers due to the inland displacement of the fresh-saline-water interfaces, referred to as saltwater intrusion (SWI; Ferguson and Gleeson, 2012; Taylor et al., 2013). Sea level rise at regional and local scales can modify the chemical balance within subterranean estuaries and increase their spatial extent by enhancing SWI, which results in enhanced load of e.g. nutrients and pollutants to  
70 the coastal ocean (Moore and Joye, 2021). In turn, increased human usage of groundwater resources is estimated to account for approximately 14% of the observed GSLR through a net transfer of freshwater from deep reservoirs into the ocean (Konikow, 2011; Church et al., 2013; Taylor et al., 2013). Increasing usage of non-renewable groundwater might further exacerbate global water depletion (Bierkens and

Wada, 2019), which is further impacted by climate variability through changes in recharge and precipitation (Thomas and Famiglietti, 2019; Beebe et al., 2022).

75

The cross-shelf extension of terrestrial coastal groundwater systems can be distinguished into two key (often interrelated) elements (see Table 1 and Fig. 1). The first comprises meteoric groundwater flux from terrestrial aquifers through the seabed (including the intertidal zone) into the coastal ocean, which is generally referred to as fresh submarine groundwater discharge (FSGD; e.g. Kohout, 1964; Taniguchi et al., 2019). The second consists of large ( $> 10$  km horizontal extent) freshened (and often brackish) groundwater reservoirs embedded in sediment and rocks below the present-day seafloor, collectively called offshore freshened groundwater (OFG; Post et al., 2013).

FSGD connects terrestrial groundwater systems to the coastal ocean on most coastlines in the world (Fig. 85 2; Luijendijk et al., 2020). FSGD is essentially the surplus of the terrestrial water budget. Most known FSGD occurs within the first few 100 meters from the coast, although its occurrence has also been reported at tens to hundreds of kilometres offshore (Manheim, 1967; Kooi et al., 2001; Bratton et al., 2010). Given the large degree of spatio-temporal variability in FSGD, estimates of regional and global fluxes are still highly uncertain (Taniguchi et al., 2019). Globally, FSGD accounts for 1–10 % of the 90 global freshwater input to the ocean (Abbott et al., 2019; Luijendijk et al., 2020). Locally, however, FSGD can be key for sustaining some marine ecosystems (Luijendijk et al., 2020).

Similar to FSGD, OFG has been observed in shelf sediments throughout the world's oceans (Fig. 2; Post et al., 2013; Micallef et al., 2021). Likewise, OFG is a potential freshwater resource, or a resource of 95 water that can be treated with desalination with comparably small energy consumption (Bakken et al., 2012), and has therefore gained increased attention over the past decade (Post et al., 2013; Micallef et al., 2021). Although OFG is generally a relic of past sea-level low stand (fossil groundwater), some reservoirs are likely hydraulically connected to the terrestrial aquifers groundwater system, as shown for the U.S. 100 Atlantic coast (Gustafson et al., 2019; Thomas et al., 2019), Canterbury Bight, New Zealand (Micallef et al., 2020; Weymer et al., 2020), and the Achziv submarine canyon in northern Israel (Paldor et al., 2020). Here, we emphasize the importance of improving our understanding of connected OFG, since its extraction as an unconventional resource for mitigating temporal water scarcity in coastal communities might cause seawater intrusion (Yu and Michael, 2019a) and distant land subsidence (Chen et al., 2007; Yu and Michael, 2019b).

105

With ongoing research in near-coastal groundwater fluxes (FSGD) and offshore reservoirs (OFG) carried out by largely different scientific communities, we address unexploited scientific and technical synergies between them. The reliance on markedly different methodologies leads to differences in scientific language, and in turn conceptually disconnects the research of both phenomena (FSGD is usually assessed 110 using geochemical tracers and hydrological observations from the intertidal zone or numerical groundwater modelling (Taniguchi et al., 2019; Luijendijk et al., 2020), whereas OFG studies often require ship-based geophysical methods (Micallef et al., 2021)). Here we present new perspectives on future research directions to improve the understanding of land-ocean connectivity through groundwater, with particular focus on joint activities of FSGD and OFG research communities. This includes *i*)

115 improving our quantitative understanding of the distribution and variability of groundwater fluxes at regional and global scales, *ii*) assessing long-term changes in groundwater sources and their expected impact on marine environments, as well as potential usage, and *iii*) evaluating conceptual and technological developments which will potentially advance joint FSGD-OFG research.

## 120 **2 Distribution and variability of groundwater fluxes**

### Fresh submarine groundwater discharge

125 The interaction between saline and fresh groundwater in coastal regions is governed by complex processes, e.g. density contrasts between fresh and saline water, tidal effects, and geological heterogeneity (Michael et al., 2016; Jiao and Post, 2019). Saline groundwater can intrude landward salinizing terrestrial aquifers (resulting in SWI). Yet, at the same time, terrestrial groundwater can cross the land-sea continuum and appear offshore as FSGD and/or OFG (Fig. 2; see e.g. Whiticar, 2002; Post et al. 2013; Jurasinski et al., 2018; Micallef et al. 2020). Groundwater flow is associated with external forcing (e.g. groundwater heads, framework geology, onshore groundwater usage, sea level) that dictates the 130 hydrostatic gradient causing fluxes to be directed inland, offshore or both. Strong distortions of hydraulic gradients can influence or even reverse groundwater flow, which, in turn, might have widespread consequences for pelagic and benthic marine ecosystems (e.g. Donis et al., 2017; Lecher and Mackey, 2018; Szymczyska et al., 2020; Santos et al., 2021), as well as for associated services such as fisheries, because both nutrients and contaminants are transported into the coastal ocean via groundwater. A recent 135 study estimated the global input of groundwater into the ocean via FSGD to be less than 1% of the surface-water runoff. However, on local scales FSGD can reach 25% of the river flux (Luijendijk et al., 2020), and saline SGD releases recycled nutrients at rates comparable to global rivers (Santos et al., 2021). The high spatial variability of this influx is partly controlled by climate at regional scale, and partly by lithological heterogeneities at local scale (Sawyer et al., 2016). Because the extrapolation of point-scale 140 measurements onto a regional, continental, or global scale is difficult, FSGD quantification heavily relies on hydrogeological modelling (Moosdorf et al., 2021), which can result in great uncertainties on large spatial scales.

### Offshore freshened groundwater

145 OFG resides beneath the seafloor along continental shelves and, in contrast to FSGD, is commonly assumed to **have lower groundwater flow velocities** (e.g. Micallef et al. 2020). Recent estimates report OFG to comprise a volume of approximately  $1*10^6$  km<sup>3</sup> (Micallef et al., 2021), which is about 10% of the Earth's liquid fresh water (Shlklomanov, 1993). Different OFG emplacement mechanisms have been proposed, from which meteoric recharge, sub- and proglacial injection, diagenesis and the decomposition 150 of gas hydrates are the most significant (Micallef et al., 2021). OFG systems may be coupled with FSGD (e.g. Paldor et al., 2020; Attias et al., 2021), and modelling shows that FSGD and OFG can occur in equilibrium with present-day sea level for a range of different stratigraphic configurations (Michael et al., 2016). However, OFG can also be decoupled from interaction with the water column (see e.g. Micallef et al., 2020). Post et al. (2013) compiled a global estimation of OFG sites based mainly on borehole 155 observations. Geophysical technologies have updated these global estimates through the detection of OFG residing within siliciclastic continental margins in the United States and New Zealand (Gustafson et al.,

2019; Micallef et al., 2020), along a carbonate coastline in Malta (Haroon et al., 2021), and offshore from the volcanic islands of Hawaii (Attias et al., 2021). These studies have improved our understanding of spatial OFG distributions, but do not bridge the knowledge gap between coastal nearshore and offshore hydrological systems. To date, continuous tracing of terrestrial aquifers along the full onshore-offshore gradient remains technically challenging (Weymer et al., 2020), and observation strategies need to be developed for specific settings. Geophysical methods employed as imaging tools to characterize the subsurface offer promising avenues towards bridging the information gap across the land-sea interface, although they are only currently available on local scales (e.g. Siemon et al., 2020; Ishizu and Ogawa, 2021). Hydrologically connected OFG systems should in principle be associated with discharging groundwater (see e.g. Weymer et al., 2020), either close to the coastline, along faults or other lithological discontinuities, or at distant locations near the shelf break. However, OFG could also seep into the marine environment on time scales of >100–1000 kyr, making it difficult to obtain observations that provide insights on its effects on biological communities if no dedicated offshore drilling is carried out.

170

### 3 Environmental impacts and resource prospects

In the terrestrial realm, the role of coastal groundwater as a habitat (Pohlman, 2011; Leitão et al., 2015; Adyasari et al., 2019) and in shaping pelagic and benthic coastal communities (Lecher and Mackey, 2018; Oberle et al., 2022) has become increasingly recognized. In contrast, the role of OFG as a fresh- or brackish water habitat within a purely marine environment remains unknown and might constitute a new frontier in ocean sciences, also in view of its potential exploitation as an unconventional source of water.

180 Human use of OFG could affect both fresh groundwater discharging into the ocean and groundwater hydraulic heads on land. FSGD has local ecological impacts on e.g. seagrass (Carruthers et al., 2005), corals (Oehler et al., 2019; Correa et al., 2021; Oberle et al., 2022), phytoplankton (Rodellas et al., 2015; Sugimoto et al., 2017; Waska and Kim, 2010), mollusc (Hwang et al., 2010), meio/macrofauna (Zipperle and Reise, 2005; Kotwicki et al., 2014; Grzelak et al., 2018; Londoño-Londoño et al., 2022a) and fish populations (Fujita et al., 2019; Pisternick et al., 2020). These influences are often triggered by nutrient and carbon inputs into the submarine environment (Santos et al., 2021; Böttcher et al., 2022). Moreover, upward fluid migration within soft seafloor sediments might fluidize them, favouring the formation of pockmarks that potentially release greenhouse gases such as carbon dioxide and methane (Whiticar, 2002; Judd and Hovland, 2009; Donis et al., 2017; Virtasalo et al., 2019; Hoffmann et al., 2020; Purkamo et al., 2022). While some of these effects have been perceived as a threat for ecosystems, for instance by inducing toxic algal blooms, adding alkalinity (Cabral et al., 2021) or harbouring dense microbial communities (Ionescu et al., 2012), they can also sustain coastal ecology and increase fishery yields. Pumping OFG that is associated with FSGD could reduce the associated landward reservoirs and eventually impact the coastal marine environment. Moreover, anthropogenic intervention on coastal sediments might impact benthic-pelagic coupling associated with FSGD (von Ahn et al., 2021).

195

Considering the manifold biogeochemical impacts of FSGD, it is difficult to assess the overall effect of different pumping approaches and particular FSGD locations in local marine ecosystems, should a connected OFG be exploited. Pumping water from a groundwater system means reducing the formation

200 pressure. The reduced pressure can communicate to the terrestrial aquifer and reduce the hydraulic head  
there (Yu and Michael, 2019b). The extent of this effect will depend on the reservoir properties as well  
as the hydraulic connectivity between terrestrial and offshore domain, which might in turn lead to SWI  
(Ferguson and Gleeson, 2012; Yu and Michael, 2019a), groundwater depletion (Bierkens and Wada,  
2019) and subsidence (Yu and Michael, 2019b). Despite the large uncertainties on the global and long-  
205 term effects of these changes for groundwater resources and associated marine ecosystems, lessons may  
be learned from the environmental effects of extensive oil exploration (Varma and Michael 2012;  
Chaussard et al., 2013).

210 Changes in FSGD volume and its chemical/biological composition could serve as an important indicator  
for changes in the coastal groundwater system, which could, in turn, also be caused by connectivity with  
OFG. FSGD can be a source of geochemical tracers (e.g. Ra and Rn; Kim and Hwang, 2002), inorganic  
nutrients (nitrate, phosphate and silicate; e.g. Waska et al., 2011; Szymczyna et al., 2012), trace metals  
(e.g. Knee and Paytan, 2011), climate-relevant trace gases (carbon dioxide, nitrous oxide, methane and  
215 carbon monoxide; e.g. Bugna et al., 1996; Chapelle and Bradley, 2007; Jurado et al., 2017; Kolker et al.,  
2021; Reading et al., 2021) and organic material (e.g. dissolved organic matter; see Kim and Kim (2017)  
and McDonough et al. (2022)) to coastal areas. The input of nutrients results in a FSGD-driven  
eutrophication of coastal areas and, thus, potentially affects coastal ecosystems (Luijendijk et al., 2020;  
Oehler et al., 2021; Santos et al., 2021). For example, large outbreaks of the macroalgae *Ulva spp.* (so-  
220 called “green tides”), which occur regularly in eutrophic coasts off China and Korea, are attributed to the  
nutrient supply by FSGD (Kwon et al., 2017; Liu et al., 2017; Cho et al., 2019; Zhao et al., 2021). Hence,  
sustained monitoring of the biogeochemical and microbially-driven transformations of key  
225 biogeochemical tracers within the subterranean estuary as well as their release to the overlying water  
column, might help tracking changes in FSGD. Furthermore, such monitoring might also facilitate  
investigating potential impacts on the productivity and ecological status of coastal environments. Beyond  
coastal nearshore environments, FSGD seems to play an important role for biogeochemical fluxes to the  
230 ocean and affects benthic and sub-seafloor ecosystems in more offshore coastal areas (Micallef et al.,  
2021 and references therein). Therefore, a thorough investigation of its dynamics in different oceanic  
basins and geological settings should be performed by future studies.

235 Furthermore, FSGD and connected OFG could be increasingly affected by ongoing environmental  
changes on the terrestrial side (e.g. by changing rain patterns or intensity; Thomas and Famiglietti, 2019),  
eutrophication (derived from increasing applications of fertilizers), urbanization of coastal areas and  
associated contamination with microplastics (Viaroli et al., 2022), chemical (e.g. pesticides,  
pharmaceuticals, and personal care products; see Knee and Paytan, 2011; Szymczyna et al., 2020) and  
biological pollutants (bacteria and viruses, including pathogenic species; see e.g. Kyle et al., 2008;  
Sorensen et al., 2021). In particular, the effects of FSGD-driven inputs of chemical/biological pollutants  
on coastal areas remain largely unknown.

## 4 Conceptual and technological approaches for assessing land-ocean groundwater connectivity

Various techniques are available to explore and identify FSGD and OFG in the offshore environment (e.g. Micallef et al., 2021) and groundwater resources on the land side (Kirsch, 2006). These techniques often reveal anomalies in the subsurface, the seafloor (e.g. pockmarks) or sea water column (e.g. salinity, geochemical tracers) associated with fresh groundwater. Often multiple techniques are applied to build confidence in interpretation of groundwater dynamics. The technologies used can be broadly categorized in four groups: *i*) geophysical imaging techniques, which detect/record physical parameters such as electrical resistivity, seismic velocity, density, temperature or structural/morphological surface anomalies, *ii*) hydrogeological approaches, including modelling and hydrological measurements (e.g. hydraulic heads, salinity, and recharge rates) *iii*) (bio)geochemical techniques, which analyse (bio)geochemical fingerprints of the fluids, and *iv*) (micro)biological sampling, which unravels biological diversity and processes associated with FSGD (see e.g. Taniguchi et al., 2019; Micallef et al., 2021; Ruiz-González et al., 2021 and references therein). These multiple approaches are often mastered by researchers within different disciplinary backgrounds including geophysics, hydrology, oceanography and biogeochemistry.

Assessing land-ocean hydraulic connectivity through groundwater requires investigating the connectivity of underlying lithologic units and their hydrological characterization. Moreover, it also requires identifying the current distribution of freshened groundwater bodies across the coastline. The occurrence of freshened groundwater along the onshore-offshore continuum may in turn be read from geochemical fingerprinting of fluid samples obtained from the different realms. Hence, the success of such a highly interdisciplinary endeavour in mapping and understanding the connectivity will depend on how well the different methodologies can be integrated. Here, we suggest overarching approaches in which synergies (both conceptual and technological) between FSGD and OFG scientific communities could contribute to an improved understanding of the dynamics of groundwater as a connecting path between land and the ocean at the coastal zone. Table 2 presents some of the most commonly used methods in groundwater studies, for which we foresee promising synergies between FSGD and OFG research.

### 270 Shoreline-crossing lithologies

Seismic reflection imaging is the method of choice for detailed subsurface mapping. Particular lithologies may be identified by the character of the seismic reflection data within a lithological unit, e.g. layered seismic facies for fine-grained marine sediments vs. chaotic patterns for coarser grained sediments (Thomas et al., 2019; Micallef et al., 2020). Co-located boreholes on seismic sections can greatly improve the identification of different facies and serve as calibration points along those sections. Of particular importance for shoreline-crossing groundwater dynamics is the possibility of seismic data to constrain the continuity of different lithological units, the presence of impermeable clay layers and faults, or other disrupting geological structures. However, seismic information in the transition zone near the coastline is not widely available due to logistical challenges for data acquisition. Land and marine seismic data have inherently different signal-to-noise ratios and imaging depths, making across-shoreline interpretation challenging. On land there are often more boreholes than offshore, which can provide data to constrain the lithology distribution. Through the integration of onshore and offshore seismic and borehole data

using geostatistical methods such as sequential indicator simulation or multiple point geostatistics (Deutsch and Pyrcz 2014), lithology distribution across the shoreline can be modelled to reduce the uncertainty of connected pathways between the terrestrial and offshore domains.

Land and marine seismic data require different seismic sources, e.g. vibroseis on land and airguns/sparkers at sea, and receivers. Noise levels are generally higher on land, whereas offshore imaging in the transition zone is hampered by seafloor multiple reflections due to the shallow water depth. 285 Generally, clastic sedimentary environments are easier to image than carbonate systems (e.g. Mountain, 2008; Lofi et al., 2013; Bertoni et al., 2020). Amphibious data acquisition, i.e. across the shoreline, is possible and can be accomplished in different ways, for instance by shooting on land and receiving at sea or vice versa. Yet, due to logistical challenges and greater expenses, amphibious sections are not a standard. In karstic carbonate or volcanic systems, the spatial occurrence of localised submarine springs 295 (rather than the diffuse discharge in siliciclastic systems) can help to characterise the onshore-offshore connectivity of aquifers (Bayari et al., 2011).

Ground penetrating radar (GPR) is suitable for detailed, near-surface lithological imaging on land. The GPR technique is based on an electromagnetic signal that is sensitive to sediment water content. Offshore, 300 hydroacoustic and seismic methods provide structural information from shallow to larger depth, but are insensitive to the water content. However, unconformable boundaries of subsurface sediment units are typically imaged as strong reflectors in both GPR and reflection seismic profiles due to the associated sharp changes in water content and density, respectively, which permits the cross-shore correlation of onshore GPR profiles with marine seismic profiles using the allostratigraphic approach (see e.g. Virtasalo 305 et al., 2019; Peterson et al., 2020).

#### *Identification of groundwater bodies*

While seismic data can reveal the geological background and are -to some extent- sensitive to the porosity of the rock, they contain no information on pore fluid salinity. The salinity of pore fluids can be explored 310 using electrical methods because the bulk electrical resistivity of a sediment rock is governed by the amount (fluid-saturated pore space) and salinity of fluid present (Archie, 1942; Keller, 1987). The better the porosity of the lithology is known, for example through seismic/lithological data, the better the pore space fluid saturation on land and the pore water salinity offshore can be assessed from bulk electrical resistivity measurements. A bulk electrical resistivity model of the subsurface can be derived from either 315 direct or alternating current electrical measurements (electromagnetic induction), where the latter allows for larger penetration depths and better resolution offshore.

On land, the highest data acquisition speed and therefore the largest areal coverage is achieved through 320 airborne electromagnetic methods (e.g. Bedrosian et al., 2016; Gottschalk et al., 2020; Siemon et al., 2020). Additional ground measurements using direct current and controlled source electromagnetic methods provide bulk electrical resistivity **models** of the subsurface at higher resolution and larger depths of penetration (e.g. Ponthai et al., 2020). Resolution in surveys of electrical resistivity on land can be augmented by conducting GPR surveys. GPR methods allow both detecting contrasts in the electrical conductivity structure (dielectric constant) contained in coastal sediments at high resolution (cm to m

325 scales), and effectively mapping the freshwater-saltwater interface at shallow depths (up to tens of m; Weymer et al., 2020).

330 Offshore, a freshened groundwater body can be identified as an electrical resistivity anomaly caused by the resistivity contrast between fresh and saline pore water. The conductive saline ocean above the seafloor strongly damps electromagnetic signals which renders airborne electromagnetic systems incapable of penetrating the seafloor at water depths larger than about 10–20 m (Goebel et al., 2019). Therefore, offshore measurements require specially adapted marine electromagnetic systems. So far, OFG exploration studies have been conducted using surface-towed (e.g. Gustafson et al., 2019; Attias et al., 2021) and/or seafloor-towed (Haroon et al., 2018; 2021; Micallef et al., 2020) systems. Both systems 335 consist of a horizontal electric source dipole followed by several electric receiving dipoles recording the inline electric field. Offsets between transmitter and receiving dipoles typically range between hundreds and several hundreds of meters, and can be adjusted according to the target depth. Sea surface-towed systems have the advantage of a greater acquisition speed, yet at the cost of lower resolution and larger source dipole moments (current amplitude times dipole length) required to compensate for the decay of 340 the source signal in the conductive ocean layer. Seafloor-towed systems have arguably better signal to noise ratios and resolution, although survey speed is much lower, and surveying is hampered by rough seafloor topography and infrastructure. Onshore-offshore acquisition with a land transmitter and offshore receiver is possible (Ishizu and Ogawa, 2021). However, to date there are no peer-reviewed published 345 studies which use this approach. Merging of a separately acquired onshore-offshore electrical resistivity section with land and marine systems is possible, although a coherent continuous picture may be hampered by different resolutions, penetration depths, noise levels and the strong 3D resistivity contrast at the shoreline (“coast effect”; Worzewski et al., 2012). Recently, joint land/water data inversion methods 350 have become available to amend that deficit (Hermans and Paepen, 2020). Furthermore, conversion of electrical resistivity sections to water saturation on land or pore water salinity (the actual target parameters), requires integration of lithological data, i.e. bulk porosity estimates and an appropriate choice of effective medium model.

355 In-situ sampling techniques are effective and simple, albeit labour- and time-intensive ways to detect freshened groundwater. These methods include pore water extraction using push-point samplers along transects or grids (Waska et al., 2019), in-situ detection of springs with infrared cameras (Röper et al., 2014), and collection of seeping groundwater with seepage meters or benthic chambers (e.g. Lee 1977, Donis et al., 2017). Although mostly applied to nearshore groundwater discharge, all above-mentioned 360 methods are adaptable to remote systems, for instance on stationary landers or ROVs (e.g. Ahmerkamp et al., 2017).

#### 360 *Groundwater flow*

365 Imaging of coastal aquifers using inversion of geophysical data constrained by groundwater transport simulations is a promising method which might greatly reduce uncertainty in FSGD rates and location (Costall et al., 2020). Other promising approaches to detect FSGD over a larger area (tens of kilometres) while also allowing an assessment of its temporal variability, are thermal radiance measurements with manned (e.g. Roxburgh, 1985; Johnson et al., 2008) or unmanned (e.g. Fischer et al., 1964; Dulai et al.,

2016; Lee et al., 2016; Mallast and Siebert, 2019) aerial and sea-going vehicles, as well as usage of satellite-based thermal infrared imagery (Londoño-Londoño et al., 2022b).

370 While geophysical methods provide the geological background and current state of onshore-offshore groundwater distribution (Weymer et al., 2015), they do not capture the dynamics and functioning of the system which are essential to determining and understanding the nature of land-sea hydrologic connectivity. Physical hydrological measurements are essential for understanding groundwater flow rates and patterns. In coastal systems with variations in fluid density, this involves characterizing groundwater  
375 head distributions and associated hydraulic gradients, as well as the salinity distributions. On land, this is typically done with measurements from groundwater wells in addition to geophysics. Offshore, these measurements are more challenging but provide critical information on the forces driving fluid flow through the onshore-offshore system. Because offshore hydrologic data is generally sparse, groundwater modelling is an essential tool to test hypotheses about system function given the geological, hydrological,  
380 and biogeochemical data available. Groundwater models that incorporate physics-based variable-density flow and salt transport, and capture the essential characteristics of the system (e.g. interconnection of geologic strata; Michael et al., 2016; Thomas et al., 2022), can be used to understand the long-timescale evolution of OFG systems towards their current state (e.g. Cohen et al., 2010; Micallef et al., 2020; Zamrsky et al., 2020), and to predict changes under expected changes in sea level and anthropogenic  
385 forcing (Yu and Michael, 2019a; 2019b). These models can not only characterize the flow in the subsurface, but also characterize the rate and distribution of FSGD and/or diffusive transport processes.

In conjunction with geophysical and hydrological data and analyses, the geochemistry of groundwater fluid samples can provide key information about the origin and age of FSGD and OFG. A combination  
390 of stable isotope and conservative tracer analysis (e.g. Hoefs, 2009; Dang et al., 2020) can be used to identify sources and estimate ages of offshore groundwater bodies, i.e. recent or fossil meteoric water (van Geldern et al., 2013), glacial meltwater (Hong et al., 2019) or methane hydrate dissociation (Dählmann and De Lange, 2003). While onshore fluid samples required for this analysis are relatively easily obtained (typically from groundwater observation wells), OFG fluid sample collection requires in-  
395 situ sampling at depth through a borehole or, if existing, knowledge of FSGD occurrences on the seafloor. FSGD sites on the seafloor can be identified through detection of morphologic depressions (pockmarks, sinkholes) through high frequency acoustic seafloor bathymetry mapping and identification of anomalous seafloor fauna and flora associated with a change in water salinity and nutrients input (e.g. Lecher and Mackey, 2018; Archana et al., 2021). Other approaches used to search FSGD sites on a regional scale  
400 include mapping radiogenic isotopes that are associated with groundwater (Burnett, 2006; Paldor et al., 2020, Ikonen et al., 2022), shallow physical imaging of resistivity anomalies, survey of small-scale magnetic susceptibility anomalies caused by preservation or diagenetic alteration of iron oxides in sediments (Müller et al., 2011), satellite infrared imagery using e.g. Landsat 8 - infrared (e.g. Wilson and Rocha, 2012; Schubert et al., 2014; Jou-Claus et al. 2021), and surface reaching fault mapping by seismic  
405 methods.

FSGD may also cause measurable anomalies in the deeper water column of offshore sites (Manheim, 1967; Attias et al., 2021). While temperature and salinity anomalies are only measurable in the immediate

410 vicinity of the FSGD location and may be obscured by natural variations in water temperature and by tidal currents, radon and radium anomalies can be traced to larger distances (e.g. Cable et al., 1996; Moore et al., 2011). This methodology works well in areas of diffuse and uniform FSGD, but it might overlook localized point sources, which can account for up to 90% of FSGD in karstic regions (Null et al., 2014).

## 415 5 Future research directions

420 In view of the increasing pressure of human activities and natural changes on groundwater resources, the fundamental role of land-ocean connectivity through groundwater on the dynamics of coastal systems requires a critical reassessment. FSGD and the associated fluxes of biogeochemical tracers might affect the physical structure, chemical composition and reactivity, as well as (micro)biology of coastal ocean ecosystems. Global and regional environmental changes (i.e. warming, eutrophication, acidification, pollution) modify processes in coastal groundwater and thereby FSGD, with largely unknown consequences for coastal marine ecosystems. Exploitation of OFG connected to terrestrial groundwater is expected to impact terrestrial groundwater flow systems. These feedback mechanisms operate over a wide range of spatial and temporal scales, ranging from molecular to global and from millisecond to 425 millennial. Thus, an overarching goal of future coastal groundwater research should aim to develop a suite of ecosystem models of land-ocean connectivity that include physical, geological, chemical and biological processes at play, and that address potential responses to dynamic interactions between nature and humans.

430 Within this framework, we recommend the following priority research tasks:

(1) assess and compare the spatio-temporal variability of physical and biogeochemical processes driving the dynamics of FSGD and OFG in different geological settings,

435 (2) characterize and quantify the geochemical/biological composition of FSGD and OFG, as well as its impacts on marine habitats and (micro)biological communities,

(3) develop an interdisciplinary framework including hydrological, geophysical, geochemical and (micro)biological and measurements to delineate groundwater fluxes (FSGD) and map reservoirs (OFG) 440 along the transition from nearshore to offshore systems,

(4) characterize the stratigraphy at the land-ocean interface to determine the potential for development of connected, active OFG systems,

445 (5) identify, quantify, and predict feedbacks between coastal groundwater dynamics and climate change to assess potential changes in volume and composition of FSGD and OFG,

(6) use numerical models and artificial intelligence to predict locations, magnitudes and connectivity of FSGD and OFG, and

450

(7) implement the knowledge gained through models and observations to improve the representation of FSGD and OFG in Earth System models.

Investigation of the land-ocean connectivity through groundwater beyond nearshore FSGD remains especially challenging because of its limited accessibility and large heterogeneity. Its future study will require representative and standardized sampling, the development of new analytical methods (e.g., in-situ offshore groundwater measurements), and new observational and experimental frameworks. These endeavours should facilitate fully representative parameterizations of FSGD-OFG connectivity in numerical models across the land-sea interface. Moreover, developing hydrologic/oceanographic models of coastal and offshore groundwater and its interactions with other system compartments (sediments, water column, subseafloor environments) will help predicting future changes of groundwater dynamics on both regional and global scales.

It is evident that only multidisciplinary research initiatives, at both local, national and international levels, can effectively address the research tasks identified in this perspective paper. Joint projects should link laboratory, field, and modelling approaches to better understand the complex interplay of the various physical, chemical and biological processes operating along the land-ocean interface. Likewise, sustained observations will help to amend the current uncertainties in temporal variability of groundwater flows. An improved understanding of land-ocean connectivity in this context will contribute to our appreciation of the crucial role of coastal groundwater in societal-relevant issues such as climate change, pollution and the overall environmental status of the coastal oceans. Future research efforts in this topic will directly address the Sustainable Development Goals 6 (“*Clean water and sanitation*”), 12 (“*Responsible consumption and production*”) and 14 (“*Life below water*”) of the United Nations (see <https://www.un.org/sustainable-development/sustainable-development-goals/>).

#### ***Author contributions***

The initial draft of this perspectives manuscript was prepared by D.L.A.M., A.H., H.W.B, E.E., M.J., N.M. and J.S.v.D. All co-authors contributed to the revision and preparation of the final version of the manuscript.

#### ***Competing interests***

The authors declare that they have no conflict of interest.

#### ***Acknowledgements***

The conception of this manuscript was fostered by the discussions during a workshop sponsored by the Future Ocean Network of Kiel University (<https://www.futureocean.org/en/>) in February 2021. D.L.A.M. was supported by the Future Ocean Network (Grant # FON2020-03). The manuscript contains contributions to the SMART (Sustainable management of offshore groundwater resources) project funded by the Helmholtz Association, the KiSNet Network funded by the German Research Foundation (DFG; Grant # MA7041/6-1), the DFG research unit “DynaDeep” (FOR 5094/1) and the DFG research training group Baltic TRANSCOAST (GRK2000/0063). A.M. has received funding from the European Research

495 **Dedication**

*This article is dedicated to Dr. Volker Liebetrau (1965–2022). Volker, our co-author and friend, passed away on February 7<sup>th</sup> 2022. Volker is remembered for his enthusiasm, commitment, and smile. His passing is a massive loss to the community, but there will be much that will still be carried on in his memory thanks to his hard work and passion.*

500

## References

Abbott, B. W., Bishop, K., Zarnetske, J. P., Minaudo, C., Chapin, F. S., Krause, S., Hannah, D. M.,  
505 Conner, L., Ellison, D., Godsey, E. S., Plont, S., Marçais, J., Kolbe, T., Huebner, A., Frei, R. J., Hampton, T., Gu, S., Buhman, M., Sara Sayedi, S., Ursache, O., Chapin, M., Henderson, K. D., and Pinay, G.: Human domination of the global water cycle absent from depictions and perceptions, *Nat. Geosci.*, 12, 533–540, 2019.

510 Adyasari, D., Hassenrück, C., Oehler, T., Sabdaningsih, A., and Moosdorf, N.: Microbial community structure associated with submarine groundwater discharge in northern Java (Indonesia), *Sci. Total Environ.*, 689, 590–601, 2019.

Ahmerkamp, S., Winter, C., Krämer, K., Beer, D. d., Janssen, F., Friedrich, J., Kuypers, M. M. M., and  
515 Holtappels, M.: Regulation of benthic oxygen fluxes in permeable sediments of the coastal ocean, *Limnol. Oceanogr.*, 62, 1935–1954, <https://doi.org/10.1002/lno.10544>, 2017.

Archana, A., Francis, C. A., and Boehm., A. B.: The Beach Aquifer Microbiome: Research Gaps and Data Needs, *Front. Environ. Sci.*, 9, 653568, doi: 10.3389/fenvs.2021.653568, 2021.

520 Archie, G.E.: The Electrical Resistivity Log as an Aid in Determining Some Reservoir Characteristics, *Transactions of AIME*, 146, 54–62, 1942.

Attias, E., Constable, S., Sherman, D., Ismail, K., Shuler, C., and Dulai, H.: Marine electromagnetic  
525 imaging and volumetric estimation of freshwater plumes offshore Hawai'I, *Geophys. Res. Lett.*, 48, e2020GL091249, <https://doi.org/10.1029/2020GL091249>, 2021.

Bakken, T. H., Ruden, F., and Mangset, L. E.: Submarine groundwater: A new concept for the supply of drinking water, *Water Resour. Manag.*, 26(4), 1015–1026, <https://doi.org/10.1007/s11269-011-9806-1>,  
530 2012.

535 Bayari, C. S., Ozyurt, N. N., Oztan, M., Bastanlar, Y., Varinlioglu, G., Koyuncu, H., et al.: Submarine and coastal karstic groundwater discharges along the southwestern Mediterranean coast of Turkey, *Hydrogeol. J.*, 19(2), 399–414, <https://doi.org/10.1007/s10040-010-0677-y>, 2011.

Bedrosian, P. A., Schamper, C., and Auken, E.: A comparison of helicopter-borne electromagnetic systems for hydrogeologic studies, *Geophys. Prospect.*, 64, doi:10.1111/1365-2478.12262, 2016.

540 Beebe, D. A., Huettemann, M. B., Webb, B. M., and Jackson, W. T. Jr.: Atmospheric groundwater forcing of a subterranean estuary: A seasonal seawater recirculation process, *Geophys. Res. Lett.*, 49, e2021GL096154, <https://doi.org/10.1029/2021GL096154>, 2022.

Bertoni, C., Lofi, J., Micallef, A., and Moe, H.: Seismic reflection methods in offshore groundwater research, *Geosciences*, 10(8), 299, 2020.

545 Bierkens, M. F. P., and Wada, Y.: Non-renewable groundwater use and groundwater depletion: a review, *Environ. Res. Lett.*, 14, 063002, 2019.

550 Böttcher, M. E., Mallast, U., Massmann, G., Moosdorf, N., Müller-Petke, M., and Waska, H.: Coastal-Groundwater interfaces (submarine groundwater discharge). In (Krause, S., Hannah, D. M., and Grimm, N., eds.) *Ecohydrological Interfaces*, 400 p., ISBN: 978-1119489672, Wiley Science, 2022.

Bratton, J. F.: The Three Scales of Submarine Groundwater Flow and Discharge across Passive Continental Margins, *The Journal of Geology*, 118, 565–575, 2010.

555 Bugna, G., Chanton, J. P., Young, J. E., Burnett, W. C., and Cable, P. H.: The importance of groundwater discharge to the methane budgets of nearshore and continental shelf waters of the northeastern Gulf of Mexico, *Geochim. Cosmochim. Acta* 60, 4735–4746, doi: 10.1016/S0016-7037(96)00290-6, 1996.

560 Burnett, W. C., Aggarwal, P. K., Aureli, A., Bokuniewicz, H., Cable, J. E., Charette, M. A., Kontar, E., Krupa, S., Kulkarni, K. M., Loveless, A., Moore, W. S., Oberdorfer, J. A., Oliveira, J., Ozyurt, N., Povinec, P., Privitera, A. M. G., Rajar, R., Ramassur, R. T., Scholten, J., Stieglitz, T., Taniguchi, M., and Turner, J. V.: Quantifying submarine groundwater discharge in the coastal zone via multiple methods, *Sci. Total Environ.*, 367(2–3), 498–543, 2006.

565 Cable, J. E., Burnett, W. C., Chanton, J. P., and Weatherly, G. L.: Estimating groundwater discharge into the northeastern Gulf of Mexico using radon-222, *Earth Plan. Sci. Lett.*, 144(3–4), [https://doi.org/10.1016/S0012-821X\(96\)00173-2](https://doi.org/10.1016/S0012-821X(96)00173-2), 1996.

570 Cabral, A., Dittmar, T., Call, M., Scholten, J., de Rezende, C. E., Asp, N., Gledhill, M., Seidel, M., and Santos, I. R.: Carbon and alkalinity outwelling across the groundwater-creek-shelf continuum off Amazonian mangroves, *Limnol. Oceanogr. Lett.*, 6, 369–378, 2021.

575 Carruthers, T. J. B., van Tussenbroek, B. I., and Dennison, W. C.: Influence of submarine springs and wastewater on nutrient dynamics of Caribbean seagrass meadows, *Estuar. Coast. Shelf S.*, 64(2–3), 191–199, 2005.

580 Chapelle, F. H., and Bradley, P. M.: Hydrologic significance of carbon monoxide concentrations in groundwater, *Groundwater*, 45, 272–280, 2007.

585 Chaussard, E., Amelung, F., Abidin, H., and Hong, S.-H.: Sinking cities in Indonesia: ALOS PALSAR detects rapid subsidence due to groundwater and gas extraction, *Remote Sens. Environ.*, 128, 150–161, <https://doi.org/10.1016/j.rse.2012.10.015>, 2013.

590 Chen, C.-T., Hu, J.-C., Lu, C.-Y., Lee, J.-C., and Chan, Y.-C.: Thirty-year land elevation change from subsidence to uplift following the termination of groundwater pumping and its geological implications in the Metropolitan Taipei Basin, Northern Taiwan, *Eng. Geol.*, 95, 30–47, 2007.

595 Cho, H.-M., Kim, G., and Shin, K.-H.: Tracing nitrogen sources fueling coastal green tides off a volcanic island using radon and nitrogen isotopic tracers, *Sci. Total Environ.*, 665, 913–919, 2019.

Church, T.M.: A groundwater route for the water cycle, *Nature*, 380, 579–580, 1996.

595 Church, J.A., Clark, P. U., Cazenave, A., Gregory, J. M. et al.: Sea Level Change. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)], 1137–1216. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

600 Cohen, D., Person, M., Wang, P., Gable, C. W., Hutchinson, D., Marksamer, A., et al: Origin and extent of fresh paleowaters on the Atlantic continental shelf, USA, *Groundwater*, 48(1), 143–158, 2010.

605 Correa, R. E., Cardenas, M. B., Rodolfo, R. S., Lapus, M. R., Davis, K. L., Giles, A. B., Fullon, J. C., Hajati, M.-C., Moosdorf, N., Sanders, C. J., and Santos, I.R.: Submarine Groundwater Discharge Releases CO<sub>2</sub> to a Coral Reef, *ACS ES&T Water*, 1(8), 1756–1764, 2021.

610 Costall, A. R., Harris, B. D., Teo, B. et al.: Groundwater Throughflow and Seawater Intrusion in High Quality Coastal Aquifers, *Sci. Rep.*, 10, 9866, <https://doi.org/10.1038/s41598-020-66516-6>, 2020.

610 Dählmann, A., and de Lange, G. J.: Fluid-sediment interactions at Eastern Mediterranean mud volcanoes: A stable isotope study from ODP Leg 160. *Earth Plan. Sci. Lett.*, 212(3–4), 377–391, [https://doi.org/10.1016/S0012-821X\(03\)00227-9](https://doi.org/10.1016/S0012-821X(03)00227-9), 2003.

615 Dang, X., Gao, M., Wen, Z., Jakada, H., Hou, G., and Liu, S.: Evolutionary process of saline groundwater influenced by palaeo-seawater trapped in coastal deltas: A case study in Luanhe River Delta, China, *Estuar. Coast. Shelf S.*, 244, 106894, <https://doi.org/10.1016/j.ecss.2020.106894>, 2020.

620 Deutsch, C. V., and Pyrcz, M. In: Pyrcz M. J., and Deutsch C. V. (eds.) *Geostatistical reservoir modeling*, 2<sup>nd</sup> edn., Oxford University Press, Oxford, 448 p., 2014.

625 Donis, D., Janssen, F., Liu, B., Wenzhöfer, F., Dellwig, O., Escher, P., Spitz, A., and Böttcher, M.E.: Biogeochemical impact of submarine ground water discharge on coastal surface sands of the southern Baltic Sea, *Est. Coast. Shelf Res.*, 189, 131–142, 2017.

630 Dulai, H., Kamenik, J., Waters, C. A., Kennedy, J., Babinec, J., Jolly, J., et al.: Autonomous long-term gamma-spectrometric monitoring of submarine groundwater discharge trends in Hawaii, *J. Radioanal. Nucl. Chem.*, 307, 1865–1870, doi: 10.1007/s10967-015-4580-9, 2016.

Ferguson, G., and Gleeson, T.: Vulnerability of coastal aquifers to groundwater use and climate change. 630 *Nature Clim. Change*, 2, 342–345, <https://doi.org/10.1038/nclimate1413>, 2012.

635 Fischer, W. A., Landis, G. H., Moxham, R. M., and Polcyn, F.: Infrared Surveys of Hawaiian Volcanoes - Aerial Surveys with Infrared Imaging Radiometer Depict Volcanic Thermal Patterns + Structural Features, *Science*, 146(364), 733–742, 1964.

640 Fujiita, K., Shoji, J., Sugimoto, R., Nakajima, T., Honda, H., Takeuchi, M., Tominaga, O., and Taniguchi, M.: Increase in Fish Production Through Bottom-Up Trophic Linkage in Coastal Waters Induced by Nutrients Supplied via Submarine Groundwater, *Front. Environ. Sci.*, 7(82), <https://doi.org/10.3389/fenvs.2019.00082>, 2019.

645 Goebel, M., Knight R., and Halkjaer, M.: Mapping saltwater intrusion with an airborne electromagnetic method in the offshore coastal environment, Monterey Bay, California, *J. Hydrol.: Reg. Stud.*, 23, <https://doi.org/10.1016/j.ejrh.2019.100602>, 2019.

650 Gottschalk, I., Knight, R., Asch, T., Abraham, J., and Cannia J.: Using an airborne electromagnetic method to map saltwater intrusion in the northern Salinas Valley, California, *Geophysics*, 85 (4), B119–B131, <https://doi.org/10.1190/geo2019-0272.1>, 2020.

655 Grzelak, K., Tamborski, J., Kotwicki, L., and Bokuniewicz, H.: Ecostructuring of marine nematode communities by submarine groundwater discharge, *Mar Environ. Res.*, 136, 106–119, doi: 10.1016/j.marenvres.2018.01.013, 2018.

Gustafson, C., Key, K., and Evans, R. L.: Aquifer systems extending far offshore on the U.S. Atlantic margin, *Scientific Reports*, 9(1), 8709, <https://doi.org/10.1038/s41598-019-44611-7>, 2019.

Haroon, A., Lippert, K., Mogilatov, V., and Tezkan, B.: First application of the marine differential electric dipole for groundwater investigations: A case study from Bat Yam, Israel, *Geophysics*, 83(2), B59–B76, 2018.

660 Haroon, A., Micallef, A., Jegen, M., Schwalenberg, K., Karstens, J., Berndt, C., et al.: Electrical resistivity anomalies offshore a carbonate coastline: Evidence for freshened groundwater? *Geophys. Res. Lett.*, 48, e2020GL091909, <https://doi.org/10.1029/2020GL091909>, 2021.

665 Hermans, T., and Paepen, M.: Combined inversion of land and marine electrical resistivity tomography for submarine groundwater discharge and saltwater intrusion characterization, *Geophys. Res. Lett.*, 47, e2019GL085877, <https://doi.org/10.1029/2019GL085877>, 2020.

Hoefs, J.: *Stable isotope geochemistry*, 203 p., Springer, Berlin/Heidelberg, 2009.

670 Hoffmann, J. J. L., Schneider von Deimling, J., Schröder, J., Schmidt, M., Scholten, J., Crutchley, G. J., Gorman, A. R.: Complex eyed pockmarks and submarine groundwater discharge revealed by acoustic data and sediment cores in Eckernförde Bay, SW Baltic Sea, *Geochem. Geophys. Geosy.*, 21(4), <https://doi.org/10.1029/2019GC008825>, 2020.

675 Hong, W.-L., Lepland, A., Himmeler, T., Kim, J. H., Chand, S., Sahy, D., et al.: Discharge of meteoric water in the eastern Norwegian Sea since the last glacial period, *Geophys. Res. Lett.*, 46, 8194–8204, <https://doi.org/10.1029/2019GL084237>, 2019.

680 Hwang, D. W., Kim, G., Lee, W. C., and Oh, H. T.: The role of submarine groundwater discharge (SGD) in nutrient budgets of Gamak Bay, a shellfish farming bay, in Korea, *J. Sea Res.*, 64(3), 224–230, 2010.

Ionescu, D., Siebert, C., Polerecky, L., Munwes, Y. Y., Lott, C., et al.: Microbial and Chemical Characterization of Underwater Fresh Water Springs in the Dead Sea, *PLoS ONE*, 7(6), e38319, <https://doi:10.1371/journal.pone.0038319>, 2012.

685 Ishizu, K., and Ogawa, Y.: Offshore-onshore resistivity imaging of freshwater using a controlled-source electromagnetic method: A feasibility study, *Geophysics*, 86, E391–E405, <https://doi.org/10.1190/geo2020-0906.1>, 2021.

690 Jiao, J., and Post, V.: *Coastal Hydrogeology*. Cambridge: Cambridge University Press. doi:10.1017/9781139344142, 2019.

Johnson, A. G., Glenn, C. R., Burnett, W. C., Peterson, R. N., and Lucey, P. G.: Aerial infrared imaging reveals large nutrient-rich groundwater inputs to the ocean, *Geophys. Res. Lett.*, 35(15), 6, 2008.

695

Jou-Claus, S., Folch, A., and Garcia-Orellana, J.: Applicability of Landsat 8 thermal infrared sensor for identifying submarine groundwater discharge springs in the Mediterranean Sea basin, *Hydrol. Earth Syst. Sci.*, 25, 4789–4805, <https://doi.org/10.5194/hess-25-4789-2021>, 2021.

700 700 Judd, A., and Hovland, M: Seabed fluid flow: The impact on geology, biology and the marine environment, 492 p., Cambridge, Cambridge University Press, 2009.

Jurado, A., Borges, A. V., and Brouyère, S.: Dynamics and emissions of N<sub>2</sub>O in groundwater: A review, *Sci. Total Environ.*, 584–585, 207–218, 2017.

705 705 Jurasinski, G., Janssen, M., Voss, M., Böttcher, M. E., Brede, M., Burchard, H., Forster, S., Gosch, L., Gräwe, U., Gründling-Pfaff, S. et al.: Understanding the Coastal ecocline: Assessing sea-land-interactions at non-tidal, low-lying coasts through interdisciplinary research, *Front. Mar. Sciences*, 5(342), 1–22, doi: 10.3389/fmars.2018.00342, 2018.

710 Keller, G. V.: Rock and Mineral Properties, in: Electromagnetic Methods in Applied Geophysics, M. Nabighian (ed.), 1, II. Series: Investigations in Geophysics, SEG. ISBN 1-56080-069-0, 1987.

715 Kim, G., and Hwang, D.-W.: Tidal pumping of groundwater into the coastal ocean revealed from submarine <sup>222</sup>Rn and CH<sub>4</sub> monitoring, *Geophys. Res. Lett.*, 29(14,1678), 10.1029/2002GL015093, 2002.

Kim, J., and Kim, G.: Inputs of humic fluorescent dissolved organic matter via submarine groundwater discharge to coastal waters off a volcanic island (Jeju, Korea), *Scientific Reports*, 7(7921), 10.1038/s-41598-017-08518-5, 2017.

720 Kirsch, R., (Ed), *Groundwater Geophysics, A Tool for Hydrogeology*, Springer ISBN 10 3-540-29383-3, 2006.

725 Knee, K. L. and Paytan, A.: Submarine Groundwater Discharge: A source of nutrients, metals, and pollutants to the coastal ocean. In: Wolanski E and McLusky DS (eds.) *Treatise on Estuarine and Coastal Science*, Vol 4, pp. 205–233, Waltham: Academic Press, 2011.

Kohout, F. A.: The flow of fresh water and salt water in the Biscayne Bay Aquifer of the Miami area, Florida. In: Cooper, H. H., Kohout, F. A., Henry, H. R., and Glover, R. E. (Eds.), *Sea Water in Coastal Aquifers*. Geological survey water-supply paper, 1613-C, USGS, Washington, D.C., pp. 12–32, 1964.

730 730 Kolker, D., Bookman, R., Herut, B., David, N., and Silverman, J.: An initial assessment of the contribution of fresh submarine ground water discharge to the alkalinity budget of the Mediterranean Sea, *J. Geophys. Res. Oceans*, 126, e2020JC017085, <https://doi.org/10.1029/2020JC017085>, 2021.

Konikow, L. F.: Contribution of global groundwater depletion since 1900 to sea-level rise, *Geophys. Res. Lett.*, 38(17), 2011.

740 Kooi, H., and Groen, K.: Offshore continuation of coastal groundwater systems; predictions using sharp-interface approximations and variable-density flow modelling, *J. Hydrol.*, 246(1–4), 19–35, [https://doi.org/10.1016/S0022-1694\(01\)00354-7](https://doi.org/10.1016/S0022-1694(01)00354-7), 2001.

745 Kotwicki, L., Grzelak, K., Czub, M., Dellwig, O., Gentz, T., Szymczyska, B., and Böttcher, M. E.: Submarine groundwater discharge to the Baltic coastal zone: Impacts on the meiofaunal community, *J. Mar. Sys.*, 129, 118–126, 2014.

750 Kwon, H. K., Kang, H., Oh, Y. H., Park, S. R., and Kim, G.: Green tide development associated with submarine groundwater discharge in a coastal harbor, Jeju, Korea, *Sci. Rep.*, 7, 6325, <https://doi.org/10.1038/s41598-017-06711-0>, 2017.

755 Kyle, J., Eydal, H., Ferris, F. et al.: Viruses in granitic groundwater from 69 to 450 m depth of the Äspö hard rock laboratory, Sweden. *ISME J.*, 2, 571–574, <https://doi.org/10.1038/ismej.2008.18>, 2008.

Ikonen, J., Hendriksson, N., Luoma, S., Lahaye, Y., and Virtasalo, J. J.: Behavior of Li, S and Sr isotopes in the subterranean estuary and seafloor pockmarks of the Hanko submarine groundwater discharge site in Finland, northern Baltic Sea, *Appl. Geochem.*, 105471, <https://doi.org/10.1016/j.apgeochem.2022.105471>, 2022.

760 Lecher, A., and Mackey, K.: Synthesizing the effects of submarine groundwater discharge on Marine Biota, *Hydrology*, 5(60), doi: 10.3390/hydrology5040060, 2018.

Lee, D. R.: A device for measuring seepage flux in lakes and estuaries, *Limnol. Oceanogr.*, 22(1), 140–147, 1977.

765 Lee, E., Yoon, H., Hyun, S. P., Burnett, W. C., Koh, D. C., Kang, K. M., et al.: Unmanned aerial vehicles (UAVs)-based thermal infrared (TIR) mapping, a novel approach to assess groundwater discharge into the coastal zone, *Limnol. Oceanogr.* 14, 725–735, doi: 10.1002/lom3.10132, 2016.

770 Leitão, F., Encarnação, J., Range, P., Schmelz, R. M., Teodósio, M.A., and Chícharo, L.: Submarine groundwater discharges create unique benthic communities in a coastal sandy marine environment, *Estuar. Coast. Shelf S.*, 163, 93–98, 2015.

Liu, K.-K., Atkinson, L., Quiñones, R., and Talaue-McManus, L. (eds.): Carbon and nutrient fluxes in continental margins: A global synthesis, Berlin & Heidelberg, 741 pp., 2010.

775 Liu, J., Su, N., Wang, X., and Du, J.: Submarine groundwater discharge and associated nutrient fluxes into the Southern Yellow Sea: A case study for semi-enclosed and oligotrophic seas - implication for green tide bloom, *J. Geophys. Res. Oceans*, 122, 139–152, doi:10.1002/2016JC012282, 2017

780 Lofi, J., Pezard, P., Bouchette, F., Raynal, O., Sabatier, P., Denchik, N., et al.: Integrated onshore-offshore investigation of a Mediterranean layered coastal aquifer, *Groundwater*, 51(4), 550–561, 2013.

785 Londoño-Londoño, J. E., Condesso de Melo, M. T., and Silva, A. C. F.: Groundwater discharge locally shapes the rocky shore macroinvertebrate community in South-Southwest Portugal, *Mar. Environ. Res.*, 179, 105672, <https://doi.org/10.1016/j.marenvres.2022.105672>., 2022a.

790 Londoño-Londoño, J. E., Condesso de Melo, M. T., Nascimento, J. N., and Silva, A. C. F.: Thermal-Based Remote Sensing Solution for Identifying Coastal Zones with Potential Groundwater Discharge, *J. Mar. Sci. Eng.*, 10(414), <https://doi.org/10.3390/jmse10030414>, 2022b.

795 Mallast, U. and Siebert, C.: Combining continuous spatial and temporal scales for SGD investigations using UAV-based thermal infrared measurements, *Hydrol. Earth Syst. Sci.*, 23, 1375–1392, <https://doi.org/10.5194/hess-23-1375-2019>, 2019.

800 Manheim, F. T.: Section of geological sciences: evidence for submarine discharge of water on the Atlantic continental slope of the southern United States, and suggestions for further search, *Transactions of the New York Academy of Sciences*, 29(7 Series II), 839–853, 1967.

805 Masaumi, K., Susumu, A., Hideo, S., and Hiroshi, S.: Reciprocal data acquisition and subsequent waveform matching for integrated onshore–offshore seismic profiling, *Geophys. J. Int.*, 212(1), 509–521, <https://doi.org/10.1093/gji/ggx374>, 2018.

810 McDonough, L. K., Andersen, M. S., Behnke, M. I. et al.: A new conceptual framework for the transformation of groundwater dissolved organic matter, *Nat. Commun.*, 13(2153), <https://doi.org/10.1038/s41467-022-29711-9>, 2022.

815 Micallef, A., Person, M., Haroon, A. et al.: 3D characterisation and quantification of an offshore freshened groundwater system in the Canterbury Bight, *Nat. Commun.*, 11, 1372, <https://doi.org/10.1038/s41467-020-14770-7>, 2020.

820 Micallef, A., Person, M., Berndt, C., Bertoni, C., Cohen, D., Dugan, B., et al.: Offshore freshened groundwater in continental margins, *Rev. Geophys.*, 58, e2020RG000706, <https://doi.org/10.1029/2020RG000706>, 2021.

825 Michael, H. A., Scott, K. C., Koneshloo, M., Yu, X., Khan, M. R., and Li, K.: Geologic influence on groundwater salinity drives large seawater circulation through the continental shelf, *Geophys. Res. Lett.*, 43, 10782–10791, <https://doi.org/10.1002/2016GL070863>, 2016.

Moore, W. S.: The Effect of Submarine Groundwater Discharge on the Ocean, *Annu. Rev. Mar. Sci.*, 2, 59–88, 2010.

825

Moore, W. S., Beck, M., Riedel, T., Rutgers van der Loeff, M., Dellwig, O., Shaw, T. J., Schnetger, B., and Brumsack, H.-J.: Radium-based pore water fluxes of silica, alkalinity, manganese, DOC, and uranium: a decade of studies in the German Wadden Sea, *Geochim. Cosmochim. Acta*, 75, 6535–6555, 2011.

830

Moore, W. S., and Joye, S. B.: Saltwater Intrusion and Submarine Groundwater Discharge: Acceleration of Biogeochemical Reactions in Changing Coastal Aquifers, *Front. Earth Sci.*, 9:600710, <https://doi.org/10.3389/feart.2021.600710>, 2021.

835

Moosdorf, N., Böttcher, M. E., Adyasari, D., Erkul, E., Gilfedder, B. S., Greskowiak, J., Jenner, A.-K., Kotwicki, L., Massmann, G., Müller-Petke, M., Oehler, T., Post, V., Prien, R., Scholten, J., Siemon, B., Ehlert von Ahn, C. M., Walther, M., Waska, H., Wunderlich, T., and Mallast, U.: A State-Of-The-Art Perspective on the Characterization of Subterranean Estuaries at the Regional Scale, *Front. Earth Sci.*, 9, 601293, <https://doi.org/10.3389/feart.2021.601293>, 2021.

840

Mountain, G.: Portable hires multi-channel seismic shot data from the New Jersey slope acquired during the r/v oceanus expedition oc270 (1995), <https://doi.org/10.1594/IEDA/307762>, 2008.

845

Müller, H., von Dobeneck, T., Nehmiz, W. et al.: Near-surface electromagnetic, rock magnetic, and geochemical fingerprinting of submarine freshwater seepage at Eckernförde Bay (SW Baltic Sea), *Geo-Mar. Lett.* 31, 123–140, <https://doi.org/10.1007/s00367-010-0220-0>, 2011.

850

Null, K. A., Knee, K. L., Crook, E. D., de Sieyes, N. R., Rebollo-Vieyra, M., Hernández-Terrones, L., and Paytan, A.: Composition and fluxes of submarine groundwater along the Caribbean coast of the Yucatan Peninsula, *Cont. Shelf Res.*, 77, 38–50, <https://doi.org/10.1016/j.csr.2014.01.011>, 2014.

Oberle, F. K. J., Prouty, N. G., Swarzenski, P. W. et al.: High-resolution observations of submarine groundwater discharge reveal the fine spatial and temporal scales of nutrient exposure on a coral reef: Faga'alu, AS. *Coral Reefs*, <https://doi.org/10.1007/s00338-022-02245-8>, 2022.

855

Oehler, T., Ramasamy, M., Mintu, E. G., Babu, S. D. S., Dähnke, K., Ankele, M., Böttcher, M. E., Santos, I. R. and Moosdorf, N.: Tropical Beaches Attenuate Groundwater Nitrogen Pollution Flowing to the Ocean, *Environ. Sci. Technol.*, 55(12), 8432–8438, 2021.

860

Oehler, T., Bakti, H., Lubis, R. F., Purwoarminta, A., Delinom, R., and Moosdorf, N.: Nutrient dynamics in submarine groundwater discharge through a coral reef (western Lombok, Indonesia), *Limnol. Oceanogr.*, 64(6), 2646–2661, 2019.

865 Paldor, A., Katz, O., Aharonov, E., Weinstein, Y., Roditi-Elasar, M., Lazar, A., and Lazar, B.: Deep submarine groundwater discharge—evidence from Achziv submarine canyon at the exposure of the Judea group confined aquifer, Eastern Mediterranean, *J. Geophys. Res. Oceans*, 125(1), e2019JC015435, 2020.

870 Peterson, C. D., Jol, H. M., Percy, D., and Perkins, R.: Use of Ground Penetrating Radar, Hydrogeochemical Testing, and Aquifer Characterization to Establish Shallow Groundwater Supply to the Rehabilitated Ni-les' tun Unit Floodplain: Bandon Marsh, Coquille Estuary, Oregon, USA, *J. Geogr. Geol.*, 12, 25–49, 2020.

875 Pisternick, T., Lilkendey, J., Audit-Manna, A., Dumur Neelayya, D., Neehaul, Y., and Moosdorf, N.: Submarine groundwater springs are characterized by distinct fish communities, *Mar. Ecol. e12610*, <https://doi.org/10.1111/maec.12610>, 2020.

Pohlman, J. W.: The biogeochemistry of anchialine caves: progress and possibilities, *Hydrobiologia*, 677(1), 33–51, 2011.

880 Pondthai, P., Everett, M. E., Micallef, A., Weymer, B. A., Faghih, Z., Haroon, A., and Jegen, M.: 3D Characterization of a Coastal Freshwater Aquifer in SE Malta (Mediterranean Sea) by Time-Domain Electromagnetics. *Water*, 12, 1566. <https://doi.org/10.3390/w12061566>, 2020.

885 Post, V. E. A., Groen, J., Kooi, H., Person, M., Ge, S., and Edmunds, W. M.: Offshore fresh groundwater reserves as a global phenomenon, *Nature*, 504(7478), 71–78, <https://doi.org/10.1038/nature12858>, 2013.

890 Purkamo, L., Milene, C., von Ahn, E., Jilbert, T., Muniruzzaman, M., Bange, H. W., Jenner, A.-K., Böttcher, M. E., and Virtasalo, J. J.: Impact of submarine groundwater discharge on biogeochemistry and microbial communities in pockmarks, *Geochem. Cosmochim. Acta*, 334, 14–44, 2022.

Reading, M. J., Tait, D. R., Maher, D. T., Jeffrey, L. C., Correa, R. E., Tucker, J. P., Shishaye, H. A., and Santos, I. R.: Submarine groundwater discharge drives nitrous oxide source/sink dynamics in a metropolitan estuary, *Limnol. Oceanogr.*, 66, 1665–1686, <https://doi.org/10.1002/lno.11710>, 2021.

895 Rocha, C., Robinson, C. E., Santos, I. R., Waska, H., Michael, H. A. and Bokuniewicz, H. J.: A place for subterranean estuaries in the coastal zone, *Est. Coast. Shelf Res.*, 250, 107167, 2021.

900 Rodellas, V., Garcia-Orellana, J., Masque, P., Feldman, M., and Weinstein, Y.: Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea, *PNAS*, 112(13), 3926–3930, 2015.

Roxburgh, I. S.: Thermal infrared detection of submarine springs associated with the Plymouth Limestone, *Hydrolog. Sci. J.*, 30(2), 185–196, 1985.

905 Röper, T., Greskowiak, J., and Massmann, G.: Detecting small groundwater discharge springs using handheld thermal infrared imagery, *Groundwater*, 52(6), 936–942, 2014.

Ruiz-González, C., Rodellas, V., and Garcia-Orellana, J.: The microbial dimension of submarine groundwater discharge: current challenges and future directions, *FEMS Microbiol. Rev.*, 45(5), fuab010, <https://doi.org/10.1093/femsre/fuab010>, 2021.

910

Santos, I. R., Chen, X., Lecher, A. L., Sawyer, A. H., Moosdorf, N., Rodellas, V., Tamborski, J., Cho, H.-M., Dimova, N., Sugimoto, R., Bonaglia, S., Li, H., Hajati, M.-C., and Li, L.: Submarine groundwater discharge impacts on coastal nutrient biogeochemistry, *Nat. Rev. Earth. Environ.*, 2, 307 – 323, <https://doi.org/10.1038/s43017-021-00152-0>, 2021.

915

Sawyer, A. H., David, C. H., and Famiglietti, J. S.: Continental patterns of submarine groundwater discharge reveal coastal vulnerabilities, *Science*, 353, 6300, 7005–707, 2016.

920

Schubert, M., Scholten, J., Schmidt, A., Comanducci, J. F., Pham, M. K., Mallast, U., and Knoeller, K.: Submarine Groundwater Discharge at a Single Spot Location: Evaluation of Different Detection Approaches, *Water*, 6, 584–601, <https://doi.org/10.3390/w6030584>, 2014.

Shlklomanov, I. A.: World fresh water resources. In: Gleick, P. H. (ed.), *Water in Crisis: A Guide to the World's Fresh Water Resources*, Oxford University Press, New York, pp. 13–24, 1993.

925

Siemon, B., Ibs-von Seht, M., Steuer, A., Deus, N., and Wiederhold, H.: Airborne Electromagnetic, Magnetic, and Radiometric Surveys at the German North Sea Coast Applied to Groundwater and Soil Investigations, *Remote Sens.*, 12, 1629, <https://doi.org/10.3390/rs12101629>, 2020.

930

Sorensen, J. P. R., Aldous, P., Bunting, S. Y., McNally, S., Townsend, B. R., Barnett, M. J., Harding, T., La Ragione, R. M., Stuart, M. E., Tipper, H. J., and Pedley, S.: Seasonality of enteric viruses in groundwater-derived public water sources, *Water Res.*, 207, <https://doi.org/10.1016/j.watres.2021.117813>, 2021.

935

Sugimoto, R., Kitagawa, K., Nishi, S., Honda, H., Yamada, M., Kobayashi, S., Shoji, J., Ohsawa, S., Taniguchi, M., and Tominaga, O.: Phytoplankton primary productivity around submarine groundwater discharge in nearshore coasts, *Mar. Ecol. Progr. Ser.*, 563, 25–33, 2017.

940

Szymczycza, B., Borecka, M., Biały-Bielińskab, A., Siedlewicz, G., and Pazdro, K.: Submarine groundwater discharge as a source of pharmaceutical and caffeine residues in coastal ecosystem: Bay of Puck, southern Baltic Sea case study, *Sci. Total Environ.*, 713, 136522, 2020.

Szymczycza, B., Vogler, S., and Pempkowiak, J.: Nutrients fluxes via submarine groundwater discharge to the Bay of Puck, Southern Baltic, *Sci. Total Environ.*, 438, 86–93, 2012.

945

Taniguchi, M., Dulai, H., Burnett, K. M., et al.: Submarine Groundwater Discharge: Updates on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects, *Front.*

950 50 Taniguchi, M., Burnett, W. C., Cable, J. E., and Turner, J. V.: Investigation of submarine groundwater discharge, *Hydrol. Process.* 16, 2115–2129. doi: 10.1002/hyp.1145, 2002.

Taylor, R., Scanlon, B., Döll, P. et al: Ground water and climate change, *Nat. Clim. Change*, 3, 322–329, 2013.

955 55 Thomas, A. T., von Harten, J., Jusri, T., Reiche, S., and Wellmann, F.: An integrated modeling scheme for characterizing 3D hydrogeological heterogeneity of the New Jersey shelf, *Mar. Geophys. Res.*, 43(2), 1–19, 2022.

960 60 Thomas, B. F., and Famiglietti, J. S.: Identifying Climate-Induced Groundwater Depletion in GRACE Observations, 9, 4124, <https://doi.org/10.1038/s41598-019-40155-y>, 2019.

965 65 Thomas, A. T., Reiche, S., Riedel, M., and Clauser, C.: The fate of submarine fresh groundwater reservoirs at the New Jersey shelf, USA, *Hydrogeol. J.*, 27(7), 2673–2694, <https://doi.org/10.1007/s10040-019-01997-y>, 2019.

970 70 Varma, S., and Michael, K.: Impact of multi-purpose aquifer utilisation on a variable-density groundwater flow system in the Gippsland Basin, Australia, *Hydrogeol. J.*, 20(1), 119–134. <https://doi.org/10.1007/s10040-011-0800-8>, 2012.

975 75 Van Geldern, R., Hayashi, T., Bottcher, M. E., Mottl, M., Barth, J. A. C., and Stadler, S: Stable isotope geochemistry of pore waters and marine sediments from the New Jersey shelf: Methane formation and fluid origin, *Geosphere*, 9(1), 96–112, <https://doi.org/10.1130/GES00859.1>, 2013.

980 80 Van Meter, K. J., Van Cappellen, P., and Basu, N. B.: Legacy nitrogen may prevent achievement of water quality goals in the Gulf of Mexico, *Science* 360, 427–430, 2018.

Viaroli, S., Lancia, M., and Re, V.: Microplastics contamination of groundwater: Current evidence and future perspectives. A review, *Sci. Total Environ.*, 824, <https://doi.org/10.1016/j.scitotenv.2022.153851>, 2022.

985 85 Virtasalo, J. J., Schröder, J. F., Luoma, S., Majaniemi, J., Mursu, J., and Scholten, J.: Submarine groundwater discharge site in the First Salpausselkä ice-marginal formation, south Finland, *Solid Earth*, 10, 405–423, <https://doi.org/10.5194/se-10-405-2019>.

von Ahn, C. M. E., Scholten, J., Malik, C., Feldens, P., Liu, B., Dellwig, O., Jenner, A.-K., Papenmeier, S., Schmiedinger, I., Zeller, M. A., and Böttcher, M. E.: A multi-tracer study of fresh submarine and surface water sources for a temperate urbanized coastal bay, *Front. Environm. Sci.*, 9, 642346, 2021.

990 Waska, H., and Kim, G.: Submarine groundwater discharge (SGD) as a main nutrient source for benthic and water-column primary production in a large intertidal environment of the Yellow Sea, *J. Sea Res.* 65, 103–113., doi: 10.1016/j.seares.2010.08.001, 2011.

995 Waska, H., and Kim, G.: Differences in microphytobenthos and macrofaunal abundances associated with groundwater discharge in the intertidal zone, *Mar. Ecol. Progr. Ser.*, 407, 159–172, 2010.

1000 Waska, H., Geskowiak, J., Ahrens, J., Beck, M., Ahmerkamp, S., Böning, P., Brusmack, H. J., Degenhardt, J., Ehlert, C., Engelen, B., Grünebaum, N., Holtappels, M., Pahnke, K., Marchant, H. K., Massmann, G., Meier, D., Schnetger, B., Schwalfenberg, K., Simon, H., Vandieken, V., Tzielinski, O., and Dittmar, T.: Spatial and temporal patterns of pore water chemistry in the inter-tidal zone of a high energy beach, *Front. Mar. Sci.*, 6, 154, doi:10.3389/fmars.2019.00154, 2019.

1005 Weymer, B. A., Everett, M. E., Smet, T. S., and Houser, C.: Review of electromagnetic induction for mapping barrier island framework geology, *Sediment. Geol.*, 321, 11–24, 2015.

1010 Weymer, B. A., Wernette, P. A., Everett, M. E., Pondthai, P., Jegen, M., and Micallef, A.: Multi-layered high permeability conduits connecting onshore and offshore coastal aquifers, *Front. Mar. Sci.*, 7(903), <https://doi.org/10.3389/fmars.2020.531293>, 2020.

1015 Whiticar, M. J.: Diagenetic relationships of methanogenesis, nutrients, acoustic turbidity, pockmarks and freshwater seepages in Eckernförde Bay, *Mar. Geol.*, 182(1–2), 29–53, [https://doi.org/10.1016/S0025-3227\(01\)00227-4](https://doi.org/10.1016/S0025-3227(01)00227-4), 2002.

1020 Wilson, J., and Rocha, C.: Regional scale assessment of Submarine Groundwater Discharge in Ireland combining medium resolution satellite imagery and geochemical tracing techniques, *Remote Sens. Environ.*, 119, 21–34, <https://doi.org/10.1016/j.rse.2011.11.018>, 2012.

1025 Worzewski, T., Jegen, M., and Swidinsky, A.: Approximation for the 2D coast effect on marine magnetotelluric data, *Geophys. J. Int.*, 189, 357–368, <https://doi.org/10.1111/j.1365-246X.2012.05385.x>, 2012.

1030 Yu, X., and Michael, H. A.: Mechanisms, configuration typology, and vulnerability of pumping-induced seawater intrusion in heterogeneous aquifers, *Adv. Wat. Resour.*, doi:10.1016/j.advwatres.2019.04.013, 2019a.

1025 Yu, X., and Michael, H. A.: Offshore pumping impacts onshore groundwater resources and land subsidence, *Geophys. Res. Lett.*, 46, 2553–2562, <https://doi.org/10.1029/2019GL081910>, 2019b.

1030 Zamrsky, D., Essink, G. H. O., Sutanudjaja, E. H., van Beek, L. R., and Bierkens, M. F.: Offshore fresh groundwater in coastal unconsolidated sediment systems as a potential fresh water source in the 21<sup>st</sup> century, *Environ. Res. Lett.*, 17(1), 014021, 2021.

1035 Zhao, S., Xu, B., Yao, Q., Burnett, W. C., Charette, M. A., Su, R., Lian., E., and Yu, Z.: Nutrient-rich submarine groundwater discharge fuels the largest green tide in the world, *Sci. Total Environ.*, 770, <https://doi.org/10.1016/j.scitotenv.2020.144845>, 2021.

Zipperle, A., and Reise, K.: Freshwater springs on intertidal sand flats cause a switch in dominance among polychaete worms, *J. Sea Res.*, 54(2), 143–150, <https://doi.org/10.1016/j.seares.2005.01.003>, 2005.

1040

1045


1050

1055

1060


1065

1070



1080 **Figure 1.** Schematic representation of known pathways for the transport of and storage of fresh and freshened groundwater between terrestrial and marine realms. Areas surrounded by dashed lines indicate groundwater reservoirs, whereas arrows represent freshwater (green) and seawater (red) fluxes. Based on Bratton et al. (2010) and Weymer et al. (2020).

1085



1090 **Figure 2.** Global distribution of reported FSGD (red circles) and OFG (blue triangles) sites. Location data from FSGD and OFG from Luijendijk et al. (2020) and Micallef et al. (2021), respectively.

## Tables

**Table 1.** Key concepts used in this manuscript.

Term	Definition
Meteoric water	Waters derived from precipitation. These waters reach the ocean either through surface flows (e.g. rivers), or as groundwater after infiltrates in soils.
Aquifer	Underground water reservoir that can consist of several layers of rock or sediments.
Groundwater	Water reservoir located beneath land surfaces.
Groundwater recharge	Replenishment of an aquifer containing groundwater from surface sources.
Fresh submarine groundwater discharge (FSGD)	Flow of fresh meteoric groundwater from terrestrial coastal aquifers through the seafloor into the ocean.
Offshore freshened groundwater (OFG)	Reservoir of fresh and brackish groundwater embedded in sediment pore waters and rocks below the seafloor.
Seawater intrusion (SWI)	Flows of marine waters into freshwater aquifers.
Non-renewable groundwater	Groundwater whose renewal (through recharge) takes place in times scales > 100 years (see Bierkins and Wada, 2019).
Fossil groundwater	Groundwater stored over millennia in isolated reservoirs below the Earth's surface.
Subterranean estuary	Coastal aquifer connected to the ocean which bears both saline and meteoric waters.

1095

**Table 2.** Commonly used methods for investigating groundwater fluxes and reservoirs. \* Current application realm.

Approach	Spatial scales	Temporal scales	Captured processes / controlling mechanisms	FSGD/OFG*
Thermal infrared sensing	cm to km	hours to months	Inflow of low-density plumes. Assessment on sea surface temperature anomalies with respect seasonal means	FSGD
Electrical ground conductivity	m to km	hours to years	Temporal variability of fresh-salt interfaces Recirculation fluxes Setting of sub-surface salt balance models	FSGD/OFG
Seafloor mapping & Sub-bottom profiling (Acoustics)	cm to km	-	Presence of seafloor depressions (e.g. "Wonky Holes") Pockmarks formation	FSGD/OFG
Electromagnetics	m to km	-	Electrical resistivity anomalies within the seafloor and water column that are indicative of active groundwater discharge	FSGD/OFG
Direct measurements of seepage rates	cm to m	hours	Quantification of fresh groundwater discharge rates	FSGD
In-situ surveys with remotely operated vehicles	m to km	-	Quantification of fresh groundwater discharge rates	FSGD
Hydrological modelling	m to km	-	Characterization of groundwater fluxes and chemical transformations Simulation of aquifer properties under hydrological changes	FSGD/OFG
Radon isotopes measurements	cm to km	days	Assessment of local sources and recent inputs based on strong gradients between groundwater and ocean Tracking of groundwater-derived greenhouse gases	FSGD
Dissolved organic matter measurements	cm to km	-	Concentration distributions and composition are used to track FSGD properties and dispersal	FSGD

Measurements of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures	km	-	Assessment of spatial distribution and C and N flows due to FSGD	FSGD/OFG
Nutrient analysis	km	-	Assessment of spatial distribution and estimation of primary production	FSGD/OFG
Water isotopes ( $\delta\text{D}$ and $\delta^{18}\text{O}$ )	m to km	months to centuries	Identification of recharge processes	FSGD
Gas measurements	cm to km	days to months	Assessment of FSGD-driven net community production Quantification of trace gas production and emissions to the atmosphere	FSGD/OFG
Phytoplankton analysis	km	-	Assessment of FSGD effects on primary production	FSGD
Benthic fauna sampling	m to km	-	Assessment of FSGD effects on benthic biomass & diversity	FSGD
Microbial ecology analyses	cm to km	-	Evaluation of abundance and diversity differences within FSGD sites	FSGD