

Quantifying land carbon cycle feedbacks under negative CO₂ emissions

V. Rachel Chimuka¹, Claude-Michel Nzotungicimpaye^{1,a} & Kirsten Zickfeld¹

¹Department of Geography, Simon Fraser University, Burnaby, BC, V5A 1S6, Canada

^a Now at Department of Geography, Planning and Environment, University of Concordia, Montréal, QC, H3G 1M8, Canada

Correspondence to: V. Rachel Chimuka (rchimuka@sfu.ca)

Abstract. Land and ocean carbon sinks play a major role in regulating atmospheric CO₂ concentration and climate. However, their future efficiency depends on feedbacks in response to changes in atmospheric CO₂ concentration and climate, namely the concentration-carbon and climate-carbon feedbacks. Since carbon dioxide removal is a key mitigation measure in emission scenarios consistent with global temperature goals in the Paris Agreement, understanding carbon cycle feedbacks under negative CO₂ emissions is essential. This study investigates land carbon cycle feedbacks under positive and negative CO₂ emissions using an Earth system model of intermediate complexity (EMIC) driven with an idealized scenario of symmetric atmospheric CO₂ concentration increase (ramp-up) and decrease (ramp-down), run in three modes. Our results show that the magnitudes of carbon cycle feedbacks are generally smaller in the atmospheric CO₂ ramp-down phase than in the ramp-up phase, except for the ocean climate-carbon feedback, which is larger in the ramp-down phase. This is largely due to carbon cycle inertia: the carbon cycle response in the ramp-down phase is a combination of the committed response to the prior atmospheric CO₂ increase and the response to decreasing atmospheric CO₂. To isolate carbon cycle feedbacks under decreasing atmospheric CO₂ and quantify these feedbacks more accurately, we propose a novel approach that uses zero emissions simulations to quantify the committed carbon cycle response. We find that the magnitudes of the concentration-carbon and climate-carbon feedbacks under decreasing atmospheric CO₂ are larger in our novel approach than in the standard approach. Accurately quantifying carbon cycle feedbacks in scenarios with negative emissions is essential for determining the effectiveness of carbon dioxide removal in drawing down atmospheric CO₂ and mitigating warming.

- Deleted:** a
- Deleted:** the magnitude of
- Deleted:** generally
- Deleted:** differs between the
- Deleted:** ramp-up
- Deleted:** and
- Deleted:** down
- Deleted:** s
- Deleted:** ese differences are
- Deleted:** likely
- Deleted:** limite system
- Deleted:** represents the
- Deleted:** both the prior positive emissions and negative emissions increasing and then
- Deleted:** concentration
- Formatted:** Subscript
- Formatted:** Subscript
- Deleted:** negative emissions
- Formatted:** Subscript
- Deleted:** reduce
- Deleted:** is inertia
- Deleted:** negative emissions
- Deleted:** This has two implications: using feedback parameters from the standard approach will (1) underestimate land and ocean carbon release under negative emissions due to changes in CO₂ concentration alone (concentration-carbon feedback), and (2) underestimate land and ocean carbon gain due to changes in climate alone (climate-carbon feedback). Given that the concentration-carbon feedback is the dominant feedback, q
- Deleted:** ing
- Deleted:** under
- Deleted:** important
- Deleted:** with the standard approach will result in the underestimation of land and ocean carbon loss under negative emissions, thereby overestimating
- Formatted:** Not Highlight
- Formatted:** Not Highlight
- Formatted:** Not Highlight
- Deleted:** negative emissions

1 Introduction

Anthropogenic CO₂ emissions have increased substantially since the preindustrial era, increasing the risk of “severe, pervasive and irreversible impacts” to the Earth system (IPCC, 2022). In an effort to reduce greenhouse gas emissions, nations adopted the Paris Agreement, which stipulated that surface warming should be kept well below 2°C above preindustrial levels and encouraged efforts to further limit it to 1.5°C (UNFCCC, 2015). Carbon dioxide removal (CDR) is a key mitigation measure in emission scenarios that are consistent with these climate goals (Ciais et al., 2013; Fuss et al., 2014; Rogelj et al., 2018; Rogelj et al., 2019; IPCC, 2022).

The land and ocean carbon sinks play a major role in regulating atmospheric CO₂ concentration by absorbing approximately half of current anthropogenic CO₂ emissions (Friedlingstein et al., 2022). However, this rate of absorption is sensitive to changes in climate and atmospheric CO₂ concentration (Cox et al., 2000; Boer & Arora, 2010; Arora et al., 2013; Boer & Arora, 2013; Arora et al., 2020). As atmospheric CO₂ concentration increases, carbon sinks will take up more carbon through air-sea exchange and CO₂ fertilization, resulting in a negative concentration-carbon cycle feedback (Boer & Arora, 2010; Arora et al., 2013; Schwinger & Tjiputra, 2018). Conversely, changing climate, in response to the increasing CO₂ concentration, will decrease the ability of carbon sinks to take up carbon, resulting in a positive climate-carbon cycle feedback (Cox et al., 2000; Jones et al., 2003; Fung et al., 2005; Friedlingstein et al., 2006; Boer & Arora, 2010; Zickfeld et al., 2011; Boer & Arora, 2013; Friedlingstein et al., 2014; Schwinger & Tjiputra, 2018).


Since the dominant feedback controlling land and ocean carbon uptake is the negative concentration-carbon feedback, the land and ocean are currently carbon sinks (Arora et al., 2020). However, the implementation of negative emissions is expected to weaken or even reverse natural carbon sinks. If negative emissions are implemented but remain lower than positive emissions (net-positive emissions), the land and ocean carbon sinks continue to take up carbon, albeit at a lower rate (Tokarska & Zickfeld, 2015; Jones et al., 2016; Melnikova et al. 2021, Koven et al., 2022). On land, the rate of carbon uptake declines because ecosystem respiration increases more than gross primary productivity increases, whereas, in the ocean, the rate of uptake declines following the declining CO₂ emissions growth rate (Melnikova et al., 2021). Once the amount of CO₂ removed from the atmosphere exceeds the amount of CO₂ added to the atmosphere (net-negative emissions), the carbon sinks are expected to weaken further and may reverse (Cao & Caldeira, 2010; Tokarska & Zickfeld, 2015; Jones et al., 2016; Melnikova et al., 2021; Canadell et al., 2022; Koven et al., 2022). Decreasing CO₂ levels will weaken the CO₂ fertilization effect, decreasing net primary productivity (NPP) more than soil respiration, resulting in a flux of carbon into the atmosphere (Cao & Caldeira, 2010; Tokarska & Zickfeld, 2015). Furthermore, the gradient in the partial pressure of CO₂ at the atmosphere-ocean interface will weaken and eventually reverse, resulting in the outgassing of CO₂ (Cao & Caldeira, 2010; Tokarska & Zickfeld, 2015). Carbon losses from the land and ocean following CDR are expected to significantly decrease the effectiveness of CDR in drawing down atmospheric CO₂ (Tokarska & Zickfeld, 2015; Jones et al., 2016; Zickfeld et al., 2021).

Deleted:

Deleted:

100 The behaviour of land carbon cycle feedbacks under positive and negative emissions is shown qualitatively in Figure 1. As the atmospheric CO₂ concentration increases under positive emissions, the land sequesters more carbon, reducing the atmospheric CO₂ concentration (Boer & Arora, 2010; Arora et al., 2013). However, under negative emissions, the declining atmospheric CO₂ concentration weakens and eventually reverses the land carbon sink, returning CO₂ to the atmosphere. The concentration-carbon feedback is negative because it promotes carbon sequestration under positive emissions and drives carbon loss under negative emissions. As the climate warms under positive emissions, the land loses carbon to the atmosphere, increasing the atmospheric CO₂ and causing further warming (Cox et al., 2000; Jones et al., 2003; Fung et al., 2005; Friedlingstein et al., 2006; Boer & Arora, 2010; Zickfeld et al., 2011; Boer & Arora, 2013; Friedlingstein et al., 2014). With cooling, the land carbon source weakens and eventually turns into a carbon sink, sequestering carbon and further cooling the climate under negative emissions. This positive climate-carbon feedback acts to amplify warming under positive emissions and enhance cooling under negative emissions.

Deleted:



110 **Figure 1: Carbon cycle feedback schematic illustrating the behaviour of the negative concentration-carbon feedback (top box) and positive climate-carbon feedback (bottom box). Each feedback loop starts with an increase (under positive emissions) or decrease (under negative emissions) in atmospheric CO₂ concentration or surface air temperature. Arrows indicate a positive coupling (change in the same direction) between components and lines with empty circles indicate a negative coupling (change in the opposite direction) between components.**

115

The goal of this study is to quantify land carbon cycle feedbacks under negative emissions. We address two research questions:

- (1) How does the magnitude of carbon cycle feedbacks under negative emissions compare to that under positive emissions?
- (2) Is the approach currently used to quantify carbon cycle feedbacks under positive emissions adequate to quantify feedbacks

120 under negative emissions? If not, how can this approach be improved upon? This study investigates carbon cycle feedbacks
under positive and negative emissions in an Earth system model of intermediate complexity (EMIC) driven with an idealized
scenario with a 1% per year increase and decrease in atmospheric CO₂ concentration. Our study adds to the small but growing
body of research on carbon cycle feedbacks under negative emissions (Schwinger & Tjiputra, 2018; Melnikova et al., 2021)
by exploring the behaviour of these feedbacks, with a focus on land processes. We propose a novel approach for quantifying
125 carbon cycle feedbacks under negative emissions and provide insight into the role of these feedbacks in determining the
effectiveness of carbon dioxide removal in reducing CO₂ levels.

2 Methodology

2.1 Model Description

The University of Victoria Earth System Climate Model (UVic ESCM, version 2.10) (**figure 2**) is a model of intermediate
130 complexity with a horizontal grid resolution of 1.8° (meridional) x 3.6° (zonal) (Weaver et al., 2001; Mengis et al., 2020). The
model consists of a simplified atmospheric model, a 3D ocean general circulation model, including ocean inorganic and organic
carbon cycle models, coupled to a dynamic-thermodynamic sea ice model, and a land surface model coupled to a vegetation
model (including permafrost) (Mengis et al., 2020). The atmosphere is a 2D energy-moisture balance model with dynamical
wind feedbacks. Atmospheric heat and freshwater are transported through diffusion and advection (Weaver et al., 2001), based
135 on wind velocities prescribed from monthly climatological wind fields from NCAR/NCEP reanalysis data (Eby et al., 2013).
The 19-layer 3D ocean general circulation model is based on the Geophysical Fluid Dynamics Laboratory (GFDL) Modular
Ocean Model Version 2 (MOM2) (Pacanowski, 1995). The coupled dynamic-thermodynamic sea ice model simulates sea ice
dynamics through elastic, viscous and plastic deformation and flow mechanisms (Weaver et al., 2001). Ocean carbon is
represented by an inorganic ocean carbon model following the Ocean Carbon Model Intercomparison Protocol (OCMIP), and
140 a NPZD (nutrient, phytoplankton, zooplankton, detritus) model of ocean biology simulating carbon uptake by the biological
pump, accounting for phytoplankton light and iron limitations (Keller et al., 2012). The land surface model, based on the
Hadley Centre Met Office Surface Exchange Scheme (MOSES), simulates the terrestrial carbon cycle and is coupled to the
Top-Down Representation of Interactive Foliage and Flora including Dynamics (TRIFFID) model which simulates vegetation
and soil carbon (Meissner et al., 2003). This model version also includes a permafrost carbon model in the soil module that
145 simulates permafrost carbon through a diffusion-based scheme (MacDougall & Knutti, 2016).




Figure 2: University of Victoria Earth System Climate Model (UVic ESCM) schematic. Energy, water and carbon exchanges between model components are represented by arrows. Figure reproduced with permission from Mengis et al. (2020).

150 2.2 Model Simulations

We performed a preindustrial spin-up simulation to equilibrate the model with the preindustrial CO₂ concentration (~285ppm). All other greenhouse gas concentrations, surface land conditions and orbital parameters were held at 1850 levels according to the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design protocol (Eyring et al., 2016). The solar forcing was set to the 1850 – 1873 mean and the volcanic forcing was held at its average over 1850 – 2014, also consistent with CMIP6 protocol (Eyring et al., 2016).

To explore how the magnitude of carbon cycle feedbacks under positive emissions differs from that under negative emissions, we ran the “CDR-reversibility” simulation from the Carbon Dioxide Removal Model Intercomparison Project (CDRMIP) (Keller et al., 2018). Starting from a preindustrial equilibrium state, atmospheric CO₂ concentration was prescribed to increase at 1% per year until quadrupling, then decline back to preindustrial levels at the same rate. [Achieving such a rapid decline in CO₂ concentration would only be possible with substantial negative CO₂ emissions \(Boucher et al., 2012\).](#) We refer to the section of the prescribed CO₂ concentration trajectory with increasing CO₂ concentration as the ramp-up phase and the section with decreasing CO₂ concentration as the ramp-down phase.

We also ran a zero emissions simulation (“Zeroemit”) for use in our novel approach for quantifying the “committed” carbon cycle [response to increasing atmospheric CO₂ during the ramp-up phase](#). This simulation was initialized from the peak

Formatted: Subscript

Deleted:

Deleted: feedbacks under negative emissions

Formatted: Subscript

170 atmospheric CO₂ concentration in the “CDR-reversibility” simulation and run in emissions-driven configuration. Emissions were set to zero at the start of the simulation, then CO₂ was allowed to evolve for 500 years.

The “CDR-reversibility” and “Zeroemit” simulations were run in three modes, following the C4MIP protocol for the quantification of carbon cycle feedbacks (Friedlingstein et al., 2006; Arora et al., 2013; Jones et al., 2016; Arora et al., 2020):

- 175 1. Fully coupled mode (FULL): the entire Earth system responds to the specified change in atmospheric CO₂ concentration or CO₂ emissions. In this mode, the land and ocean carbon sinks are subject to changing atmospheric CO₂ concentration and temperature.
2. Biogeochemically coupled mode (BGC): the land and ocean carbon sinks are subject to changing atmospheric CO₂ concentration but not changing temperature. This is achieved by prescribing a specified time-invariant CO₂ concentration to the radiation module (preindustrial CO₂ concentration for the “CDR-reversibility” simulation and quadruple the preindustrial CO₂ concentration for the “Zeroemit” simulation), while the land and ocean carbon cycle modules see an evolving atmospheric CO₂ concentration.
- 180 3. Radiatively coupled mode (RAD): the land and ocean carbon sinks are subject to changes in temperature but no change in atmospheric CO₂ concentration. The land and ocean carbon cycle modules see a specified time invariant CO₂ concentration (preindustrial CO₂ concentration in the “CDR-reversibility” simulation and quadruple the preindustrial CO₂ concentration in the “Zeroemit” simulation), while the radiation module sees changing atmospheric CO₂ concentration.

185 In both the “CDR-reversibility” and “Zeroemit” simulations, non-CO₂ forcings are held fixed at their preindustrial values.

2.3 Approaches to Carbon Cycle Feedback Quantification

190 In the first approach (referred to as the “standard” approach), we use the “CDR-reversibility” simulation to quantify carbon cycle feedbacks under increasing and decreasing atmospheric CO₂ concentration. Although this simulation is highly idealized, the ramp-up phase is standardly used to quantify carbon cycle feedbacks under positive emissions, and therefore, allows easier comparison of these results to other literature. The ramp-up phase represents the response to increasing atmospheric CO₂ alone. However, the ramp-down phase represents the response to both the prior increasing CO₂ and decreasing CO₂ because the latter is prescribed when the system is still in a transient (that is, time-evolving) state, responding to the prior atmospheric CO₂ increase (Zickfeld et al., 2016; Keller et al., 2018). As a result, carbon cycle feedbacks quantified from the ramp-down phase do not represent the response to decreasing atmospheric CO₂ alone.

200 Our second and novel approach, therefore, aims to improve the quantification of carbon cycle feedbacks under decreasing CO₂ by isolating the carbon cycle response to decreasing CO₂ alone. We use an experimental design utilizing both the “CDR-reversibility” and “Zeroemit” simulations. Since the “Zeroemit” simulation quantifies the “committed” or jagged response to

Formatted: ... [1]

Deleted: t

Deleted: climate

Deleted: The radiation module sees changing atmospheric CO₂ concentration.

Deleted: alone... This is achieved by prescribing a e radiation module stays fixedsees ... [2]

Deleted: stays fixed

Deleted: sees

Deleted: at the...CO₂ level ... [3]

Deleted: from which the simulation is initialized i.e.,

Formatted: Subscript

Deleted: climate

Deleted: alone... The land and ocean carbon cycle modulesinks...see a specified time invariantfixed...CO₂ concentration (: ...reindustrial CO₂ concentration in the “CDR-reversibility” simulation and quadruple the preindustrial CO₂ concentration in the “Zeroemit” simulation), while t. T ... [4]

Formatted: Subscript

Formatted: Subscript

Formatted: Normal, No bullets or numbering

Deleted:

Deleted: positive and negative emissions... Although this simulation is highly idealized, the ramp-up phase is standardly used to quantify carbon cycle feedbacks under positive emissions, and therefore, allows easier comparison of these results to other literature. The ramp-up phase represents the response to positive ... [5]

Formatted: Subscript

Deleted: emissions

Formatted: Subscript

Deleted: positive emissions

Deleted: negative emissions

Deleted: ...cause the negative emissions are ... [6]

Formatted: Subscript

Formatted: Subscript

Deleted: CO₂ decrease...is prescribed when the system is st ... [7]

Deleted:

Formatted: Subscript

Deleted: to negative emissions

Formatted: Subscript

Formatted: Subscript

Deleted: negative emissions ...y isolating the carbon cycle ... [8]

Deleted: “...agged” ... [9]

270 the prior positive emissions, the first 140 years of this simulation was subtracted from the ramp-down phase of the “CDR-reversibility” simulation to isolate the response to **decreasing CO₂ alone**. A similar approach was used in Zickfeld et al. (2016) to **quantify the temperature response to decreasing atmospheric CO₂**. The main assumption made here is that of linearity, that is, we assume that the committed carbon cycle response to the prior **CO₂ increase** and the carbon cycle response to **CO₂ decrease** combine linearly to the total carbon cycle response in the ramp-down phase. From our approach – referred to as the “**inertia corrected**” approach – we quantify carbon cycle feedbacks and compare them to those from the first approach.

2.4 Carbon Cycle Feedback Metrics

We use integrated flux-based feedback parameters (Friedlingstein et al., 2006) to quantify carbon cycle feedbacks in both approaches. **In this framework, changes in land and ocean carbon are** expressed as the sum of two terms: a term representing the change in land (ocean) carbon in response to changes in atmospheric CO₂, and a term representing the change in land (ocean) carbon in response to changes in surface air temperature:

$$\Delta C_x = \beta_x \Delta C_A + \gamma_x \Delta T \quad [1]$$

285 **with the subscript X representing land or ocean**. The concentration-carbon feedback parameter β quantifies the carbon cycle response to changes in CO₂ concentration in units of PgC ppm⁻¹, whereas the climate-carbon feedback parameter γ quantifies the carbon cycle response to changes in climate in units of PgC °C⁻¹.

290 The change in land (ocean) carbon due to **changing** atmospheric CO₂ concentration is determined using the biogeochemically coupled (BGC) simulation. In this simulation, the land and ocean only respond to changes in the CO₂ concentration, and therefore, this simulation can be used to quantify the concentration-carbon feedback parameter β . Warming is still observed in these simulations because the water use efficiency of vegetation increases at higher CO₂ concentrations and changes in albedo due to shifts in vegetation structure and spatial distribution, result in a small warming effect (Cox et al., 2004, Boer & Arora, 2013; Arora et al., 2013). However, this warming is considered negligible **in this framework (Friedlingstein et al., 2006)**. Assuming that $\Delta T = 0$ in Eq. [1], the change in land (ocean) carbon due to changes in atmospheric CO₂ concentration is expressed as:

$$\Delta C_x = \beta_x \Delta C_A \quad [2]$$

300 Equation [2] can then be rearranged to solve for the concentration-carbon feedback parameter, β as follows:

- Deleted: negative emissions
- Formatted: Subscript
- Deleted: positive emissions
- Formatted: Subscript
- Formatted: Subscript
- Deleted: negative emissions
- Deleted: Ramp-down – Zeroemi
- Deleted: t
- Deleted: Framework
- Deleted: , under both positive and negative emissions
- Deleted: Using the index X to represent
- Deleted: The total change in
- Deleted: land or
- Deleted: (
- Deleted: ocean
- Deleted:) carbon
- Deleted: the
- Deleted: (
- Deleted:)
- Deleted: is
- Deleted: L
- Deleted: L
- Deleted: L
- Deleted: [
- Deleted: $\Delta C_o = \beta_o \Delta C_A + \gamma_o \Delta T$ [2]
- Deleted: the increasing
- Deleted: in this feedback framework
- Deleted: (
- Deleted:)
- Deleted: and (2),
- Deleted: L
- Deleted: L
- Deleted: [
- Deleted: 3a
- Deleted: $\Delta C_o = \beta_o \Delta C_A$ [4a]
- Deleted: s
- Deleted: (3a) and (4a)
- Deleted:

$$\beta_x = \frac{\Delta C_x}{\Delta C_o} \quad [3]$$

Deleted: L

Deleted: L

Deleted: [3b] $\beta_o = \frac{\Delta C_o}{\Delta C_A}$

Deleted: 4b

The change in land (ocean) carbon due to climate change is determined using the radiatively coupled (RAD) simulation. In this simulation, the land and ocean only respond to changes in climate, and therefore, this simulation can be used to quantify the climate-carbon feedback parameter γ . The change in land (ocean) carbon due to climate change is expressed as:

$$\Delta C_x = \gamma_x \Delta T \quad [4]$$

Deleted: L

Deleted: L

Deleted: [

Deleted: 5a

Deleted: $\Delta C_o = \gamma_o \Delta T$ [6a]

Deleted: ns (5a) and (6a)

Deleted:

Deleted: L

Deleted: L

Deleted: [5b] $\gamma_o = \frac{\Delta C_o}{\Delta T}$ [

Deleted: 6b

Deleted: C

Deleted: CLCLIM

Deleted: L

Deleted: L

Deleted: L

Deleted: L

Deleted: L

Deleted: [

Deleted: 7

Deleted: $\Delta C_o^{CLIM} = \Delta C_o^{FC} - \Delta C_o^{BGC} = \gamma_o (\Delta T^{FC} - \Delta T^{BGC})$ [8]

Formatted: English (UK), Subscript

Formatted: Justified

Deleted: (

Deleted: 5b

Deleted:), (6b)

Deleted: & Tjiputra

Deleted: , 2018

Deleted: under positive emissions

Deleted: in the ramp-up phase

Formatted: Subscript

Deleted: under negative emissions

$$\gamma_x = \frac{\Delta C_x}{\Delta T} \quad [5]$$

Equation [4] can then be rearranged to solve for the climate-carbon feedback parameter γ as follows:

An alternative method for quantifying the change in land (ocean) carbon due to climate change uses the fully coupled and biogeochemically coupled simulations (Arora et al., 2013). Here, we refer to this method as the FULL-BGC method. Here, the change in land (ocean) carbon in the biogeochemically coupled simulation (BGC) is subtracted from that in the fully coupled simulation (FULL) and expressed as the product of the climate-carbon feedback parameter, and the difference between the surface air temperature changes in the two simulations:

$$\Delta C_x = \Delta C_x^{FULL} - \Delta C_x^{BGC} = \gamma_x (\Delta T^{FULL} - \Delta T^{BGC}) \quad [6]$$

Equation [6] can then be rearranged to solve for the climate-carbon feedback parameter γ as follows:

$$\gamma_x = \frac{C_x^{FULL} - \Delta C_x^{BGC}}{\Delta T^{FULL} - \Delta T^{BGC}} \quad [7]$$

The resulting feedback parameters differ from those quantified from the RAD mode (Eq. [5]) alone due to nonlinearities in carbon cycle feedbacks (Zickfeld et al., 2011; Schwinger et al., 2014).

Feedback parameters under increasing atmospheric CO₂ (ramp-up phase) are computed at the peak atmospheric CO₂ concentration (quadruple the preindustrial level) using changes in carbon pools, atmospheric CO₂ concentration and surface air temperature computed relative to preindustrial levels. Feedback parameters under decreasing atmospheric CO₂ (ramp-down

400 phase) are computed at the return to preindustrial levels (end of ramp-down phase) using changes in carbon pools, atmospheric CO₂ concentration and surface air temperature computed relative to the time of peak atmospheric CO₂.

405 In the ramp-up phase, feedback parameters are positive for land or ocean carbon gain and negative for land or ocean carbon loss. Note that the signs we refer to here are not the signs of the feedback but rather the signs of the feedback parameters, which are generally opposite to the sign of the feedback because they are computed from the perspective of the land and ocean, whereas the sign of the feedback is determined from the perspective of the atmosphere. In the ramp-down phase, both atmospheric CO₂ concentration and surface air temperature decline relative to their values at the end of the ramp-up phase, resulting in a negative denominator (see Eq. [3], [5], [7]). Therefore, the sign convention is reversed: feedback parameters are negative for a gain in land or ocean carbon (positive numerator divided by negative denominator) and positive for a loss in land or ocean carbon (negative numerator divided by negative denominator).

410 2.4.1 Isolating Carbon Cycle Feedbacks under Negative Emissions

415 When a CO₂ decrease is prescribed from a transient state, the land and ocean carbon pools not only respond to this CO₂ decrease, but also to the prior CO₂ trajectory due to inertia in these systems (Zickfeld et al., 2016). The land (ocean) carbon cycle responses in the ramp-down phase can, therefore, be expressed as the sum of two terms: one term driven by the sensitivities of land (ocean) to the CO₂ and temperature decrease during the ramp-down phase, ('SENS' for sensitivity) and an inertia term that represents the lagged response to past atmospheric CO₂ and climate changes, ('LAG');

$$\Delta C_x = \Delta C_x^{SENS} + \Delta C_x^{LAG} \quad [8]$$

420 The carbon pool response to the CO₂ and temperature decrease can then be isolated as follows:

$$\Delta C_x^{SENS} = \Delta C_x - \Delta C_x^{LAG} \quad [9]$$

425 ΔC_x^{SENS} is driven by the sensitivities to changes in atmospheric CO₂ (β) and temperature (γ) in the ramp-down phase and can be linearly decomposed in the same way as the land and ocean carbon response in the standard framework (Eq. [1]):

$$\Delta C_x^{SENS} = \Delta C_x - \Delta C_x^{LAG} = \beta_x (\Delta C_A) + \gamma_x (\Delta T) \quad [10]$$

430 Here, ΔC_A and ΔT refer to the changes in atmospheric CO₂ and temperature in the ramp-down phase of the CDR-reversibility simulation relative to their values at the end of the ramp-up phase. This framework becomes identical to the standard framework (section 2.4) in cases where a change in atmospheric CO₂ is applied from a state of equilibrium, i.e., $\Delta C_x^{LAG} = 0$.

- Deleted: in the ramp-down phase
- Deleted: Under positive emissions
- Moved (insertion) [1]
- Deleted: The signs we refer to here, however, are not the (... [10])
- Deleted: under positive emissions
- Deleted: our feedback parameters
- Deleted: Under negative emissions
- Deleted: however,
- Deleted: peak
- Deleted: values at the end of the ramp-up phase
- Formatted (... [11])
- Deleted: (...)]b)... [(4...b...)]... [(...5b...)]...and (6b)). (... [12])
- Moved up [1]: The signs we refer to here, however, are not the
- Deleted: the Response to
- Deleted: (Ramp-down – Zeroemit Approach)
- Deleted: (time-evolving)
- Deleted: representing
- Deleted: s... to the CO₂ and temperature decrease during th (... [13])
- Deleted: history
- Deleted: When negative emissions are applied from a trans (... [15])
- Formatted (... [14])
- Deleted: $\Delta C_L = \Delta C_L^{NE} + \Delta C_L^{INERTIA}$ [9] (... [16])
- Deleted: ¶
- Deleted: In this Using zero emissions simulations to quant (... [17])
- Deleted: response to
- Deleted: negative emissions
- Deleted: by taking the difference between the ramp-down (... [18])
- Deleted: for a particular mode e.g., the fully coupled mod (... [19])
- Deleted: ¶
- Deleted: DEC
- Deleted: The response to the CO₂ decrease
- Deleted: beta
- Deleted: gamma
- Deleted: alone...nd can beis then (... [22])
- Formatted (... [21])
- Deleted: into terms
- Deleted: [2]
- Deleted: representing the response to changes in CO₂ conc (... [24])
- Formatted (... [23])
- Deleted: ¶
- Deleted: DCA and DT
- Formatted (... [26])
- Deleted: 4...41 (... [27])
- Deleted: DC*LAG=0
- Formatted (... [28])

Eq. [10] can be rewritten for the biogeochemically ($\Delta T = 0$) and radiatively coupled simulations ($\Delta C_A = 0$), respectively:

$$\Delta C_X^{SENS} = \Delta C_X - \Delta C_X^{LAG} = \beta_X (\Delta C_A) \quad [11]$$

$$\Delta C_X^{SENS} = \Delta C_X - \Delta C_X^{LAG} = \gamma_X (\Delta T) \quad [12]$$

Rearranging the equations above allows for the calculation of the feedback parameters, which measure the sensitivity of the land and ocean carbon response to changes in CO₂ concentration and temperature in the ramp-down phase:

$$\beta_X = \frac{\Delta C_X - \Delta C_X^{LAG}}{\Delta C_A} \quad [13]$$

$$\gamma_X = \frac{\Delta C_X - \Delta C_X^{LAG}}{\Delta T} \quad [14]$$

The lagged responses of land and ocean carbon pools ΔC_X^{LAG} are calculated from the “Zeroemit” simulations run in the respective mode (biogeochemically coupled for the calculation of β and radiatively coupled for the calculation of γ), and are then subtracted from the responses of the ramp-down phase of the CDR-reversibility simulations run in the same mode. The land (ocean) carbon changes, surface air temperature and CO₂ concentration changes are computed relative to the year of peak CO₂ concentration (year 140 in the CDR-reversibility simulation; year 1 in the zero emissions simulation).

3 Results

3.1 “CDR-reversibility” Carbon Cycle Feedback Analysis

Our results focus on the ramp-down phase of the “CDR-reversibility” simulation and compare the system response in this phase to that in the ramp-up phase. While the prescribed atmospheric CO₂ concentration for the “CDR-reversibility” simulations is the same, the temperature response differs by mode (figure 3(a, b)). In the FULL and RAD modes, surface air temperature increases approximately linearly with increasing atmospheric CO₂ concentration, continues to increase for approximately half a decade after atmospheric CO₂ concentration peaks, then decreases with decreasing CO₂ concentration. Surface air temperature declines more slowly in the ramp-down phase due to the thermal inertia of the ocean, and therefore, does not return to preindustrial levels by the end of the ramp-down phase. The temperature response in the FULL mode is consistent with earlier studies (Boucher et al., 2012; Zickfeld et al., 2016; MacDougall, 2019; Ziehn et al., 2020; Park & Kug, 2022). Surface air temperature in the BGC mode changes only marginally: surface air temperature increases slightly with increasing CO₂ concentration and decreases as the CO₂ concentration decreases. This temperature change is driven by

- Deleted: s
- Deleted: 3
- Deleted: [14]
- Deleted: This
- Deleted: (DT=0)
- Deleted: (DCA=0)
- Deleted: as follows
- Deleted: ¶
- Deleted: $\Delta C_L^{NE} = \Delta C_L - \Delta C_L^{INERTIA} = \beta_L (\Delta C_A)$ [23a]¶
- Deleted: $\Delta C_O^{NE} = \Delta C_O - \Delta C_O^{INERTIA} = \beta_O (\Delta C_A)$ [24a]¶
- Deleted: $\Delta C_L^{NE} = \Delta C_L - \Delta C_L^{INERTIA} = \gamma_L (\Delta T)$ [25a]¶
- Deleted: $\Delta C_O^{NE} = \Delta C_O - \Delta C_O^{INERTIA} = \gamma_O (\Delta T)$ [26a]¶
- Deleted: ¶
- Deleted: The feedback parameters are then computed by r
- Deleted: to the CO₂ decrease alone
- Deleted: as follows
- Deleted: L
- Deleted: CL
- Deleted: L
- Deleted: INERTIA
- Deleted: CA
- Deleted: [23b] $\beta_O = \frac{\Delta C_O - \Delta C_O^{INERTIA}}{\Delta C_A}$
- Deleted: 2
- Deleted: 4b
- Deleted: L
- Deleted: ΔCL
- Deleted: L
- Deleted: INERTIA
- Deleted: T
- Deleted: [25b] $\gamma_O = \frac{\Delta C_O - \Delta C_O^{INERTIA}}{\Delta T}$
- Deleted: 26b
- Deleted: s

biophysical responses to changing atmospheric CO₂, in particular, changes in evaporative fluxes as plants adjust stomatal conductance based on atmospheric CO₂ levels. Biophysical effects are also responsible for the difference in warming between the FULL and RAD modes (Arora et al., 2020). The temperature response in the ramp-up phase of the FULL, BGC and RAD modes is consistent with Arora et al. (2020) while the temperature response in the ramp-up and ramp-down phases of all three modes is consistent with Schwinger & Tjiputra (2018).





Figure 3: a. Prescribed atmospheric CO₂ concentration anomaly b. surface air temperature change (SAT) c. atmosphere to land CO₂ flux and d. land e. vegetation and f. soil carbon changes in the fully coupled (FULL), biogeochemically coupled (BGC) and radiatively coupled (RAD) “CDR-reversibility” simulations. Panels a, b, and d - f are calculated relative to 1850 (preindustrial). Carbon fluxes for the three modes are shown in the bottom panels (g, h, i). NPP = net primary productivity, LLF = leaf litter flux and SR = soil respiration. Solid lines represent the ramp-up phase and dot-dashed lines represent the ramp-down phase. The vertical dotted lines mark the beginning and end of the ramp-down phase.

3.1.1 Land Carbon Change in the FULL Mode

Figure 3(d) shows land carbon changes as a function of time. In the FULL mode, land carbon increases, stabilizes, then begins to decrease 7 years before the peak atmospheric CO₂ concentration is reached. Similar carbon change patterns are observed for the soil carbon pool, which starts decreasing roughly 20 years before the peak in atmospheric CO₂ concentration, but vegetation carbon decreases 2 years after the peak atmospheric CO₂ concentration (figure 3(e, f)). Our results are qualitatively consistent with Ziehn et al. (2020). However, they differ from other studies (MacDougall, 2019; Arora et al., 2020) wherein the land carbon pool remains a carbon sink in the ramp-up phase. MacDougall (2019) shows that the soil carbon sink switches



Deleted:

Deleted: carbon

Formatted: Subscript

Deleted: the

Deleted: gains

Deleted: at a decreasing rate,

Deleted: slowly lose

Deleted: carbon

Deleted:

Deleted: uptake and loss

Deleted: losing carbon

Deleted: loss begins

into a source later in the ramp-up phase than our results show. Furthermore, other studies (Boucher et al., 2012; Zickfeld et al., 2016) show that both vegetation and soil carbon sinks persist throughout the ramp-up phase.

745

Here, ~~land carbon decreases~~ throughout the ramp-down phase (**figure 3(d)**) whereas, earlier studies show continued ~~increase in the~~ land carbon ~~pool~~ in the early ramp-down phase (Boucher et al., 2012; Zickfeld et al., 2016; Park & Kug, 2021). Changes in land carbon are governed by the balance between net primary productivity (NPP) and soil respiration. ~~The increase in the~~ ~~land carbon pool~~ is driven by the CO₂ fertilization effect: photosynthesis is enhanced under increasing CO₂ concentration, increasing NPP (**figure 3(g)**) (Arora et al. 2013). Soil respiration also increases with warming (**figure 3(g)**). Initially, soil respiration remains below NPP, but the rate of increase of NPP declines faster and soil respiration exceeds NPP towards the end of the ramp-up phase. This occurs due to the different response timescales of NPP and soil respiration: NPP depends on atmospheric CO₂ changes, whereas soil respiration depends on temperature change, which lags behind the change in CO₂ concentration (Cao & Caldeira, 2010). In the ramp-down phase, NPP decreases as the CO₂ fertilization effect weakens, whereas soil respiration continues to increase for a year before decreasing at a slower rate than NPP, driven by decreasing surface air temperature and soil carbon.

750

755

3.1.2 Land Carbon Change in the BGC Mode

In the BGC mode, ~~land carbon increases~~ in the ramp-up phase, ~~continues to increase~~ until 16 years after the peak in CO₂ concentration, then ~~decreases~~ (**figure 3(d)**). A similar lag is observed for both vegetation and soil carbon pools, but the soil carbon sink persists for five years longer than the vegetation carbon sink (**figure 3(e, f)**). ~~Land carbon increases~~ in the ramp-up phase due to the CO₂ fertilization effect, which increases NPP (**figure 3(h)**) (Arora et al. 2013). In the UVic ESCM, soil respiration depends on soil temperature, moisture, and carbon content (Cox et al., 2001; Mengis et al., 2020). Since changes in surface air temperature in the BGC mode are small (**figure 3(b)**), changes in the first two factors are negligible and soil carbon content is the main driver of soil respiration changes. Soil respiration increases with increasing soil carbon, but NPP remains higher, resulting in ~~an increase in the land carbon pool~~ in the ramp-up phase (**figure 3(h)**). In the ramp-down phase, NPP decreases as the CO₂ fertilization effect weakens, whereas soil respiration continues to increase before decreasing at a slower rate than NPP, following changes in soil carbon (**figure 3(h)**). NPP declines below soil respiration, and ~~land carbon begins to decrease~~.

760

765

3.1.3 Land Carbon Change in the RAD Mode

~~Land carbon decreases~~ in the ramp-up phase of the RAD mode, ~~continues to decrease~~ until roughly 30 years after the peak in atmospheric CO₂ concentration, then switches into a carbon sink (**figure 3(d)**). Both vegetation and soil carbon pools exhibit a similar lag, but the vegetation carbon pool remains a carbon source for a decade longer than the soil carbon pool (**figure 3(e, f)**). ~~Land carbon decreases~~ in the ramp-up phase because NPP decreases as plant respiration rates increase (see **figure S1**), whereas soil respiration increases with warming (**figure 3(i)**) consistent with earlier literature (Arora et al., 2020). NPP later

770

Deleted: the
Deleted: loses
Deleted: uptake
Deleted: followed by land carbon loss
Deleted: Carbon gain

Deleted: the
Deleted: sequesters
Deleted: remains a carbon sink
Deleted: switches into a source of carbon
Deleted: The land sequesters carbon
Deleted:

Deleted: land carbon uptake

Deleted: the
Deleted: switches into a carbon source

Deleted: The land loses carbon
Deleted: remains a carbon source

Deleted: The land loses carbon

increases due to vegetation shifts that occur on decadal to centennial timescales (see figure S2) but remains lower than soil respiration. In the ramp-down phase, NPP increases (figure 3(i)) as gross primary productivity increases and plant respiration decreases with cooling, then later declines as gross primary productivity declines, because cooler temperatures negatively impact vegetation growth in the high latitudes (see figures S1, S3). Soil respiration decreases steadily with declining surface air temperature, and after a few decades, declines below NPP, and the land carbon pool begins to grow again.

Deleted: switches into a carbon sink

3.1.4 Ocean Carbon Change in the FULL, BGC and RAD Modes

In the FULL mode, the ocean carbon pool grows at a steady rate, then begins to slowly lose carbon roughly three decades after the peak in atmospheric CO₂ concentration (figure 4(a)). In the ramp-up phase, the partial pressure of CO₂ in the atmosphere increases, strengthening the partial pressure gradient and driving an influx of CO₂ into the ocean (figure 4(b)). In the ramp-down phase, the gradient in partial pressure weakens and eventually reverses, and the ocean carbon sinks switches into a source. Earlier studies forced with the “CDR-reversibility” simulation also show ocean carbon uptake in the ramp-up phase (MacDougall, 2019; Arora et al., 2020) followed by delayed carbon loss in the ramp-down phase (Boucher et al., 2012; Zickfeld et al., 2016).

Deleted: gains

The ocean exhibits a delayed response in the ramp-down phase of the BGC and RAD modes consistent with Schwinger & Tjiputra (2018). In the BGC mode, ocean carbon increases in the ramp-up phase, continues to increase for approximately half a century after the peak atmospheric CO₂ concentration, then switches into a source of carbon (figure 4(a)). The partial pressure gradient of CO₂ strengthens in the ramp-up phase, driving CO₂ uptake, then weakens and reverses in the ramp-down phase, promoting carbon loss, but the magnitude of the flux is larger than in the FULL mode (figure 4(b)). In the RAD mode, ocean carbon decreases in the ramp-up phase, continues to decrease for over a century in the ramp-down phase, then switches into a weak carbon sink (figure 4(a)). The ocean outgasses in the ramp-up phase possibly due to climate effects on ocean circulation and the solubility pump (Cox et al., 2000; Fung et al., 2005; Friedlingstein et al., 2006; Zickfeld et al., 2011). In the ramp-down phase, the ocean remains a carbon source for over a century before switching into a weak carbon sink. Ocean carbon changes in the BGC and RAD modes are also driven by the concentration-carbon and climate-carbon feedbacks. An in-depth discussion of the mechanisms behind the ocean carbon response is beyond the scope of this paper.

Formatted: Justified

Deleted: the

Deleted: takes up

Deleted:


Deleted: remains a carbon sink


Deleted: the

Deleted: loses

Deleted: remains a carbon source

Deleted: ¶





830 Figure 4: a. Ocean carbon change and b. atmosphere to ocean CO_2 flux in the fully coupled (FULL), biogeochemically coupled (BGC) and radiatively coupled (RAD) “CDR-reversibility” simulations. Ocean carbon change is calculated relative to 1850 (preindustrial). Solid lines represent the ramp-up phase and dot-dashed lines represent the ramp-down phase. The vertical dotted lines mark the beginning and end of the ramp-down phase.

Deleted: carbon
Formatted: Subscript

3.1.5 Sensitivity of Land and Ocean Carbon Pools

To assess the sensitivity of land and ocean carbon pools to changes in atmospheric CO_2 and temperature, we plot carbon changes in the BGC mode as a function of atmospheric CO_2 concentration (figure 5) and carbon changes in the RAD mode as a function of surface air temperature (figure 6). The trajectory of carbon change differs in the ramp-up and ramp-down phases of the BGC mode (figure 5), a behavior referred to as hysteresis. Hysteresis in the land carbon pool is primarily driven by the soil carbon pool, although the contribution from the vegetation carbon pool is also significant (figure 5(a, c, d)). The width of the hysteresis – measured as the vertical distance between the ramp-up and ramp-down trajectories – initially increases, then
840 decreases (figure 5(a - d)), except in the vegetation carbon pool where the width of the hysteresis increases throughout the ramp-down phase (figure 5(c)). The land and ocean carbon pools in the RAD mode also exhibit hysteresis (figure 6). The hysteresis in the land carbon pool is dominated by the soil carbon pool (figure 5(d)), and the width of the hysteresis appears to increase throughout the ramp-down phase for all carbon pools except the vegetation carbon, which shows nearly constant hysteresis. The observed hysteresis in the land and ocean carbon pools in the BGC and RAD modes is likely largely due to
845 climate system inertia: the carbon cycle response in the ramp-down phase is a combination of the response to both increasing and decreasing CO_2 concentrations.


Deleted: under negative emissions, that is,
Deleted: ,
Deleted: negative
Deleted: emissions and the prior positive emissions
Formatted: Subscript

Despite the restoration of preindustrial atmospheric CO_2 levels in the BGC mode, the land and ocean carbon pools do not return to their preindustrial states. At the end of the ramp-down phase, the land carbon pool holds approximately 250 PgC more than at preindustrial, with 80 PgC remaining in vegetation and 170 PgC remaining in the soil (figure 5(a, c, d), due to
850 time lags associated with vegetation and soil carbon turnover. The ocean carbon pool holds much more carbon (615PgC) than at preindustrial (figure 5(b)). In the RAD mode, the land and ocean carbon lost in the ramp-up phase is not completely regained

Deleted: whereas t
Deleted:

860 in the ramp-down phase, though this response would be expected given the asymmetric surface air temperature response in
 this mode. By the end of the RAD mode, the land carbon pool holds approximately 300 PgC less than at preindustrial, with
 the vegetation carbon pool accounting for 70 PgC and the soil carbon pool accounting for the remaining 230PgC (figure 6(a,
 c, d)). The ocean holds only 70PgC less than at preindustrial, but unlike the land carbon pool, a miniscule amount of ocean
 carbon is regained in the ramp-down phase (figure 6(b)).

865 Previous studies have shown carbon cycle hysteresis in the FULL mode of the “CDR-reversibility” simulation (Boucher et al.,
 2012; Zickfeld et al., 2016; Jeltsch-Thömmes et al., 2020; Park & Kug, 2022), consistent with our results (see figure S4).
 However, in most of these studies, the vegetation and soil carbon pools do not return to their preindustrial states by the end
 of the ramp-down phase (Boucher et al., 2012; Zickfeld et al., 2016; Park & Kug, 2022). Our results for the FULL mode of the
 870 “CDR-reversibility” simulation show that the vegetation and soil carbon pools are very close to their preindustrial states by
 the end of the ramp-down phase (see figure S4), consistent with Ziehn et al. (2020), who show a near-return to the preindustrial
 state in the vegetation carbon pool.




875 **Figure 5:** a. Land b. ocean c. vegetation and d. soil carbon pool changes as a function of atmospheric CO₂ concentration, taken from
 the biogeochemically coupled (BGC) “CDR-reversibility” simulation ramp-up and ramp-down phases, and “inertia corrected”
 approach. All values are calculated relative to 1850 (preindustrial).

Deleted: not

Deleted: b,

Deleted: 5



Deleted:

Formatted: Caption, Left

Deleted: “inertia corrected”

Deleted: approach

Deleted:

Figure 5: a. Land b. ocean c. vegetation and d. soil carbon pool changes as a function of atmospheric CO₂ concentration, taken from the biogeochemically coupled (BGC) “CDR-reversibility” simulation ramp-up and ramp-down phases, and “ramp-down – zeroemit” simulation. All values are calculated relative to 1850 (preindustrial).




Figure 6: a. Land b. ocean c. vegetation and d. soil carbon pool changes as a function of surface air temperature change, taken from the radiatively coupled (RAD) “CDR-reversibility” simulation ramp-up and ramp-down phases, and [“inertia corrected” approach](#). All values are calculated relative to 1850 (preindustrial).

Deleted: ramp-down – zeroemit

Deleted: simulation

900

905 3.1.6 Carbon Cycle Feedback Parameters quantified from “CDR-reversibility” simulations

Table 1 shows the carbon cycle feedback parameters quantified using the Friedlingstein et al. (2006) carbon cycle feedback framework (see Section 2.4). The concentration-carbon feedback parameter (β), which quantifies the concentration-carbon feedback, is computed as the change in land or ocean carbon per unit change in atmospheric CO_2 concentration in the BGC mode. The climate-carbon feedback parameter (γ) quantifies the climate-carbon feedback as the change in land or ocean carbon per unit change in surface air temperature in the RAD mode (referred to as the RAD approach). An alternative approach to

910 quantifying the climate-carbon feedback involves taking the difference between the fully coupled and biogeochemically coupled simulations and computing the change in land or ocean carbon per unit change in surface air temperature from that difference (referred to here as the FULL-BGC approach).

Deleted:

915 In the “CDR-reversibility” simulation, the magnitudes of β and γ for both land and ocean are smaller [in the ramp-down phase](#) (under negative emissions) than [in the ramp-up phase](#) (under positive emissions), except the ocean climate-carbon feedback parameter, which is larger. (Table 1). Climate-carbon feedback parameters calculated using the FULL-BGC approach (shown in parentheses) are consistent in sign with those calculated using the RAD approach, but the magnitudes of these feedback parameters are larger (see Figure S5 for hysteresis figures for this approach). Carbon cycle feedback parameters are smaller

in the ramp-down phase because the land and ocean carbon pools show a lagged response to changes in CO₂ concentration and climate in the early ramp-down phase. In the ocean, this lagged response to changes in climate is much greater, and carbon loss continues throughout the ramp-down phase (shown by the positive ocean climate-carbon feedback parameter). As a result, feedback parameters in the ramp-down phase are underestimated. Improving this quantification could be achieved by quantifying and removing this inertia.

Deleted: under negative emissions

Deleted: under negative emissions

Deleted: under negative emissions

Simulations(s) used for calculation of feedback parameters	Positive Emissions (Ramp-up)				Negative Emissions (Ramp-down)			
	β_L	β_o	γ_L	γ_o	β_L	β_o	γ_L	γ_o
	(PgC ppm ⁻¹)		(PgC °C ⁻¹)		(PgC ppm ⁻¹)		(PgC °C ⁻¹)	
“CDR-reversibility” simulation taken at 4xCO ₂ for positive emissions and at return to preindustrial for negative emissions	0.96	0.88	-117.8 (-121.5)	-7.36 (-22.7)	0.68	0.16	-56.4 (-67)	10.8 (31.1)
“Inertia corrected” approach taken at 4xCO ₂ for positive emissions and at return to preindustrial for negative emissions	0.96	0.88	-117.8	-7.36	0.80	0.84	-157.1	-18.1

Deleted: Ramp-up – Zeroemit

Table 1: Carbon cycle feedback parameters under positive and negative emissions quantified at 4xCO₂ (quadruple the preindustrial CO₂ level) from the “CDR-reversibility” simulation and using the proposed “inertia corrected” approach. Feedback parameters for negative emissions are positive for land or ocean carbon loss and negative for land or ocean carbon gain, opposite to the sign convention for feedbacks under positive emissions. Values shown in parentheses were calculated using the FULL-BGC approach for quantifying climate-carbon feedbacks (see Eq. 7). Feedback parameters quantified from the “CDR-reversibility” simulation can also be derived from Figures 5 and 6 respectively by taking the slope of the land or ocean response at the same time points at which they are computed.

Deleted: Ramp-up – Zeroemit

Deleted: (

Deleted:) and (8)

3.2 Isolating Carbon Cycle Feedbacks under Negative Emissions

3.2.1 “Zeroemit” Simulation: Quantifying Climate System Inertia

Zero emissions simulations quantify committed changes due to the prior CO₂ trajectory. Changes in atmospheric CO₂ concentration in zero emissions simulations are driven by the carbon sinks, which in turn are influenced by the CO₂ concentration and climate. Following cessation of emissions, the CO₂ concentration in the FULL mode declines steadily, mainly driven by ocean carbon uptake consistent with results from MacDougall et al. (2020) (figure 7(a)). The CO₂ concentration in the BGC mode declines more than in the FULL mode because both land and ocean remain carbon sinks. In the RAD mode, the CO₂ concentration increases as both land and ocean carbon decrease, releasing CO₂ into the atmosphere. Changes in atmospheric CO₂ concentration, together with changes in ocean heat uptake and surface albedo, drive changes in surface air temperature. In the FULL mode, the warming effect of declining ocean heat uptake dominates over the cooling



Deleted: positive emissions

Formatted: Subscript

Deleted: e

effect of declining CO₂ concentration resulting in continued warming (MacDougall et al., 2020) (figure 7(b); figure S6). The decline in CO₂ concentration is partly offset by permafrost carbon release from the soil (figure 7(e)). Surface air temperature in the RAD mode increases more than in the FULL mode because the CO₂ concentration increases, causing further warming. Surface air temperature remains relatively constant in the BGC mode. In the FULL mode, the land switches into a source of carbon after emissions cease, consistent with the behaviour of the UVic ESCM in the Zero Emissions Commitment Model Intercomparison Project (ZECMIP) (MacDougall et al., 2020) (figure 7(c)). Vegetation carbon continues to increase (figure 7(d)) whereas soil carbon decreases (figure 7(e)). The ocean remains a carbon sink after cessation of emissions (figure 7(f)). In the BGC mode, the ocean remains a strong carbon sink after CO₂ emissions are set to zero, whereas land carbon initially increases, then decreases (figure 7(c, f)). Vegetation carbon increases throughout the zero emissions phase whereas soil carbon initially increases, then slowly decreases (figure 7(d, e)). Both land and ocean carbon decrease in the RAD mode (figure 7(c, f)) with both vegetation and soil carbon pools driving this decrease (figure 7(d, e)).

- Deleted: The v
- Deleted: pool
- Deleted: take up carbon
- Deleted: the
- Deleted: switches into a source of carbon
- Deleted: the
- Deleted: takes up
- Deleted: carbon, stabilizes,
- Deleted: becomes a weak carbon sink again
- Deleted: The v
- Deleted: pool takes up
- Deleted: carbon
- Deleted: the
- Deleted: takes up carbon
- Deleted: , stabilizes
- Deleted: releases CO₂
- Deleted: release CO₂ to the atmosphere
- Deleted: the land carbon release



- Deleted: change

Figure 7: a. Atmospheric CO₂ concentration anomaly b. surface air temperature anomaly c. land carbon change d. vegetation carbon change e. soil carbon change and f. ocean carbon change for the zero emissions simulations relative to 1850 (preindustrial). ALL = the CDR-reversibility ramp-up phase from which all modes are initialized; BGC = biogeochemically coupled, RAD = radiatively coupled and FULL = fully coupled. Solid lines are for the ramp-up phase; dashed lines are for the zero emissions phase.

995 3.2.2 “**Inertia Corrected**” Approach: Isolating the Response to Negative Emissions

The “**inertia corrected**” approach uses the zero emissions simulations described in the previous section to isolate the response to negative emissions in the “CDR-reversibility” simulations by taking the difference between the ramp-down phase of the RAD (BGC) “CDR-reversibility” simulation and the RAD (BGC) zero emissions simulation. In the BGC mode, despite our attempt to reduce climate system inertia in our novel approach, carbon pools do not return to their preindustrial states at the time atmospheric CO₂ returns to preindustrial levels (**figure 5**). In the RAD mode, all carbon pools **eventually** gain more carbon than they held at preindustrial (**figure 6**).

The “**inertia corrected**” approach removes the initial carbon increase in the “CDR-reversibility” BGC mode (**figure 5**) and removes the initial carbon decrease in the “CDR-reversibility” RAD mode (**figure 6**) reducing the width of the hysteresis. Zickfeld et al. (2016) used zero emissions to isolate the response to negative emissions and observed a reduction in the initial carbon change at the beginning of the ramp-down phase consistent with our results. **In our approach, the hysteresis may persist because of the different configurations in which the “CDR-reversibility” and “zeroemit” simulations were run, that is, that the former were run with prescribed atmospheric CO₂ concentration, whereas the latter were emissions-driven, may also impact the quantification of the inertia. Another possibility may be irreversible changes in vegetation distribution in the “CDR-reversibility” ramp-down phase that are caused by state changes rather than inertia. When the CO₂ decrease is prescribed, the earth system is in a state of elevated CO₂ concentration and surface air temperature, which may lead to a different vegetation response than to an equivalent CO₂ increase applied from a preindustrial state (Zickfeld et al., 2021). Alternatively, the remaining hysteresis may show that the linearity assumption made in this experiment is not satisfied; the linearity assumption made here is that the total carbon cycle response in the ramp-down phase is a linear combination of the committed response, following increasing CO₂ concentration and temperature, and the response driven by the decrease in atmospheric CO₂ and temperature in the ramp-down phase (see Section 2.4.1: Eq. 83).**

After isolating the response to negative emissions alone in the “**inertia corrected**” approach, the magnitudes of β_L and β_O are smaller **in the ramp-down phase** as compared to their respective magnitudes **in the ramp-up phase**, but the magnitudes of γ_L and γ_O become larger **in the ramp-down phase** (**Table 1**). **In the ramp-down phase**, the magnitudes of β and γ from our novel approach are larger compared to those from the “CDR-reversibility” simulation, implying greater land and ocean carbon loss due to changes in CO₂ concentration alone and greater land and ocean carbon gain due to changes in climate alone. For example, a decrease in atmospheric CO₂ of one ppm would result in the loss of 0.68 PgC of land carbon in the standard approach and 0.80 PgC of land carbon in our approach due to changes in CO₂ concentration alone, whereas, cooling by one degree, would result in land carbon gain of 56.4 PgC in the standard approach and almost three times as much (157.1 PgC) in our approach due to changes in climate alone.

Deleted: Ramp-down – Zeroemit

Deleted: Ramp-down – Zeroemit

Deleted: Ramp-down – Zeroemit

Deleted: held fixed at the peak value held fixed at the peak value

Deleted: One possible reason why the hysteresis

Deleted: persists may

Deleted: negative emissions are applied

Formatted: Subscript

Deleted: amount of positive emissions

Formatted: Subscript

Deleted: carbon cycle

Deleted: s

Deleted: to

Deleted: past positive emissions and the carbon cycle response to negative emissions

Deleted: and decreasing

Formatted: Subscript

Formatted: Subscript

Deleted: combine linearly to the total carbon cycle response in the ramp-down phase in this experimental design

Deleted: (

Deleted: 9

Deleted:)

Deleted: and (10), with prescribed atmospheric CO₂, the quantification of the inertia reduces uptake relative to a case with atmospheric CO₂ prescribed to decline as in the ramp-down phase relative to a case with prescribed atmospheric CO₂

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Deleted: Ramp-down – Zeroemit

Deleted:

Deleted: under negative emissions

Deleted: under positive emissions

Deleted: under negative emissions

Deleted: Under negative emissions

Deleted: under negative emissions

4 Discussion and conclusions

Our results from the “CDR-reversibility” simulation show that, due to changes in CO₂ concentration alone, carbon pools take up carbon in the ramp-up phase, continue to take up carbon in the early ramp-down phase, then switch into sources of carbon.

1060 Due to changes in climate alone, carbon pools lose carbon in the ramp-up phase, continue to lose carbon in the ramp-down phase, then switch into carbon sinks. Furthermore, the land and ocean carbon pools do not return to their preindustrial states at the end of both modes, suggesting that land and ocean carbon changes in the ramp-up phase are irreversible on centennial timescales. The differences in the magnitudes of carbon cycle feedbacks in the ramp-up and ramp-down phases, as quantified by feedback parameters, are likely largely due to climate system inertia. This inertia generally reduces the magnitude of both

1065 feedbacks [in the ramp-down phase \(under negative emissions\)](#) relative to feedbacks [in the ramp-up phase \(under positive emissions\)](#), implying reduced land and ocean carbon loss due to changes in CO₂ concentration alone and reduced land carbon gain due to the changes in climate. The exception is the ocean that continues to lose carbon [in the ramp-down phase](#), implying increased carbon loss due to changes in climate alone.

Deleted: under negative emissions

Deleted: under negative emissions

1070 To quantify the carbon cycle inertia, that is, the response to [the prior increasing CO₂ trajectory](#), we ran zero emissions simulations in fully coupled, biogeochemically coupled and radiatively coupled modes. Consistent with previous studies, the ocean continues to sequester carbon in the fully coupled zero emissions simulation (MacDougall et al., 2020). The terrestrial biosphere switches into a carbon source after emissions cease. Carbon uptake, largely by the ocean sink, decreases the atmospheric CO₂ concentration. Surface air temperature increases due to the interplay between declining CO₂ concentration and ocean heat uptake (Matthews & Caldeira, 2008; Solomon et al., 2009; Arora et al., 2013). While the carbon cycle response

1075 is consistent with the behaviour of the UVic ESCM in the Zero Emissions Commitment Model Intercomparison Project (ZECMIP) (MacDougall et al., 2020), the UVic ESCM response in ZECMIP is noticeably different from the rest of the Earth system models. On centennial times, the UVic ESCM is the only model with a positive zero emissions commitment. However, most of the other models do not represent permafrost carbon. The carbon pools in the biogeochemically coupled

Deleted: positive emissions

Formatted: Subscript

Deleted:

1080 and radiative coupled zero emissions simulations also exhibit inertia: the land and ocean [carbon pools](#) continue to [grow](#) after cessation of emissions in the biogeochemically coupled simulation, whereas both carbon pools [reduce](#) in the radiatively coupled simulation.

Deleted: sequester

Deleted: carbon

Deleted: release CO₂

Deleted: prior positive emissions and negative emissions

Deleted: (

Deleted: 9

Deleted:) and (10)

Formatted: Subscript

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Deleted: under negative emissions

Deleted: approach

1085 Assuming linearity in the response to [increasing and decreasing CO₂ concentrations](#) (see Section 2.4.1: Eq. 68), we subtract the zero emissions simulations from the “CDR-reversibility” simulations, to isolate the response to negative emissions alone. We find that [in the ramp-down phase](#), the magnitudes of β and γ from our novel approach are generally larger as compared to those from the “CDR-reversibility” simulation, implying greater land and ocean carbon loss due to changes in CO₂ concentration and greater land and ocean carbon gain due to changes in climate if feedback parameters from our approach are applied instead. Furthermore, land and ocean carbon changes in the ramp-up phase remain irreversible in our [simulations](#).

1105 A similar feedback analysis was conducted for ocean carbon cycle feedbacks using the Norwegian Earth System Model (NorESM) (Schwinger & Tjiputra, 2018). Schwinger and Tjiputra calculated ocean concentration-carbon and climate-carbon feedback parameters using the same carbon cycle feedback framework and “CDR-reversibility” simulations used here. Their results also show a lagged ocean carbon response to [the prior increasing CO₂ trajectory in the ramp-down phase](#), and as a result, the magnitude of both carbon cycle feedbacks is smaller [in the ramp-down phase than in the ramp-up phase](#).

Deleted: positive emissions

Formatted: Subscript

Deleted: under negative missions than under positive emissions

1110 We compare carbon cycle feedback parameters, quantified from the “CDR-reversibility” [ramp-up phase](#) to model means and standard deviations from CMIP5 and CMIP6 – the fifth and sixth phases of the Coupled Model Intercomparison Project – respectively (Arora et al., 2020) (see **Table S1**). The concentration-carbon feedback parameter for land (β_L) is generally consistent with those from CMIP5 and CMIP6, while the ocean concentration-carbon feedback parameter (β_O) lies slightly above the CMIP6 range ([mean \$\pm\$ 1 standard deviation](#)). The land climate-carbon feedback parameter (γ_L) lies well above the 1115 CMIP5 and CMIP6 ranges, implying a stronger sensitivity to warming relative to CMIP5 and CMIP6 models. The ocean climate-carbon feedback parameter (γ_O) [lies slightly above the ranges for](#) CMIP5 and CMIP6. We have included in the supplement feedback parameters at twice the preindustrial CO₂ concentration (2xCO₂), which are more relevant, in terms of atmospheric CO₂ levels and warming, for real-world mitigation scenarios (**Table S2**).

Deleted: under positive emissions

Deleted: simulation

Deleted: is consistent with those from

1120 We use the UVic ESCM, an EMIC, due to the number of simulations and length of model integration required in this study. Compared to comprehensive Earth system models, EMICs generally have coarser resolution and represent less Earth system processes at a lower level of detail. Moreover, the version of the UVic ESCM used here does not represent the nitrogen cycle on land and its coupling to the carbon cycle, which has ramifications for the estimated magnitude of carbon cycle feedbacks. Models without a nitrogen cycle exhibit greater land carbon gain under [increasing CO₂ concentrations](#) relative to other CMIP5 1125 and CMIP6 models, that is, the concentration-carbon feedback parameter is more positive (**Table S1**). They also exhibit greater carbon loss under [increasing CO₂ concentrations](#), that is, the climate-carbon feedback parameter is more negative. Therefore, the magnitude of both carbon cycle feedbacks in this study is generally larger under [increasing CO₂ concentrations](#), relative to other CMIP5 and CMIP6 models with a nitrogen cycle. Due to the exclusion of the nitrogen cycle, the UVic ESCM is expected to exhibit greater land carbon gain due to changes in climate alone under [decreasing CO₂ concentrations](#) relative to CMIP5 1130 and CMIP6 models with a nitrogen cycle. Nitrogen [mineralization will likely decline](#) as surface air temperature declines, reducing land carbon gain due to changes in climate alone in a model with the nitrogen cycle. The direction of land carbon change due to changes in CO₂ concentration alone is less certain. With the consideration of nitrogen limitation, the already weakened CO₂ fertilization effect under declining CO₂ concentrations could be further constrained, exacerbating the carbon loss due to changes in CO₂ concentration alone. However, this may be counteracted by an enhanced rate of photosynthesis as 1135 declining CO₂ concentrations decrease carbon-nitrogen ratios.

Deleted: positive emissions

Formatted: Subscript

Deleted: positive emissions

Deleted: positive emissions

Deleted: negative emissions

Deleted: re

Each of the two approaches used here to quantify carbon cycle feedback parameters has its benefits and drawbacks. Because the “CDR-reversibility” simulation is commonly used in literature (Schwinger & Tjiputra, 2018; Keller et al., 2018; Zickfeld et al., 2016), it allows easier comparison of results across models. However, research shows that this idealized scenario may delay the land sink-to-source transition, and underestimate ocean carbon uptake and the strength of the permafrost carbon feedback (MacDougall, 2019).

In their 2016 paper, Zickfeld et al. used zero emissions simulations to correct for the thermal and carbon cycle inertia in a suite of “CDR-reversibility” simulations, similar to our novel approach in this study. This reduced, but did not eliminate the climate system inertia, consistent with our results. Although our approach does not eliminate the inertia, it provides a more accurate estimate of the magnitude of carbon cycle feedbacks in the ramp-down phase by reducing the response to the prior CO₂ trajectory, bringing the estimate closer to a quantification of carbon cycle feedbacks under negative emissions alone. The remaining inertia may be associated with the different configurations in which the “CDR-reversibility” and “zeroemit” simulations were run: the former were run in concentration-driven mode whereas, the latter were emissions-driven. Therefore, changes in land and ocean carbon fluxes affect the atmospheric CO₂ concentration in the zero emissions simulations, but not in the “CDR-reversibility” simulations. Alternatively, the remaining inertia may be related to irreversible changes in vegetation distribution in the “CDR-reversibility” simulations. Lastly, the linearity assumption made in this experimental design may not hold, that is, the total carbon cycle response in the ramp-down phase may not be a linear combination of the committed response following increasing CO₂ concentration and temperature, and the response driven by the decrease in atmospheric CO₂ and temperature in the ramp-down phase. If the responses to increasing and decreasing CO₂ concentrations are not additive, then the zero emissions simulations may not quantify and remove all the inertia in the “CDR-reversibility” simulations.

Carbon cycle feedbacks under negative emissions have been quantified from the ramp-down phase of the “CDR-reversibility” simulation. However, this approach underestimates the magnitudes of carbon cycle feedbacks because the response in the ramp-down phase includes climate system inertia effects that generally weaken both feedbacks. Our novel approach aims to reduce the inertia in the ramp-down phase, thereby improving the quantification of carbon cycle feedbacks under negative emissions. We find that the magnitudes of the concentration-carbon and climate-carbon feedbacks under negative emissions are larger in our approach as compared to the standard approach. The concentration-carbon feedback drives greater land and ocean carbon release under negative emissions in our approach than in the standard approach. The climate-carbon feedback promotes more land and ocean carbon sequestration in our approach than in the standard approach. This has two implications: using feedback parameters from the standard approach will (1) underestimate land and ocean carbon release under negative emissions due to changes in CO₂ concentration alone (concentration-carbon feedback), and (2) underestimate land and ocean carbon gain due changes in climate alone (climate-carbon feedback). Given that the concentration-carbon feedback is the dominant feedback, quantifying carbon cycle feedbacks under negative emissions from the “CDR-reversibility” simulation

Deleted: The main limitation is that carbon cycle feedback parameters quantified for the ramp-down phase include carbon cycle inertia effects, making this approach inaccurate for quantifying carbon cycle feedbacks under negative emissions.

Deleted: under negative emissions

Deleted: positive emissions

Formatted: Subscript

Formatted: Not Highlight

Deleted: We hypothesize that

Deleted:

Formatted: Not Highlight

Deleted: Alternatively

Formatted: Not Highlight

Deleted: .

Deleted: prior positive emissions and negative emissions

Formatted: Subscript

Deleted: Lastly, the remaining inertia may be associated with the different configurations in which the “CDR-reversibility” and “zeroemit” simulations were run: the former were run in concentration-driven mode whereas, the latter were emissions-driven. Therefore, changes in land and ocean carbon fluxes affect the atmospheric CO₂ concentration in the zero emissions simulations, but not in the “CDR-reversibility” simulations.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

will result in the underestimation of carbon loss under negative emissions, thereby overestimating the effectiveness of negative emissions in drawing down CO₂.

1200

Future research should test the robustness of these results in a multi-model framework. A first step could be analyzing the “CDR-reversibility” simulations in three modes (biogeochemically coupled, radiatively coupled and fully coupled) in the next CMIP phase. In addition, [increasing and decreasing CO₂ trajectories](#) could be applied from an equilibrium state to overcome issues related to climate system inertia.

Deleted: positive and negative

Deleted: emissions

1205 **5 Code/Data Availability**

The UVic ESCM data will be made available after publishing and the model code for UVic ESCM 2.10 is available at <http://terra.seos.uvic.ca/model/2.10/>.

6 Author contribution

1210 K.Z. developed the research question and worked with C.N. on the initial data analysis. V.R.C ran the model simulations and worked with K.Z. to analyse and interpret the model data and write the manuscript. C.N. also helped revise the manuscript.

7 Competing interests

The authors declare no competing interests.

8 Acknowledgements

1215 This research was funded by the Natural Sciences and Engineering Research Council (NSERC) Discovery Grant Program. Computing resources were provided by the Digital Research Alliance of Canada (formerly Compute Canada).

References

- Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon–Concentration and Carbon–Climate Feedbacks in CMIP5 Earth System Models, *J. Climate*, 26, 5289–5314, <https://doi.org/10.1175/JCLI-D-12-00494.1>, 2013.
- 1220 Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C., Krasting, J., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T.: Carbon-concentration and carbon-climate feedbacks in

- CMIP6 models, and their comparison to CMIP5 models, *Biogeosciences*, 17, 4173–4222, doi.org/10.5194/bg-17-4173-2020, 2020.
- Boer, G. J. & Arora, V.: Geographic Aspects of Temperature and Concentration Feedbacks in the Carbon Budget, *J. Clim.*, 23(3), 775–784, doi: 10.1175/2009JCLI3161.1, 2010.
- 1230 Boer, G. J. & Arora, V.: Feedbacks in emission-driven and concentration-driven global carbon budgets, *J. Clim.*, 32(10), 3326–3341, doi: 10.1175/JCLI-D-12-00365.1, 2013.
- Boucher, O., Halloran, P. R., Burke, E. J., Doutriaux-Boucher, M., Jones, C. D., Lowe, J., Ringer, M. A., Robertson, E., and Wu, P.: Reversibility in an earth system model in response to CO₂ concentration changes, *Environ. Res. Lett.*, 7, 024013, doi: 10.1088/1748-9326/7/2/024013, 2012.
- 1235 Cao, L. & Caldeira, K.: Atmospheric carbon dioxide removal: Long term consequences and commitment, *Environ. Res. Lett.*, 5, 024011, doi: 10.1088/1748-9326/5/2/024011, 2010.
- Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks. In *Climate Change 2021: The Physical Science Basis*.
- 1240 Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. P. An, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekci, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, doi:10.1017/9781009157896.007, 2021
- 1245 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S. and Thornton, P.: Carbon and Other Biogeochemical Cycles, in: Working Group I Contribution to the Intergovernmental Panel on Climate Change Fifth Assessment Report *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., Cambridge University Press, 2013.
- 1250 Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., & Totterdell, I.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 184–187, doi: 10.1038/35041539, 2000.
- Cox, P.: Description of the TRIFFID Dynamic Global Vegetation Model, Hadley Centre Technical Note # 24, UK Met Office, available at: https://digital.nmla.metoffice.gov.uk/IO_cc8f146a-d524-4243-88fc-e3a3bcd782e7/ (last access: June 2022), 2001
- 1255 Eby, M., Weaver, A. J., Alexander, K., Zickfeld, K., Abe-Ouchi, A., Cimatoribus, A., Crespin, E., Drijfhout, S. S., Edwards, N. R., Eliseev, A. V., Feulner, G., Fichefet, T., Forest, C. E., Goose, H., Holden, P. B., Joos, F., Kawamiya, M., Kicklighter, D., Kienert, H., Matsumoto, K., Mokhov, I. I., Monier, E., Olsen, S. M., Pedersen, J. O. P., Perrette, M., Philippon-Berthier, G., Ridgwell, A., Schlosser, A., Schneider von Deimling, T., Shaffer, G., Smith, R. S., Spahni, R., Sokolov, A. P., Steinacher, M., Tachiiri, K., Tokos, K., Yoshimiri, M., Zeng, N., and Zhao, F.: Historical and idealized climate model experiments: An intercomparison of Earth system models of intermediate complexity, *Clim. Past*, 9(3), 1111–1140, doi:10.5194/cp-9-1111-2013, 2013.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, doi.org/10.5194/gmd-9-1937-2016, 2016.
- 1265 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., Zeng, A. N., and Friedlingstein, P.: Climate–Carbon Cycle Feedback Analysis: Results from the C4 MIP Model Intercomparison, *J. Clim.*, 19, 3337–3353, doi.org/10.1175/JCLI3800.1, 2006.
- 1270 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneeth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L.

Deleted:

Deleted:

- 1275 P., Cronin, M., Evans, W., Falk, S., Feely, R.A., Gasser, T., Gelen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, O., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A., Jerslid, A., Kadono, K., Kato, E., Kennedy, D., Goldewijk, K. K., Knauer, J., Korsbakken, Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzel, N., Monacci, N. M., Nakaoka, S., Niwa, Y., Obrien, K., Ono, T., Palmer, P.I., Pan, N., Pierrot, D., Poccock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tillbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Waranne, A. W., Wright, R., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2022, *Earth Syst. Sci. Data*, 14, 4811–4900, doi: 10.5194/essd-14-4811-2022, 2022.
- 1280 Friedlingstein, P., Meinshausen, M., Arora, V., Jones, C., Anav, A., Liddicoat, S., & Knutti, R.: Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks, *J. Clim.*, 27(2), 511–526, doi: 10.1175/JCLI-D-12-00579.1, 2014.
- 1285 Fung, I.Y., Doney, S. C., Lindsay, K., & Jasmin J. G.: Evolution of carbon sinks in a changing climate, *PNAS*, 102(32), 11201–11206, doi: 10.1073/pnas.0504949102, 2005.
- Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M. Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quéré, C., Raupach, M. R., Sharifi, A., Smith P., and Yamagata, Y.: Betting on negative emissions, *Nat. Clim. Change*, 1-3, doi.org/10.1038/nclimate2392, 2014.
- 1290 IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability, Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)], Cambridge University Press, In Press, 2022.
- 1295 Jeltsch-Thömmes, A., Stocker, T. F., & Joos, F.: Hysteresis of the Earth system under positive and negative CO₂ emissions. *Environmental Research Letters*, 15(12), 124026. doi: 10.1088/1748-9326/abc4af, 2020.
- Jones, C. D., Arora, V., Friedlingstein, P., Bopp, L., Brovkin, V., Dunne, J., Graven, H., Hoffman, F., Ilyina, T., John, J. G., Jung, M., Kawamiya, M., Koven, C., Pongratz, J., Raddatz, T., Randerson, J. T., and Zaehle, S.: C4MIP – The Coupled Climate-Carbon Cycle Model Intercomparison Project: experimental protocol for CMIP6, *Geosci. Model Dev.*, 9, 2853–2880, doi: 10.5194/gmd-9-2853-2016, 2016.
- 300 Jones, C. D., Ciais, J., Davis, S. J., Friedlingstein, P., Gasser, T., Peters, G. P. ... Wiltshire, A.: Simulating the Earth System response to negative emissions, *Environ. Res. Lett.*, 11, 095012, doi: 10.1088/1748-9326/11/9/095012, 2016.
- Jones, C. D., Cox, P. M., Essery, R. L. H., Roberts, D. L., and Woodage, M. J.: Strong carbon cycle feedbacks in a climate model with interactive CO₂ and sulphate aerosols, *Geophys. Res. Lett.*, 30(9), 1479, doi:10.1029/2003GL016867, 2003.
- 1305 Keller, D. P., Lenton, A., Scott, V., Vaughan, N. E., Bauer, N., Ji, D. ... Zickfeld, K.: The Carbon Dioxide Removal Model Intercomparison Project (CDR-MIP): Rationale and experimental protocol for CMIP6, *Geosci. Model Dev.*, 11, 1133 – 1160, doi: 10.5194/gmd-11-1133-2018, 2018.
- Keller, D. P., Oeschles, A. & Eby, M.: A new marine ecosystem model for the University of Victoria earth system climate model, *Geosci. Model Dev.*, 5(5), 1195–1220, doi: 10.5194/gmd-5-1195-2012, 2012.
- 1310 Koven, C. D., Arora, V. K., Cadule, P., Fisher, R. A., Jones, C. D., Lawrence, D. M., Lewis, J., Lindsay, K., Mathesius, S., Meinshausen, M., Mills, M., Nicholls, Z., Sanderson, B. M., Séférian, R., Swart, N. C., Wieder, W. R., and Zickfeld, K.: Multi-century dynamics of the climate and carbon cycle under both high and net negative emissions scenarios, *Earth Syst. Dynam.*, 13, 885–909, https://doi.org/10.5194/esd-13-885-2022, 2022.
- MacDougall, A. H.: Limitations of the 1 % experiment as the benchmark idealized experiment for carbon cycle intercomparison in C4MIP, *Geosci. Model Dev.*, 12, 597–611, doi: 10.5194/gmd-12-597-2019, 2019.
- 1315 MacDougall, A. H. & Knutti, R.: Projecting the release of carbon from permafrost soils using a perturbed parameter ensemble modelling approach, *Biogeosciences*, 13, 2123–2136, doi: 10.5194/bg-13-2123-2016, 2016.
- MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., Matthews, H. D., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., Eby, M., Eliseev, A. V., Hajima, T., Holden, P. B., Jeltsch-Thömmes, A., Koven, C., Mengis, N., 1320 Menviel, L., Michou, M., Mokhov, I. I., Oka, A., Sewinger, J., Séférian, R., Shaffer, G., Sokolov, A., Tachiiri, K., Tjiputra,

Deleted:

Deleted:

Deleted:

- J., Wiltshire, A. and Ziehn, T.: Is there warming in the pipeline? A multi-model analysis of zero emissions commitment of CO₂, *Biogeosciences*, 17, 2987-3016, doi: 10.5194/bg-17-2987-2020, 2020.
- 1325 Matthews, H. D. and Caldeira, K.: Stabilizing climate requires near-zero emissions, *Geophys. Res. Lett.*, 35, L04705, doi: 10.1029/2007GL032388, 2008.
- Meissner, K. J., Weaver, A. J., Matthews, H. D. & Cox, P. M.: The role of land surface dynamics in glacial inception: a study with the UVic Earth System Model, *Clim. Dyn.*, 21(7–8), 515–537, doi: 10.1007/s00382-003-0352-2. 2003.
- 1330 Melnikova, I., Boucher, O., Cadule, P., Ciais, P., Gasser, T., Quilcaille, Y., Shiogama, H., Tachiiri, K., Yokohata, T. and Tanaka, K.: Carbon cycle response to temperature overshoot beyond 2°C: An analysis of CMIP6 models. *Earth's Future*, 9, e2020EF001967. <https://doi.org/10.1029/2020EF001967>, 2021.
- Mengis, N., Keller, D. P., MacDougall, A., Eby, M., Wright, N., Meissner, K. J. Oschlies, A., Schmittner, A., MacIsaac, A. J., Matthews, H. D., and Zickfeld, K.: Evaluation of the University of Victoria Earth System Climate Model version 2.10 (UVic ESCM 2.10), *Geosci. Model Dev. Discuss.*, 1-28, doi: 10.5194/gmd-13-4183-2020, 2020.
- 1335 Pacanowski, R. C.: MOM 2 Documentation, users guide and reference manual, GFDL Ocean Group Technical Report 3, Geophys. Fluid Dyn. Lab., Princet. Univ. Princeton, NJ, 1995.
- Park, S. & Kug, J.: A decline in atmospheric CO₂ levels under negative emissions may enhance carbon retention in the terrestrial biosphere, *Commun. Earth. Environ.* 3, 289, doi: 10.1038/s43247-022-00621-4, 2022
- 1340 Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J., and Séférian, R.: Estimating and tracking the remaining carbon budget for stringent climate targets, *Nature*, 571, 335–342, doi: 10.1038/s41586-019-1368-z, 2019.
- Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., Mundaca, L., Seferian, R., and Vilarino, M. V.: Mitigation Pathways Compatible with 1.5 °C in the Context of Sustainable Development, in: *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*, edited by: Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock, T., Tignor, M., and Waterfield, T., available at: <https://www.ipcc.ch/report/sr15/mitigation-pathways-compatible-with-1-5c-in-the-context-of-sustainable-4-development/>, 2018.
- 1350 Schwinger, J. and Tjiputra, J.: Ocean Carbon Cycle Feedbacks Under Negative Emissions, *Geophys. Res. Lett.*, 45, 5062–5070, doi: 10.1029/2018GL077790, 2018
- Solomon, S., Plattner, G.-K., Knutti, R., and Friedlingstein, P.: Irreversible climate change due to carbon dioxide emissions, *P. Natl. Acad. Sci.*, 106, 1704–1709, doi: 10.1073/pnas.0812721106, 2009.
- 1355 Tokarska, K. B. & Zickfeld, K.: The effectiveness of net negative carbon dioxide emissions in reversing anthropogenic climate change, *Environ. Res. Lett.*, 094013, doi: 10.1088/1748-9326/10/9/094013, 2015.
- UNFCCC.: Adoption of the Paris Agreement. Retrieved from <https://unfccc.int/sites/default/files/resource/docs/2015/cop21/eng/109r01.pdf>, 2022.
- 1360 Weaver, A. J., Eby, M., Wiebe, E. C., Bitz, C. M., Duffy, P. B., Ewen, T. L., ... Fanning, A. F.: The UVic Earth System Climate Model: Model description, climatology, and applications to past, present and future climates, *Atmosphere Ocean*, 39, 361–428, 2001.
- Zickfeld, K., Azevedo, D., Mathesius, S. & Matthews, H. D.: Asymmetry in the climate-carbon cycle response to positive and negative CO₂ emissions, *Nat. Clim. Chang.* 11, 613–617, doi: 10.1038/s41558-021-01061-2, 2021.
- Zickfeld, K., Eby, M., Matthews, H. D., Schmittner, A., and Weaver, A. J.: Nonlinearity of Carbon Cycle Feedbacks, *J. Climate*, 24, 4255–4275, doi: 10.1175/2011JCLI3898.1, 2011.
- 1365 Zickfeld, K., MacDougall, A. H., & Matthews, H. D.: On the proportionality between global temperature change and cumulative CO₂ emissions during periods of net negative CO₂ emissions, *Environ. Res. Lett.*, 11(5), 055006, doi: 10.1088/1748-9326/11/5/055006, 2016.

1370 Ziehn, T., Lenton, A. & Law, R.: An assessment of land-based climate and carbon reversibility in the Australian Community
Climate and Earth System Simulator, *Mitig. Adapt. Strateg. Glob. Change*, 25, 713–731, doi: 10.1007/s11027-019-09905-1,
2020.

Page 6: [1] Formatted Kirsten Zickfeld 4/14/23 1:07:00 PM

Subscript

Page 6: [1] Formatted Kirsten Zickfeld 4/14/23 1:07:00 PM

Subscript

Page 6: [2] Deleted Kirsten Zickfeld 4/14/23 1:09:00 PM

1.

Page 6: [2] Deleted Kirsten Zickfeld 4/14/23 1:09:00 PM

2.

Page 6: [3] Deleted Vimbikayi Rachel Chimuka 3/30/23 10:42:00 AM

3.

Page 6: [3] Deleted Vimbikayi Rachel Chimuka 3/30/23 10:42:00 AM

4.

Page 6: [4] Deleted Kirsten Zickfeld 4/14/23 1:10:00 PM

5.

Page 6: [4] Deleted Kirsten Zickfeld 4/14/23 1:10:00 PM

6.

Page 6: [4] Deleted Kirsten Zickfeld 4/14/23 1:10:00 PM

7.

Page 6: [4] Deleted Kirsten Zickfeld 4/14/23 1:10:00 PM

8.

Page 6: [4] Deleted Kirsten Zickfeld 4/14/23 1:10:00 PM

9.

Page 6: [5] Deleted Kirsten Zickfeld 4/21/23 4:26:00 PM

Page 6: [5] Deleted Kirsten Zickfeld 4/21/23 4:26:00 PM

Page 6: [6] Deleted Vimbikayi Rachel Chimuka 4/23/23 7:24:00 AM

Page 6: [6] Deleted Vimbikayi Rachel Chimuka 4/23/23 7:24:00 AM

Page 6: [7] Deleted Kirsten Zickfeld 4/21/23 4:29:00 PM

▲
Page 6: [8] Deleted Kirsten Zickfeld 4/21/23 4:30:00 PM

▼
▲
Page 6: [8] Deleted Kirsten Zickfeld 4/21/23 4:30:00 PM

▼
▲
Page 6: [9] Deleted Vimbikayi Rachel Chimuka 4/23/23 7:07:00 AM

▼
▲
Page 6: [9] Deleted Vimbikayi Rachel Chimuka 4/23/23 7:07:00 AM

▼
▲
Page 9: [10] Deleted Kirsten Zickfeld 4/14/23 6:13:00 PM

▼
▲
Page 9: [10] Deleted Kirsten Zickfeld 4/14/23 6:13:00 PM

▼
▲
Page 9: [10] Deleted Kirsten Zickfeld 4/14/23 6:13:00 PM

▼
▲
Page 9: [11] Formatted Vimbikayi Rachel Chimuka 4/4/23 10:29:00 AM

Not Highlight

▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [12] Deleted Vimbikayi Rachel Chimuka 4/15/23 6:04:00 AM

▼
▲
Page 9: [13] Deleted **Kirsten Zickfeld** **4/15/23 9:34:00 AM**

▼
▲
Page 9: [13] Deleted **Kirsten Zickfeld** **4/15/23 9:34:00 AM**

▼
▲
Page 9: [13] Deleted **Kirsten Zickfeld** **4/15/23 9:34:00 AM**

▼
▲
Page 9: [13] Deleted **Kirsten Zickfeld** **4/15/23 9:34:00 AM**

▼
▲
Page 9: [14] Formatted **Kirsten Zickfeld** **4/15/23 9:29:00 AM**

Subscript

▲
Page 9: [15] Deleted **Vimbikayi Rachel Chimuka** **4/2/23 3:45:00 PM**

▼
▲
Page 9: [16] Deleted **Vimbikayi Rachel Chimuka** **4/4/23 11:19:00 AM**

▼
▲
Page 9: [17] Deleted **Vimbikayi Rachel Chimuka** **4/16/23 9:35:00 AM**

▼
▲
Page 9: [18] Deleted **Kirsten Zickfeld** **4/15/23 9:40:00 AM**

▼
▲
Page 9: [19] Deleted **Vimbikayi Rachel Chimuka** **4/2/23 3:40:00 PM**

▼
▲
Page 9: [20] Deleted **Vimbikayi Rachel Chimuka** **4/4/23 11:23:00 AM**

▼
▲
Page 9: [21] Formatted **Kirsten Zickfeld** **4/15/23 9:49:00 AM**

Subscript

▲
Page 9: [22] Deleted **Kirsten Zickfeld** **4/15/23 9:45:00 AM**

▼
▲
Page 9: [22] Deleted **Kirsten Zickfeld** **4/15/23 9:45:00 AM**

▼
▲
Page 9: [23] Formatted **Vimbikayi Rachel Chimuka** **4/5/23 12:36:00 AM**

Not Highlight

▲
Page 9: [24] Deleted **Kirsten Zickfeld** **4/15/23 9:46:00 AM**

Subscript

▲
Page 9: [27] Deleted **Vimbikayi Rachel Chimuka** **4/16/23 10:50:00 AM**

▼

▲
Page 9: [27] Deleted **Vimbikayi Rachel Chimuka** **4/16/23 10:50:00 AM**

▼

▲
Page 9: [28] Formatted **Kirsten Zickfeld** **4/15/23 10:02:00 AM**

Subscript

▲
Page 15: [29] Deleted **Vimbikayi Rachel Chimuka** **4/15/23 8:06:00 AM**

▼