

1 **Title:** Contemporary biodiversity pattern is affected by climate change at multiple temporal scales in
2 steppe on the Mongolian Plateau

3 **Running Title:** Climate change and plant biodiversity

4 **List of Authors:**

5 Zijing Li^{1,2}, Zhiyong Li^{3,*}, Xuze Tong⁴, Lei Dong^{1,2}, Ying Zheng^{1,2}, Jinghui Zhang³, Bailing Miao⁵,

6 Lixin Wang³, Liqing Zhao³, Lu Wen³, Guodong Han⁶, Frank Yonghong Li³, Cunzhu Liang³

7 **Institutional affiliations:**

8 ¹Yinshanbeilu Grassland Eco-hydrology National Observation and Research Station, China Institute
9 of Water Resources and Hydropower Research, Beijing 100038, China

10 ²Institute of Water Resources for Pastoral Area Ministry of Water Resources, Hohhot 010020, Inner
11 Mongolia

12 ³Key Laboratory of Ecology and Resource Use of the Mongolian Plateau, Ministry of Education of
13 China; Collaborative Innovation Center for Grassland Ecological Security, Ministry of Education of
14 China; School of Ecology and Environment, Inner Mongolia University, Hohhot 010021, P. R. China

15 ⁴ Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural
16 Sciences, Beijing 100081, P. R. China

17 ⁵ Inner Mongolia Meteorological Institute, Hohhot 010051, P. R. China

18 ⁶Key Laboratory of Grassland Resources of the Ministry of Education, Key Laboratory of Forage
19 Cultivation, Processing and High Efficient Utilization of the Ministry of Agriculture and Rural
20 Affairs, Inner Mongolia Key Laboratory of Grassland Management and Utilization, College of
21 Grassland, Resources and Environment, Inner Mongolia Agricultural University, Hohhot 010011, P.
22 R. China

23 **Contact Information:** *Corresponding author. E-mail: zylee007@imu.edu.cn; tel: +86 136 7475
24 7329.

25

26 **Abstract**

27 Present and historical climate conditions jointly determine contemporary biodiversity patterns and
28 ecosystem functions. However, it remains unclear how contemporary climate and paleoclimate
29 changes together affect the three dimensions of biodiversity (i.e., taxonomic diversity, functional
30 diversity and phylogenetic diversity) and their relationship with ecosystem functions. Here, we
31 assess the impact of current climate, paleoclimate and its anomalies on contemporary biodiversity
32 and ecosystem functions. We estimated the taxonomic, functional and phylogenetic diversity of
33 grassland on the Mongolian Plateau using vegetation survey data and trait information. We then used
34 Random Forest and Structural Equation models to assess the relative importance of the present, the
35 Mid-Holocene and the Last-Glacial Maximum climate as well as paleoclimate changes as
36 determinants of diversity and aboveground biomass. Our results showed that paleoclimate changes
37 and modern climate jointly determined contemporary biodiversity patterns, while community
38 biomass was mainly affected by modern climate, namely the aridity index. Modern aridity and
39 temperature were two major influences on all three dimensions of biodiversity. Mid-Holocene
40 climate anomalies had a strong effect on species richness and phylogenetic diversity, while
41 functional diversity was mainly affected by temperature anomalies since the Last Glacial Maximum.
42 These findings suggest that contemporary biodiversity patterns may be affected by processes at
43 divergent temporal scales. Our results show that simultaneously exploring the response of the three
44 dimensions of biodiversity in different periods of climate change and the theoretical framework for

45 its impact on community biomass is helpful to provide a more comprehensive understanding of
46 patterns of biodiversity and its relationship with ecosystem functions.

47 **Keywords**

48 Aridity index, ecosystem functions, functional diversity, paleoclimate anomaly, phylogenetic
49 diversity, taxonomic diversity

50 **1. Introduction**

51 Global climate changes, such as increasing temperature and changing precipitation, are altering
52 biodiversity patterns at an unprecedented rate, and have serious consequences for ecosystem
53 functions related to biodiversity changes (Mottl et al., 2021; Nolan et al., 2018; Ye et al., 2019).
54 Ecosystem functioning associated with biodiversity, and contemporary diversity patterns and their
55 relationships may be regulated by modern climate, past climate and climate change (Fordham et al.,
56 2020; Eiserhardt et al., 2015; Svenning et al., 2015).

57 Ecosystem functions are strongly influenced by biodiversity (Isbell et al., 2011; Hooper et al.,
58 2005), but understanding the effects of different dimensions of biodiversity (i.e., taxonomic diversity,
59 functional diversity and phylogenetic diversity) on ecosystem functions is still a challenge in ecology
60 (Van Der Plas, 2019; Cadotte et al., 2008; Swenson, 2011). Traditionally, most studies of biodiversity
61 and ecosystem functions have predominantly focused on the relationship between taxonomic
62 diversity (i.e., species richness) and biomass or productivity. However, only considering taxonomic
63 diversity has the limitation that it may ignore important ecological processes such as the formation of
64 function traits or the evolutionary history of species (Flynn et al., 2011; Swenson, 2011). The
65 importance of taking into account other aspects of biodiversity, including functional diversity and
66 phylogenetic diversity, has increasingly been recognized. The effects of plant functional diversity on

67 ecosystem functioning are largely driven by variations in plant functional traits, such as adult plant
68 height and leaf size (Diaz et al., 2016). Phylogenetic diversity (PD) is a key driver of community
69 assembly and ecosystem functions (Srivastava et al., 2012; Cavender-Bares et al., 2009). Previous
70 research had shown that PD may be a better metric than species richness or functional diversity for
71 predicting plant biomass (Cadotte et al., 2008). Communities with higher PD have stronger
72 anti-interference ability because of their evolutionary potential to adapt to changing environmental
73 conditions. However, the role of multi-dimensional biodiversity in driving ecosystem functions in
74 response to climate change still needs further research. Elucidating the impacts of multi-dimensional
75 biodiversity on biomass in a changing world can deepen our understanding of the direct and indirect
76 effects of changes in climate and biodiversity on terrestrial ecosystems, and improve predictions of
77 the ecological consequences of global climate changes.

78 Changes in community biodiversity and composition caused by climate change have been
79 explored in various ecosystems (Avolio et al., 2021), including tropical forests in the Amazon
80 (Esquivel-Muelbert et al., 2019), alpine meadows on the Qinghai-Tibet Plateau (Liu et al., 2018a)
81 and a heterogeneous California grassland (Harrison et al., 2015). Dispersal limitation and
82 environmental filtering (i.e., climatic conditions and paleoclimate change) play an important role in
83 sorting species from the global species pool and in shaping large-scale diversity patterns (Kubota et
84 al., 2018; Liu et al., 2021). Many studies have found that climate changes in the Mid-Holocene and
85 the Quaternary interglacial period are the main driving factors of current species distribution,
86 functional traits and community phylogeny (Svenning and Skov, 2007b). The current patterns of
87 species richness and distribution of European flora are jointly affected by contemporary climatic
88 conditions and climate change during the Late Quaternary glacial-interglacial period (Svenning and

89 Skov, 2007a; Svenning and Skov, 2007b). Contemporary plant functional traits and composition are
90 highly sensitive to climate and influenced by climate from tens of thousands of years ago, with
91 important consequences for ecosystem functions (Blonder et al., 2018; Butler et al., 2017). For
92 example, in Europe patterns of plant functional diversity exhibited prominent glacial-interglacial
93 climate change imprints (Ordonez and Svenning, 2015; Ordonez and Svenning, 2017). This is partly
94 due to the non-random removal of functional combinations that may have occurred during the
95 Last-Glacial Maximum (LGM; ~21,000 years ago), increasing functional differences between native
96 and migratory species, leading to smaller and dispersed functional spaces (Ordonez and Svenning,
97 2017). Evidence derived from the fossil pollen sequence and ecological datasets has shown that the
98 acceleration of biodiversity changes began millennia ago, whereas vegetation changes during the
99 Late Pleistocene to Early Holocene were driven primarily by changing climate (Mottl et al., 2021). In
100 the South American tropics, the species composition and structure of Last Glacial Maximum forests
101 were quite different from those of today, and biomass was also lower than that of contemporary
102 forests (Mayle et al., 2009).

103 Furthermore, the effects of climate change on biodiversity should depend partly on climate
104 displacement rate (climate-change velocity) (Sandel et al., 2011). For example, phylogenetic
105 clustering increased with increasing intensity of Quaternary glacial-interglacial climatic oscillations
106 in South America and Africa (Kissling et al., 2012). There is evidence that vegetation changes in
107 Northern Europe were non-linear and varied greatly among regions in the early-Mid Holocene
108 (Seddon et al., 2014). Paleoclimate simulations since 21,000 years B.P. suggest that there are great
109 similarities between past and future temperature changes in Eurasian temperate grasslands (Fordham
110 et al., 2020). During the last glacial-interglacial transition period, global warming and related climate

111 change led to changes in ecosystems to a degree comparable to the predicted warming under future
112 high-emission scenarios (Nolan et al., 2018). Therefore, studying the impact of past climate and its
113 effects on contemporary biodiversity patterns will help us understand the potential effects of future
114 climate changes. However, most studies exploring the impact of present climate and paleoclimate
115 drivers on ecosystem functions have focused on forest ecosystems, while grassland ecosystems
116 remain understudied. In addition, it is not yet clear whether taxonomy, function and phylogenetic
117 diversity will vary with climate gradients and climate changes, and the intensity of any such changes
118 remain unclear. Therefore, there is an urgent need to consider the impact of current climate,
119 paleoclimate and its anomalies on contemporary biodiversity and ecosystem functions.

120 The Mongolian Plateau is located in the arid and semi-arid area of eastern Eurasia. Our previous
121 researches ~~has~~ have shown that climate is an important direct factor mediating the relationship
122 between species richness and aboveground biomass in the Mongolian plateau steppe (Li et al., 2021;
123 Li et al., 2020). However, in the Late Quaternary, the Mongolian Plateau experienced severe
124 alternating dry and wet climatic fluctuations (Maestre et al., 2021; Tian et al., 2017), which may have
125 a significant impact on current plant diversity and ecosystem function. However, the impacts of
126 climate in different periods and climate legacy effects on aboveground biomass of steppes in the
127 Mongolian Plateau, as well as whether these effects are biodiversity dependent, remain to be
128 evaluated. To address these knowledge gaps, we coupled data from a field survey of 152 sites in the
129 Mongolian Plateau (Fig. S1) to existing databases on present and past climates to explore present
130 climate and the legacy effects of climate during the Last Glacial Maximum and the Mid-Holocene on
131 current biodiversity and biomass.

132 Here, we aim to evaluate the relative importance of taxonomic, functional, and phylogenetic

133 diversity on community biomass of steppe in the Mongolian Plateau, and to assess whether and how
134 climate in different periods and their changes affect the three dimensions of diversity, and their
135 relationships with community biomass. We specifically ask: (1) Do the three dimensions of diversity
136 affect community biomass, and if so, to what extent? (2) Whether and how climate at divergent
137 temporal scales and paleoclimatic changes affect different dimensions of biodiversity? And (3) If
138 there are effects, how do they affect the biodiversity-biomass relationship?

139 **2. Materials and Methods**

140 **2.1 Study Area**

141 We conducted field surveys in the grasslands of the Mongolian Plateau, including Inner Mongolia
142 Autonomous Region in China and Mongolia, during June to August of 2014~2018 (Fig. S1). The
143 Mongolian Plateau is located in the eastern part of Eurasia, with a geographic range of 37.61 °to
144 53.35 °N, 87.83 °to 125.95 °E, and an elevation range from 85 m to 4203 m a.s.l. (above sea level).
145 The mean annual temperature (MAT) of the study region ranges from -12 °C to 10 °C and mean
146 annual precipitation (MAP) ranges from 26 mm to 641 mm.

147 **2.2 Data Collection**

148 **2.2.1 Sites Sampled and Vegetation Investigation**

149 We investigated 152 grassland sites set in flat areas with little human interference and a relatively
150 homogeneous environment. “Little human interference” mainly refers to the selection of areas with
151 less human activities such as grazing for investigation. Specific indicators include the following
152 aspects: 1) Avoid choosing areas with manure of cattle and sheep; 2) Dominant species such as
153 Leymus chinensis, Stipa spp. are eaten less by livestock; 3) There were fewer degradation indicators.
154 It mainly refers to the decrease in the number of established species or dominant species in the

155 grassland, while the degeneration indicator plants such as *Convolvulus ammannii*, *Potentilla acaulis*,
156 *Artemisia frigida*, *Stellera chamaejasme* and *Cynanchum hancockianum* appeared in large numbers.
157 These sites were separated by about 50 km. According to the methods and protocols for plant
158 community inventories proposed by Fang et al. (Fang et al., 2009), most of the vegetation surveys in
159 this study were conducted by setting one 10 m x 10 m quadrat at the respective site. However, in a
160 few areas with sparse vegetation and large heterogeneity, we set three 10 m × 10 m quadrats to
161 ensure the accuracy of the survey data. ~~One or three 10 m × 10 m quadrats were set in each site (in a~~
162 ~~few areas with sparse vegetation and large heterogeneity, three 10 m × 10 m quadrats were set to~~
163 ~~ensure the accuracy of the survey data).~~ Within each quadrat, five 1 m × 1 m plots were placed at
164 each corner and at the center of the quadrat to investigate vegetation. A total of 296 vascular plant
165 species were recorded.

166 We measured species height, density, coverage and standing biomass of each species in each
167 plot. Three 1 m × 1 m plots along the diagonal line 10 m × 10 m quadrat were selected from the five
168 plots to measure the standing biomass of each species. Plant species were cut at ground level and
169 separately bagged, then oven-dried at 65 °C to constant weight, and weighed for dry matter.

170 **2.2.2 Climate Data**

171 Mean annual temperature (MAT, °C) and mean annual precipitation (MAP, mm) were obtained from
172 the CHELSA database (Climatologies at High Resolution for the Earth's Land Surface Areas,
173 <http://chelsa-climate.org/>) (Karger et al., 2017), a data set consisting of monthly temperature and
174 precipitation climatology from 1979 to 2013. Potential evapotranspiration (PET) was obtained from
175 the CGIAR-CSI (CGIAR Consortium for Spatial Information, <http://www.cgiar-csi.org>) based on
176 latitude and longitude data for each site (Trabucco and Zomer, 2019). The resolution of both is 30 arc

177 seconds. Aridity was measured using the aridity index (AI, $AI = MAP / PET$). Higher AI values
 178 indicate lower aridity and higher humidity. Paleoclimate data were downloaded from Worldclim
 179 (<http://worldclim.com/paleo-climate1>) with a resolution of 2.5 minutes. The database was made
 180 available by CMIP5 (Coupled Model Intercomparison Project Phase 5) and was calibrated based on
 181 the current climate using WorldClim 1.4. The MAT and MAP of the Mid-Holocene (MID, about
 182 6000 years ago) and the Last Glacial Maximum (LGM, about 22,000 years ago) were simulated
 183 based on CCMS4 (Community Climate System Model version 4). Climate anomaly (i.e., present-day
 184 values minus paleoclimate values) was used to indicate the degree of climate change since the MID
 185 and the LGM (Sandel et al., 2011). Temperature-change velocity from the LGM to the present
 186 (velocity, m/yr) was obtained from DRYAD (<https://datadryad.org>) (Sandel et al., 2011). The climate
 187 variables used in this study are shown in Table 1.

188

Table 1 Climate variables included in this study

	Climate variables	Abbreviation
The present climate (1979-2013)	Mean annual precipitation of the present	MAP
	Mean annual temperature of the present	MAT
	Aridity index of the present	AI
The Mid-Holocene climate	Mean annual precipitation of the Mid-Holocene	MAP_{mid}
	Mean annual temperature of the Mid-Holocene	MAT_{mid}
The Last Glacial Maximum climate	Mean annual precipitation of the Last Glacial Maximum	MAP_{lgm}
	Mean annual temperature of the Last Glacial Maximum	MAT_{lgm}
Paleoclimate change	MAP anomaly from the Mid-Holocene to the present	$AMAP_{mid}$
	MAT anomaly from the Mid-Holocene to the present	$AMAT_{mid}$

MAP anomaly from the Last Glacial Maximum to the present	AMAP _{l_{gm}}
MAT anomaly from the Last Glacial Maximum to the present	AMAT _{l_{gm}}
Temperature-change velocity from the LGM to the present	Velocity

189 **2.2.3 Plant functional traits**

190 The functional traits of each plant species include growth form, life form, phyllotaxy, single or
191 compound leaf, petiole, inflorescence, fruit type, flowering period, fruit ripening period, length of the
192 flowering period, water ecotypes, leaf length, leaf width and plant height (Table S1). These traits
193 were looked up in the Flora of Inner Mongolia (Third Edition) or Flora of China (Online Edition)
194 (<http://www.iplant.cn/frps>). The trait data for leaf length and leaf width were the mean values
195 provided in Flora of China (Fig. S2). For example, if the leaf length of *Leymus chinensis* was
196 described as 7-18 cm in the Flora of China, then leaf length was defined as 12.5 cm.

197 **2.3 Aboveground biomass**

198 The aboveground biomass of the community was calculated based on the dry matter accumulation
199 value (g/m^2) of all plant samples in each plot, and the average aboveground biomass of each plot in
200 each quadrat was calculated to represent the aboveground biomass of the plant community (g/m^2).

201 **2.4 Diversity calculations**

202 **2.4.1 Taxonomic diversity**

203 We used species richness (SR), Shannon-Wiener index and Pielou index as measures of community
204 taxonomic diversity. The species richness (number of species/ m^2) at each site was calculated as the
205 average number of species per plot. The Shannon-Wiener index and Pielou index were calculated
206 using the ‘diversity’ function in the “*vegan*” R package (Oksanen et al., 2020).

207 **2.4.2 Functional diversity**

208 The functional richness (FRic), functional evenness (FEve), functional divergence (FDiv), functional
209 dispersion (FDis) and Rao index (Rao's Q) were calculated based on the Gower distance of
210 functional traits using the 'dbFD' function in the 'FD' R package (Lalibert é et al., 2015).

211 **2.4.3 Phylogenetic diversity**

212 All plant species names recorded in surveys were checked in the plant list (www.theplantlist.org) to
213 obtain acceptable species names. Based on the APG III system (The Angiosperm Phylogeny Group,
214 2009), we entered the taxonomic information (family/genus/species) of all species into the
215 Phylomatic online plant database (Phylomatic Version 3, <http://phylodiversity.net/phyloomatic/>),
216 selected storedtree = "zanne2014 (Zanne et al., 2014)", and output a phylogenetic tree with branch
217 length and differentiation time (Fig. S2). Faith's phylogenetic diversity (PD, the sum of the
218 phylogenetic branch lengths), the mean pairwise distance (MPD) and the mean nearest taxon
219 distance (MNTD) between species in a community were calculated using the 'mpd' function in the
220 '*picante*' R package (Kembel et al., 2010).

221 **2.5 Data analysis**

222 First, we conducted a Random Forest model to estimate the relative importance of the three
223 dimensions of biodiversity to community biomass. The Random Forest model can alleviate
224 multicollinearity and complex interactions between independent variables. We built 5000 regression
225 trees in Random Forest, and each tree was fitted with 2/3 of the data. The other 1/3 of the data was
226 used to estimate the importance of each predictor variable, which was represented by the increase in
227 mean square error (MSE). In the Random Forest model, the variables that ranked first in terms of
228 taxonomic, functional and phylogenetic diversity were selected as proxies for the three dimensions of
229 biodiversity.

230 To explore the influence of climate in different periods and climate anomalies on biodiversity,
231 we also used the Random Forest model to analyze the relative importance of the present climate, the
232 Mid-Holocene climate, the Last Glacial Maximum climate, and paleoclimate change on the three
233 dimensions of biodiversity. To evaluate the relative importance of climate drivers of biodiversity, we
234 calculated the ratio between the increase in MSE of the predictor and the sum of all the increases in
235 MSE, which is expressed as a percentage.

236 To study the effects of climate in different periods and climate anomalies on biodiversity and
237 aboveground biomass, we performed piecewise structural equation models (pSEM) to test the direct
238 and indirect causal relationships between the climate in different periods and paleoclimate change,
239 species diversity, functional diversity, phylogenetic diversity and biomass. Model construction
240 procedures involved the following three stages. First, based on the results of the Random Forest
241 model, we identified the climate variables that significantly affected biodiversity and divided them
242 into composite variables. The model contained two composite variables that potentially represent
243 collections of variables in terms of the present climate and paleoclimate change. We calculated
244 Pearson correlations between the climate factors (Fig. S3) and deleted the predictors with high
245 correlation coefficients (>0.85 , $P < 0.05$) to avoid multicollinearity. Before pSEM analysis, all
246 selected predictors were standardized. Second, principal component analyses (PCAs) were used for
247 the composite variables with multiple predictors. For each composite variable, the first principal
248 component (PC1) explained 64~71% of the total variance and was used in the subsequent pSEM
249 analysis (Table S2). Third, pSEM was developed from the full conceptual model (Fig. S4). We fitted
250 the component models of the piecewise SEM as linear models and reported the standardized
251 coefficient for each path from each component model. We used Shipley's d-separation test and

252 Fisher's C statistic to evaluate the overall fit of the pSEM, ~~as well as Fisher's C statistic and Akaike~~
253 information criteria (AIC) to select the best model and AIC. We used standardized path coefficients to
254 measure the direct, indirect, and total effect sizes of the present climate and paleoclimate changes on
255 AGB.


256 We conducted all analyses in R x64 4.0.5. The Random Forest model was conducted using the
257 'randomForest' function in the 'randomForest' package (Cutler et al., 2007; Breiman et al., 2018),
258 and the significance of the variables was tested using the 'rp.importance' function in the 'rfPermute'
259 package (Archer, 2016). PCA was performed using the 'princomp' function in the 'psych' package in
260 R (Revelle, 2021). Piecewise structural equation modeling was conducted and tested using the
261 'piecewiseSEM' package (Lefcheck, 2020; Lefcheck and Freckleton, 2015).

262 **3. Results**

263 **3.1 Effects of biodiversity on aboveground biomass**

264 Among the 11 biodiversity variables, SR, PD and Pielou index identified by the RF model were the
265 three most significant predictors influencing aboveground biomass (Fig. 1a). Taken together,
266 taxonomic diversity represented by SR, Pielou and Shannon-Wiener index collectively contributed
267 43.78% to the RF model for aboveground biomass, while phylogenetic diversity represented by PD
268 (including MNTD and MPD) and functional diversity represented by FRic (including FDiv, FEve,
269 FDis and RaoQ) contributed 30.02% and 26.2%, respectively (Fig. 1b).

270 Based on the ranking results from the Random Forest model, SR, PD, and FRic were selected as
271 agents of taxonomic, phylogenetic, and functional diversity, respectively, and used for subsequent
272 analysis.



273

274

275

276

277

278

279

280

281

282


283

284

Fig. 1 The importance (a) and relative contributions (b) of the three dimensions of biodiversity to community biomass. An increase in mean square error (MSE) denotes an increase in the percentage mean square error. Significance levels are as follows: *: $P < 0.05$ and ***: $P < 0.001$.

3.2 Paleoclimate and present climate predictors of contemporary diversity

Random Forest models explained 35% (cross-validation $R^2 = 0.35$, $P < 0.001$), 31% (cross-validation $R^2 = 0.31$, $P < 0.001$) and 31% (cross-validation $R^2 = 0.31$, $P < 0.001$) of the variance in SR, PD and FRic, respectively (Fig. 2). Temperatures in different periods, including MAT_{mid} , MAT and MAT_{lgm} , were the main factors driving SR and PD ($P < 0.05$, Fig. 2a and 2b), followed by climate anomalies in various periods. However, the current climate (i.e., AI, MAT and MAP) had a stronger impact on FRic ($P < 0.05$, Fig. 2c).



285

286

287

288

289

290

291

292


293

294

Fig. 2 The importance of climate variables in different periods and paleoclimate change to species richness (a), phylogenetic diversity (b) and functional richness (c). An increase in mean square error (MSE) denotes an increase in the percentage mean square error. Significance levels are as follows: *:

$$P < 0.05 \text{ and } ***: P < 0.001.$$

In terms of relative contributions, the current climate and the paleoclimate changes had greater impacts on the three dimensions of biodiversity. Paleoclimate change was the best predictor of SR (33.90%) and PD (32.10%). Strikingly, the present climate (38.68%) showed a larger contribution to the prediction of FRic than paleoclimate change (26.06%; Fig. 3).



295

296

Fig. 3 Relative contributions of present climate, Mid-Holocene, Last Glacial Maximum and

297

paleoclimate change as drivers of species richness, phylogenetic diversity and functional richness.

298

3.3 Influence of climate factors on biodiversity and aboveground biomass

299

The present climate, MAT of the Mid-Holocene and the Last Glacial Maximum, and paleoclimate

300

change explained 38% ($R^2 = 0.38$), 33% ($R^2 = 0.33$), 36% ($R^2 = 0.36$) and 29% ($R^2 = 0.29$) of the

301

variances in SR, PD, FRic and AGB, respectively (Fig. 4). The present climate had direct and

302

indirect significant associations with AGB via all biodiversity variables (Fig. 4). The present climate

303

was significantly and positively associated with SR ($r_{\partial} = 0.46$, $P < 0.05$), PD ($r_{\partial} = 0.35$, $P < 0.05$)

304


and FRic ($r_{\partial} = 0.58$, $P < 0.05$, Fig. 4). However, the paleoclimate change had direct significant

305

negative effects on SR ($r_{\partial} = -0.28$, $P < 0.05$), PD ($r_{\partial} = -0.32$, $P < 0.05$) and FRic ($r_{\partial} = -0.25$, $P < 0.05$,

306

Fig. 4). MAT_{mid} and MAT_{lgm} had no significant effects on SR, PD, FRic and AGB (Fig. 4).



307

308 Fig. 4 SEMs fitted to the effects of climate in different periods and paleoclimate change on diversity

309 and AGB. The present climate is PCA component 1 of the AI and MAT. The paleoclimate change

310 represents a composite variable including MAT anomaly from the Mid-Holocene to the present,

311 MAT and MAP anomaly from the Last Glacial Maximum to the present. Numbers adjacent to arrows

312 represent the standardized path coefficients (r_{∂}). R^2 indicates the proportion of variance explained.

313 Blue (positive paths) and orange (negative paths) solid arrows indicate significant paths ($P < 0.05$),

314 and gray dashed arrows indicate non-significant paths ($P > 0.05$).

315

316 For the present climate, the direct effects of AI on SR ($r_{\partial} = 0.33$, $P < 0.05$), PD ($r_{\partial} = 0.27$, $P <$

317 0.05), FRic ($r_{\partial} = 0.49$, $P < 0.05$), and AGB ($r_{\partial} = 0.37$, $P < 0.05$) were all significantly positive, while

318 the direct effects of MAT on the three dimensions of biodiversity (SR, $r_{\partial} = -0.34$; PD, $r_{\partial} = -0.36$;


319 FRic, $r_{\partial} = -0.29$; $P < 0.05$ in all cases) were all significantly negative (Fig. 5). In addition, SR ($r_{\partial} =$

320 -0.25 , $P < 0.05$) and PD ($r_{\partial} = -0.25$, $P < 0.05$) were negatively affected by MAT anomaly from the

321 Mid-Holocene to the present ($AMAT_{mid}$), but FRic ($r_{\partial} = -0.23$, $P < 0.05$) was mainly driven by MAT

322 anomaly from the Last Glacial Maximum to the present ($AMAT_{lgm}$; Fig. 5). AI accounted for the

323 largest proportion of variation in AGB (80.06%), followed by MAT at 10.81% (Table 2). The direct
 324 effect ($r_{\hat{c}} = 0.37$) of AI on AGB was greater than the indirect effect ($r_{\hat{c}} = 0.12$, Table 2).



325
 326 Fig. 5 The effects of present climate (AI and MAT) and paleoclimate change (AMAT_{mid} and
 327 AMAT_{lgm}) on diversity and AGB. Numbers adjacent to arrows represent the standardized path
 328 coefficients. R^2 indicates the proportion of variance explained. Blue (positive paths) and orange
 329 (negative paths) solid arrows indicate significant paths ($P < 0.05$), and gray dashed arrows indicate
 330 non-significant paths ($P > 0.05$).

331 Table 2 The direct, indirect, total effects and the relative contribution of the present climate and
 332 paleoclimate change on AGB in the structural equation model

<u>Variables</u>	<u>Direct effect</u>	<u>Indirect effect</u>	<u>Total effect</u>	<u>Relative contribution (%)</u>
<u>AI</u>	<u>0.37</u>	<u>0.12</u>	<u>0.49</u>	<u>80.06</u>
<u>MAT</u>	<u>0</u>	<u>-0.07</u>	<u>-0.07</u>	<u>10.81</u>
<u>AMAT_{lgm}</u>	<u>0</u>	<u>-0.03</u>	<u>-0.03</u>	<u>4.95</u>
<u>AMAT_{mid}</u>	<u>0</u>	<u>-0.03</u>	<u>-0.03</u>	<u>4.18</u>

333 **4. Discussion**

334 **4.1 The effects of different dimensions of biodiversity on aboveground biomass**

335 Our research showed that, without considering the influence of climatic conditions, taxonomic
336 diversity was the most important factor influencing the aboveground biomass of Mongolian Plateau
337 grasslands, followed by phylogenetic diversity and functional diversity. However, a large number of
338 studies have emphasized the importance of phylogenetic diversity and functional diversity for
339 ecosystem functions (Srivastava et al., 2012; Cadotte et al., 2008; Swenson et al., 2012). For example,
340 research on subtropical forests in northern China showed that functional diversity is more important
341 than taxonomic diversity in controlling aboveground biomass, which is related to the acquisitive
342 resource use strategy of functional traits (Hanif et al., 2019). However, in Mongolian Plateau
343 grasslands taxonomic diversity may be more critical. Selection effects and compensatory effects are
344 the main mechanisms through which biodiversity impacts on biomass. The grassland community
345 structure of the Mongolian Plateau is relatively simple, and biomass mainly depends on the number
346 or dominance of dominant species. However, functional traits considered in this study, such as leaf
347 size and plant height, do not have a strong impact on biomass. The Mongolian Plateau is located in
348 an arid and semi-arid area, and drought-tolerant species invest a lot in seed resources, leaf structure
349 and root biomass (Wolf et al., 2021). Studies have shown that these traits are negatively correlated
350 with aboveground biomass, while functional traits consistent with growth strategies, such as leaf
351 nitrogen concentration and photosynthetic utilization efficiency, can promote aboveground biomass
352 (Wolf et al., 2021).

353 Secondly, phylogenetic diversity also had a significant impact on aboveground biomass.

354 Phylogenetic diversity contains the evolutionary information of species ecology. It is influenced by

355 the average degree of correlation between species and the number of existing species, and can reflect
356 the evolutionary sequence of characteristics related to habitat and resource utilization (Srivastava et
357 al., 2012). If species make better use of all available resources, productivity may increase. Therefore,
358 our research suggests that protecting taxonomic diversity and conserving evolutionarily different
359 species are equally important in effectively maintaining high-yielding communities.

360 **4.2 Relationship between paleoclimate anomalies and current biodiversity patterns**

361 Although a rich body of research has explored the factors affecting biodiversity patterns and the
362 relationships between biodiversity and biomass (Van Der Plas, 2019; Tilman et al., 2012) , it is
363 not clear how current climate and paleoclimate together affect the relationship, especially in
364 grassland communities. Here, our results showed that paleoclimate changes and present climate
365 factors were critical variables influencing contemporary biodiversity patterns. This illustrates the
366 importance of historical factors, such as evolution, as well as contemporary factors, such as climate,
367 in determining the geographic ranges of species and the accumulation of species in the region (Fine,
368 2015). Previous studies aiming to explain patterns of biodiversity have focused on the current
369 climate (Harrison et al., 2015), energy (Kreft and Jetz, 2007), soil attributions (Ulrich et al., 2014),
370 human activities (Hautier et al., 2015; Newbold et al., 2015) or water availability (Jiao et al., 2021;
371 Liu et al., 2021). Modern climate research on biodiversity and ecosystem functions has been widely
372 confirmed (Walther et al., 2002; Liu et al., 2021; Yang et al., 2011), but our research shows that the
373 pattern of biodiversity is also affected by past climates, and especially climate anomalies. Based on
374 climate-related traits, climate anomalies filtered the regional species pool, which in turn affected
375 contemporary biodiversity patterns.

376 Species richness is affected by the significant positive effects of AI, and the significant negative

377 effects of MAT and $AMAT_{mid}$. Jansson's research showed that the total number of vascular plant
378 species decreased with increasing temperature change since the LGM in mainland areas (Jansson,
379 2003), which is inconsistent with our results. We found that the species richness of Mongolian
380 Plateau grassland is more affected by temperature changes since the Mid-Holocene. In Europe, the
381 richness of widespread species was largely controlled by the modern climate, while the LGM climate
382 and climatic heterogeneity were codominant with modern climate as controls of richness for
383 restricted and intermediate species, which appear to still be associated with their glacial refugia
384 (Svenning and Skov, 2007b).

385 Phylogenetic diversity (PD) is usually used to quantify the uniqueness of a region's evolution.
386 On longer timescales, the effects of climate on speciation and extinction may cumulatively affect
387 current PD patterns (Svenning et al., 2015). Our research showed that PD was negatively correlated
388 with MAT and $AMAT_{mid}$, but not significantly related to temperature anomalies during the Last
389 Glacial Maximum. However, the PD of global forest ecosystems is positively associated with
390 temperature and precipitation and is negatively associated with Quaternary precipitation change
391 (Kubota et al., 2018). This indicates the importance of paleoprecipitation variation in shaping the
392 pattern of contemporary phylogenetic diversity in forest ecosystems, possibly by survival and
393 recolonization from refugia along hydrological gradients (Blonder et al., 2018). Additionally,
394 Quaternary glacial-interglacial climate oscillations led to the extinction of trees in many parts of the
395 globe, for example in Australia (Macphail et al., 1995) and Europe (Svenning, 2003), affecting the
396 PD of global forest ecosystems. ~~Liu et al.~~ It was showed-shown that herbaceous plants were mainly
397 affected by climate change since Mid-Holocene, while woody plants were mainly affected by climate
398 change since the LGM, which may be due to the differences in morphology of herbaceous and

399 woody plants (Liu et al., 2018b; Wang et al., 2021). Woody plants with large conduits are vulnerable
400 to drought or freezing and cause clogging of conduits (Zanne et al., 2014), and thus, ~~were more~~
401 ~~are~~ susceptible to cold Quaternary ~~climatic conditions~~~~climate oscillations~~. ~~In contrast, herbs~~~~Herbs~~
402 can mitigate the effects of extreme weather by producing underground buds and stems, or ~~by~~ wilting
403 ~~of~~ aboveground parts (Qian et al., 2017).

404 Our results suggested that functional richness was negatively related to Last Glacial Maximum
405 climate anomaly. This may be due to the intolerance of plant traits to past climatic conditions
406 (Bhagwat and Willis, 2008) and the postglacial migration lag (Svenning et al., 2008). In the Late
407 Quaternary, the Mongolian Plateau experienced severe alternating dry and wet climate fluctuations
408 (Tian et al., 2017), and likely experienced a non-random removal of functional combinations and
409 change in functional diversity (Ordonez and Svenning, 2017). Previous research has indicated that
410 the magnitude of past glacial-to-interglacial warming was sufficient at most locations across the
411 globe (including the Mongolian Plateau) to drive moderate to large changes in vegetation
412 composition and structure (Nolan et al., 2018). When climate changes, inappropriate species may be
413 quickly ruled out, and appropriate species may migrate more slowly, resulting in a negative
414 correlation between the paleoclimate anomaly and contemporary FD in the Mongolian Plateau. This
415 is also limited by the spreading ability or migration potential of plants after the glaciers retreated,
416 when species with inappropriate characteristics disappeared from the community and were not
417 replaced by other species (Blonder et al., 2018). Studies in Europe indicate that the lagging effect of
418 glacial-interglacial climate instability on functional diversity is >10,000 years (Ordonez and
419 Svenning, 2015).

420 **4.3 Effects of climate on the relationship between biodiversity and biomass**

421 The influence of climate on relationship between contemporary diversity and biomass may be
422 caused by [a combination of](#) climate change and human activities in the Mongolian Plateau grasslands.
423 In the eastern part of the Mongolian Plateau, the interval of 35230 to 25150 yr B.P. was a
424 forest-steppe landscape under humid conditions, represented by Pinaceae, Compositae and
425 Chenopodiaceae (Tian et al., 2017). Subsequently, the forest patches gradually shrank, the vegetation
426 transitioned to the steppe landscape, and drought began to intensify. During the Last Glacial
427 Maximum period, the vegetation types on the Mongolian Plateau were mainly polar, alpine and
428 temperate deserts (Ray and Adams, 2001), and the main plants were Compositae and
429 Chenopodiaceae. By the Holocene, xerophytes had declined sharply, and plants of humid climates
430 had increased. The humidity increased during the early Holocene, and generally humid conditions
431 lasted from 10,400 until 7000 yr BP and the region experienced the Lavliakan humid phase
432 (Lioubimtseva, 2004; Yin et al., 2011), with more hygrophilous species, such as Cyperaceae (Tian et
433 al., 2017). This climate condition was also conducive to the development and spread of Compositae
434 and Gramineae families in the steppes of Central Asia (Lioubimtseva, 2004). Since approximately
435 5200 yr BP to present, the climate has become more arid, with corresponding vegetation change and
436 strong aeolian activity (Yin et al., 2011).

437 As far as aboveground biomass is concerned, after taking into account climatic factors, it is
438 mainly directly affected by aridity index, while the three dimensions of biodiversity have no
439 significant impact on it, which is consistent with our previous research (Li et al., 2020). A
440 meta-analysis of 40 global grasslands plant communities showed that the response of plant diversity
441 to climate warming was an important factor in determining grassland aboveground biomass (Shao et
442 al., 2021). Similarly, paleoclimate changes also have no direct effects on aboveground biomass,

443 which indirectly affects aboveground biomass through regulating biodiversity. This is mainly
444 because paleoclimatic filtering can alter the historical legacies of species and functional diversity
445 (removal or accumulation over time of species and functional states) through evolutionary, extinction,
446 or recolonization (Svenning et al., 2015). Legacies in the species and functional traits pool further
447 affect ecosystem function, leading to changes in biomass (Svenning et al., 2015).

448 In addition, the relationship between biodiversity and biomass is also affected by human
449 activities (Buisson et al., 2022). Human history has become profoundly intertwined with grassland
450 communities, from the evolution of the *Homo* in savannas 2 million years ago (~~Ma~~) to the emergence
451 of species domestication in agricultural societies 10,000 years ago (Strömberg and Staver, 2022).
452 Palaeoecological and archaeological evidence shows that people have shaped most of terrestrial
453 nature for at least 12,000 years, including burning, hunting, cultivation and domestication (Ellis et al.,
454 2021). In the Mongolian Plateau, especially in the Inner Mongolia region of northern China, grazing
455 appeared at ~5.7–5.5 ka B.P. and further intensified again after ~4.2–4.0 ka B.P. (Huang et al., 2021).
456 A large number of studies have shown that the intensification of grazing activities also causes
457 changes in biodiversity and biomass (Liang et al., 2021; Herrero-Jáuregui and Oesterheld, 2018;
458 D'Áz et al., 2007). Therefore, there is an urgent need for future research to develop systematic
459 analytical frameworks to assess the importance of paleoclimate and climate change in shaping
460 contemporary biodiversity and ecosystems, particularly under the influence of human activities.

461 **5. Conclusion**

462 We conclude that unique evolutionary and ecological histories played key roles in explaining
463 biodiversity patterns in the Mongolian Plateau. When climatic conditions are not considered,
464 taxonomic diversity, phylogenetic diversity and functional diversity all have effects on community

465 biomass, and taxonomic diversity has a more obvious impact on biomass. However, the climate after
466 the Last Glacial Maximum has left a strong legacy affecting contemporary biodiversity patterns.
467 These findings extend our understanding of the spatial and temporal scale of climate effects on
468 biodiversity and aboveground biomass, providing additional confidence that the paleoclimate had a
469 key role in shaping contemporary biodiversity patterns.

470

471 **Authors' Contributions**

472 Z. Li analyzed the data and wrote the first draft of the manuscript. Zh. Li, J. Zhang, G. Han, F.Y. Li
473 and C. Liang contributed to later versions of the manuscript and all authors provided constructive
474 comments to improve the manuscript. F.Y. Li and C. Liang designed and led the field survey. Z. Li,
475 Zh. Li, X. Tong, J. Zhang, L. Dong, Y. Zheng, B. Miao, W. Ma, L. Zhao, L. Wang, L. Wen, C. Liang
476 and F.Y. Li conducted the experiment and collected the data.

477 **Data available**

478 The data that support the findings of this study are available on request from the corresponding
479 author.

480 **Conflict of interest statement**

481 The authors declare that they have no conflict of interest.

482 **Acknowledgements**

483 This study was supported by National Natural Science Foundation of China (31960261), the Basic
484 Scientific Research Foundation Special Project of the Institute of Water Resources and Hydropower
485 Research (MK2023J01), Science and Technology Program of Inner Mongolia Autonomous Region
486 of China (2020GG0126), the Natural Science Foundation of Inner Mongolia Autonomous Region of

487 China (2020MS03026), Major Science and Technology Project in Inner Mongolia Autonomous
488 Region (zdzx2018020), Inner Mongolia Autonomous Region Science and Technology Achievement
489 Transformation Project (2019CG069), the Innovative Team of the Ministry of Education of China
490 (IRT_17R59), and Yinshanbeilu Grassland Eco-hydrology National Observation and Research
491 Station, China Institute of Water Resources and Hydropower Research (No. YSS202108).

492 **References**

- 493 Archer, E.: rfPermute: Estimate Permutation p-Values for Random Forest Importance Metrics, 2016.
- 494 Avolio, M. L., Komatsu, K. J., Collins, S. L., Grman, E., Koerner, S. E., Tredennick, A. T., Wilcox, K. R.,
495 Baer, S., Boughton, E. H., Britton, A. J., Foster, B., Gough, L., Hovenden, M., Isbell, F., Jentsch, A., Johnson,
496 D. S., Knapp, A. K., Kreyling, J., Langley, J. A., Lortie, C., McCulley, R. L., McLaren, J. R., Reich, P. B.,
497 Seabloom, E. W., Smith, M. D., Suding, K. N., Suttle, K. B., and Tognetti, P. M.: Determinants of community
498 compositional change are equally affected by global change, *Ecol Lett*, 10.1111/ele.13824, 2021.
- 499 Bhagwat, S. A. and Willis, K. J.: Species persistence in northerly glacial refugia of Europe: a matter of chance
500 or biogeographical traits?, *Biogeogr*, 35, 464-482, 10.1111/j.1365-2699.2007.01861.x, 2008.
- 501 Blonder, B., Enquist, B. J., Graae, B. J., Kattge, J., Maitner, B. S., Morueta-Holme, N., Ordonez, A., Simova,
502 I., Singarayer, J., Svenning, J. C., Valdes, P. J., and Violle, C.: Late Quaternary climate legacies in
503 contemporary plant functional composition, *Glob Change Biol*, 24, 4827-4840, 10.1111/gcb.14375, 2018.
- 504 Breiman, L., Cutler, A., Liaw, A., and MatthewWiener: Package ‘randomForest’: Breiman and Cutler's
505 Random Forests for Classification and Regression, 2018.
- 506 Buisson, E., Archibald, S., Fidelis, A., and Suding, K. N.: Ancient grasslands guide ambitious goals in
507 grassland restoration, *Science*, 377, 594-598, 10.1126/science.abo4605, 2022.
- 508 Butler, E. E., Datta, A., Flores-Moreno, H., Chen, M., Wythers, K. R., Fazayeli, F., Banerjee, A., Atkin, O. K.,
509 Kattge, J., Amiaud, B., Blonder, B., Boenisch, G., Bond-Lamberty, B., Brown, K. A., Byun, C., Campetella,
510 G., Cerabolini, B. E. L., Cornelissen, J. H. C., Craine, J. M., Craven, D., de Vries, F. T., D áz, S., Domingues,
511 T. F., Forey, E., González-Melo, A., Gross, N., Han, W., Hattingh, W. N., Hickler, T., Jansen, S., Kramer, K.,
512 Kraft, N. J. B., Kurokawa, H., Laughlin, D. C., Meir, P., Minden, V., Niinemets, Ü., Onoda, Y., Peñuelas, J.,
513 Read, Q., Sack, L., Schamp, B., Soudzilovskaia, N. A., Spasojevic, M. J., Sosinski, E., Thornton, P. E.,
514 Valladares, F., van Bodegom, P. M., Williams, M., Wirth, C., and Reich, P. B.: Mapping local and global
515 variability in plant trait distributions, *P Natl Acad Sci USA*, 114, E10937-E10946, 10.1073/pnas.1708984114,
516 2017.
- 517 Cadotte, M. W., Cardinale, B. J., and Oakley, T. H.: Evolutionary history and the effect of biodiversity on
518 plant productivity, *P Natl Acad Sci USA*, 105, 17012-17017, 10.1073/pnas.0805962105, 2008.

519 Cavender-Bares, J., Kozak, K. H., Fine, P. V., and Kembel, S. W.: The merging of community ecology and
520 phylogenetic biology, *Ecol Lett*, 12, 693-715, 10.1111/j.1461-0248.2009.01314.x, 2009.

521 Cutler, D. R., Edwards, J. T. C., Beard, K. H., Cutler, A., Hess, K. T., Gibson, J., and Lawler, J. J.: Random
522 forest for classification ecology, *Ecology*, 88, 2783-2792, 10.1890/07-0539.1 2007.

523 Diaz, S., Kattge, J., Cornelissen, J. H., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C.,
524 Prentice, I. C., Garnier, E., Bonisch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J.,
525 Gillison, A. N., Zanne, A. E., Chave, J., Wright, S. J., Sheremet'ev, S. N., Jactel, H., Baraloto, C., Cerabolini,
526 B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J. S., Gunther, A., Falczuk, V., Ruger, N.,
527 Mahecha, M. D., and Gorne, L. D.: The global spectrum of plant form and function, *Nature*, 529, 167-171,
528 10.1038/nature16489, 2016.

529 D'Áz, S., Lavorel, S., McIntyre, S. U. E., Falczuk, V., Casanoves, F., Milchunas, D. G., Skarpe, C., Rusch, G.,
530 Sternberg, M., Noy-Meir, I., Landsberg, J., Zhang, W. E. I., Clark, H., and Campbell, B. D.: Plant trait
531 responses to grazing - a global synthesis, *Glob Change Biol*, 13, 313-341, 10.1111/j.1365-2486.2006.01288.x,
532 2007.

533 Eiserhardt, W. L., Borchsenius, F., Plum, C. M., Ordonez, A., and Svenning, J. C.: Climate-driven extinctions
534 shape the phylogenetic structure of temperate tree floras, *Ecol Lett*, 18, 263-272, 10.1111/ele.12409, 2015.

535 Ellis, E. C., Gauthier, N., Klein Goldewijk, K., Bliege Bird, R., Boivin, N., Diaz, S., Fuller, D. Q., Gill, J. L.,
536 Kaplan, J. O., Kingston, N., Locke, H., McMichael, C. N. H., Ranco, D., Rick, T. C., Shaw, M. R., Stephens,
537 L., Svenning, J. C., and Watson, J. E. M.: People have shaped most of terrestrial nature for at least 12,000
538 years, *P Natl Acad Sci USA*, 118, 10.1073/pnas.2023483118, 2021.

539 Esquivel-Muelbert, A., Baker, T. R., Dexter, K. G., Lewis, S. L., Brienen, R. J. W., Feldpausch, T. R., Lloyd, J.,
540 Monteagudo-Mendoza, A., Arroyo, L., Alvarez-Davila, E., Higuchi, N., Marimon, B. S., Marimon-Junior, B.
541 H., Silveira, M., Vilanova, E., Gloor, E., Malhi, Y., Chave, J., Barlow, J., Bonal, D., Davila Cardozo, N.,
542 Erwin, T., Fauset, S., Herault, B., Laurance, S., Poorter, L., Qie, L., Stahl, C., Sullivan, M. J. P., Ter Steege, H.,
543 Vos, V. A., Zuidema, P. A., Almeida, E., Almeida de Oliveira, E., Andrade, A., Vieira, S. A., Aragao, L.,
544 Araujo-Murakami, A., Arets, E., Aymard, C. G., Baraloto, C., Camargo, P. B., Barroso, J. G., Bongers, F.,
545 Boot, R., Camargo, J. L., Castro, W., Chama Moscoso, V., Comiskey, J., Cornejo Valverde, F., Lola da Costa,
546 A. C., Del Aguila Pasquel, J., Di Fiore, A., Fernanda Duque, L., Elias, F., Engel, J., Flores Llampazo, G.,
547 Galbraith, D., Herrera Fernandez, R., Honorio Coronado, E., Hubau, W., Jimenez-Rojas, E., Lima, A. J. N.,
548 Umetsu, R. K., Laurance, W., Lopez-Gonzalez, G., Lovejoy, T., Aurelio Melo Cruz, O., Morandi, P. S., Neill,
549 D., Nunez Vargas, P., Pallqui Camacho, N. C., Parada Gutierrez, A., Pardo, G., Peacock, J., Pena-Claros, M.,
550 Penuela-Mora, M. C., Petronelli, P., Pickavance, G. C., Pitman, N., Prieto, A., Quesada, C., Ramirez-Angulo,
551 H., Rejou-Mechain, M., Restrepo Correa, Z., Roopsind, A., Rudas, A., Salomao, R., Silva, N., Silva Espejo, J.,
552 Singh, J., Stropp, J., Terborgh, J., Thomas, R., Toledo, M., Torres-Lezama, A., Valenzuela Gamarra, L., van de
553 Meer, P. J., van der Heijden, G., van der Hout, P., Vasquez Martinez, R., Vela, C., Vieira, I. C. G., and Phillips,
554 O. L.: Compositional response of Amazon forests to climate change, *Glob Change Biol*, 25, 39-56,
555 10.1111/gcb.14413, 2019.

556 Fang, J.-y., Wang, X.-p., Shen, Z.-h., Tang, Z.-y., He, J.-s., Yu, D., Jiang, Y., Wang, Z.-h., Zheng, C.-y., Zhu,
557 J.-l., and Guo, Z.-d.: Methods and protocols for plant community inventory, *Biodiversity Science*, 17, 533-548,

558 10.3724/sp.J.1003.2009.09253, 2009.

559 Fine, P. V. A.: Ecological and Evolutionary Drivers of Geographic Variation in Species Diversity, *Annu Rev*
560 *Ecol Evol S*, 46, 369-392, 10.1146/annurev-ecolsys-112414-054102, 2015.

561 Flynn, D. F. B., Mirotchnick, N., Jain, M., Palmer, M. I., and Naeem, S.: Functional and phylogenetic
562 diversity as predictors of biodiversity-ecosystem-function relationships, *Ecology*, 92, 1573-1581,
563 10.1890/10-1245.1, 2011.

564 Fordham, D. A., Jackson, S. T., Brown, S. C., Huntley, B., Brook, B. W., Dahl-Jensen, D., Gilbert, M. T. P.,
565 Otto-Bliesner, B. L., Svensson, A., Theodoridis, S., Wilmshurst, J. M., Buettel, J. C., Canteri, E., McDowell,
566 M., Orlando, L., Pilowsky, J., Rahbek, C., and Nogues-Bravo, D.: Using paleo-archives to safeguard
567 biodiversity under climate change, *Science*, 369, 10.1126/science.abc5654, 2020.

568 Hanif, M. A., Yu, Q., Rao, X., and Shen, W.: Disentangling the Contributions of Plant Taxonomic and
569 Functional Diversities in Shaping Aboveground Biomass of a Restored Forest Landscape in Southern China,
570 *Plants (Basel)*, 8, 612, 10.3390/plants8120612, 2019.

571 Harrison, S. P., Gornish, E. S., and Copeland, S.: Climate-driven diversity loss in a grassland community, *P*
572 *Natl Acad Sci USA*, 112, 8672-8677, 10.1073/pnas.1502074112, 2015.

573 Hautier, Y., Tilman, D., Isbell, F., Seabloom, E. W., Borer, E. T., and Reich, P. B.: Anthropogenic
574 environmental changes affect ecosystem stability via biodiversity, *Science*, 348, 336-340,
575 10.1126/science.aaa1788, 2015.

576 Herrero-Jáuregui, C. and Oesterheld, M.: Effects of grazing intensity on plant richness and diversity: a
577 meta-analysis, *Oikos*, 127, 757-766, 10.1111/oik.04893, 2018.

578 Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D. M.,
579 Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., J., V., and Wardle, D. A.: Effects of
580 biodiversity on ecosystem functioning: a consensus of current knowledge, *Ecol Monogr*, 75, 3-35,
581 10.1890/04-0922 2005.

582 Huang, X., Zhang, J., Ren, L., Zhang, S., and Chen, F.: Intensification and Driving Forces of Pastoralism in
583 Northern China 5.7 ka Ago, *Geophys Res Lett*, 48, e2020GL092288, 2021.

584 Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W. S., Reich, P. B., Scherer-Lorenzen, M., Schmid,
585 B., Tilman, D., van Ruijven, J., Weigelt, A., Wilsey, B. J., Zavaleta, E. S., and Loreau, M.: High plant diversity
586 is needed to maintain ecosystem services, *Nature*, 477, 199-202, 10.1038/nature10282, 2011.

587 Jansson, R.: Global patterns in endemism explained by past climatic change, *P Roy Soc B-Biol Sci*, 270,
588 583-590, 10.1098/rspb.2002.2283, 2003.

589 Jiao, W., Wang, L., Smith, W. K., Chang, Q., Wang, H., and D'Odorico, P.: Observed increasing water
590 constraint on vegetation growth over the last three decades, *Nat Commun*, 12, 3777,
591 10.1038/s41467-021-24016-9, 2021.

592 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder,
593 H. P., and Kessler, M.: Climatologies at high resolution for the earth's land surface areas, *Scientific Data*, 4,
594 170122, 10.1038/sdata.2017.122, 2017.

595 Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., Blomberg, S. P.,
596 and Webb, C. O.: Picante: R tools for integrating phylogenies and ecology, *Bioinformatics*, 26, 1463-1464,
597 10.1093/bioinformatics/btq166, 2010.

598 Kissling, W. D., Eiserhardt, W. L., Baker, W. J., Borchsenius, F., Couvreur, T. L., Balslev, H., and Svenning, J.
599 C.: Cenozoic imprints on the phylogenetic structure of palm species assemblages worldwide, *P Natl Acad Sci*
600 *USA*, 109, 7379-7384, 10.1073/pnas.1120467109, 2012.

601 Kreft, H. and Jetz, W.: Global patterns and determinants of vascular plant diversity, *P Natl Acad Sci USA*, 104,
602 5925-5930, 10.1073/pnas.0608361104, 2007.

603 Kubota, Y., Kusumoto, B., Shiono, T., Ulrich, W., and Duarte, L.: Environmental filters shaping angiosperm
604 tree assembly along climatic and geographic gradients, *J Veg Sci*, 29, 607-618, 10.1111/jvs.12648, 2018.

605 Laliberté, E., Legendre, P., and Shipley, B.: Package 'FD': Measuring functional diversity (FD) from multiple
606 traits, and other tools for functional ecology, 2015.

607 Lefcheck, J.: Package 'piecewiseSEM': Piecewise Structural Equation Modeling, 2020.

608 Lefcheck, J. S. and Freckleton, R.: piecewiseSEM: Piecewise structural equation modelling in r for ecology,
609 evolution, and systematics, *Methods Ecol Evol*, 7, 573-579, 10.1111/2041-210x.12512, 2015.

610 Li, Z., Li, Z., Tong, X., Zhang, J., Dong, L., Zheng, Y., Ma, W., Zhao, L., Wang, L., Wen, L., Dang, Z.,
611 Tuvshintogtokh, I., Liang, C., and Li, F. Y.: Climatic humidity mediates the strength of the species richness–
612 biomass relationship on the Mongolian Plateau steppe, *Sci Total Environ*, 718, 137252,
613 10.1016/j.scitotenv.2020.137252, 2020.

614 Li, Z., Liang, M., Li, Z., Mariotte, P., Tong, X., Zhang, J., Dong, L., Zheng, Y., Ma, W., Zhao, L., Wang, L.,
615 Wen, L., Tuvshintogtokh, I., Gornish, E. S., Dang, Z., Liang, C., Li, F. Y., and Schöb, C.: Plant functional
616 groups mediate effects of climate and soil factors on species richness and community biomass in grasslands of
617 Mongolian Plateau, *J Plant Ecol*, 14, 679-691, 10.1093/jpe/rtab021, 2021.

618 Liang, M., Liang, C., Hautier, Y., Wilcox, K. R., and Wang, S.: Grazing-induced biodiversity loss impairs
619 grassland ecosystem stability at multiple scales, *Ecol Lett*, 24, 2054-2064, 10.1111/ele.13826, 2021.

620 Lioubimtseva, E.: Climate change in arid environments: revisiting the past to understand the future, *Prog Phys*
621 *Geog*, 28, 502-530, 10.1191/0309133304pp422oa, 2004.

622 Liu, D., Zhang, C., Ogaya, R., Fernández-Martínez, M., Pugh, T. A. M., and Peñuelas, J.: Increasing climatic
623 sensitivity of global grassland vegetation biomass and species diversity correlates with water availability, *New*
624 *Phytol*, 230, 1761-1771, 10.1111/nph.17269, 2021.

625 Liu, H., Mi, Z., Lin, L., Wang, Y., Zhang, Z., Zhang, F., Wang, H., Liu, L., Zhu, B., Cao, G., Zhao, X., Sanders,

626 N. J., Classen, A. T., Reich, P. B., and He, J. S.: Shifting plant species composition in response to climate
627 change stabilizes grassland primary production, *P Natl Acad Sci USA*, 115, 4051-4056,
628 10.1073/pnas.1700299114, 2018a.

629 Liu, Y., Su, X., Shrestha, N., Xu, X., Wang, S., Li, Y., Wang, Q., Sandanov, D., and Wang, Z.: Effects of
630 contemporary environment and Quaternary climate change on drylands plant diversity differ between growth
631 forms, *Ecography*, 42, 334-345, 10.1111/ecog.03698, 2018b.

632 MacPhail, M. K., Colhoun, E. A., and Fitzsimons, S. J.: Key Periods in the Evolution of the Cenozoic
633 Vegetation and Flora in Western Tasmania: the Late Pliocene, *Australian Journal of Botany*, 43,
634 10.1071/bt9950505, 1995.

635 Maestre, F. T., Benito, B. M., Berdugo, M., Concostrina-Zubiri, L., Delgado-Baquerizo, M., Eldridge, D. J.,
636 Guirado, E., Gross, N., Kefi, S., Le Bagousse-Pinguet, Y., Ochoa-Hueso, R., and Soliveres, S.: Biogeography
637 of global drylands, *New phytol*, 231, 540-558, 10.1111/nph.17395, 2021.

638 Mayle, F. E., Burn, M. J., Power, M., and Urrego, D. H.: Past Climate Variability in South America and
639 Surrounding Regions, Springer2009.

640 Mottl, O., Flantua, S. G. A., Bhatta, K. P., Felde, V. A., Giesecke, T., Goring, S., Grimm6, E. C., Haberle, S.,
641 Hooghiemstra, H., Ivory, S., Kuneš, P., Wolters, S., Seddon, A. W. R., and Williams, J. W.: Global acceleration
642 in rates of vegetation change over the past 18,000 years, *Science*, 372, 860-864, 10.1126/science.abg1685,
643 2021.

644 Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., Borger, L., Bennett, D. J.,
645 Choimes, A., Collen, B., Day, J., De Palma, A., Diaz, S., Echeverria-Londono, S., Edgar, M. J., Feldman, A.,
646 Garon, M., Harrison, M. L., Alhousseini, T., Ingram, D. J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L.,
647 Kleyer, M., Correia, D. L., Martin, C. D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H. R., Purves, D. W.,
648 Robinson, A., Simpson, J., Tuck, S. L., Weiher, E., White, H. J., Ewers, R. M., Mace, G. M., Scharlemann, J.
649 P., and Purvis, A.: Global effects of land use on local terrestrial biodiversity, *Nature*, 520, 45-50,
650 10.1038/nature14324, 2015.

651 Nolan, C., Overpeck, J. T., Allen, J. R. M., Anderson, P. M., Betancourt, J. L., Binney, H. A., Brewer, S., Bush,
652 M. B., Chase, B. M., Cheddadi, R., Djamali, M., Dodson, J., Edwards, M. E., Gosling, W. D., Haberle, S.,
653 Hotchkiss, S. C., Huntley, B., Ivory, S. J., Kershaw, A. P., Kim, S.-H., Latorre, C., Leydet, M., Lézine, A.-M.,
654 Liu, K.-B., Liu, Y., Lozhkin, A. V., McGlone, M. S., Marchant, R. A., Momohara, A., Moreno, P. I., Müller, S.,
655 Otto-Bliesner, B. L., Shen, C., Stevenson, J., Takahara, H., Tarasov, P. E., Tipton, J., Vincens, A., Weng, C.,
656 Xu, Q., Zheng, Z., and Jackson, S. T.: Past and future global transformation of terrestrial ecosystems under
657 climate change, *Science*, 361, 920-923, 10.1126/science.aan5360, 2018.

658 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R., O'Hara, R. B.,
659 Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H.: Package 'vegan': Community
660 Ecology Package, 2020.

661 Ordonez, A. and Svenning, J.-C.: Geographic patterns in functional diversity deficits are linked to
662 glacial-interglacial climate stability and accessibility, *Global Ecol Biogeogr*, 24, 826-837, 10.1111/geb.12324,

663 2015.

664 Ordonez, A. and Svenning, J. C.: Consistent role of Quaternary climate change in shaping current plant
665 functional diversity patterns across European plant orders, *Sci Rep-UK*, 7, 42988, 10.1038/srep42988, 2017.

666 Qian, H., Jin, Y., and Ricklefs, R. E.: Phylogenetic diversity anomaly in angiosperms between eastern Asia
667 and eastern North America, *P Natl Acad Sci USA*, 114, 11452-11457, 10.1073/pnas.1703985114, 2017.

668 Ray, N. and Adams, J. M.: A GIS-based Vegetation Map of the World at the Last Glacial Maximum
669 (25,000-15,000 BP). *Internet Archaeology*, 11, 1-44, 10.11141/ia.11.2, 2001.

670 Revelle, W.: Package ‘psych’: Procedures for Psychological, Psychometric, and Personality Research, 2021.

671 Sandel, B., Arge, L., Dalsgaard, B., Davies, R. G., Gaston, K. J., Sutherland, W. J., and Svenning, J. C.: The
672 influence of Late Quaternary climate-change velocity on species endemism, *Science*, 334, 660-664,
673 10.1126/science.1210173, 2011.

674 Seddon, A. W. R., Macias-Fauria, M., and Willis, K. J.: Climate and abrupt vegetation change in Northern
675 Europe since the last deglaciation, *Holocene*, 25, 25-36, 10.1177/0959683614556383, 2014.

676 Shao, J., Zhou, X., Groenigen, K. J., Zhou, G., Zhou, H., Zhou, L., Lu, M., Xia, J., Jiang, L., Hungate, B. A.,
677 Luo, Y., He, F., Thakur, M. P., and Mayfield, M.: Warming effects on grassland productivity depend on plant
678 diversity, *Global Ecol Biogeogr*, 31, 588-598, 10.1111/geb.13441, 2021.

679 Srivastava, D. S., Cadotte, M. W., MacDonald, A. A., Marushia, R. G., and Mirotnick, N.: Phylogenetic
680 diversity and the functioning of ecosystems, *Ecol Lett*, 15, 637-648, 10.1111/j.1461-0248.2012.01795.x, 2012.

681 Strömberg, C. A. E. and Staver, A. C.: The history and challenge of grassy biomes-Grassy biomes are >20
682 million years old but are undervalued and under threat today, *Science*, 377, 592-593, 10.1126/science.add1347,
683 2022.

684 Svenning, J.-C.: Deterministic Plio-Pleistocene extinctions in the European cool-temperate tree flora, *Ecol*
685 *Lett*, 6, 646-653, 10.1046/j.1461-0248.2003.00477.x, 2003.

686 Svenning, J.-C. and Skov, F.: Could the tree diversity pattern in Europe be generated by postglacial dispersal
687 limitation?, *Ecol Lett*, 10, 453-460, 10.1111/j.1461-0248.2007.01038.x, 2007a.

688 Svenning, J.-C. and Skov, F.: Ice age legacies in the geographical distribution of tree species richness in
689 Europe, *Global Ecol Biogeogr*, 16, 234-245, 10.1111/j.1466-822x.2006.00280.x, 2007b.

690 Svenning, J.-C., Normand, S., and Skov, F.: Postglacial dispersal limitation of widespread forest plant species
691 in northern Europe, *Ecography*, 31, 316-326, 10.1111/j.2008.0906-7590.05206.x, 2008.

692 Svenning, J.-C., Eiserhardt, W. L., Normand, S., Ordonez, A., and Sandel, B.: The Influence of Paleoclimate
693 on Present-Day Patterns in Biodiversity and Ecosystems, *Annu Rev Ecol Evol S*, 46, 551-572,
694 10.1146/annurev-ecolsys-112414-054314, 2015.

695 Swenson, N. G.: The role of evolutionary processes in producing biodiversity patterns, and the

696 interrelationships between taxonomic, functional and phylogenetic biodiversity, *Am J Bot*, 98, 472-480,
697 10.3732/ajb.1000289, 2011.

698 Swenson, N. G., Erickson, D. L., Mi, X., Bourg, N. A., Forero-Montaña, J., Ge, X., Howe, R., Lake, J. K., Liu,
699 X., Ma, K., Pei, N., Thompson, J., Uriarte, M., Wolf, A., Wright, S. J., Ye, W., Zhang, J., Zimmerman, J. K.,
700 and Kress, W. J.: Phylogenetic and functional alpha and beta diversity in temperate and tropical tree
701 communities, *Ecology*, 93, S112-S125, 2012.

702 The Angiosperm Phylogeny Group: An update of the Angiosperm Phylogeny Group classification for the
703 orders and families of flowering plants APG II, *Bot J Linn Soc*, 161, 105-121, 10.1111/boj.12385, 2009.

704 Tian, F., Wang, Y., Chi, Z., Liu, J., Yang, H., Jiang, N., and Tang, W.: Late Quaternary vegetation and climate
705 reconstruction based on pollen data from southeastern Inner Mongolia, China, *Rev Palaeobot Palyno*, 242,
706 33-42, 10.1016/j.revpalbo.2017.03.003, 2017.

707 Tilman, D., Reich, P. B., and Isbell, F.: Biodiversity impacts ecosystem productivity as much as resources,
708 disturbance, or herbivory, *P Natl Acad Sci USA*, 109, 10394-10397, 10.1073/pnas.1208240109 2012.

709 Trabucco, A. and Zomer, R. J.: Global Aridity Index and Potential Evapo-Transpiration (ET₀) Climate
710 Database v2, 10.6084/m9.figshare.7504448.v3, 2019.

711 Ulrich, W., Soliveres, S., Maestre, F. T., Gotelli, N. J., Quero, J. L., Delgado-Baquerizo, M., Bowker, M. A.,
712 Eldridge, D. J., Ochoa, V., Gozalo, B., Valencia, E., Berdugo, M., Escobar, C., Garcia-Gomez, M., Escudero,
713 A., Prina, A., Alfonso, G., Arredondo, T., Bran, D., Cabrera, O., Cea, A., Chaieb, M., Contreras, J., Derak, M.,
714 Espinosa, C. I., Florentino, A., Gaitan, J., Muro, V. G., Ghiloufi, W., Gomez-Gonzalez, S., Gutierrez, J. R.,
715 Hernandez, R. M., Huber-Sannwald, E., Jankju, M., Mau, R. L., Hughes, F. M., Miriti, M., Moneris, J.,
716 Muchane, M., Naseri, K., Pucheta, E., Ramirez-Collantes, D. A., Raveh, E., Romao, R. L., Torres-Diaz, C.,
717 Val, J., Veiga, J. P., Wang, D., Yuan, X., and Zaady, E.: Climate and soil attributes determine plant species
718 turnover in global drylands, *J Biogeogr*, 41, 2307-2319, 10.1111/jbi.12377, 2014.

719 Van der Plas, F.: Biodiversity and ecosystem functioning in naturally assembled communities, *Biol Rev*, 94,
720 1220-1245, 10.1111/brv.12499, 2019.

721 Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M.,
722 Hoegh-Guldberg, O., and Bairlein, F.: Ecological responses to recent climate change, *Nature*, 416, 389-395,
723 10.1038/416389a, 2002.

724 Wang, Q., Li, Y., Zou, D., Su, X., Cai, H., Luo, A., Jiang, K., Zhang, X., Xu, X., Shrestha, N., and Wang, Z.:
725 Phylogenetic niche conservatism and variations in species diversity–climate relationships, *Ecography*, 44,
726 1856-1868, 10.1111/ecog.05759, 2021.

727 Wolf, A. A., Funk, J. L., Selmants, P. C., Morozumi, C. N., Hernandez, D. L., Pasari, J. R., and Zavaleta, E. S.:
728 Trait-based filtering mediates the effects of realistic biodiversity losses on ecosystem functioning, *P Natl Acad
729 Sci USA*, 118, 10.1073/pnas.2022757118, 2021.

730 Yang, H., Wu, M., Liu, W., Zhang, Z. H. E., Zhang, N., and Wan, S.: Community structure and composition in
731 response to climate change in a temperate steppe, *Glob Change Biol*, 17, 452-465,

- 732 10.1111/j.1365-2486.2010.02253.x, 2011.
- 733 Ye, J. S., Delgado-Baquerizo, M., Soliveres, S., and Maestre, F. T.: Multifunctionality debt in global drylands
734 linked to past biome and climate, *Glob Change Biol*, 25, 2152-2161, 10.1111/gcb.14631, 2019.
- 735 Yin, Y., Liu, H., He, S., Zhao, F., Zhu, J., Wang, H., Liu, G., and Wu, X.: Patterns of local and regional grain
736 size distribution and their application to Holocene climate reconstruction in semi-arid Inner Mongolia, China,
737 *Palaeogeogr Palaeocl*, 307, 168-176, 10.1016/j.palaeo.2011.05.011, 2011.
- 738 Zanne, A. E., Tank, D. C., Cornwell, W. K., Eastman, J. M., Smith, S. A., FitzJohn, R. G., McGlenn, D. J.,
739 O'Meara, B. C., Moles, A. T., Reich, P. B., Royer, D. L., Soltis, D. E., Stevens, P. F., Westoby, M., Wright, I. J.,
740 Aarssen, L., Bertin, R. I., Calaminus, A., Govaerts, R., Hemmings, F., Leishman, M. R., Oleksyn, J., Soltis, P.
741 S., Swenson, N. G., Warman, L., and Beaulieu, J. M.: Three keys to the radiation of angiosperms into freezing
742 environments, *Nature*, 506, 89-92, 10.1038/nature12872, 2014.
- 743