

Associate Editor

Comments to the author:

Two reviewers have assessed your manuscript and provided useful feedback. One reviewer mentioned concerns about your methodology, please make sure to respond to specifically their comments in your revised manuscript.

We appreciate the comments. All general and specific comments from the two reviewers were carefully considered and taken care of. Specifically, reviewer 2 had concerns on the methodology. In the revised manuscript, we added detailed information about the setting up of the static chambers (including temperature control and chamber sealing), offered details about gas chromatography, and provided necessary background information of our study region. We also explained in the response why we excluded the effects of climatic factors and considered vegetation as a major influencing factor. As for the concern about upscaling site CO₂ flux to global CO₂ flux, comparison with CO₂ flux of global forest and verification of the carbon offset model, we have carefully considered the reviewer's suggestion and re-evaluated these parts. Considering these parts were only loosely connected to the main arguments of this manuscript, and we were not able to give comprehensive evaluation of the global riparian carbon uptake potential due to data limitation (e.g.: systematic differences in global riparian vegetation, soil and hydrology), we removed these parts and relevant sections in methodology (section 2.3) and result (section 3.5, section 3.6). The discussion sections were then substantially restructured corresponding to the change, focusing on two main results of our study: 1) increased carbon emission from riparian area due to flooding and 2) the role of post-disturbance survived vegetation in riparian carbon sequestration after the flooding period. At the same time, we took the reviewer 2's suggestion to use accumulated CO₂ for analysis and added a new section 2.2.4 "Annual riparian and river CO₂ emission calculation" to show how we use the measured CO₂ flux to evaluate the carbon balance of the whole region, with assumption about different flooding seasons, flooding characteristics, and riparian characteristics. This section not only shows how we calculate the accumulated CO₂ flux, but also shows how we evaluate whether the riparian area and fluvial area overall acted as a carbon sink or carbon source. The assumption we made for calculation have also been added at the beginning of the results part to make it more fluent to readers. We believe these changes make the main arguments of our manuscript more focused and the manuscript more coherent, consistent, and evidence-based.

Review 1 had main concerns that the terms used to describe the measurements in different seasons may lead to ambiguity. We agreed with the reviewer, so in the revised manuscript, we used "pre-flooding season", "flooding season", and "post-flooding season" to describe the measurements.

All other methodology, table, figure, language, unit and terminology-related problems raised by the two reviewers have been carefully considered and taken care of in the revised manuscript. See detailed responses to general and specific comments of the reviewers below.

Reviewer #1:

This study evaluates how carbon fluxes of river riparian areas seasonally respond to flooding by investigating the CO₂ and dissolved carbon fluxes during pre- and post-flooding seasons. The methods were generally well established and results were generally well discussed. In particular, authors discussed how vegetation and the post-flooding change of riparian vegetation species and richness affect/determines net carbon fluxes in river riparian areas. I think this could be a valuable contribution to this area.

We appreciate the positive comments on our manuscript. Below, we have addressed all issues raised by the reviewer. Please see below for details.

Note: The line numbers shown in the bracket in our response are referring to the line numbers in the revised manuscript with tracked changes.

One thing that might be ambiguous to audience is that authors use terms like “pre-flooding”, “flooding” and “post-flooding” to describe their measurements in different seasons. This might be a bit misleading because floods often refer to short-term events. Considering authors evaluated these carbon fluxes on a seasonal base, I’d suggest authors changes these expressions to pre-flooding seasons, flooding seasons and post-flooding seasons instead. I have only a few minor comments on the rest of the manuscript. See below,

Thanks for pointing this out. We agree that the original expressions can lead to misunderstanding. We have checked throughout the manuscript and used pre-flooding season, flooding season, and post-flooding season to describe our measurements. See below for a few examples,

“... we quantified the vertical CO₂ fluxes at the soil-air interface and water-air interface **during the flooding season and non-flooding seasons (pre-flooding season and post-flooding season)** based on two-year in-situ measurements along the Lijiang River.” (Line 124-126)

“...we find that the fluvial area turned from carbon sources in **pre-flooding season and during flooding season** to a carbon sink **in post-flooding season.**” (Line 323-325)

“The species richness index decreased from 2.945 **in pre-flooding season** to 1.695 **in post-flooding season** (Table 1, Table A3).” (Line 342-343)

Line 174: in different seasons

Corrected accordingly. See below,

“...we directly measured the CO₂ fluxes in the riparian area with and without vegetation (bare soil) **in different seasons.**” (Line 269)

Line 175: we first reviewed the diel cycle.

In the revised manuscript, the paragraph has been restructured and the idea is conveyed in another way. See below,

“We assume that diel CO₂ flux follows similar patterns as measured on the selected days during the pre-flooding season and post-flooding season. Based on this assumption, we compared the diel CO₂ flux of pre-flooding season and post-flooding season. In order to evaluate the effect of vegetation on riparian CO₂ flux, we directly measured the CO₂ fluxes in the riparian area with and without vegetation (bare soil) in different seasons. Significant diel variations in CO₂ fluxes were observed in the riparian area...” (Line 266-269)

Line 176: change “the terrestrial area” to “riparian soils” or “riparian area”, same below

We agree that “riparian area” or “riparian soils” would be more appropriate. We have changed all “terrestrial area” into “riparian area”, which indicates the measurements on land. See examples below,

“In April, carbon sequestration in the **riparian area** with vegetation was observed between 10:00 to 14:00 hours; while in October, the carbon sequestration is observed between 6:00 to 14:00 hours (Fig. 1).” (Line 275-277)

“The **riparian area** is composed of vegetation area and bare soil.” (Line 291)

Line 183: “terrestrial area with vegetation” to “riparian area with vegetation”

Relevant expressions have been modified accordingly.

“Thus, in post-flooding season, **the riparian area with vegetation** sequestered carbon for a longer time. Indeed, the vegetation area's all-day CO₂ flux was 0.358 g·m⁻² d⁻¹ in April...” (Line 277-278)

Line 200-203: the sentence repeat line 186-188. Consider delete one of them

Thanks. The repeated sentence is deleted, and the paragraph is restructured. See below,

“The riparian area is composed of vegetation area and bare soil. During the field investigation, we found the vegetation coverage in Lijiang riparian area is about 60%. Using vegetation coverage as the weight, we can get the accumulated CO₂ flux of riparian area (Section 2.2.4, equation (3)). Within a day, the carbon sequestration in the riparian area peaked at 14:00 (April: -62.680 mg·m⁻² h⁻¹; October: -68.813 mg·m⁻² h⁻¹), and the maximum carbon emission occurred at 18:00 (April: 36.347 mg·m⁻² h⁻¹; October: 14.110 mg·m⁻² h⁻¹; Fig. 1). In both April and October, the all-day carbon fluxes in the riparian area were negative, indicating that the riparian area acted as a carbon sink in non-flooding season (April: -0.156 g·m⁻² d⁻¹, October: -0.500 g·m⁻² d⁻¹). The carbon uptake in October, which represented the post-flooding season, was higher. Overall, we found that in the post-flooding season, the riparian vegetation can sequester CO₂ for a longer time and

fix a higher amount of carbon. **Thus, even though the all-day CO₂ flux of bare soil changed from -0.927 g·m⁻² d⁻¹ to -0.231 g·m⁻² d⁻¹, showing a reduced capacity of carbon sequestration after flooding, the whole riparian area still turned out to be a carbon sink in the post-flooding season.**” (Line 291-309)

Line 241: is the water-air carbon flux (Fig. 2b).

The figure number is added to the sentence. See below,

“During the flooding, the riparian areas with and without vegetation were submerged, so only the carbon fluxes from the water-air interfaces were measured **(Fig.1b)**.” (Line 321-322)

Line 285-286: no need to capitalize first letters

Corrected accordingly. See below,

“We measured total organic carbon (TOC) and total inorganic carbon (TIC) in riparian soils and the fluvial area during different periods.” (Line 367)

Line 313: needs further analysis

Thanks. The sentence has been modified. See below,

“Also, besides the contribution of recovered vegetation, our data shows that bare soil also contributes to the carbon neutralization, but the mechanism for bare soil to capture carbon still **needs further analysis**.” (Line 498-500)

Line 332: the riparian zone thus has ...

In response to comments from reviewer #2, we have re-organized the results and discussion sections of the manuscript. This section (Section 3.6) has been removed from the revised manuscript. See also response to Reviewer#2 on Section 3.6.

Reviewer #2:

The authors studied the effects of flooding on carbon emissions and sinks in the riparian zone. They measured field CO₂ flux and developed a model to simulate the riparian carbon emissions. Although multiple methods applied, I am not convinced by the data yet. The introduction is not comprehensive, methods lack some important information, the results and discussions are not convincing. At the current stage, I think it is not suitable for publishing. There are also some comments need to be considered:

We appreciate these comments. In the revised manuscript, we have substantially revised the introduction, methods, and result and discussion sections.

In the introduction, we added contents related to the anoxic production of methane (see response to the comment on line 39-44 below) and particularly the impacts of riparian soil property and vegetation on riparian carbon fluxes and related modeling works (see response to the comments on introduction below), and therefore offered a more thorough overview of the recent progresses on flooding and riparian carbon fluxes.

In methods, we added necessary information (e.g.: sediment and soil) of our study region in section 2.1, added details about the setting up of the static chambers in section 2.2.1, and offered details about the gas chromatography used in this study in the new section 2.2.2. We removed section 2.3 (model scenario prediction method) in response to the restructure of results and discussion (see response to the comments on section 2.3.2 and section 3.5 for the restructure). We then added a new section 2.2.4 to explain how we get the accumulated CO₂ flux and calculate the annual CO₂ emission of riparian area and fluvial area as a whole (see response to the comments on equation 2, line 189-194 below).

In results and discussion, considering the estimation at global scale and the comparison with global forest CO₂ flux are only loosely connected to the main arguments of the manuscript and that we were not able to give a comprehensive evaluation of the carbon uptake potential for global riparian areas, we removed related sections in result (section 3.5, 3.6) and discussion (section 4.4) in the revised manuscript. Following this change, the discussion section has been substantially restructured to focus on two main results of our study: 1) increased carbon emission from riparian area due to flooding and 2) the role of post-disturbance survived vegetation in riparian carbon sequestration after the flooding period (see response to the comments on section 2.3.2 and section 3.5 below).

The abstract and conclusion part were revised correspondingly.

We believe these changes have greatly improved the quality of present manuscript, and made our manuscript more compact, organized and coherently presented.

Note: The line numbers shown in the bracket in our response are referring to the line numbers in the revised manuscript with tracked changes.

Please correct the reference “Xunhua et al., 1998”. The first name and last name were switched.

Thanks for pointing this out. We have corrected the reference in the revised manuscript. See below,

“The increasing atmospheric CO₂ originating from fossil fuel combustion and industrial activities can be regulated by plant metabolism (photosynthesis and respiration) and soil microbial activities (**Zheng et al., 1998**).” (Line 37)

Lines 39-44, flooding submergence could cause anoxic conditions, which favors the reduction reactions. Thus, methane should be released more than carbon dioxide and are more important?

We agree with the reviewer that flooding submergence causes anoxic conditions which could stimulate methane production and emission. In revised version of the manuscript, we have modified the second paragraph and added the following parts related to methane production in riparian zones to the introduction. See below for details.

“Riparian zones are of great importance in carbon cycling, which is associated with the production and consumption of CO₂ and methane (CH₄) (Zhang et al., 2016; Allen et al., 2007; Liu et al., 2021).” (Line 43-45)

“Also, the short-term anaerobic conditions caused by flooding may increase the production of methane because of the strengthened methanogenesis in riparian soils (Hassanzadeh et al., 2019; Hondula et al., 2021; Morse et al., 2012; Le Mer and Roger, 2001; Thorp et al., 2006).” (Line 55-57)

“This raises the possibility of elevated carbon (including methane and carbon dioxide) emissions and reduced carbon sequestration from riparian zones...” (Line 70-71)

“Considering an overall small contribution of CH₄ to the carbon balance of riparian zones (Liu et al., 2021; Vidon et al., 2019), only CO₂ fluxes were measured in analysis.” (Line 126-128)

Line 58, a space was missed in the unit mg·m⁻² h⁻¹.

Corrected accordingly.

“Liu et al. (2021) demonstrated that high plant and soil respiration in riparian wetlands lead to large amounts of CO₂ emission in wet season (335-2790 **mg·m⁻² h⁻¹**) than in dry season (72 - 387 **mg·m⁻² h⁻¹**).” (Line 53-55)

Introduction is too short to summarize the recent research progress in riparian carbon cycle. More information is needed about riparian soil properties, CO₂ and CH₄ emissions, vegetation, and some modelling work.

Thanks for the comments. In the revised version of the manuscript, we have substantially changed the introduction section of the manuscript. We added more discussion related to riparian CO₂ and CH₄ emissions, how flooding disturbance would be affected by the flooding characteristics and the properties of riparian zone (e.g., soil properties, vegetation), and some modeling works on flooding and riparian carbon fluxes (in paragraph 2-5, Line 43-78, 114-122). At the same time, we removed contents that are only loosely related to the main content of the paper (e.g., line 45-51 or 3rd paragraph of the original manuscript, which discussed the impact of disturbances in systems other than the riparian system). We believe these changes offered a better overview of the recent progresses in the field and improved quality of the introduction substantially. See below for details:

Added discussion on how flooding characteristics affect riparian carbon cycling:

“Flooding disturbance strongly influences the biotic characteristics of riparian assemblages (Anderson et al., 2020) as well as the carbon cycle. Flooding could increase soil respiration and enzymatic degradation rate (Wilson et al., 2011). It was found that the rate of CO₂ emission in riparian wetlands is higher than that in neighbouring hillslope grasslands (Anderson et al., 2020). Liu et al. (2021) demonstrated that high plant and soil respiration in riparian wetlands lead to large amounts of CO₂ emission in wet season (335-2790 mg·m⁻²·h⁻¹) than in dry season (72 - 387 mg·m⁻²·h⁻¹). (Liu et al., 2021). Also, the short-term anaerobic conditions caused by flooding may increase the production of methane because of the strengthened methanogenesis in riparian soils (Hassanzadeh et al., 2019; Hondula et al., 2021; Morse et al., 2012; Le Mer and Roger, 2001; Thorp et al., 2006).

The influence of flooding disturbance would also depend on the flooding characteristics and the properties of riparian soils. Hirota et al. (2007) found that temporal variations of the greenhouse gases fluxes were strongly manipulated by water-level fluctuations in the sandy shore and by soil temperature in the salt marsh (Hirota et al., 2007). The duration of flooding was also considered an important factor for riparian carbon dynamics and microbial community structure (Wilson et al., 2011). The spatial heterogeneity of soil properties would also affect the composition and diversity of bacterial communities in riparian zones and thus may influence the riparian carbon cycle under flooding disturbance (Wang et al., 2019b; Wilson et al., 2011).” (Line 50-64)

Added discussion on how vegetation affects riparian carbon fluxes:

“Strong seasonality for different greenhouse gas emissions has been detected in previous studies (Gaughan and Waylen, 2012; Allen et al., 2007). With flooding disturbance, riparian vegetation plays an indispensable role in sequestering carbon (Maraseni and Mitchell, 2016) and the variations in riparian vegetation communities are expected to define the ecological role of riparian zones in carbon cycle. During flooding season, flooding submergence may impede gas diffusion and decrease light intensity, leading to high mortality and limited growth of plant species (Colmer et al., 2009). This raises the possibility of elevated carbon (including methane and carbon dioxide) emissions and reduced carbon sequestration from riparian zones, shifting the role of riparian zones from a carbon sink to a carbon source. Conversely, as riparian species adapt to flooding submergence and recover from flooding, riparian zones may gradually return to the initial status or even promote CO₂ capture. Previous studies found that riparian vegetation may increase their leaf gas exchange in response to submergence stress so as to cope with oxygen limitation (Huang et al., 2017; Mommer et al., 2006; Liu et al., 2020). Besides, inundation depth increased reed density, height, leaf area index and biomass, and thus decreased the global warming potential during the growing season (Zhao

et al., 2020). Therefore, riparian zone may oscillate between carbon source and sink depending on flooding. It raises the open question of whether riparian zones quantitatively promote or hinder carbon capture overall.” (Line 66-78)

Added discussion on riparian carbon modeling works:

“However, the current research on the riparian carbon sequestration under flooding disturbance remains poorly constrained. There has been some modelling work about the riparian carbon stock, but fewer on the carbon flux. For instance, Dybala et al., (2019) modelled the change in carbon stock as a function of vegetation age, considering effects of climate and whether or not the riparian forest had been actively planted (Dybala et al., 2019). One limitation for models like Riparian Ecosystem Management Model (REMM) or other riparian models is that they require a large amount of site specific parameters, many of which are often modeled using other models as inputs (Vidon et al., 2019). In addition to climatic factors, factors such as floodplain width, flow regime, frequency of inundation, and the presence of dams, diversions, and levees also need to be considered when modelling the riparian carbon flux with the disturbance of flooding (Sutfin et al., 2016).” (Line 114-122)

Line 86, province?

Corrected accordingly.

“Our study site is in the downstream of the 164 kilometres long Lijiang River in the Pearl River Basin in northwestern Guangxi Zhuang Autonomous Province, Southwest China (25° 06' N, 110° 25'; Fig. A1).” (Line 140)

The information of sediment or soil should be added, including soil pH, total organic carbon/nitrogen content, etc.

Thanks for the suggestion. Related information on soil and sediment characteristics of the study region has been added in section 2.1. See below for details.

“Lijiang River has a typical karst landscape, with widely exposed carbonate rocks (Wang et al., 2019b). The river from Guilin to Yangshuo is the most typical karst development area. The river channel is composed of sand and pebbles, and the soil type is red loam with high sand content (Wang et al., 2019b).” (Line 140-144)

“The soils of the Lijiang River riparian zone were sand-based, with sand contents ranging from 74.99% to 88.44%; silt and clay contents are lower, accounting for approximately 10% (Wang et al., 2019b; Lu and Wang, 2015). With the decrease of inundation frequency, the sand content is found to decrease while the clay and silt content increased gradually (Wang et al., 2019b). Soil pH is around 6.99 to 7.71, and soil total nitrogen is around 0.93 to 1.40 (g·kg⁻¹) (Wang et al., 2019b). Different vegetation zones can further influence the chemical properties of soils (Lu and Wang, 2015).” (Line 148-152)

Section 2.2.1, the setup of static chamber and gas sampling are not clear for me. Do you have a base for the chamber? How to seal the chamber during the non-flood periods? The gas samples were taken every four hours, and the chamber were always closed during this time? If it is so, did the temperature inside of the chamber change a lot? How did you calibrate the flux data with temperature?

With respect to setting up of the static chamber and gas sampling, the following details have been added to section 2.2.1, in which we explained the design details of the floating and terrestrial chambers, sealing of the chambers, temperature controls and gas sampling procedures. We believe the information is able to provide a clearer view of the field methods we used for the study. See below for details.

“... On the river, floating static chambers were used (Sun et al., 2012) and were set up on shallow water and deep water. The floating static chamber was a cylindrical chamber (of radius 50cm and height 65cm) with a floating ring (about 20cm) around the bottom of the chamber to keep it floating on the water, and was thus sealed by the water. On the land during non-flooding seasons, the terrestrial static chambers (length 50 cm, width 50 cm, and height 50 cm) were used and were set up on riparian areas with vegetation and without vegetation. The terrestrial static chamber was put on a stainless-steel underside base (length 50 cm, width 50 cm and height 15 cm) instead of setting directly on the ground. The underside base increased the chamber’s size and prevented damage to the vegetation inside (Sun et al., 2013). There was a groove on the top of the underside base, and the upper portion of chamber was designed to be put into this groove. By adding water to the groove, the whole setting was sealed (Sun et al., 2012, 2013). The floating static chamber and the terrestrial static chamber both were covered by foam and reflective aluminium, which can easily reflect the heat from sunlight and thus prevent rapid temperature changes or temperature becoming too high in the chamber (Søvik and Kløve, 2007). Also, the chambers contained two exhaust fans, a thermometer and a tube inside. A syringe was used to collect gas samples from the tube at intervals of 0, 10, 20 and 30 minutes. For 24-hour monitoring, samples were taken every 4 hours (a total 6 times per day starting at 10:00 and finishing at 06:00 the next day) in one day in April, August, and October (covering pre-flooding season, flooding season, and post-flooding season) in 2014 (both riparian area and river) and 2016 (river). In other words, diel data was taken at the 0, 10min 20min and 30min of 10:00, 14:00, 18:00, 22:00, 2:00 and 6:00.” (Line 160-175)

The detailed information of gas chromatography should be introduced.

As suggested, we added the following information about the gas chromatography method, followed by a reference to the operating guide. See below for details. Also, in response to changes in other places of the manuscript, we made the Section 2.2.3 “Measurement of gas concentration and hydro-environment condition” as Section 2.2.2, and the original Section 2.2.2 “Vegetation inventory and flooding tolerant experiment” as Section 2.2.3 in the revised manuscript, so that the methodology part is more coherent.

“Gas samples were collected by a syringe from the tube of chamber and were instantly transferred to airtight glass bottles (20ml, Agilent5190-2286). All samples were analysed within three days. The CO₂ concentration was measured using gas chromatography (Agilent7890A) equipped with an electron capture detector (ECD)

and a flame ionization detector (FID) (Agilent Technologies, 2010). The measurements were conducted by Pony Testing International Group Co. Ltd (300887:CH). Standard CO₂ gases (with concentration of 0 and 10000 pm, respectively) were used for method calibration.” (Line 177-181)

The statistical method of multiple comparisons should be given.

In this analysis, the two-way ANOVA method was used to examine the effects of vegetation and time on the riparian carbon fluxes. In section 2.2.5, we explained the usage, data requirement, null assumptions and software of the method. See below for details. The corresponding results were presented in Appendix Table A1-2, 4.

“For riparian areas, two-way repeated-measurement ANOVA were employed to examine the effects of vegetation (bare soil vs. land with vegetation; between-subject factor) and time (measuring times in one day, within-subject factor) on the CO₂ flux in two sampling stages (April: pre-flooding and October: post-flooding). For aquatic habitats (fluvial area), two-way repeated-measures ANOVAs were used to examine the effects of sampling position (with vegetation vs. without vegetation or under water surface; between-subject factor) and time (measuring times in one day; within-subject factor) on CO₂ flux in sampling stages (April: pre-flooding, August: during flooding, and October: post-flooding). The *p*-values were calculated with the null hypothesis that the CO₂ flux of riparian areas or aquatic habits is not influenced by the factors mentioned. Before analyses, homogeneity of variance and normality are also examined. All data analyses were performed by the SPSS statistical software package (<https://www.ibm.com/products/spss-statistics>, version 22.0, Chicago, IL, USA). The effects were considered significant if *p*-value < 0.05.” (Line 226-235)

Equation 2, the unit of each parameter should be clarified. How did you calculate the D and p value?

In the revised manuscript, we replaced this section with a new section (section 2.2.4 Annual riparian and river CO₂ emission calculation). In this new section, instead of presenting the carbon offset model, we explained how the measured carbon fluxes were accumulated to obtain carbon fluxes for different periods (pre-flooding, flooding and post-flooding) (see response to the comments on the daily CO₂ flux, lines 189-194 and line 357-361). Equation 2 was therefore removed from the revised manuscript.

Section 2.3.2, why did you compare with the CO₂ flux of global forests?

We appreciate the reviewer’s comment on this. In revising the manuscript, we have carefully considered and re-evaluated the estimation of carbon uptake potential from global riparian zones and the comparison with global forest CO₂ flux. Considering this part was only loosely connected to the main arguments of the manuscript and that we were not able to give a comprehensive evaluation of the carbon uptake potential for global riparian areas (e.g., taking into account the systematic differences in global riparian vegetation, soil and hydrology), we removed this part and related sections in result (section 3.6) and discussion (section 4.4) in the revised manuscript. Following this change, we have substantially restructured the discussion section of the manuscript, focusing now on two main results of our study: 1) increased carbon emission from riparian

area due to flooding and 2) the role of post-disturbance survived vegetation in riparian carbon sequestration after the flooding period. We feel these changes make main arguments of our manuscript more focused and the manuscript more consistent and evidence-based. See below for changes to the discussion,

“4.1 Increased carbon emission during flooding periods of the riparian zone

Hydrological flow has been found to be an essential factor within the carbon cycle of riparian ecosystems (Zarnetske et al., 2018). Our data suggest that flooding not only affects carbon emission from the fluvial channel but also the carbon fluxes of the riparian area. With regard to carbon emission from the fluvial channel, our data show that carbon emission of water-air interface significantly increased and showed a net emission of CO₂ in both the daytime and night-time (all-day CO₂ flux: 0.291 g·m⁻² d⁻¹ in April, 2.560 g·m⁻² d⁻¹ in August). This is probably due to the increased lateral carbon flux from terrestrial areas to rivers due to flooding. Research found that when water flows through ecosystem, it would pick up dissolved organic carbon from vegetation and soils, transporting the carbon from riparian ecosystem to streams (Raymond and Saiers, 2010). A large amount of carbon could be transported to the river because of enhanced hydrological connectivity between the fluvial channel and its riparian areas during flooding periods (Zarnetske et al., 2018).

When comparing the CO₂ flux of shallow-water area (with aquatic vegetation) and deep-water area (without vegetation) (Fig. A2), it is also found that shallow-water released less carbon in pre-flooding season and captured more carbon in post-flooding season than deep-water area (pre-flooding: 0.090 g·m⁻² d⁻¹ in shallow water, 0.492 g·m⁻² d⁻¹ in deep water; post-flooding: -0.880 g·m⁻² d⁻¹ in shallow water, -0.545 g·m⁻² d⁻¹ in deep water). However, during the flooding season, both the shallow-water and deep-water areas had a carbon flux of about 2.55 g·m⁻² d⁻¹, probably because of an enhanced input of carbon from riparian vegetation and soils to the waters.

In addition to increased hydrologic connectivity between the riparian area and fluvial channel of the river, enhanced carbon emission also results from enhanced substrate availability during flooding (Hirota et al., 2007). Previous work also reported that the extensive root system of riparian species with strong taproots and well-developed fibrous roots could force the species to demand more oxygen and accelerate root respiration and CO₂ emissions from the neighbouring rhizosphere (Elias et al., 2015). In submerged areas, the CO₂ may be transported to water and then released to the atmosphere as the carbon flux of water surface. Especially, the recovery of some C₄ riparian species after periodic flooding also contributed to the higher gas transportability and abundant substrate for CO₂ emission compared to the performance of C₃ species (Still et al., 2003). In addition to riparian vegetation, inundation could also increase the decomposition of stored organic matter (Denef *et al.*, 2001, Marín-Muñiz *et al.*, 2015) and soil respiration (Anderson *et al.*, 2020, Ou *et al.*, 2019). A previous study found that after 25 days of soil moisture enhancement, the anaerobiosis stimulates CO₂ loss by 1.5 times more than the normal soil moisture environment (Huang & Hall, 2017). Flooding leads to elevated soil moisture for weeks or even months, and thus an accelerated CO₂ supply to the inundated channel.” (Line 436-465)

“4.2 Post-disturbance survived vegetation as a critical factor for riparian systems to sequester carbon

We observed that the carbon sequestration of riparian area and fluvial area as a whole was greatly enhanced after the flooding period, to the point that the overall carbon flux was negative. In consistence with our analysis, Kathilankal et al. (2008) proposed that tidal inundation caused a mean reduction of 49 % in the marsh-atmosphere carbon (CO₂) flux compared to non-flooded conditions (Kathilankal *et al.*, 2008). Our

study offers proof that the hydrological flow is a determining factor on whether the riparian ecosystem is a net carbon source or sink.

One possible reason is that the vegetation's recovery after flooding enhances its ability to sequester more CO₂ for photosynthesis. The post-flooding succession of vegetation suggests that not all riparian plants can survive submergence and to become efficient carbon sinks. Indeed, species richness decreased after flooding, which indicates a decrease of the interspecific competition, giving a chance to species that can quickly recover from submergence. The dominant species changed from *C. dactylon* to *C. aciculatus* after flooding disturbance. Although the individual biomass and number of *C. aciculatus* did not increase, existing literature suggests that the leaf maximum net photosynthesis rate may increase significantly after severe submergence in the riparian zones of Lijiang (Huang *et al.*, 2017, Jie *et al.*, 2012). For the clonal plants, its physiological integration allowing them to survive submergence and spread rapidly after de-submergence. Luo *et al.* (2014), studying *Alternanthera philoxeroides* (alligator weed) after 30 days of submergence, found that connections between submerged and non-submerged ramets enhance the performance of the submerged ramets; and the de-submerged ramets had high soluble sugar concentrations, suggesting high metabolic activities (Luo *et al.*, 2014). Wei *et al.* (2018) also found that after 30 days of submergence, stolon connection significantly increased growth, biomass allocation to roots and photosynthetic capacities of the submerged ramets, and increased growth and photosynthetic capacities of the unsubmerged ramets (Wei *et al.*, 2018). Also, flooding could promote CO₂ use efficiency and the ability of the plant to use low light (Wang *et al.*, 2019a). The enhanced photosynthetic capacity is believed to be one of the physiological strategies for species growing in critical zones with flooding disturbance. Moreover, human impacts can no longer be ignored on the riparian ecosystem (Ren *et al.*, 2019), suggesting vegetation that can recover quickly and densely is essential to allow riparian zones to be efficient carbon sinks.

Our results suggest, on an annual scale, riparian area behaves either as a net source or sink of carbon depending on the relative importance between enhanced emission during flooding and the strength of post-disturbance carbon absorbance. Assuming the carbon flux rates of flooding season and non-flooding seasons were the same as we have measured on the selected days (Section 2.2.4, Fig.1-2), we estimated that the riparian area and the fluvial area as a whole can achieve carbon neutralization ($C_{annual}=0$) only when flooding days are fewer than 15 days. Therefore, the relative ratio of flooding to non-flooding days are essential factors to determine whether the riparian area is a net source or sink on an annual scale, and future long-term, high-frequency measurements are required to monitor the carbon dynamics of the riparian zone. Also, besides the contribution of recovered vegetation, our data shows that bare soil also contributes to the carbon neutralization, but the mechanism for bare soil to capture carbon still needs further analysis.

Nowadays, the risk and the number of global flooding events are expected to rise significantly with global warming (Hirabayashi *et al.*, 2013). This means that the annual carbon cycle of riparian area and fluvial area as a whole is subject to a more variant and stronger impact from flooding. Previous research found that with a warmer climate, there would be a large increase in flood frequency in Southeast Asia, Peninsular India, eastern Africa and the northern half of the Andes (Hirabayashi *et al.*, 2013). Our research highlights that flooding disturbance would not only cause large carbon emission during the flooding season, but can also promote carbon sequestration in the post-flooding season. It is therefore necessary to consider the dynamic effect of flooding on ecosystems' carbon cycle especially under global climate change.” (Line 467-508)

Where is the daily CO₂ flux in different months?

Explanation about the assumption of calculation has been added at the beginning of Section 3.1. The rest of Section 3.1 is restructured and the sentences that may lead to misunderstanding have been removed. In accordance with this change and the removal of Section 2.3 (see response to Section 3.5), we have also added a new Section 2.2.4 “Annual riparian and river CO₂ emission calculation” to show how we use the measured CO₂ flux to evaluate the carbon balance of the whole region (see response to the comment on Lines 189-194), with assumption about different flooding seasons, flooding characteristics, riparian characteristics. See below for details.

“We assume that diel CO₂ flux follows similar patterns as measured on the selected days during the pre-flooding and post-flooding season. Based on this assumption, we compared the diel CO₂ flux of pre-flooding season and post-flooding season. In order to evaluate the effect of vegetation on riparian CO₂ flux, we directly measured the CO₂ fluxes in the riparian area with and without vegetation (bare soil) in different seasons.” (Line 266-269)

Lines 180-183, is that due to the effects of climate? Without exclusion of the climate effects, I think you can't reach the conclusion that “in post-flooding season, the terrestrial area with vegetation sequesters carbon for a longer time”.


We have checked the climate factors of Lijiang River to see whether the change is due to the effects of climate. The climate factors in April (before flooding season) and October (after flooding season) are actually very similar. Based on the latitude and longitude of the studied area, it is easy to calculate that the sunrise time, sunset time and daylength of April are around 5:40, 18:14, and 12.5 hours; of October are 6:17, 17:43, and 11.42 hours. Thus, the sunrise time is later in October, and the daylength time is even shorter than October. Therefore, we do not think the longer carbon sequestration time occurred in this study is relevant to daytime change in Lijiang riparian area.

The temperature conditions are also similar (Minimum average temperature: April: 16°C, October: 17°C, Maximum average temperature: April: 23°C, October: 26 °C). As for the precipitation, the rainfall in April (224mm) is even higher than October (85mm), which is not beneficial for plant growth and carbon sequestration. Thus, we do not think the enhanced carbon sequestration ability or the longer carbon sequestration time is caused by the season change itself in the studied area.

We suggest that vegetation is one of the most dynamic biotic factors affecting riparian carbon cycle and stronger and longer carbon accumulation during the post-flooding period could be due to a systematic change in riparian vegetation composition after flooding. As has been discussed in discussion (section 4.2), survived riparian vegetation had a higher tolerance to water submergence and high carbon uptake capacity. The carbon uptake of vegetation reflects the effects of a series of abiotic changes, which include not only the disturbance factor like flooding, but also the climatic factors.

Figure 1, the standard deviation/error of the data should be provided.

Thanks for the suggestion. Standard deviations of data were added. See right,



Lines 189-194, you should calculate accumulated CO₂ emissions rather use the average value. Otherwise, I think you can't get the conclusion that "the riparian zone acted overall as a carbon sink".

In the revised version of manuscript, we added a new Section 2.2.4 "Annual riparian and river CO₂ emission calculation" where we explained how we calculated carbon fluxes of riparian area and fluvial area as a whole using coverage as the weight. Following the suggestion, the annual riparian and fluvial CO₂ emission are

now calculated as the sum of emission in pre-flooding season, flooding season and post-flooding season, instead of averaged values of the measured fluxes. See below,

In Results:

“The riparian area is composed of vegetation area and bare soil. During the field investigation, we found the vegetation coverage in Lijiang riparian area is about 60%. Using vegetation coverage as the weight, we can get the accumulated CO₂ flux of riparian area (Section 2.2.4, equation(3)).” (Line 291-295, Results 3.1)

“In both April and October, the all-day carbon fluxes in the riparian area were negative, indicating that the riparian area acted as a carbon sink in non-flooding season (April: -0.156 g·m⁻² d⁻¹, October: -0.500 g·m⁻² d⁻¹). The carbon uptake in October, which represented the post-flooding season, was higher. Overall, we found that in the post-flooding season, the riparian vegetation can sequester CO₂ for a longer time and fix a higher amount of carbon.” (Line 299-303, Results 3.1)

“Based on the vegetation coverage and the ratio of riparian area width to river width in flooding season, we can accumulate the CO₂ flux of riparian area and the river as a whole (Section 2.2.4). The CO₂ flux of the whole region was 1.833 g·m⁻² d⁻¹ in pre-flooding season, and -0.592 g·m⁻² d⁻¹ in post-flooding season, which indicated that the whole region turned from a carbon source to a carbon sink after flooding.” (Line 329-332, Results 3.2)

In Methods:

“2.2.4 Annual riparian and river CO₂ emission calculation

We are interested in whether or under what conditions the riparian area and the fluvial area as a whole can achieve carbon neutralization (which means the net carbon emission is zero) at the annual level with flooding disturbance. We take flooding disturbance into account by dividing the whole year into pre-flooding season, flooding season, and post-flooding season. We assume that flooding events happen at an annual timescale and consider the time that flooding would happen as flooding season. The occurrence of extreme weather like rainstorms or frost is not considered here. The riparian area refers to area that would be submerged during flooding. The field investigation showed that the riparian area in the non-flooding seasons (pre-flooding season and post-flooding season) was about 25% of the river width in the flooding season, and the vegetation coverage is about 60%. Thus, the annual riparian CO₂ emission is calculated as the sum of emissions in pre-flooding season, flooding season and post-flooding season, by the following equation:

$$C_{annual} = \sum C_{i,j} = \sum W_{i,j} * D_j * a_{i,j} \quad (2)$$

Where C_{annual} is the annual expected carbon emission ($C_{annual} = 0$ means the whole region reaches carbon neutralization at the annual level), $C_{i,j}$ is the annual CO₂ emission of river or riparian area in different seasons ($i=1, 2$ refer to river and riparian area respectively, $j=1, 2, 3$ refer to pre-flooding season, flooding season, and post-flooding season respectively), $W_{i,j}$ is the width of river or riparian area in different seasons, D_j is the days of corresponding season, and $a_{i,j}$ is the all-day CO₂ flux of river or riparian area in different seasons. Specially, during flooding season, the width of riparian area ($W_{1,2}$) is 0 meter because all the riparian area is submerged. The all-day CO₂ flux of riparian area in pre- ($a_{2,1}$) or post-flooding season ($a_{2,3}$) is calculated by the following equation:

$$a = a_{veg} * p + a_{soil} * (1 - p) \quad (3)$$

Where a_{veg} is the all-day CO₂ flux of vegetation area, a_{soil} is the CO₂ flux of bare soil area, and p is the

vegetation coverage.” (Line 205-224)

Table 3, the whole plant species can be shown in the supplementary data.

Thanks for the suggestion. Appendix Table 3 has been added to show the whole plant species. See below,

Appendix Table A3. The whole plant species in pre-flooding season (surveyed in April) and post-flooding season (surveyed in October).

Pre-flooding season	Post-flooding season
<i>Aster tataricus</i>	<i>Alternanthera philoxeroides</i>
<i>Astragalus sinicus</i>	<i>Aster tataricus</i>
<i>Athyrium sinense</i>	<i>Astragalus sinicus</i>
<i>Cardamine hirsuta</i>	<i>Cardamine hirsuta</i>
<i>Carex duriuscula</i> subsp. <i>stenophylloides</i>	<i>Carex polycephala</i> var. <i>simplex</i>
<i>Carex polycephala</i> var. <i>simplex</i>	<i>Chrysopogon aciculatus</i>
<i>Chrysopogon aciculatus</i>	<i>Cynodon dactylon</i>
<i>Cichorium endivia</i>	<i>Oxalis corymbosa</i>
<i>Conyza canadensis</i>	<i>Polygonum hydropiper</i>
<i>Cynodon dactylon</i>	<i>Polygonum lapathifolium</i>
<i>Digitaria ciliaris</i>	<i>Stellaria media</i>
<i>Hemarthria altissima</i>	
<i>Lindernia antipoda</i>	
<i>Oxalis corymbosa</i>	
<i>Poa annua</i>	
<i>Polygonum hydropiper</i>	
<i>Polygonum lapathifolium</i>	
<i>Polygonum muricatum</i>	
<i>Potentilla chinensis</i>	
<i>Salvia plebeia</i>	
<i>Stellaria media</i>	
<i>Urena lobata</i>	
<i>Viola philippica</i>	
<i>Vitex negundo</i>	

Section 3.5, how did you verify the model data?

In the revised manuscript, in response to the main restructuring of the manuscript's discussion (see response to section 2.3.2 above), we have decided to present this as a subsection of section 4.2 (Post-disturbance survived vegetation as a critical factor for riparian systems to sequester carbon). Therefore, instead of presenting this as a separate section and a model that needs verification, this part intends to give a general idea of how flooding and post-disturbance vegetation recovery interplay to control net carbon balance of the riparian zone on an annual scale. We argue that this change makes the manuscript more coherent and evidence-based. See related changes in discussion below.

“Our results suggest, on an annual scale, riparian area behaves either as a net source or sink of carbon depending on the relative importance between enhanced emission during flooding and the strength of post-disturbance carbon absorbance. Assuming the carbon flux rates of flooding season and non-flooding seasons were the same as we have measured on the selected days (Section 2.2.4, Fig.1-2), we estimated that the riparian area and the fluvial area as a whole can achieve carbon neutralization ($C_{annual}=0$) only when flooding days are fewer than 15 days. Therefore, the relative ratio of flooding to non-flooding days are essential factors to determine whether the riparian area is a net source or sink on an annual scale, and future long-term, high-frequency measurements are required to monitor the carbon dynamics of the riparian zone. Also, besides the contribution of recovered vegetation, our data shows that bare soil also contributes to the carbon neutralization, but the mechanism for bare soil to capture carbon still needs further analysis.” (Line 491-500)

The unit should be kept the same through the manuscript. For example, the CO₂ flux is expressed as g·m⁻² d⁻¹, g·m⁻² year⁻¹, and mg·m⁻² h⁻¹.

We have checked throughout the manuscript, and we ensure that all CO₂ fluxes were expressed in the unit of mg·m⁻² h⁻¹ when describing the CO₂ flux of different measuring time within a diel cycle, and in the unit of g·m⁻² d⁻¹ when describing the integrated all-day CO₂ flux.

Figure 5, the unit should be Gg·m⁻² year⁻¹?

In response to the main re-structuring of the manuscript's discussion (see response to comments on section 2.3.2), Figure 5 has been removed from the revised manuscript.

Section 3.6, how did you upscale the site CO₂ flux to a global/regional CO₂ flux? Did you consider the effects of temperature, vegetation, seasonal changes, variations of soils etc.? If not, it is hard to believe the data.

We appreciate this comment. We have carefully reevaluated this part. Considering this part was only loosely connected to main arguments of the manuscript and that we were not able to give a comprehensive evaluation

of the carbon uptake potential for global riparian areas (e.g., taking into account the systematic differences in global riparian vegetation, soil and hydrology), this part and related sections in result (section 3.6) and discussion (section 4.4) were removed in the revised manuscript. See also response to section 2.3.2 above.

Lines 357-361, again, you couldn't conclude a net emission by using the average flux data.

As suggested in the comment to Line 189-194, we used the accumulated rather than averaged carbon flux to evaluate the carbon balance of the whole area. We added a new section 2.2.4 in the revised manuscript to explain the calculation. See also response to Line 189-194.

Section 4.3, I didn't find any data of microbiology. So please delete this part.

In response to the comment on section 2.3.2, we have substantially restructured the discussion and this part was removed from the main discussion sections in the revised manuscript. See also response to comment to section 2.3.2 above.

Reference mentioned in the response:

Allen, D. E., Dalal, R. C., Rennenberg, H., Meyer, R. L., Reeves, S., and Schmidt, S.: Spatial and temporal variation of nitrous oxide and methane flux between subtropical mangrove sediments and the atmosphere, *Soil Biology and Biochemistry*, 39, 622–631, <https://doi.org/10.1016/j.soilbio.2006.09.013>, 2007.

Anderson, N. J., Heathcote, A. J., Engstrom, D. R., and Globocarb data contributors: Anthropogenic alteration of nutrient supply increases the global freshwater carbon sink, *Sci. Adv.*, 6, eaaw2145, <https://doi.org/10.1126/sciadv.aaw2145>, 2020.

Colmer, T. D., Voesenek, L. a. C. J., Colmer, T. D., and Voesenek, L. a. C. J.: Flooding tolerance: suites of plant traits in variable environments, *Functional Plant Biol.*, 36, 665–681, <https://doi.org/10.1071/FP09144>, 2009.

Dyballa, K. E., Matzek, V., Gardali, T., and Seavy, N. E.: Carbon sequestration in riparian forests: A global synthesis and meta-analysis, *Global Change Biology*, 25, 57–67, <https://doi.org/10.1111/gcb.14475>, 2019.

Elias, E., Steele, C., Havstad, K., Steenwerth, K., Chambers, J., Deswood, H., Kerr, A., Albert, R., Schwartz, M., Stine, P., and Steele, R.: Southwest Regional Climate Hub and California Subsidiary Hub assessment of climate change vulnerability and adaptation and mitigation strategies, 2015.

Gaughan, A. E. and Waylen, P. R.: Spatial and temporal precipitation variability in the Okavango–Kwando–Zambezi catchment, southern Africa, *Journal of Arid Environments*, 82, 19–30, <https://doi.org/10.1016/j.jaridenv.2012.02.007>, 2012.

Hassanzadeh, Vidon, Gold, Pradhanang, and Lowder: RZ-TRADEOFF: A New Model to Estimate Riparian Water and Air Quality Functions, *Water*, 11, 769, <https://doi.org/10.3390/w11040769>, 2019.

Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H., and Kanae, S.: Global flood risk under climate change, *Nature Clim Change*, 3, 816–821, <https://doi.org/10.1038/nclimate1911>, 2013.

Hirota, M., Senga, Y., Seike, Y., Nohara, S., and Kunii, H.: Fluxes of carbon dioxide, methane and nitrous oxide in two contrastive fringing zones of coastal lagoon, Lake Nakaumi, Japan, *Chemosphere*, 68, 597–603,

<https://doi.org/10.1016/j.chemosphere.2007.01.002>, 2007.

Hondula, K. L., Jones, C. N., and Palmer, M. A.: Effects of seasonal inundation on methane fluxes from forested freshwater wetlands, *Environ. Res. Lett.*, 16, 084016, <https://doi.org/10.1088/1748-9326/ac1193>, 2021.

Huang, D., Wang, D., Ren, Y., Qin, Y., and Wu, L.: Responses of leaf traits to submergence stress and analysis of the economic spectrum of plant species in an aquatic-terrestrial ecotone, the Li River, *Acta Ecol. Sin.*, 37, <http://dx.doi.org/10.5846/stxb201508281789>, 2017.

Le Mer, J. and Roger, P.: Production, oxidation, emission and consumption of methane by soils: A review, *European Journal of Soil Biology*, 37, 25–50, [https://doi.org/10.1016/S1164-5563\(01\)01067-6](https://doi.org/10.1016/S1164-5563(01)01067-6), 2001.

Liu, R., Liang, S., Long, W., and Jiang, Y.: Variations in Leaf Functional Traits Across Ecological Scales in Riparian Plant Communities of the Lijiang River, Guilin, Southwest China, *trcs*, 11, <https://doi.org/10.1177/1940082918804680>, 2020.

Liu, X., Lu, X., Yu, R., Sun, H., Xue, H., Qi, Z., Cao, Z., Zhang, Z., and Liu, T.: Greenhouse gases emissions from riparian wetlands: an example from the Inner Mongolia grassland region in China, *Biogeosciences*, 18, 4855–4872, <https://doi.org/10.5194/bg-18-4855-2021>, 2021.

Luo, F.-L., Chen, Y., Huang, L., Wang, A., Zhang, M.-X., and Yu, F.-H.: Shifting effects of physiological integration on performance of a clonal plant during submergence and de-submergence, *Annals of Botany*, 113, 1265–1274, <https://doi.org/10.1093/aob/mcu057>, 2014.

Maraseni, T. N. and Mitchell, C.: An assessment of carbon sequestration potential of riparian zone of Condamine Catchment, Queensland, Australia, *Land Use Policy*, 54, 139–146, <https://doi.org/10.1016/j.landusepol.2016.02.013>, 2016.

Mommer, L., Lenssen, J. P. M., Huber, H., Visser, E. J. W., and de Kroon, H.: Ecophysiological Determinants of Plant Performance under Flooding: A Comparative Study of Seven Plant Families, *Journal of Ecology*, 94, 1117–1129, 2006.

Morse, J. L., Ardón, M., and Bernhardt, E. S.: Greenhouse gas fluxes in southeastern U.S. coastal plain wetlands under contrasting land uses, *Ecological Applications*, 22, 264–280, <https://doi.org/10.1890/11-0527.1>, 2012.

Raymond, P. A. and Saiers, J. E.: Event controlled DOC export from forested watersheds, *Biogeochemistry*, 100, 197–209, <https://doi.org/10.1007/s10533-010-9416-7>, 2010.

Ren, Y., Wang, D., and Li, X.: Impacts of Human Disturbances on Riparian Herbaceous Communities in a Chinese Karst River, *Nature Environment and Pollution Technology*, 18, 1107–1118, 2019.

Søvik, A. K. and Kløve, B.: Emission of N₂O and CH₄ from a constructed wetland in southeastern Norway, *Science of The Total Environment*, 380, 28–37, <https://doi.org/10.1016/j.scitotenv.2006.10.007>, 2007.

Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: Global distribution of C₃ and C₄ vegetation: Carbon cycle implications: C₄ PLANTS AND CARBON CYCLE, *Global Biogeochem. Cycles*, 17, 6-1-6–14, <https://doi.org/10.1029/2001GB001807>, 2003.

Sun, Q., Shi, K., Damerell, P., Whitham, C., Yu, G., and Zou, C.: Carbon dioxide and methane fluxes: Seasonal dynamics from inland riparian ecosystems, northeast China, *Science of The Total Environment*, 465, 48–55, <https://doi.org/10.1016/j.scitotenv.2013.01.036>, 2013.

Sun, Q.-Q., Whitham, C., Shi, K., Yu, G.-H., and Sun, X.-W.: Nitrous oxide emissions from a waterbody in the Nenjiang basin, China, *Hydrology Research*, 43, 862–869, <https://doi.org/10.2166/nh.2012.060>, 2012.

Sutfin, N. A., Wohl, E. E., and Dwire, K. A.: Banking carbon: a review of organic carbon storage and physical factors influencing retention in floodplains and riparian ecosystems: BANKING CARBON, *Earth Surf. Process. Landforms*, 41, 38–60, <https://doi.org/10.1002/esp.3857>, 2016.

Thorp, J. H., Thoms, M. C., and DeLong, M. D.: The riverine ecosystem synthesis: biocomplexity in river networks across space and time, *River Research and Applications*, 22, 123–147, <https://doi.org/10.1002/rra.901>, 2006.

Vidon, P. G., Welsh, M. K., and Hassanzadeh, Y. T.: Twenty Years of Riparian Zone Research (1997–2017): Where to Next?, *Journal of Environmental Quality*, 48, 248–260, <https://doi.org/10.2134/jeq2018.01.0009>, 2019.

Wang, J., Wang, D., Ren, Y., and Wang, B.: Coupling relationships between soil microbes and soil nutrients under different hydrologic conditions in the riparian zone of the Lijiang River, *Acta Ecol. Sin.*, 39, <https://doi.org/10.5846/stxb201803260595>, 2019a.

Wang, J., Wang, D., and Wang, B.: Soil Bacterial Diversity and its Determinants in the Riparian Zone of the Lijiang River, China, *Current Science*, 117, 1324, <https://doi.org/10.18520/cs/v117/i8/1324-1332>, 2019b.

Wei, G.-W., Shu, Q., Luo, F.-L., Chen, Y.-H., Dong, B.-C., Mo, L.-C., Huang, W.-J., and Yu, F.-H.: Separating effects of clonal integration on plant growth during submergence and de-submergence, *Flora*, 246–247, 118–125, <https://doi.org/10.1016/j.flora.2018.08.004>, 2018.

Wilson, J. S., Baldwin, D. S., Rees, G. N., and Wilson, B. P.: The effects of short-term inundation on carbon dynamics, microbial community structure and microbial activity in floodplain soil, *River Research and Applications*, 27, 213–225, <https://doi.org/10.1002/rra.1352>, 2011.

Zarnetske, J. P., Bouda, M., Abbott, B. W., Saiers, J., and Raymond, P. A.: Generality of Hydrologic Transport Limitation of Watershed Organic Carbon Flux Across Ecoregions of the United States, *Geophys. Res. Lett.*, 45, <https://doi.org/10.1029/2018GL080005>, 2018.

Zhang, T., Huang, X., Yang, Y., Li, Y., and Dahlgren, R. A.: Spatial and temporal variability in nitrous oxide and methane emissions in urban riparian zones of the Pearl River Delta, *Environ Sci Pollut Res*, 23, 1552–1564, <https://doi.org/10.1007/s11356-015-5401-y>, 2016.

Zhao, M., Han, G., Wu, H., Song, W., Chu, X., Li, J., Qu, W., Li, X., Wei, S., Eller, F., and Jiang, C.: Inundation depth affects ecosystem CO₂ and CH₄ exchange by changing plant productivity in a freshwater wetland in the Yellow River Estuary, *Plant Soil*, 454, 87–102, <https://doi.org/10.1007/s11104-020-04612-2>, 2020.

Zheng, X., Wang, M., Wang, Y., Shen, R., Li, J., J., H., M., K., Li, L., and Jin, J.: Comparison of manual and automatic methods for measurement of methane emission from rice paddy fields, *Adv. Atmos. Sci.*, 15, 569–579, <https://doi.org/10.1007/s00376-998-0033-5>, 1998.