

# Mapping soil organic carbon fractions for Australia, their stocks and uncertainty

Mercedes Román Dobarco<sup>1</sup>, Alexandre M.J-C. Wadoux<sup>1</sup>, Brendan Malone<sup>2</sup>, Budiman Minasny<sup>1</sup>, Alex B. McBratney<sup>1</sup>, Ross Searle<sup>3</sup>

5 <sup>1</sup>Sydney Institute of Agriculture & School of Life and Environmental Sciences, The University of Sydney, 1 Central Avenue, Eveleigh, 2015, NSW, Australia

<sup>2</sup>CSIRO Agriculture and Food, Black Mountain, ACT, Australia

<sup>3</sup>CSIRO Agriculture and Food, 306 Carmody Road, St Lucia, QLD, Australia.

\*Correspondance to: Mercedes Román Dobarco (mercedes.romandobarco@sydney.edu.au)

10 **Abstract.** Soil organic carbon (SOC) is the largest terrestrial carbon pool. SOC is composed of a continuum set of compounds with different chemical composition, origin and susceptibilities to decomposition, that are commonly separated into pools characterised by different responses to anthropogenic and environmental disturbance. Here we map the contribution of three SOC fractions to the total SOC content of Australia's soils. The three SOC fractions: mineral-associated organic carbon (MAOC), particulate organic carbon (POC) and pyrogenic organic carbon (PyOC), represent SOC composition with distinct

15 turnover rates, chemistry, and pathway formation. Data for MAOC, POC, and PyOC were obtained with near- and mid-infrared spectral models calibrated with measured SOC fractions. We transformed the data using an isometric log-ratio transformation (ilr) to account for the closed compositional nature of SOC fractions. The resulting, back-transformed ilr components were mapped across Australia. SOC fraction stocks for the 0-30 cm were derived with maps of total organic carbon concentration, bulk density, coarse fragments and soil thickness. Mapping was done by quantile regression forest fitted with the ilr

20 transformed data and a large set of environmental variables as predictors. The resulting maps along with the quantified uncertainty show the unique spatial pattern of SOC fractions in Australia. MAOC dominated the total SOC with an average of 59%  $\pm$  17%, whereas 28%  $\pm$  17% was PyOC and 13%  $\pm$  11% was POC. The allocation of TOC into the MAOC fractions increased with depth. SOC vulnerability (i.e., POC/[MAOC + PyOC]) was greater in areas with Mediterranean and temperate climate. TOC and the distribution among fractions were the most influential variables on SOC fraction uncertainty. Further,

25 the diversity of climatic and pedological conditions suggests that different mechanisms will control SOC stabilisation and dynamics across the continent, as shown by the model covariates importance metric. We estimated the total SOC stocks (0-30 cm) to be 13 Pg MAOC, 2 Pg POC and 5 Pg PyOC, which is consistent with previous estimates. The maps of SOC fractions and their stocks can be used for modelling SOC dynamics and forecasting changes in SOC stocks as response to land use change, management, and climate change.

30

Keywords: Soil organic carbon, SOC fractions, particulate organic carbon, mineral-associated organic carbon, pyrogenic organic carbon, digital soil mapping.

## 1 Introduction

35 Soils are the main organic carbon pool in terrestrial ecosystems, storing around two thirds of the total C. Global soil organic carbon (SOC) stock is estimated to be around 1500 PgC for the first metre of soil (Jobbagy and Jackson, 2000), with other estimates ranging from 504 to 3000 PgC (Scharlemann et al., 2014). Changes in SOC storage and dynamics can alter the ecosystem C balance and determine whether soils become C sources or sinks from local to global scale (Friedlingstein et al., 2020). SOC is strongly linked to most soil properties and functions (e.g., nutrient and water storage and cycling, habitat provisioning and biodiversity) (Van Leeuwen et al., 2019), and hence is used as a general indicator of soil quality/capacity 40 (Schoenholtz et al., 2000; Bunemann et al., 2018) and soil health/condition.

Soil organic matter consists of a continuum of compounds with different chemical compositions, origin (aboveground litter, dead roots, rhizodeposition, microbial-derived), degree of microbial processing and decomposition, and turnover times (Lehmann and Kleber, 2015). SOC is the main component of soil organic matter and varies in spatial and temporal dynamics. SOC is protected against microbial decomposition by several stabilisation mechanisms which have been generally grouped 45 into: 1) selective preservation due to biochemical recalcitrance, 2) chemical stabilisation via interactions between organic compounds and mineral surfaces or metal cations, and 3) physical protection by the inaccessibility of decomposers to the organic matter (Sollins et al., 1996; Von Lutzow et al., 2006; Rowley et al., 2018). Spatial inaccessibility and interactions between the mineral surfaces of silt and clay-sized particles and organic compounds are considered the major mechanisms for mid and long-term SOC stability (Von Lutzow et al., 2006), whereas an increasing body of literature questions the relevance 50 of biochemical recalcitrance for long-term persistence of SOC (Schmidt et al., 2011). However, the hierarchy between stabilization mechanisms varies with the pedoclimatic context, land use and SOC fraction.

A myriad of physical, chemical, or combined fractionation methods have been designed for separating SOC into operational pools characterized by specific stabilization mechanisms, chemical composition, and distinct turnover rates, and yet it is difficult to isolate fractions that correspond to functional SOC pools (Lutzow et al., 2007). Some fractionation schemes were 55 adapted to quantify conceptual SOC pools from established C dynamics models, e.g., the Rothamsted C model (RothC, Jenkinson and Rayner (1977)) from measured SOC fractions (Poeplau et al., 2013; Zimmermann et al., 2007). However, there can be some discrepancies between the predicted SOC pools when the model is initialized with modelled SOC pools from equilibrium runs or from measured SOC fractions (Poeplau et al., 2013). Other biogeochemical models have been conceptualized and calibrated with functional and measurable SOC fractions to overcome these differences (Robertson et al., 60 2019), but often require the determination of many SOC fractions. A comparison of several fractionation schemes (Poeplau et al., 2018) suggests that size separation into silt + clay (<53  $\mu\text{m}$ ) (i.e., mineral-associated organic carbon (MAOC)), and sand-sized particles (>53  $\mu\text{m}$ ) (i.e., particulate organic carbon (POC)), may suffice to differentiate pools with distinct turnover rates,

chemistry, and pathway formation (Lavallee et al., 2020). MAOC is predominantly composed of low molecular weight molecules of microbial origin (e.g., microbial metabolites, necromass) (Miltner et al., 2012; Kallenbach et al., 2016; Liang et al., 2019), leachates from plant litter, and rhizodeposition (Villarino et al., 2021), which are protected through sorption to mineral surfaces or occluded inside microaggregates (Lavallee et al., 2020). POC is mainly composed of partially decomposed plant litter and roots and fungal-derived compounds (Baisden et al., 2002; Gregorich et al., 2006; Geng et al., 2019; Villarino et al., 2021), that can be found free or inside macroaggregates (Rabbi et al., 2014). MAOC has a lower C:N ratio, a higher proportion of microbial-derived and proteinaceous compounds (Kleber et al., 2007; Knicker, 2011) and a longer mean residence time (decades to centuries) than POC (mean residence time of years to decades) (Baisden et al., 2002; Gregorich et al., 2006; Von Lützow et al., 2007; Heckman et al., 2022). Separating SOC into POC and MAOC has been proposed as a simple and effective way to conceptualize and model SOC dynamics (Lavallee et al., 2020), and has been applied to predict the vulnerability of SOC to future climate scenarios (Lugato et al., 2022).

In Australia, the long history of burning suggests pyrogenic organic carbon (PyOC) is an additional important component of SOC (Lehmann et al., 2008). PyOC refers to charred residues derived from incomplete combustion of organic matter (also known as charcoal or black carbon) (Lutfalla et al., 2017). PyOC is comprised by a continuum of organic compounds thermally altered by fire, and its chemical composition and pool size depends on the technique used for its determination (Zimmerman and Mitra, 2017). PyOC is considered a relatively stable SOC fraction with a turnover time that ranges from decades to centuries (Singh et al., 2012), protected from decomposition by the biochemical recalcitrance of condensed aromatic C (Lavallee et al., 2019). However, turnover rates previously assessed from centuries to multi-millenia may have overestimated its persistence in soil (Singh et al., 2012; Lutfalla et al., 2017). PyOC can be found in both POC and MAOC fractions (Lavallee et al., 2019). Beyond the biochemical recalcitrance, which would be the main stabilization mechanism of PyOC in the POC fraction (e.g., free PyOC non-associated to clay and silt sized mineral particles), PyOC can also interact with mineral phases and be protected inside microaggregates or through adsorption (Burgeon et al., 2021). Zimmerman and Mitra (2017) suggest that PyOC (e.g., naturally occurring or added as biochar) may promote the stabilization of non-PyOC by enhancing the creation of microaggregates and sorption onto existing organo-mineral complexes. PyOC represents on average 14-26% of total SOC and can reach up to 60-80% of SOC (Lehmann et al., 2008; Reisser et al., 2016). Globally, PyOC stocks are not controlled by fire intensity and return interval, but are explained by soil properties and climate (Reisser et al., 2016). However, in systems with local records of fire history PyOC content increased in sites with high-frequency fires (Reisser et al., 2016). In Australia, fire is an important driver of ecological processes, and bush-fire frequency has increased in recent decades (Dutta et al., 2016). Hence, PyOC can represent a large proportion of total SOC in some Australian regions (Lehmann et al., 2008).

Mapping and quantifying the stocks of SOC fractions can be used for modelling SOC dynamics and forecasting changes in SOC stocks as a response to land-use change, management, and climate change (Lugato et al., 2022; Xu et al., 2011; Wiesmeier et al., 2016). High-resolution maps of SOC fractions can inform agricultural management at the farm scale and be incorporated in the design of climate change and soil security policies at the state and national level (Dangal et al., 2022). The objective of this study was to map the contribution of SOC fractions (MAOC, POC, and PyOC) to the total SOC in the top 30 cm of the

soil and update the Soil and Landscape Grid of Australia (SLGA) (Grundy et al., 2015) products for SOC fraction stocks (Viscarra Rossel et al., 2019). These digital soil maps will be part of the v2.SLGA for Australia, following the principles of transparency, reproducibility, and updatability as new data become available (Malone and Searle, 2021).

## 100 **2 Materials and Methods**

The methodology implemented in this study consists of two main steps: 1) prediction of SOC fractions across the soil spectral libraries available for Australia with different spectral models calibrated with measured SOC fraction data from the Australian Soil Carbon Research Program (SCaRP) (Baldock et al., 2013a), and 2) mapping of the proportions of SOC fractions after applying the isometric log-ratio transformation (ilr) to account for the compositional nature of the data (i.e., the proportion of SOC fractions sum to 100%). Previous studies modelled SOC fractions stocks or concentrations (Sanderman et al., 2021; Viscarra Rossel et al., 2019), but we found some difficulties in implementing such an approach (see Sect. 2.5) and hence mapped proportions of SOC. Finally, we calculated SOC fraction stocks for the 0-30 cm topsoil using data from SLGA maps of total organic carbon (TOC) concentration (Wadoux et al., 2022), bulk density (Viscarra Rossel et al., 2015), coarse fragments (this study) and soil thickness (Malone and Searle, 2020).

### 110 **2.1 Study area**

The study area covers the continent of Australia, including Tasmania and near-shore small islands. In Australia there are six major climatic regions following the Köppen classification. In the north, there are hot humid summers in equatorial, tropical and subtropical regions. Summers are hot and dry, with mild to cold winters in grasslands and desert regions in the interior. Temperate areas in the south coastal band have cold winters and warm summers, that are mild at higher elevations and latitudes. Vast areas of the continent are very dry, with precipitation increasing towards the coast and elevation in the mountains of the eastern uplands. Australia is characterised by flat and low relief vast areas where the tectonic stability and lack of glaciation have preserved a deeply weathered mantle that dates from the Tertiary (McKenzie et al., 2004). The distribution of soils in many regions depends on the stripping of this weathered mantle, while in other areas the dominant drivers of landforms and soils are water, fluvial, and aeolian erosion and depositional processes (McKenzie et al., 2004). Younger soils are found mainly in the east, along the Great Divide (McKenzie et al., 2004). The dominant soil orders according to the Australian Soil Classification (Isbell et al., 1997) are Kandosol (30 % of the area), Tenosol (20 %), Vertosol (14 %), Rudosol (8 %), Calcarosol (8 %) and Chromosol (7 %) (Searle, 2021). In 2015-2016 most of the area was dedicated to primary production, with 48 % of the continent used for livestock grazing (42 % on native vegetation and 6 % modified pastures), less than 5 % dedicated to dryland and irrigated cropping. About 9.5% of Australia is allocated to nature conservation and 18 % are protected managed resources (ABARES, 2022).

## 2.2 SCaRP dataset – sampling design and SOC fractionation scheme

The soil sampling design and SOC fractionation protocol of SCaRP are described in detail in Sanderman et al. (2011) and Baldock et al. (2013c). The objective of SCaRP was to characterize SOC stocks and composition of agricultural topsoils (0-30 cm) and their response to agricultural practices. Plots of 25 m<sup>2</sup> were laid in 4526 sites representative of the different combinations of agricultural management and soil types across Australia between 2008 and 2012. At each plot, soil samples were collected randomly at 10 nodes of a 5 m x 5 m grid with a soil corer ( $\geq 4$  cm diameter) by depth interval (0-10, 10-20, 20-30 cm) and composited by layer. In addition, at least three measurements of bulk density by depth interval were taken at each plot, although the methods varied between states to accommodate the soil type. A subset of 312 samples representative of the range of TOC content (1.2 - 90.9 mg C g<sup>-1</sup> soil) found among the whole dataset (N = 24,495 samples) and different soil types and biomes were subject to fractionation following the protocol described in detail in Baldock et al. (2013c). The size and chemical fractionation scheme divided SOC into MAOC, POC and PyOC. MAOC and PyOC were originally defined as humic organic carbon and resistant organic carbon (ROC) in Baldock et al. (2013c), and although the ROC fraction may not correspond completely with PyOC we changed the terminology to match the recent literature. A 10-g aliquot of air-dried soil  $\leq 2$  mm was dispersed with 5 g L<sup>-1</sup> sodium hexametaphosphate and separated into coarse ( $>50$   $\mu\text{m}$ ) and fine ( $<50$   $\mu\text{m}$ ) fractions with wet sieving using an automated sieve shaker system (Baldock et al., 2013c). The TOC concentrations of the coarse and fine fractions were analysed with high-temperature oxidative combustion after the removal of inorganic carbon with 5-6 % H<sub>2</sub>SO<sub>3</sub> if carbonates were present (method 6B3a, Rayment and Lyons (2011)). Solid-state <sup>13</sup>C nuclear magnetic resonance (<sup>13</sup>C NMR) spectroscopy analyses were conducted on both the coarse ( $>50$   $\mu\text{m}$ ) and fine ( $<50$   $\mu\text{m}$ ) fractions. <sup>13</sup>C NMR is a semi-quantitative method that is commonly used to measure the proportion of aromatic C compounds in soil and organic matter samples. The proportion of poly-aryl C and aryl C that could be defined as lignin was determined and used as an estimate of PyOC. We note that whereas the chemical signature of the poly-aryl C is consistent with, and likely dominated by charcoal and charred plant residues, it may also indicate the presence of compounds non-pyrogenic origin (Baldock et al., 2013c). POC and MAOC contents (mg C-fraction g<sup>-1</sup> soil) were calculated by subtracting the proportion of PyOC in each fraction with the following equations (Baldock et al., 2013c):

$$\text{POC} = (2000 - 50 \mu\text{m OC})(1 - f\text{PyOC}_{2000})\text{MF}_{2000} \quad (1)$$

$$\text{MAOC} = (\leq 50 \mu\text{m OC})(1 - f\text{PyOC}_{50})\text{MF}_{50} \quad (2)$$

$$\text{PyOC} = (2000 - 50 \mu\text{m OC})(f\text{PyOC}_{2000})\text{MF}_{2000} + (\leq 50 \mu\text{m OC})(f\text{PyOC}_{50})\text{MF}_{50} \quad (3)$$


where 2000-50  $\mu\text{m}$  OC is the measured TOC content in the coarse fraction (mg C g<sup>-1</sup> 2000-50  $\mu\text{m}$  fraction),  $f\text{PyOC}_{2000}$  is the proportion of TOC attributed to poly-aryl C in the coarse fraction,  $\text{MF}_{2000}$  is the proportion of soil mass found in the coarse fraction (g 2000-50  $\mu\text{m}$  fraction / g  $\leq 2$ mm soil),  $\leq 50$   $\mu\text{m}$  OC is the measured TOC content in the fine fraction (mg C g<sup>-1</sup>  $\leq 50$   $\mu\text{m}$  fraction),  $f\text{PyOC}_{50}$  is the proportion of TOC attributed to poly-aryl C in the fine fraction, and  $\text{MF}_{50}$  is the proportion of soil mass found in the fine fraction (g  $\leq 50$   $\mu\text{m}$  fraction / g  $\leq 2$ mm soil).

### 2.3 Spectral datasets and harmonization

160 Four spectral soil datasets were combined in this study: SCaRP (4526 sites and 24495 soil specimens; Baldock et al. 2013b),  
the Australian Soil Archive mid-infrared (MIR) spectral library (703 sites and 4300 soil specimens; AusSpecMIR) (Hicks et  
al., 2015), an additional MIR library of specimens from the Australian Soil Archive that were not included in the Hicks et al.  
(2015) work (346 sites and 429 soil specimens; AusSpecMIR2) and the Australian Soil Archive visible and near-infrared (vis-  
NIR) spectral library (4533 sites and 25174 soil specimens; AusSpecNIR). The majority but not all of these vis-NIR data were  
165 described in (Viscarra Rossel and Hicks, 2015).

The AusSpecNIR library also contained a subset (309) of the soil specimens described in Viscarra Rossel and Hicks (2015)  
which contain measures of soil carbon fractions from selected SCaRP soil specimens and associated vis-NIR spectra. The  
spectra were collected using the same instrument that was used to derive the Australian Soil Archive vis-NIR spectral library.  
The AusSpecNIR library contained further data from the Terrestrial Ecosystem Research Network (TERN) Ecosystem  
170 Surveillance plots (Sparrow et al., 2020; Malone et al., 2020), increasing the representation of non-agricultural soils  
(rangelands, forests, woodlands) of the combined datasets. (Figure 1b). With these TERN data, vegetation and soils are  
sampled in more than 800 permanent plots laid with stratified random sampling across bioregions (i.e., zones with similar  
landform, vegetation, and climate, analogous to ecoregion) (Sparrow et al., 2020). Soil samples are taken from three depth  
intervals (0-10, 10-20, 20-30 cm) in nine locations within a 100 m x 100 m plot (Sparrow et al., 2020), for a total of 15,157  
175 soil specimens (with associated vis-NIR spectra) at 5,711 georeferenced sampling points.

Excepting those with soil carbon fraction measurements, multiple soil samples (approximately 700) were represented both in  
AusSpecMIR and AusSpecNIR, and duplicates were removed from the AusSpecNIR for the subsequent modelling processes.



180 **Figure 1: a) Location of SCaRP soil samples subject to fractionation (n=312) by Baldock et al. (2013a). b) Soil spectral datasets combined for mapping soil organic carbon fractions. TERN samples are part of the AusSpecNIR library.**

The ScaRP spectral library was produced with a Nicolet 6700 FTIR spectrometer (Thermo Fisher Scientific Inc., Waltham, MA, USA) equipped with a KBr beam-splitter, a DTGS detector and an AutoDiff Automated diffuse reflectance accessory (Pike Technologies, Madison, WI, USA). Spectra were acquired over the range 8000–400  $\text{cm}^{-1}$  with a resolution of 8  $\text{cm}^{-1}$ . The spectrum of each soil sample was the average of 60 scans. The AusSpecMIR spectral library was created with a Bruker FT-IR Vertex70 spectrometer. This instrument is fitted with a mercury cadmium telluride detector that is liquid  $\text{N}_2$  cooled to improve signal-to-noise ratio and has a spectral range of 7500–600  $\text{cm}^{-1}$  at 4  $\text{cm}^{-1}$  resolution. AusSpecMIR2 were measured with a Bruker FT-IR Invenio spectrometer with a similar liquid  $\text{N}_2$  cooling system and spectral range of 7500–600  $\text{cm}^{-1}$  at 4  $\text{cm}^{-1}$  resolution. Soil samples from AusSpecMIR and AusSpecMIR2 were scanned in quadruplicate. Diffuse reflectance spectra for the AusSpecNIR library were collected with a Labspec® vis–NIR spectrometer (PANalytical Inc., Boulder, CO, USA) for the range 350–2500 nm at 1 nm resolution. Each spectrum was the average of 30 scans and up to 4 spectra were collected for each soil sample.

SCaRP spectra were baseline-corrected and mean-centered prior subsequent analyses but were not subject to additional pre-processing. Pre-processing of the AusSpecNIR spectra consisted of spectral trimming (453–2500 nm), a Savitsky-Golay smoothing filter, conversion of reflectance to absorbance, and standard normal variate transformation. AusSpecMIR and AusSpecMIR2 were both pre-processed with the following steps: 1) spectral resolution and range harmonisation. All spectra were resampled using a smoothing spline function to a common 2  $\text{cm}^{-1}$  resolution. The spectral range was set to 6500  $\text{cm}^{-1}$  to 598  $\text{cm}^{-1}$ , 2) Savitsky-Golay smoothing filter with 22  $\text{cm}^{-1}$  local neighbourhood, 3) conversion from reflectance to absorbance units, and 4) standard normal variate transformation.

## 2.4 Spectral models and predictions of SOC fractions

SOC fractions contents ( $\text{mg C g}^{-1}$  soil) of ScaRP had previously been estimated with partial least squares regression (PLSR) models by Baldock et al. (2013b). Baldock et al. (2013b) reported cross-validation results (squared-root transformed SOC fraction contents) with  $R^2$  values of 0.84, 0.88 and 0.85, and RMSE of 0.43, 0.40 and 0.32 for POC, MAOC and PyOC respectively.

SOC fractions content for AusSpecMIR and AusSpecMIR2 were estimated with a new set of spectral predictive functions (Malone and Wadoux 2021). Spectral harmonisations using piecewise direct standardisation (Bouveresse and Massart, 1996; Ge et al., 2011) was used to align spectra collected for ScaRP and AusSpecMIR2 to AusSpecMIR spectrometer. Approximately 200 soil samples from the ScaRP dataset were measured with the Bruker FT-IR Vertex70 spectrometer, following the estimate of parameters to harmonise the ScaRP library with AusSpecMIR. Similarly, the AusSpecMIR2 was harmonised with AusSpecMIR with PDS using a standard of 300 soil samples measured with both instruments.

Soil spectral model functions were developed with the 312 ScaRP samples with data on SOC fractions concentration ( $\text{mg C SOC fraction g}^{-1}$  soil), total organic carbon (TOC) concentration ( $\text{mg C g}^{-1}$  soil), and used PDS harmonised spectra. The contents of SOC fractions were converted into percentages of TOC (summing up to 100 %) and modelled as compositional data. Hence, the ilr transformation was applied to the proportions of SOC fractions. The ilr generates a D-1 dimensional

215 Euclidean vector, where  $D$  is the number of variables in the original variables (Aitchison, 1986). PLSR models with bootstrapping were used to predict the  $\ln$ -transformed SOC fractions. Out-of-bag validation statistics were calculated from the average of 50 model realisations on the back-transformed data. The average RMSE for POC, MAOC and PyOC were 6.2, 7.7, and 6.4 %, respectively. The Lin's concordance correlation coefficient ( $\rho_c$ ) values were 0.86, 0.76 and 0.70, and  $R^2$  of 0.74, 0.61, and 0.52. When the percentages were back-transformed into SOC fraction contents (mg C-SOC fraction  $\text{g}^{-1}$  soil), the  
220 RMSE were 2.1, 2.7, and 1.8 mg C  $\text{g}^{-1}$  soil,  $\rho_c$  of 0.92, 0.95, and 0.90, and  $R^2$  of 0.85, 0.88, and 0.77 for POC, MAOC and PyOC respectively.


SOC fraction contents for the AusSpecNIR dataset were predicted with PLSR models calibrated with data on TOC concentration and SOC fractions contents from the 309 SCaRP samples. Before modelling, the spectra were trimmed to the 453-2500 nm range, processed with a Savitsky-Golay smoothing filter and transformed from reflectance to absorbance. We  
225 also applied the standard normal variate transformation and then wavelet coefficients were derived. Similarly, as for the MIR datasets, spectral predictive functions were developed for  $\ln$ -transformed SOC fraction compositional data, and validation statistics were calculated for the back-transformed data. The validation statistics for POC, MAOC and PyOC models were respectively RMSE of 5.0, 6.4, and 5.8 %,  $\rho_c$  of 0.92, 0.84, and 0.76, and  $R^2$  of 0.83, 0.71, and 0.61 when validated as proportions of TOC. When the validation statistics were calculated for SOC fraction contents, the RMSE were 3.2, 4.0 and  
230 2.3 mg C  $\text{g}^{-1}$  soil and  $\rho_c$  of 0.81, 0.87, 0.80, and  $R^2$  of 0.65, 0.75 and 0.64 for POC, MAOC and PyOC.

Fitted spectral models based on either MIR and vis-NIR data were then extended across to all data cases of the respective soil spectral libraries.

## 2.5 SOC fractions data processing and depth standardisation

235 The georeferenced SOC fraction data from all libraries was projected to WGS 84 (EPSG:4326) and collated. The data were filtered and processed to harmonize units, duplicates and potentially wrong data entries (e.g., missing upper or lower horizon depths) were excluded. SOC predictions of the quadruplicated scans per soil sample were averaged, as well as multiple soil samples per horizon by sampling location. SOC fraction concentrations (mg C-SOC fraction  $\text{g}^{-1}$  soil) were transformed to proportion of total SOC (% SOC). SOC potential vulnerability ( $V_p$ ) was defined as the ratio of POC to the sum of MAOC and  
240 PyOC (Baldock et al., 2018), i.e., the proportion of the less protected SOC fraction (POC) to the SOC fractions with stability by physico-chemical protection and recalcitrance, and longer turnover rates. SOC fractions and  $V_p$  data were standardized to the first three depth intervals of the GlobalSoilMap specifications (0-5 cm, 5-15 cm, 15-30 cm) (Arrouays et al., 2014) with equal-area quadratic splines (Bishop et al., 1999) (Figure 2). For sampling locations with just one sampled horizon, this was converted to the GlobalSoilMap depth intervals (assigned to a single or several depth intervals) via aggregation with the *slab*  
245 function of the *aqp* R package (Beaudette et al., 2022). We calculated the mean and standard deviation of SOC fractions and  $V_p$  by GlobalSoilMap depth interval and by biome (Olson et al., 2001). Extreme values of  $V_p$  were constrained to a maximum of 5 before calculating the summary statistics.





250 **Figure 2: Location of the spectral predictions standardized for the depths 0-5 cm, 5-15 cm, and 15-30 cm used as input data for digital soil mapping. Contribution of SOC fractions to total organic carbon (% TOC): mineral-associated SOC (MAOC), particulate organic carbon (POC), pyrogenic organic carbon (PyOC) and soil organic carbon vulnerability (Vp).**

Modelling SOC fraction concentrations directly was the preferred option tested in preliminary work. We compared the sum of the predicted SOC fraction contents with measured TOC. The Pearson's  $r$  correlation coefficient was 0.56, but the sum of SOC fractions showed some extreme values (Figure S1). The mismatch between the sum of SOC fractions and measured TOC is most likely derived from the prediction error of the spectral models, and in minor extent from the TOC recoveries of the laboratory SOC fractionation data. We adjusted the SOC fraction contents with measured TOC data extracted from the Soil Data Federator (Searle et al., 2021) and the Biome of Australian Soil Environments (BASE) contextual data (Bissett et al., 2016). However, the resulting dataset had a reduced spatial coverage and the maps exhibited unrealistic patterns. Hence, we preferred to use all available data and map the proportion of SOC fractions and SOC vulnerability.

The values of SOC proportions were modelled considering their compositional nature (i.e., multivariate data with positive values that sum up to a constant, in the case of SOC fractions 100%). The set of all compositional observations ( $S^D$ ) is a simplex sample space, a subset of the real space  $\mathbb{R}^D$ . The ilr-transformation (Egozcue et al., 2003) transposes  $S^D$  into multidimensional real space ( $\mathbb{R}^{D-1}$ ), without the collinearity problems associated with other transformations, such as the centered log-ratio transformation (clr). Ilr is based on the choice of an orthonormal basis on the hyperplane in  $\mathbb{R}^D$  formed by the clr transformation. The ilr-transformation equation is defined:

$$z_i = ilr(x_i) = \sqrt{\frac{i}{i+1}} \ln \left( \frac{\sqrt{\prod_{j=1}^i x_j}}{x_{i+1}} \right), i = 1, 2, \dots, D - 1 \quad (4)$$

And the inverse equation is defined as follows (Filzmoser and Hron, 2008):

$$x_i = \frac{\exp(y_i)}{\sum_{i=1}^D \exp(y_i)} \quad i = 1, 2, \dots, D \quad (5)$$

$$y_i = \sum_{j=i}^D \frac{z_j}{\sqrt{j(j+1)}} - \sqrt{\frac{i-1}{i}} z_{i=1} \quad \text{with } z_0 = z_D = 0 \text{ for } i = 1, 2, \dots, D \quad (6)$$

We applied the ilr-transformation with the *ilr* function of the *compositions* R package (Van Den Boogaart et al., 2022), generating two variables hereafter referred to as *ilr1* and *ilr2*. The predictions (*ilr1* and *ilr2*) were back-transformed into MAOC, POC, PyOC with the *ilrInv* function.


## 2.6 Coarse fragments

Data on the abundance of coarse fragments (particles > 2 mm) and gravimetric content (% weight) were extracted using the TERN Soil Data Federator (<https://esoil.io/TERNLandscapes/Public/Pages/SoilDataFederator/SoilDataFederator.html>) managed by CSIRO (Searle et al., 2021). The Soil Data Federator is a web API that compiles soil data from different institutions and government agencies throughout Australia. The abundance (% volume) is assessed visually in the field as part of the soil profile description using standards described in the Australian Soil and Land Survey field Handbook (National Committee on Soil and Terrain, 2009). The abundance of rock fragments per soil horizon on the cut surface of the soil profile was grouped into six categories: very few (0-2 %), few (2-10 %), common (2-20 %), many (20-50 %), abundant (50-90 %) and very

abundant (> 90%). The gravimetric content (% mass) is measured in the laboratory as percent mass of coarse fragments (particles > 2 mm) from the whole soil. Here, we take the profile surface abundance of coarse fragments as a proxy for volumetric coarse fragments ( $CF_{Vol}$ ). The data was cleaned and processed to exclude duplicates and wrong data entries (e.g., missing values). The observations of  $CF_{Vol}$  (%) were converted into GlobalSoilMap depth intervals with the *slab* function of the *aqp* R package (Beaudette et al., 2022), assigning the most probable class to each depth interval. The gravimetric coarse fragments were also standardized to the GlobalSoilMap depth intervals with equal-area quadratic splines (Bishop et al., 1999). Observations of gravimetric coarse fragment content ( $CF_{Mass}$ ) were transformed into volumetric with Eq.(4):

$$CF_{Vol} (\%) = \frac{Vol_{CF}}{Vol_{WhSoil}} \frac{Mass_{CF} / \rho_{CF}}{Mass_{WhSoil} / \rho_{WhSoil}} = \frac{CF_{Mass} \times \rho_{WhSoil}}{\rho_{CF}} \quad (7)$$

where  $\rho_{WhSoil}$  is the bulk density prediction for bulk soil from SLGA (Viscarra Rossel et al., 2015),  $\rho_{CF}$  is assumed to be 2.65 g cm<sup>-3</sup> (Hurlbut and Klein (1977) in McKenzie et al. (2002)) and  $CF_{Vol}$  is the volumetric coarse fragment content (continuous), which was assigned to the corresponding class. This resulted in  $CF_{Vol}$  observations for 110,308 locations (Figure 3).



295

**Figure 3: Calibration data on coarse fragments (% volume) classes for the 0-5, 5-15, 15-30 cm depth intervals.**

## 2.7 Environmental covariates

We selected 61 spatially-exhaustive covariates (90 m grid cell size) for Australia made available by TERN, describing soil-forming factors and soil properties (Table 2), to calibrate *scorpan* models (McBratney et al., 2003). A *scorpan* model establishes quantitative relationships between soil properties or classes and other soil properties (*s*), climate (*c*), organisms (*o*), relief (*r*), parent material (*p*), age (*a*), and spatial position (*n*), i.e.,  $soil = f(s, c, o, r, p, a, n)$  (McBratney et al., 2003). We used the soil properties of clay and sand content for each GSM layer due to the relevant role of soil texture in SOC stabilization. We selected 15 climatic covariates since climate is a relevant driver of SOC storage from the global to subregional scale (Wiesmeier et al., 2019), influencing both the SOC decomposition and the C input into the soil. The organisms factor was represented by 22

305

covariates related to net primary productivity (e.g., Normalized Difference Vegetation Index (NDVI), enhanced vegetation index (EVI)) and vegetation type which influence the C allocation patterns, quantity and pathways of C input (e.g., aboveground vs. belowground). Long-term average NDVI calculated with Landsat 5 surface reflectance data was processed with Google Earth Engine. Eight relief covariates were surrogates for elevation, water flow, erosion processes, and sediment transport and accumulation, derived from the 3-second Shuttle Radar Topographic Mission (SRTM) (Farr et al., 2007). Gamma radiometrics, total magnetic intensity and weathering intensity index (Wilford, 2012a) were used as proxies for parent material and age. Gamma radiometrics inform on regolith's mineralogy and surface geochemistry. The concentrations and ratios of three radioelements (potassium (K), thorium (Th), uranium (U)) are indicators of the lithology and degree of weathering (Wilford and Minty, 2007). The weathering intensity index map (Wilford, 2012) was generated from field-based data on the degree of bedrock weathering and multivariate analysis using gamma radiometrics and relief covariates as predictors. The covariates were void-filled by a combination of plane fitting and inverse distance weighting (< 0.5% of pixels in the study area). The plane fitting method was used to compute the average value of the neighbouring pixels, and otherwise an inverse distance weighting algorithm with default parameters was applied when the error in the plane fitting was large. This is the default implementation from ArcGIS software 10.5.

320

[Table 2 is at the end of the manuscript]

## 2.8 Modelling SOC fractions, SOC vulnerability and coarse fragments

Quantile regression forests (Meinshausen, 2006) is a generalization of the popular machine-learning algorithm random forests (Breiman, 2001). Random forests is based on an ensemble of regression trees. Each decision tree is fitted on a bootstrap sample of the original data. Further randomness is incorporated in each individual regression tree by selecting a subset of variables in each node for which the split is made. Whereas random forests report the mean from the observations allocated in each final node, quantile regression forests keeps all values (Meinshausen, 2006), thus estimates of conditional quantiles can be made (Meinshausen, 2006). In digital soil mapping (DSM), quantile regression forests was applied previously by Vaysse and Lagacherie (2017).

We fitted quantile regression forest models for  $ilr_1$ ,  $ilr_2$ , and  $V_p$  by GSM depth interval with the following settings:  $n_{tree} = 500$  (number of trees), default values of  $n_{odesize} = 5$  (minimum number of observations in terminal nodes) and  $m_{try} = 7$  (number of predictor variables subset as candidates to make the split at each node). The models were calibrated with the *ranger* package (Wright and Ziegler, 2017) for the R environment. We fitted probability random forest models for coarse fragments (Malley et al., 2012), setting  $n_{odesize} = 10$ . In probability random forest models, each tree predicts the class probability for each sample, and these are averaged for the forest probability estimate. The models were evaluated with 10-fold cross-validation. Variable importance was calculated with permutation (Breiman, 2001) on models fitted with 5000 trees and all observations. After the regression trees are constructed, the values of a variable of interest are randomly permuted and the

error for the out-of-bag data is estimated. The variable importance is given by the percent increase in error compared with the out-of-bag predictions leaving all variables intact.

- 340 We checked the spatial structure of the regression model residuals with cross- and auto- variograms. The spatial cross-correlation was somewhat important at a short range (approximately 20 km) for the 15-30 cm depth interval, but with a high nugget-to-sill ratio in the fitted variogram models. For the 0-15 cm and 5-15 cm depth, only the residuals of ilr2 had some spatial structure. Overall, we considered the spatial structure of the residuals negligible for the effects of mapping at the continental scale, and hence modelled the spatial distribution of SOC fractions with quantile regression models only.
- 345 For mapping, the expected mean values of the SOC fractions, quantile regression forest models for ilr1 and ilr2 were fitted with all available observations, predicted at 90 m resolution and back-transformed into MAOC, POC and PyOC. Similarly, the mean and prediction interval for Vp were predicted with a quantile regression forest model fitted with all observations, setting the 95<sup>th</sup> and 5<sup>th</sup> percentiles as prediction interval limits at a 90% confidence level. The prediction intervals for the SOC fractions were calculated from the full conditional probability distributions of ilr1 and ilr2 inferred from the quantile regression
- 350 forest models. In the model cross-validation, for each observation, a set of 500 values was generated with simple random sampling with replacement from both ilr1 and ilr2 models. Each of the 500 pairs was back-transformed into proportions of MAOC, POC and PyOC (% of SOC). The lower and upper prediction interval limits were calculated as the 5<sup>th</sup> and 95<sup>th</sup> percentiles from the empirical distribution of the 500 values of the SOC fractions. For mapping, we used 100 simulations instead of 500 to reduce the computational time. We also mapped the probability of each CF<sub>vol</sub> class at 90 m resolution.

## 355 2.9 Validation statistics

Model performance was assessed with a random 10-fold cross-validation for the ilr-variables, back-transformed SOC fractions predictions and Vp. The validation statistics included the root mean square error (RMSE), the bias or mean error (ME), the coefficient of determination ( $R^2$ ), and Lin's concordance correlation coefficient ( $\rho_c$ ) (Lin, 1989). For variable  $z$  at a location  $s_i$ , the validation statistics are calculated as:

$$360 \quad RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (z(s_i) - \hat{z}(s_i))^2} \quad (8)$$

$$ME = \frac{1}{n} \sum_{i=1}^n z(s_i) - \hat{z}(s_i) \quad (9)$$

$$R^2 = 1 - \frac{\sum_{i=1}^n (z(s_i) - \hat{z}(s_i))^2}{\sum_{i=1}^n (z(s_i) - \bar{z})^2} \quad (10)$$

$$\rho_c = \frac{2\rho\sigma_z\sigma_{\hat{z}}}{\sigma_z^2 + \sigma_{\hat{z}}^2 + (\bar{z} - \bar{\hat{z}})^2} \quad (11)$$

- where  $z(s_i)$  and  $\hat{z}(s_i)$  are observed and predicted values of  $z$  at the location  $s_i$  ( $i = 1, \dots, n$ ),  $\bar{z}$  and  $\bar{\hat{z}}$  are the means of the
- 365 observed and predicted values, respectively,  $\sigma_z^2$  and  $\sigma_{\hat{z}}^2$  are their respective variances, and  $\rho$  is the correlation between predicted and observed values. The concordance evaluates both the accuracy and the precision of the prediction, it can range between  $-1$  and  $1$ , and a value closer to  $1$  indicates a better agreement between predictions and observations.

We assessed the validity of the uncertainty estimates with the prediction interval coverage probability (PICP) (Shrestha and Solomatine, 2006). The PICP is calculated as:

$$370 \quad PICP = \frac{\text{count}(LPL_i < z(s_i) < UPL_i)}{n} \times 100, \quad (12)$$

where  $n$  is the number of observations, and the numerator the counts that an observation  $z(s_i)$  fits within its prediction interval limits. If the prediction uncertainty was correctly estimated, for example a 90% confidence level should have a PICP value close to 90% (approximately 90% of the observed values fall within the 90% prediction interval).

The  $CF_{vol}$  models were validated with 10-fold cross-validation, assigning the class with the highest probability to each observation prediction. Using the predicted and observed class values, we computed an error matrix. The error matrix is a two-way contingency table composed of the observed and predicted class, for all points within the validation dataset. From the error matrix, we calculated kappa indices: overall accuracy, user's accuracy, producer's accuracy.

The overall accuracy is the fraction of locations correctly classified. It is calculated as:

$$p = \sum_{i=1}^U N_{UU} / N, \quad (13)$$

380 where  $U$  is the total number of classes. The user's accuracy represents the fraction of the class  $u$  that is correctly classified (i.e. mapped class  $u$  in the validation dataset is also observed as class  $u$ ):

$$p_u = \frac{N_{uu}}{N_{u+}} \quad (14)$$

Producer's accuracy is similar to the user's accuracy but is calculated on the columns marginal or the error matrix. It is the fraction of observations  $u$  for which the prediction is also class  $u$ . It is obtained as follow:

$$385 \quad r_u = N_{uu} / N_{+u} \quad (15)$$

For more information on kappa statistics for evaluating map accuracy we refer to Congalton (1991) and Brus et al. (2011).

## 2.10 Mapping SOC fraction densities and stocks

The expected SOC fraction density for each GSM depth interval  $i$  was calculated with the following equation, using the map of TOC concentration from v1.2.SLGA (Wadoux et al., 2022), bulk density for the whole soil (Viscarra Rossel et al., 2015), and volumetric coarse fragments:

$$SOC_{fraction\ i\ density} (Mg\ C\ ha^{-1}\ cm^{-1}) =$$

$$SOC_{fraction\ i} \left( \frac{mg\ C_{SOC\ fraction\ i}}{mg\ C} \right) \times TOC \left( \frac{mg\ C}{g\ soil < 2\ mm} \right) \times Bulk\ density_{whole} \left( \frac{g\ soil}{cm^3} \right) \times (1 - Coarse\ Fragments_{vol} (\% / 100)) \left( \frac{soil < 2\ mm}{g\ soil} \right) \times correction\ units \left( \frac{10^8\ cm^2}{ha} \frac{Mg}{10^9\ mg} \right) \quad (16)$$

$$Coarse\ Fragments_{vol} = \sum_{u=1}^6 CF\ probability_u \times CF\ mid_u \quad (17)$$

395 where  $CF\ probability_u$  is the predicted probability and  $CF\ mid_u$  the midpoint of the  $CF_{vol}$  class  $u$ . The SOC stock density may be overestimated, especially in soils with high content of rock fragments, due to using the bulk density of whole soil and not of fine soil (Poeplau et al., 2017). We then calculated the SOC fraction stocks for the 0-30 cm depth interval using the median predictions of soil thickness (Malone and Searle, 2020) as a constraint in areas with shallow soils (< 30 cm). We

explored differences among SOC fraction stocks ( $Mg\ C\ ha^{-1}$ ) by biome and land use (natural or agriculture) by taking a regular  
 400 sample of 500,000 pixels across Australia.

The uncertainty of the SOC fraction density for each GSM depth interval was estimated with simulations. We generated a  
 sample of 500 values from the conditional probability distributions of  $ilr1$  and  $ilr2$  and back-transformed those into proportions  
 of MAOC, POC and PyOC (% of SOC). Similarly, we generated a sample of 500 TOC values with the quantile regression  
 model. To account for the uncertainty of the coarse fragment volume, we generated a sample of 500 values where: 1) the  
 405 number of samples within each  $CF_{vol}$  class was proportional to the predicted class probability, and 2) assuming a continuous  
 uniform distribution within each class. The sample for bulk density was generated assuming a normal distribution, where the  
 standard deviation was calculated from the prediction interval limits,  $sd = UPL - LPL / 2 \times z$ , with z-score = 1.64 for a 90%  
 prediction interval. We calculated the SOC fraction density for each simulation and calculated the mean and upper and lower  
 prediction interval limits with the 95<sup>th</sup> and 5<sup>th</sup> percentiles.

410 The uncertainty of SOC fraction stocks for 0-30 cm was calculated in a similar way for a subset of 98,400 pixels, incorporating  
 the uncertainty of soil thickness (Malone and Searle, 2020). A sample of 500 values for soil thickness was generated assuming  
 a normal distribution, where the standard deviation was calculated from the prediction interval limits (10<sup>th</sup> and 90<sup>th</sup> percentiles),  
 $sd = UPL - LPL / 2 \times z$ , with z-score = 1.28 for a 80% prediction interval. For each sample, we calculated the corresponding  
 thickness for each GlobalSoilMap layer ( $j = 1, 2, 3$ ), and calculated the SOC fraction stock:

415  $SOC_{fraction\ i\ stock} (Mg\ C\ ha^{-1}) =$

$$\sum_{j=1}^3 SOC_{fraction\ i} \left( \frac{mg\ C_{SOC\ fraction\ i}}{mg\ C} \right) \times TOC \left( \frac{mg\ C}{g\ soil < 2\ mm} \right) \times Bulk\ density_{whole} \left( \frac{g\ soil}{cm^3} \right) \times (1 -$$

$$Coarse\ Fragments_{vol} (\%/100)) \left( \frac{soil < 2\ mm}{g\ soil} \right) \times thickness_j (cm) \times correction\ units \left( \frac{10^8\ cm^2}{ha} \frac{Mg}{10^9\ mg} \right) \quad (18)$$

The 95<sup>th</sup> and 5<sup>th</sup> percentiles of the 500 simulations were calculated to estimate the upper and lower prediction interval limits  
 of the SOC fraction stocks. The mean of the 500 simulations were compared with predictions of SOC fraction stocks for 0-30  
 420 cm at those locations by Viscarra Rossel et al. (2019).


We performed a variance-based, global sensitivity analysis on every pixel to identify which variables account most for the  
 uncertainty of SOC fractions' density and stocks and how they vary spatially. We calculated the first-order and total effects  
 Sobol's sensitivity indices (Saltelli et al., 2008). The first-order sensitivity index  $S_i$  represents the main effect contribution of  
 each input factor  $X_i$  (SOC fraction %, TOC,  $CF_{vol}$ , and bulk density) on the variance of the output of model Y (SOC fraction  
 425 density). It is calculated as the quotient between the variance of the conditional expectation of the output with respect to an  
 input factor, and the unconditional variance of the output (Saltelli et al., 2008).  $S_i$  ranges between 0 and 1, and a high value  
 indicates that factor  $X_i$  is an important contributor to the variance of the output (if we knew the true value of  $X_i$  we would  
 significantly reduce the variance of Y). The total effect index  $S_{Ti}$  includes nonadditive features of the model, by accounting  
 for the first-order effect of  $X_i$  and its interactions with other factors on the variance of Y. When  $S_{Ti}$  is close to 0, the factor is  
 430 non-influential on the model output variance. A comprehensive explanation of the variance-based methods for global

sensitivity analysis can be found in Saltelli et al. (2008). We calculated the indices with the method by Martinez (2011), with the function `sobolmartinez` of the sensitivity package in R (Iooss et al., 2021).

### 3 Results

#### 3.1 Variation of the proportions of SOC fractions with depth and biome

- 435 The boxplots of the SOC fraction data predicted with spectral models showed that most of the TOC was found in the MAOC fraction with a mean ( $\pm$ standard deviation) of  $59\% \pm 16\%$ , whereas  $28\% \pm 18\%$  of TOC was PyOC and  $13\% \pm 11\%$  was POC. The allocation of TOC into the MAOC fractions increased with depth (Figure 4), with a mean ( $\pm$ standard deviation) of  $56\% \pm 17\%$ ,  $59\% \pm 17\%$  and  $62\% \pm 18\%$  at 0-5 cm, 5-15 cm and 15-30 cm respectively. Conversely, the proportion of SOC stored as POC decreased with depth, from  $15\% \pm 11\%$  and  $13\% \pm 11\%$  in the top 0-5 cm and 5-15 cm to  $11\% \pm 11\%$  at 15-30 cm.
- 440 The percentage of SOC in the PyOC fraction remained relatively constant with depth, with values around  $29\% \pm 17\%$  (Figure 4). Hence, in average, carbon vulnerability decreased slightly with depth, with  $V_p = 0.2 \pm 0.2$  at 0-5 cm and 5-15 cm, and  $V_p = 0.1 \pm 0.3$  at 15-30 cm.



- 445 **Figure 4: Distribution of total organic carbon (TOC) among three soil organic carbon (SOC) fractions in the spectral dataset. Mineral-associated SOC (MAOC), particulate organic carbon (POC), and pyrogenic organic carbon (PyOC).**

- The distribution of TOC among fractions showed similar patterns across biomes (MAOC  $\gg$  PyOC  $>$  POC), except for montane grasslands and shrublands where 67% of TOC was stored as PyOC while only 27% was stored as MAOC (Table 3 and Figure S2). However, there were only 160 observations located in this biome. The proportion of TOC as POC was greater in Mediterranean forests and scrub, temperate forests, and tropical and sub-tropical forests. In Mediterranean systems, the proportion of MAOC was around 10% less than across the other biomes, (excluding montane grasslands), whereas the smallest
- 450



proportion of PyOC was found in tropical and subtropical forests. Hence, the  $V_p$  was highest in the Mediterranean biome followed by temperate forests, and tropical and subtropical forests, decreasing with depth across all biomes (Table 3).

455

**Table 3: Summary statistics, mean ( $\pm$ standard deviation) of SOC fractions and SOC vulnerability ( $V_p$ ) by biome. Spectral dataset used as input for digital soil mapping. N: number of observations. Mineral-associated SOC (MAOC), particulate organic carbon (POC), and pyrogenic organic carbon (PyOC).**

Biome	Depth	N	MAOC (% SOC)	POC (% SOC)	PyOC (% SOC)	$V_p$
Montane Grasslands and Shrublands	0-5 cm	160	26 $\pm$ 9	7 $\pm$ 7	67 $\pm$ 14	0.1 $\pm$ 0.1
	5-15 cm	160	27 $\pm$ 10	6 $\pm$ 5	67 $\pm$ 15	0.1 $\pm$ 0.1
	15-30 cm	152	28 $\pm$ 13	5 $\pm$ 4	68 $\pm$ 16	0.1 $\pm$ 0.1
Temperate Broadleaf & Mixed Forests	0-5 cm	2865	54 $\pm$ 10	21 $\pm$ 8	25 $\pm$ 8	0.3 $\pm$ 0.2
	5-15 cm	2871	58 $\pm$ 10	16 $\pm$ 7	26 $\pm$ 9	0.2 $\pm$ 0.2
	15-30 cm	2722	64 $\pm$ 11	10 $\pm$ 7	26 $\pm$ 10	0.1 $\pm$ 0.2
Temperate Grasslands, Savannas & Shrublands	0-5 cm	750	58 $\pm$ 18	11 $\pm$ 7	31 $\pm$ 21	0.1 $\pm$ 0.1
	5-15 cm	762	60 $\pm$ 19	9 $\pm$ 6	30 $\pm$ 20	0.1 $\pm$ 0.1
	15-30 cm	703	62 $\pm$ 20	8 $\pm$ 7	31 $\pm$ 21	0.1 $\pm$ 0.1
Mediterranean Forests, Woodlands & Scrub	0-5 cm	4671	49 $\pm$ 17	20 $\pm$ 14	31 $\pm$ 20	0.3 $\pm$ 0.3
	5-15 cm	4695	52 $\pm$ 18	18 $\pm$ 14	30 $\pm$ 20	0.3 $\pm$ 0.3
	15-30 cm	4229	55 $\pm$ 21	16 $\pm$ 15	30 $\pm$ 21	0.3 $\pm$ 0.5
Deserts & Xeric Shrublands	0-5 cm	2427	65 $\pm$ 16	8 $\pm$ 7	26 $\pm$ 18	0.1 $\pm$ 0.1
	5-15 cm	2421	66 $\pm$ 17	7 $\pm$ 7	26 $\pm$ 18	0.1 $\pm$ 0.1
	15-30 cm	2187	67 $\pm$ 17	7 $\pm$ 7	26 $\pm$ 18	0.1 $\pm$ 0.2
Tropical & Subtropical Grasslands, Savannas & Shrublands	0-5 cm	3283	61 $\pm$ 14	10 $\pm$ 7	28 $\pm$ 15	0.1 $\pm$ 0.1
	5-15 cm	3236	64 $\pm$ 14	9 $\pm$ 6	28 $\pm$ 15	0.1 $\pm$ 0.1
	15-30 cm	2851	67 $\pm$ 15	7 $\pm$ 6	25 $\pm$ 15	0.1 $\pm$ 0.1
Tropical & Subtropical Moist Broadleaf Forests	0-5 cm	242	62 $\pm$ 19	18 $\pm$ 11	20 $\pm$ 20	0.3 $\pm$ 0.4
	5-15 cm	243	65 $\pm$ 19	17 $\pm$ 10	18 $\pm$ 20	0.2 $\pm$ 0.4
	15-30 cm	234	70 $\pm$ 18	15 $\pm$ 10	15 $\pm$ 18	0.2 $\pm$ 0.4

### 460 3.2 Cross-validation statistics for SOC fractions

The cross-validation statistics indicated that both ilr-fractions models had similar RMSE and bias, although the  $R^2$  and  $\rho_c$  were better for ilr2 than ilr1. Model performance indices decreased with depth but were overall good, with  $R^2 \geq 0.54$  for ilr1 and  $R^2 \geq 0.68$  for ilr2. Similarly, the concordance coefficients indicated good agreement between predictions and observations, with  $\rho_c \geq 0.70$  for ilr1 and  $\rho_c \geq 0.81$  for ilr2. The uncertainty was somewhat underestimated (Table 4) but close to a PICP of 90%.

465 The cross-validation of the back-transformed SOC fractions indicated that the concordance coefficient and  $R^2$  followed the trend  $PyOC > MAOC > POC$  (Table 4). Contrarily, the RMSE was greatest for MAOC with values between 8 % and 10 %, and were somewhat smaller for PyOC and POC, between 8 % and 6 %. SOC fractions predictions had a small bias, with an average underprediction of MAOC and overprediction of POC and PyOC. The accuracy plots (Supplementary material Figure S3) indicate that the uncertainty was overestimated for MAOC and PyOC, and underestimated for POC, for all probability

470 intervals, although the uncertainty was adequately estimated for the 90% prediction interval (Table 4). Cross-validation  
 statistics for SOC vulnerability were worse than for the SOC fractions, with  $R^2$  ranging between 0.39 to 0.56 and concordance  
 coefficient between 0.58 and 0.72, which is possibly due to some extreme values in the calibration dataset. A PICP around  
 86% indicates that the uncertainty was somewhat underestimated.

475

**Table 4: Cross-validation statistics for ilr1, ilr2, mineral-associated SOC (MAOC), particulate organic carbon (POC), pyrogenic  
 organic carbon (PyOC) and SOC vulnerability (Vp) by depth interval. Mean error (ME), root mean squared error (RMSE),  
 coefficient of determination ( $R^2$ ), Lin's concordance correlation coefficient ( $\rho_c$ ) and prediction interval coverage probability (PICP).**

Ilr-transformed SOC fractions	Depth	ME	RMSE	$R^2$	$\rho_c$	PICP (%)
ilr1	0-5 cm	-0.01	0.45	0.60	0.75	85
	5-15 cm	-0.01	0.42	0.63	0.77	86
	15-30 cm	-0.01	0.56	0.54	0.70	86
ilr2	0-5 cm	0.01	0.46	0.73	0.84	86
	5-15 cm	0.01	0.47	0.74	0.85	87
	15-30 cm	0.00	0.58	0.68	0.81	86
Vp	0-5 cm	0	0.21	0.39	0.58	86
	5-15 cm	0	0.17	0.56	0.72	86
	15-30 cm	0	0.21	0.56	0.72	86
Soil property	Depth	ME (%)	RMSE (%)	$R^2$	$\rho_c$	PICP (%)
MAOC	0-5 cm	1.2	8.4	0.74	0.85	95
	5-15 cm	1.3	8.8	0.73	0.85	95
	15-30 cm	2.0	10.4	0.68	0.82	95
POC	0-5 cm	-0.9	6.6	0.67	0.80	85
	5-15 cm	-0.8	6.2	0.65	0.79	86
	15-30 cm	-1.2	7.1	0.59	0.74	85
PyOC	0-5 cm	-0.3	7.7	0.80	0.89	94
	5-15 cm	-0.5	7.9	0.79	0.89	94
	15-30 cm	-0.8	8.8	0.76	0.87	94

### 480 3.3 Cross-validation statistics for coarse fragment classes


The overall accuracy of predicting coarse fragments classes was 67% for the 0-5 cm, 66% for the 5-15 cm, and 63% at the 15-  
 30 cm depth interval. The Kappa statistics were 0.39, 0.38, and 0.37, respectively, which indicate some agreement. The  
 producer's accuracy was around 90 % for the 'very few' class across the three depths, but the omission error was significant  
 for the remaining classes, especially for 'common' and 'very abundant' with values around 15 %. The user's accuracy was  
 485 smaller than 50% for the classes 'few', 'common', and 'many' coarse fragments but improved somewhat for 'very abundant',  
 'abundant', and 'few' (Table 5). The confusion matrices are provided in the Supplementary material (Table S1-S3).

**Table 5: Cross-validation statistics for classification of coarse fragments (% volume).**


	Depth	Very few (< 2 %)	Few (2-10 %)	Common (10-20 %)	Many (20-50 %)	Abundant (50-90 %)	Very abundant (> 90%)
Producer's accuracy (%)	0-5	93	39	17	35	22	15
	5-15	92	40	16	34	21	12
	15-30	92	38	13	31	39	8
User's accuracy (%)	0-5	76	47	42	44	58	56
	5-15	74	47	43	43	53	63
	15-30	71	46	42	43	53	52

490 **3.4 Variable importance for predicting proportions of SOC fractions and coarse fragments**

The environmental covariates important for both ilr1 and ilr2 models were parent material and climate, whereas relief and organisms were secondary factors (Figure 5). The rank of the 10 most important variables indicates that parent material, soil properties and relief were more relevant for ilr1, especially gravity, gamma radiometrics K, clay and elevation (DEM), whereas for ilr2 the importance of gamma radiometrics variables and gravity was more important for the depth intervals 5-15 cm and 495 15-30 cm. On the contrary, climate variables were more critical for ilr2 models, especially precipitation seasonality (PTS1 and PTS2), potential evaporation (EPA and EPX), and annual temperature (TNM and TXM), while annual temperature range (TRA) was more important for the ilr1 model. Amongst the relief covariates, DEM, topographic wetness index (TWI) and roughness were the most relevant variables for both ilr1 and ilr2. Non-photosynthetic vegetation and mean EVI were the only covariates representing the soil-forming factor organisms with some importance in the models. The most significant variables 500 predicting the distribution of coarse fragments were gamma radiometrics and other proxies of parent material (Th, U, total dose, and ratio Th/K), followed by some covariates of climate (EPX, TRA), and relief (roughness, slope, DEM) (Figure 6).



505 **Figure 5: Variable importance of random forest models for predicting ilr-transformed SOC fractions. Variable importance was calculated from permutation as per Breiman (2001).**




**Figure 6: Variable importance of random forest probability models for coarse fragments classes. Variable importance was estimated with Gini index impurity (Sandri and Zuccolotto, 2008).**

510

### 3.5 Maps of SOC fraction proportions, SOC vulnerability and coarse fragments

The spatial predictions of SOC fractions allocated most of SOC to the MAOC fraction across most of Australia (Figure 7.a and Figures S4-S6). The POC proportion was greater in the Mediterranean and temperate areas along the coast and north Queensland. There were three main regions with a high proportion of PyOC: in the north-west around Prince Regent River and Dampier Peninsula, in the west south of the Gascoyne River, and in some parts of the south-centre, e.g., east of Lake Frome. These predictions are likely driven by some calibration data with high PyOC values at these locations (Figure 2). The prediction intervals for the SOC fractions were wide (Figures S4-S6), although based on the accuracy plots (Figure S3) was estimated relatively well at the 90% confidence level. Therefore, SOC vulnerability was higher in areas with Mediterranean and temperate climate and in the centre-east (Figure 7.b). The latter is likely an artifact since that area is largely occupied by salt lakes (Kati-Thanda /Lake Eyre) and the Strzelecki desert. The coarse fragment class with highest probability across Australia was “very few” (< 2 %), and the estimated volume of coarse fragments was higher in the west and north-west Australia as well as along the Great Divide (Figures S7-S9).

520




525 **Figure 7:**a) Composite of the contribution of the three SOC fractions to SOC for the depth intervals 0-5, 5-15, 15-30 cm. The colours indicate the dominant fractions with MAOC in red, POC in green and PyOC in blue. b) SOC vulnerability for the three depth intervals. SOC vulnerability is in the log<sub>10</sub> scale for better differentiation.

### 3.6 SOC fraction stocks and sensitivity analysis

530 The spatial patterns of SOC fractions density are mainly determined by the spatial gradient in TOC concentration. Thus, irrespective of the dominant fraction, SOC density follows a climatic gradient and is higher in the south-west, east, Tasmania, and some regions in north Australia (Figures S10-S12). PyOC density was predicted higher along the Snowy Mountains, as the proportion of PyOC in montane grasslands and shrublands in the calibration dataset was around 67% (Table 3) and the TOC concentration was high (Figure S10). POC density was higher in southwestern Australia, east of Tasmania and in the southeast and east. MAOC density showed similar patterns and had higher values than POC. Ultimately, the SOC fraction stocks in the 0-30 cm (Figure 8) were not high in the east of Tasmania because these are organic soils, and the estimated median soil thickness was estimated for mineral soils. Similarly, the SOC stocks in the Snowy mountains were constrained by the shallow soil thickness. The total stock of SOC fractions for Australia (0-30 cm) were 13 Pg MAOC, 2 Pg POC and 5 Pg


535

PyOC. The predictions of SOC fractions stocks were generally smaller but correlated with the previous SOC fraction stocks estimated by Viscarra Rossel et al. (2019) (Figures S21 and S22). The Pearson correlation coefficients were respectively  $r = 0.72$  for MAOC,  $r = 0.75$  for POC, and  $r = 0.77$  for PyOC (Figure S21). The differences between the predictions of the present study and Viscarra Rossel et al. (2019) were higher in areas of shallow soils (Figure S22). We mapped the uncertainty of the SOC fractions density ( $\text{mg C cm}^{-3}$ ) by depth interval with simulations (Figures S11-S13) and estimated the uncertainty of SOC fractions stocks (0-30 cm) in a subset of pixels (Figure 9).



545 **Figure 8: SOC fraction stocks (0-30 cm) ( $\text{Mg C ha}^{-1}$ ). The values are represented in the log<sub>10</sub> scale.**

MAOC stocks were generally higher in tropical and subtropical moist broadleaf forests and temperate broadleaf and mixed forests (Figure 10). In some biomes (temperate and tropical grasslands, savannas and shrublands, and desert and xeric shrublands), MAOC stocks were higher in agricultural soils (which included rainfed and irrigated pastures), than in natural areas. In the Mediterranean biome, MAOC stocks were similar in natural and agricultural regions, where only 1.6% of the agricultural area (pastures and cropping) is irrigated (Abares, 2022). POC stocks followed similar trends than MAOC, although in the Mediterranean biome, POC was greater in agricultural soils than in natural areas. Montane grasslands and shrublands had maximum values of PyOC, although the median PyOC stocks were greater in the natural and agricultural temperate biome, followed by tropical and subtropical forests (Figure 10).




555

**Figure 9: Uncertainty of SOC fraction stocks ( $\text{Mg C ha}^{-1}$ ) for the 0-30 cm calculated with 500 simulations in a subset of 98,400 pixels.**

The two variables with most influence on the uncertainty of SOC fraction density were TOC concentration and SOC fraction % (Figures S14-S19). The total Sobol indices varied spatially depending on the SOC fraction, with TOC being the most relevant in interior areas with smaller SOC stocks, and both variables having similar influence towards the coast in the southern

560

half of the continent. Total Sobol indices show that bulk density was not very influential on the uncertainty of the SOC fraction density. This may be related to the relatively small prediction interval of the bulk density maps, calculated with bootstrapping (Viscarra Rossel et al., 2015). Coarse fragments also had a small influence as indicated by the small Sobol indices across most of Australia except in zones where a higher volume of coarse fragments had higher probability, e.g., western and northern Australia (Figures S7-S9). A sensitivity analysis was also performed for the total SOC fraction stocks (0-30 cm) at a subset of pixels to determine the influence of soil thickness on the uncertainty. The results of the first and total Sobol indices indicates that soil thickness had high influence on the SOC fraction stocks in areas of shallow soils, since we only estimated the stocks for the 0-30 cm (Figure S20). The variables with higher total Sobol indices for SOC fraction stocks were generally TOC concentration and the distribution among SOC fractions in the 5-15 cm and 15-30 cm.



570

**Figure 10: SOC fraction stock (Mg C ha<sup>-1</sup>) by biome and land use. Mineral-associated SOC (MAOC), particulate organic carbon (POC), pyrogenic organic carbon (PyOC).**




## 4 Discussion

### 4.1 Differences in SOC allocation into fractions across biomes

575 The trend of lower MAOC proportion with increasing sand content was observed across all biomes (Figure 11). Sand content is generally higher in the western half of Australia, coinciding with the Mediterranean and desert biomes, which have a mean ( $\pm$  standard deviation) sand content of  $73\% \pm 14\%$  and  $71\% \pm 10\%$ , respectively, in the calibration dataset. The smaller capacity of coarse-textured soils to stabilize SOC through organo-mineral associations may partly cause the lower proportion of MAOC and higher POC in Mediterranean soils (Figure 10 and Figure 11). Similarly, Doetterl et al. (2015) found that more SOC was stored as POC in arid environments where biochemical weathering is limited, due to a lower capacity for physico-chemical protection.

580 Conversely, the proportion of SOC found as MAOC is around 60% in Australian temperate grasslands, savannas and shrublands, 50% in Mediterranean forests and woodlands, ~54-64% in temperate forests and 65-65% in tropical (Table 3). Sokol et al. (2022) reported a high proportion of MAOC in temperate grasslands (~70%) was due to higher NPP and microbial decomposition favouring MAOC formation. They also found that savannas, temperate and tropical forests had a relatively high proportion of MAOC (~64%), whereas shrublands had a lower proportion of MAOC (~42%). Some differences are possibly explained by a higher percentage of SOC as PyOC in some systems (e.g., montane grasslands, tropical and subtropical savannas), in comparison to two fractions (MAOC and POC) (Sokol et al., 2022), and differences in the definition of biomes.



590 **Figure 11: Allocation of SOC into the mineral-associated organic carbon fraction (MAOC) plotted vs. sand content (%) by biome, from calibration data.**

### 4.2 Abiotic and biotic predictors of SOC allocation into SOC fractions at the continental scale

Climate and parent material were the main soil-forming factors for predicting the partition of SOC into fractions in continental Australia. Climate is a major driver of SOC storage (Wiesmeier et al., 2019) and partition into SOC fractions at continental scale (Bui et al., 2009). Climate influences weathering of primary minerals, the development of reactive secondary minerals

over long-term pedogenetic processes, and the chemistry of the soil solution, which in turn condition the formation of organo-mineral associations (Kleber et al., 2015). At the same time, climate also controls SOC decomposition rates and input of organic matter through net primary productivity. At the global scale, climate is a major driver of the abundance, persistence, and distribution of SOC among fractions (Heckman et al., 2022), with different effects by fractions and depth. Mean annual temperature had a strong effect on POC. The content of free POC decreased with an increase of MAT in topsoils (0-30 cm), whereas free POC increased slightly with MAT in subsoils, and occluded POC increased with MAT at all depths. In contrast, the wetness index (ratio of annual precipitation to potential evapotranspiration) had a stronger effect on MAOC, suggesting that under wetter conditions, weathering and increasing reactivity of minerals with depth, together with the downward transport of organic matter, enhances the formation of organo-mineral associations (Kleber et al., 2015). Parent material alone did not have a significant effect on the partition of SOC into MAOC at the global scale, but in interaction with the wetness index (Heckman et al., 2022).

In this study we could not verify the influence of soil geochemistry and mineralogy (metal ions, sesquioxides) on the proportion of SOC fractions due to lack of samples with extensive laboratory analysis. Soil geochemical properties have been found to control SOC storage at continental and regional scales (Doetterl et al., 2015b), and are involved in the stabilisation of SOC with different mechanisms depending on the climatic context and soil pH (Lutzow et al., 2006; Rasmussen et al., 2018). In Australia, soil physico-chemical properties, and particularly extractable iron, were the most important predictors of SOC storage at the continental scale (Li et al., 2020). Multiple soil chemical properties can be estimated fairly well with mid-infrared spectral models (Ng et al., 2022). Therefore, future research could expand on this study by investigating the relationships (and their spatial patterns) between soil chemical properties (exchangeable Ca and Mg, oxalate and dithionite extractable Fe and Al) and MAOC in the context of Australia, where the soil pH is quite acidic even under arid and semi-arid conditions.

Among vegetation variables, only EVI and the fractional cover of non-photosynthetic vegetation were important predictors for the distribution of SOC among fractions. The fraction of non-photosynthetic vegetation may be indicative of the type of C input into the soil (e.g., woody debris), which influences the subsequent decomposition and transformation pathways of organic matter (Cotrufo et al., 2015) and the allocation into SOC fractions (Heckman et al., 2022).

#### 620 **4.3 Vulnerability of SOC fractions to climate change**

We estimated the total stock of SOC fractions for Australia (0-30 cm) in 13 Pg MAOC, 2 Pg POC and 5 Pg PyOC, which give a total of 20 Pg SOC. This value is smaller than a previous baseline of SOC stock of 25 Pg SOC for Australian topsoils, but within its prediction interval (19-31.8 Pg C) (Viscarra Rossel et al., 2014). Differences in the total SOC stock may be partly due to differences in soil thickness (Figure S22) as well as in TOC predictions between the previous version of SLGA v1 (Viscarra Rossel et al., 2015; Viscarra Rossel et al., 2014) and the current SLGA v1.2 (Wadoux et al., 2022). Both maps show similar patterns and range of TOC values, and hence differences in SOC fraction and total SOC stock may be mostly caused by differences in the DSM framework (e.g., calculating SOC fraction densities prior to spatialization, accounting or not with coarse fragments, etc.) and input soil layers (soil thickness).

Our calculations estimated that about 64% of the total SOC is stored as mineral-associated SOC, which is consistent with other  
630 studies at global and continental scales (Heckman et al., 2022). In Australian topsoils, we estimated that only 10% of the SOC  
stock is stored as POC and 26% as PyOC. POC is generally more responsive than MAOC to management practices and to  
global change (Rocci et al., 2021). Our PyOC estimate is higher than the world average (14% of SOC (Reisser et al., 2016))  
but is consistent with a study in Australia (14-33%, (Lehmann et al., 2008)). Since the 1970s, there is an upward trend in “fire  
weather” conditions in Australia linked to anthropogenic climate change (Harris and Lucas, 2019), which may modify the  
635 proportion and stock of PyOC.

There is great uncertainty about the effects of an increase in temperature on SOC fractions stocks and dynamics. There is large  
evidence supporting that the temperature sensitivity of decomposition is higher for stable SOC fractions (Conant et al., 2008;  
Jia et al., 2020) or SOC pools with longer mean residence time (Li et al., 2013), although other studies indicate no differences  
or opposite trends in sensitivity between SOC fractions (Von Lützow and Kögel-Knabner, 2009). Since most of SOC is stored  
640 as MAOC, an increase in MAOC decomposition rates with temperature (when soil moisture is not limiting) may turn some  
soils into a C source (Li et al., 2013). Contrarily, coarse-textured soils with a lower capacity for physico-chemical protection  
and a greater proportion of POC may be more vulnerable to SOC loss with an increase in temperature than fine-textured soils  
(Hartley et al., 2021). Heckman et al. (2022) found a decrease in SOC persistence among all SOC fractions with higher mean  
annual temperature at the global scale, and a decrease in SOC abundance in free POC in the surface (0-30 cm). But the increase  
645 in temperature did not affect the abundance of occluded POC and MAOC, which may be less vulnerable to warming. Similarly,  
a recent meta-analysis (Rocci et al., 2021) found a reduction of POC (% SOC) with warming, suggesting that soils with higher  
POC contribution may be more vulnerable to SOC loss. In the case of Australia, this would mean that coarse-textured soils  
from Mediterranean and temperate ecosystems may be more vulnerable to an increase in temperature.

Other climatic and hydrological conditions linked to climate change may also affect SOC fractions stocks in Australia. Changes  
650 in the precipitation regime (e.g., intensity and frequency of droughts, extreme precipitation events and flooding) can affect the  
SOC fraction stocks by either limiting or enhancing C input into the soil (effects on NPP), as well as modifying decomposition  
or SOC losses by increased erosion. Rocci et al. (2021) did not find clear effects on the partition of SOC among fractions with  
an increase in precipitation, although they found a negative tendency for POC and a positive tendency on MAOC. The effect  
of wind erosion on SOC loss will depend on particle size distribution and soil cover, with vulnerable soils losing 3.6 Mg C ha<sup>-1</sup>  
655 <sup>1</sup> in south-western Australia (Harper et al., 2010). While wind erosion may deplete locally the soil of clay and silt-size particles,  
and light SOC fractions (light MAOC and light POC), and facilitate mineralization by disruption of aggregates, aeolian  
transport and deposition of may contribute to SOC enrichment in other regions (Webb et al., 2012). The influence of water  
erosion on SOC fractions will vary with agricultural practices, with the latter sometimes having a stronger effect on POC than  
erosion depending on hillslope position (Zhao et al., 2022). SOC desestabilization and stabilization processes vary along the  
660 hillslope with changes in particle size distribution, degree of weathering, and abundance of secondary minerals (Doetterl et al.,  
2015a).

#### 4.4 Components in the calculation of SOC fraction stocks: current limitations and priorities for improving their quantification

The maps of total Sobol indices inform which variables are more influential on the uncertainty of SOC fraction densities and stocks across Australia, and can guide the priorities for mapping locally, regionally or at a continental scale the different components of SOC fraction stocks. TOC was generally the main variable of influence for SOC fraction density and stock, and methods for measuring it efficiently and more economically on-farm and at the laboratory are experiencing continuous development. TOC concentration in Australian ecosystems has been underestimated by previous SOC maps in temperate forests (Bennett et al., 2020). It is then possible that in this study the SOC fraction stocks for these ecosystems are underestimated for two reasons: random forests models tend to underestimate the high values (Wadoux et al., 2022), and the SOC fraction data used to calibrate the spectral models did not represent forest ecosystems.

The percentage of TOC in the three fractions was generally the second variable of influence on the uncertainty of SOC fractions density. Still, it could be the most influential variable in areas with moderate to low SOC density. Soil thickness was the most influential variable on SOC fraction uncertainty in areas of shallow soils.

Several sources of error in the SOC fraction predictions were not accounted for in the sensitivity analysis, like the error propagated from the fractionation scheme, the different spectral models, digital soil mapping models, or the fact that the fractionation in the original dataset was applied to agricultural soils and some pastures but lack forest soils. There is also an under representation of some biomes and land cover types (e.g., tropical savannas) in the soil samples used for fractionation, which decreases the reliability of the SOC fraction stock predictions in some areas. Ideally, the spectral models could be improved by increasing the representation of different natural ecosystems (especially forests and woodlands) across biomes, which may have very different mechanisms of stabilization.

Another limitation of the current study is linked to the fractionation method used for the determination of PyOC.  $^{13}\text{C}$  NMR has been used in the past to determine the ROC in the soil (Baldock et al., 2013c). ROC has a chemical composition that is not incompatible with that of charcoal (or that is dominated in its majority by charcoal and charred plant residuals) but there is a potential presence of other poly-aryl carbon compounds that do not have a pyrogenic origin. However, the benzene polycarboxylic acid (BPCA) approach is a preferred method that gives a more realistic estimate of concentrations of PyOC in soil. Hence, future studies could carry a comparison between PyOC determined with BCPA and  $^{13}\text{C}$  NMR in Australian soil samples (Dymov et al., 2021).

We used estimates of rock fragment volume in the calculation of SOC fraction stocks, which can overestimate the stocks when the bulk density is for the whole soil instead of that of the fine soil ( $< 2$  mm) (Poeplau et al., 2017). The error due to combining volumetric coarse fragments and bulk density of the whole soil is not captured in the sensitivity analysis. Neglecting the content of coarse fragments can significantly overestimate the SOC fraction stocks in soils with non-negligible stoniness ( $> 5$  %), more than doubling the actual stocks in soils with  $> 30$ % rock fragments (Poeplau et al., 2017). We anticipated that the inaccuracy of the coarse fragments' maps and the broad range within each category would contribute significantly to the error and uncertainty in the SOC fraction stocks estimates. This was true in areas with non-negligible stoniness ( $> 2$ %), as indicated by

the total Sobol indices (Figures S14-S19). However, compared with the distribution among SOC fractions and TOC concentration, coarse fragments were not the most relevant variable influencing SOC fraction density.

## Conclusions

700 SOC fractions are crucial as input for dynamic SOC models. These maps of MAOC, POC and PyOC can be used as input for modelling SOC stocks under different management strategies and identify areas where SOC stocks could be augmented more efficiently (i.e., areas with higher SOC sequestration potential/SOC deficit) (Meyer et al., 2017; Martin et al., 2021). Land management practitioners could then optimize the spatial allocation of different agricultural practices while maintaining several soil functions and services, mainly food security and climate change mitigation and adaptation.

705 The main covariates predicting the distribution of SOC among fractions at the continental scale were identified as climate and parent material. Yet, a comprehensive and homogeneous dataset that examines the soil geochemical properties (e.g., exchangeable Ca and Mg, extractable Fe and Al, CEC ascribed to minerals, etc.) controlling SOC stabilisation processes is lacking in Australia. The diversity of climatic and pedological conditions suggests that different mechanisms will control SOC stabilization and dynamics across the continent, as observed in other regions (Rasmussen et al., 2018). The link between mycorrhizal associations (Averill et al., 2014; Jo et al., 2019) and soil microbial community composition (e.g., N<sub>2</sub>-fixing organisms), soil stoichiometry and vegetation communities (Bui and Henderson, 2013), and their effects on SOC fractions (Cotrufo et al., 2019) should be further investigated. For example, it is possible that in native ecosystems with higher soil C:N ratio and recalcitrant litter, there may be a high proportion of SOC as POC, whereas the MAOC fraction may not be C-saturated.

715 The uncertainty on the spatial predictions of SOC fraction stocks was driven mainly by TOC and the proportion of SOC fractions predictions, which in turn rely on spectral predictive models developed with soil samples originating mainly from agricultural soils. However, the sensitivity analysis allows targeting variables that should be prioritised at the local and regional scale to reduce the uncertainties of SOC fraction stock estimates. Future works shall include more efforts into sampling for measuring TOC, fractionation on underrepresented regions, or developing local spectral models for predicting SOC fractions.

## Author contributions

720 MRD, AW, BMa, BMi, AMB, RS contributed to the conceptualization and methodology of the study; MRD, BMa, RS, AW curated data and verified the reproducibility of the scripts; MRD, AW, BMa carried the formal analyses; MRD, BMa, AW, BMi prepared the manuscript with contributions from AMB and RS, MRD and BMa reviewed the manuscript with contributions from all co-authors.

## 725 Acknowledgments

The authors acknowledge the Sydney Informatics Hub and the use of the University of Sydney's high performance computing cluster, Artemis, that enabled the computations for mapping at 90 m resolution for continental Australia. We also acknowledge the high-performance computing facilities from CSIRO that were used for mapping at 90 m resolution the SOC fraction densities. Some data was sourced from Terrestrial Ecosystem Research Network (TERN) infrastructure, which is enabled by  
730 the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS). This work was partly funded by the Terrestrial Ecosystem Research Network (TERN), an Australian Government NCRIS-enabled project. MRD, AMJCW BM and AMcB acknowledge support from the Research Portfolio at the University of Sydney. AMcB acknowledges support via the Australian Research Council's Laureate Program entitled 'A calculable approach to securing Australia's soil'

### Data availability

735 The observed data on total organic carbon and coarse fragments are publicly available in their majority from the Soil Data Federator (<https://esoil.io/TERNLandscapes/Public/Pages/SoilDataFederator/SoilDataFederator.html>). The covariate data is available at <https://esoil.io/TERNLandscapes/Public/Pages/COGs/>. The R scripts used to develop the maps and figures for the paper are publicly available at <https://github.com/AusSoilsDSM/SLGA/tree/main/Production/DSM>. The spectral models and SOC fraction predictions are available upon reasonable request. The maps can be downloaded from  
740 <https://data.tern.org.au/landscapes/slga/NationalMaps>.

### References

- Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES). Land use: <https://www.agriculture.gov.au/abares/aclump/land-use>, last access: 20/09/2022.
- Aitchison, J.: Monographs on statistics and applied probability, 1986.
- 745 Arrouays, D., McBratney, A. B., Minasny, B., Hempel, J. W., Heuvelink, G., MacMillan, R., Hartemink, A., Lagacherie, P., and McKenzie, N. J.: The GlobalSoilMap project specifications, GlobalSoilMap, 494, 9-12, 2014.
- Averill, C., Turner, B. L., and Finzi, A. C.: Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage, *Nature*, 505, 543-545, 2014.
- Baisden, W. T., Amundson, R., Cook, A. C., and Brenner, D. L.: Turnover and storage of C and N in five density fractions from California annual grassland surface soils, *Global Biogeochemical Cycles*, 16, 64-61-64-16, <https://doi.org/10.1029/2001GB001822>, 2002.
- 750 Baldock, J., Beare, M., Curtin, D., and Hawke, B.: Stocks, composition and vulnerability to loss of soil organic carbon predicted using mid-infrared spectroscopy, *Soil Research*, 56, 468-480, 2018.
- Baldock, J., Sanderman, J., Macdonald, L., Allen, D., Cowie, A., Dalal, R., Davy, M., Doyle, R., Herrmann, T., Murphy, D.,  
755 and Robertson, F.: Australian Soil Carbon Research Program. v2. CSIRO. Data Collection [dataset], <https://doi.org/10.25919/5ddfd6888d4e5>, 2013a.
- Baldock, J. A., Hawke, B., Sanderman, J., and Macdonald, L. M.: Predicting contents of carbon and its component fractions in Australian soils from diffuse reflectance mid-infrared spectra, *Soil Research*, 51, 577-583, 10.1071/Sr13077, 2013b.

- 760 Baldock, J. A., Sanderman, J., Macdonald, L. M., Puccini, A., Hawke, B., Szarvas, S., and McGowan, J.: Quantifying the allocation of soil organic carbon to biologically significant fractions, *Soil Research*, 51, 561-576, 10.1071/Sr12374, 2013c.
- Beaudette, D. E., Roudier, P., and Brown, A.: aqp: Algorithms for Quantitative Pedology. R package version 1.42 [code], 2022.
- 765 Bennett, L. T., Hinko-Najera, N., Aponte, C., Nitschke, C. R., Fairman, T. A., Fedrigo, M., and Kasel, S.: Refining benchmarks for soil organic carbon in Australia's temperate forests, *Geoderma*, 368, ARTN 114246  
10.1016/j.geoderma.2020.114246, 2020.
- Bishop, T. F. A., McBratney, A. B., and Laslett, G. M.: Modelling soil attribute depth functions with equal-area quadratic smoothing splines, *Geoderma*, 91, 27-45, Doi 10.1016/S0016-7061(99)00003-8, 1999.
- Bouveresse, E. and Massart, D. L.: Improvement of the piecewise direct standardisation procedure for the transfer of NIR spectra for multivariate calibration, *Chemometrics and Intelligent Laboratory Systems*, 32, 201-213,  
770 [https://doi.org/10.1016/0169-7439\(95\)00074-7](https://doi.org/10.1016/0169-7439(95)00074-7), 1996.
- Breiman, L.: Random forests, *Mach Learn*, 45, 5-32, Doi 10.1023/A:1010933404324, 2001.
- Brus, D. J., Kempen, B., and Heuvelink, G. B. M.: Sampling for validation of digital soil maps, *Eur J Soil Sci*, 62, 394-407,  
<https://doi.org/10.1111/j.1365-2389.2011.01364.x>, 2011.
- 775 Bui, E., Henderson, B., and Viergever, K.: Using knowledge discovery with data mining from the Australian Soil Resource Information System database to inform soil carbon mapping in Australia, *Global Biogeochemical Cycles*, 23,  
<https://doi.org/10.1029/2009GB003506>, 2009.
- Bui, E. N. and Henderson, B. L.: C:N:P stoichiometry in Australian soils with respect to vegetation and environmental factors, *Plant Soil*, 373, 553-568, 10.1007/s11104-013-1823-9, 2013.
- 780 Bunemann, E. K., Bongiorno, G., Bai, Z. G., Creamer, R. E., De Deyn, G., de Goede, R., Fleskens, L., Geissen, V., Kuyper, T. W., Mader, P., Pulleman, M., Sukkel, W., van Groenigen, J. W., and Brussaard, L.: Soil quality - A critical review, *Soil Biol Biochem*, 120, 105-125, 10.1016/j.soilbio.2018.01.030, 2018.
- Burgeon, V., Fouché, J., Leifeld, J., Chenu, C., and Cornélis, J.-T.: Organo-mineral associations largely contribute to the stabilization of century-old pyrogenic organic matter in cropland soils, *Geoderma*, 388, 114841,  
<https://doi.org/10.1016/j.geoderma.2020.114841>, 2021.
- 785 Conant, R. T., Steinweg, J. M., Haddix, M. L., Paul, E. A., Plante, A. F., and Six, J.: Experimental warming shows that decomposition temperature sensitivity increases with soil organic matter recalcitrance, *Ecology*, 89, 2384-2391,  
<https://doi.org/10.1890/08-0137.1>, 2008.
- Congalton, R. G.: A review of assessing the accuracy of classifications of remotely sensed data, *Remote Sensing of Environment*, 37, 35-46, [https://doi.org/10.1016/0034-4257\(91\)90048-B](https://doi.org/10.1016/0034-4257(91)90048-B), 1991.
- 790 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and mineral-associated organic matter, *Nat Geosci*, 12, 989+, 10.1038/s41561-019-0484-6, 2019.
- Cotrufo, M. F., Soong, J. L., Horton, A. J., Campbell, E. E., Haddix, Michelle L., Wall, D. H., and Parton, W. J.: Formation of soil organic matter via biochemical and physical pathways of litter mass loss, *Nat Geosci*, 8, 776-779, 10.1038/ngeo2520, 2015.
- 795 Dangal, S. R. S., Schwalm, C., Cavigelli, M. A., Gollany, H. T., Jin, V. L., and Sanderman, J.: Improving Soil Carbon Estimates by Linking Conceptual Pools Against Measurable Carbon Fractions in the DAYCENT Model Version 4.5, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002622, <https://doi.org/10.1029/2021MS002622>, 2022.
- Doetterl, S., Cornelis, J. T., Six, J., Bodé, S., Opfergelt, S., Boeckx, P., and Van Oost, K.: Soil redistribution and weathering controlling the fate of geochemical and physical carbon stabilization mechanisms in soils of an eroding landscape, *Biogeosciences*, 12, 1357-1371, 10.5194/bg-12-1357-2015, 2015a.
- 800 Doetterl, S., Stevens, A., Six, J., Merckx, R., Van Oost, K., Pinto, M. C., Casanova-Katny, A., Munoz, C., Boudin, M., Venegas, E. Z., and Boeckx, P.: Soil carbon storage controlled by interactions between geochemistry and climate, *Nat Geosci*, 8, 780+, 10.1038/Ngeo2516, 2015b.
- Donohue, R., McVicar, T., and Roderick, M.: Fraction of Photosynthetically Active Radiation (FPAR) - AVHRR (5),  
805 Terrestrial Ecosystem Research Network [dataset], 2021.
- Dutta, R., Das, A., and Aryal, J.: Big data integration shows Australian bush-fire frequency is increasing significantly, *Roy Soc Open Sci*, 3, ARTN 150241  
10.1098/rsos.150241, 2016.

- 810 Dymov, A. A., Startsev, V. V., Gorbach, N. M., Pausova, I. N., Gabov, D. N., and Donnerhack, O.: Comparison of the Methods for Determining Pyrogenically Modified Carbon Compounds, *Eurasian Soil Science*, 54, 1668-1680, 10.1134/S1064229321110065, 2021.
- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., and Roth, L.: The shuttle radar topography mission, *Reviews of geophysics*, 45, 2007.
- 815 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quere, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragao, L. E. O. C., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschutzer, P., Lefevre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, 820 J. E. M. S., Nakaoka, S. I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rodenbeck, C., Schwinger, J., Seferian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W. P., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth Syst Sci Data*, 12, 3269-3340, 10.5194/essd-12-3269-2020, 2020.
- 825 Gallant, J. C. and Austin, J. M.: Derivation of terrain covariates for digital soil mapping in Australia, *Soil Research*, 53, 895-906, <https://doi.org/10.1071/SR14271>, 2015.
- Gallant, J. C. and Dowling, T. I.: A multiresolution index of valley bottom flatness for mapping depositional areas, *Water Resour Res*, 39, 10.1029/2002WR001426, 2003.
- 830 Ge, Y., Morgan, C. L. S., Grunwald, S., Brown, D. J., and Sarkhot, D. V.: Comparison of soil reflectance spectra and calibration models obtained using multiple spectrometers, *Geoderma*, 161, 202-211, <https://doi.org/10.1016/j.geoderma.2010.12.020>, 2011.
- Geng, J., Cheng, S. L., Fang, H. J., Pei, J., Xu, M., Lu, M. Z., Yang, Y., Cao, Z. C., and Li, Y. N.: Different Molecular Characterization of Soil Particulate Fractions under N Deposition in a Subtropical Forest, *Forests*, 10, ARTN 914 10.3390/f10100914, 2019.
- 835 Gregorich, E. G., Beare, M. H., Mckim, U. F., and Skjemstad, J. O.: Chemical and biological characteristics of physically uncomplexed organic matter, *Soil Sci Soc Am J*, 70, 975-985, 2006.
- Grundy, M. J., Rossel, R. A. V., Searle, R. D., Wilson, P. L., Chen, C., and Gregory, L. J.: Soil and Landscape Grid of Australia, *Soil Research*, 53, 835-844, <https://doi.org/10.1071/SR15191>, 2015.
- 840 Harper, R. J., Gilkes, R. J., Hill, M. J., and Carter, D. J.: Wind erosion and soil carbon dynamics in south-western Australia, *Aeolian Research*, 1, 129-141, <https://doi.org/10.1016/j.aeolia.2009.10.003>, 2010.
- Harris, S. and Lucas, C.: Understanding the variability of Australian fire weather between 1973 and 2017, *Plos One*, 14, e0222328, 10.1371/journal.pone.0222328, 2019.
- Hartley, I. P., Hill, T. C., Chadburn, S. E., and Hugelius, G.: Temperature effects on carbon storage are controlled by soil stabilisation capacities, *Nat Commun*, 12, 6713, 10.1038/s41467-021-27101-1, 2021.
- 845 Harwood, T.: 9s climatology for continental Australia 1976-2005: BIOCLIM variable suite. v1. [dataset], <https://doi.org/10.25919/5dce30cad79a8>, 2019.
- Heckman, K., Pries, C. E. H., Lawrence, C. R., Rasmussen, C., Crow, S. E., Hoyt, A. M., von Fromm, S. F., Shi, Z., Stoner, S., McGrath, C., Beem-Miller, J., Berhe, A. A., Blankinship, J. C., Keiluweit, M., Marin-Spiotta, E., Monroe, J. G., Plante, A. F., Schimel, J., Sierra, C. A., Thompson, A., and Wagai, R.: Beyond bulk: Density fractions explain heterogeneity in global 850 soil carbon abundance and persistence, *Global Change Biol*, 28, 1178-1196, 10.1111/gcb.16023, 2022.
- Hicks, W., Rossel, R. V., and Tuomi, S.: Developing the Australian mid-infrared spectroscopic database using data from the Australian Soil Resource Information System, *Soil Research*, 53, 922-931, 2015.
- Hurlbut, C. S. and Klein, C.: *Manual of mineralogy (after James D. Dana)*, Wiley1977.
- 855 Iooss, B., Da Veiga, S., Janon, A., and Pujol, G.: sensitivity: Global Sensitivity Analysis of Model Outputs. R package version 1.27.0., 2021.
- Isbell, R. F., McDonald, W. S., and Ashton, L. J.: *Concepts and rationale of the Australian Soil Classification*, CSIRO Land and Water1997.



- Jenkinson, D. S. and Rayner, J. H.: The turnover of soil organic matter in some of the Rothamsted classical experiments, *Soil science*, 123, 298-305, 1977.
- 860 Jia, Y., Kuzyakov, Y., Wang, G., Tan, W., Zhu, B., and Feng, X.: Temperature sensitivity of decomposition of soil organic matter fractions increases with their turnover time, *Land Degrad Dev*, 31, 632-645, <https://doi.org/10.1002/ldr.3477>, 2020.
- Jo, I., Fei, S. L., Oswalt, C. M., Domke, G. M., and Phillips, R. P.: Shifts in dominant tree mycorrhizal associations in response to anthropogenic impacts, *Sci Adv*, 5, 2019.
- 865 Jobbagy, E. G. and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to climate and vegetation, *Ecol Appl*, 10, 423-436, Doi 10.2307/2641104, 2000.
- Johansen, K., Gill, T., Trevithick, R., Armston, J., Scarth, P., Flood, N., and Phinn, S.: Landsat based Persistent Green-Vegetation Fraction for Australia, Proceedings of the 16th Australasian Remote Sensing and Photogrammetry Conference, Joint Remote Sensing Research, P.: Seasonal fractional cover - Landsat, JRSRP algorithm, Australia coverage (1.0.0), Terrestrial Ecosystem Research Network [dataset], 2021.
- 870 Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls, *Nat Commun*, 7, 1-10, 2016.
- Kleber, M., Sollins, P., and Sutton, R.: A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces, *Biogeochemistry*, 85, 9-24, 10.1007/s10533-007-9103-5, 2007.
- Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., and Nico, P. S.: Mineral-Organic Associations: Formation, 875 Properties, and Relevance in Soil Environments, *Adv Agron*, 130, 1-140, 10.1016/bs.agron.2014.10.005, 2015.
- Knicker, H.: Soil organic N - An under-rated player for C sequestration in soils?, *Soil Biol Biochem*, 43, 1118-1129, 10.1016/j.soilbio.2011.02.020, 2011.
- Lane, R. J. L., Wynne, P. E., Poudjom Djomani, Y. H., Stratford, W. R., Barretto, J. A., and Caratori Tontini, F.: 2019 Australian National Gravity Grids [dataset], 2020.
- 880 Lavalley, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century, *Global Change Biol*, 26, 261-273, 10.1111/gcb.14859, 2020.
- Lavalley, J. M., Conant, R. T., Haddix, M. L., Follett, R. F., Bird, M. I., and Paul, E. A.: Selective preservation of pyrogenic carbon across soil organic matter fractions and its influence on calculations of carbon mean residence times, *Geoderma*, 354, 113866, <https://doi.org/10.1016/j.geoderma.2019.07.024>, 2019.
- 885 Lehmann, J. and Kleber, M.: The contentious nature of soil organic matter, *Nature*, 528, 60-68, 10.1038/nature16069, 2015.
- Lehmann, J., Skjemstad, J., Sohi, S., Carter, J., Barson, M., Falloon, P., Coleman, K., Woodbury, P., and Krull, E.: Australian climate-carbon cycle feedback reduced by soil black carbon, *Nat Geosci*, 1, 832-835, 10.1038/ngeo358, 2008.
- Li, D., Schädel, C., Haddix, M. L., Paul, E. A., Conant, R., Li, J., Zhou, J., and Luo, Y.: Differential responses of soil organic carbon fractions to warming: Results from an analysis with data assimilation, *Soil Biology and Biochemistry*, 67, 24-30, 890 <https://doi.org/10.1016/j.soilbio.2013.07.008>, 2013.
- Li, J. Q., Nie, M., Powell, J. R., Bissett, A., and Pendall, E.: Soil physico-chemical properties are critical for predicting carbon storage and nutrient availability across Australia, *Environ Res Lett*, 15, ARTN 094088 10.1088/1748-9326/ab9f7e, 2020.
- Liang, C., Amelung, W., Lehmann, J., and Kästner, M.: Quantitative assessment of microbial necromass contribution to soil 895 organic matter, *Global Change Biol*, 25, 3578-3590, <https://doi.org/10.1111/gcb.14781>, 2019.
- Lin, L. I.-K.: A concordance correlation coefficient to evaluate reproducibility, *Biometrics*, 45, 255-268, 10.2307/2532051, 1989.
- Lugato, E., Lavalley, J. M., Haddix, M. L., Panagos, P., and Cotrufo, M. F.: Different climate sensitivity of particulate and mineral-associated soil organic matter (vol 14, pg 295, 2021), *Nat Geosci*, 15, 509-509, 10.1038/s41561-022-00945-y, 2022.
- 900 Lutfalla, S., Abiven, S., Barré, P., Wiedemeier, D. B., Christensen, B. T., Houot, S., Kätterer, T., Macdonald, A. J., van Oort, F., and Chenu, C.: Pyrogenic Carbon Lacks Long-Term Persistence in Temperate Arable Soils, *Front Earth Sc-Switz*, 5, 10.3389/feart.2017.00096, 2017.
- Lutzow, M. v., Kogel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., Marschner, B., and von Lutzow, M.: SOM fractionation methods: relevance to functional pools and to stabilization mechanisms, *Soil Biol Biochem*, 39, 2183- 905 2207, 10.1016/j.soilbio.2007.03.007, 2007.

- Lutzow, M. v., Kogel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., Flessa, H., and v. Lutzow, M.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions - a review, *Eur J Soil Sci*, 57, 426-445, 10.1111/j.1365-2389.2006.00809.x, 2006.
- 910 Lymburner, L., Tan, P., Mueller, N., Thackway, R., Thankappan, M., Islam, A., Lewis, A., Randall, L., and Senarath, U.: The National Dynamic Land Cover Dataset [dataset], 2010.
- Malley, J. D., Kruppa, J., Dasgupta, A., Malley, K. G., and Ziegler, A.: Probability machines, *Methods of information in medicine*, 51, 74-81, 2012.
- Malone, B. and Searle, R.: Improvements to the Australian national soil thickness map using an integrated data mining approach, *Geoderma*, 377, ARTN 114579
- 915 10.1016/j.geoderma.2020.114579, 2020.
- Malone, B. and Searle, R.: Updating the Australian digital soil texture mapping (Part 2\*): spatial modelling of merged field and lab measurements, *Soil Research*, 59, 435-451, 10.1071/Sr20284, 2021.
- Malone, B., Stockmann, U., Tuomi, S., and Sparrow, B.: TERN Surveillance monitoring program: Soil vis-NIR spectral library with accompanying soil measurement data for 367 specimens. v1. [dataset], <https://doi.org/10.25919/9bya-9545>, 2020.
- 920 Martin, M. P., Dimassi, B., Dobarco, M. R., Guenet, B., Arrouays, D., Angers, D. A., Blache, F., Huard, F., Soussana, J. F., and Pellerin, S.: Feasibility of the 4 per 1000 aspirational target for soil carbon: A case study for France, *Global Change Biol*, 27, 2458-2477, 10.1111/gcb.15547, 2021.
- Martinez, J.-M.: Analyse de sensibilite globale par decomposition de la variance, GdR Ondes and GdR MASCOT-NUM, Institut Henri Poincare, Paris, France, January, 13th, 20112011.
- 925 Masek, J. G., Vermote, E. F., Saleous, N. E., Wolfe, R., Hall, F. G., Huemmrich, K. F., Gao, F., Kutler, J., and Lim, T.-K.: A Landsat surface reflectance dataset for North America, 1990-2000, *IEEE Geoscience and Remote Sensing Letters*, 3, 68-72, 2006.
- McBratney, A. B., Santos, M. L. M., and Minasny, B.: On digital soil mapping, *Geoderma*, 117, 3-52, 10.1016/S0016-7061(03)00223-4, 2003.
- 930 McKenzie, N., Coughlan, K., and Cresswell, H.: Soil physical measurement and interpretation for land evaluation, CSIRO publishing2002.
- McKenzie, N., Jacquier, D., Isbell, R., and Brown, K.: Australian soils and landscapes: an illustrated compendium, CSIRO publishing2004.
- Meinshausen, N.: Quantile regression forests, *Journal of Machine Learning Research*, 7, 983-999, 2006.
- 935 Meyer, N., Bornemann, L., Welp, G., Schiedung, H., Herbst, M., and Amelung, W.: Carbon saturation drives spatial patterns of soil organic matter losses under long-term bare fallow, *Geoderma*, 306, 89-98, 10.1016/j.geoderma.2017.07.004, 2017.
- Miltner, A., Bombach, P., Schmidt-Brucken, B., and Kastner, M.: SOM genesis: microbial biomass as a significant source, *Biogeochemistry*, 111, 41-55, 10.1007/s10533-011-9658-z, 2012.
- 940 Ng, W., Minasny, B., Jeon, S. H., and McBratney, A.: Mid-infrared spectroscopy for accurate measurement of an extensive set of soil properties for assessing soil functions, *Soil Security*, 6, 100043, <https://doi.org/10.1016/j.soisec.2022.100043>, 2022.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., and Kassem, K. R.: Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity, *BioScience*, 51, 933-938, 10.1641/0006-945 3568(2001)051[0933:Teotwa]2.0.Co;2, 2001.
- Poeplau, C., Vos, C., and Don, A.: Soil organic carbon stocks are systematically overestimated by misuse of the parameters bulk density and rock fragment content, *Soil*, 3, 61-66, 2017.
- Poeplau, C., Don, A., Dondini, M., Leifeld, J., Nemo, R., Schumacher, J., Senapati, N., and Wiesmeier, M.: Reproducibility of a soil organic carbon fractionation method to derive RothC carbon pools, *Eur J Soil Sci*, 64, 735-746, 10.1111/ejss.12088,
- 950 2013.
- Poeplau, C., Don, A., Six, J., Kaiser, M., Benbi, D., Chenu, C., Cotrufo, M. F., Derrien, D., Giocchini, P., Grand, S., Gregorich, E., Griepentrog, M., Gunina, A., Haddix, M., Kuzyakov, Y., Kuhnle, A., Macdonald, L. M., Soong, J., Trigalet, S., Vermeire, M. L., Rovira, P., van Wesemael, B., Wiesmeier, M., Yeasmin, S., Yevdokimov, I., and Nieder, R.: Isolating organic carbon fractions with varying turnover rates in temperate agricultural soils - A comprehensive method comparison, *Soil Biol Biochem*, 125, 10-26, 10.1016/j.soilbio.2018.06.025, 2018.
- 955

- Rabbi, S. M. F., Linser, R., Hook, J. M., Wilson, B. R., Lockwood, P. V., Daniel, H., and Young, I. M.: Characterization of soil organic matter in aggregates and size-density fractions by solid state <sup>13</sup>C CPMAS NMR spectroscopy, *Communications in Soil Science and Plant Analysis*, 45, 1523-1537, 10.1080/00103624.2014.904335, 2014.
- 960 Rasmussen, C., Heckman, K., Wieder, W. R., Keiluweit, M., Lawrence, C. R., Berhe, A. A., Blankinship, J. C., Crow, S. E., Druhan, J. L., Pries, C. E. H., Marin-Spiotta, E., Plante, A. F., Schadel, C., Schimel, J. P., Sierra, C. A., Thompson, A., and Wagai, R.: Beyond clay: towards an improved set of variables for predicting soil organic matter content, *Biogeochemistry*, 137, 297-306, 10.1007/s10533-018-0424-3, 2018.
- Rayment, G. E. and Lyons, D. J.: *Soil chemical methods: Australasia*, CSIRO publishing 2011.
- 965 Reisser, M., Purves, R. S., Schmidt, M. W. I., and Abiven, S.: Pyrogenic Carbon in Soils: A Literature-Based Inventory and a Global Estimation of Its Content in Soil Organic Carbon and Stocks, *Front Earth Sc-Switz*, 4, ARTN 80 10.3389/feart.2016.00080, 2016.
- Robertson, A. D., Paustian, K., Ogle, S., Wallenstein, M. D., Lugato, E., and Cotrufo, M. F.: Unifying soil organic matter formation and persistence frameworks: the MEMS model, *Biogeosciences*, 16, 1225-1248, 10.5194/bg-16-1225-2019, 2019.
- 970 Rocci, K. S., Lavallee, J. M., Stewart, C. E., and Cotrufo, M. F.: Soil organic carbon response to global environmental change depends on its distribution between mineral-associated and particulate organic matter: A meta-analysis, *Sci Total Environ*, 793, 148569, <https://doi.org/10.1016/j.scitotenv.2021.148569>, 2021.
- Rowley, M. C., Grand, S., and Verrecchia, E. P.: Calcium-mediated stabilisation of soil organic carbon, *Biogeochemistry*, 137, 27-49, 10.1007/s10533-017-0410-1, 2018.
- 975 Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M., and S., T.: *Variance-based methods, in: Global Sensitivity Analysis. The Primer*, John Wiley & Sons, Ltd, 2008.
- Sanderman, J., Baldock, J., Hawke, B., Macdonald, L., Puccini, A., and Szarvas, S.: *National soil carbon research programme: field and laboratory methodologies*, 2011.
- Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil organic carbon fractions in the Great Plains of the United States: an application of mid-infrared spectroscopy, *Biogeochemistry*, 156, 97-114, 10.1007/s10533-021-00755-1, 2021.
- 980 Sandri, M. and Zuccolotto, P.: A bias correction algorithm for the Gini variable importance measure in classification trees, *Journal of Computational and Graphical Statistics*, 17, 611-628, 2008.
- Scharlemann, J. P. W., Tanner, E. V. J., Hiederer, R., and Kapos, V.: Global soil carbon: understanding and managing the largest terrestrial carbon pool, *Carbon Manag*, 5, 81-91, 10.4155/Cmt.13.77, 2014.
- 985 Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kogel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49-56, 10.1038/nature10386, 2011.
- Schoenholtz, S. H., Van Miegroet, H., and Burger, J. A.: A review of chemical and physical properties as indicators of forest soil quality: challenges and opportunities, *Forest Ecology and Management*, 138, 335-356, Doi 10.1016/S0378-1127(00)00423-0, 2000.
- 990 Searle, R.: *Australian Soil Classification Map [dataset]*, 10.25901/edyr-wg85, 2021.
- Searle, R., Stenson, M., Wilson, P. L., Gregory, L. J., Singh, R., and Malone, B. P.: *Soil data, united, will never be defeated – The SoilDataFedorator*, Joint Australian and New Zealand Soil Science Societies Conference, Cairns, QLD 2021.
- 995 Shrestha, D. L. and Solomatine, D. P.: Machine learning approaches for estimation of prediction interval for the model output, *Neural Networks*, 19, 225-235, <https://doi.org/10.1016/j.neunet.2006.01.012>, 2006.
- Singh, N., Abiven, S., Torn, M. S., and Schmidt, M. W. I.: Fire-derived organic carbon in soil turns over on a centennial scale, *Biogeosciences*, 9, 2847-2857, 10.5194/bg-9-2847-2012, 2012.
- Sokol, N. W., Whalen, E. D., Jilling, A., Kallenbach, C., Pett-Ridge, J., and Georgiou, K.: Global distribution, formation and fate of mineral-associated soil organic matter under a changing climate: A trait-based perspective, *Funct Ecol*, 36, 1411-1429, <https://doi.org/10.1111/1365-2435.14040>, 2022.
- 1000 Sollins, P., Homann, P., and Caldwell, B. A.: Stabilization and destabilization of soil organic matter: Mechanisms and controls, *Geoderma*, 74, 65-105, Doi 10.1016/S0016-7061(96)00036-5, 1996.
- Sparrow, B. D., Foulkes, J. N., Wardle, G. M., Leitch, E. J., Caddy-Retalic, S., van Leeuwen, S. J., Tokmakoff, A., Thurgate, N. Y., Guerin, G. R., and Lowe, A. J.: A Vegetation and Soil Survey Method for Surveillance Monitoring of Rangeland 1005 Environments, *Frontiers in Ecology and Evolution*, 8, 10.3389/fevo.2020.00157, 2020.

- Terrain, N. C. o. S. a.: Australian soil and land survey field handbook, Third Edition, Australian Soil and Land Survey Handbooks Series 1, 1, CSIRO PUBLISHING, Melbourne, Australia2009.
- van den Boogaart, K. G., Tolosana-Delgado, R., and Bren, M.: compositions: Compositional Data Analysis. R package version 2.0-4., 2022.
- 1010 van Leeuwen, J. P., Creamer, R. E., Cluzeau, D., Debeljak, M., Gatti, F., Henriksen, C. B., Kuzmanovski, V., Menta, C., Peres, G., Picaud, C., Saby, N. P. A., Trajanov, A., Trinsoutrot-Gattin, I., Visioli, G., and Rutgers, M.: Modeling of Soil Functions for Assessing Soil Quality: Soil Biodiversity and Habitat Provisioning, *Frontiers in Environmental Science*, 7, ARTN 113 10.3389/fenvs.2019.00113, 2019.
- 1015 Vaysse, K. and Lagacherie, P.: Using quantile regression forest to estimate uncertainty of digital soil mapping products, *Geoderma*, 291, 55-64, 2017.
- Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition: A key factor for soil organic matter formation in stable fractions, *Sci Adv*, 7, eabd3176, doi:10.1126/sciadv.abd3176, 2021.
- Viscarra Rossel, R. and Hicks, W.: Soil organic carbon and its fractions estimated by visible–near infrared transfer functions, *Eur J Soil Sci*, 66, 438-450, 2015.
- 1020 Viscarra Rossel, R. A., Webster, R., Bui, E. N., and Baldock, J. A.: Baseline map of organic carbon in Australian soil to support national carbon accounting and monitoring under climate change, *Global Change Biol*, 20, 2953-2970, 10.1111/gcb.12569, 2014.
- Viscarra Rossel, R. A., Chen, C., Grundy, M. J., Searle, R., Clifford, D., and Campbell, P. H.: The Australian three-dimensional soil grid: Australia’s contribution to the <i>GlobalSoilMap</i> project, *Soil Research*, 53, 845-864, <https://doi.org/10.1071/SR14366>, 2015.
- 1025 Viscarra Rossel, R. A., Lee, J., Behrens, T., Luo, Z., Baldock, J., and Richards, A.: Continental-scale soil carbon composition and vulnerability modulated by regional environmental controls, *Nat Geosci*, 12, 547-+, 10.1038/s41561-019-0373-z, 2019.
- von Lütow, M., Kogel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., and Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions - a review, *Eur J Soil Sci*, 57, 426-445, 10.1111/j.1365-2389.2006.00809.x, 2006.
- 1030 von Lütow, M. and Kögel-Knabner, I.: Temperature sensitivity of soil organic matter decomposition—what do we know?, *Biol Fert Soils*, 46, 1-15, 10.1007/s00374-009-0413-8, 2009.
- von Lütow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., and Marschner, B.: SOM fractionation methods: Relevance to functional pools and to stabilization mechanisms, *Soil Biology and Biochemistry*, 39, 2183-2207, <https://doi.org/10.1016/j.soilbio.2007.03.007>, 2007.
- 1035 Wadoux, A. M. J.-C., Román Dobarco, M., Malone, B., Minasny, M., McBratney, A. B., and Searle, R.: Baseline high-resolution maps of organic carbon content in Australian soils. [dataset], 2022.
- Webb, N. P., Chappell, A., Strong, C. L., Marx, S. K., and McTainsh, G. H.: The significance of carbon-enriched dust for global carbon accounting, *Global Change Biol*, 18, 3275-3278, <https://doi.org/10.1111/j.1365-2486.2012.02780.x>, 2012.
- 1040 Wiesmeier, M., Poeplau, C., Sierra, C. A., Maier, H., Frühauf, C., Hübner, R., Kühnel, A., Spörlein, P., Geuß, U., Hangen, E., Schilling, B., von Lütow, M., and Kögel-Knabner, I.: Projected loss of soil organic carbon in temperate agricultural soils in the 21st century: effects of climate change and carbon input trends, *Sci Rep-Uk*, 6, 32525, 10.1038/srep32525, 2016.
- 1045 Wiesmeier, M., Urbanski, L., Hobley, E., Lang, B., von Lütow, M., Marin-Spiotta, E., van Wesemael, B., Rabot, E., Liess, M., Garcia-Franco, N., Wollschlager, U., Vogel, H. J., and Kogel-Knabner, I.: Soil organic carbon storage as a key function of soils - A review of drivers and indicators at various scales, *Geoderma*, 333, 149-162, 10.1016/j.geoderma.2018.07.026, 2019.
- Wilford, J.: A weathering intensity index for the Australian continent using airborne gamma-ray spectrometry and digital terrain analysis, *Geoderma*, 183, 124-142, 10.1016/j.geoderma.2010.12.022, 2012a.
- 1050 Wilford, J.: A weathering intensity index for the Australian continent using airborne gamma-ray spectrometry and digital terrain analysis, *Geoderma*, 183-184, 124-142, <https://doi.org/10.1016/j.geoderma.2010.12.022>, 2012b.
- Wilford, J. and Minty, B.: The use of airborne gamma-ray imagery for mapping soils and understanding landscape processes. In ‘Digital soil mapping: an introductory perspective’. (Eds P Lagacherie, AB McBratney, M Voltz) pp. 207–218, 2007.
- Wilford, J. R. and Kroll, A.: Complete Radiometric Grid of Australia (Radmap) v4 2019 with modelled infill [dataset], 2020.
- Wilson, J. P. and Gallant, J. C.: Secondary topographic attributes, *Terrain analysis: Principles and applications*, 87-131, 2000.
- 1055 Wright, M. N. and Ziegler, A.: ranger: A Fast Implementation of Random Forests for High

Dimensional Data in C++ and R, Journal of Statistical Software, 77, 1-17, 10.18637/jss.v077.i01, 2017.

Xu, X., Liu, W., and Kiely, G.: Modeling the change in soil organic carbon of grassland in response to climate change: Effects of measured versus modelled carbon pools for initializing the Rothamsted Carbon model, Agriculture, Ecosystems & Environment, 140, 372-381, <https://doi.org/10.1016/j.agee.2010.12.018>, 2011.

1060 Zevenbergen, L. W. and Thorne, C. R.: Quantitative analysis of land surface topography, Earth Surf Proc Land, 12, 47-56, 1987.

Zhao, P., Li, L., Lin, L., Zhai, G., Cruse, R. M., and Wang, E.: Response of surface soil nutrients and organic carbon fractions to tillage erosion vs. water erosion in an agricultural landscape, Soil Sci Soc Am J, 00, 1– 13, <https://doi.org/10.1002/saj2.20461>, 2022.

1065 Zimmerman, A. R. and Mitra, S.: Trial by Fire: On the Terminology and Methods Used in Pyrogenic Organic Carbon Research, Front Earth Sc-Switz, 5, 10.3389/feart.2017.00095, 2017.

Zimmermann, M., Leifeld, J., Schmidt, M. W. I., Smith, P., and Fuhrer, J.: Measured soil organic matter fractions can be related to pools in the RothC model, Eur J Soil Sci, 58, 658-667, 10.1111/j.1365-2389.2006.00855.x, 2007.

1070 **Table 2: List of environmental covariates with unit and associated reference when applicable. All covariates are in geographic coordinates with 3 arc second grid cell (about 90 m x 90 m) resolution with coordinate system WGS84 (EPSG:4326) and extent: 112.99958°E - 153.99958°E; 10.0004°S - 44.00042°S.**

Factor	Covariate	Predictor variable	Unit	Reference
Soil	Clay	Depth-specific soil clay content (0-5 cm, 5-15 cm, 15-30 cm)	percent	Malone and Searle (2021)
	Sand	Depth-specific soil sand content (0-5 cm, 5-15 cm, 15-30 cm)	percent	Malone and Searle (2021)
Climate	ADM	Mean annual aridity index (annual precipitation/annual potential evaporation)	index	Harwood (2019)
	EPA	Annual potential evaporation	mm	Harwood (2019)
	EPI	Minimum monthly potential evaporation	mm	Harwood (2019)
	EPX	Maximum monthly potential evaporation	mm	Harwood (2019)
	Prescott	Prescott Index generated by using Prescott index = 0.445P / E0.75	index	-
	PTA	Annual precipitation	mm	Harwood (2019)
	PTI	Minimum monthly precipitation	mm	Harwood (2019)
	PTS1	Precipitation seasonality 1- solstice seasonality composite factor ratio	ratio	Harwood (2019)
	PTS2	Precipitation seasonality 2- equinox seasonality composite factor ratio	ratio	Harwood (2019)
	PTX	Maximum monthly precipitation	mm	Harwood (2019)
	RSM	Short-wave solar radiation - annual mean (SRAD data)	MJ/m2/day	Harwood (2019)
	TNM	Minimum temperature – Annual mean	°C	Harwood (2019))
	TRA	Annual temperature range	°C	Harwood (2019)
	TXM	Maximum temperature - Annual mean	°C	Harwood (2019)
WDA	Annual atmospheric water deficit (annual precipitation – annual potential evaporation)	mm	Harwood (2019)	
Organisms/ Vegetation	NDVI_Q1	Landsat 5 long-term average NDVI (January-March) 1986-2011	unitless	U.S. Geological Survey Landsat 5 Surface Reflectance Tier 1. Masek et al. (2006)
	NDVI_Q2	Landsat 5 long-term average NDVI (April-June) 1986- 2011	unitless	U.S. Geological Survey Landsat 5 Surface Reflectance Tier 1. Masek et al. (2006)
	NDVI_Q3	Landsat 5 long-term average NDVI (July-September) 1986-2011	unitless	U.S. Geological Survey Landsat 5 Surface

	NDVI_Q4	Landsat 5 long-term average NDVI (October-December) 1986 - 2011	unitless	Reflectance Tier 1. Masek et al. (2006) U.S. Geological Survey Landsat 5 Surface Reflectance Tier 1. Masek et al. (2006)
	FC_Max_BS	Landsat Fractional cover - Bare Soil -Maximum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Max_NPV	Landsat Fractional cover - Non Photosynthetic Vegetation - Maximum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Max_PV	Landsat Fractional cover - Photosynthetic Vegetation - Maximum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Mean_BS	Landsat Fractional cover - Bare Soil - Mean (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Mean_NPV	Landsat Fractional cover - Non Photosynthetic Vegetation - Mean (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Mean_PV	Landsat Fractional cover - Photosynthetic Vegetation - Mean (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Min_BS	Landsat Fractional cover - Bare Soil Minimum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Min_NPV	Landsat Fractional cover - Non Photosynthetic Vegetation - Minimum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_Min_PV	Landsat Fractional cover - Photosynthetic Vegetation - Minimum (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_SD_BS	Landsat Fractional cover - Bare Soil - Standard deviation (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_SD_NPV	Landsat Fractional cover - Non Photosynthetic Vegetation - Standard deviation (1987 – 2019)	percent	Joint Remote Sensing Research Program (2021)
	FC_SD_PV	Landsat Fractional cover - Bare Soil - Standard deviation (1987 – 2019)	percent	Joint Remote Sensing Research (2021)
	FPAR_Max	Fraction of Photosynthetically Active Radiation (FPAR) - AVHRR - Maximum Value in time series	percent	Donohue et al. (2021)
	FPAR_Mean	Fraction of Photosynthetically Active Radiation (FPAR) - AVHRR - Mean Value in time series	percent	Donohue et al. (2021)
	FPAR_Median	Fraction of Photosynthetically Active Radiation (FPAR) - AVHRR - Median Value in time series	percent	Donohue et al. (2021)
	FPAR_Min	Fraction of Photosynthetically Active Radiation (FPAR) - AVHRR - Minimum Value in time series	percent	Donohue et al. (2021)
	LC_EVI	National Dynamic Land Cover Dataset Mean of enhanced vegetation index (EVI) for the timeseries 2000 – 2008	unitless	Lymburner et al. (2010)
	Pers_Green_Veg	Landsat 2000-2010 Persistent Green-Vegetation Fraction	unitless	Johansen et al. (2012)
Relief	DEM	Elevation 3 Second - Shuttle Radar Topography Mission	meter	Farr et al. (2007)
	MRRTF	Multi-resolution Ridgetop Flatness	unitless	Gallant and Austin (2015)
	MRVBF	Multiresolution Index of Valley Bottom Flatness	unitless	Gallant and Dowling (2003)
	Plan_curv	Plan curvature	unitless	Wilson and Gallant (2000)
	Prof_curv	Profile curvature	unitless	Wilson and Gallant (2000)
	Roughness	Relief roughness	unitless	Wilson and Gallant (2000)
	Slope	Slope	percent	Zevenbergen and Thorne (1987)
	TWI	Topographic wetness index	unitless	Wilson and Gallant (2000)
Parent material/ Age	Gravity	Gravity Anomaly Grid of Australia	unitless	Lane et al. (2020)
	Dose	Radiometric grid of Australia (Radmap) v4 2019 - Filtered dose	unitless	Wilford and Kroll (2020)
	K	Radiometric grid of Australia (Radmap) v4 2019 - Potassium	percent	Wilford and Kroll (2020)
	Th	Radiometric grid of Australia (Radmap) v4 2019 - Thorium	ppm	Wilford and Kroll (2020)

U	Radiometric grid of Australia (Radmap) v4 2019 - Uranium	ppm	Wilford and Kroll (2020)
Th_K	Radiometric grid of Australia (Radmap) v4 2019 - Thorium Potassium ratio	ratio	Wilford and Kroll (2020)
U2_Th	Radiometric grid of Australia (Radmap) v4 2019 - Uranium squared to Thorium ratio	ratio	Wilford and Kroll (2020)
U_K	Radiometric grid of Australia (Radmap) v4 2019 - Uranium Potassium ratio	ratio	Wilford and Kroll (2020)
U_Th	Radiometric grid of Australia (Radmap) v4 2019 - Uranium Thorium ratio	ratio	Wilford and Kroll (2020)
WII	Weathering intensity index	unitless	Wilford (2012b)

---