

1 Reviews and Syntheses: Understanding the impacts of peatland catchment
2 management on DOM concentration and treatability

3 Jennifer Williamson^{1*}, Chris Evans¹, Bryan Spears², Amy Pickard², Pippa J. Chapman³, Heidrun
4 Feuchtmayr⁴, Fraser Leith⁵, Susan Waldron⁶, Don Monteith⁴

5

6 ¹UK Centre for Ecology & Hydrology, Environment Centre Wales, Deiniol Road, Bangor, Gwynedd, LL57 2UW

7 ²UK Centre for Ecology ~~and~~ & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB

8 ³ School of Geography, Faculty of Environment, University of Leeds, Leeds, LS2 9JT

9 ⁴UK Centre for Ecology & Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster, LA1 4AP

10 ⁵Scottish Water, 6 Castle Drive, Dunfermline, KY11 8GG

11 ⁶School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ

12 **Corresponding Author* (jwl@ceh.ac.uk)

13

14

15 **Abstract**

16 In the UK, most large reservoirs constructed for public water supply are in upland areas and situated
17 in catchments characterised that contain at least some by with organic-rich soils including peatlands ,
18 and that are often considered to be in sub-standard poor condition. Such catchments leach large
19 amounts of dissolved organic matter (DOM) leaching from these soils imparts a brownish colour to
20 water, with water draining peatlands tending to release the most. High and rising DOM concentrations
21 in these regions and raises treatment challenges for the water industry since excessive post-treatment
22 concentrations result in the generation of potentially harmful disinfection by-products in drinking
23 water. The primary method for maintaining sufficiently low pre-disinfection DOM concentrations is
24 chemical coagulation, but,

25
26 In the UK, in the past 15 years water companies have increasingly considered whether the
27 capacity for catchment upland catchment soil/peat restoration measures can interventions to slow
28 down or even reverse rising source water DOM concentrations improve raw water quality at source
29 and thus reduce, reducing the need for more costly and complex engineering solutions, in treatment
30 works. There remains considerable uncertainty around the efficacy of such effectiveness of these
31 catchment engineering-based measures, and a comprehensive overview of the research in this area
32 remains lacking. Here we review the peer-reviewed evidence for the effectiveness of four catchment
33 management options in for controlling DOM release from peat soils: for upland organic soil-
34 dominated catchments that are being considered by the water industry as options for controlling DOM
35 releases. These are ditch blocking, revegetation, reducing forest cover, and cessation of managed
36 burning.

37 Results of plot scale investigations into effects of ditch blocking on DOM leaching ditch blocking are
38 currently available but largely equivocal, while there is a paucity of information regarding impacts at
39 spatial scales of more direct relevance to water managers. There is some, although limited Although
40 not widely studied, the available evidence suggests that terrestrial the main vegetation type may
41 influence species present impacts DOM concentrations and treatability. The presence of plantation
42 forestry on peat soils is generally associated with elevated increasing DOM concentrations, although
43 canopy removal/reducing forest cover has little short-term benefit and can even exacerbate
44 concentrations further increase concentrations. Although not widely studied, the available evidence
45 suggests that Sphagnum mosses produce DOM that is more easily removed via conventional
46 treatment processes compared to vascular plants such as heather and grass species. We found
47 surprisingly little published research around the extent to which manipulation of in-reservoir
48 processes might be used to mitigate or exacerbate changes in inflowing DOM as part of a catchment
49 management approach.

50 This review concluded that catchment management measures have rarely been monitored with
51 downstream water quality as the focus, and that restoration impacts vary across sites. To mitigate the
52 uncertainty surrounding restoration effects on DOM, measures should be undertaken on a site-
53 specific basis, where the scale, effect size and duration of the intervention are considered in relation
54 to subsequent biogeochemical processing that occurs in the reservoir, the treatment capacity of the
55 water treatment works and future projected DOM trends.


56

57

58 **Introduction**

59 Peatland restoration has become an integral part of the UK environment strategy, particularly in the
60 drive toward Net Zero (HM Government, 2021). It is founded on the potential to achieve multiple
61 benefits that include improving biodiversity, enhancing carbon sequestration, and controlling water
62 runoff and quality, in catchments that are deemed to have been degraded by anthropogenic stressors.
63 Nearly three quarters of the storage capacity of drinking water reservoirs in the UK is sourced from
64 peatland areas (Xu et al., 2018). Peatlands release particularly high amounts of organic matter as The
65 dissolved organic matter (DOM) into drainage waters, and DOM concentrations of these water from
66 draining from peatlands tend to be relatively high, and have been rising since the 1980s (e.g. Naden
67 and McDonald, 1989; Robson and Neal, 1996; Harriman et al., 2001; Freeman et al., 2001; Worrall et
68 al., 2004). Mean DOM concentrations in UK Upland Waters Monitoring Network (UWMN) surface
69 waters, most of which are dominated by organic-rich soils, have approximately doubled over the last
70 three decades being approximately double those seen in the late 1980s (Figure 1). At the sub-
71 catchment scale, Chapman et al. (2010) found that water colour increased by between 22 and 155
72 percent over a 20 year period between 1986 and 2006. This phenomenon has now been observed
73 across much of industrialised North America and north-west Europe, and appears to largely result
74 largely from an long-term increase in the solubility of terrestrial organic matter as soils recover from
75 the effects of acid rain (Monteith et al., 2007; De Wit et al., 2021; Monteith et al., 2023). One
76 consequence of these changes is that water treatment works in some regions are having to adjust to
77 much higher source water DOM concentrations than they were originally designed to cope with, since
78 most were built at a time of much higher atmospheric deposition. Atmospheric deposition of
79 pollutants across the UK uplands has now declined to a very low level, and it is expected that Once
80 the trend in declining precipitation ionic strength has stabilised further in future, changes in DOM
81 export will be increasingly affected by other factors including temperature, changes in precipitation
82 seasonality and intensity and marine ion deposition (Monteith et al., 2023). Rising levels of DOM in
83 waters draining many of these peatland catchments pose considerable water treatment challenges,
84 with respect to increasing treatment costs and risks of regulatory failure (see Figure 1). It has been
85 proposed that peatland restoration measures might help slow or even reverse these DOM trends,
86 along with other important benefits including increased terrestrial carbon storage, water retention
87 and improvements in upland biodiversity but while some of the benefits of peatland restoration are
88 now becoming clear (e.g. Glenk and Martin-Ortega, 2018), evidence for impacts on water quality have
89 been more difficult to glean.

90



91

92 Figure 1: Mean (+/- Standard error) annual dissolved organic carbon (DOC) concentrations from [the](#)
 93 [236 UWMN UK Upland Water Monitoring Network](#) sites. These sites are predominately situated in the
 94 north and west of the UK – see www.uwmn.uk for more details.

95

96 Although consumption of DOM in drinking water is not directly harmful to people, coloured water
 97 reduces customer satisfaction (Ritson et al., 2014) and can be indicative of further problems.
 98 Indirectly, elevated DOM concentrations have implications for human health due to their potential
 99 influence on treatment processes and the production of carcinogenic disinfectant by-products (DBPs)
 100 such as trihalomethanes (THMs) [during chemical disinfection](#), which are regulated by the Drinking
 101 Water Inspectorate (DWI) due to their potential carcinogenic properties (Ding and Chu 2017).
 102 ~~Chlorination, a standard disinfection process in most UK WTWs, leaves free chlorine in the water supply~~
 103 ~~supply as a residual disinfectant. Free chlorine reacts with DOM remaining in the water supply~~
 104 ~~following coagulation and filtration to form DBPs, including THMs. Chloramination, the treatment of~~
 105 ~~drinking water with chlorine and ammonia to form chloramine, has been used as a method of reducing~~
 106 ~~THM formation. However, it has been found that chloramination promotes the formation of~~
 107 ~~nitrogenous DBPs (e.g. Bond et al., 2011; Lavonen et al., 2013), which are more carcinogenic than~~
 108 ~~THMs (Ding and Chu, 2017) and are likely to be regulated in the future.~~ DOM also may hamper the
 109 efficacy of chlorine as a disinfectant while simultaneously acting as a substrate for bacterial regrowth
 110 (Prest et al., 2016), thus increasing the risk of regulatory failure from bacterial contamination and the
 111 subsequent loss of customer trust.

112 The composition of DOM can have a large influence on the performance of the water treatment
 113 processes and the formation of DBPs upon chlorination (Matilainen et al., 2010). DOM in water
 114 draining peatland areas tends to be predominantly hydrophobic, and relatively photoreactive and
 115 biologically recalcitrant. It is relatively easily removed by conventional coagulation and filtration
 116 during drinking water treatment due to the presence of charged functional groups (Matilainen et al.,
 117 2010). Hydrophilic DOM, on the other hand, is mostly produced within the waterbodies by
 118 phytoplankton [activity](#) (Imai et al., 2002), and is biologically labile but less easily degraded by sunlight
 119 (Berggren and Del Giorgio, 2015; Berggren et al., 2018). The relative balance of hydrophobic to
 120 hydrophilic DOM in water is referred to as hydrophobicity, and is conventionally assessed in the water
 121 treatment system using Specific UV Absorbance measurements at 254 nm (SUVA₂₅₄), i.e. absorbance
 122 at 254 nm per unit dissolved organic carbon concentration (Weishaar et al., 2003). Values greater than


123 4 $\text{L mg}^{-1} \text{m}^{-1}$ indicate hydrophobic dominance, while values less than 2 $\text{L mg}^{-1} \text{m}^{-1}$ show the DOM is
124 primarily hydrophilic and will not be effectively removed using conventional coagulation and filtration
125 alone (Matilainen et al., 2010).

126 Higher concentrations of DOM in raw water necessitate a greater amount of treatment to provide
127 potable water to customers (Monteith et al., 2021). This may include larger coagulant dosages, shorter
128 filter run times, and longer and more frequent cleaning of filtration units, and result in higher energy
129 costs, higher sludge removal costs and an increase in direct and indirect (energy-related) greenhouse
130 gas (GHG) emissions from the treatment process (Jones et al., 2016). ~~Overall, the cost of DOM removal
131 in UK water supplies is estimated to be hundreds of millions of pounds, and has risen sharply in recent
132 years as a direct consequence of rising DOM concentrations.~~ Major additional costs are incurred
133 where capital investment is needed to upgrade treatment infrastructure designed for lower
134 concentration ranges experienced in the past (Monteith et al., 2021).

135 Peatland restoration, (i.e. physical interventions -to return them to a more natural state i.e. high water
136 table and active peat-forming vegetation) has been suggested as a catchment scale method for
137 reducing DOM concentrations in water draining peatlands (IUCN Peatland Programme). The primary
138 restoration methods undertaken to date in the UK uplands are: blocking of peatland drainage to raise
139 the water table, revegetation of bare peat with peatland species, removal of plantation forestry to
140 allow peatland species to recolonise and water tables to rise, and cessation of managed burning to
141 encourage growth of peatland plant species (Figure 2) (IUCN Peatland Programme). It is important,
142 therefore, for water industry decision makers to understand the extent to which peatland restoration
143 could make a positive contribution to reducing DOM concentrations of raw water and thus relieve
144 stresses on the treatment system and potentially remove the need for major additional capital
145 investment in treatment plant. This work reviews the available peer-reviewed literature and provides
146 a qualitative assessment of the impacts of peatland restoration on DOM concentrations and
147 treatability of raw drinking water.

148

Anthropogenic land use pressures affecting dissolved organic matter (DOM) export from peat



149

150 Figure 2: Schematic showing anthropogenic pressures on peatland catchments, and the potential
151 peatland management processes covered in this review.

152

153 2. Evidence for the efficacy of catchment management approaches in the reduction of DOM

154 ~~To answer the question “will peatland catchment management will reduce DOM concentrations in~~
155 ~~raw water” we and explored determine the evidence within the peer-reviewed scientific literature~~
156 ~~base for the efficacy of catchment management approaches within peatland dominated drinking~~
157 ~~water catchments to influence reduce DOM concentrations in the soils and waters of peatland~~
158 ~~catchments, water abstracted for drinking water treatment. This was achieved by applying we used a~~
159 ~~standard set of Boolean search terms within to search Web of Science and Google Scholar, hence~~
160 ~~including only peer reviewed literature in the review. The se terms were: (“dissolved organic matter”~~
161 ~~OR “dissolved organic carbon” OR “DOM” OR “DOC” OR “colour”) AND (“peatland” OR “bog” OR “fen”~~
162 ~~OR “moor”) AND (“ditch blocking” OR “forest” OR “plantation” OR “managed burning”). Initial results,~~
163 ~~including titles and abstracts, were rapidly reviewed to determine whether the information within the~~
164 ~~papers was relevant, both in terms of subject matter and in region (limited to temperate peatlands),~~
165 ~~then relevant papers were read in full and included in the review.~~

166 2.1. Ditch blocking

167 ~~Extensive areas of upland peatlands across the UK uplands were drained in the mid-20th century in an~~
168 ~~attempt to increase agricultural productivity. Following pPeatland drainage reduces, the resulting~~
169 ~~reductions in water tables, resulting in a loss of peat forming plant species. The, and consequent~~
170 ~~drying and cracking of peat surfaces, exposesd previously permanently saturated organic matter to~~
171 ~~oxidative processes, making themit more vulnerable to erosion and dissolution into DOM (e.g. Clark~~
172 ~~et al., 2009). Extensive efforts have been made by the water industry and organisations concerned~~
173 ~~with peatland conservation to block ditches in an attempt to restore the hydrological, biogeochemical~~
174 ~~and ecological functions of these landscapes (IUCN Peatland Programme 2023) (Figures 2 & 3).~~

175 ~~Search results of the scientific literature showed that the impact of ditch blocking on DOM~~
176 ~~concentrations had been assessed in pore waters and in ditches at streams at the sites being restored.~~
177 ~~Of the five plot-scale studies of peat soil water relevant to UK peatlandsidentified found during this~~
178 ~~review, four out of five (Table 1) reported significant changes in DOM concentrations within peat soil~~
179 ~~pore water (i.e. plot scale). The studies investigated effects between after five and twenty years~~
180 ~~following ditch blocking, and reported with a cross-study average 34% reduction in DOC concentration~~
181 ~~(range 0 to 69%) (Wallage et al., 2006; Holl et al., 2009; Haapalehto et al., 2014; Strack et al., 2015;~~
182 ~~Menberu et al., 2017). While therefore suggesting a general tendency for ditch blocking to reduce~~
183 ~~pore water DOM concentrations, these studies do not necessarily imply that effects will be translated~~
184 ~~through to surface waters and ultimately to the point of abstraction.~~

185 ~~Changes observed in DOM concentrations Aat a the drainage ditch scale, results are more variable~~
186 ~~than those for pore waters (Table 1, Figure 3). The ten eleven studies reviewed showed a mean 108%~~
187 ~~increase in DOM concentrations following ditch blocking, although this figure is skewed by the large~~
188 ~~increases reported by Worrall et al. (2007b) and Haapalehto et al. (2014) (100% increase immediately~~
189 ~~following ditch blocking and 50-75% increases after ten and five years respectively); the median~~
190 ~~change is zero. Importantly, no significant change in DOM concentration was reported in over half of~~
191 ~~these studies (O'brien et al., 2008; Gibson et al., 2009; Armstrong et al., 2010; Wilson et al., 2011;~~
192 ~~Evans et al., 2018; Pickard et al., 2022). Likewise, a recent study monitoring study DOM concentrations~~
193 ~~found no reduction in DOM concentrations in the restored site compared to the ditched site six years~~
194 ~~after ditch blocking, while both drained and restored site DOM concentrations remained elevated~~
195 ~~compared to the non-drained controlsix years after ditch blocking on a blanket bog there was found~~

196 ~~no reduction in DOM concentrations in the restored site compared to the ditched site (and both~~
 197 ~~drained and restored site DOM concentrations remained elevated compared to the non drained~~
 198 ~~control~~ (Pickard et al., 2022). Differences between studies in apparent effect size may in part be
 199 related to experimental design, including whether the work included a simultaneous control and the
 200 time period over which post-restoration monitoring was carried out.

201 Studies of DOM flux changes following ditch blocking report an average 24% reduction (range 0 – 88%
 202 reduction) in DOM flux, primarily attributed to decreased water fluxes from the restoration site.
 203 ~~However, the measurement and reporting of~~ water fluxes (and hence DOM fluxes) at a site- or
 204 catchment-scale requires careful consideration of the potential for dominant water flow pathways to
 205 be altered following ditch blocking. For example, Holden et al. (2017) showed that damming of
 206 drainage ditches in North Wales ~~did reduce~~ discharge along the original ditch lines ~~following blanket~~
 207 ~~bog re-wetting~~, but that most, or all, of the displaced flow instead left the peatland via overland flow
 208 or near-surface through-flow. Subsequent reporting from the same experiment demonstrated that
 209 DOM concentrations in water displaced along these surficial pathways were approximately the same
 210 as those in water travelling along the ditches, with the result that ditch-blocking was not found to have
 211 any clear effect on either DOM concentrations or fluxes at the catchment scale (Evans et al., 2018).
 212 ~~Studies of DOM flux changes following ditch blocking report an average 24% reduction (range 0 – 88%~~
 213 ~~reduction) in DOM flux, primarily attributed to decreased water fluxes from the restoration site.~~

214 **Table 1: Summary of the impacts of drainage ditch blocking on DOM concentrations and fluxes from peatlands, reported**
 215 **in increasing time since ditch blocking. BA = Before/After, CI = Control/Intervention**


Reference	Location	Sampling scale	Concentration or flux measured	Time since ditch blocking	Experimental Design	Change since ditch blocking
Worrall et al. (2007b)	UK, blanket bog	Ditches	DOM concentration	7 months	BACI	100% increase in DOM concentration.
Turner et al. (2013)	UK, blanket bog	0 and 1 st order ditches	DOM concentration and flux	1 year	BACI	DOM concentration decreased by 2.5% compared to control, DOM flux decreased by 2.2 – 9.2% as a result of decreased water export.
Gibson et al. (2009)	UK, blanket bog	Ditches	DOM concentration and flux	1 year	CI	DOM concentrations unchanged, water flux decreased by 39% meaning DOM flux also declined by the same amount.
Wilson et al. (2011)	UK, blanket bog	Ditches and headwater streams	DOM concentration and flux	2 years	BACI	DOM concentrations unchanged, fluxes were 88% lower in streams draining ditch-blocked catchments due to much lower estimated water export.
O'brien et al. (2008)	UK, blanket bog	Headwater streams	DOM flux and water colour	2 years	BACI	Water colour was unchanged. Fluxes decreased by 24% in streams as a result of decreasing water export.
Menberu et al. (2017)	Finland fen, pine mire and spruce mire	Pore water	DOM concentration	3 years	BACI	41% reduction in DOM concentration.

Evans et al. (2018)	UK, blanket bog	Ditches	DOM concentration	4 years	BACI	No change in DOM concentration
Wallage et al. (2006)	UK, blanket bog	Pore water	DOM concentration	5 years	CI	DOM concentration lower in porewaters adjacent to blocked ditches (69% lower compared to open ditches)
Haapalehto et al. (2014)	Finland, raised bog	Pore water	DOM concentration	5 years and 10 years	Chronosequence	DOM concentration approx. 10% lower in sites 5 years post restoration and 25% lower in sites 10 years post restoration
Haapalehto et al. (2014)	Finland, raised bog	Ditches	DOM concentration	5 years and 10 years	Chronosequence	Concentrations approx. 75% higher in sites 5 years post restoration and 50% higher in sites 10 years post restoration
Armstrong et al. (2010)	UK, blanket bog	Ditches	DOM flux	7 years	CI	No change in DOM flux
Strack et al. (2015)	Canada, bog	Pore water and ditch water	DOM concentration	10 years	CI	No change in pore water DOM concentration. Ditch water DOM concentrations were similar in spring and summer and up to 30% lower in the restored site in autumn.
Armstrong et al. (2010)	UK, blanket bog	Ditches from a survey in Northern England and Northern Scotland	DOM concentration	6 months to 18 years	Survey	DOM concentrations 28% lower on average in blocked drains compared to unblocked drains.
Holl et al. (2009)	Germany, ex-fenland extraction site	Pore water	DOM concentration	20 years	CI	DOM concentrations 37% lower at restored site compared to drained site.
Urbanova et al. (2011)	Czech Republic, bog	Pore water	DOM concentration	NA comparison between drained and intact sites	CI	No difference in DOM concentration between intact and moderately degraded site, 50% higher DOM concentrations at highly degraded site.
Pickard et al. (2022)	UK, blanket bog	Headwater streams	DOM concentration	6-8 years	CI	No difference in DOM concentration between drained and restored sites. DOM concentrations significantly higher (50% increase) in drained and restored sites compared to non-drained controls.

216

217

218



219

220 **Figure 3: Percentage change in DOM concentration following ditch blocking. Grey circles show DOM percentage change in**
 221 **peatland pore waters, and black circles show DOM percentage change in ditches and streams.**

222

223 ~~We identified N~~ine studies ~~that to date~~ have assessed the potential impact of ditch blocking on DOM
 224 treatability and hence the ease of treatability within a conventional water treatment works. ~~They~~
 225 ~~found that t~~he majority of studies at UK and continental European ditch blocking locations, along
 226 with results from their experimental work, showed little effect of ditch blocking on DOM treatability
 227 as measured by commonly reported metrics such as SUVA, E2:E3 ratios (ratio of light absorbance at
 228 250 and 365 nm) and E4:E6 ratios (ratio of light absorbance at 465 and 665 nm) (Glatzel et al., 2003;
 229 Strack et al., 2015; Gough et al., 2016; Lundin et al., 2017; Peacock et al., 2018). While none of the
 230 studies included direct measures of DOM hydrophobic and hydrophilic fractions, one measured THM
 231 formation potential and found no change between water samples taken from drained and rewetted
 232 blanket bog mesocosms (Gough et al., 2016), suggesting that in the short term ditch blocking may not
 233 reduce THM formation following water treatment.

234 More broadly, therefore, while the evidence suggests that ditch blocking may reduce DOM
 235 concentrations within pore waters (Table 3, [Figure 3](#)), there is no published evidence for such activities
 236 to have successfully influenced DOM concentrations in runoff at a catchment scale, and thus at a level
 237 of potential relevance to raw water supply to treatment works. It is important to note, however, that
 238 catchment-scale studies are hugely challenging logistically and financially to design and maintain, ~~and~~
 239 ~~are currently very rare over timescales suitable to detect land management effects on water quality.~~

240



241

242 **Figure 34:** Drainage ditches before (left) and after (right) blocking on a blanket bog in North Wales, the ditches run down
 243 the slope and individual dams can be seen crossing the ditches (Photos: Chris Evans).

244

245 **2.2. Re-vegetation of bare peat**

246 Exposure of bare peat following anthropogenic disturbance has been an extensive problem in a
 247 number of UK peatland regions, most notably in the Peak District (Pilkington et al., 2015). The
 248 subsequent erosion of the peat has caused significant problems for the water industry because of the
 249 high particulate loads from the catchment to the downstream reservoirs. There have been significant
 250 efforts in recent years to revegetate some of the most degraded upland peatland areas in order to
 251 stabilise these systems (Pilkington et al., 2015).

252 Published research on the impacts of revegetation of peatland areas on DOM is limited, but Qassim
 253 et al. (2014) found that pore water DOM concentrations were higher in revegetated sites compared
 254 to bare peat areas and vegetated controls over a five-year period. The initial revegetation mix in this
 255 work was a nurse crop of *Agrostis* sp., *Deschampsia flexuosa* and *Festuca* sp. applied in combination
 256 with additions of lime and fertiliser to ensure grass growth. Heather brash was also applied to stabilise
 257 the peat surface and provide a seed source of peatland species. The use of lime is likely to have
 258 increased DOM solubility through a reduction in acidity of the peat (Evans et al., 2012), and the re-
 259 establishment of vegetation may have increased the production of ‘new’ DOM via root leachate and
 260 fresh litter decomposition. Particulate losses from peatland systems decreased following stabilisation
 261 of the peat surface through revegetation irrespective of gully blocking activities (Pilkington et al.,
 262 2015), as overland flow velocities are lower on vegetated peat than bare peat (Holden et al., 2008).
 263 However, the same study (Pilkington et al., 2015), and more recent assessments of the effects of
 264 revegetation on DOM concentrations (Stimson et al., 2017; Alderson et al., 2019), found no long-term
 265 changes in DOM concentrations following revegetation at the headwater catchment scale.

266 Radiocarbon (¹⁴C) measurements of DOM in UK upland waters indicate that the principal source of
 267 DOM in waters draining relatively undisturbed soils is recent primary production, probably formed
 268 within the last few years (Evans et al., 2014). It follows, therefore, that plant productivity, and plant
 269 tissue composition and degradability, which depend both on ambient environmental conditions and
 270 species composition, may be important factors, both for DOM concentrations and the treatability of
 271 the DOM produced. In a laboratory-based extraction experiment, DOM leached from *Sphagnum* was
 272 more easily removed by a conventional coagulation process and decomposed more rapidly than DOM
 273 leached from *Molinia caerulea* or *Calluna vulgaris* litter. In addition, *M. caerulea* and *C. vulgaris* litter

274 released more DOM per unit dry weight compared to Sphagnum litter (Ritson et al., 2016). At the field
275 scale, published results are less clear cut: one study found that DOM concentrations in pore waters
276 were higher in areas of blanket bog dominated by *C. vulgaris* compared to areas dominated by sedges
277 or Sphagnum species (Armstrong et al., 2012). In contrast, Parry et al. (2015) found no correlation
278 between dominant vegetation type (differentiated into ericoids, grasses, sedges and bare peat) and
279 stream water DOM concentrations in headwater catchments. This may reflect the greater [biotic](#)
280 heterogeneity of peatland environments at the catchment scale in comparison to single species
281 experiments.

282 The evidence available to date suggests that while revegetation of peatland sites has stabilised bare
283 peat surfaces (e.g. Pilkington et al., 2015), and is likely to have reduced particulate organic matter loss,
284 it has not changed DOM export from peat headwater catchments. Laboratory based work has shown
285 that the species present could impact DOM treatability, with *Sphagnum* derived DOM being more
286 easily treatable than *M. caerulea* or *C. vulgaris* litter (Ritson et al., 2016). This suggests that catchment
287 management via revegetation should aim to achieve high cover of *Sphagnum* species compared to
288 vascular plants to maximise DOM treatability (Table 3), ~~although this arguably an inevitable~~
289 ~~consequence of restoring bog functionality. However, as with other restoration measures, there is~~
290 ~~currently little in the peer-reviewed literature to demonstrate the effectiveness of this at a catchment~~
291 ~~scale.~~

292

293 **2.3. Plantation forestry / deforestation**

294 It has long been recognised that forestry activities can have detrimental impacts on reservoir water
295 quality and treatability. For example, in 1984 it was shown that drainage and deforestation resulted
296 in large sedimentation issues at Crai Reservoir in south Wales (Stretton, 1984 cited in: Hudson et al.
297 1997), ~~while large pulses of nutrients (N and P) to upland streams were observed can also occur~~
298 ~~after forest-felling (Neal, 2002). This review covers the impact of ground preparation and forest~~
299 ~~planting, in-situ forest growth, and forest removal (including forest to bog restoration) on peat on~~
300 ~~DOM concentration and quality.~~

301 To reduce the impacts of forest operations on sediment and nutrient loss and consequent raw water
302 quality in the UK, the Forest and Water Guidelines now state that no more than 20% of a drinking
303 water catchment should be felled in any 3 year period (Forestry Commission, 2017). In addition to
304 this, although primarily to conserve soil carbon stocks rather than for improved water quality, the
305 2000 Forestry Commission guidance note on forest and peatland habitats (Patterson and Anderson,
306 2000) states that approval will no longer be given for forestry planting or regeneration on active raised
307 bog or inactive raised bogs that could be restored to active bog, and areas of active blanket bog greater
308 than 25 ha area and > 45 – 50 cm depth.

309 A recent review for Yorkshire Water (Chapman et al., 2017) noted that conventional conifer site
310 preparation on peat, peaty gley and peaty podzol soils would be expected to increase DOM
311 concentrations. This would be largely due to the implemented drainage reducing the height of the
312 water table and consequently increasing the production of DOM via increased aeration of the peat
313 surface (Clark et al., 2009). Jandl et al. (2007), in their review of studies of the effect of forest
314 management on soil carbon sequestration, highlighted two Finnish studies where DOM
315 concentrations increased following drainage ditch installation but returned to pre-drainage levels later
316 in the forest cycle, while Schelker et al. (2012) observed increased colour in sites being prepared for
317 forestry in northern Sweden. Furthermore, Rask et al. (1998) reported an increase in colour in streams

318 draining peat dominated catchments following afforestation in Finland, while in Sweden afforestation
 319 has also been linked to long-term increases in water colour (Skerlep et al., 2019). At a regional to
 320 national scale in the UK, recent work suggests that the presence of plantation forestry on peat soils [is](#)
 321 [associated with higher increases](#) DOM concentrations in streams and rivers compared to peat soils
 322 [supporting with](#) semi-natural vegetation (Williamson et al., 2021).

323

324 Table 2: UK studies reporting DOM concentration monitoring of forestry activities on peat. Note that
 325 where percentage differences are preceded by ~ concentrations were not explicitly listed in text,
 326 figures and tables or supplementary information so are estimated from graphs.

Paper	Location	Forestry activity monitored	Scale	Timescale of monitoring	% difference
Muller and Tankere-Muller (2012)	Flow Country	Felling compared to blanket bog	Stream (upstream and downstream)	1 year post felling	-6%
Zheng et al. (2018)	Central Scotland	Felling compared to windfarm on blanket bog	Stream	1 year ~ 8 years after felling	~ 100%
Muller et al. (2015)	Flow Country	Felling compared to blanket bog	Stream	3 months before ~ 1 year after	No difference
Shah and Nisbet (2019)	Central Scotland (raised bog)	Before / after felling	Stream	1 year before and up to 8 years after	0%, 29% & 51% (mean 27%)
Cummins and Farrell (2003)	Ireland	Before / after felling	Stream	5 years	~0 – 100%
Gaffney et al. (2020)	Flow Country	Before / after felling and felling compared to blanket bog	Stream	2 years	No significant difference
Muller et al. (2015)	Flow Country	Before / after felling	Ditch	3 months before ~ 1 year after	~ 75%
Gaffney et al. (2018)	Flow Country	Before / after felling	Ditch	1 year post felling	~ 150%
Cummins and Farrell (2003)	Ireland	Before / after felling	Ditch	5 years	~50%
Gaffney et al. (2018)	Flow Country	Felling compared to blanket bog	Ditch	0 – 17 years post felling. 1 year of measurement	~500%
Muller and Tankere-Muller (2012)	Flow Country	Felling compared to blanket bog	Ditch	1 year post felling	30-325% (overall average 159%)

Gough et al. (2012)	North Wales	Presence / absence of forestry	Pore waters	<u>1 off sampling</u>	-19% - 111% (average 45%)
Howson et al. (2021)	Flow Country	Presence / absence of forestry	Pore waters	<u>~ 20 months</u>	~ 66%
Howson et al. (2021)	Central Scotland (raised bog)	Presence / absence of forestry	Pore waters	<u>~ 20 months</u>	~14%
Flynn et al. (2022)	Ireland	Presence / absence of forestry	Pore waters	<u>~ 2 years</u>	~400%
Gaffney et al. (2018)	Flow Country	Presence / absence of forestry	Ditch	<u>0 – 17 years post felling 1 year of measurement</u>	~ 100%
Flynn et al. (2022)	Ireland	Presence / absence of forestry	Stream	<u>~ 2 years</u>	No significant difference
Shah et al. (2021)	Flow Country	Presence / absence of forestry – time series	Stream	<u>25 years</u>	No significant difference
Cummins and Farrell (2003)	Ireland	Presence / absence of forestry	Stream	<u>5 years</u>	~25%

327

328 The presence of ~~conifersforestry~~ on peat soils in a UK and Irish context is associated with higher pore
329 water DOM concentrations across the four studies covered in this review (Table 2), with a mean
330 difference of approximately 130%. The exception to this pattern was found in spruce plantations in
331 north Wales where DOM concentrations in pore waters were 19% lower than in adjacent blanket bog
332 (Gough et al., 2012). We found only one study (Gaffney et al., 2018) ~~that compared~~ DOM
333 concentrations ~~in drainage ditches at a ditch scale~~ between forested and intact blanket bog areas,
334 with DOM concentrations ~~being~~ approximately 100% higher in ~~ditches draining the the former forested~~
335 ~~areas. At the stream scale~~ The presence of forestry on peat had less clear cut impacts on ~~streamwater~~
336 DOM concentrations, with two out of three studies reporting no significant difference between
337 streams draining catchments with forestry and intact blanket bogs (Shah et al., 2021; Flynn et al.,
338 2022), and the third showing an DOM concentrations approximately 25% higher in a stream draining
339 a forested catchment compared to a blanket bog catchment (Cummins and Farrell, 2003).

340 Tree felling tends to ~~produce larger increases~~ ~~cause an increase~~ in DOM, though the effects are not
341 universal across studies and locations. ~~At the stream scale~~ Three of five studies ~~of streamwater DOM~~
342 ~~concentrations~~ reported increases following felling (Cummins and Farrell, 2003; Zheng et al., 2018;
343 Shah and Nisbet, 2019), with a mean increase of approximately 43%, although the two studies in the
344 ~~Thurso catchment~~ ~~Flow Country~~ showed no change (Muller et al., 2015) and a 6% decrease in
345 concentrations (Muller and Tankere-Muller, 2012), which was attributed to the success of buffer strips
346 between the plantation and the monitored stream. ~~At the ditch scale~~ The mean increase in DOM
347 concentrations ~~in ditches~~ was nearly 200% (ranging from a 50% increase to a 500% increase, see Table

2) (Cummins and Farrell, 2003; Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2018). Most studies measuring DOM concentrations from forestry on peat were relatively short-term in timeframe, lasting two years or shorter. Only two studies monitored DOM concentrations for five years or longer.

There has been comparatively little research on the effects of forest presence on the treatability of DOM, although Gough et al. (2012) evaluated DOM concentrations and SUVA₂₅₄ values in waters draining catchments forested with different tree species. They found that pore water leachates from pine and larch plantation yielded particularly high DOM concentrations relative to a blanket bog control (19 and 13 mg L⁻¹, respectively, compared to 9 mg L⁻¹). Leachates also had lower SUVA₂₅₄ values (1.2 and 2.4 respectively, compared to 3.3 L mg⁻¹ m⁻¹). This would suggest that DOM leaching from plantations dominated by these tree types may be less easily treatable than DOM from blanket bogs. Similarly, samples taken from Scottish blanket and raised bog sites (Howson et al., 2021) found that SUVA₂₅₄ values were lower from forested sites, again suggesting that forestry on peat results in less aromatic, hydrophobic DOM that may be less easily removed via conventional coagulation.

Recently there have been attempts to restore previously afforested fen and bog peatlands in parts of Europe and North America under what is often referred to as 'forest-to-bog' restoration (Chimner et al., 2017; Andersen et al., 2017). Although still a relatively new practice within the UK, this type of restoration has been carried out for 18 years in the Flow Country in northern Scotland, and national policies on peat restoration may lead to its expansion in the future. Some of the studies listed in Table 2 (Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2018; Shah and Nisbet, 2019; Gaffney et al., 2020; Howson et al., 2021; Shah et al., 2021) monitored the impacts of felling as part of ongoing forest-to-bog restoration monitoring, with the main differences in management being that the trees were felled to waste (the practice of leaving felled trees *in-situ* to rot) and there was less ground disturbance at the site compared with the use of machinery to extract felled timber (Gaffney, 2017). However, the practice of felling trees to waste has been suggested to provide a potential additional DOM source as the trees slowly decompose (Muller et al., 2015), with mulched fallen trees providing a major source of water soluble DOM (Howson et al., 2021).

As bog vegetation regenerated after such restoration in the Flow Country, DOM concentrations reduced from elevated levels towards those seen in forest control areas. T, although the time frame for complete recovery to pre-intervention levels is to date inconsistent, with some areas still showing elevated DOM in the restoration sites relative compared to the control sites after 17 years (Gaffney et al., 2018). However, in others, DOM concentrations had returned to those seen in intact blanket bog within the same time frame (Howson et al., 2021), or were showing inconsistent effects across sub-catchments, with the most upstream catchments showing increased DOM concentrations compared to bog controls, an effect not seen further downstream (Pickard et al., 2022). Other studies have reported shorter-term perturbation increases in DOM (~4-5 years) following forest-to-bog restoration, including within an assessment of forest to bog restoration of a Scottish lowland raised bog area, Flanders Moss, where stream water baseline DOM levels were reached within two years at one site (Shah, 2018). In a Finnish study of the impacts of forest to mire restoration, a short-term peak in pore water DOM concentration following initial restoration activity was followed by a return to reference concentrations within six years (Menberu et al., 2017).

Management of peatland for conifer plantation In summary, coniferous afforestation of peatlands increases DOM concentrations in pore waters and streams, both during site establishment, potentially during the forest growth, and again as the trees are felled (by up to 500%) (summarised in Table 3). Forest-to-bog restoration as a method of land management produces short-term increases in DOM concentrations while trees are felled and brash remaining on site decomposes. However, given a long

394 enough timeframe, DOM concentrations appear to reduce back towards levels seen from comparable
395 control locations. ~~From a water company perspective it is important to~~ ~~Water companies should~~ note
396 that this time frame can be up to 20 years in blanket bogs, ~~i.e. a time frame~~ considerably longer than
397 the standard funding cycle.

398 2.4. Managed burning

399 Managed burning of peatland vegetation (Figures ~~2 & 45~~) (primarily ~~the burning of Calluna sp. heather~~
400 ~~as part of~~ ~~for~~ grouse moor management) is a contentious issue within peatland conservation and
401 management (e.g. Davies et al., 2016) and has been extensively reviewed over the past decade,
402 particularly in relation to the impacts on DOM (Worrall et al., 2010; Holden et al., 2012; e.g. Brown et
403 al., 2015), and most recently by Harper et al. (2018). There is little evidence within these reviews to
404 suggest that DOM concentrations or colour increase within ~~peat~~ pore waters ~~at the plot scale~~ following
405 managed burns. A recent study showed no change in DOM concentrations following low and high
406 intensity burning (Grau-Andres et al., 2019), and in previous studies ~~plot scale pore water~~ DOM
407 concentrations were unchanged (Clay et al., 2009; Clay et al., 2012; Worrall et al., 2013) or decreased
408 (Worrall et al., 2007a). At the catchment scale, ~~positive correlations between the extent of burning~~
409 ~~and DOM concentrations and water colour have been interpreted as causal~~ ~~it has been suggested that~~
410 ~~managed burning contributes to increases in water colour and DOM concentrations~~ (Clutterbuck and
411 Yallop, 2010; Yallop et al., 2010; Ramchunder et al., 2013) ~~although this has been questioned- in the~~
412 ~~literature~~ (Holden et al., 2012) ~~by others~~. Burning as a management practice is designed to ensure that
413 there is a mosaic of ~~variously different~~ aged heather habitat so it seems plausible that these effects
414 are linked to changes in vegetation cover. As previously discussed *C. vulgaris* produced higher amounts
415 of DOM than *Sphagnum* in the laboratory (Ritson et al., 2016) and at plot scale (Armstrong et al.,
416 2012). It is also worth noting that Evans et al. (2017b) found that a wildfire in Northern Ireland resulted
417 in a temporary reduction of DOM concentrations in a downstream monitoring lake, which was
418 attributed to re-acidification of catchment soils following the fire.



419
420 **Figure 45: Burning of vegetation on peat in North Wales (Photo: Chris Evans).**

421
422 Table 3: summary of the published impacts of catchment management activities on DOM concentrations and treatability,
423 focussing on those studies relevant in a UK and Irish context. Numbers in brackets refer to the number of studies showing
424 that effect in each case, while the overall impacts on DOM concentration and treatability for water treatment are shown
425 as +/- (positive/neutral/negative) for concentrations and treatability respectively.

<u>Catchment intervention</u>	<u>Impact on DOM concentration</u>	<u>Impact on DOM treatability</u>
<u>Ditch blocking (=/=)</u>	Increase (2) (<u>Worrall et al., 2007b; Haapalehto et al., 2014</u>) No change (8) (<u>O'brien et al., 2008; Gibson et al., 2009; Armstrong et al., 2010; Wilson et al., 2011; Urbanova et al., 2011; Turner et al., 2013; Strack et al., 2015; Evans et al., 2018</u>) Decrease (5) (<u>Wallage et al., 2006; Holl et al., 2009; Armstrong et al., 2010; Haapalehto et al., 2014; Menberu et al., 2017</u>)	No change (5) (<u>Glatzel et al., 2003; Strack et al., 2015; Gough et al., 2016; Lundin et al., 2017; Peacock et al., 2018</u>)
<u>Revegetation (to grass species) (=/-)</u>	Increase (2) (<u>Qassim et al., 2014; Ritson et al., 2016</u>) No change (4) (<u>Parry et al., 2015; Pilkington et al., 2015; Stimson et al., 2017; Alderson et al., 2019</u>)	Decrease (1) (<u>Ritson et al., 2016</u>)
<u>Revegetation (to heather) (-/-)</u>	Increase (2) (<u>Armstrong et al., 2012; Ritson et al., 2016</u>) No change (1) (<u>Parry et al., 2015</u>)	Decrease (1) (<u>Ritson et al., 2016</u>)
<u>Revegetation (to Sphagnum) (+/+)</u>	Decrease (1) (<u>Armstrong et al., 2012</u>)	Improve (1) (<u>Ritson et al., 2016</u>)
<u>Forest presence (-/-)</u>	Increase (5) (<u>Cummins and Farrell, 2003; Gough et al., 2012; Gaffney et al., 2018; Howson et al., 2021; Flynn et al., 2022</u>) No change (2) (<u>Shah et al., 2021; Flynn et al., 2022</u>)	Decrease (2) (<u>Gough et al., 2012; Howson et al., 2021</u>)
<u>Clearfell and forest-to-bog conversion (-/-)</u>	Increase (6) (<u>Cummins and Farrell, 2003; Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2018; Zheng et al., 2018; Shah and Nisbet, 2019</u>) No change (3) (<u>Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2020</u>)	Decrease (1) (<u>Zheng et al., 2018</u>)
<u>Managed burning (-/no evidence)</u>	Increase (3) (<u>Clutterbuck and Yallop, 2010, Yallop et al., 2010, Ramchunder et al., 2013</u>) No change (4) (<u>Clay et al., 2009; Clay et al., 2012; Worrall et al., 2013; Grau-Andres et al., 2019</u>) Decrease (1) (<u>Worrall et al., 2007a</u>)	

426 **Table 3: summary of the published impacts of catchment management activities on DOM concentrations and treatability,**
427 **focussing on those studies relevant in a UK and Irish context. Numbers in brackets refer to the number of studies showing**
428 **that effect in each case. Colour coding shows whether the overall conclusion is that effects are positive (green), no /**
429 **limited change (yellow), or negative (red).**

Catchment intervention	Impact on DOM concentration	Impact on DOM treatability
Ditch blocking	Increase (2) (Worrall et al., 2007b; Haapalehto et al., 2014) No change (8) (O'Brien et al., 2008; Gibson et al., 2009; Armstrong et al., 2010; Wilson et al., 2011; Urbanova et al., 2011; Turner et al., 2013; Strack et al., 2015; Evans et al., 2018) Decrease (5) (Wallage et al., 2006; Holl et al., 2009; Armstrong et al., 2010; Haapalehto et al., 2014; Menberu et al., 2017)	No change (5) (Glatzel et al., 2003; Strack et al., 2015; Gough et al., 2016; Lundin et al., 2017; Peacock et al., 2018)
Revegetation to grass species	Increase (2) (Qassim et al., 2014; Ritson et al., 2016) No change (4) (Parry et al., 2015; Pilkington et al., 2015; Stimson et al., 2017; Alderson et al., 2019)	Decrease (1) (Ritson et al., 2016)
Revegetation to heather	Increase (2) (Armstrong et al., 2012; Ritson et al., 2016) No change (1) (Parry et al., 2015)	Decrease (1) (Ritson et al., 2016)
Revegetation to <i>Sphagnum</i>	Decrease (1) (Armstrong et al., 2012)	Improve (1) (Ritson et al., 2016)
Forest presence	Increase (5) (Cummins and Farrell, 2003; Gough et al., 2012; Gaffney et al., 2018; Howson et al., 2021; Flynn et al., 2022) No change (2) (Shah et al., 2021; Flynn et al., 2022)	Decrease (2) (Gough et al., 2012; Howson et al., 2021)
Clearfell and forest to bog conversion	Increase (6) (Cummins and Farrell, 2003; Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2018; Zheng et al., 2018; Shah and Nisbet, 2019) No change (3) (Muller and Tankere-Muller, 2012; Muller et al., 2015; Gaffney et al., 2020)	Decrease (1) (Zheng et al., 2018)
Managed burning	Increase (3) (Clutterbuck and Yallop, 2010; Yallop et al., 2010; Ramchunder et al., 2013) No change (4) (Clay et al., 2009; Clay et al., 2012; Worrall et al., 2013; Grau-Andres et al., 2019) Decrease (1) (Worrall et al., 2007a)	

430

431 [Discussion and conclusions](#)

432 **3: ~~Catchment management impacts on downstream DOM processing~~Discussion and conclusion**

433 ~~As indicated by~~ Table 3 summarises the range and extent of the current peer-reviewed evidence for
434 the impacts of peatland restoration on DOM concentrations in raw water and the treatability of the
435 DOM present. However, ~~there remain~~ considerable knowledge gaps remain regarding in the area of
436 the effects of peatland restoration on raw water DOM concentrations and treatability. Our thorough
437 screening of the literature revealed remarkably few published studies in this area, to the extent that
438 generalisations of the effects of most of the interventions examined must be taken with considerable
439 cautions.

440 The available literature does indicate that ~~This review highlights that~~ both revegetation of bare peat
441 (particularly to *Sphagnum* dominated bog) and ditch blocking ~~is have been~~ associated with decreased
442 DOM concentrations within pore waters and ditches at the location restoration occurs. ~~The available~~
443 evidence also suggests, again at this local scale, that ~~plantation forestry presence and felling tend to~~
444 lead to increasing DOM concentrations and potentially reduced treatability of exported DOM.
445 However, ~~and in contrast to much more widely reported positive impacts of these restoration actions~~
446 with respect to carbon sequestration, soil particulate losses, flood management and upland
447 biodiversity, ~~the evidence that such impacts may translate to comparable changes within the wider~~
448 catchments for impacts at the stream scale of more relevance to drinking water resources is generally
449 lacking ~~more equivocal~~.

450 There is arguably much stronger evidence pointing to the risks posed by the afforestation of peatlands,
451 and the subsequent management of such plantations, with ~~The available evidence also suggests, again~~
452 at this local scale, that ~~plantations forestry presence and felling tending to lead to increasing DOM~~
453 concentrations and potentially reduced treatability of exported DOM. In the published literature we
454 have been unable to find experimental evidence incorporating local changes in water chemistry in the
455 vicinity of interventions with downstream DOM processing to show whether water quality effects are
456 detectable at the point of abstraction for water treatment works. This extension beyond the plot and
457 hillslope scale represents a significant gap in current understanding, as DOM processing continues
458 within the aquatic environment downstream of peatland restoration sites.

459 Robust quantification of the impacts of catchment management on DOM concentration and
460 treatability at the point of abstraction clearly represents a major current evidence gap. The size of the
461 research challenge with respect to the necessary spatial and temporal scale and need for robust
462 Before-After-Control Impact (BACI) of any field experiment cannot be underestimated, and perhaps
463 explains in part the current dearth of reliable information. This is particularly pertinent when changes
464 in water chemistry may take a number of years to be seen, depending on catchment dynamics and
465 within reservoir processes. Our review has highlighted ~~that that in-reservoir biogeochemical processes~~
466 ~~should be considered alongside~~ catchment land management approaches ~~by the water industry to~~
467 ~~maximise the potential for upstream solutions to rising DOM concentrations in source waters have~~
468 ~~not been followed downstream to monitor their impacts to the wider catchment.~~

469 The general paucity of evidence to support widespread terrestrial-catchment focussed interventions
470 specifically to manage source water DOM concentrations and treatability leads then to the question
471 as to whether there are other water quality management options that could be applied within
472 reservoirs.

473 DOM in ~~s not conservatively mixed through~~ rivers and lakes ~~but~~ is subject to both biotic and abiotic
474 processing, which change both concentrations and chemical structure (e.g. Tranvik et al., 2009) ~~and~~
475 ~~hence affect treatability. For example DOM is lost to ~~Loss pathways for DOM include:~~ respiration~~
476 (Koehler et al., 2012; Stets et al., 2010), sedimentation (Einola et al., 2011; Von Wachenfeldt and

477 Tranvik, 2008), photo-oxidisation (via UV radiation) (Moody et al., 2013; Koehler et al., 2014) and
478 flocculation with naturally-occurring aluminium and iron (Mcknight et al., 1992; Koehler et al., 2014).

479 More importantly for treatability, however, DOM is generated within lakes and reservoirs via
480 photosynthesis (production of algal exudates and release via cell lysis) and through processing of
481 particulate matter (Tranvik et al., 2009) so that DOM concentrations at the point of abstraction from
482 reservoirs represent the sum of these removal and generation processes. Consequently, the resulting

483 DOM tends to be produced via these processes is relatively transparent and hydrophilic in comparison
484 with DOM generated by organic rich soils, and thus presents different challenges for treatment,
485 particularly as the hydrophilic DOM is not easily removed through coagulation (Matilainen et al., 2010)
486 and may lead to the need for additional capital investment in order to effectively reduce residual DOM
487 in drinking water.

488 Importantly, in-reservoir aAlgal production, and hence within-reservoir generation of DOM, is often
489 limited by the availability of phosphorus, nitrogen or both. Hence, waterbodies with high
490 concentrations of inorganic nutrients, either delivered externally from their catchments or re-released
491 internally from sediments, are likely to generate additional DOM within the water column
492 (Feuchtmayr et al., 2019; Evans et al., 2017a). Further, evidence is growing on the importance of lake
493 and reservoir bed sediments as a direct source of DOM to the water column, with reducing conditions
494 occurring during stratification of lakes and reservoirs causing redissolution of previously sedimented
495 organic matter (Peter et al., 2017).

496 In their assessment of DOM in lake inflows and outflows, including those of several reservoirs, Evans
497 et al. (2017a) concluded that any measures that can reduce N and P export from the catchment or
498 release from sediments, or which can strip nutrients from the water column, could provide effective
499 mitigation for high DOM concentrations by reducing algal DOM production. For example, measures
500 for reducing nutrient loading to lakes from the catchment (Spears and May, 2015) and bed sediments
501 (Spears et al., 2016) can be effective in reducing algal biomass in UK lakes - although the effects on
502 algal DOM production in relation to drinking water treatment require further assessment. To date,
503 this option has rarely been considered in relation to DOM-related treatment issues, although nutrient
504 management is often considered in relation to other (taste and odour) related treatment issues. The
505 available evidence therefore suggests that measures to reduce taste and odour problems could deliver
506 co-benefits in relation to DOM levels.

507 It is pertinent, therefore, to consider ~~A future research focus should therefore include answering the~~
508 ~~question of~~ whether measures which reduce in-reservoir DOM production, and/or favour in-reservoir
509 DOM removal, may be as – or perhaps more – effective than measures aimed at reducing DOM export
510 from the terrestrial catchment. For lakes acting as DOM sources, management regimes that reduce
511 nutrient (primarily N and P) inputs from catchments and/or internal loading of nutrients and DOM
512 from sediment to the water column may be more effective than those focussed on reducing inflowing
513 DOM concentrations directly. Restricting nutrient inputs is also likely to reduce organic nitrogen
514 concentrations relative to organic carbon concentrations, which has the added benefit of reducing the
515 formation potential of nitrogenous DBPs. In addition, Birk et al. (2020) suggest that rising DOM loading
516 from the catchment may act to dampen algal responses to nutrients through light limitation of primary
517 production within some European lakes. If, by extension, this also limits in-reservoir DOM production
518 then catchment interventions that relieve DOM load, but not nutrient load, may result in an increase
519 in in-reservoir DOM production. Even in the case of less nutrient-rich water bodies, it appears that
520 reducing N and P loadings would be beneficial for water treatment as this is likely to restrict additional
521 DOM formation.

522 In summary, our review demonstrates that catchment management initiatives, while providing clear
523 overall restoration benefits for peatlands, have yet to deliver a generalised solution to the challenge
524 of stabilising or reversing DOM increases in drinking water sources, although there is some evidence
525 that catchment interventions may provide benefits for DOM export in specific cases. Catchment
526 management measures that reduce in-reservoir DOM production, or favour in-reservoir DOM
527 removal, may be as or more effective, particularly with respect to more nutrient rich systems. More
528 generally, it seems clear that catchment management should be considered part of the response
529 strategy to rising DOM levels, and as part of a process to improve the resilience of source waters, not
530 a panacea. It is therefore important that the water industry also develops effective tools to predict
531 likely future DOM levels resulting from a combination of large-scale and catchment-scale drivers, to
532 ensure that investments in both catchment management measures and DOM treatment
533 infrastructure are correctly targeted, integrated, timely and cost-effective.

537 **4- Conclusions**

538 ~~Increasing DOM concentrations in reservoirs draining catchments dominated by peat soils are a cause~~
539 ~~for concern for water companies, from both regulatory compliance and treatment cost perspectives.~~
540 ~~To a large extent this increase appears to be a long-term large-scale phenomenon, driven by~~
541 ~~improvements in air quality, and thus beyond the direct control of catchment managers. While it is~~
542 ~~likely that atmospheric deposition driven changes in DOM are beginning to level off it is also feasible~~
543 ~~that future climate change could also contribute to further increases in concentrations. The~~
544 ~~production of DOM in peat soils, for example, is known to be highly sensitive to soil temperature (Clark~~
545 ~~et al., 2009) while long-term increases in precipitation have also been linked with DOM increases (De~~
546 ~~Wit et al., 2021).~~

547 ~~To date, catchment management initiatives, while providing clear overall restoration benefits for~~
548 ~~peatlands, do not appear to have produced a generalised solution to the challenge of stabilising or~~
549 ~~reversing DOM increases in drinking water sources, although there is some evidence that catchment~~
550 ~~interventions may provide benefits for DOM export in specific cases. We have identified some areas~~
551 ~~where there is mounting evidence for the importance of certain catchment interventions. In~~
552 ~~particular, short-term effects of forest felling and harvesting activities have repeatedly shown to have~~
553 ~~detrimental effects on DOM concentrations. Catchment interventions may also provide co-benefits~~
554 ~~such as reductions in sediment and particulate organic carbon loadings to reservoirs, reductions in~~
555 ~~greenhouse gas emissions and enhancement of biodiversity, which may justify the implementation of~~
556 ~~measures when all benefits are combined, even if the direct benefits for DOM alone may not.~~

557 ~~Our review of the published literature highlights a major current evidence gap of importance to the~~
558 ~~water industry: the quantification of the impacts of catchment management on DOM concentration~~
559 ~~and treatability at the point of abstraction. The size of the research challenge with respect to the~~
560 ~~necessary spatial and temporal scale and need for robust Before-After-Control Impact (BACI) of any~~
561 ~~field experiment cannot be underestimated, and perhaps explains in part the current dearth of reliable~~
562 ~~information. This is particularly pertinent when changes in water chemistry may take a number of~~
563 ~~years to be seen, depending on catchment dynamics and within-reservoir processes. Our review has~~
564 ~~highlighted that in-reservoir biogeochemical processes should be considered alongside catchment~~

565 ~~land management approaches by the water industry to maximise the potential for upstream solutions~~
566 ~~to rising DOM concentrations in source waters.~~

567 ~~Catchment management measures that reduce in-reservoir DOM production, or favour in-reservoir~~
568 ~~DOM removal, may be as or more effective, particularly with respect to more nutrient rich systems.~~
569 ~~More generally, it seems clear that catchment management should be considered part of the response~~
570 ~~strategy to rising DOM levels, and as part of a process to improve the resilience of source waters, not~~
571 ~~a panacea. It is therefore important that the water industry also develops effective tools to predict~~
572 ~~likely future DOM levels resulting from a combination of large-scale and catchment-scale drivers, to~~
573 ~~ensure that investments in both catchment management measures and DOM treatment~~
574 ~~infrastructure are correctly targeted, integrated, timely and cost-effective.~~

575

576 The authors declare that they have no conflict of interests.

577

578 The work was conceptualised by DM and CE, funding acquisition was by DM, CE and BS, JW carried
579 out the initial review and wrote the manuscript with contribution from all authors.

580

581 Acknowledgements:

582 We thank staff from Scottish Water, United Utilities, Yorkshire Water, Irish Water and Dŵr Cymru
583 Welsh Water for their informative discussions and comments on early drafts of this manuscript.
584 Discussions with Nadeem Shah and Tom Nisbet regarding work being undertaken by Forest Research
585 are also gratefully acknowledged, as are the comments from 2 reviewers of an earlier draft of this
586 manuscript. The artwork in Figure 2 is by Andy Sier (UKCEH). This work was funded by a NERC
587 Environmental Risks to Infrastructure Innovation Programme grant NE/R009198/1, UKRI SPF UK
588 Climate Resilience programme – Project no. NE/S016937/2, NERC LTSM LOCATE (Land Ocean Carbon
589 Transfer, NE/N018087/1) and Scottish Water.

590

591

592 **References**

- 593 Alderson, D. M., Evans, M. G., Shuttleworth, E. L., Pilkington, M., Spencer, T., Walker, J., and Allott, T.
594 E. H.: Trajectories of ecosystem change in restored blanket peatlands, *Sci. Total Environ.*, 665, 785-
595 796, 10.1016/j.scitotenv.2019.02.095, 2019.
- 596 Andersen, R., Farrell, C., Graf, M., Muller, F., Calvar, E., Frankard, P., Caporn, S., and Anderson, P.: An
597 overview of the progress and challenges of peatland restoration in Western Europe, *Restoration*
598 *Ecology*, 25, 271-282, <https://doi.org/10.1111/rec.12415>, 2017.
- 599 Armstrong, A., Holden, J., Luxton, K., and Quinton, J. N.: Multi-scale relationship between peatland
600 vegetation type and dissolved organic carbon concentration, *Ecological Engineering*, 47, 182-188,
601 10.1016/j.ecoleng.2012.06.027, 2012.
- 602 Armstrong, A., Holden, J., Kay, P., Francis, B., Foulger, M., Gledhill, S., McDonald, A. T., and Walker,
603 A.: The impact of peatland drain-blocking on dissolved organic carbon loss and discolouration of
604 water; results from a national survey, *J. Hydrol.*, 381, 112-120, 10.1016/j.jhydrol.2009.11.031, 2010.
- 605 Berggren, M. and del Giorgio, P. A.: Distinct patterns of microbial metabolism associated to riverine
606 dissolved organic carbon of different source and quality, *J. Geophys. Res.-Biogeosci.*, 120, 989-999,
607 10.1002/2015jg002963, 2015.
- 608 Berggren, M., Klaus, M., Selvam, B. P., Strom, L., Laudon, H., Jansson, M., and Karlsson, J.: Quality
609 transformation of dissolved organic carbon during water transit through lakes: contrasting controls
610 by photochemical and biological processes, *Biogeosciences*, 15, 457-470, 10.5194/bg-15-457-2018,
611 2018.
- 612 Birk, S., Chapman, D., Carvalho, L., Spears, B. M., Andersen, H. E., Argillier, C., Auer, S., Baattrup-
613 Pedersen, A., Banin, L., Beklioğlu, M., Bondar-Kunze, E., Borja, A., Branco, P., Bucak, T., Buijse, A. D.,
614 Cardoso, A. C., Couture, R.-M., Cremona, F., de Zwart, D., Feld, C. K., Ferreira, M. T., Feuchtmayr, H.,
615 Gessner, M. O., Gieswein, A., Globevnik, L., Graeber, D., Graf, W., Gutiérrez-Cánovas, C., Hanganu, J.,
616 Işkın, U., Järvinen, M., Jeppesen, E., Kotamäki, N., Kuijper, M., Lemm, J. U., Lu, S., Solheim, A. L.,
617 Mischke, U., Moe, S. J., Nöges, P., Nöges, T., Ormerod, S. J., Panagopoulos, Y., Phillips, G., Posthuma,
618 L., Pouso, S., Prudhomme, C., Rankinen, K., Rasmussen, J. J., Richardson, J., Sagouis, A., Santos, J. M.,
619 Schäfer, R. B., Schinegger, R., Schmutz, S., Schneider, S. C., Schülting, L., Segurado, P., Stefanidis, K.,
620 Sures, B., Thackeray, S. J., Turunen, J., Uyarra, M. C., Venohr, M., von der Ohe, P. C., Willby, N., and
621 Hering, D.: Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems,
622 *Nature Ecology & Evolution*, 4, 1060-1068, 10.1038/s41559-020-1216-4, 2020.
- 623 Bond, T., Huang, J., Templeton, M. R., and Graham, N.: Occurrence and control of nitrogenous
624 disinfection by-products in drinking water - A review, *Water Res.*, 45, 4341-4354,
625 10.1016/j.watres.2011.05.034, 2011.
- 626 Brown, L. E., Holden, J., Palmer, S. M., Johnston, K., Ramchunder, S. J., and Grayson, R.: Effects of fire
627 on the hydrology, biogeochemistry, and ecology of peatland river systems, *Freshw. Sci.*, 34, 1406-
628 1425, 10.1086/683426, 2015.
- 629 Chapman, P. J., Moody, C. S., Grayson, R., and Palmer, S. M.: Factors controlling water colour on the
630 North York Moors (Part 1), University of Leeds, 2017.
- 631 Chapman, P. J., McDonald, A. T., Tyson, R., Palmer, S. M., Mitchell, G., and Irvine, B.: Changes in
632 water colour between 1986 and 2006 in the headwaters of the River Nidd, Yorkshire, UK,
633 *Biogeochemistry*, 101, 281-294, 10.1007/s10533-010-9474-x, 2010.
- 634 Chimner, R. A., Cooper, D. J., Wurster, F. C., and Rochefort, L.: An overview of peatland restoration in
635 North America: where are we after 25 years?, *Restoration Ecology*, 25, 283-292,
636 <https://doi.org/10.1111/rec.12434>, 2017.
- 637 Clark, J. M., Ashley, D., Wagner, M., Chapman, P. J., Lane, S. N., Evans, C. D., and Heathwaite, A. L.:
638 Increased temperature sensitivity of net DOC production from ombrotrophic peat due to water table
639 draw-down, *Global Change Biology*, 15, 794-807, 10.1111/j.1365-2486.2008.01683.x, 2009.

640 Clay, G. D., Worrall, F., and Aebischer, N. J.: Does prescribed burning on peat soils influence DOC
641 concentrations in soil and runoff waters? Results from a 10 year chronosequence, *J. Hydrol.*, 448,
642 139-148, 10.1016/j.jhydrol.2012.04.048, 2012.

643 Clay, G. D., Worrall, F., and Fraser, E. D. G.: Effects of managed burning upon dissolved organic
644 carbon (DOC) in soil water and runoff water following a managed burn of a UK blanket bog, *J.*
645 *Hydrol.*, 367, 41-51, 10.1016/j.jhydrol.2008.12.022, 2009.

646 Clutterbuck, B. and Yallop, A. R.: Land management as a factor controlling dissolved organic carbon
647 release from upland peat soils 2 Changes in DOC productivity over four decades, *Sci. Total Environ.*,
648 408, 6179-6191, 10.1016/j.scitotenv.2010.08.038, 2010.

649 Cummins, T. and Farrell, E. P.: Biogeochemical impacts of clearfelling and reforestation on blanket-
650 peatland streams - II. major ions and dissolved organic carbon, *For. Ecol. Manage.*, 180, 557-570,
651 10.1016/s0378-1127(02)00649-7, 2003.

652 Davies, G. M., Kettridge, N., Stoof, C. R., Gray, A., Ascoli, D., Fernandes, P. M., Marrs, R., Allen, K. A.,
653 Doerr, S. H., Clay, G. D., McMorrow, J., and Vandvik, V.: The role of fire in UK peatland and moorland
654 management: the need for informed, unbiased debate, *Philos. Trans. R. Soc. B-Biol. Sci.*, 371,
655 10.1098/rstb.2015.0342, 2016.

656 de Wit, H. A., Stoddard, J. L., Monteith, D. T., Sample, J. E., Austnes, K., Couture, S., Fölster, J.,
657 Higgins, S. N., Houle, D., Hruška, J., Krám, P., Kopáček, J., Paterson, A. M., Valinia, S., Van Dam, H.,
658 Vuorenmaa, J., and Evans, C. D.: Cleaner air reveals growing influence of climate on dissolved organic
659 carbon trends in northern headwaters, *Environmental Research Letters*, 16, 104009, 10.1088/1748-
660 9326/ac2526, 2021.

661 Ding, S. and Chu, W.: Recent advances in the analysis of nitrogenous disinfection by-products, *Trends*
662 *in Environmental Analytical Chemistry*, 14, 19-27, <https://doi.org/10.1016/j.teac.2017.04.001>, 2017.

663 Einola, E., Rantakari, M., Kankaala, P., Kortelainen, P., Ojala, A., Pajunen, H., Makela, S., and Arvola,
664 L.: Carbon pools and fluxes in a chain of five boreal lakes: A dry and wet year comparison, *J.*
665 *Geophys. Res.-Biogeosci.*, 116, 10.1029/2010jg001636, 2011.

666 Evans, C. D., Futter, M. N., Moldan, F., Valinia, S., Frogbrook, Z., and Kothawala, D. N.: Variability in
667 organic carbon reactivity across lake residence time and trophic gradients, *Nature Geoscience*, 10,
668 832-+, 10.1038/ngeo3051, 2017a.

669 Evans, C. D., Malcolm, I. A., Shilland, E. M., Rose, N. L., Turner, S. D., Crilly, A., Norris, D., Granath, G.,
670 and Monteith, D. T.: Sustained Biogeochemical Impacts of Wildfire in a Mountain Lake Catchment,
671 *Ecosystems*, 20, 813-829, 10.1007/s10021-016-0064-1, 2017b.

672 Evans, C. D., Peacock, M., Green, S. M., Holden, J., Chapman, P. J., Lebron, I., Callaghan, N., Grayson,
673 R., and Baird, A.: The impact of ditch blocking on fluvial carbon export from a UK blanket bog,
674 *Hydrological Processes*, 32, 2141-2154, 2018.

675 Evans, C. D., Jones, T. G., Burden, A., Ostle, N., Zielinski, P., Cooper, M. D. A., Peacock, M., Clark, J.
676 M., Oulehle, F., Cooper, D., and Freeman, C.: Acidity controls on dissolved organic carbon mobility in
677 organic soils, *Global Change Biology*, 18, 3317-3331, 10.1111/j.1365-2486.2012.02794.x, 2012.

678 Evans, C. D., Page, S. E., Jones, T., Moore, S., Gauci, V., Laiho, R., Hruska, J., Allott, T. E. H., Billett, M.
679 F., Tipping, E., Freeman, C., and Garnett, M. H.: Contrasting vulnerability of drained tropical and
680 high-latitude peatlands to fluvial loss of stored carbon, *Global Biogeochemical Cycles*, 28, 1215-1234,
681 10.1002/2013gb004782, 2014.

682 Feuchtmayr, H., Pottinger, T. G., Moore, A., De Ville, M. M., Caillouet, L., Carter, H. T., Pereira, M. G.,
683 and Maberly, S. C.: Effects of brownification and warming on algal blooms, metabolism and higher
684 trophic levels in productive shallow lake mesocosms, *Sci. Total Environ.*, 678, 227-238,
685 <https://doi.org/10.1016/j.scitotenv.2019.04.105>, 2019.

686 Flynn, R., Mackin, F., McVeigh, C., and Renou-Wilson, F.: Impacts of a mature forestry plantation on
687 blanket peatland runoff regime and water quality, *Hydrological Processes*, 36, e14494,
688 <https://doi.org/10.1002/hyp.14494>, 2022.

689 Forestry Commission: The UK Forestry Standard, Edinburgh2017.

690 Freeman, C., Evans, C. D., Monteith, D. T., Reynolds, B., and Fenner, N.: Export of organic carbon
691 from peat soils, *Nature*, 412, 785-785, 10.1038/35090628, 2001.

692 Gaffney, P.: The effects of bog restoration in formerly afforested peatlands on water quality and
693 aquatic carbon fluxes, Environmental Research Institute, University of the Highlands and Islands,
694 2017.

695 Gaffney, P. P. J., Hancock, M. H., Taggart, M. A., and Andersen, R.: Measuring restoration progress
696 using pore- and surface-water chemistry across a chronosequence of formerly afforested blanket
697 bogs, *J. Environ. Manage.*, 219, 239-251, <https://doi.org/10.1016/j.jenvman.2018.04.106>, 2018.

698 Gaffney, P. P. J., Hancock, M. H., Taggart, M. A., and Andersen, R.: Restoration of afforested
699 peatland: Immediate effects on aquatic carbon loss, *Sci. Total Environ.*, 742, 140594,
700 <https://doi.org/10.1016/j.scitotenv.2020.140594>, 2020.

701 Gibson, H. S., Worrall, F., Burt, T. P., and Adamson, J. K.: DOC budgets of drained peat catchments:
702 implications for DOC production in peat soils, *Hydrological Processes*, 23, 1901-1911,
703 10.1002/hyp.7296, 2009.

704 Glatzel, S., Kalbitz, K., Dalva, M., and Moore, T.: Dissolved organic matter properties and their
705 relationship to carbon dioxide efflux from restored peat bogs, *Geoderma*, 113, 397-411,
706 [https://doi.org/10.1016/S0016-7061\(02\)00372-5](https://doi.org/10.1016/S0016-7061(02)00372-5), 2003.

707 Glenk, K. and Martin-Ortega, J.: The economics of peatland restoration, *Journal of Environmental*
708 *Economics and Policy*, 7, 345-362, 10.1080/21606544.2018.1434562, 2018.

709 Gough, R., Holliman, P. J., Fenner, N., Peacock, M., and Freeman, C.: Influence of Water Table Depth
710 on Pore Water Chemistry and Trihalomethane Formation Potential in Peatlands, *Water Environ.*
711 *Res.*, 88, 107-117, 10.2175/106143015x14362865227878, 2016.

712 Gough, R., Holliman, P. J., Willis, N., Jones, T. G., and Freeman, C.: Influence of habitat on the
713 quantity and composition of leachable carbon in the O2 horizon: Potential implications for potable
714 water treatment, *Lake Reserv. Manag.*, 28, 282-292, 10.1080/07438141.2012.741187, 2012.

715 Grau-Andres, R., Davies, G. M., Waldron, S., Scott, E. M., and Gray, A.: Increased fire severity alters
716 initial vegetation regeneration across Calluna-dominated ecosystems, *J. Environ. Manage.*, 231,
717 1004-1011, 10.1016/j.jenvman.2018.10.113, 2019.

718 Haapalehto, T., Kotiaho, J. S., Matilainen, R., and Tahvanainen, T.: The effects of long-term drainage
719 and subsequent restoration on water table level and pore water chemistry in boreal peatlands, *J.*
720 *Hydrol.*, 519, 1493-1505, 10.1016/j.jhydrol.2014.09.013, 2014.

721 Harper, A. R., Doerr, S. H., Santin, C., Froyd, C. A., and Sinnadurai, P.: Prescribed fire and its impacts
722 on ecosystem services in the UK, *Sci. Total Environ.*, 624, 691-703, 10.1016/j.scitotenv.2017.12.161,
723 2018.

724 Harriman, R., Watt, A. W., Christie, A. E. G., Collen, P., Moore, D. W., McCartney, A. G., Taylor, E. M.,
725 and Watson, J.: Interpretation of trends in acidic deposition and surface water chemistry in Scotland
726 during the past three decades, *Hydrol. Earth Syst. Sci.*, 5, 407-420, 10.5194/hess-5-407-2001, 2001.

727 HM Government: Net Zero Strategy: Build Back Greener, 2021.

728 Holden, J., Chapman, P. J., Palmer, S. M., Kay, P., and Grayson, R.: The impacts of prescribed
729 moorland burning on water colour and dissolved organic carbon: A critical synthesis, *J. Environ.*
730 *Manage.*, 101, 92-103, 10.1016/j.jenvman.2012.02.002, 2012.

731 Holden, J., Kirkby, M. J., Lane, S. N., Milledge, D. G., Brookes, C. J., Holden, V., and McDonald, A. T.:
732 Overland flow velocity and roughness properties in peatlands, *Water Resour. Res.*, 44,
733 10.1029/2007wr006052, 2008.

734 Holden, J., Green, S. M., Baird, A. J., Grayson, R. P., Dooling, G. P., Chapman, P. J., Evans, C. D.,
735 Peacock, M., and Swindles, G.: The impact of ditch blocking on the hydrological functioning of
736 blanket peatlands, *Hydrological Processes*, 31, 525-539, 10.1002/hyp.11031, 2017.

737 Holl, B. S., Fiedler, S., Jungkunst, H. F., Kalbitz, K., Freibauer, A., Drosler, M., and Stahr, K.:
738 Characteristics of dissolved organic matter following 20 years of peatland restoration, *Sci. Total*
739 *Environ.*, 408, 78-83, 10.1016/j.scitotenv.2009.08.046, 2009.

740 Howson, T., Chapman, P. J., Shah, N., Anderson, R., and Holden, J.: A comparison of porewater
741 chemistry between intact, afforested and restored raised and blanket bogs, *Sci. Total Environ.*, 766,
742 144496, <https://doi.org/10.1016/j.scitotenv.2020.144496>, 2021.

743 Hudson, J. A., Gilman, K., and Calder, I. R.: Land use and water issues in the uplands with reference
744 to the Plynlimon study, *Hydrol. Earth Syst. Sci.*, 1, 389-397, 10.5194/hess-1-389-1997, 1997.

745 Imai, A., Fukushima, T., Matsushige, K., Kim, Y.-H., and Choi, K.: Characterization of dissolved organic
746 matter in effluents from wastewater treatment plants, *Water Res.*, 36, 859-870,
747 [https://doi.org/10.1016/S0043-1354\(01\)00283-4](https://doi.org/10.1016/S0043-1354(01)00283-4), 2002.

748 Jandl, R., Lindner, M., Vesterdal, L., Bauwens, B., Baritz, R., Hagedorn, F., Johnson, D. W., Minkinen,
749 K., and Byrne, K. A.: How strongly can forest management influence soil carbon sequestration?,
750 *Geoderma*, 137, 253-268, <https://doi.org/10.1016/j.geoderma.2006.09.003>, 2007.

751 Jones, T. G., Evans, C. D., and Freeman, C.: The greenhouse gas (GHG) emissions associated with
752 aquatic carbon removal during drinking water treatment, *Aquat. Sci.*, 78, 561-572, 10.1007/s00027-
753 015-0458-8, 2016.

754 Koehler, B., von Wachenfeldt, E., Kothawala, D., and Tranvik, L. J.: Reactivity continuum of dissolved
755 organic carbon decomposition in lake water, *J. Geophys. Res.-Biogeosci.*, 117,
756 10.1029/2011jg001793, 2012.

757 Koehler, B., Landelius, T., Weyhenmeyer, G. A., Machida, N., and Tranvik, L. J.: Sunlight-induced
758 carbon dioxide emissions from inland waters, *Global Biogeochemical Cycles*, 28, 696-711,
759 10.1002/2014gb004850, 2014.

760 Lavonen, E. E., Gonsior, M., Tranvik, L. J., Schmitt-Kopplin, P., and Kohler, S. J.: Selective Chlorination
761 of Natural Organic Matter: Identification of Previously Unknown Disinfection Byproducts,
762 *Environmental Science & Technology*, 47, 2264-2271, 10.1021/es304669p, 2013.

763 Lundin, L., Nilsson, T., Jordan, S., Lode, E., and Strömberg, M.: Impacts of rewetting on peat,
764 hydrology and water chemical composition over 15 years in two finished peat extraction areas in
765 Sweden, *Wetlands Ecology and Management*, 25, 405-419, 10.1007/s11273-016-9524-9, 2017.

766 Matilainen, A., Vepsäläinen, M., and Sillanpää, M.: Natural organic matter removal by coagulation
767 during drinking water treatment: A review, *Advances in Colloid and Interface Science*, 159, 189-197,
768 <https://doi.org/10.1016/j.cis.2010.06.007>, 2010.

769 McKnight, D. M., Bencala, K. E., Zellweger, G. W., Aiken, G. R., Feder, G. L., and Thorn, K. A.: Sorption
770 of dissolved organic carbon by hydrous aluminum and iron oxides occurring at the confluence of
771 Deer Creek with the Snake River, Summit County, Colorado, *Environmental Science & Technology*,
772 26, 1388-1396, 10.1021/es00031a017, 1992.

773 Menberu, M., Mattila, H., Tahvanainen, T., Kotiaho, J. S., Hokkanen, R., Klove, B., and Ronkanen, A.:
774 Changes in pore water quality after peatland restoration: assessment of a large scale replicated
775 before-after-control-impact study in Finland, *Water Resour. Res.*, 53, 8327-8343, 2017.

776 Monteith, D., Pickard, A. E., Spears, B. M., and Feuchtmayr, H.: An introduction to the FREEDOM-
777 BCCR project: FREEDOM-BCCR briefing note 1 to the water industry, 2021.

778 Monteith, D. T., Henrys, P. A., Hruška, J., de Wit, H. A., Krám, P., Moldan, F., Posch, M., Räike, A.,
779 Stoddard, J. L., Shilland, E. M., Pereira, M. G., and Evans, C. D.: Long-term rise in riverine dissolved
780 organic carbon concentration is predicted by electrolyte solubility theory, *Science Advances*, 9,
781 eade3491, doi:10.1126/sciadv.ade3491, 2023.

782 Monteith, D. T., Stoddard, J. L., Evans, C. D., de Wit, H. A., Forsius, M., Hogasen, T., Wilander, A.,
783 Skjelkvale, B. L., Jeffries, D. S., Vuorenmaa, J., Keller, B., Kopacek, J., and Vesely, J.: Dissolved organic
784 carbon trends resulting from changes in atmospheric deposition chemistry, *Nature*, 450, 537-U539,
785 10.1038/nature06316, 2007.

786 Moody, C. S., Worrall, F., Evans, C. D., and Jones, T. G.: The rate of loss of dissolved organic carbon
787 (DOC) through a catchment, *J. Hydrol.*, 492, 139-150, 10.1016/j.jhydrol.2013.03.016, 2013.

788 Muller, F. L. L. and Tankere-Muller, S. P. C.: Seasonal variations in surface water chemistry at
789 disturbed and pristine peatland sites in the Flow Country of northern Scotland, *Sci. Total Environ.*,
790 435, 351-362, 10.1016/j.scitotenv.2012.06.048, 2012.

791 Muller, F. L. L., Chang, K.-C., Lee, C.-L., and Chapman, S. J.: Effects of temperature, rainfall and
792 conifer felling practices on the surface water chemistry of northern peatlands, *Biogeochemistry*, 126,
793 343-362, 10.1007/s10533-015-0162-8, 2015.

794 Naden, P. S. and McDonald, A. T.: Statistical modeling of water color in the uplands - the Upper Nidd
795 catchment 1979-1987, *Environmental Pollution*, 60, 141-163, 10.1016/0269-7491(89)90224-8, 1989.

796 Neal, C.: Nutrient concentrations and fluxes for podzolic and gley soils at Plynlimon, mid-Wales:
797 implications for modelling inorganic nitrogen and phosphorus in upland UK environments, *Hydrol.*
798 *Earth Syst. Sci.*, 6, 403-420, 10.5194/hess-6-403-2002, 2002.

799 O'Brien, H. E., Labadz, J. C., Butcher, D. P., Billett, M. F., and Midgley, N. G.: Impact of catchment
800 management upon dissolved organic carbon and stream flows in the Peak District, Derbyshire, UK.,
801 10th BHS National Hydrology Symposium, Exeter,

802 Parry, L. E., Chapman, P. J., Palmer, S. M., Wallage, Z. E., Wynne, H., and Holden, J.: The influence of
803 slope and peatland vegetation type on riverine dissolved organic carbon and water colour at
804 different scales, *Sci. Total Environ.*, 527, 530-539, 10.1016/j.scitotenv.2015.03.036, 2015.

805 Patterson, G. and Anderson, R.: *Forests and Peatland Habitats*, Forestry Commission, Edinburgh,
806 2000.

807 Peacock, M., Jones, T. G., Futter, M. N., Freeman, C., Gough, R., Baird, A. J., Green, S. M., Chapman,
808 P. J., Holden, J., and Evans, C. D.: Peatland ditch blocking has no effect on dissolved organic matter
809 (DOM) quality, *Hydrological Processes*, 32, 3891-3906, <https://doi.org/10.1002/hyp.13297>, 2018.

810 Peter, S., Agstam, O., and Sobek, S.: Widespread release of dissolved organic carbon from anoxic
811 boreal lake sediments, *Inland Waters*, 7, 151-163, 10.1080/20442041.2017.1300226, 2017.

812 Pickard, A. E., Branagan, M., Billett, M. F., Andersen, R., and Dinsmore, K. J.: Effects of peatland
813 management on aquatic carbon concentrations and fluxes, *Biogeosciences*, 19, 1321-1334,
814 10.5194/bg-19-1321-2022, 2022.

815 Pilkington, M., Walker, J., Maskill, R., Allott, T. E. H., and Evans, M.: *Restoration of blanket bogs;
816 flood risk reduction and other ecosystem benefits*, Edale, 2015.

817 Prest, E. I., Hammes, F., van Loosdrecht, M. C. M., and Vrouwenvelder, J. S.: Biological Stability of
818 Drinking Water: Controlling Factors, Methods, and Challenges, *Frontiers in Microbiology*, 7,
819 10.3389/fmicb.2016.00045, 2016.

820 Qassim, S. M., Dixon, S. D., Rowson, J. G., Worrall, F., Evans, M. G., and Bonn, A.: A 5-year study of
821 the impact of peatland revegetation upon DOC concentrations, *J. Hydrol.*, 519, 3578-3590,
822 10.1016/j.jhydrol.2014.11.014, 2014.

823 Ramchunder, S. J., Brown, L. E., and Holden, J.: Rotational vegetation burning effects on peatland
824 stream ecosystems, *J. Appl. Ecol.*, 50, 636-648, 10.1111/1365-2664.12082, 2013.

825 Rask, M., Nyberg, K., Markkanen, S.-L., and Ojala, A.: Forestry in catchments: effects on water
826 quality, plankton, zoobenthos and fish in small lakes, *Boreal Environment Research*, 3, 75-86, 1998.

827 Ritson, J. P., Graham, N. J. D., Templeton, M. R., Clark, J. M., Gough, R., and Freeman, C.: The impact
828 of climate change on the treatability of dissolved organic matter (DOM) in upland water supplies: A
829 UK perspective, *Sci. Total Environ.*, 473-474, 714-730,
830 <https://doi.org/10.1016/j.scitotenv.2013.12.095>, 2014.

831 Ritson, J. P., Bell, M., Brazier, R. E., Grand-Clement, E., Graham, N. J. D., Freeman, C., Smith, D.,
832 Templeton, M. R., and Clark, J. M.: Managing peatland vegetation for drinking water treatment, *Sci*
833 *Rep*, 6, 10.1038/srep36751, 2016.

834 Robson, A. J. and Neal, C.: Water quality trends at an upland site in Wales, UK, 1983-1993,
835 *Hydrological Processes*, 10, 183-203, 10.1002/(sici)1099-1085(199602)10:2<183::Aid-
836 hyp356>3.0.Co;2-8, 1996.

837 Schelker, J., Eklof, K., Bishop, K., and Laudon, H.: Effects of forestry operations on dissolved organic
838 carbon concentrations and export in boreal first-order streams, *J. Geophys. Res.-Biogeosci.*, 117,
839 10.1029/2011jg001827, 2012.

840 Flanders Moss Peatland Restoration: <https://www.forestry.gov.uk/fr/bee-atugla>, last access:
841 12/04/2018.

842 Shah, N. and Nisbet, T.: The effects of forest clearance for peatland restoration on water quality, *Sci.*
843 *Total Environ.*, 693, 133617, 10.1016/j.scitotenv.2019.133617, 2019.

844 Shah, N. W., Nisbet, T. R., and Broadmeadow, S. B.: The impacts of conifer afforestation and climate
845 on water quality and freshwater ecology in a sensitive peaty catchment: A 25 year study in the upper
846 River Halladale in North Scotland, *For. Ecol. Manage.*, 502, 119616,
847 <https://doi.org/10.1016/j.foreco.2021.119616>, 2021.

848 Skerlep, M., Steiner, E., Axelsson, A. L., and Kritzberg, E. S.: Afforestation driving long-term surface
849 water browning, *Global Change Biology*, 26, 1390-1399, 10.1111/gcb.14891, 2019.

850 Spears, B. M. and May, L.: Long-term homeostasis of filterable un-reactive phosphorus in a shallow
851 eutrophic lake following a significant reduction in catchment load, *Geoderma*, 257-258, 78-85,
852 <https://doi.org/10.1016/j.geoderma.2015.01.005>, 2015.

853 Spears, B. M., Mackay, E. B., Yasserli, S., Gunn, I. D. M., Waters, K. E., Andrews, C., Cole, S., De Ville,
854 M., Kelly, A., Meis, S., Moore, A. L., Nürnberg, G. K., van Oosterhout, F., Pitt, J.-A., Madgwick, G.,
855 Woods, H. J., and Lüring, M.: A meta-analysis of water quality and aquatic macrophyte responses
856 in 18 lakes treated with lanthanum modified bentonite (Phoslock®), *Water Res.*, 97, 111-121,
857 <https://doi.org/10.1016/j.watres.2015.08.020>, 2016.

858 Stets, E. G., Striegl, R. G., and Aiken, G. R.: Dissolved organic carbon export and internal cycling in
859 small, headwater lakes, *Global Biogeochemical Cycles*, 24, 10.1029/2010gb003815, 2010.

860 Stimson, A. G., Allott, T. E. H., Boulton, S., and Evans, M. G.: Fluvial organic carbon composition and
861 concentration variability within a peatland catchment-Implications for carbon cycling and water
862 treatment, *Hydrological Processes*, 31, 4183-4194, 10.1002/hyp.11352, 2017.

863 Strack, M., Zuback, Y., McCarter, C., and Price, J.: Changes in dissolved organic carbon quality in soils
864 and discharge 10 years after peatland restoration, *J. Hydrol.*, 527, 345-354,
865 10.1016/j.jhydrol.2015.04.061, 2015.

866 Stretton, C.: Water supply and forestry - a conflict of interests: Cray reservoir, a case study, *J Inst*
867 *Water Eng Scient*, 38, 323 - 330, 1984.

868 Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay,
869 K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M.,
870 McCallister, S. L., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W.
871 H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A.
872 M., von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon
873 cycling and climate, *Limnology and Oceanography*, 54, 2298-2314,
874 10.4319/lo.2009.54.6_part_2.2298, 2009.

875 Turner, E. K., Worrall, F., and Burt, T. P.: The effect of drain blocking on the dissolved organic carbon
876 (DOC) budget of an upland peat catchment in the UK, *J. Hydrol.*, 479, 169-179,
877 10.1016/j.jhydrol.2012.11.059, 2013.

878 Urbanova, Z., Pícek, T., and Barta, J.: Effect of peat re-wetting on carbon and nutrient fluxes,
879 greenhouse gas production and diversity of methanogenic archaeal community, *Ecological*
880 *Engineering*, 37, 1017-1026, 10.1016/j.ecoleng.2010.07.012, 2011.

881 von Wachenfeldt, E. and Tranvik, L. J.: Sedimentation in boreal lakes - The role of flocculation of
882 allochthonous dissolved organic matter in the water column, *Ecosystems*, 11, 803-814,
883 10.1007/s10021-008-9162-z, 2008.

884 Wallage, Z. E., Holden, J., and McDonald, A. T.: Drain blocking: An effective treatment for reducing
885 dissolved organic carbon loss and water discolouration in a drained peatland, *Sci. Total Environ.*,
886 367, 811-821, 10.1016/j.scitotenv.2006.02.010, 2006.

887 Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R., and Mopper, K.: Evaluation of
888 specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of
889 dissolved organic carbon, *Environmental Science & Technology*, 37, 4702-4708, 10.1021/es030360x,
890 2003.

891 Williamson, J. L., Tye, A., Lapworth, D. J., Monteith, D., Sanders, R., Mayor, D. J., Barry, C., Bowes, M.,
892 Bowes, M., Burden, A., Callaghan, N., Farr, G., Felgate, S., Fitch, A., Gibb, S., Gilbert, P., Hargreaves,

893 G., Keenan, P., Kitidis, V., Juergens, M., Martin, A., Mounteney, I., Nightingale, P. D., Pereira, M. G.,
894 Olszewska, J., Pickard, A., Rees, A. P., Spears, B., Stinchcombe, M., White, D., Williams, P., Worrall, F.,
895 and Evans, C.: Landscape controls on riverine export of dissolved organic carbon from Great Britain,
896 *Biogeochemistry*, 10.1007/s10533-021-00762-2, 2021.

897 Wilson, L., Wilson, J., Holden, J., Johnstone, I., Armstrong, A., and Morris, M.: Ditch blocking, water
898 chemistry and organic carbon flux: Evidence that blanket bog restoration reduces erosion and fluvial
899 carbon loss, *Sci. Total Environ.*, 409, 2010-2018, 10.1016/j.scitotenv.2011.02.036, 2011.

900 Worrall, F., Armstrong, A., and Adamson, J. K.: The effects of burning and sheep-grazing on water
901 table depth and soil water quality in a upland peat, *J. Hydrol.*, 339, 1-14,
902 10.1016/j.jhydrol.2006.12.025, 2007a.

903 Worrall, F., Armstrong, A., and Holden, J.: Short-term impact of peat drain-blocking on water colour,
904 dissolved organic carbon concentration, and water table depth, *J. Hydrol.*, 337, 315-325,
905 10.1016/j.jhydrol.2007.01.046, 2007b.

906 Worrall, F., Rowson, J., and Dixon, S.: Effects of managed burning in comparison with vegetation
907 cutting on dissolved organic carbon concentrations in peat soils, *Hydrological Processes*, 27, 3994-
908 4003, 10.1002/hyp.9474, 2013.

909 Worrall, F., Clay, G. D., Marrs, R., and Reed, M.: Impacts of burning management on peatlands, 2010.

910 Worrall, F., Harriman, R., Evans, C. D., Watts, C. D., Adamson, J., Neal, C., Tipping, E., Burt, T., Grieve,
911 I., Monteith, D., Naden, P. S., Nisbet, T., Reynolds, B., and Stevens, P.: Trends in dissolved organic
912 carbon in UK rivers and lakes, *Biogeochemistry*, 70, 369-402, 10.1007/s10533-004-8131-7, 2004.

913 Xu, J., Morris, P. J., Liu, J., and Holden, J.: Hotspots of peatland-derived potable water use identified
914 by global analysis, *Nature Sustainability*, 1, 246-253, 10.1038/s41893-018-0064-6, 2018.

915 Yallop, A. R., Clutterbuck, B., and Thacker, J.: Increases in humic dissolved organic carbon export
916 from upland peat catchments: the role of temperature, declining sulphur deposition and changes in
917 land management, *Clim. Res.*, 45, 43-56, 10.3354/cr00884, 2010.

918 Zheng, Y., Waldron, S., and Flowers, H.: Fluvial dissolved organic carbon composition varies spatially
919 and seasonally in a small catchment draining a wind farm and felled forestry, *Sci. Total Environ.*, 626,
920 785-794, 10.1016/j.scitotenv.2018.01.001, 2018.

921