

Trawling effects on biogeochemical processes are mediated by fauna in high energy biogenic reef-inhabited coastal sediments

Justin C. Tiano^{1,2}, Jochen Depestele³, Gert Van Hoey^{2,3}, João Fernandes^{4,5}, Pieter van Rijswijk¹, Karline Soetaert^{1,2}

5 ¹Royal Netherlands Institute for Sea Research (NIOZ) and Utrecht University, Department of 5 Estuarine and Delta Systems, PO box 140, 4400 AC, Yerseke, The Netherlands

²Ghent University, Department of Biology, Marine Biology Section, Krijgslaan 281/S8, 9000 Ghent, 9 Belgium

³Research Institute for Agriculture, Fisheries and Food (ILVO), Fisheries Research Group, Ankerstraat 1, 8400 Oostende, Belgium

10 ⁴CETEMARES, Instituto Politécnico de Leiria, Avenida do Porto de Pesca 30, 2520-620 Peniche, Portugal

⁵Lab. MAREFOZ, Incubadora de Empresas da Figueira da Foz, R. Acácias lote 40A, 3090-380 Figueira da Foz, Portugal

Correspondence to: Justin C. Tiano (Justin.Tiano@nioz.nl)

Abstract. Dynamic, sandy environments are generally less vulnerable to mechanical stress compared to silty, low energy habitats. Biogenic reef communities, however, may provide an exception to this. This study explores the physical, biological and biogeochemical effects of bottom trawl fishing on a coastal ecosystem dominated by the tube building polychaete, *Lanice conchilega*. Two specific gears types, both used to exploit North Sea sole (*Solea solea*), were compared: electric pulse trawls and tickler chain rigged beam trawls. We detected a ~1 cm bathymetric deepening after trawling associated with significant losses in benthic chlorophyll *a* caused from both fishing gears. Tickler chain trawls significantly reduced sediment oxygen consumption (57%), total organic matter mineralization (56%), denitrification (61%), nitrification (60%), and total benthos densities (52%) while pulse trawls had no statistically significant impact on these parameters. Before trawling, significant relationships could be found between *L. conchilega* and very fine sand fractions, oxygen and nitrate fluxes, macrobenthos densities and species richness, however, the trawl disturbances from both gears disrupted these connections. Our results suggest a reduced average effect for pulse compared to tickler chain beam trawls for several ecological and biogeochemical characteristics though their impact was still significant for *L. conchilega* and associated species. This study also suggests that faunal-mediated ecosystem functions in *L. conchilega* dominated habitats may be sensitive to relatively shallow sediment penetration from trawl gears and should be considered when assessing habitat vulnerability.

1 Introduction

Marine sediment ecosystems are important drivers for benthic pelagic coupling (Mermillod-Blondin and Rosenberg, 2006) and carbon mineralization (Seiter et al., 2005). Biogenic reef habitats in particular, such as those created by the tube-building polychaete, *Lanice conchilega*, are able to capture substantial amounts of organic carbon compared to adjacent sediments (De

Smet et al., 2016a) resulting in dense and biologically diverse benthic communities (Van Hoey et al., 2008). This species may be found in various North Sea habitats but generally prefer shallow muddy and fine sands where it may exhibit densities up to 1,000 individuals per m² (Van Hoey et al., 2008). Bottom trawling activities are a common occurrence in North Sea *L. conchilega* dominated habitats (Rabaut, 2009) and can present a considerable threat to most biogenic reef ecosystems (Kaiser et al., 2002; Fariñas-Franco et al., 2018). Biological resilience to bottom trawl impacts have, however, been linked to both *L. conchilega* reefs (Rabaut et al., 2008) and the dynamic sandy habitats they are often found in (Kaiser et al., 2006; Sciberras et al., 2016; Rijnsdorp et al., 2020a). Consequences of trawling on more holistic ecosystem processes such as benthic pelagic coupling, however, have not been thoroughly examined in these systems.

Bottom fishing affects benthic carbon cycling by displacing bottom dwelling organisms (Hiddink et al., 2017; Sciberras et al., 2018), altering sediment structures leading to the impairment of biogeochemical processes (Depestele et al., 2018; Ferguson et al., 2020), and resuspending fresh organic material from the sediment surface, which can result in reduced mineralization of organic carbon in seabed sediments (Tiano et al., 2019; Paradis et al., 2020; Bradshaw et al., 2021; Morys et al., 2021). Conversely, trawling may also increase benthic carbon mineralization through depositional effects or if refractory organic fractions are mixed with and reactivated by fresh organic matter (van de Velde et al., 2018). Trawl-induced mortality to benthic organisms can simplify benthic communities and can lead to a loss in functional diversity (Tillin et al., 2006; Tiano et al., 2020). The removal (significant or partial) of an abundant ecosystem engineer such as *L. conchilega* and associated fauna would likely have major impacts on biogeochemical and community dynamics (Rabaut et al., 2008; Van Hoey et al., 2008; De Smet et al., 2016a; Foshtomi et al., 2018).

The severity of bottom trawl impact depends on the size and type of fishing gear used and speed at which it drags over the seabed (Eigaard et al., 2016; Depestele et al., 2016; Depestele et al., 2018; Rijnsdorp et al., 2020a, 2020b). In the southern North Sea, beam trawls rigged with tickler chains mechanically stimulate flatfish up from the seafloor, facilitating their capture. An alternative fishing gear for flatfish called “pulse trawls” have been used to expose the seabed to electrical fields, causing the immobilization of target fish for oncoming bottom nets (Soetaert et al., 2015). Previous comparisons conducted in the Frisian Front, a relatively homogenous habitat known for high abundances and diversity of benthic organisms (Dauwe et al., 1998; Dewick et al., 2002), have found a reduced impact of pulse compared to tickler rigged trawls on physical (Depestele et al., 2018) and biogeochemical parameters (Tiano et al., 2019) but impacts on benthic communities were similar for both gears (Tiano et al., 2020). Dynamic nearshore habitats, in contrast, may provide a more challenging environment for detecting significant anthropogenic alterations.

Regarding to sensitivity to direct impacts and recovery potential, coarse sediment ecosystems characterized by high levels of natural disturbance typically display more resistance and resilience to bottom fishing compared to muddy and more hydrodynamically static ecosystems (Kaiser et al., 2002; Bolam et al., 2014; Sciberras et al., 2016; Rijnsdorp et al., 2020a). Macrobenthos found in sandy, well-sorted sediments, tend to be less sensitive to trawl disturbance than in muddy poorly sorted habitats (Bolam et al., 2014; van Denderen et al., 2015). Communities dominated by three-dimensional biological features,

65 such biogenic reefs, seem to be an exception to this and can be vulnerable to trawl disturbance regardless of hydrodynamic energy level or substrate type (Kaiser et al., 2002; Grabowski et al., 2014).

The current study uses a combination of water column, sonar, sediment profile imagery, box core sampling, sediment incubation and porewater nutrient extraction techniques to provide a comprehensive assessment of bottom trawling on ecosystem functioning. This integrated data collection allows us to explore the links between pure mechanical trawling effects, 70 disturbance-induced changes to faunal-mediated biogeochemical functions and their collective impact on benthic pelagic coupling. Our main objective was to measure the acute ecosystem effects of electric pulse and tickler chain beam trawls in a high-energy coastal system. Furthermore, the high abundances of *Lanice conchilega* in the region, allowed us to quantify the effect of this species on ecological and biogeochemical parameters, and how this may change due to fishing disturbance.

2 Methods

75 2.1 Study area

This study took place in shallow subtidal waters (8 – 10 m depth), 10 km away from the Netherlands coast close to the Dutch and Belgian EEZ (exclusive economic zone) border (Fig. 1). This region, known as the “Vlakte van de Raan”, is an area known for its subtidal sandbanks, which create a mosaic of habitat areas of varying ecological value (Degraer and Hostens, 2016, Pecceu et al., 2016). The experimental areas were characterized by “fine sand” with an average median grain size of 200.5 μm 80 and an average chlorophyll *a* content (0 – 10 cm depth) of 13.1 $\mu\text{g g}^{-1}$ (Fig. S1 in the Supplement). The exact location for this study within the Vlakte van de Raan (Fig. 1), was chosen due to its relatively low levels of bottom trawl disturbance (no flatfish or shrimp trawl disturbance in one year as verified with vessel monitoring data) and the high macrobenthic densities and biomass (Craeymeersch et al., 2006).

2.2 Experimental trawling design

85 Three experimental treatments were conducted: (1) tickler chain beam trawling (hereafter referred to as “Tickler” treatment or tickler trawls), (2) pulse trawling with and (3) without electricity turned on (hereafter referred to as “Pulse-ON” and “Pulse-OFF” treatment). Each treatment was repeated three times in spatially separated areas (Pulse-ON = A, B, C; Pulse-OFF = D, E, F, Tickler = G, H, I), resulting in a total of nine experimental plots of 50 by 300 m (15,000 m² per plot; Fig. 1). The locations and orientation of the treatment plots were chosen based on their position within a bathymetric depression. Plots and untrawled 90 adjacent sites were designated to represent northwest, central and southeast areas within this depression and were spatially separated by at least 100 m to minimize effects of trawl induced sediment resuspension and deposition. Individual plots were also subdivided by three different sampling stations representing the northwest, central, and southeast locations within the plot. Three additional untrawled stations located adjacent to the treatment plots (AD1, AD2, AD3) were chosen within the experimental area (Fig. 1). Samples taken from AD stations deviated within 80 m of a single sampling point.

95 Pulse trawls (Delmaco; Soetaert et al., 2015) were deployed from the commercial fishing vessel (FV) *TH6 Johanna*
Cornelia (24 m) and tickler trawls from the FV *YE238 Geertruida* (22 m). The vessels were relatively small trawlers (<221
kW engine) and allowed to fish within 12 nm of the coast. Both pulse trawl (~ 1.1 tonne in air) and tickler trawl (~ 2.4 tonne
in air) gears measured 4 m in width and had one trawl shoe at either side of its beam. The fishing vessels made a total of six
trawl passes over each experimental plot at a speed of ~ 5 knots (pulse) and ~ 6 knots (tickler). Experimental fishing with the
100 pulse trawl was performed on 12 June 2018 while the tickler trawl was used on 15 of June 2018 (Table S1 in the Supplement).
All experimental equipment was deployed from the research vessel (RV) *Simon Stevin*.

2.3 Water column turbidity

To measure turbidity during trawling events, the RV deployed an SBE 25plus Sealogger (CTD) with two optical backscatter
point sensors (OBS) while suspended continuously at 5 m depth during experimental trawling activity. For each treatment plot
105 measured (A, B, C, D, E, G, and H), the RV strategically positioned itself against the current and the ongoing trawling ~ 80 m
adjacent to the corresponding plots (due to logistic constraints plots F and I were not measured).

2.4 Bathymetry and backscatter

Bathymetry and backscatter data were collected using a Kongsberg EM2040 single head multi-beam echo sounder (MBES)
mounted on the RV (Depestele et al., 2018). All data were collected with a MBES frequency of 300 kHz, a speed of 8 knots
110 and at orientation parallel to the longest side of the experimental plots to avoid backscatter bias from the survey line directions
(Lurton et al., 2018). Bathymetrical and backscatter data were collected before (T0) and after (T1) trawling (Table S1 in the
Supplement). The RV followed the FV at an approximate distance of 300 m during one single haul in each of the experimental
plots to register the bathymetrical data. After six passes in each experimental plot, the RV collected MBES backscatter data
within 12 hours.

115 A high resolution (0.5 x 0.5 m) digital elevation model of the seabed was created from the MBES data using Qimera
(QPS). Trawl tracks were located from the GPS-derived position of the vessel path and visual detection using backscatter and
bathymetrical information. Transects of 12 m width were positioned every 20 m along the trawl track in GIS software (N = 74
for Pulse-ON, N = 100 for Pulse-OFF and N = 94 for Tickler; Fig. S2 in the Supplement). For the transects with visible trawl
marks, the mean water depth was calculated for five locations inside and outside the trawl tracks. The water depth was
120 estimated at these same locations at T0. The difference between mean water depths inside and outside the track was used to
assess if the trawl caused bathymetrical changes in the track.

A 1 x 1 m backscatter mosaic was created in FMGeocoder Toolbox (QPS FMGT). Backscatter data were sampled
by using transects for comparison of T0 and T1 and using all backscatter data in each experimental plot. Values outside 1.5
times the interquartile range were removed as outliers (1.5% of all values).

125 **2.5 Sediment profile imagery**

Sediment Profile Imagery (SPI) was used to characterize surface and subsurface sediment features using a Nikon D7100 Digital Camera (Ocean Imaging Systems, North Falmouth, MA, USA). The SPI is a device (lowered onto the sediment from the RV) that captures images of the sediment-water interface in a vertical cross-section and can provide information on benthic fauna, physical disturbance, and chemical features (e.g. redox depth) within the upper sediment (Rosenburg et al., 2003). Three replicate images were taken at each sampling station (3 stations per treatment plot; Fig. 1). One hundred eighty SPI photos were captured before (T0) and after (T1) trawling in the treatment plots (A – I) and also at the beginning (T0) and end (T1) of the experimental period for untrawled adjacent stations (AD1 – 3; Table S1 in the Supplement). Information for prism penetration depth (depth at which the camera penetrates the sediment) and visual observations of fauna, sediment characteristics (appearance of fine/coarse sediment, sand ripples etc.) and oxidation, expressed as % anoxic sediment (assessed by sediment color [black = anoxic]) were obtained using SpiArcBase software (Romero-Ramires et al., 2013). Individual SPI images can be found in Sect. S4 of the Supplement.

2.6 Benthic samples

Samples for sediment, macrofauna, porewater nutrients, and biogeochemical incubations were collected using cylindrical NIOZ box corers (50 cm height, 32 cm diameter; Netherlands Institute for Sea Research, Texel). For timesteps T0 and T1 for the fishing treatments, three replicate box cores were taken inside individual plots (northwest, central, and southeast locations). Samples were collected at untrawled adjacent stations AD1 and AD2 at the beginning (T0) and end (T1) of the experimental period, however, AD3 samples were only collected at the T1 timestep. This amounted to 69 total box cores collected and processed for this study. During experimental fishing activity, box cores in trawled sediments were collected within 0.5 h after 6 consecutive trawling passes were made in each treatment plot. Detailed sampling information can be found in Table S1 of the Supplement.

2.6.1 Sediment parameters

Two out of the three box cores were subsampled for sediment parameters from each trawl treatment plot and untrawled adjacent station. Sediment samples were subdivided at 0–0.5, 0.5–1, 1–1.5, 1.5–2, 2–2.5, 2.5–3, 3–4, 4–5, 5–6, 6–8, 8–10 cm slices taken from 3.5 cm inner diameter subcores. All samples were stored in a -20 °C freezer before freeze-drying and sieving (1 mm) for particle analysis. Sediment grain sizes were obtained via laser diffraction using a Malvern Mastersizer 2000 (McCave et al., 1986). Chlorophyll *a* (chl *a*) pigments were extracted using acetone and analyzed with UV spectrophotometry (Ritchie, 2006).

2.6.2 Porewater nutrients

155 Porewater was collected in the same box cores used for sediment parameters using 10 cm diameter sub cores with vertical sampling ports. Porewater was extracted at 0, 1, 2, 3, 5, 7 and 10 cm depth using rhizon samplers (Rhizophere Research Products; Seeberg-Elverfeldt et al., 2005; Dickens et al., 2007; Shobolt 2010). Porewater samples were added to 10 mL polystyrene vials and stored in -20 °C prior to analysis. To acquire nutrients, samples were thawed and analyzed using a SEAL QuAAtro segmented flow analyzer (Jodo et al., 1992) to determine concentrations of ammonium (NH_4^+), nitrite (NO_2^-), nitrate (NO_3^-), phosphate (PO_4^{3-}), and silicate ($\text{Si}(\text{OH})_4$).

160 2.6.3 Benthic biogeochemical fluxes

Incubation cores (15 cm inner diameter by 30 cm height) were collected from all box core samples (3x per treatment plot and reference station) with overlying water. These were subsequently placed in a water bath filled with surface seawater on board the RV, and aerated ~ 6 hours to acclimatize. Magnetic stirrers, fixed to the lid of incubation cores, ensured homogenized water column conditions. After the acclimatization period, cores were sealed from air contact for 4 hours to measure rates of sediment community oxygen consumption (SCOC) in the dark. During this period, oxygen concentrations were monitored every 30 s using optode sensors placed in the overlying water of the incubation cores (FireStingO₂, Pyroscience). At the end of the O₂ incubations, cores were re-opened and aerated to remain at above 90% O₂ saturation for the ongoing nutrient incubations. Ten mL water samples were collected at the overlying water of incubation cores at 0, 4, 8, 16 hour time intervals to obtain flux information on NH_4^+ , NO_2^- , NO_3^- , PO_4^{3-} , and $\text{Si}(\text{OH})_4$. Water samples for nutrient fluxes were stored and analyzed using the same methods described for porewater nutrients. Flux estimates were obtained by fitting linear regressions on nutrient concentrations changes over time and multiplying the regression coefficient with the height of the overlying water (Tiano et al., 2019).

2.6.4 Mass budget modelling

175 Rates for total mineralization of organic matter, denitrification, nitrification and the percentages of nitrogen and phosphorous removal were estimated using the measured sediment-water exchange fluxes of O₂, NH_4^+ , NO_2^- , NO_3^- , PO_4^{3-} calculated from the core incubations. These flux values were used to create an integrated mass budget of the solutes within the sediment column (Soetaert et al., 2001; Braeckman et al., 2010). A linear inverse model is implemented that estimates the rates of nitrification, denitrification, total mineralization and nutrient removal using the R package “limsolve” (Soetaert et al., 2009). More detailed methodology can be found in Sect. S5 in the Supplement.

180 2.6.5 Macrofauna collection

After the flux measurements, incubation cores were rinsed through a 1 mm sieve to collect the benthic macrofauna inside them. Faunal samples were preserved in 8% formalin seawater on the RV. Sorting and identification of preserved macrofauna samples took place at the Flanders Institute for Agricultural and Fisheries Research (ILVO) to obtain measurements for taxon densities and biomass (blotted wet weights) following the accredited procedure: BELAC ISO17025 norm (ILVO-DIER-185 ANIMALAB; Certificate N°:BELAC T-315).

2.7 Statistical analysis

All figures and statistical analyses were carried out using R (R Core Team 2021).

2.7.1 Water column

Differences between water column turbidity during Pulse-ON, Pulse-OFF and Tickler treatments were assessed with one way 190 ANOVA's and Tukey HSD pairwise tests. Assumptions of normality and homogeneity of variances were verified using Shapiro-Wilk and Levene's tests. If parametric assumptions were violated, data underwent a natural log-transformation prior to analysis.

2.7.2 MBES information

Bathymetrical differences between water depths inside and outside the track locations $\Delta_{i,T,G}$ were calculated for each transect 195 i at each time interval T (T_0 = before and T_1 = after trawling) for each trawl gear type G (1). A paired Wilcoxon signed-rank test was used to test if Delta-values were significantly different between T_1 and T_0 for each gear type. Significant positive Delta-values indicated that the water depth inside the trawl track was higher than outside the trawl track.

$$\Delta_{i,T,G} = (\text{water depth}_{\text{inside}} - \text{water depth}_{\text{outside track location}})_{i,T,G} \quad (1)$$

The indention I of each transect i was calculated by subtracting the delta value at T_0 from the Delta-value at T_1 (2).

200
$$I_{i,G} = (\Delta_{T_1} - \Delta_{T_0})_{i,G} \quad (2)$$

A resulting positive value indicated a deepening of the trawl track. The indention of the trawl was compared between trawl types using a Kruskal-Wallis Rank sum Test.

Statistical differences in backscatter values were compared for the combination of time and trawl types using experimental plots as replicates in a Kruskal-Wallis Rank sum Test. The experimental plots were also compared using the

205 Kruskal-Wallis Rank sum Test. A Dunn test with Benjamini-Hochberg method was applied for pairwise comparisons between sites.

2.7.3 SPI and benthic sample analysis

Linear mixed effects models (LMM) were used to investigate significant differences before and after fishing (T0 – T1) for: SPI measurements, sediment parameters, porewater nutrient concentrations, biogeochemical fluxes, mass budget model results and ecological characteristics (macrobenthos densities, biomass, species richness) using the *lmer*-function in the R package: “lme4” (Bates et al., 2015). For a given variable and treatment (example: oxygen flux and tickler treatment), a full-model was specified using the “timestep” (T0 - T1) as a fixed effect variable, “temperature” as a co-variate, and “station” as a random effect variable. A reduced-model was created to consider only the random effect variable (station). The full and reduced models were tested against each other using a partial F test. This approach was taken to assess the effect of fishing disturbance (timestep = before-after statistical comparison) in respect to temperature while minimizing spatial autocorrelation between stations.

2.7.4 Impact of *Lanice conchilega*

Robust regression analysis, using the R package: “robustbase” (Maechler et al., 2020), was conducted to investigate the effect of *L. conchilega* densities and biomass on physical, biogeochemical and faunal parameters in trawled and untrawled samples. This alternative to least squares techniques is an iterative analysis which down-weighs the influence of outliers on regression coefficients and is less affected by violations of linear regression assumptions (Koller and Stahel 2011). To reduce possible bias from the spatial differences in *L. conchilega* densities/biomass between treatment plots, all pre-trawled and post-trawled data were pooled together for this analysis.

3 Results

3.1 Water column conditions during trawling

225 During the experimental period, the water column exhibited mixed thermal conditions (no thermocline) with temperatures ranging between 15.8 – 17.0°C. Turbidity measured by the OBS sensors detected significantly higher amounts of water column SPM concentrations (mg L^{-1}) during tickler trawling (mean \pm standard deviation = 62.5 ± 17.1 , $p < 0.001$) compared to Pulse-ON (19.6 ± 5.3) and Pulse-OFF treatments (13.3 ± 5.8 ; Fig. 2). Data from a “Meetnet Vlaamse Banken” buoy (located in ~ 7 km southwest of the experimental location) showed current speeds ranging from 0.08 to 1.21 m s^{-1} during pulse trawling and 0.23 to 0.85 m s^{-1} during tickler trawling. On the day before both pulse trawl treatments, water currents reached a maximum speed of 1.23 m s^{-1} compared to 1.5 m s^{-1} the day before the tickler treatment (Fig. S3 in the Supplement).

3.2 Trawl effects on bathymetry

Delta-values were significantly different before and after trawling (T0-T1) for each trawl type (Pulse-ON: $p < 0.05$, Pulse-OFF: $p < 0.001$, Tickler: $p < 0.001$; Fig. S4 in the Supplement). The track locations were deepened, on average, by 1 cm after
235 trawling (Pulse-ON: 0.99 ± 3.92 ; Pulse-OFF: 1.17 ± 1.84 ; Tickler: 1.11 ± 2.74). The indentation of the trawl track was not significantly different between trawl treatments.

3.3 Trawl effects on backscatter

The mean backscatter values of the track locations in all treatments exhibited reduced values after 1 trawl pass ($\Delta T0 - T1$: Pulse-ON = -0.20; Pulse-OFF = -0.23; Tickler = -0.30) though there was no statistical difference between the three trawl types
240 (Table S2 in the Supplement). Comparison of mean backscatter values over entire plots (after 6 trawl passes) showed dB values ranging between -26.64 and -25.40 (Table S2 in the Supplement). The mean backscatter strength did not differ by more than 2 dB between the treatments at T0 and T1 and was not significantly different between any of the time-treatment interactions (Fig. S5 in the Supplement).

3.4 Sediment profile image analysis

245 *L. conchilega* tubeworms were identified in 72 % of images taken and were often associated with heterogeneous patterns of oxidized sediment (Fig. 3b). Mechanical disturbance was evident in certain T1 images showing “flattened” tubes and/or sediment (Fig. 3c). The percentage of anoxic sediment was significantly lower in T1 images from the tickler treatment compared to T0 images ($p < 0.01$) but did not differ significantly from T0 – T1 in other trawl treatments (Table 1). Anoxic (dark) sediment was significantly higher in images from the untrawled AD1 taken at the end of the experimental period (T1)
250 compared to the beginning (T0; $p < 0.01$). The sediment penetration depth of the SPI prism did not differ significantly between T0 – T1 for any treatment (Table 1).

3.5 Trawl effects on sediment parameters

Very fine sand fractions (62.5 – 125 μm) and chl *a* content in the sediment column (0 – 10 cm) were significantly lower in Tickler T1 compared to T0 samples ($p < 0.01$; $p < 0.001$; Table 2). For the Pulse-ON treatment, T1 silt fractions and chl *a*
255 content were significantly lower than before trawling ($p < 0.01$; Table 2). No significant differences in sediment parameters between T0 – T1 were found in the Pulse-OFF treatment. Particle size distributions of surface sediments (0 – 1 cm depth) between T0 – T1 for the Tickler and Pulse-ON treatment were characterized by a shift towards larger grain size fractions after trawling (Fig. 4).

3.6 Trawl effects on biogeochemical fluxes and nutrient profiles

260 Average SCOC rates ($\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$) were significantly reduced between T0 (172 ± 109) and T1 (74 ± 40) for the Tickler treatment ($p < 0.001$; Fig. 6a; Table 1). No other consistent or significant patterns were found for nutrient fluxes between T0 – T1 fishing treatments. Average and absolute values for nutrient fluxes can be found in Table 1 and Table S5 in the Supplement.

265 Average porewater nitrate concentrations within the sediment column (0 – 10 cm) were significantly higher after trawling in Tickler and Pulse-OFF treatments compared to pre-trawled sediments (Tickler, $p < 0.01$; Pulse-OFF, $p < 0.01$; Table 1). Ammonium concentrations in the sediment significantly increased from T0 to T1 ($p < 0.01$) after tickler trawling (Table 1). Mean phosphate concentrations were elevated after trawling for all experimental treatments but this was significant only in Pulse-ON ($p < 0.01$) and Pulse-OFF ($p < 0.001$; Table 1).

3.7 Mass budget calculations

270 The Tickler treatment exhibited significantly lower values for total mineralization ($p < 0.001$), denitrification ($p < 0.05$), and nitrification ($p < 0.05$) in T1 compared to T0 samples (Table 1). For all fishing treatments, the percentage of nitrogen removed after mineralization ranged between 80 and 95 % while sedimentary phosphorus removal varied between 91 and 99 %, however, no changes between T0 – T1 were statistically significant for individual fishing treatments.

3.8 Trawl effects on macrofauna density and macrobenthos composition

275 Sixty-nine taxonomical groups were identified in this study with *L. conchilega* dominating benthic macrobenthos densities (54%). The bivalve, *Kurtiella bidentata*, and polychaetes, *Magelona spp.*, and *Eumida sanguinea* made up 13%, 4% and 3% of total macrofauna density respectively. Though they only contributed 1% of total macrobenthos density, the razor clam, *Ensis leei*, represented 45% of total biomass collected followed by *L. conchilega* (22%), *Spisula subtruncata* (bivalve, 14%), and *Echinocardium cordatum* (heart urchins, 4%).

280 Total macrofaunal densities (individuals m^{-2}) decreased from T0 (Tickler: $11,782 \pm 7,360$; Pulse-ON: $9,783 \pm 14,072$; Pulse-OFF: $6,241 \pm 5,120$) to T1 (Tickler: $4,949 \pm 3,428$; Pulse-ON: $3,336 \pm 3,529$; Pulse-OFF: $5,483 \pm 3,623$) but were only statistically significant after the Tickler treatment ($p < 0.05$; Fig. 5b). Reference samples varied greatly but held the highest average densities from all the treatments (T0 = $19,174 \pm 17,493$; T1 = $22,993 \pm 17,945$; Fig. 5b).

285 *L. conchilega* maintained the highest average densities (individuals m^{-2}) in the reference stations (T0 = $10,703 \pm 9,943$; T1 = $14,707 \pm 11,336$). *L. conchilega* densities declined between T0 and T1 for Pulse-ON (T0 = $5,801 \pm 9,352$; T1 = $1,587 \pm 3,022$) the Pulse-OFF (T0 = $3,081 \pm 3,711$; T1 = $2,251 \pm 3,547$) and Tickler treatments (T0 = $5,880 \pm 3,838$; T1 = $2,172 \pm 1,913$), but was only significant for the Pulse-ON treatment ($p < 0.05$). Amongst other abundant taxa, *Abra alba* (bivalve) density declined

significantly after tickler trawling ($T_0 = 418 \pm 632$; $T_1 = 231 \pm 420$, $p < 0.05$) and *E. sanguinea* was significantly lower after the Pulse-ON treatment ($T_0 = 555 \pm 1,315$, $T_1 = 93 \pm 195$, $p < 0.001$).

290 **3.9 Effects of *Lanice conchilega* in trawled and untrawled sediments**

L. conchilega was found in 68 out of 69 (98.5 %) benthic sediment samples collected in this study. In pre-trawled sediments, *L. conchilega* density explained 85 % of variation in the abundances of the other taxa ($p < 0.001$) and while biomass explained 38 % of variation in species richness ($p < 0.001$; Fig. 6, Table 2). Tubeworm densities in pre-trawled samples were linked with smaller sediment grain sizes, accounting for 33 % of variability in the percentage of very fine sand ($p < 0.05$) while biomass explained 18 % of variability in silt content ($p > 0.01$) (Table 2). Densities of *L. conchilega* significantly predicted 87 % of SCOC in undisturbed sediments ($p < 0.001$). Biomass of *L. conchilega* (compared to density) was a slightly better predictor of SCOC and explained 93 % of variation in O_2 fluxes ($p < 0.001$) as well as 21 % of nitrate effluxes ($p < 0.01$) in untrawled sediments (Fig. 6; Table 2). No significant relationships were found after trawling (Fig. 6, Table 2). Due to lack of sufficient data, regressions between $T_0 - T_1$ within individual fishing treatments (Pulse-ON, Pulse-OFF, Tickler) did not yield consistent results.

4 Discussion

The data gathered in this study exhibited a large amount of spatial variability reflecting the dynamic nature of the Vlakte van de Raan ecosystem (Degraer and Hostens, 2016). Despite the challenge of detecting disturbance in such a variable system, we were able to provide a comprehensive assessment of acute bottom trawling with our 3 x 3 plot design. The dominance of *L. conchilega* in this habitat provided an additional opportunity to explore the effects of tubeworm abundance on environmental parameters in disturbed and undisturbed conditions.

This study follows up on similar research conducted in the Frisian Front region of the North Sea (Depestele et al., 2018, Tiano et al., 2019; Tiano et al., 2020) with each study (including the current one) collecting information in the same season (early June) of its respective year. The offshore Frisian Front ecosystem (~ 60 km from the Dutch Coast), however, experiences much lower levels of tidal disturbance, exhibits smaller grain sizes, and holds completely different macrofaunal assemblages and biogeochemical characteristics compared to the Vlakte van de Raan (van Raaphorst et al., 1992; Dewicke et al., 2002; Degraer and Hostens, 2016; Depestele et al., 2018). By comparing the findings from these two contrasting areas, we can better understand how different ecosystems are affected by bottom disturbance. To our knowledge, this is the first coastal habitat study that simultaneously investigates acute beam trawl impacts on biological (Rabaut et al., 2008), physical (Depestele et al., 2016) and biogeochemical dynamics.

4.1 SPM during experimental trawling

The water column was 73-78% more turbid during tickler trawling compared to both pulse trawl treatments. As tickler trawls tow at faster speeds compared to pulse trawls (Poos et al., 2020), they are predicted to mobilize more sediment (Rijnsdorp et al., 2021). However, due to the higher current speeds evident in the day prior to tickler trawling, we cannot be certain that this increased turbidity was the result from enhanced resuspension caused from bottom trawling alone. It is possible that trawling coupled with current induced resuspension enhanced SPM measurements during the Tickler treatment. This would corroborate with the findings in Depestele et al., (2016) where no significant differences in sediment resuspension of pulse and tickler trawling was found in a similar coastal area. In the Frisian Front, where Tiano et al., (2019) reported conspicuous SPM peaks attributed to individual trawl passes, water current speeds were 62 – 88 % lower during fishing compared to the Vlakte van de Raan study. The more ambiguous link between trawling and SPM in our study (no obvious turbidity peak during trawl passes) suggests that the water column effects of trawling are more difficult to detect in areas with higher natural current speeds. These effects may be diluted through current induced dispersion of resuspended particles and/or higher levels of natural turbidity.

4.2 Changes to the physical environment

A single trawl pass for all three treatments yielded a bathymetrical indentation of ~ 1 cm in the sediment, similar to the 0.9 cm found after tickler trawling in coarse sediment off the Dutch coast (Depestele et al., 2016). Our study, however, did not replicate previous findings, which found greater bathymetrical effects for tickler beam trawling in finer North Sea sediments (Depestele et al., 2018). This is likely due to differences in grain size which are smaller in the Frisian Front (aforementioned study), compared to the Vlakte van de Raan (current study). Backscatter data, which produced similar values between T0-T1 in our study, can approximate sediment characteristics and trawling disturbance (Roche et al., 2018), but do not purely reflect sediment grain size and can be distorted by “scatterers” such as shell debris and biogenic reefs, which may interfere with the acoustic signals (Degraer et al., 2008; Feldens et al., 2018).

Tubeworm reefs in untrawled sediments were positively associated with smaller grain size fractions (Fig. 3; Table 2). This pattern is commonly found with *L. conchilega* as high tube densities will slow down bottom current velocities leading to the settling of smaller particles (Van Hoey et al., 2008; De Smet et al., 2015; Foshtomi et al., 2018). Trawl induced coarsening of sediments found in *L. conchilega* dominated habitats may occur through both the resuspension and winnowing of finer grained particles from direct trawling impacts (Palanques et al., 2014; Mengual et al., 2016; Depestele et al., 2018; Tiano et al., 2019) and the alteration of sediment characteristics caused by the rapid extraction of tube structures from the seabed.

The shift in particle sizes and changes in observed anoxic sediment after trawling, suggests that the largest sediment alterations were attributed to the Tickler treatment followed by the Pulse-ON treatment with no significant effects detected after Pulse-OFF trawling (Fig. 4; Table 1). As electricity from pulse trawls does not affect physical sediment characteristics (Tiano et al., 2021), differences between Pulse-ON and Pulse-OFF treatments may have been due the variable mechanical impacts of pulse trawls (Depestele et al., 2018) or the environmental variability of the site (Degraer and Hostens, 2016).

4.3 Changes to biogeochemical characteristics

We detected significant declines in chl *a* from surface sediment (0 – 1 cm) of 63% and 37% after tickler and pulse trawling (Pulse-ON) respectively. These percent changes are lower relative to the reductions measured after tickler (83 %) and pulse (43 %) trawling in the siltier Frisian Front sediments (Tiano et al., 2019) even with the three to six times higher sediment chl *a* content in the more coastal Vlake van de Raan system. Sandy dynamic habitats, like the Vlake van de Raan, experience higher bed shear stress and are generally less vulnerable to trawling impacts compared to muddier more static environments like the Frisian Front (Kaiser et al., 2006; Queirós et al., 2006; Allen and Clarke, 2007; Sciberras et al., 2016; Rijnsdorp et al., 2020a).

Measured SCOC values in the present study were exceptionally high indicating biogeochemically active sediments (Stratmann et al., 2019). There are several factors driving high SCOC in the Vlake van de Raan including enhanced water temperatures during the experimental period, increased oxidation by bioirrigation from *L. conchilega* (evidenced in SPI images; Fig. 3), and the breakdown of fresh organic matter. High levels of the latter (implied by chl *a* concentrations; Table 1) coupled with elevated faunal and microbial activity due to increased temperatures, is the likely mechanism for the high SCOC and mineralization found in this study.

Despite the lower trawl penetration into the sediment (~ 1 cm Vlake van de Raan vs. ~ 4 cm Frisian Front; Depestele et al., 2019), our study produced a stronger statistically significant reduction in SCOC (Tickler: 57 %) compared to Tiano et al., (2019; Tickler: 41 %, Pulse: 33 %). Significant reductions in total mineralization, denitrification, and nitrification after the Tickler treatment suggests that trawling reduced benthic nutrient removal and recycling functions in this study. Furthermore, higher percentages in the anoxic surface area from the SPI images taken in adjacent sites (significant for AD1) may suggest increased mineralization in undisturbed areas during the course of the experiments (Table S5 in the Supplement). Mass budget estimates for mineralization were also higher on average at T1 for AD1 and AD2, though this was not statistically significant. If the benthic metabolism in untrawled areas did increase during the experimental period, the trawling effects detected in this study (lower metabolism after trawling) may be considered to be conservative estimates.

The large changes in SCOC may be explained by the decrease in fauna such as *L. conchilega*, which can be highly influential to SCOC and other biogeochemical fluxes (Braeckman et al., 2010). In addition to respiration, when *L. conchilega* (and other infauna) irrigate their burrows, they can drastically expand the surface area available for sediment-oxygen exchange (De Smet et al., 2016b; Kristensen and Kostka, 2013), with major impacts to biogeochemical dynamics (Olsgard et al., 2008; Braeckman et al., 2010). These results also suggest that areas with *L. conchilega* may be vulnerable to relatively shallow (~ 1 cm) seabed disturbances as their tubes extend from over 10 cm within the sediment to above the seabed surface. Contrary to this, faunal-mediated biogeochemical functions in the Frisian Front are relegated to organisms such as the burrowing mud shrimp, *Callinassa subterranea*, that do not exhibit any protruding sediment surface structures and reside deeper within the seabed (Amaro et al., 2007; Tiano et al., 2020). These organisms are probably less affected by trawling despite their habitats displaying softer sediments which are prone to greater trawl penetration (Depestele et al., 2019; Tiano et al., 2019, 2020).

L. conchilega in untrawled sediments had a clear effect on O₂ and NO₃ fluxes, though these relationships disappeared after disturbance (Fig. 6, Table 2). The change in these relationships probably came from a combination of reduced average tubeworm densities (consistently lower after trawling but only significant after the Pulse-ON treatment) and direct trawl-induced habitat alterations. These results support the assertion by Hale et al., (2017), that trawling decouples faunal-mediated biogeochemical processes.

The Pulse-ON treatment, which led to the biggest decrease (72%) and lowest *L. conchilega* densities after T1, was also the only experimental treatment that did not produce a significant increase in porewater nitrate concentrations. Previous research suggests a density dependent enhancement by *L. conchilega* on denitrification/nitrification processes (Aller, 1988; Braeckman et al., 2010; Foshtomi et al., 2018). Trawling has also been shown to decrease denitrification by altering the sediment layers that optimize denitrification and reducing total mineralization rates (Ferguson et al., 2020; De Borger et al., 2021). It is possible that both mechanical trawl disturbance and a loss (Pulse-ON) or retention (Pulse-OFF, Tickler) of functional species roles after trawling led to the enhanced sediment nitrate concentrations, though it is difficult to pinpoint the exact cause of this increase.

4.4 Benthic community observations

Our study found a significant decrease in *L. conchilega* densities with the Pulse-ON treatment and not in the other trawl treatments, however, it is unlikely that electrical stimulation caused this reduction. Previous research has not found any direct mortality of fish or benthos associated with electrical stimulation used in the pulse trawl fishery (Polet et al., 2005; van Marlen et al., 2009; Soetaert et al., 2015b; Soetaert et al., 2016; ICES, 2018, 2019, 2020; Boute et al., 2020). Moreover, electric stimulation from flatfish pulse trawls cannot physically remove *L. conchilega* tubes from the sediment (Boute et al., 2020; ICES 2020; Tiano et al., 2021). Our results, however, point to higher physical disturbance occurring in the Pulse-ON areas as being the likely culprit for reduced *L. conchilega* densities (Fig. 5, Table 2).

The Pulse-ON treatment also caused a significant decrease of *Eumida sanguinea*, a species closely associated to *L. conchilega* (Rabaut et al., 2008) though other taxa, such as *Abra alba*, only showed notable decreases in response to tickler trawling. Nonetheless, when accounting for total macrobenthos densities, the only significant effect came from the Tickler treatment (58% decrease) reflecting the more consistent effect associated with tickler beam trawl gears (Depestele et al., 2018; Tiano et al., 2019). Pre-trawled habitats in this study linked *L. conchilega* to the densities of other taxa and species richness, though this relationship was no longer detected after experimental trawling (Fig. 6, Table 2). This suggests that, although tubeworm densities were impacted to some extent, trawling disproportionately affected the corresponding taxa. Rabaut et al., (2008) reported significant changes to an intertidal *L. conchilega* dominated community after beam trawling but not to the tubeworms densities themselves. Our results show a similar removal of species associated with *L. conchilega* (*E. sanguinea* in particular), though, our findings of decreased tubeworm densities support evidence that *L. conchilega* shows a certain level of sensitivity to physical sediment disturbances (Callaway et al., 2010). *L. conchilega* has been found to successfully recolonize

previously disturbed habitats within 1-2 years suggesting high resilience but not necessarily resistance to anthropogenic impacts (Callaway et al., 2010).

415 **5 Conclusions**

This study found various effects from conventional tickler rigged beam trawls and electric pulse trawl techniques. Though pulse trawls (and any other bottom fishing technique) can undoubtedly cause significant habitat alterations, these effects tend to be quite variable (Tiano et al., 2019 and this study). The more consistent effect of tickler trawls are, on average, higher than that of pulse trawls as shown in the sediment, geochemical, and species community level changes seen in this study. However, we also show that pulse trawls may be at least equally as impactful to densities of certain benthos. It is likely that the lower average penetration depth of the lighter pulse trawl gears are less damaging to certain benthic communities compared to tickler trawls (Hiddink et al., 2017; Depestele et al., 2018), however, this reduced effect is dependent on species assemblages and sediment type. Compared to any difference in direct impacts between pulse and beam trawls (due to different sediment penetration depths), the potential for a reduced environmental footprint from pulse trawls (compared to beam trawls) stems more from its higher catch efficiency which leads to reduced time needed to catch their quotas (less time fishing) when exploiting North Sea sole (*Solea solea*; Rijnsdorp et al., 2020b; ICES, 2020).

The high SCOC values found at our experimental location imply that this community is an important site for carbon degradation (Stratmann et al., 2019). The trawl-induced declines in SCOC and total mineralization suggest that some disturbed locations in this community lost over half of their ability to mineralize organic matter. This highlights the propensity for bottom trawling to reduce carbon cycling in marine sediments and shift the mineralization processes to the overlying water (Tiano et al., 2019). With enhanced mineralization rates in the water column (due to higher oxic conditions compared to inside the sediment), trawling may lead to the removal of sedimentary carbon, increased nitrogen retention in the seabed (decreased denitrification), and a reallocation of nutrients away from trawled sediments (Tiano et al., 2019; Ferguson et al., 2020; De Borger et al., 2021; Tiano et al., 2021).

Results in this study show that even highly dynamic coastal areas can be susceptible to significant trawl-induced changes depending on the type of benthic community present. Biogenic reef communities like the ones found in the Vlakte van de Raan, can be highly productive habitats with enhanced biogeochemical characteristics. With these features, the potential for disturbance-induced changes becomes greater if keystone species such as *L. conchilega* are affected. When assessing the vulnerability of marine habitats, environmental managers and policy makers need to consider abundant ecosystem engineers as they perform important ecological and biogeochemical functions.

Data availability

Data and R code can be made available on request to the corresponding author.

Author contribution

JT, JD, GvH, and KS devised the study, and contributed to the manuscript. JT, PvR and KS collected and analyzed data for
445 water column, nutrient flux, porewater nutrients, and sediment. JD, GvH and JF collected and analyzed data for sediment
profile imagery, macrofauna, and sonar (bathymetry and backscatter).

Competing interests

GvH was a member of the editorial board for this special issue of Biogeosciences: “Towards an understanding and assessment
of human impact on coastal marine environments”, however, the peer-review process was guided by an independent editor,
450 and no other authors have competing interests nor financial benefits to declare resulting from the publication of this manuscript.

Acknowledgements

*We would like to thank the crews of the TH6 for YE238 for their excellent work and coordination in making multiple trawled
areas. We extend a huge thank you to the crew of the Simon Stevin for helping us collect our data and putting up with our
annoying requests to do more sampling and the students at ILVO who helped process our macrofauna samples. This work
455 would not have been possible without the collaboration with the Flanders Marine Institute (VLIZ) who provided the RV Simon
Stevin, along with the multiple data collection devices (multi-beam, CTD, SPI etc.). Thank you, Chiu Cheng, Bea Vega Garre,
and Wesley Kreeft for making our lives easier during the campaign by helping with sample collection. Finally, we send a
special thank you to Peter van Breugel, Yvonne Maas, Jan Peene, Jurian Brassers, Yke Timmermans and any other members
of the NIOZ analytical lab for processing our sediment and biogeochemical samples. This research is partially funded by the
460 European Maritime and Fisheries Fund (EMFF) and the Netherlands Ministry of Agriculture Nature and Food Quality (LNV).*

References

- Allen, J. I., & Clarke, K. R.: Effects of demersal trawling on ecosystem functioning in the North Sea: A modelling study.
Mar. Ecol. Prog. Ser., 336, 63–75. <https://doi.org/10.3354/meps336063>, 2007
- Aller, R. C.: Benthic fauna and biogeochemical processes in marine sediments: the role of burrow structures. In: Blackburn
465 TH, Sørensen J (eds) Nitrogen cycling in coastal marine environments. Scope, Chichester, p 301–338, 1988
- Amaro, T. P. F., Duineveld, G. C. A., Bergman, M. J. N., Witbaard, R., & Scheffer, M.: The consequences of changes in
abundance of *Callianassa subterranea* and *Amphiura filiformis* on sediment erosion at the Frisian Front (south-eastern North
Sea). *Hydrobiologia*, 589(1), 273–285. <https://doi.org/10.1007/s10750-007-0750-2>, 2007

- Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C.: Fitting linear mixed-effects models using lme4. *J. Stat. Softw.*, 470 67(1). <https://doi.org/10.18637/jss.v067.i01>, 2015
- Bolam, S. G., Coggan, R. C., Eggleton, J., Diesing, M., & Stephens, D.: Sensitivity of macrobenthic secondary production to trawling in the English sector of the Greater North Sea : A biological trait approach. *J. Sea Res.*, 85, 162–177. <https://doi.org/10.1016/j.seares.2013.05.003>, 2014
- Boute, P.G., Soetaert, M., Reid Navarro, J.A., Lankheet, M. J.: Effects of electrical pulse stimulation on behaviour and survival of marine benthic invertebrates. *Front. Mar. Sci.*, (February). <https://doi.org/10.3389/fmars.2020.592650>, 2020 475
- Bradshaw, C., Jakobsson, M., Brüchert, V., Bonaglia, S., Mörth, C.-M., Muchowski, J., Stranne, C., & Sköld, M.: Physical Disturbance by Bottom Trawling Suspends Particulate Matter and Alters Biogeochemical Processes on and Near the Seafloor. *Front. Mar. Sci.*, 8(August). <https://doi.org/10.3389/fmars.2021.683331>, 2021
- Braeckman, U., Provoost, P., Gribsholt, B., Van Gansbeke, D., Middelburg, J. J., Soetaert, K., Vincx, M., & Vanaverbeke, 480 J.: Role of macrofauna functional traits and density in biogeochemical fluxes and bioturbation. *Mar. Ecol. Prog. Ser.*, 399, 173–186. <https://doi.org/10.3354/meps08336>, 2010
- Callaway, R., Desroy, N., Dubois, S., Fournier, J., Frost, M., Godet, L., Hendrick, V. J., & Rabaut, M.: Ephemeral Bio-engineers or Reef-building Polychaetes : How Stable are Aggregations of the Tube Worm *Lanice conchilega* (Pallas, 1766)?. *Integr. Comp. Biol.*, 50(2), 237–250. <https://doi.org/10.1093/icb/icq060>, 2010
- 485 Craeymeersch, J. A., Escaravage, V., Steenbergen, J., Wijsman, J., Wijnhoven, S., & Kater, B.: De bodemfauna in het Nederlands deel van de Scheldemonding, Issue Ministerie van infrastructuur en Waterstaat Report, Institute for Marine Resources and Ecosystem Studies, Netherlands, 105 pp., 2006
- Dray, S., Dufour, A.: Ade4: Analysis of Ecological Data. *Exploratory and Euclidean Methods in Environmental Sciences*, 22(4), 2007
- 490 Dauwe, B., Herman, P. M. J., & Heip, C. H. R.: Community structure and bioturbation potential of macrofauna at four North Sea stations with contrasting food supply. *Mar. Ecol. Prog. Ser.*, 173(1978), 67–83. <https://doi.org/10.3354/meps173067>, 1998
- De Borger, E., Tiano, J., Braeckman, U., Rijnsdorp, A., & Soetaert, K.: Impact of bottom trawling on sediment biogeochemistry: a modelling approach. *Biogeosciences*, April, 1–32. <https://doi.org/10.5194/bg-2020-328>, 2021
- 495 De Smet, B., Van Oevelen, D., Vincx, M., Vanaverbeke, J., & Soetaert, K.: *Lanice conchilega* structures carbon flows in soft-bottom intertidal areas. *Mar. Ecol. Prog. Ser.*, 552(June), 47–60. <https://doi.org/10.3354/meps11747>, 2016a
- De Smet, B., Braeckman, U., Soetaert, K., Vincx, M., & Vanaverbeke, J.: Predator effects on the feeding and bioirrigation

- activity of ecosystem-engineered *Lanice conchilega* reefs. *J. Exp. Mar. Biol. Ecol.*, 475, 31–37.
<https://doi.org/10.1016/j.jembe.2015.11.005>, 2016b
- 500 De Smet, B., D'Hondt, A. S., Verhelst, P., Fournier, J., Godet, L., Desroy, N., Rabaut, M., Vincx, M., & Vanaverbeke, J.: Biogenic reefs affect multiple components of intertidal soft-bottom benthic assemblages: The *Lanice conchilega* case study. *Estuar. Coast.*, 152, 44–55. <https://doi.org/10.1016/j.ecss.2014.11.002>, 2015
- Degraer, S., Hostens, K.: Assessment of the conservation value of the Vlakte van de Raan Site of Community Interest. MARECO report 16/02. Royal Belgian Institute of Natural Sciences, Operational Directorate Natural Environment, Aquatic and Terrestrial Ecology, Marine Ecology and Management, Brussels. 16 pp., 2016.
- 505 Degraer, S., Moerkerke, G., Rabaut, M., Van Hoey, G., Du Four, I., Vincx, M., Henriet, J. P., & Van Lancker, V.: Very-high resolution side-scan sonar mapping of biogenic reefs of the tube-worm *Lanice conchilega*. *Remote Sens. Environ.*, 112(8), 3323–3328. <https://doi.org/10.1016/j.rse.2007.12.012>, 2008.
- Depestele, J., Degrendele, K., Esmaili, M., Ivanovi, A., Kro, S., Neill, F. G. O., Parker, R., Polet, H., Roche, M., Teal, L. R., Vanelslander, B., & Rijnsdorp, A. D.: Comparison of mechanical disturbance in soft sediments due to tickler-chain SumWing trawl vs . electro-fitted PulseWing trawl. *ICES J. Mar. Sci.*, fsy124, 1–18. <https://doi.org/10.1093/icesjms/fsy124>, 2018
- 510 Depestele, J., Ivanovic, A., Degrendele, K., Esmaili, M., Polet, H., Roche, M., Summerbell, K., Teal, L. R., Vanelslander, B., & O'Neil, F. G.: Measuring and assessing the physical impact of beam trawling. *ICES J. Mar. Sci.*, 73, i15–i26. <https://doi.org/10.1093/icesjms/fst176>, 2018
- 515 Dewicke, A., Rottiers, V., Mees, J., & Vincx, M.: Evidence for an enriched hyperbenthic fauna in the Frisian front (North Sea). *J. Sea Res.*, 47(2), 121–139. [https://doi.org/10.1016/S1385-1101\(02\)00106-5](https://doi.org/10.1016/S1385-1101(02)00106-5), 2002
- Dickens, G. R., Koelling, M., Smith, D. C., & Schnieders, L.: Rhizon sampling of pore waters on scientific drilling expeditions: An example from the IODP expedition 302, Arctic Coring Expedition (ACEX). *Sci. Drill.*, 4, 22–25. <https://doi.org/10.2204/iodp.sd.4.08.2007>, 2007
- 520 Eigaard, O. R., Bastardie, F., Breen, M., Dinesen, G. E., Hintzen, N. T., Laffargue, P., Mortensen, L. O., Nielsen, J. R., Nilsson, H. C., Neill, F. G. O., Smith, C., Sørensen, T. K., Polet, H., Reid, D. G., Sala, A., Sko, M., Tully, O., Zengin, M., & Rijnsdorp, A. D.: Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. *ICES J. Mar. Sci.*, 73(Effects of Fishing on Benthic Fauna, Habitat and Ecosystem Function), 2016
- 525 Fariñas-Franco, J. M., Allcock, A. L., & Roberts, D.: Protection alone may not promote natural recovery of biogenic habitats of high biodiversity damaged by mobile fishing gears. *Mar. Environ. Res.*, 135(September 2017), 18–28. <https://doi.org/10.1016/j.marenvres.2018.01.009>, 2018

- Feldens, P., Schulze, I., Papenmeier, S., Schönke, M., & von Deimling, J. S.: Improved interpretation of marine sedimentary environments using multi-frequency multibeam backscatter data. *Swiss. J.*, 8(6).
530 <https://doi.org/10.3390/geosciences8060214>, 2018
- Ferguson, A. J. P., Oakes, J., & Eyre, B. D.: Bottom trawling reduces benthic denitrification and has the potential to influence the global nitrogen cycle. *Limnol. Oceanogr.*. <https://doi.org/10.1002/lol2.10150>, 2020
- Foshtomi, M. Y., Leliaert, F., Derycke, S., Willems, A., Vincx, M., & Vanaverbeke, J.: The effect of bio-irrigation by the polychaete *Lanice conchilega* on active denitrifiers: Distribution, diversity and composition of *nosZ* gene. *PLoS ONE*, 13(2),
535 1–25. <https://doi.org/10.1371/journal.pone.0192391>, 2018
- Grabowski, J. H., Bachman, M., Demarest, C., Eayrs, S., Harris, B. P., Malkoski, V., Packer, D., & Stevenson, D.: Assessing the Vulnerability of Marine Benthos to Fishing Gear Impacts. *Rev. Fish. Sci. Aquac.*, 22(2), 142–155.
<https://doi.org/10.1080/10641262.2013.846292>, 2014
- Hale, R., Godbold, J. A., Sciberras, M., Dwight, J., Wood, C., Hiddink, J. G., & Solan, M.: Mediation of macronutrients and
540 carbon by post-disturbance shelf sea sediment communities. *Biogeochemistry*, 135(1–2), 121–133.
<https://doi.org/10.1007/s10533-017-0350-9>, 2017
- Hiddink, J. G., Jennings, S., Sciberras, M., Szostek, C. L., Hughes, K. M., & Ellis, N.: Global analysis of depletion and recovery of seabed biota after bottom trawling disturbance. *P. Natl. Acad. Sci. USA*, 114, 8301–8306.
<https://doi.org/10.1073/pnas.1618858114>, 2017
- 545 ICES: Working Group on Electrical Trawling (WGELECTRA). *ICES Scientific Reports*, 2:37, 108 pp.,
<https://doi.org/10.17895/ices.pub.5619>, 2020
- ICES: Working Group on Electrical Trawling (WGELECTRA). *ICES Scientific Reports*. 1:71, Ghent, Belgium, 81 pp.,
<http://doi.org/10.17895/ices.pub.5619>, 2019
- ICES. 2018. Report of the Working Group on Electric Trawling (WGELECTRA). *ICES Report WGELECTRA 2018 17 - 19*
550 April 2018. IJmuiden, the Netherlands. 155pp., 2018
- Jodo, M., Kawamoto, K., Tochimoto, M., & Coverly, S. C.: Determination of nutrients in seawater by analysis rate and reduced interference on segmented-flow analysis with higher ammonia. *J. Autom. Chem.*, 14(5), 163–167, 1992
- Kaiser, M. J., Clarke, K. R., Hinz, H., Austen, M. C. V., Somerfield, P. J., & Karakassis, I.: Global analysis of response and recovery of benthic biota to fishing. *Mar. Ecol. Prog. Ser.*, 311, 1–14, 2006
- 555 Kaiser, M. J., Collie, J. S., Hall, S. J., Jennings, S., & Poiner, I. R.: Modification of marine habitats by trawling activities: prognosis and solutions. *Fish. Fish.*, 3, 114–136, 2002

- Kristensen, E., & Kostka, J. E.: Macrofaunal Burrows and Irrigation in Marine Sediment: Microbiological and Biogeochemical Interactions. *Interactions Between Macro- and Microorganisms in Marine Sediments, The Ecogeomorphology of Tidal Marshes Coastal and Estuarine Studies 59*, Copyright 2004 by the American Geophysical Union, pp., 125–157. <https://doi.org/10.1029/CE060p0125>. 2013
- 560
- Koller and Stahel: Sharpening Wald-type inference in robust regression for small samples. *Computational Statistics and Data Analysis*. 55, 2504-2515, 2011
- Lurton, X., Eleftherakis, D., and Augustin, J. M.: Analysis of seafloor backscatter strength dependence on the survey azimuth using multibeam echosounder data. *Mar. Geophys. Res.*, 39: 183-203, 2018.
- 565
- Maechler M, Rousseeuw P, Croux C, Todorov V, Ruckstuhl A, Salibian-Barrera M, Verbeke T, Koller M, Conceicao EL, Anna di Palma M : robustbase: Basic Robust Statistics. R package version 0.93-6, <http://robustbase.r-forge.r-project.org/>, 2018
- McCave, I. N., Bryant, R. J., Cook, H. F., & Coughanowr, C. A.: Evaluation of a laser-diffraction-size analyzer for use with natural sediments. *Res. Meth. Pap.*, 56, 561–564, 1986
- 570
- Mengual, B., Cayocca, F., Le Hir, P., Draye, R., Laffargue, P., Vincent, B., & Garlan, T.: Influence of bottom trawling on sediment resuspension in the ‘Grande-Vasière’ area (Bay of Biscay, France). *Ocean Dynamics*, 66(9), 1181–1207. <https://doi.org/10.1007/s10236-016-0974-7>, 2016
- Mermillod-Blondin, F., Rosenberg, R.: Ecosystem engineering: the impact of bioturbation on biogeochemical processes in marine and freshwater benthic habitats. *Aquat. Sci.*, 68 (4): 434-442. 2006
- 575
- Morys, C., Brüchert, V., & Bradshaw, C.: Impacts of bottom trawling on benthic biogeochemistry in muddy sediments: Removal of surface sediment using an experimental field study. *Mar. Environ. Res.*, 169(December 2020), 105384. <https://doi.org/10.1016/j.marenvres.2021.105384>, 2021
- Nieuwenhuize, J., Maas, Y. E. M., & Middelburg, J. J.: Rapid analysis of organic carbon and nitrogen in particulate materials. *Mar. Chem.*, 45, 217–224, 1994
- 580
- Olsgard, F., Schaanning, M. T., Widdicombe, S., Kendall, M. A., & Austen, M. C.: Effects of bottom trawling on ecosystem functioning. *J. Exp. Mar. Biol. Ecol.*, 366(1–2), 123–133. <https://doi.org/10.1016/j.jembe.2008.07.036>, 2008
- Paradis, S., Goni, M., Masqué, P., Duran, R., Aronja-Camas, M., Palanques, A., & Puig, P.: Persistence of Biogeochemical Alterations of Deep-Sea Sediments by Bottom Trawling *Geophys. Res. Lett.*, 48(e2020GL091279). <https://doi.org/10.1029/2020GL091279>, 2021.
- 585
- Palanques, A., Puig, P., Guillén, J., Demestre, M., & Martín, J.: Effects of bottom trawling on the Ebro continental shelf

- sedimentary system (NW Mediterranean). *Cont. Shelf Res*, 72, 83–98. <https://doi.org/10.1016/j.csr.2013.10.008>, 2014
- Pecceu, E., Hostens, K., & Maes, F.: Governance analysis of MPAs in the Belgian part of the North Sea. *Mar. Policy*, 71, 265–274. <https://doi.org/10.1016/j.marpol.2015.12.017>, 2016
- Polet, Hans; Delanghe, Fernand ; Verschoore, R.: On electrical fishing for brown shrimp (*Crangon crangon*) I. Laboratory experiments. *Fish. Res.*, 72, 1–12, 2005
- Poos, J., Hintzen, N. T., van Rijssel, J. C., & Rijnsdorp, A.: Efficiency changes in bottom trawling for flatfish species as a result of the replacement of mechanical stimulation by electric stimulation. *ICES J. Mar. Sci.*, October. <https://doi.org/10.1093/icesjms/fsaa126>, 2020
- Queirós, A. M., Hiddink, J. G., Kaiser, M. J., & Hinz, H.: Effects of chronic bottom trawling disturbance on benthic biomass, production and size spectra in different habitats. *J. Exp. Mar. Biol. Ecol.*, 335(1), 91–103. <https://doi.org/10.1016/j.jembe.2006.03.001>, 2006
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2021
- Rabaut, M.: *Lanice conchilega*, fisheries and marine conservation, PhD Thesis. Ghent University. Ghent, Belgium, 354 pp. 2009
- Rabaut, M., Braeckman, U., Hendrickx, F., Vincx, M., & Degraer, S.: Experimental beam-trawling in *Lanice conchilega* reefs: Impact on the associated fauna. *Fish. Res.*, 90(1–3), 209–216. <https://doi.org/10.1016/j.fishres.2007.10.009>, 2008
- Rijnsdorp, A. D., Depestele, J., Molenaar, P., Eigaard, O. R., Ivanović, A., & O’Neill, F. G.: Sediment mobilization by bottom trawls: A model approach applied to the Dutch North Sea beam trawl fishery. *ICES J. Mar. Sci.*, 78(5), 1574–1586. <https://doi.org/10.1093/icesjms/fsab029>, 2021
- Rijnsdorp, A D, Hiddink, J. G., van Denderen, P. D., Hintzen, N. T., Eigaard, O. R., Valanko, S., Bastardie, F., Bolam, S. G., Boulcott, P., Egekvist, J., Garcia, C., van Hoey, G., Jonsson, P., Laffargue, P., Nielsen, J. R., Piet, G. J., Sköld, M., & van Kooten, T.: Different bottom trawl fisheries have a differential impact on the status of the North Sea seafloor habitats. *ICES J. Mar. Sci.*. <https://doi.org/10.1093/icesjms/fsaa050>, 2020a
- Rijnsdorp, A. D., Depestele, J., Eigaard, O. R., Hintzen, N. T., Ivanovic, A., Molenaar, P., O’Neill, F. G., Polet, H., Poos, J. J., & van Kooten, T.: Mitigating seafloor disturbance of bottom trawl fisheries for North Sea sole *Solea solea* by replacing mechanical with electrical stimulation. *PLoS ONE*, 15(11 November). <https://doi.org/10.1371/journal.pone.0228528>, 2020b
- Ritchie, R. J.: Consistent sets of spectrophotometric chlorophyll equations for acetone, methanol and ethanol solvents. *Photosynth. Res.*, 89(1), 27–41. <https://doi.org/10.1007/s11120-006-9065-9>, 2006

- 615 Roche, M., Degrendele, K., Vrignaud, C., Loyer, S., Le Bas, T., Augustin, J.-M., and Lurton, X.: Control of the repeatability of high frequency multibeam echosounder backscatter by using natural reference areas. *Mar. Geophys. Res.*, 39: 89-104, 2018
- Romero-Ramirez, A., Gremare, A., Desmalades, M., & Duchene, J. C.: Semi-automatic analysis and interpretation of sediment profile images. *Environ. Modell. Softw.*, 47, 42–54. <https://doi.org/10.1016/j.envsoft.2013.04.008>, 2013
- 620 Rosenberg, R., Nilsson, H., Grémare, A., & Amouroux, J.: Effects of demersal trawling on marine sedimentary habitats analysed by sediment profile imagery. *J. Exp. Mar. Biol. Ecol.*, 285–286, 465–477. [https://doi.org/10.1016/S0022-0981\(02\)00577-4](https://doi.org/10.1016/S0022-0981(02)00577-4), 2003
- Sciberras, M., Hiddink, J. G., Jennings, S., Szostek, C. L., Hughes, K. M., Kneafsey, B., Clarke, L. J., Ellis, N., Rijnsdorp, A. D., McConnaughey, R. A., Hilborn, R., Collie, J. S., Pitcher, C. R., Amoroso, R. O., Parma, A. M., Suuronen, P., &
- 625 Kaiser, M. J.: Response of benthic fauna to experimental bottom fishing: A global meta-analysis. *Fish. Fish.*, February, 1–18. <https://doi.org/10.1111/faf.12283>, 2018
- Sciberras, M., Parker, R., Powell, C., Robertson, C., Krüger, S., Bolam, S., & Geert Hiddink, J.: Impacts of bottom fishing on the sediment infaunal community and biogeochemistry of cohesive and non-cohesive sediments. *Limnol. Oceanogr.*, 61(6), 2076–2089. <https://doi.org/10.1002/lno.10354>, 2016
- 630 Seeberg-Elverfeldt, J., Schluter, M., Feseker, T., & Kolling, M.: Rhizon sampling of porewaters near the sediment-water interface of aquatic systems. *Limnol. Oceanogr.-Methods*, 3, 361–371. [https://doi.org/10.1016/S0012-821x\(02\)01064-6](https://doi.org/10.1016/S0012-821x(02)01064-6) Doi 10.1016/S0012-821x(02)01064-6, 2005
- Seiter, K., Hensen, C., and Zabel, M.: Benthic carbon mineralization on a global scale, *Global Biogeochem. Cycles*, 19, GB1010, doi:10.1029/2004GB002225, 2005
- 635 Shotbolt, L.: Pore water sampling from lake and estuary sediments using Rhizon samplers. *J. Paleolimnol.*, 44(2), 695–700. <https://doi.org/10.1007/s10933-008-9301-8>, 2010
- Soetaert, K., Herman, P. M. J., Middelburg, J. J., Heip, C., Smith, C. L., Tett, P., & Wild-Allen, K. Numerical modelling of the shelf break ecosystem: Reproducing benthic and pelagic measurements. *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 48(14–15), 3141–3177. [https://doi.org/10.1016/S0967-0645\(01\)00035-2](https://doi.org/10.1016/S0967-0645(01)00035-2), 2001
- 640 Soetaert, K., Van den Meersche, K., van Oevelen D.: *limSolve*: Solving linear inverse models. R package version 1.5.1., 2009
- Soetaert, M., De Haan, D., Verschueren, B., Decostere, A., Puvanendran, V., Saunders, J., Polet, H., & Chiers, K.: Atlantic Cod Show a Highly Variable Sensitivity to Electric-Induced Spinal Injuries. *Mar. Coast. Fish.*, 8(1), 412–424.

<https://doi.org/10.1080/19425120.2016.1180332>, 2016

645 Soetaert, Maarten, Chiers, K., Duchateau, L., Polet, H., Verschueren, B., & Decostere, A.: Determining the safety range of electrical pulses for two benthic invertebrates: brown shrimp (*Crangon crangon* L.) and ragworm (*Alitta virens* S.). *ICES J. Mar. Sci.*, 72(3), 973–980, 2015

Soetaert, Maarten, Decostere, A., Polet, H., Verschueren, B., & Chiers, K.: Electrotrawling: A promising alternative fishing technique warranting further exploration. *Fish. Fish.*, 16(1), 104–124. <https://doi.org/10.1111/faf.12047>, 2015

650 Stratmann, T., Soetaert, K., Wei, C. L., Lin, Y. S., & van Oevelen, D.: The SCOC database, a large, open, and global database with sediment community oxygen consumption rates. *Sci. Data*, 6(1), 242. <https://doi.org/10.1038/s41597-019-0259-3>, 2019

Tiano, J. C., Borger, E. De, Flynn, S. O., Cheng, C. H., Oevelen, D. Van, & Soetaert, K.: Physical and electrical disturbance experiments uncover potential bottom fishing impacts on benthic ecosystem functioning. *J. Exp. Mar. Biol. Ecol.*, 655 545(August), 151628. <https://doi.org/10.1016/j.jembe.2021.151628>, 2021

Tiano, J. C., van der Reijden, K. J., O'Flynn, S., Beauchard, O., van der Ree, S., van der Wees, J., Ysebaert, T., & Soetaert, K.: Experimental bottom trawling finds resilience in large-bodied infauna but vulnerability for epifauna and juveniles in the Frisian Front. *Mar. Environ. Res.*, 159(March), 104964. <https://doi.org/10.1016/j.marenvres.2020.104964>, 2020

Tiano, J. C., Witbaard, R., Bergman, M. J. N., Rijswijk, P. Van, Tramper, A., Oevelen, D. Van, & Soetaert, K.: Acute 660 impacts of bottom trawl gears on benthic metabolism and nutrient cycling. *ICES J. Mar. Sci.*. <https://doi.org/10.1093/icesjms/fsz027>, 2019

Tillin, H. M., Hiddink, J. G., Jennings, S., & Kaiser, M. J.: Chronic bottom trawling alters the functional composition of benthic invertebrate communities on a sea-basin scale. *Mar. Ecol. Prog. Ser.*, 318, 31–45, 2006

van de Velde, S., Lancker, V. Van, Hidalgo-martinez, S., & Berelson, W. M.: Anthropogenic disturbance keeps the coastal 665 seafloor biogeochemistry in a transient state. , December 2017, 1–10. <https://doi.org/10.1038/s41598-018-23925-y>, 2018

van Denderen, P. D. Van, Bolam, S. G., Hiddink, J. G., Jennings, S., Kenny, A., Rijnsdorp, A. D., and van Kooten, T.: Similar effects of bottom trawling and natural disturbance on composition and function of benthic communities across habitats. *Mar. Ecol. Prog. Ser.* 541, 31–43. <https://doi.org/10.3354/meps11550>, 2015

Van Hoey, G., Guilini, K., Rabaut, M., Vincx, M., & Degraer, S.: Ecological implications of the presence of the tube- 670 building polychaete *Lanice conchilega* on soft-bottom benthic ecosystems. *Mar. Biol.*, 154(6), 1009–1019. <https://doi.org/10.1007/s00227-008-0992-1>, 2008

van Marlen, B., de Haan, D., van Gool, A., & Burggraaf, D.: The effect of pulse stimulation on biota - Research in relation to

ICES advice - Progress report with preliminary results, Institute for Marine Resources & Ecosystem Studies (IMARES), Report number C103/09, Wageningen, Netherlands, 53 pp., 2009

- 675 Van Raaphorst, W., Kloosterhuis, H. T., Berghuis, E. M., Gieles, A. J. M., Malschaert, J. F. P., & Van Noort, G. J.: Nitrogen cycling in two types of sediments of the Southern North sea (Frisian front, broad fourteens): field data and mesocosm results. *Neth. J. Sea Res.* 28(4), 293–316. [https://doi.org/10.1016/0077-7579\(92\)90033-B](https://doi.org/10.1016/0077-7579(92)90033-B), 1992




Figure 1: Geographic location of the study site (left). Visual descriptions for experimental plots in Pulse-ON (A-C), Pulse-OFF (D-F), and Tickler (G-I) trawl treatments, untrawled adjacent stations (AD1, AD2, AD3), sediment profile imagery (SPI) locations, and box core sampling locations (right). Experimental plots were sectioned into northwest, central and southeast stations.







Figure 2: Water column turbidity during the time of trawling for Pulse-ON plots (a-c), Pulse-OFF plot (d) and Tickler plots (e-f).



690 **Figure 3: Examples of sediment profile imagery from the study showing a dense *Lanice conchilega* tubeworm reef (a), bioirrigation-induced sediment oxidation (b) and evidence of trawl disturbance: flattened tubeworms and sediment (c).**




695 **Figure 4: Particle size distributions for the surface sediments (top 1 cm) in experimentally trawled samples. Solid lines represent the distribution of sediment size fractions in Pulse-ON, Pulse-OFF, and Tickler, plots before trawling (T0) and dashed lines represent distributions after trawl disturbances (T1).**



700

Figure 5: Oxygen consumption (a) and macrobenthos densities (b) measured from treatment plots before (T0) and after (T1) fishing. Y-axes are on a log-scale.



705 **Figure 6: Robust regression analysis of sediment community oxygen consumption (SCOC; $\text{mmol m}^{-2} \text{d}^{-1}$) versus *Lanice conchilega* biomass (grams of wet weight m^{-2}) (top) and densities of other taxa versus *L. conchilega* densities (individuals m^{-2}) (bottom) in pre-trawled and trawled sediments. Note x-axis scales are different.**

Tables

Table 1: Results for sediment and biogeochemical parameters between T0-T1 for Tickler, Pulse-ON, and Pulse-OFF treatments (mean \pm SD). Nitrogen (N) and phosphorus (P) removal are presented as the percentage of inorganic N or P removed from the sediments after being formed from organic matter mineralization.

	Tickler				Pulse-ON				Pulse-OFF			
	T0	n	T1	n	T0	n	T1	n	T0	n	T1	n
<i>Sediment: very fine sand (%), chlorophyll a ($\mu\text{g g}^{-1}$), silt (%), median grain size (D50: μm)</i>												
Very fine sand	11 \pm 4.3	64	9.2\pm4.5**	65	8.3 \pm 4.2	53	7.4 \pm 4.2	64	11 \pm 3.4	55	11 \pm 2.8	63
Chl <i>a</i>	19 \pm 31	64	8.4\pm17***	65	10 \pm 8.9	53	9.1\pm16**	64	12 \pm 16	55	19 \pm 30	63
Silt	13 \pm 17	64	11 \pm 14	65	8.4 \pm 6.6	53	7.7\pm12**	64	8.4 \pm 9.1	55	11 \pm 16	63
D50	189 \pm 61	64	207 \pm 72	65	211 \pm 42	53	220 \pm 63	64	192 \pm 29	55	181 \pm 44	63
<i>Sediment profile imagery (SPI): prism penetration (mm), anoxic surface area (%)</i>												
Penetration	51 \pm 23	26	67 \pm 41	22	48 \pm 18	23	49 \pm 12	19	53 \pm 20	23	54 \pm 17	21
Anoxic surface	24 \pm 23	26	16\pm24***	22	3.8 \pm 6.1	23	5.0 \pm 10	19	22 \pm 18	23	27 \pm 28	21
<i>Porewater nutrients (mmol m⁻³)</i>												
NH ₄ ⁺	26 \pm 17	35	47\pm38**	35	74 \pm 59	34	81 \pm 59	36	92 \pm 83	36	80 \pm 61	36
NO ₃ ⁻	4.1 \pm 2.7	35	6.4\pm4.1**	35	14 \pm 20	34	6.1 \pm 8.2	36	4.9 \pm 7.3	36	6.8\pm4.9**	36
PO ₄ ³⁻	7.6 \pm 5.8	35	8.7 \pm 9.0	35	5.2 \pm 6.7	34	8.2\pm7.4**	36	9.4 \pm 15.8	36	13\pm9.9***	36
Si(OH) ₄	69 \pm 52	35	105 \pm 95	35	91 \pm 93	34	98 \pm 72	36	116 \pm 89	36	137 \pm 89	36
<i>Nutrient Fluxes (mmol m⁻²d⁻¹)</i>												
O ₂	-172 \pm 109	9	-74\pm40***	9	-144 \pm 199	9	-77 \pm 89	9	-72 \pm 35	9	-73 \pm 44	9
NH ₄ ⁺	1.3 \pm 1.3	9	1.6 \pm 1.6	9	1.1 \pm 2.1	9	0.6 \pm 0.3	9	0.6 \pm 0.8	9	0.5 \pm 0.3	9
NO ₃ ⁻	0.01 \pm 0.01	9	7e-3 \pm 0.01	9	1e-3 \pm 0.02	9	0.02 \pm 0.02	9	6e-3 \pm 0.02	9	9e-3 \pm 8e-3	9
PO ₄ ³⁻	0.03 \pm 0.02	9	0.04 \pm 0.02	9	0.01 \pm 0.03	9	0.03 \pm 0.03	9	0.05 \pm 0.07	9	0.03 \pm 0.03	9
Si(OH) ₄	1.1 \pm 0.95	9	1.8 \pm 0.76	9	0.75 \pm 0.63	9	1.1 \pm 0.90	9	1.4 \pm 1.2	9	1.1 \pm 0.70	9
<i>Mass Budgets: total mineralization and denitrification (mmol C m⁻²d⁻¹), nitrification (mmol N m⁻²d⁻¹), nitrogen (N) and phosphorus (P) removal (%)</i>												
Mineralization	156 \pm 98	9	68\pm36**	9	130 \pm 180	9	70 \pm 80	9	65 \pm 32	9	66 \pm 40	9
Denitrification	28 \pm 19	9	11\pm6.8*	9	23 \pm 31	9	12 \pm 14	9	11 \pm 5.6	9	12 \pm 7.5	9
Nitrification	22 \pm 16	9	8.6\pm5.5*	9	18 \pm 25	9	10 \pm 12	9	9.2 \pm 4.4	9	9.5 \pm 6.0	9
% N removal	86 \pm 24	9	82 \pm 16	9	95 \pm 5.6	9	89 \pm 9.5	9	94 \pm 6.4	9	93 \pm 5.1	9
% P removal	96 \pm 5.1	9	92 \pm 4.6	9	99 \pm 2.6	9	92 \pm 8.1	9	93 \pm 12	9	94 \pm 5.3	9

Bold signifies that T1 is significantly different compared to T0. *p < 0.05; **p < 0.01; ***p < 0.001.

Table 2: Robust regression coefficients (slope, p-value, multiple R-squared) showing the relationship of *Lanice conchilega* densities and biomass versus sedimentary faunal and biogeochemical flux parameters in trawled and untrawled sediments

PARAMETERS		<i>Pre-trawled</i>			<i>Trawled</i>			
		<i>slope</i>	<i>p</i>	R ²	<i>slope</i>	<i>p</i>	R ²	
<i>L. conchilega</i> DENSITIES	Sediment	Median grain size	-0.0012	0.29	0.07	-0.0068	0.18	0.11
		% Very fine sand	0.0002	0.01	0.33	0.0005	0.16	0.12
		% Silt	0.0003	0.25	0.08	0.0015	0.14	0.13
	Fauna	Density of other taxa	0.460	<0.001	0.85	0.1030	0.07	0.05
		Species richness	0.0003	0.005	0.27	0.0004	0.13	0.09
	Fluxes	SCOC	0.021	<0.001	0.87	0.0020	0.78	0.01
NO ₃ flux		-3.2e-7	0.65	0.01	-8.8e-8	0.89	6.9e-4	
<i>L. conchilega</i> BIOMASS	Sediment	Median grain size	0.0053	0.78	0.00	-0.163	0.17	0.11
		% Very fine sand	0.0022	0.27	0.09	0.010	0.19	0.11
		% Silt	0.0056	0.009	0.18	0.044	0.054	0.21
	Fauna	Density of other taxa	1.262	<0.001	0.39	-0.571	0.25	0.01
		Species richness	0.0166	<0.001	0.38	0.003	0.34	0.15
	Fluxes	SCOC	0.427	<0.001	0.93	0.056	0.75	0.01
NO ₃ flux		<0.0001	0.002	0.21	<-0.0001	0.82	0.01	

Bolded values show significant relationships