

## Response to Reviewer 1

The article “Gap geometry, seasonality and associated losses of biomass – combining UAV imagery and field data from a Central Amazon forest” studies gap formation on an 18ha field plot in the Amazon, using both remote sensing (photogrammetry/Structure from Motion) and field data. It provides an interesting look into canopy dynamics at one particular tropical forest site and a comparison (or validation) between field-based methods and remote sensing, which is crucial in linking traditional approaches with modern technology. Due to its substantial field sampling effort, the study can relate gap formation to different tree mortality modes and to associated biomass losses, thus linking ecological processes to the carbon cycle, which should be of great interest to readers of Biogeosciences. I also found the paper generally very well written, with well thought-through methods and clear and concise descriptions.

There are, however, a few changes/issues that I would recommend the authors to consider before publication. I will highlight a few larger aspects first, and then provide line-by-line comments in a classic review style.

*We thank Reviewer 1 for this supportive revision and constructive comments. We will consider all aspects for preparing our revised manuscript. Please, find below a point-by-point answer to all questions.*

**1/ Definition of gap:** My impression is that the definition of gaps is not entirely consistent in the study. On the one hand, Brokaw’s definition of gaps as extending down to 2m in canopy height seems to be used (l.161), but on the other hand, the authors argue several times that there are undetected “understory gaps”, or gaps that are not visible in the upper canopy. Specifically, they attribute the differences between UAV imagery and field data to the UAV imagery not being able to detect such subtle changes below the canopy. But if we use the Brokaw definition, that should not be the case, as any gap would, by necessity, be a hole in the upper canopy and extend down to the ground, no? Could it be that the authors implicitly use treefall events or other canopy characteristics as part of their gap definition in their field-based studies? Could this also explain why gaps created by standing dead trees were the main difference? The definition aspect also affects what should be considered the “truth” for the validation – field-based assessments certainly offer more information to interpret gap formation (is it a branch fall or a tree fall? etc.), but to automatically consider them the truth (l.209) is not evident to me. Could one not argue that the 3D canopy height models derived from photogrammetry (or even better, lidar) can more accurately quantify height changes than visual/manual assessments?

*As suggest by Reviewer 1, we will clarify the definitions of gap and revise the text for consistence.*

*We used the definition by Brokaw (i.e., gap in the forest canopy extending from the upper stratum to an average height of two meters above ground) to compute and measure gaps in the field (Brokaw, 1982). This is a classical and efficient method, which allows comparing our findings with those from fundamental work conducted in other tropical forests.*

*Gaps were identified through detailed inspection of dead trees and fallen branches. For the identification of mortality events and description of mortality modes, we followed the protocol of previous studies developed in our study region (Marra et al. 2014, 2018; Negron-Juarez et al. 2011; Ribeiro et al. 2016). To minimize errors, the inspections were conducted across relatively narrow strips of forest (less than 10 m), and were always led by ASLP. A field campaign was undertaken to quantify and mark all pre-existing gaps (i.e., baseline).*

*Our remote-sensing approach provides detailed data on the upper canopy of the forest, but no information on the understory. Gaps were defined as disturbed patches with total area  $> 5 \text{ m}^2$  and with reductions of canopy height greater than 10 m. These thresholds were established based on the nominal resolution of our processed imagery (1 m) and the scale at which the forest inventories were conducted, i.e., tree level. Overall, the fall of branches and/or standing dead trees produced severe damage mostly in the upper canopy while the understory remained intact. Therefore, upper canopy gaps detected remotely were not always detected on the ground using the definition by Brokaw. This pattern shows that apart from covering relatively large areas at low costs, UAV photogrammetry is an efficient method for monitoring gap dynamics - including the detection of upper canopy disturbance usually not visible from the forest ground.*

*We believe our approach combining field with remote sensing data provides interesting insights on concepts and methods for quantifying gaps and their effects on forest dynamics. Classical methods based on field observations are efficient to detect gaps extending from the upper canopy to the understory of the forest, and are crucial for validating remote tools, and for quantifying and modeling associated losses of biomass. High-resolution photogrammetry allows for more precise measurements of the features of the gaps, including those restricted to upper canopy and/or causing minor damage. Although more frequent, small-scale disturbances not implying tree mortality such as defoliation and branch fall are often neglected in forest inventories. Our study brings novel knowledge on the contribution of these events to ecosystem processes such as carbon cycle. Quantifying the size-distribution of gaps and their landscape importance is crucial to understand how forests respond to shifts in the disturbance regimes triggered by climate change and land use.*


**2/ Study area size and gap size frequency distributions (GSFD):** Having such a detailed comparison between field based and remotely sensed gap structure is an important feat, based on substantial field work, so it makes sense that the authors focused on a plot size of “only” 18ha. However, this limits the analysis somewhat when it comes to assessing GSFD and the “landscape scale” patterns the authors are interested in. As expected for 18ha, sample sizes are very small (32 gaps in total, but only 14 gaps that co-occur in both field and remote sensing data). I am sceptical that such sample sizes yield much information on which distribution actually fits better, and I would expect the fitted Weibull, exponential and power law distributions to be so uncertain in their parameters (the power law exponent has an uncertainty of  $2.137 \pm 0.913$ , which is huge) that there is not much sense in comparing the fit of different GSFDs (one single data point might already shift the goodness of fit). If the authors would like to keep this analysis, I suggest they explicitly use confidence intervals / simulations of data generation to assess how reliably these distributions can actually be differentiated with so few gaps, or maybe focus less on which distribution fits better and more on the field-remote sensing comparison. They should also provide a careful discussion that does not place too much emphasis on the different AIC values (which have generally low delta, anyways). More generally, if this type of analysis is carried out, I would also highly recommend the additional fitting of a lognormal distribution, which comes about through similar generative processes as power law models and is often an equally good fit.

We thank the Reviewer 1 for this valuable comment. We used a bootstrap with 1000 interactions for calculating the confidence interval of the different fits and will include the information in the revised figure. We also tested lognormal as part of our analyses.

```
bsfitsizedistcontdata <-
function(sizedf,minsize=c(10,25),maxsize=c(720,900,NA),fitfcn=c("exp","pow","weib","logn"),
nbootstraps=1000,alpha1=0.05,alpha2=0.01)
```

# nbootstraps is the number of bootstrap interactions to run to get CIs

We also used the Kolmogorov-Smirnov statistic to evaluate the goodness of fit (not only the AIC). We also tested lognormal as part of our analyses. The lognormal distribution did not fit for values greater than 25 m<sup>2</sup> in the UAV data, and the K-S statistic was also adequate (these are the largest values for a good fit). As requested by the reviewer, we propose to replace the following Figure S2 and Table S2:



**Figure S2.** Size distribution of gaps at the INVENTA plot, Central Amazon, Brazil, for the period from September 18<sup>th</sup>, 2018 to January 19<sup>th</sup>, 2021. Size distribution of gaps detected with UAV photogrammetry (a) and field surveys (c). Modeled distribution of gaps larger than 10 m<sup>2</sup> detected with UAV imagery (b) and field data (d).

**Table 1.** Summary of fitting measures of the exponential, Power-law, Weibull and Lognormal functions describing the size distribution of gaps identified on the INVENTA plot, Central Amazon, Brazil.

Detection method	Minimum size (m <sup>2</sup> )	Distribution	$\lambda$ (95 % CI)	$\alpha$ (95 % CI)	K-S	Log likelihood	$\Delta$ AIC
------------------	--------------------------------	--------------	---------------------	--------------------	-----	----------------	--------------

UAV imagery	10	Exponential	0.014 (0.008 - 0.031)		0.2489	-152.989	9.888
	10	Power-law	1.650 (1.529 - 1.831)		0.2242	-152.035	7.981
	10	Weibull	0.512 (0.266 - 1.460)	21.544 (0.770 - 65.311)	0.1166	-148.08	2.077
	<b>10</b>	<b>Lognormal</b>	<b>3.551 (2.137 - 3.995)</b>		<b>0.8466</b>	<b>-147.04</b>	<b>0</b>
	25	Exponential	0.013 (0.007 - 0.032)		0.2879	-117.622	13.41204
<b>25</b>	<b>Power</b>	<b>2.094 (1.901 - 2.501)</b>		<b>0.0760</b>	<b>-110.92</b>	<b>0</b>	
25	Weibull	0.157 (0.078 - 1.685)	0.0002 (0.0000 - 62.147)	0.0868	-110.747	1.661607	
25	Lognormal	-	-	-	-	-	
Field data	9	Exponential	0.017 (0.009 - 0.032)		0.1658	-146.783	4.9063
	9	Power-law	1.614 (1.494 - 1.766)		0.2659	-153.006	17.3533
	9	Weibull	0.744 (0.487 - 2.236)	40.370 (17.893 - 68.309)	0.1463	-145.61	4.5635
	<b>9</b>	<b>Lognormal</b>	<b>3.742 (3.3510 - 4.0823)</b>		<b>0.8368</b>	<b>-143.33</b>	<b>0</b>
	25	Exponential	0.017 (0.008 - 0.0418)		0.2147	-111.135	4.5447
	25	Power-law	2.137 (1.797 - 2.710)		0.1578	-109.306	0.8869
	25	Weibull	0.414 (0.131 - 2.982)	4.703 (0.000 - 68.937)	0.1437	-108.28	0.8305
	25	Lognormal	3.3953 (- 17.2998 - 4.1895)		0.9996	-107.86	0

*To our knowledge, this is the first study merging field and remote sensing data for assessing the geometry of gaps and computing their contribution to processes regulating carbon cycle in Amazon. Our study region is covered with old-growth terra-firme forests including a topographic/edaphic gradient comprised of plateaus, slopes and valleys. Our 18-ha plot is monitored since the year 2000; all the trees >10 cm DBH are recorded and tagged (~10.500 individuals). Apart from relying on this long-term inventory, our study also used LiDAR and rainfall data available for the same plot. Flying a contiguous area allowed us to reduce costs and optimized the logistics of field campaigns, which was crucial for the success of a 28-months monitoring.*

*Our relatively large plot allowed for a robust quantification of tree mortality and associated losses of biomass across an environmental gradient. A total of 32 gaps were formed during the entire studied period; 18 were formed during a 14-month period for which field and remote data were acquired. However, our results shall not be extrapolated beyond our study region. A regional assessment of the size distribution and geometry patterns of gaps require the inclusion of further sites reflecting existing variations in forest attributes and disturbance regimes. We will improve the discussion on this aspect in the revised version of the manuscript.*


**3/ Precipitation and gap formation:** This part of the paper, while relevant, is not really motivated in the introduction, and more effort should be spent on explaining why it is relevant to suppose a link between precipitation and gap formation, and why presumably more direct drivers of gap formation (wind or even lightning) were not used. It is understandable that such data may not be available, but nothing in the introduction/methods section explains why precipitation is interesting. I would also remove the analysis of extreme rainfall events, because this seems like a filtering of the data that could be done with many thresholds (90<sup>th</sup> / 95<sup>th</sup> percentile, etc.), and with only 3 years and 8 data points for extreme rainfall (Figure 8), I doubt that the correlation the authors found tells us much about the system.

*We agree with the Reviewer 1 and will improve the text explaining the links between precipitation and gap formation. In our study region, extreme wind gusts and precipitation are important mechanisms of tree damage and mortality (Chambers et al. 2013, Magnabosco Marra et al. 2018, Negrón-Juárez et al. 2018, 2023). A study monitoring tree mortality over five decades in a Central Amazon forest found that trees died more frequently in wet months, even during drought years (Aleixo et al., 2019). A positive correlation between precipitation and tree mortality was also reported for our study area (Fontes et al., 2018). A regional study based on 12 years of satellite data found that major windthrows (visible on Landsat) in Central Amazon occurred more frequently between September and February, months characterized by heavy rainfall, than the rest of the year (Negrón-Juárez et al., 2017). These background studies support that a greater number of gaps can be expected during the rainy season. We will revise the introduction to clarify this aspect.*

*We also understand that a comprehensive assessment of the influence of precipitation on gap formation requires longer-term data addressing seasonal and interannual variability. While this aspect could not be fully investigated with our 28-month dataset, it is part of an ongoing study which will incorporate a larger monitoring period and thus a higher number of extreme rainfall events and potential associated gaps. We will also aggregate background information on this aspect in the introduction and point out the limitations of our assessment. In addition, we suggest moving Figure 8 to the Supplementary Material.*

**4/ Remote sensing vs. field data in assessing mechanisms of gap formation and biomass loss:** My impression was that section 3.3 would be one of the most interesting sections for readers of Biogeosciences, and that the authors could extend their analysis here a little bit without too much effort. For example, I would relate released biomass to overall plot biomass. There could also be an interesting comparison of released biomass visible from gaps, to overall biomass released from tree mortality, also counting understory mortality (if these data exist). Finally, since they have such a comprehensive data set, the authors could also compare other aspects of gaps between the different mortality modes (branch fall, snapped, etc.). I would suggest a look at the metrics the authors already calculated (gap geometry), but also previous and surrounding canopy height, and maybe also gap closure rates, with a focus on the values from remote sensing. Maybe, the authors could also use the RGB signature of the orthophotos as an additional metric to compare between

mortality modes. Such an analysis would provide some hints on whether remote sensing/photogrammetry could distinguish different modes of mortality/gap formation/biomass losses, or at least separate one specific mode (standing dead). These are only suggestions and would, of course, only be indicative due to the small



sample sizes, but I think they might be very interesting for future studies/Biogeosciences readers and be in line with the authors' objectives to assess how much we can learn from remote sensing compared to field-based assessments.

*Biomass estimations are indeed an important aspect of our study, and we thank the Reviewer 1 for stimulating this discussion. As suggested, we did calculations to compare the biomass released in gaps vs. the stocks of old-growth forests in our study region. Apart from discussing these numbers, we will include a new figure in the Supplement (Fig. S3) to show the size distribution of gaps formed by branch fall and the different modes of tree mortality.*

**Figure S3.** Size distribution of gaps formed by branch fall and different modes of tree mortality.

*We agree that our RGB chronosequence can support research beyond the scope of this manuscript. For instance, we are investigating the geometric and reflectance signature of gaps created by branch fall and modes of tree mortality (i.e., standing dead, snapping and uprooting). This investigation may contribute to more reliable assessments of tree mortality events related to opposing climate extremes such as storms and droughts.*

*Our sample unit in this study is a gap event, which is not dependent on the size of the plot or subplot size. The 18-ha plot was selected because (i) it is covered with an old-growth forest; (ii) includes a topographic/edaphic gradient (see answer to the second comment of Reviewer 2); (iii) has an infrastructure of trails and (iv) forest inventory, LiDAR and rainfall data.*

**Line-by-line comment:**

3: Is the title actually accurate? Gap geometry and seasonality do not seem to be such important results/aspects of this study, so maybe rethink/rephrase it?

*We agree and suggest the following title: "Gap attributes and associated losses of biomass - combining UAV imagery and field data from a Central Amazon forest".*

37: What is a multi-temporal process? Maybe rephrase?

*We suggest rephrasing as following: "The maintenance of these stocks depends on dynamic processes that regulate the growth and mortality of trees"*

41: Even though this may not be fully relevant to the paper, maybe droughts could be mentioned as another major extreme event?

*We agree that droughts are another extreme event that can influence gap dynamics by causing water stress and consequently increasing the frequency of branch fall and standing dead trees. We will include a comment on that in the revised version of the manuscript.*

50: This may be a definition question and not crucial, but in the context of tropical forests, gaps that are thousands of hectares in size (or tens of squarekilometers) seem unlikely, or probably not what tropical ecologists would commonly classify as gaps (e.g. one or several large canopy trees falling and leaving a gap in the canopy). Such a definition seems more common in fire-dominated boreal ecosystems. Maybe you could add one sentence specifically on tropical gap sizes. Also, this would be more in line with the extent of your sample plot.

*We thank Reviewer 1 for this important comment regarding the definition of gap size. This scenario of giant gaps is also common in the Amazon. Although with some surviving trees, gaps opened by extreme rain and wind in Amazonia can have a total area greater than 3,000 hectares (Nelson et al., 1994; Espírito-Santo et al., 2014; Negrón-Juárez et al., 2010, 2018, 2023). Recent evidences suggest hotspots for the occurrence of blowdowns (Negrón-Juárez et al., 2023; Urquiza Muñoz et al., 2021) and that the frequency of these events is regulated by atmospheric phenomena that are highly sensitive to climate change, such as the potential energy available for convection (CAPE) (Feng et al. 2023). We will include information on that aspect in our revised manuscript.*

62-74: My impression is that this part of the paper jumps quite a lot between points, i.e. from the advantages of remote sensing, citing lidar remote sensing studies such as Dalagnol et al. 2021, to different definitions of gaps, to the problems of optical remote sensing. My question would be: Is the discussion of Landsat needed here, as UAV operates on a very different scale. A more interesting point might be how UAV photogrammetry differs from ALS/UAV lidar (e.g. no within-canopy structure, no ground model, but likely cheaper, more flexible [although limited by meteorological conditions]).

*Our goal here was to discuss how UAV imagery could expand on previous assessments of windthrow tree-mortality, which up to date is mainly based on Landsat imagery (Nelson et al., 1994; Espírito-Santo et al., 2014; Negrón-Juárez et al., 2010, 2018, 2023). Although allowing for long-term studies at the regional scale, Landsat is only sensible for detecting relatively large gaps (>1,000 m<sup>2</sup>), which in Central Amazon usually involve the death of more than eight trees (Negrón-Juárez et al., 2011; Chambers et al., 2013).*

90: what are "traceable" modes of tree mortality? Or what would be "untraceable" ones?

*We will rewrite the sentence for clarification. Recurrent field surveys following the routine and protocol established in previous studies (Magnabosco Marra et al., 2014, 2018; Ribeiro et al., 2016; Fontes et al., 2018) allowed us to clearly distinguish branch fall, standing dead, snapped and uprooted trees. For example, a standing dead tree may lose part or all of its crown shortly after dying. Thus, when field surveys are not conducted frequently, it is not possible to distinguish*

*the mechanism of gap formation. A snapped tree is characterized by mechanical disruption of the stem (breakage or cracking). If checked in the field within a short period of time after its occurrence, it is still possible to find leaves attached the crown, thin wood/fibers at the snapping point/height and sap (e.g., resins and latex). Uprooted trees have exposed roots usually still attached to the trunk/crown. Branch fall gaps are formed when branches of living trees are broken and/or fall after dying. We will edit the correspondent text for clarification.*

90: The last question with regard to rainfall is very adhoc and not really set up in the introduction. I would provide justification in the introduction on why precipitation should be relevant for gap formation. Would wind be a more important variable?

*Thunderstorms may propagate extreme wind gusts and rain. For instance, Araujo et al. (2021) found a significant relationship between gap formation and precipitation (extreme rainfall events, i.e. the rainfall rate in  $\text{mm h}^{-1}$  of the 98.2th). We agree with Reviewer 1 and propose including more information to support why we expect precipitation to be an important aspect regulating gap dynamics and geometry.*

113-136: I am no expert in SfM/photogrammetry, but this seems well-described and a good workflow. I have one question: How did you deal with different meteorological conditions during planned flights (fog/rain)? Did you, for example, postpone scans during rainy days? Could this affect your results? How consistent was the timing of the acquisitions on average? I don't think this would be a major problem, but it would be good to mention this somewhere here.

*We thank the Reviewer 1 for this question. The flights were always carried around 09:00h and on the absence of fog and rain. The flights last approximately 15 minutes, which allowed us to acquire images under similar conditions of light. When possible, the flights were carried at cloudy conditions and diffuse light, which improves the visibility of the canopy while reducing shadow. We will add these details in the Supplement.*

155-172: This seems like a substantial effort and great, important work! Just out of interest: since you seem to have access to EBA project's overlapping lidar data, is there a reason why you did not predelineate initial gap distributions from the lidar derived canopy height models?

*The LiDAR data was acquired in 2016. Although at a higher spatial resolution, these data could not be used as a reliable baseline for our study, which started approximately 2.5 years later. Instead, we carried a detailed field survey to collect the coordinates and mark all existing gaps with a plastic-colored stick.*

197-209: This also makes a lot of sense. However, I would move the information from the last sentence (i.e. field value is considered true value) to the beginning to make it clearer to the readers what is considered the validation. I was wondering, however, whether in this case field data can actually be considered the true data? One could make a point that remote sensing (but maybe less so photogrammetry) actually provides a more accurate quantification of the 3D canopy canopy than visual/field-based assessments can. How would you justify your decision?

*We agree with Reviewer 1 and will carry the suggested change. The Brokaw definition was used to identify and describe gaps in the field. Apart from allowing comparison with previous studies, this method is compatible with our measurements of forest structure and losses of biomass. We also used the field data to validate the occurrence of gaps identified and measured remotely.*

215-223: While it is common to fit these distributions and the approach is methodologically sound, does this make sense here? 18ha is a very small area when

it comes to gap delineation, so even without looking at the results, one would assume that your sample size is going to be so low that the inferred distributions are not telling us a lot (and the results bear this out, with 32 or 14 gaps in total). At the very least, I would expect simulations to construct confidence/credibility intervals that show how much variability there is and how uncertain the differences between the different distribution types are. My guess is that it would be very hard to come to any clear conclusion across 18ha. Also, would it not make sense to also test a lognormal distribution? The lognormal distribution is usually the one closest to the power law and comes about through very similar generative processes, so if you fit distributions.

*See answers related to the comment 2.*

225-229: Very interesting! I find the idea of quantifying released biomass very appealing.

*We appreciate this positive comment and agree that the field data is a highlight of our study.*

232: This process of calculating gap area formation rates sounds very complicated. Could you not just take the number/area of gaps that formed between each image acquisition and then divide the number/area by the time between each image? Assuming that images are taken at roughly the same intervals, that should give you a very sound estimate, no? Or am I missing something?

*Apart from relating number of gaps per unit of area, we aimed at analyzing the correlation between precipitation (measured as accumulated rainfall) and gap frequency. We will improve the text for clarification.*

253: "which indicates that there was no traceable change in the upper canopy of the forest". This is probably more a discussion sentence anyways, but I find this problematic. According to the definition (Brokaw) you use, a gap is an "an opening in the forest canopy extending from the upper stratum to an average height of two meters above ground." So by definition something in the upper canopy has to change – either you don't pick it up in the photogrammetry data (maybe one of the processing algorithms is smoothing the canopy too much), or, alternatively, your field-based assessment wrongly found a change in the upper canopy. This could also be an interesting question about gap definitions: should a standing dead tree already be classified as a gap, because light is reaching down almost without obstruction to 2m? How do you interpret this?

*We moved the text to the discussion. A gap easily identified in the field but not recognizable in the images could be one formed by the fall of lower branches from an emergent/canopy tree or the fall of a relatively small standing dead tree. These contribute relatively small area/volume. The respective changes could then be smoothed and gaps not detected. Since dense tropical forests have several vertical strata, the definition by Brokaw only fits when gaps extend to the upper canopy; this was not the case of relatively small damage promoted by branch fall. A standing dead tree was classified as gap when the light penetrated the understory of the forest leading to detectable changes in height/volume.*

260: I'm not sure the p-value is the best way to assess this here. Looking at Figure 3, one would guess that, at the large end of the gap spectrum, UAV seems to find larger gaps than field-based assessments (a difference of ca. 830 m<sup>2</sup> to 580m<sup>2</sup> for the largest gap seems substantial and larger than I would have expected). How did you

derive the p-value? Did you log-transform the data beforehand (if you assume power-law/lognormal scaling, for example, that would be necessary, I assume)

*We log transformed these variables before performing the paired t-test. In R, the procedure was performed as:*

```
## Filtering pairs that were captured gap ##
pairs <- subset(gap, classe_RS == 1 & classe_BRK == 1 , select=c("Area_RS_m2",
"Area_Brokaw_m2"))
## log-transform ##
log_area_RS <- log(pairs$Area_RS_m2)
log_area_BRK <- log(pairs$Area_Brokaw_m2)
log_dif <- log_area_RS - log_area_BRK
shapiro.test(log_dif)
## Paired t-test ##
t.test(log_area_BRK, log_area_RS, paired = TRUE, alternative = "two.sided")
```

285: My takeaway from Table 2 would actually be that all distributions perform similarly (the dAIC is typically very low), and my guess would be that, if you account for the uncertainty of the small sample size, you cannot really differentiate between any of them here. I would highly recommend to test this! One interesting question is whether the field data have a slightly different exponent/shape, with a steeper decline at the largest gap areas (in line with the visual assessment). But, of course, sample sizes are very low.

*We thank Reviewer 1 for this valuable comment. We agree that the interpretation of calculated uncertainties is limited and will make this aspect clearer in the discussion. We will also discuss the small differences of the fits for the field and remote approaches. The power-law fit had the steeper slope among other functions and, in comparison with our field data, indicated a lower frequency of gaps greater than 100 m.*

315: I like this idea of calculating the biomass loss, and that at least one branch fall exceeded some of the uprooted/snapped tree losses. Could you put this into context of how much total biomass is stocked in the plot? I.e. what percentage is lost by gaps?

*We thank the Reviewer 1 for this suggestion. The losses of biomass in our studied gaps ( $1.35 \text{ Mg ha}^{-1} \text{ year}^{-1}$ ) account for 0.88% of the stocks in an old-growth forest contiguous to our plot ( $355.67 \pm 34.53 \text{ Mg ha}^{-1}$  (mean  $\pm$  standard deviation) (Amaral et al., 2019).*

334: This is not my favourite figure (and analysis). There are very few data points, and while I understand the general reasoning, it seems a bit like one could also pick a different percentile of extreme rainfall events, and the pattern might disappear. I suggest you remove this Figure and analysis.

*As previously mentioned in our answer to the comment 3, we suggest moving this figure to the Supplement. We will also improve the discussion on these results and include a couple of sentences exploring the limitations of our dataset for addressing the effects of the precipitation on gap dynamics.*

351-353: As noted above (and sorry for the repetition), there seems an inconsistency in the gap definition in the paper. If gaps are defined as openings in the upper canopy that clearly reach down to 2m (Brokaw), it does not make much sense to me to say that there are “no clear signs of opening in the upper canopy”. Could it be that your field-based gap definition is slightly wider than the one you apply with the remote sensing/photogrammetry data, and is implicitly based around whether a tree has fallen? I am not saying that this is necessarily wrong, but that could explain divergences between both methods, because unless your photogrammetry approach overly smoothes the canopy, there is no a priori reason why it should not detect openings in the Brokaw sense, no? In this respect, I would also expect 2-3 sentences here on the problem of which of the two data sets (remote sensing or field) is the actual truth!

*We suggest editing the text as indicate below:*

*“Field data acquired by using the definition by Brokaw et al. 1982 was considered as ground truth. Apart from simple and precise, these data allow comparing our findings with those from fundamental work conducted in other tropical forests. The Brokaw's definition has been used in ecological studies of gaps and to evaluate it with more recent remote-sensing tools (such as UAV photogrammetry) was one of the goals from our this work. To accomplish with this goal, we used a confusion matrix for assessing the accuracy of our remote method of gap detection. Further, we calculated the percentiles of accuracy (a), precision (p), recall (r), and F1 Score (F) (Eqs. 1 – 4) (Dalagnol et al., 2021), where TP is true positive, TN is true negative, FP is false positive and FN is false negative:*

$$\text{Accuracy (a)} = ((TP+TN) /n) *100 \quad (1)$$

$$\text{Precision (p)} = (TP/(TP+FP)) *100 \quad (2)$$

$$\text{Recall (r)} = (TP/(TP+FN)) *100 \quad (3)$$

$$\text{Score F1 (F)} = (((2*p*r) / (p + r))) *100 \quad (4)$$

*The total number of correct detections is expressed as percentile. The p percentile indicates the ratio of positive predictions performed correctly based on all positive predictions (including false ones). The r percentile is used to access the ratio of correct positive-predictions in relation to all positive predictions. The F1 Score (F) is the harmonic mean between p and r, i.e., the mean between the errors of commission and omission; higher F-values indicate higher agreement between gaps identified in the imagery data and observed in the field (ground truth).”*

358: Unless I have missed it, I am not sure that the study shows how much gaps contribute to landscape patterns of biomass. It would help to put the losses into context of the whole-plot biomass stocks (cf. above), but I would still be wary of calling this “landscape” patterns. 18ha is probably not on the scale where landscape effects can be assessed, particularly, because power law-type distributions imply that you will have very few, very large gaps, and your plot may just accidentally miss out on extreme events / the long tails of the GSFD distribution (blowdowns/multiple emergent/canopy trees falling).

*As previously suggested, we will express rates of biomass losses in terms of area. We will also improve the explanation on the characteristics of our plot and its potential to integrate part of the landscape heterogeneity typical of our study region.*

362: again, what is an “understory gap”?

*We used this term for describing gaps restricted to the forest understory and that could not be in our UAV data. Meanwhile, our UAV data enable for the detection of height reductions of 10 m,*

*which allowed us to detect small gaps restricted to the higher portions of the canopy. We will edit for clarification.*

379: That the area of the gaps did not vary between methods is not entirely correct (cf. my comments above on this particular p-value), and even if we were to solely rely on the p-value, I would rephrase to say there was no evidence for strong variation in the area between the two methods (although in my opinion, there is some, limited evidence for divergences between the two methods in terms of gap area).

*We will improve the discussion on this topic and point out the limitations of our results.*

389: Cf. my comments before. I don't think, we can conclude that power-law distribution is the best distribution here, cf. also the large confidence interval of  $2.137 \pm 0.913$ ! That is huge uncertainty!

*Thank you for the input regarding the uncertainty of the data. We will make this clear in the text and simplify the discussion regarding the distribution fits. Instead, we will focus on the differences between the remote and field detection.*

401: It seems to me that in many cases (not just your study), Weibull laws actually fit gap size frequency distributions better than power laws. I would discuss here what that would mean: it is more difficult to interpret (more parameters, not just one nice exponent), and it probably means that there is a change in generative mechanisms in gap formation across scales, which could make a lot of sense, because we probably shift from tree to branch level below a certain size threshold. You could also discuss this in the context of the typical tree size in your plot!

*Disturbances that are relatively small ( $>5 \text{ m}^2$ ) but generates height losses  $>10 \text{ m}$  were detected. This includes branch fall or crown damage not resulting on individual mortality. We will improve the discussion regarding the size distribution of gaps and the resolution of our imagery.*

430-437: This motivation for rainfall patterns – correlation with extreme winds or lighting – should come much earlier in the paper (ideally in the introduction), so that the reader understands why these patterns are studied.

*As previously mentioned in the third comment by Reviewer 1, we will provide further information on the motivation of our study.*

448-449: I fully agree with your statement that forest inventories are fundamental, but I am not sure you showed conclusively that “mechanisms of formation could only be distinguished using field data”. A very strong addition in my opinion (in Section 3.3.) would be to compare various attributes of the different gap types (branch/snapped dead/uprooted/standing dead), i.e. area/perimeter ratios, average height of lost canopy, average height within gap, canopy closure rate after gap formation (or even canopy changes before gap formation), and to really test whether there are indicators of how to differentiate different gap formation processes from remote sensing. My guess is, as you state, that you cannot reliably distinguish, for example, between uprooting and snapping, but it could still be that you would find a specific signature for standing dead trees (I imagine you could also use the RGB values of the orthophotos to identify them), or maybe you could find a

difference between branch falls and tree falls? This would be very interesting ecologically!

*We agree with this suggestion. However, this aspect is not within the scope of the present study. We are generating classifiers based on spectral, texture, and structural features of gap, and plan at presenting the related results in a next study.*

450-457: My impression is that the concluding paragraph is describing too many aspects at the same time (Weibull, mechanisms of gap formation, regional variation). Maybe focus on one priority, which seems to me the mechanisms/drivers of gap formation, and centre the paragraph around this notion?

*We thank the Reviewer 1 for this constructive comment. We propose to revise the Conclusions section as following:*

*By combining high temporal and spatial resolution UAV imagery with detailed field data on the mechanisms of gap formation, we could reliably assess the geometry and related losses of biomass for a closed-canopy Amazon forest. Although a larger proportion of canopy gaps could be detected from orthomosaics, their mechanisms of formation could only be distinguished using field data. Our results highlight that detailed forest inventories are fundamental for evaluating remote sensing products and metrics [linking ecological processes to the carbon cycle](#). Future studies are needed to generate proxies for distinguishing mechanisms of gap formation and their landscape importance.*

## References

Aleixo, I., Norris, D., Hemerik, L., Barbosa, A., Prata, E., Costa, F., and Poorter, L.: Amazonian rainforest tree mortality driven by climate and functional traits, *Nat. Clim. Chang.*, 9, 384–388, <https://doi.org/10.1038/s41558-019-0458-0>, 2019.

Amaral, M., Lima, A., Higuchi, F., dos Santos, J., and Higuchi, N.: Dynamics of Tropical Forest Twenty-Five Years after Experimental Logging in Central Amazon Mature Forest, *Forests*, 10, 89, <https://doi.org/10.3390/f10020089>, 2019.

Araujo, R. F., Grubinger, S., Celes, C. H. S., Negrón-Juárez, R. I., Garcia, M., Dandois, J. P., and Muller-Landau, H. C.: Strong temporal variation in treefall and branchfall rates in a tropical forest is related to extreme rainfall: Results from 5 years of monthly drone data for a 50 ha plot, *Biogeosciences*, 18, 6517–6531, <https://doi.org/10.5194/bg-18-6517-2021>, 2021.

Brokaw, N. V. L.: The Definition of Treefall Gap and Its Effect on Measures of Forest Dynamics, *Biotropica*, 14, 158, <https://doi.org/10.2307/2387750>, 1982.

Chambers, J. Q., Negrón-Juárez, R. I., Marra, D. M., Di Vittorio, A., Tews, J., Roberts, D., Ribeiro, G. H. P. M., Trumbore, S. E., and Higuchi, N.: The steady-state mosaic of disturbance and succession across an old-growth central Amazon forest landscape, *Proc. Natl. Acad. Sci. U. S. A.*, 110, 3949–3954, <https://doi.org/10.1073/pnas.1202894110>, 2013.

Espírito-Santo, F. D. B., Gloor, M., Keller, M., Malhi, Y., Saatchi, S., Nelson, B., Junior, R. C. O., Pereira, C., Lloyd, J., Frolking, S., Palace, M., Shimabukuro, Y. E., Duarte, V., Mendoza, A. M., López-González, G., Baker, T. R., Feldpausch, T. R., Brienen, R. J. W., Asner, G. P., Boyd, D. S., and Phillips, O. L.: Size and frequency of natural forest disturbances and the Amazon forest carbon balance, *Nat. Commun.*, 5, 1–6, <https://doi.org/10.1038/ncomms4434>, 2014.

Feng, Y., Negrón-Juárez, R. I., Romps, D. M., and Chambers, J. Q.: Amazon windthrow disturbances are likely to increase with storm frequency under global warming, *Nat. Commun.*, 14, 101, <https://doi.org/10.1038/s41467-022-35570-1>, 2023.

Fontes, C. G., Chambers, J. Q., and Higuchi, N.: Revealing the causes and temporal distribution of tree mortality in Central Amazonia, *For. Ecol. Manage.*, 424, 177–183, <https://doi.org/10.1016/j.foreco.2018.05.002>, 2018.

Magnabosco Marra, D., Chambers, J. Q., Higuchi, N., Trumbore, S. E., Ribeiro, G. H. P. M., Dos Santos, J., Negrón-Juárez, R. I., Reu, B., and Wirth, C.: Large-scale wind disturbances promote tree diversity in a Central Amazon forest, *PLoS One*, 9, <https://doi.org/10.1371/journal.pone.0103711>, 2014.

Magnabosco Marra, D., Trumbore, S. E., Higuchi, N., Ribeiro, G. H. P. M., Negrón-Juárez, R. I., Holzwarth, F., Rifai, S. W., dos Santos, J., Lima, A. J. N., Kinupp, V. F., Chambers, J. Q., and Wirth, C.: Windthrows control biomass patterns and functional composition of Amazon forests, *Glob. Chang. Biol.*, 24, 5867–5881, <https://doi.org/10.1111/gcb.14457>, 2018.

Negrón-Juárez, R., Magnabosco-Marra, D., Feng, Y., Urquiza-Muñoz, J. D., Riley, W. J., and Chambers, J. Q.: Windthrow characteristics and their regional association with rainfall, soil, and surface elevation in the Amazon, *Environ. Res. Lett.*, 18, 014030, <https://doi.org/10.1088/1748-9326/acaf10>, 2023.

Negrón-Juárez, R. I., Chambers, J. Q., Guimaraes, G., Zeng, H., Raupp, C. F. M., Marra, D. M., Ribeiro, G. H. P. M., Saatchi, S. S., Nelson, B. W., and Higuchi, N.: Widespread Amazon forest tree mortality from a single cross-basin squall line event, *Geophys. Res. Lett.*, 37, 1–5, <https://doi.org/10.1029/2010GL043733>, 2010.

Negrón-Juárez, R. I., Chambers, J. Q., Marra, D. M., Ribeiro, G. H. P. M., Rifai, S. W., Higuchi, N., and Roberts, D.: Detection of subpixel treefall gaps with Landsat imagery in Central Amazon forests, *Remote Sens. Environ.*, 115, 3322–3328, <https://doi.org/10.1016/j.rse.2011.07.015>, 2011.

Negrón-Juárez, R. I., Jenkins, H. S., Raupp, C. F. M., Riley, W. J., Kueppers, L. M., Marra, D. M., Ribeiro, G. H. P. M., Monteiro, M. T. F., Candido, L. A., Chambers, J. Q., and Higuchi, N.: Windthrow variability in central Amazonia, *Atmosphere (Basel)*, 8, 1–17, <https://doi.org/10.3390/atmos8020028>, 2017.

Negrón-Juárez, R. I., Holm, J. A., Marra, D. M., Rifai, S. W., Riley, W. J., Chambers, J. Q., Koven, C. D., Knox, R. G., McGroddy, M. E., Di Vittorio, A. V., Urquiza-Muñoz, J., Tello-Espinoza, R., Muñoz, W. A., Ribeiro, G. H. P. M., and Higuchi, N.: Vulnerability of Amazon forests to storm-driven tree mortality, *Environ. Res. Lett.*, 13, <https://doi.org/10.1088/1748-9326/aabe9f>, 2018.

Nelson, B. W., Kapos, V., Adams, J. B., Oliveira, W. J., and Braun, O. P. G.: Forest Disturbance by Large Blowdowns in the Brazilian Amazon, *Ecology*, 75, 853–858, <https://doi.org/10.2307/1941742>, 1994.

Ribeiro, G. H. P. M., Chambers, J. Q., Peterson, C. J., Trumbore, S. E., Marra, D. M., Wirth, C., Cannon, J. B., Negrón-Juárez, R. I., Lima, A. J. N., Paula, E. V. C. M. De, Santos, J., and Higuchi, N.: Forest Ecology and Management Mechanical vulnerability and resistance to snapping and uprooting for Central Amazon tree species, *For. Ecol. Manage.*, 380, 1–10, <https://doi.org/10.1016/j.foreco.2016.08.039>, 2016.

Urquiza Muñoz, J. D., Magnabosco Marra, D., Negrón-Juarez, R. I., Tello-Espinoza, R., Alegría-Muñoz, W., Pacheco-Gómez, T., Rifai, S. W., Chambers, J. Q., Jenkins, H. S., Brenning, A., and Trumbore, S. E.: Recovery of Forest Structure Following Large-Scale Windthrows in the Northwestern Amazon, *Forests*, 12, 667, <https://doi.org/10.3390/f12060667>, 2021.