

Effects of water table level and nitrogen deposition on methane and nitrous oxide emissions in an alpine peatland

Wantong Zhang^{1,2,4}, Zhengyi Hu², Joachim Audet⁴, Thomas A. Davidson⁴, Enze Kang^{1,3}, Xiaoming Kang^{1,3}, Yong Li^{1,3}, Xiaodong Zhang^{1,3}, Jinzhi Wang^{1,3, *}

¹Institute of Wetland Research, Chinese Academy of Forestry, Beijing Key Laboratory of Wetland Services and Restoration, Beijing 100091, China

²Sino-Danish Centre for Education and Research, University of Chinese Academy of Sciences, Beijing 100049, China

³Sichuan Zoige Wetland Ecosystem Research Station, Tibetan Autonomous Prefecture of Aba 624500, China

⁴Department of Ecoscience and Arctic Research Centre (ARC), Aarhus University, C.F. Møllers Allé, 8000 Aarhus, Denmark

*Correspondence to: Jinzhi Wang (wangjz04@126.com)

Abstract

Alpine peatlands are recognized as a major natural contributor to the budgets of atmospheric methane (CH₄) but as a weak nitrous oxide (N₂O) source. Anthropogenic activities and climate change have put these fragile nitrogen (N)-limited peatlands under pressure by altering water table (WT) levels and enhancing N deposition. The response of greenhouse gas (GHG) emissions from these peatlands to these changes is uncertain. To address this knowledge gap, we conducted a mesocosm experiment in 2018 and 2019 investigating individual and interactive effects of three WT levels (WT₋₃₀, 30 cm below soil surface; WT₀, 0 cm at the soil surface; WT₁₀, 10 cm above soil surface) and multiple levels of N deposition (0, 20, 40, 80 and 160 kg N ha⁻¹ yr⁻¹) on growing season CH₄ and N₂O emissions in the Zoige alpine peatland, Qinghai-Tibetan Plateau. We found that the elevated WT levels increased CH₄ emission, while N deposition had non-linear effects (with stimulation at moderate levels, but inhibition at higher levels). In contrast no clear pattern of the effect of WT levels on the cumulative N₂O emission was evident, while N deposition led to a consistent and linear increase (emission factor: 2.3%-2.8%), and this was dependent on the WT levels. Given the current N deposition in the Zoige alpine peatland (1.08-17.81 kg N·ha⁻¹), our results suggested that the CH₄ and N₂O emissions from the alpine peatlands could greatly increase in response to the possible doubling N deposition in the future. We believe that our results provide insights into how interactions between climate change and human disturbance will alter CH₄ and N₂O emissions from this globally important habitat.

Keywords Cumulative GHG emissions; multi-level N enrichment; critical threshold; non-linear effect; Qinghai-Tibetan Plateau

1. Introduction

Peatlands only cover ca. 3% of the land surface of the Earth but store one third of the global carbon pool (Yu et al., 2010). In pristine peatlands, the shallow water table (WT) and waterlogged conditions allow accumulation of organic matter and favour anaerobic production of methane (CH₄) and nitrous oxide (N₂O). Traditionally, this nitrogen-limited ecosystem is recognized as major CH₄ sources and weak N₂O sources (Frolking et al., 2011). Nevertheless, these conditions could be markedly changed by

anthropogenic disturbance and climate change, and growing evidence shows that peatlands are experiencing drainage and increasing nitrogen deposition (Chen et al., 2013; Evans et al., 2021). Consequently, the magnitude of CH₄ and N₂O emissions from peatlands may be severely altered, particularly the high-altitude or alpine peatlands that are especially vulnerable and highly sensitive to the climate change and anthropogenic activities (Squeo et al., 2006).

Large-scale artificial drainage of peatlands was initiated hundreds of years ago and escalated in the 20th century (Evans et al., 2021). As a result, about 10-20% of the global peatlands were primarily drained for the purposes of agriculture, peat extraction and forestry (Frolking et al., 2011). The resulting lower WT altered the anaerobic conditions of the peat soil and led to oxidative loss of peat (Laine et al., 2019; Wilson et al., 2016). Generally, the drainage decreased the CH₄ efflux and increased CO₂ and N₂O emissions (Cao et al., 2017). The increase in N₂O emissions from drained peatlands is often small, but may potentially reach a high level at sufficient nutrient input, especially when the soil is fertilized (Laine et al., 2019). Ecological restoration has been proposed as a measure to conserve the drained or degraded peatlands, particularly to meet the demand for mitigation of greenhouse gas (GHG) emissions outlined in the Paris Agreement (Evans et al., 2021). Numerous studies have reported a remarkably decreased CO₂ efflux in rewetted or restored peatlands, but the rising WT levels have also augmented the emissions of CH₄ and N₂O (Audet et al., 2013; Järveoja et al., 2016).

Atmospheric N deposition, primarily caused by anthropogenic activities (i.e. fossil fuels combustion, fertilizer application), has increased consistently during the past decades (Gomez-Casanovas et al., 2016, IPCC, 2013), and it is predicted to increase two- or three-fold in terrestrial ecosystems by the end of the century (Lamarque, 2005). The increasing N deposition could alleviate the N stress on peatlands, but the N effects on CH₄ and N₂O emissions are unclear (Deng et al., 2019). Thus, positive (Juutinen et al., 2018), negative (Gao et al., 2014) or neutral (Wang et al., 2017) effects of N deposition on CH₄ emissions in peatlands have been observed. We speculate that the contrasting results probably are a result of the prevailing environmental conditions and the N addition rate. Besides CH₄ emission, N deposition generally stimulates N₂O emissions from peatlands due to the increasing supply of N substrate (Wang et al., 2017). However, previous studies have also shown that a higher N input leads to a transition of the grassland into a state of declining N saturation as well as a reduction of the sensitivity of the GHG exchange to the continuously increasing N deposition (Gomez-Casanovas et al., 2016). To eliminate the possible gap resulting from the N addition rate, multiple levels of N deposition are required to study the possible linear or non-linear effects of deposition on GHG emissions.

Numerous studies have reported on the individual effects of WT and N deposition on GHG emissions in peatlands (Evans et al., 2021; Saiz et al., 2021). To our knowledge, only a few studies (Gao et al., 2014; Wang et al., 2017) exist that focus on their interactive effects on peatland GHG emissions. Gao et al. (2014) found that N addition in peatlands decreased CH₄ emissions but increased N₂O emissions without any significant interaction with WT levels. Wang et al. (2017) observed no interactive effects of a lower WT and increasing N deposition on GHG emissions in an alpine wetland. The above-mentioned studies were, however, limited to a single level of N addition and associated water addition. The response of GHG emissions in peatlands to the gradients of N deposition and WT levels remains to be elucidated, in particular at the N saturation stage, even though it may be a key factor

in shaping GHG emissions. The large uncertainties regarding the interactive effects of N deposition and WT levels on GHG emissions severely hamper obtaining a reliable estimation of the response of peatlands to climate change and anthropogenic activities.

To address this knowledge gap we conducted a mesocosm investigation to study the influence of three WT levels (from drained to inundated) and multi-level N deposition (from non-addition to 160 kg N ha⁻¹ yr⁻¹) on the soil CH₄ and N₂O emissions in the Zoige alpine peatland, located on the eastern edge of the Qinghai-Tibetan Plateau. Being the largest and highest swamp wetland area in China, its sensitivity to the global climate change and human activities is high (Chen et al., 2013). Exposure to a potential influence of drainage, restoration or increasing N deposition (Yang et al., 2017; Zhang et al., 2011) may increase the risk of high GHG emissions from this area. In this study, we aim to address the following two questions: i) **do increasing rates of N deposition consistently stimulate CH₄ and N₂O emissions?** ii) if there is interaction between N deposition and WT level, how do they combine to influence CH₄ and N₂O emissions in the alpine peatland?

2. Methods and Materials

2.1 Study site

This study was conducted in the Zoige alpine wetland, situated on the eastern edge of the Qinghai-Tibetan Plateau, Southeast China, during the 2018 and 2019 growing seasons. This alpine wetland covers an area of 6180 km², which is 31.5% of the whole Zoige plateau. The mean annual temperature is 1.4°C, with a maximum of 9.1 to 11.4°C in July and a minimum of -8.2 to -10.6°C in January, while the average annual precipitation is approximately 650 mm (Chen et al., 2013; Yang et al., 2014). Over the past four decades, the mean annual air temperature has increased by 0.4°C per decade, while the total annual precipitation has decreased by 22 mm per decade (Chen et al., 2013; Yang et al., 2014). Data on precipitation and air temperature in this study were obtained from the closest meteorological station belonging to the Chinese National Meteorological Information Center (www.nmic.gov.cn) and are shown in Figure S1. The depth of peat in the vertical profile is around 1.2 m, soil pH is 6.8-7.2 and soil bulk density around 0.78 g m⁻³ (Zhang et al., 2020). The plant growing season ranges from June to September, and the dominant plants are *Carex muliensis*, *Lancea tibetica*, *Potentilla anserina* L. and *Trollius farreri* Stapf.

2.2 Experimental design

Our experiment was carried out at the Sichuan Zoige Wetland Ecosystem Research Station, Tibetan Autonomous Prefecture of Aba (33°57'N, 102°52'E, 3500 m a.s.l.). A homogeneous swamp wetland was selected for collection of soil and plants to be used in the mesocosm. Forty-five tanks (0.6 m length × 0.6 m width × 0.6 m height) were kept aboveground and filled with intact soil cores and vegetation (Figure S2). The bottom of the tanks was welded, and the outside of the tanks was wrapped with polystyrene foam to avoid heat exchange with the surroundings.

The experimental treatments consisted of five levels of added N and three water table levels and applied in a factorial design (5 N addition × 3 water table). The treatments were replicated three times, giving a total of 45 experimental plots. Based on previous

studies indicating water levels effects on GHG emission in the Zoige peatland (Cui et al., 2017; Yang et al., 2017), three water table levels (WT₋₃₀, 30 cm below the soil surface; WT₀, 0 cm at the soil surface; and WT₁₀, 10 cm above the soil surface) were selected. To maintain the water table at the selected three levels, we developed a water table control system composed of three water table detectors, a manostat, a relay and micropumps. Three water table detectors were placed in the PVC pipe (diameter 3 cm) of each tank at the exact water table level and at +2 cm and -2 cm water table. When the water table was below the -2 cm detector, the pump switched on, supplying the tanks with local tap water until the water table reached the +2 cm detector. To prevent the water table from becoming too high due to pump water or precipitation, four small holes (diameter 1 cm, and two holes for two sides) were drilled at the precise position of the water table in each tank to allow water overflow.

The current N deposition in the Zoige area is 1.08-17.81 kg N·ha⁻¹·yr⁻¹, NH₄⁺ and NO₃⁻ being the main component, and N deposition is expected to increase further in the future (Han et al., 2019). NH₄NO₃ was adopted as N source to simulate the different stages of the response of alpine peatlands to multi-level N deposition, and five N addition levels were established for each water table level, namely 0 (N₀), 20 (N₂₀), 40 (N₄₀), 80 (N₈₀), and 160 (N₁₆₀) kg N ha⁻¹ yr⁻¹. The three lowest levels (N₀, N₂₀ and N₄₀) cover the gradient of current and near-future deposition levels while the two highest levels (N₈₀ and N₁₆₀) represent levels of N-enrichment resulting from extreme deposition levels possibly combined with N input from fertilization or livestock excreta. The annual added N doses were further divided into four portions and applied at the beginning of every month from June to September in 2018 and 2019. 25% of the added N was dissolved into 1 L water and sprayed evenly upon the surface of each plot. while the control plot only received 1 L water (Wang et al., 2017).

2.3 GHG sampling and measurements

We measured the CH₄ and N₂O fluxes with the sampling events of 1-3 times per month during the growing seasons in 2018 and 2019 in our study. In total, 16 sampling occasions of individual fluxes were recorded for CH₄ and N₂O. In each tank, CH₄ and N₂O fluxes were measured using static opaque chambers and gas chromatography (GC) (Zhang et al., 2017). The cubic chamber was made of stainless steel (0.5 m length × 0.5 m width × 0.5 m height; without bottom). At the top surface of the chamber, there were two ports for headspace gas sampling and enclosed air temperature measurements, respectively. A dry battery-powered fan was placed in the chamber to avoid stratification of the gases during sampling. Meanwhile, 45 square collars (0.5 m length × 0.5 m width × 0.2 m height) were produced and buried into the middle part of the soil core in each tank at about 0.2 m depth. Before placing the chambers on top of the collars to collect gas samples, the collars were sealed with water to ensure minimum gas exchange between chamber and atmosphere.

Gas samples (20 mL) were collected from each chamber using a rubber tube connected to the valve of the chamber and a syringe at 10 min intervals over a 20-minute period (0, 10 and 20 min). The samples were then injected into a pre-evacuated 10 mL vacuum vial (Aladdin, Shanghai, China). The samples were kept cold and dark until the CH₄ and N₂O fluxes were determined via GC (Agilent 7890A, Agilent Co., Santa Clara, CA, USA) within 72 hours. The GC was equipped with a flame ionization detector

(FID) to analyse the CH₄ concentration and an electron capture detector (ECD) to analyse the N₂O concentration. The carrier gas was N₂, and the operation temperature for the FID was set at 250 °C and ECD at 300 °C. The CH₄ and N₂O fluxes were calculated as follows.

$$F = \frac{M}{V_0} \frac{P}{P_0} \frac{T}{T_0} \frac{dc}{dt} H$$

where dc/dt is the slope of the linear regression for the gas concentration gradient through time; M is the molecular mass of CH₄ or N₂O; P is the atmospheric pressure at the sampling site; T is the absolute temperature during sampling; V_0 , P_0 , and T_0 are the gas mole volume, atmospheric pressure, and absolute temperature under standard conditions, respectively; and H is the chamber height.

Each linear regression was assessed individually, and the estimates of the data quality of the fluxes were uniformly dependent on R-squared values. However, the coefficients of determinations (R^2) of the linear regression for CH₄ and N₂O were sometimes low (<0.4), particularly when the fluxes were low. We did not want to create bias against these low fluxes and therefore kept them if the CO₂ concentration (data not shown) showed a good linear trend with time (Laine et al., 2019). Apart from these fluxes, values were generally accepted only if the R-squared values of the linear regressions were equal to or greater than 0.8 and 0.7 for CH₄ and N₂O (Lafuente et al., 2020; Laine et al., 2019), respectively. The CH₄ flux had 5.29% discarded values, while the N₂O flux had 3.70% discarded values.

2.4 Analysis of soil properties

To determine soil properties, soil samples were collected in late September, considered as the end of the growing seasons in 2018 and 2019. Three sub-samples of soil were collected from each tank at the top 5 cm depth and then bulked into a composite sample representing a reliable replicate. The collected soil samples were stored under cold and dark conditions and then forwarded to the laboratory within three days. The soil samples were passed through a 2 mm sieve and air dried for the determination of soil pH, soil organic carbon (SOC) and total nitrogen (TN). Soil pH was measured at a soil:water ratio of 1:2.5 with a pH electrode (PHS 29, China). SOC and TN were determined via dry combustion using an Elementar Vario Max CN analyzer (Hanau, Germany). SWC was determined by using a TDR300 moisture meter (Spectrum Technologies Inc., Plainfield, Illinois, USA).

2.5 Statistical analysis

Generalized least square (GLS) ANOVA was used to assess the effect of WT and N (fixed factor) on the cumulative CH₄ and N₂O emissions in 2018 and in 2019 (respectively), and the soil properties were determined via the R package *nlme* (Pekár et al., 2016; Tiemeyer et al., 2016). We also used the GLS method to compare the effects of N deposition on CH₄ and N₂O emissions at each WT level in each year, followed by a Tukey HSD test. The GLS model included an autoregressive structure, accommodated for unequal time of sampling, and a variance function allowing for unequal variance in the fixed factors (Wanyama et al., 2019).

A Generalized Additive Model (GAM) was used to fit the relationship between the cumulative CH₄ emissions and N deposition

dosages at different water table levels. Via the R package “mgcv” (Wood, 2017), we used method “gam” to perform the GAM analysis and method “predict.gam” to see the response value of GHG emissions along the N deposition gradient from 0 to 160 kg N ha⁻¹ yr⁻¹. Compared to the linear models, GAM directly and smoothly fitted the non-linear relationship between the response variable and the multiple explanatory variables, regardless of the data distribution (Chen et al., 2021).

The cumulative GHG emissions in the growing seasons of each year were calculated by linear interpolation between sampling events using the trapezoidal rule (Goldberg et al., 2010). In addition to the cumulative GHG emissions between the first and the last sampling event, the GHG emissions from 1st June to the first sampling and from the last sampling to 30th September were taken into consideration.

Statistical analysis was carried out using R (version 3.4.3) (R Core Team, 2017), and graphs were drawn using OriginPro 9.8.0.200. Final *p* values were Bonferroni adjusted to mitigate the risk of type I error. A significance level of $\alpha = 0.05$ was used for all statistical tests.

3. Results

3.1 Soil properties

During the two years of the growing season mesocosm experiment, the SWC varied from 63.5% to 81.1% and was only significantly affected by the water table levels (Table 1 and 2). The higher WT levels significantly increased the SWC in both years. The soil pH varied within the range 7.3 to 7.8 and was only significantly affected by N deposition. Large variability of SOC was observed within the range 215.9 g kg⁻¹ to 296.1 g kg⁻¹, and both the WT and N treatments showed significant effects on SOC, without any interactive effects. Compared with the control treatment without N deposition, N deposition increased SOC by 1.4% to 31.1% (except WT₀ N₁₆₀ in 2019). Soil TN varied between 12.9 g kg⁻¹ and 19.1 g kg⁻¹ and was elevated by N deposition, whereas no significant response to the WT treatments was observed. N deposition enhanced soil TN by 1.3% to 110.5% compared to the N₀ treatment at each WT level.

Table 1. Soil properties in the different water table (WT) treatments and nitrogen deposition (N) levels in 2018 and 2019.

		2018			2019		
		WT ₋₃₀	WT ₀	WT ₁₀	WT ₋₃₀	WT ₀	WT ₁₀
SWC (%)	N ₀	65.8±2.5	67.4±2.6	69.8±1.5	62.8±1.7	70.4±1.0	81.1±1.6
	N ₂₀	66.8±1.8	74.2±1.6	74.2±1.9	63.5±2.0	71.3±0.4	79.0±2.0
	N ₄₀	67.3±1.5	73.2±3.0	71.0±1.7	64.4±1.9	71.6±1.9	79.8±1.6
	N ₈₀	67.4±2.2	72.0±0.8	73.1±1.1	67.8±0.8	69.6±1.4	77.4±2.1
	N ₁₆₀	64.4±1.3	68.0±2.2	72.8±1.8	68.1±0.7	72.2±1.8	81.1±1
pH	N ₀	7.6±0	7.7±0.1	7.7±0.1	7.7±0	7.8±0.1	7.8±0.1
	N ₂₀	7.5±0	7.7±0	7.4±0.1	7.7±0.1	7.5±0.2	7.6±0.1

	N ₄₀	7.3±0	7.6±0.1	7.6±0.1	7.5±0.1	7.6±0.2	7.7±0.2
	N ₈₀	7.6±0	7.4±0.1	7.5±0.1	7.4±0.1	7.6±0.1	7.4±0
	N ₁₆₀	7.5±0	7.6±0	7.3±0.1	7.5±0.1	7.4±0.1	7.5±0
SOC (g kg ⁻¹)	N ₀	231.3±5.4	237±24.3	246.6±16.0	215.9±3.6	227.3±14.4	218.3±14.0
	N ₂₀	296.1±5.7	285.8±8.4	279.2±23.4	228.7±9.6	249.9±12.0	273.3±11.8
	N ₄₀	292.3±14.1	281.2±18.7	262.8±20.9	241.8±6.7	281.0±17.8	253.3±17.5
	N ₈₀	265.4±17.7	294.3±7.7	291.4±9.3	240.9±12	230.6±10.8	286.2±9.1
	N ₁₆₀	275.6±7.0	276.8±10.1	266.8±32.4	254.6±18.2	226.8±13.7	251.5±19.1
TN (g kg ⁻¹)	N ₀	17.6±0.8	16.1±1.1	18.7±0.8	12.9±0.9	14.4±0.3	14.7±2.1
	N ₂₀	18.7±0.7	19.1±0.6	19.3±0.8	21.9±2.2	21.2±3.2	23.3±5.0
	N ₄₀	18.4±1.1	18.8±0.9	19.2±0.4	19.1±1.5	21.4±4.0	15.0±2.7
	N ₈₀	18.3±0.8	19.4±0.2	19.7±0.2	18.6±1.1	16.2±1.0	31.0±2.6
	N ₁₆₀	18.1±0.8	18.2±0.4	19.0±0.4	19.5±2.0	21.7±6.6	21.9±4.6

Each value represents mean ± SE (n=3). SWC, soil water content; SOC, soil organic carbon; TN, total nitrogen.

Table 2. Effects of water table (WT) and nitrogen (N) deposition levels and their interactions on soil properties using generalized least square (GLS) models.

	SWC		pH		SOC		TN	
	F	P	F	P	F	P	F	P
WT	19.4	<0.001***	0.34	0.7103	9.92	<0.001***	2.08	0.1319
N	0.64	0.6352	6.78	<0.001***	5.18	0.001**	4.49	0.002**
WT×N	0.25	0.9807	0.35	0.944	0.91	0.5147	0.74	0.6526

Bold font denotes significant values. The statistical significance levels used were: *: 0.01 < P < 0.05; **: 0.001 < P < 0.01; ***: P < 0.001. SWC: soil water content; SOC: soil organic carbon; TN: total nitrogen.

3.2 Methane

The Zoige alpine peatland acted as a net source of CH₄ in the WT₀ and WT₁₀ treatments throughout the two growing seasons, although the CH₄ flux was almost 0 in the WT₋₃₀ treatment. Temporal variability of the CH₄ flux was observed (Figure 1). The cumulative CH₄ emissions of the growing season ranged from -0.26 to 29.26 g CH₄-C m⁻² in 2018 and from -0.35 to 16.36 g CH₄-C m⁻² in 2019 (Figure 2). During the two years, the WT treatments and their interaction with the N treatments had significant effects on the cumulative CH₄ emissions, while N deposition only had significant effects in 2019 (Table 3). Along the WT level gradient from WT₋₃₀ to WT₁₀, the cumulative CH₄ emissions increased markedly. The response of the cumulative CH₄ emissions to N deposition was non-linear under WT₀ and WT₁₀ conditions (Figure 3), with the highest value observed in the N₂₀ treatment.

Compared to the N_0 treatment, the N_{80} and N_{160} treatments remarkably decreased the cumulative CH_4 emissions by 36.5% to 97.4%, while N_{40} was in the same order of magnitude as in the N_0 treatment. The GAM results showed that the cumulative CH_4 emissions could be explained by N deposition for 55.9% under WT_0 conditions and for 45.4% under WT_{10} conditions. The modelling results also indicated that the critical thresholds for the highest cumulative CH_4 emissions were 14.41 g C m^{-2} with $20.9 \text{ kg ha}^{-1} \text{ yr}^{-1}$ N deposition under WT_0 conditions and 21.60 g C m^{-2} with $16.2 \text{ kg ha}^{-1} \text{ yr}^{-1}$ N deposition under WT_{10} conditions.




Figure 1. Temporal variation of growing-season CH_4 and N_2O fluxes under five levels of nitrogen deposition (0, 20, 40, 80 and 160 $\text{kg N ha}^{-1} \text{ yr}^{-1}$) and three water table levels in 2018 and 2019. Error bars represent the SE ($n=3$).




Figure 2. Effects of nitrogen deposition levels on cumulative CH₄ and N₂O emissions at three water table levels during the growing seasons in 2018 and 2019. Error bars represent the SE (n=3). Different letters above the bars indicate statistically significant differences ($P < 0.05$).

Table 3. The individual and interactive effects of water table (WT) and nitrogen (N) deposition levels on cumulative CH₄ and N₂O emissions in 2018 and 2019 using generalized least square (GLS) models.

		CH ₄		N ₂ O	
		F	<i>P</i>	F	<i>P</i>
2018					
	WT	24.88	<0.001***	36.68	<0.001***
	N	1.37	0.27	239.38	<0.001***
	WT×N	15.15	<0.001***	4.28	0.002**
2019					
	WT	615.89	<0.001***	351.26	<0.001***
	N	5.99	0.001**	989.75	<0.001***
	WT×N	18.01	<0.001***	5.23	<0.001***

Bold font denotes significant values. The statistical significance levels used were: *:0.01<P < 0.05; **: 0.001< P < 0.01; ***: P<0.001.




Figure 3. The relationship between cumulative greenhouse gas (CH₄ and N₂O) emissions and nitrogen deposition levels at three water table levels. A linear model was used to estimate cumulative N₂O emission at five nitrogen deposition levels, while a generalized additive model (GAM) was used to assess the response of cumulative CH₄ emission to nitrogen deposition levels. The statistical significance levels used were: * = $P < 0.05$ and > 0.01 ; ** = $P < 0.01$ and > 0.001 ; *** $P < 0.001$; NS = not significant ($P > 0.05$). The shaded areas indicate 95% confidence intervals.

3.3 Nitrous oxide

The Zoige alpine peatland acted as a net N₂O source during the growing seasons, the N₂O fluxes showing clear temporal variability in 2018 and 2019. The N₂O flux tended to be highest in early September 2018 and in mid-August 2019, while the lowest flux was observed at the start or the end of the growing seasons (Figure 1). The cumulative N₂O emissions ranged from 0.02 to 0.49 g N₂O-N m⁻² in 2018 and from 0.05 to 0.50 g N₂O-N m⁻² in 2019. The cumulative N₂O emissions were significantly affected by the WT levels, N deposition and their interaction (Table 3). N deposition significantly increased the cumulative N₂O emission by 28.9% to 1974.6%, most significantly in the N₁₆₀ treatment. However, there was no clear effect of WT levels on N₂O emissions. We observed a significantly positive and linear relationship between the cumulative N₂O emissions and N application doses, and the slope and intercept of the linear correlation depended on the WT levels (Figure 3). The linear results also showed that the 1 kg N ha⁻¹ addition increased the cumulative N₂O emission by 0.0024, 0.0028 and 0.0023 g N₂O-N m⁻² under WT₋₃₀, WT₀ and WT₁₀ conditions, respectively.

4. Discussion

4.1 Effects of WT and N deposition on CH₄ emission

The cumulative CH₄ emissions from the Zoige alpine peatland in our study ranged from -0.35 to 29.26 g CH₄-C m⁻² across the two years, which is in the same order of magnitude as the cumulative CH₄ emissions (25.4-29.6 g CH₄-C m⁻²) from an alpine wetland of the Qinghai-Tibetan Plateau in a previous study (Wang et al., 2017). As expected, WT levels had a significant positive effect on CH₄ emissions, with higher WT levels increased CH₄ emissions. This corresponds with the results of previous studies, due to the manipulative effects of WT levels on the soil redox conditions (Hoyos-Santillan et al., 2019; Wang et al., 2017). With higher WT levels, SWC increased and likely formed more anaerobic conditions conducive to CH₄ production, leading to elevated CH₄ emissions (Evans et al., 2021; Hoyos-Santillan et al., 2019; Zhang et al., 2020).

In the current study, a moderate level of N deposition positively stimulated the CH₄ emissions, but subsequently the positive effect declined with further N addition. However, this non-linear effect of N deposition was only observed at high WT levels, and the close to zero emissions of CH₄ at low WT levels were not affected by N deposition. To our knowledge, the non-linear effects of N deposition on CH₄ fluxes in the upland ecosystems has been reported in numerous studies (Li et al., 2012; Lafuente et al., 2020; Qu et al., 2021), however, there have been very few attempts in peatlands. Song et al. (2013) found a similar pattern of non-linear effects of N addition (0-240 kg N ha⁻¹ yr⁻¹) on CH₄ fluxes in a wetland with the highest CH₄ flux observed at 60 kg ha⁻¹ yr⁻¹ N addition, but unfortunately, this N effect was not significant. Wu et al. (2022) conducted a global meta-analysis and found the N addition (0-300 kg N ha⁻¹ yr⁻¹) could consistently increase the CH₄ emissions in the global wetlands, but the highest value occurred at 50-100 kg N ha⁻¹ yr⁻¹ N addition. The results of the aforementioned studies were not in agreement with the present study, due to the differences between the threshold values and the inhibitory or promotive effects of further N addition on CH₄ emission. However, in accordance with the previous studies, the relatively low level of N addition probably alleviated N constraints on microbial metabolism in soils and increased CH₄ production (Currey et al., 2009; Deng et al., 2019). With the further N addition, large amounts of available NO₃⁻ might have led negative and inhibitory effects on the methanogenic activity due to the competition of NO₃⁻-reducing bacteria with methanogens (Liu et al., 2020).

The interactive effects of WT levels and N deposition on the cumulative CH₄ emissions were distinct in our study (Table 3 and Figure 3). We found that the WT levels were more likely to determine the direction and magnitude of CH₄ emissions from alpine peatlands than N deposition. This is in line with a previous study (Evans et al., 2020), which indicated that WT was the overriding factor to control GHG emissions from the peatlands at a global scale. The N deposition non-linearly affected the CH₄ emissions, and the scenario for maximum CH₄ emissions was roughly ca. 20 kg N ha⁻¹ yr⁻¹, which could be slightly changed by the WT levels. To the best of our knowledge, this is the first mesocosm experiments in an alpine peatland comparing the interactive effects of multi-level N deposition and WT levels on CH₄ emissions, which makes it challenging to put in context the scope of these results, and we have not found any theory in the previous studies which could best explain the underlying mechanism. We speculate that the N

deposition supplied N substrate and WT levels were associated with N utilization by microorganisms. Precisely, the higher WT levels promoted diffusion of the added N in the water-filled soil pore, and N thus becoming readily accessible in the microbial process to support CH₄ production (Wang et al., 2017).

4.2 Effects of WT and N deposition on N₂O emission

The mesocosms in the Zoige alpine peatland were consistently acting as N₂O sources, and the cumulative N₂O emissions (0.02-0.49 g m⁻² in 2018 and 0.05-0.50 g m⁻² in 2019) did not show significant differences between the two years. The cumulative N₂O emission from the Zoige peatland in our study was relatively higher than that in a previous report (0.08-0.2 g m⁻²), which focused on the drainage or lower water table level (Cao et al., 2018). Unlike the CH₄ emissions, we did not observe a clear pattern for the effects of WT levels on N₂O emissions in our study. This is different from the numerous previous studies, which all confirmed the positive effects of WT on N₂O emissions in peatlands, due to its positive effects on anaerobic denitrification and N₂O production (Gao et al. 2014; Regina et al., 1999; Regina et al., 1996). However, Wang et al. (2017) reported that elevated WT levels from drained to inundated conditions had no effects on N₂O fluxes in an alpine wetland of the Qinghai-Tibetan Plateau, which is in line with our study. The possible reason for this could be due to the large differences of habitat types, soil properties or precipitations among the regional heterogeneous peatlands.

N deposition stimulated N₂O emissions across the two years in our study, and same results were also found in the previous reports (Gao et al., 2014; Deng et al., 2019). The N deposition increased soil TN ($F = 4.49$, $P = 0.002$) in our study and is likely to supply more N substrate (NH₄⁺ and NO₃⁻) in soil (Zhu et al., 2020). The consequently increased N substrate could potentially activate the microbial process of N₂O production and increase N₂O emissions (Yue et al., 2021). Additionally, contrary to the CH₄ emissions, the N deposition showed consistently linear effects on N₂O emissions in the peatland, regardless of the WT levels. This is not in line with a previous study (Song et al., 2013), which indicated that N addition ranging from 0-120 kg N ha⁻¹ yr⁻¹ did not show significant effects on N₂O emissions in peatlands, while high level (240 kg N ha⁻¹ yr⁻¹) largely increased the N₂O emissions. However, a global meta-analysis showed that N₂O emissions from wetlands demonstrated significant positive response to N enrichment (Deng et al., 2019), which agreed with our study. The possible explanation could be that the current level of N deposition in the Zoige peatland did not meet the N-saturation stage, after which the sensitivity of N₂O emission from the ecosystem to further N addition would decline (Gomez-casanovas et al., 2016).

The N₂O emission from the alpine peatland was likely primarily determined by N deposition rather than by WT levels. We confirmed the occurrence of an interactive effect of WT and N deposition on N₂O emissions, but it was neither synergistic nor antagonistic. N deposition had linear positive effects on N₂O emission, and WT level did not alter this linear relationship but slightly changed the slope and intercept. The N₂O emissions could be generated from both the aerobic nitrification and anaerobic denitrification processes. This is confirmed by the previous study (Bateman et al., 2005), which indicate that the denitrification was the major contributor to N₂O emission at water-filled pore space (WFPS) > 60% while nitrification was the major one at WFPS <

60%. The overall production of N_2O could not be uniquely manipulated by the WT levels, though the WT levels could influence the redox conditions and the microbial processes. Additionally, the Zoige peatland being a N-limited ecosystem (Squeo et al., 2006), we observed the overwhelming effects of N deposition on N_2O emission in the peatland, rather than WT levels.

The growing-season N_2O emission in the current study increased by 0.0023-0.0028 g N_2O -N m^{-2} in response to the additional 1 kg NH_4NO_3 -N ha^{-1} yr^{-1} deposition. This is slightly lower than the levels from previous studies (Gong et al., 2019), which showed that 1 kg annual N ha^{-1} addition led to an increase of ca. 0.0076 g N_2O -N m^{-2} during the growing season in a boreal peatland. This could be attributed to the relatively low air temperature at this particular alpine peatland, which hampered the microbial N_2O production. Furthermore, IPCC (2013) suggested that the default emission factor (the fraction of nitrogen added that is released as N_2O) is 1%, indicating that 1 kg annual N ha^{-1} addition may increase N_2O emissions by 0.01 kg N_2O -N ha^{-1} (or 0.001 g N_2O -N m^{-2}). The relatively higher emission factor for N_2O -N in our study was probably due to the high dose of N addition.

4.3 Implications and limitations

To what extent the magnitude of non- CO_2 emissions from alpine peatlands could be upscaled in response to the increasing N deposition, and how this effect interacts with the WT levels is still high uncertain. Our study demonstrated that the CH_4 and N_2O emissions from the alpine peatland system could reach a new state if the projected increases in N deposition continued, in particular the CH_4 emission could be disproportionally increased. In the Zoige peatland, the current level of N deposition was 1.08-17.81 kg N ha^{-1} yr^{-1} (Han et al., 2019), and the N deposition was predicted to be a possible doubling or tripling by the end of the century (Lamarque, 2005). Given the highest CH_4 emissions occurred at the level of ca. 20 kg N ha^{-1} yr^{-1} deposition and the consistent linear correlation between N_2O emissions and N deposition levels in our study, we infer that the CH_4 and N_2O emissions had not yet peaked, and the increasing N deposition could still result in the alpine peatland becoming a potential hotspot of GHG emissions in the future. Moreover, the fluctuation of WT levels resulting from anthropogenic drainage or rewetting could affect the GHG emissions but not suppress the N deposition-induced trend. However, our study is limited to a specific habitat, and thus the upscaling of this conclusion contains large uncertainty. To the best of our knowledge, very few advances heretofore have been achieved to elaborate this general conclusion. We believe that our results are useful for predicting the GHG emissions from alpine peatlands in response to the climate change and anthropogenic activities in the future.

It should be noted that some levels of N deposition (80 or 160 kg N ha^{-1} yr^{-1}) in our study were much higher than the local N deposition (1.08-17.81 kg N ha^{-1} yr^{-1}). This should not affect our general conclusion, because the non-linear and linear effects of N deposition on CH_4 and N_2O emissions (respectively) were primarily dependent on the low levels of N deposition (0-40 kg N ha^{-1} yr^{-1}), and the higher levels did not alter the relationship pattern. Meanwhile, note must be made that we did not measure the net ecosystem CO_2 exchange and wintertime GHG fluxes, which might hamper estimating the annual carbon budget from GHG emissions and SOC change. However, our study focused on the growing-season non- CO_2 emissions from the peatland at different WT levels under the future scenarios of increasing N deposition, and also the non-growing-season GHG emissions had only a minor

contribution to the yearly budget due to the low temperature and microbial activities (Peng et al., 2019). In addition, the low frequency of GHG sampling in 2018 could cause uncertainties in the temporal variation and cumulative emissions of CH₄ and N₂O, and this might result in bias in the present result. The monthly measurements of GHG fluxes from peatlands have already been found in the previous study (Cao et al., 2018), and we also increased the sampling frequency in 2019 to better support the current conclusion. However, further monitoring of GHG fluxes from the peatland would still be required to eliminate the uncertainties.

5. Conclusion

This study explored the response of non-CO₂ GHG emissions in the alpine peatland to the increasing N deposition at different WT levels. We found that CH₄ emissions were determined by N deposition, WT levels and their interactive effects. A modest input of N deposition and high WT levels both stimulated CH₄ emissions. N₂O emissions were remarkably sensitive to N deposition, which consistently and linearly increased the N₂O emissions, irrespective of WT levels. The projected increasing N deposition suggested an inevitable risk for higher CH₄ and N₂O emissions from the alpine peatland in the future.

Data availability

All data are available from the corresponding author by request.

Author contributions

Wantong Zhang: Conceptualization, Data curation, Software, Writing - original draft, Writing - review & editing, Validation, Formal analysis. **Zhengyi Hu:** Conceptualization, Supervision. **Joachim Audet:** Data curation, Supervision, Writing – review & editing. **Thomas A. Davidson:** Supervision, Data curation, Writing – review & editing. **Enze Kang:** Investigation. **Xiaoming Kang:** Investigation, Project administration. **Yong Li:** Investigation. **Xiaodong Zhang:** Investigation, Project administration. **Jinshi Wang:** Investigation, Conceptualization, Project administration, Writing - review & editing.

Funding

This study was financially supported by the National Natural Science Foundation of China (41877421 and 31770511).

Competing interests

The contact author has declared that neither they nor their co-author has any competing interests.

References

- Audet J., Elsgaard L., Kjaergaard C., Larsen S. E. , Hoffmann C. C.: Greenhouse gas emissions from a Danish riparian wetland before and after restoration, *Ecol. Eng.*, 57, 170-182. <https://doi.org/10.1016/j.ecoleng.2013.04.021>, 2013.
- Bateman E. J., Baggs E.M.: Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space. *Biology and Fertility of Soils* 41:379-388. <https://doi.org/10.1007/s00374-005-0858-3>, 2005
- Cao R., Xi X., Yang Y., Wei X., Wu X. , Sun S.: The effect of water table decline on soil CO₂ emission of Zoige peatland on eastern

- Tibetan Plateau: A four-year in situ experimental drainage, *Appl. Soil Ecol.*, 120, 55-61. <https://doi.org/10.1016/j.apsoil.2017.07.036>, 2017.
- Cao R., Chen Y., Wu X., Zhou Q., Sun S.: The effect of drainage on CO₂, CH₄ and N₂O emissions in the Zoige peatland: a 40-month in situ study. *Mires Peat*, 21, 1-15. <https://doi.org/10.19189/MaP.2017.OMB.292>, 2018.
- Chen H., Zhu Q., Peng C., Wu N., Wang Y., Fang X., Gao Y., Zhu D., Yang G., Tian J., Kang X., Piao S., Ouyang H., Xiang W., Luo Z., Jiang H., Song X., Zhang Y., Yu G., Zhao X., Gong P., Yao T., Wu J.: The impacts of climate change and human activities on biogeochemical cycles on the Qinghai-Tibetan Plateau, *Global Change Biol.*, 19, 10, 2940-55. <https://doi.org/10.1111/gcb.12277>, 2013.
- Chen Q., Wang S., Ni Z., Guo Y., Liu X., Wang G., Li H.: No-linear dynamics of lake ecosystem in responding to changes of nutrient regimes and climate factors: Case study on Dianchi and Erhai lakes, China, *Sci. Total Environ.*, 781. <https://doi.org/10.1016/j.scitotenv.2021.146761>, 2021.
- Cui L., Kang X., Li W., Hao Y., Zhang Y., Wang J., Yan L., Zhang X., Zhang M., Zhou J., Kardol P.: Rewetting decreases carbon emissions from the Zoige alpine peatland on the Tibetan plateau, *Sustainability*, 9, 6. <https://doi.org/10.3390/su9060948>, 2017.
- Currey P. M., Johnson D., Sheppard L. J., Leith I. D., Toberman H., Van Der Wal R., Dawson L. A., Artz R. R. E.: Turnover of labile and recalcitrant soil carbon differ in response to nitrate and ammonium deposition in an ombrotrophic peatland, *Global Change Biol.*, 16, 8, 2307-2321. <https://doi.org/10.1111/j.1365-2486.2009.02082.x>, 2009.
- Deng L., Huang C., Kim D. G., Shangguan Z., Wang K., Song X., Peng C.: Soil GHG fluxes are altered by N deposition: New data indicate lower N stimulation of the N₂O flux and greater stimulation of the calculated C pools, *Global Change Biol.* <https://doi.org/10.1111/gcb.14970>, 2019.
- Evans C. D., Peacock M., Baird A. J., Artz R. R. E., Burden A., Callaghan N., Chapman P. J., Cooper H. M., Coyle M., Craig E., Cumming A., Dixon S., Gauci V., Grayson R. P., Helfter C., Heppell C. M., Holden J., Jones D. L., Kaduk J., Levy P., Matthews R., McNamara N. P., Misselbrook T., Oakley S., Page S., Rayment M., Ridley L. M., Stanley K. M., Williamson J. L., Worrall F., Morrison R.: Overriding water table control on managed peatland greenhouse gas emissions, *Nature.*, <https://doi.org/10.1038/s41586-021-03523-1>, 2021.
- Frolking S., Talbot J., Jones M. C., Treat C. C., Kauffman J. B., Tuittila E.-S., Roulet N.: Peatlands in the Earth's 21st century climate system, *Environ. Rev.*, 19, 371-396. <https://doi.org/10.1139/a11-014>, 2011.
- Gao Y., Chen H., Zeng X.: Effects of nitrogen and sulfur deposition on CH₄ and N₂O fluxes in high-altitude peatland soil under different water tables in the Tibetan Plateau, *Soil Sci. Plant Nutr.*, 60, 3, 404-410. <https://doi.org/10.1080/00380768.2014.893812>, 2014.
- Goldberg S. D., Knorr K.-H., Blodau C., Lischeid G., Gebauer G.: Impact of altering the water table height of an acidic fen on N₂O and NO fluxes and soil concentrations, *Global Change Biol.*, 16, 1, 220-233. <https://doi.org/10.1111/j.1365-2486.2009.02015.x>, 2010.
- Gomez-Casanovas N., Hudiburg T. W., Bernacchi C. J., Parton W. J., Delucia E. H.: Nitrogen deposition and greenhouse gas emissions from grasslands: uncertainties and future directions, *Global Change Biol.*, 22, 4, 1348-1360. <https://doi.org/10.1111/gcb.13187>, 2016.
- Gong Y., Wu J., Vogt J., Le T. B.: Warming reduces the increase in N₂O emission under nitrogen fertilization in a boreal peatland, *Sci. Total Environ.*, 664, 72-78. <https://doi.org/10.1016/j.scitotenv.2019.02.012>, 2019.
- Han Y., Dong S., Zhao Z., Sha W., Li S., Shen H., Xiao J., Zhang J., Wu X., Jiang X., Zhao J., Liu S., Dong Q., Zhou H., Yeomans J. C.: Response of soil nutrients and stoichiometry to elevated nitrogen deposition in alpine grassland on the Qinghai-Tibetan Plateau, *Geoderma*, 343, 263-268. <https://doi.org/10.1016/j.geoderma.2018.12.050>, 2019.
- Hoyos-Santillan J., Lomax B. H., Large D., Turner B. L., Lopez O. R., Boom A., Sepulveda-Jauregui A., Sjögersten S.: Evaluation of vegetation communities, water table, and peat composition as drivers of greenhouse gas emissions in lowland tropical peatlands, *Sci. Total Environ.*, 688, 1193-1204. <https://doi.org/10.1016/j.scitotenv.2019.06.366>, 2019.
- IPCC: Climate Change: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.
- Järveoja J., Peichl M., Maddison M., Soosaar K., Vellak K., Karofeld E., Teemusk A., Mander Ü.: Impact of water table level on annual carbon and greenhouse gas balances of a restored peat extraction area, *Biogeosciences*, 13, 9, 2637-2651.

- <https://doi.org/10.5194/bg-13-2637-2016>, 2016.
- Juutinen S., Moore T. R., Bubier J. L., Arnkil S., Humphreys E., Marincak B., Roy C. , Larmola T.: Long-term nutrient addition increased CH₄ emission from a bog through direct and indirect effects, *Sci. Rep.*, 8, 1, 3838. <https://doi.org/10.1038/s41598-018-22210-2>, 2018.
- Lafuente A., Recio J., Ochoa-Hueso R., Gallardo A., Perez-Corona M. E., Manrique E. , Duran J.: Simulated nitrogen deposition influences soil greenhouse gas fluxes in a Mediterranean dryland, *Sci. Total Environ.*, 737, 139610. <https://doi.org/10.1016/j.scitotenv.2020.139610>, 2020.
- Laine A. M., Mehtatalo L., Tolvanen A., Frolking S. , Tuittila E. S.: Impacts of drainage, restoration and warming on boreal wetland greenhouse gas fluxes, *Sci. Total Environ.*, 647, 169-181. <https://doi.org/10.1016/j.scitotenv.2018.07.390>, 2019.
- Lamarque J. F.: Assessing future nitrogen deposition and carbon cycle feedback using a multimodel approach: Analysis of nitrogen deposition, *J. Geophys. Res.*, 110, D19. <https://doi.org/10.1029/2005jd005825>, 2005.
- Liu Z., Li D., Zhang J., Saleem M., Zhang Y., Ma R., He Y., Yang J., Xiang H. , Wei H.: Effect of simulated acid rain on soil CO₂, CH₄ and N₂O emissions and microbial communities in an agricultural soil, *Geoderma*, 366. <https://doi.org/10.1016/j.geoderma.2020.114222>, 2020.
- Pekár S., Brabec M. , Bshary R.: Marginal models via gls: A convenient yet neglected tool for the analysis of correlated data in the behavioural sciences, *Ethology*, 122, 8, 621-631. <https://doi.org/10.1111/eth.12514>, 2016.
- Peng, H., Guo, Q., Ding, H., Hong, B., Zhu, Y., Hong, Y., et al.: Multi-scale temporal variation in methane emission from an alpine peatland on the Eastern Qinghai-Tibetan Plateau and associated environmental controls. *Agr. Forest. Meteorol.*, 276-277, <https://doi.org/10.1016/j.agrformet.2019.107616>, 2019.
- Qu S., Xu R., Yu J., Li F., Wei D. , Borjigidai A.: Nitrogen deposition accelerates greenhouse gas emissions at an alpine steppe site on the Tibetan Plateau, *Sci. Total Environ.*, 765, 144277. <https://doi.org/10.1016/j.scitotenv.2020.144277>, 2021.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2017
- Regina K., Nykanen H., Silvola J., Martikainen P. J.: Fluxes of nitrous oxide from boreal peatlands as affected by peatland type, water table level and nitrification capacity, *Biogeochemistry*, 35:401-418. <https://doi.org/10.1007/bf02183033>, 1996
- Regina K., Silvola J., Martikainen P. J.: Short-term effects of changing water table on N₂O fluxes from peat monoliths from natural and drained boreal peatlands. *Global Change Biol.*, 5, 183-189. <https://doi.org/10.1046/j.1365-2486.1999.00217.x>, 1999.
- Saiz E., Sgouridis F., Drijfhout F. P., Peichl M., Nilsson M. B. , Ullah S.: Chronic atmospheric reactive nitrogen deposition suppresses biological nitrogen fixation in peatlands, *Environ. Sci. Technol.*, 55, 2, 1310-1318. <https://doi.org/10.1021/acs.est.0c04882>, 2021.
- Song C., Wang L., Tian H., Liu D., Lu C., Xu X., Zhang L., Yang G., Wan Z.: Effect of continued nitrogen enrichment on greenhouse gas emissions from a wetland ecosystem in the Sanjiang Plain, Northeast China: A 5 year nitrogen addition experiment, *J. Geophys. Res.-Biogeo.*, 118, 741-751. <https://doi.org/10.1002/jgrg.20063>, 2013
- Squeo F. a., Warner B. g., aravena R. , espinoza D.: Bofedales: high altitude peatlands of the central Andes, *Rev. Chil. Hist. Nat.*, 79, 245-255. <http://dx.doi.org/10.4067/S0716-078X2006000200010>, 2006.
- Tiemeyer B., Albiac Borraz E., Augustin J., Bechtold M., Beetz S., Beyer C., Drosler M., Ebli M., Eickenscheidt T., Fiedler S., Forster C., Freibauer A., Giebels M., Glatzel S., Heinichen J., Hoffmann M., Hoper H., Jurasinski G., Leiber-Sauheitl K., Peichl-Brak M., Rosskopf N., Sommer M. , Zeitz J.: High emissions of greenhouse gases from grasslands on peat and other organic soils, *Global Change Biol.*, 22, 12, 4134-4149. <https://doi.org/10.1111/gcb.13303>, 2016.
- Wang H., Yu L., Zhang Z., Liu W., Chen L., Cao G., Yue H., Zhou J., Yang Y., Tang Y. , He J. S.: Molecular mechanisms of water table lowering and nitrogen deposition in affecting greenhouse gas emissions from a Tibetan alpine wetland, *Global Change Biol.*, 23, 2, 815-829. <https://doi.org/10.1111/gcb.13467>, 2017.
- Wanyama L., Pelster D. E., Butterbach-Bahl K., Verchot L. V., Martius C. , Rufino M. C.: Soil carbon dioxide and methane fluxes from forests and other land use types in an African tropical montane region, *Biogeochemistry*, 143, 171–190. <https://doi.org/10.1007/s10533-019-00555-8>, 2019.
- Wilson D., Farrell C. A., Fallon D., Moser G., Muller C., Renou-Wilson F.: Multiyear greenhouse gas balances at a rewetted temperate peatland, *Global Change Biol.*, 22, 12, 4080-4095. <https://doi.org/10.1111/gcb.13325>, 2016.
- Wood S. N.: Generalized Additive Models: An Introduction with R (2nd edition). Chapman and Hall/CRC.

- <https://doi.org/10.1201/9781315370279>, 2017.
- Wu J., Cheng X., Xing W., Liu G.: Soil-atmosphere exchange of CH₄ in response to nitrogen addition in diverse upland and wetland ecosystems: A meta-analysis, *Soil Biol Biochem*, <https://doi.org/10.1016/j.soilbio.2021.108467>, 2022.
- Yang G., Chen H., Wu N., Tian J., Peng C., Zhu Q., Zhu D., He Y., Zheng Q., Zhang C.: Effects of soil warming, rainfall reduction and water table level on CH₄ emissions from the Zoige peatland in China, *Soil Biol. Biochem.*, 78, 83-89. <https://doi.org/10.1016/j.soilbio.2014.07.013>, 2014.
- Yang G., Wang M., Chen H., Liu L. F., Wu N., Zhu D., Tian J. Q., Peng C. H., Zhu Q. A., He Y. X.: Responses of CO₂ emission and pore water DOC concentration to soil warming and water table drawdown in Zoige Peatlands, *Atmos. Environ.*, 152, 323-329. <https://doi.org/10.1016/j.atmosenv.2016.12.051>, 2017.
- Yu Z., Loisel J., Brosseau D. P., Beilman D. W., Hunt S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophys. Res. Lett.*, 37, 13. <https://doi.org/10.1029/2010gl043584>, 2010.
- Yue P., Zuo X., Li K., Cui X., Wang S., Misselbrook T., Liu X.: The driving effect of nitrogen-related functional microorganisms under water and nitrogen addition on N₂O emission in a temperate desert, *Sci. Total Environ.*, 772, 145470. <https://doi.org/10.1016/j.scitotenv.2021.145470>, 2021.
- Zhang L., Hou L., Guo D., Li L., Xu X.: Interactive impacts of nitrogen input and water amendment on growing season fluxes of CO₂, CH₄, and N₂O in a semiarid grassland, Northern China, *Sci. Total Environ.*, 578, 523-534. <https://doi.org/10.1016/j.scitotenv.2016.10.219>, 2017.
- Zhang W., Wang J., Hu Z., Li Y., Yan Z., Zhang X., Wu H., Yan L., Zhang K., Kang X.: The Primary Drivers of Greenhouse Gas Emissions Along the Water Table Gradient in the Zoige Alpine Peatland, *Water Air Soil Poll.*, 231, 5. <https://doi.org/10.1007/s11270-020-04605-y>, 2020.
- Zhang Y., Wang G., Wang Y.: Changes in alpine wetland ecosystems of the Qinghai-Tibetan plateau from 1967 to 2004, *Environ. Monit. Assess.*, 180, 1-4, 189-99. <https://doi.org/10.1007/s10661-010-1781-0>, 2011.
- Zhu J., Chen Z., Wang Q., Xu L., He N., Jia Y., Zhang Q., Yu G.: Potential transition in the effects of atmospheric nitrogen deposition in China, *Environ. Pollut.*, 258, 113739. <https://doi.org/10.1016/j.envpol.2019.113739>, 2020.