

# Effects of water table level and nitrogen deposition on methane and nitrous oxide emissions in an alpine peatland

Wantong Zhang<sup>1,2,4</sup>, Zhengyi Hu<sup>2</sup>, Joachim Audet<sup>4</sup>, Thomas A. Davidson<sup>4</sup>, Enze Kang<sup>1,3</sup>, Xiaoming Kang<sup>1,3</sup>, Yong Li<sup>1,3</sup>, Xiaodong Zhang<sup>1,3</sup>, Jinzhi Wang<sup>1,3, \*</sup>

<sup>1</sup>Institute of Wetland Research, Chinese Academy of Forestry, Beijing Key Laboratory of Wetland Services and Restoration, Beijing 100091, China

<sup>2</sup>Sino-Danish Centre for Education and Research, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>Sichuan Zoige Wetland Ecosystem Research Station, Tibetan Autonomous Prefecture of Aba 624500, China

<sup>4</sup>Department of Ecoscience and Arctic Research Centre (ARC), Aarhus University, C.F. Møllers Allé, 8000 Aarhus, Denmark

\*Correspondence to: Jinzhi Wang (wangjz04@126.com)

## Abstract

Alpine peatlands are recognized as a major natural contributor to the budgets of atmospheric methane (CH<sub>4</sub>) but as a weak nitrous oxide (N<sub>2</sub>O) source. Anthropogenic activities and climate change have put these fragile nitrogen (N)-limited peatlands under pressure by altering water table (WT) levels and enhancing N deposition. The response of greenhouse gas (GHG) emissions from these peatlands to these changes is uncertain. To address this knowledge gap, we conducted a mesocosm experiment in 2018 and 2019 investigating individual and interactive effects of three WT levels (WT<sub>-30</sub>, 30 cm below soil surface; WT<sub>0</sub>, 0 cm at the soil surface; WT<sub>10</sub>, 10 cm above soil surface) and multiple levels of N deposition (0, 20, 40, 80 and 160 kg N ha<sup>-1</sup> yr<sup>-1</sup>) on growing season CH<sub>4</sub> and N<sub>2</sub>O emissions in the Zoige alpine peatland, Qinghai-Tibetan Plateau. We found that the elevated WT levels increased CH<sub>4</sub> emission, while N deposition had non-linear effects (with stimulation at moderate levels, but inhibition at higher levels). In contrast no clear pattern of the effect of WT levels on the cumulative N<sub>2</sub>O emission was evident, while N deposition led to a consistent and linear increase (emission factor: 2.3%-2.8%), and this was dependent on the WT levels. Given the current N deposition in the Zoige alpine peatland (1.08-17.81 kg N·ha<sup>-1</sup>), our results suggested that the CH<sub>4</sub> and N<sub>2</sub>O emissions from the alpine peatlands could greatly increase in response to the possible doubling N deposition in the future. We believe that our results provide insights into how interactions between climate change and human disturbance will alter GHG emissions from this globally important habitat.

**Keywords** Cumulative GHG emissions; multi-level N enrichment; critical threshold; non-linear effect; Qinghai-Tibetan Plateau

## 1. Introduction

Peatlands only cover ca. 3% of the land surface of the Earth but store one third of the global carbon pool (Yu et al., 2010). In pristine peatlands, the shallow water table (WT) and waterlogged conditions allow accumulation of organic matter and favour anaerobic production of methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). Traditionally, this nitrogen-limited ecosystem is recognized as major CH<sub>4</sub> sources and weak N<sub>2</sub>O sources (Frolking et al., 2011). Nevertheless, these conditions could be markedly changed by

anthropogenic disturbance and climate change, and growing evidence shows that peatlands are experiencing drainage and increasing nitrogen deposition (Chen et al., 2013; Evans et al., 2021). Consequently, the magnitude of CH<sub>4</sub> and N<sub>2</sub>O emissions from peatlands may be severely altered, particularly the high-altitude or alpine peatlands that are especially vulnerable and highly sensitive to the climate change and anthropogenic activities (Squeo et al., 2006).

Large-scale artificial drainage of peatlands was initiated hundreds of years ago and escalated in the 20<sup>th</sup> century (Evans et al., 2021). As a result, about 10-20% of the global peatlands were primarily drained for the purposes of agriculture, peat extraction and forestry (Frolking et al., 2011). The resulting lower WT altered the anaerobic conditions of the peat soil and led to oxidative loss of peat (Laine et al., 2019; Wilson et al., 2016). Generally, the drainage decreased the CH<sub>4</sub> efflux and increased CO<sub>2</sub> and N<sub>2</sub>O emissions (Cao et al., 2017). The increase in N<sub>2</sub>O emissions from drained peatlands is often small, but may potentially reach a high level at sufficient nutrient input, especially when the soil is fertilized (Laine et al., 2019). Ecological restoration has been proposed as a measure to conserve the drained or degraded peatlands, particularly to meet the demand for mitigation of greenhouse gas (GHG) emissions outlined in the Paris Agreement (Evans et al., 2021). Numerous studies have reported a remarkably decreased CO<sub>2</sub> efflux in rewetted or restored peatlands, but the rising WT levels have also augmented the emissions of CH<sub>4</sub> and N<sub>2</sub>O (Audet et al., 2013; Järveoja et al., 2016).

Atmospheric N deposition, primarily caused by anthropogenic activities (i.e. fossil fuels combustion, fertilizer application), has increased consistently during the past decades (Gomez-Casanovas et al., 2016, IPCC, 2013), and it is predicted to increase two- or three-fold in terrestrial ecosystems by the end of the century (Lamarque, 2005). The increasing N deposition could alleviate the N stress on peatlands, but the N effects on CH<sub>4</sub> and N<sub>2</sub>O emissions are unclear (Deng et al., 2019). Thus, positive (Juutinen et al., 2018), negative (Gao et al., 2014) or neutral (Wang et al., 2017) effects of N deposition on CH<sub>4</sub> emissions in peatlands have been observed. We speculate that the contrasting results probably are a result of the prevailing environmental conditions and the N addition rate. Besides CH<sub>4</sub> emission, N deposition generally stimulates N<sub>2</sub>O emissions from peatlands due to the increasing supply of N substrate (Wang et al., 2017). However, previous studies have also shown that a higher N input leads to a transition of the grassland into a state of declining N saturation as well as a reduction of the sensitivity of the GHG exchange to the continuously increasing N deposition (Gomez-Casanovas et al., 2016). To eliminate the possible gap resulting from the N addition rate, multiple levels of N deposition are required to study the possible linear or non-linear effects of deposition on GHG emissions.

Numerous studies have reported on the individual effects of WT and N deposition on GHG emissions in peatlands (Evans et al., 2021; Saiz et al., 2021). To our knowledge, only a few studies (Gao et al., 2014; Wang et al., 2017) exist that focus on their interactive effects on peatland GHG emissions. Gao et al. (2014) found that N addition in peatlands decreased CH<sub>4</sub> emissions but increased N<sub>2</sub>O emissions without any significant interaction with WT levels. Wang et al. (2017) observed no interactive effects of a lower WT and increasing N deposition on GHG emissions in an alpine wetland. The above-mentioned studies were, however, limited to a single level of N addition and associated water addition. The response of GHG emissions in peatlands to the gradients of N deposition and WT levels remains to be elucidated, in particular at the N saturation stage, even though it may be a key factor

in shaping GHG emissions. The large uncertainties regarding the interactive effects of N deposition and WT levels on GHG emissions severely hamper obtaining a reliable estimation of the response of peatlands to climate change and anthropogenic activities.

To address this knowledge gap we conducted a mesocosm investigation to study the influence of three WT levels (from drained to inundated) and multi-level N deposition (from non-addition to 160 kg N ha<sup>-1</sup> yr<sup>-1</sup>) on the soil CH<sub>4</sub> and N<sub>2</sub>O emissions in the Zoige alpine peatland, located on the eastern edge of the Qinghai-Tibetan Plateau. Being the largest and highest swamp wetland area in China, its sensitivity to the global climate change and human activities is high (Chen et al., 2013). Exposure to a potential influence of drainage, restoration or increasing N deposition (Yang et al., 2017; Zhang et al., 2011) may increase the risk of high GHG emissions from this area. In this study, we aim to address the following two questions: i) with the N deposition consistently increasing, do the positive effects of N deposition on CH<sub>4</sub> and N<sub>2</sub>O emissions persist? ii) if there is interaction between N deposition and WT level, how do they combine to influence CH<sub>4</sub> and N<sub>2</sub>O emissions in the alpine peatland?

## 2. Methods and Materials

### 2.1 Study site

This study was conducted in the Zoige alpine wetland, situated on the eastern edge of the Qinghai-Tibetan Plateau, Southeast China, during the 2018 and 2019 growing seasons. This alpine wetland covers an area of 6180 km<sup>2</sup>, which is 31.5% of the whole Zoige plateau. The mean annual temperature is 1.4°C, with a maximum of 9.1 to 11.4°C in July and a minimum of -8.2 to -10.6°C in January, while the average annual precipitation is approximately 650 mm (Chen et al., 2013; Yang et al., 2014). Over the past four decades, the mean annual air temperature has increased by 0.4°C per decade, while the total annual precipitation has decreased by 22 mm per decade (Chen et al., 2013; Yang et al., 2014). Data on precipitation and air temperature in this study were obtained from the closest meteorological station belonging to the Chinese National Meteorological Information Center ([www.nmic.gov.cn](http://www.nmic.gov.cn)) and are shown in Figure S1. The depth of peat in the vertical profile is around 1.2 m, soil pH is 6.8-7.2 and soil bulk density around 0.78 g m<sup>-3</sup> (Zhang et al., 2020). The plant growing season ranges from June to September, and the dominant plants are *Carex muliensis*, *Lancea tibetica*, *Potentilla anserina* L. and *Trollius farreri* Stapf.

### 2.2 Experimental design

Our experiment was carried out at the Sichuan Zoige Wetland Ecosystem Research Station, Tibetan Autonomous Prefecture of Aba (33°57'N, 102°52'E, 3500 m a.s.l.). A homogeneous swamp wetland was selected for collection of soil and plants to be used in the mesocosm. Forty-five tanks (0.6 m length × 0.6 m width × 0.6 m height) were kept aboveground and filled with intact soil cores and vegetation (Figure S2). The bottom of the tanks was welded, and the outside of the tanks was wrapped with polystyrene foam to avoid heat exchange with the surroundings.

The experimental treatments consisted of five levels of added N and three water table levels and applied in a factorial design (5 N addition × 3 water table). The treatments were replicated three times, giving a total of 45 experimental plots. Based on previous

94 studies indicating water levels effects on GHG emission in the Zoige peatland (Cui et al., 2017; Yang et al., 2017), three water table  
95 levels (WT<sub>-30</sub>, 30 cm below the soil surface; WT<sub>0</sub>, 0 cm at the soil surface; and WT<sub>10</sub>, 10 cm above the soil surface) were selected.  
96 To maintain the water table at the selected three levels, we developed a water table control system composed of three water table  
97 detectors, a manostat, a relay and micropumps. Three water table detectors were placed in the PVC pipe (diameter 3 cm) of each  
98 tank at the exact water table level and at +2 cm and -2 cm water table. When the water table was below the -2 cm detector, the pump  
99 switched on, supplying the tanks with local tap water until the water table reached the +2 cm detector. To prevent the water table  
100 from becoming too high due to pump water or precipitation, four small holes (diameter 1 cm, and two holes for two sides) were  
101 drilled at the precise position of the water table in each tank to allow water overflow.

102 The current N deposition in the Zoige area is 1.08-17.81 kg N·ha<sup>-1</sup>·yr<sup>-1</sup>, NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> being the main component, and N  
103 deposition is expected to increase further in the future (Han et al., 2019). NH<sub>4</sub>NO<sub>3</sub> was adopted as N source to simulate the different  
104 stages of the response of alpine peatlands to multi-level N deposition, and five N addition levels were established for each water  
105 table level, namely 0 (N<sub>0</sub>), 20 (N<sub>20</sub>), 40 (N<sub>40</sub>), 80 (N<sub>80</sub>), and 160 (N<sub>160</sub>) kg N ha<sup>-1</sup> yr<sup>-1</sup>. The three lowest levels (N<sub>0</sub>, N<sub>20</sub> and N<sub>40</sub>) cover  
106 the gradient of current and near-future deposition levels while the two highest levels (N<sub>80</sub> and N<sub>160</sub>) represent levels of N-enrichment  
107 resulting from extreme deposition levels possibly combined with N input from fertilization or livestock excreta. The annual added  
108 N doses were further divided into four portions and applied at the beginning of every month from June to September in 2018 and  
109 2019. 25% of the added N was dissolved into 1 L water and sprayed evenly upon the surface of each plot. while the control plot  
110 only received 1 L water (Wang et al., 2017).

### 111 2.3 GHG sampling and measurements

112 We measured the CH<sub>4</sub> and N<sub>2</sub>O fluxes with the sampling events of 1-3 times per month during the growing seasons in 2018  
113 and 2019 in our study. In total, 16 sampling occasions of individual fluxes were recorded for CH<sub>4</sub> and N<sub>2</sub>O. In each tank, CH<sub>4</sub> and  
114 N<sub>2</sub>O fluxes were measured using static opaque chambers and gas chromatography (GC) (Zhang et al., 2017). The cubic chamber  
115 was made of stainless steel (0.5 m length × 0.5 m width × 0.5 m height; without bottom). At the top surface of the chamber, there  
116 were two ports for headspace gas sampling and enclosed air temperature measurements, respectively. A dry battery-powered fan  
117 was placed in the chamber to avoid stratification of the gases during sampling. Meanwhile, 45 square collars (0.5 m length × 0.5 m  
118 width × 0.2 m height) were produced and buried into the middle part of the soil core in each tank at about 0.2 m depth. Before  
119 placing the chambers on top of the collars to collect gas samples, the collars were sealed with water to ensure minimum gas exchange  
120 between chamber and atmosphere.

121 Gas samples (20 mL) were collected from each chamber using a rubber tube connected to the valve of the chamber and a  
122 syringe at 10 min intervals over a 20-minute period (0, 10 and 20 min). The samples were then injected into a pre-evacuated 10 mL  
123 vacuum vial (Aladdin, Shanghai, China). The samples were kept cold and dark until the CH<sub>4</sub> and N<sub>2</sub>O fluxes were determined via  
124 GC (Agilent 7890A, Agilent Co., Santa Clara, CA, USA) within 72 hours. The GC was equipped with a flame ionization detector

(FID) to analyse the CH<sub>4</sub> concentration and an electron capture detector (ECD) to analyse the N<sub>2</sub>O concentration. The carrier gas was N<sub>2</sub>, and the operation temperature for the FID was set at 250 °C and ECD at 300 °C. The CH<sub>4</sub> and N<sub>2</sub>O fluxes were calculated as follows.

$$F = \frac{M P T}{V_0 P_0 T_0} \frac{dc}{dt} H$$

where  $dc/dt$  is the slope of the linear regression for the gas concentration gradient through time;  $M$  is the molecular mass of CH<sub>4</sub> or N<sub>2</sub>O;  $P$  is the atmospheric pressure at the sampling site;  $T$  is the absolute temperature during sampling;  $V_0$ ,  $P_0$ , and  $T_0$  are the gas mole volume, atmospheric pressure, and absolute temperature under standard conditions, respectively; and  $H$  is the chamber height.

Each linear regression was assessed individually, and the estimates of the data quality of the fluxes were uniformly dependent on R-squared values. However, the coefficients of determinations ( $R^2$ ) of the linear regression for CH<sub>4</sub> and N<sub>2</sub>O were sometimes low (<0.4), particularly when the fluxes were low. We did not want to create bias against these low fluxes and therefore kept them if the CO<sub>2</sub> concentration (data not shown) showed a good linear trend with time (Laine et al., 2019). Apart from these fluxes, values were generally accepted only if the R-squared values of the linear regressions were equal to or greater than 0.8 and 0.7 for CH<sub>4</sub> and N<sub>2</sub>O (Lafuente et al., 2020; Laine et al., 2019), respectively. The CH<sub>4</sub> flux had 5.29% discarded values, while the N<sub>2</sub>O flux had 3.70% discarded values.

#### 2.4 Analysis of soil properties

To determine soil properties, soil samples were collected in late September, considered as the end of the growing seasons in 2018 and 2019. Three sub-samples of soil were collected from each tank at the top 5 cm depth and then bulked into a composite sample representing a reliable replicate. The collected soil samples were stored under cold and dark conditions and then forwarded to the laboratory within three days. The soil samples were passed through a 2 mm sieve and air dried for the determination of soil pH, soil organic carbon (SOC) and total nitrogen (TN). Soil pH was measured at a soil:water ratio of 1:2.5 with a pH electrode (PHS 29, China). SOC and TN were determined via dry combustion using an Elementar Vario Max CN analyzer (Hanau, Germany). SWC was determined by using a TDR300 moisture meter (Spectrum Technologies Inc., Plainfield, Illinois, USA).

#### 2.5 Statistical analysis

Generalized least square (GLS) ANOVA was used to assess the effect of WT and N (fixed factor) on the cumulative CH<sub>4</sub> and N<sub>2</sub>O emissions in 2018 and in 2019 (respectively), and the soil properties were determined via the R package *nlme* (Pekár et al., 2016, Tiemeyer et al., 2016). We also used the GLS method to compare the effects of N deposition on CH<sub>4</sub> and N<sub>2</sub>O emissions at each WT level in each year, followed by a Tukey HSD test. The GLS model included an autoregressive structure, accommodated for unequal time of sampling, and a variance function allowing for unequal variance in the fixed factors (Wanyama et al., 2019).

A Generalized Additive Model (GAM) was used to fit the relationship between the cumulative CH<sub>4</sub> emissions and N deposition

dosages at different water table levels. Via the R package “mgcv” (Wood, 2017), we used method “gam” to perform the GAM analysis and method “predict.gam” to see the response value of GHG emissions along the N deposition gradient from 0 to 160 kg N ha<sup>-1</sup> yr<sup>-1</sup>. Compared to the linear models, GAM directly and smoothly fitted the non-linear relationship between the response variable and the multiple explanatory variables, regardless of the data distribution (Chen et al., 2021).

The cumulative GHG emissions in the growing seasons of each year were calculated by linear interpolation between sampling events using the trapezoidal rule (Goldberg et al., 2010). In addition to the cumulative GHG emissions between the first and the last sampling event, the GHG emissions from 1st June to the first sampling and from the last sampling to 30th September were taken into consideration.

Statistical analysis was carried out using R (version 3.4.3) (R Core Team, 2017), and graphs were drawn using OriginPro 9.8.0.200. Final *p* values were Bonferroni adjusted to mitigate the risk of type I error. A significance level of  $\alpha = 0.05$  was used for all statistical tests.

### 3. Results

#### 3.1 Soil properties

During the two years of the growing season mesocosm experiment, the SWC varied from 63.5% to 81.1% and was only significantly affected by the water table levels (Table 1 and 2). The higher WT levels significantly increased the SWC in both years. The soil pH varied within the range 7.3 to 7.8 and was only significantly affected by N deposition. Large variability of SOC was observed within the range 215.9 g kg<sup>-1</sup> to 296.1 g kg<sup>-1</sup>, and both the WT and N treatments showed significant effects on SOC, without any interactive effects. Compared with the control treatment without N deposition, N deposition increased SOC by 1.4% to 31.1% (except WT<sub>0</sub> N<sub>160</sub> in 2019). Soil TN varied between 12.9 g kg<sup>-1</sup> and 19.1 g kg<sup>-1</sup> and was elevated by N deposition, whereas no significant response to the WT treatments was observed. N deposition enhanced soil TN by 1.3% to 110.5% compared to the N<sub>0</sub> treatment at each WT level.

Table 1. Soil properties in the different water table (WT) treatments and nitrogen deposition (N) levels in 2018 and 2019.

		2018			2019		
		WT <sub>-30</sub>	WT <sub>0</sub>	WT <sub>10</sub>	WT <sub>-30</sub>	WT <sub>0</sub>	WT <sub>10</sub>
SWC (%)	N <sub>0</sub>	65.8±2.5	67.4±2.6	69.8±1.5	62.8±1.7	70.4±1.0	81.1±1.6
	N <sub>20</sub>	66.8±1.8	74.2±1.6	74.2±1.9	63.5±2.0	71.3±0.4	79.0±2.0
	N <sub>40</sub>	67.3±1.5	73.2±3.0	71.0±1.7	64.4±1.9	71.6±1.9	79.8±1.6
	N <sub>80</sub>	67.4±2.2	72.0±0.8	73.1±1.1	67.8±0.8	69.6±1.4	77.4±2.1
	N <sub>160</sub>	64.4±1.3	68.0±2.2	72.8±1.8	68.1±0.7	72.2±1.8	81.1±1
pH	N <sub>0</sub>	7.6±0	7.7±0.1	7.7±0.1	7.7±0	7.8±0.1	7.8±0.1
	N <sub>20</sub>	7.5±0	7.7±0	7.4±0.1	7.7±0.1	7.5±0.2	7.6±0.1

	N <sub>40</sub>	7.3±0	7.6±0.1	7.6±0.1	7.5±0.1	7.6±0.2	7.7±0.2
	N <sub>80</sub>	7.6±0	7.4±0.1	7.5±0.1	7.4±0.1	7.6±0.1	7.4±0
	N <sub>160</sub>	7.5±0	7.6±0	7.3±0.1	7.5±0.1	7.4±0.1	7.5±0
SOC (g kg <sup>-1</sup> )	N <sub>0</sub>	231.3±5.4	237±24.3	246.6±16.0	215.9±3.6	227.3±14.4	218.3±14.0
	N <sub>20</sub>	296.1±5.7	285.8±8.4	279.2±23.4	228.7±9.6	249.9±12.0	273.3±11.8
	N <sub>40</sub>	292.3±14.1	281.2±18.7	262.8±20.9	241.8±6.7	281.0±17.8	253.3±17.5
	N <sub>80</sub>	265.4±17.7	294.3±7.7	291.4±9.3	240.9±12	230.6±10.8	286.2±9.1
	N <sub>160</sub>	275.6±7.0	276.8±10.1	266.8±32.4	254.6±18.2	226.8±13.7	251.5±19.1
TN (g kg <sup>-1</sup> )	N <sub>0</sub>	17.6±0.8	16.1±1.1	18.7±0.8	12.9±0.9	14.4±0.3	14.7±2.1
	N <sub>20</sub>	18.7±0.7	19.1±0.6	19.3±0.8	21.9±2.2	21.2±3.2	23.3±5.0
	N <sub>40</sub>	18.4±1.1	18.8±0.9	19.2±0.4	19.1±1.5	21.4±4.0	15.0±2.7
	N <sub>80</sub>	18.3±0.8	19.4±0.2	19.7±0.2	18.6±1.1	16.2±1.0	31.0±2.6
	N <sub>160</sub>	18.1±0.8	18.2±0.4	19.0±0.4	19.5±2.0	21.7±6.6	21.9±4.6

178 Each value represents mean ± SE (n=3). SWC, soil water content; SOC, soil organic carbon; TN, total nitrogen.

179

180 Table 2. Effects of water table (WT) and nitrogen (N) deposition levels and their interactions on soil properties using generalized  
181 least square (GLS) models.

	SWC		pH		SOC		TN	
	F	P	F	P	F	P	F	P
WT	19.4	<b>&lt;0.001***</b>	0.34	0.7103	9.92	<b>&lt;0.001***</b>	2.08	0.1319
N	0.64	0.6352	6.78	<b>&lt;0.001***</b>	5.18	<b>0.001**</b>	4.49	<b>0.002**</b>
WT×N	0.25	0.9807	0.35	0.944	0.91	0.5147	0.74	0.6526


182 Bold font denotes significant values. The statistical significance levels used were: \*: 0.01<P < 0.05; \*\*: 0.001< P < 0.01; \*\*\*:  
183 P<0.001. SWC: soil water content; SOC: soil organic carbon; TN: total nitrogen.

184

### 185 3.2 Methane

186 The Zoige alpine peatland acted as a net source of CH<sub>4</sub> in the WT<sub>0</sub> and WT<sub>10</sub> treatments throughout the two growing seasons,  
187 although the CH<sub>4</sub> flux was almost 0 in the WT<sub>-30</sub> treatment. Temporal variability of the CH<sub>4</sub> flux was observed (Figure 1). The  
188 cumulative CH<sub>4</sub> emissions of the growing season ranged from -0.26 to 29.26 g CH<sub>4</sub>-C m<sup>-2</sup> in 2018 and from -0.35 to 16.36 g CH<sub>4</sub>-  
189 C m<sup>-2</sup> in 2019 (Figure 2). During the two years, the WT treatments and their interaction with the N treatments had significant effects  
190 on the cumulative CH<sub>4</sub> emissions, while N deposition only had significant effects in 2019 (Table 3). Along the WT level gradient  
191 from WT<sub>-30</sub> to WT<sub>10</sub>, the cumulative CH<sub>4</sub> emissions increased markedly. The response of the cumulative CH<sub>4</sub> emissions to N  
192 deposition was non-linear under WT<sub>0</sub> and WT<sub>10</sub> conditions (Figure 3), with the highest value observed in the N<sub>20</sub> treatment.

193 Compared to the  $N_0$  treatment, the  $N_{80}$  and  $N_{160}$  treatments remarkably decreased the cumulative  $CH_4$  emissions by 36.5% to 97.4%,  
 194 while  $N_{40}$  was in the same order of magnitude as in the  $N_0$  treatment. The GAM results showed that the cumulative  $CH_4$  emissions  
 195 could be explained by N deposition for 55.9% under  $WT_0$  conditions and for 45.4% under  $WT_{10}$  conditions. The modelling results  
 196 also indicated that the critical thresholds for the highest cumulative  $CH_4$  emissions were  $14.41 \text{ g C m}^{-2}$  with  $20.9 \text{ kg ha}^{-1} \text{ yr}^{-1}$  N  
 197 deposition under  $WT_0$  conditions and  $21.60 \text{ g C m}^{-2}$  with  $16.2 \text{ kg ha}^{-1} \text{ yr}^{-1}$  N deposition under  $WT_{10}$  conditions.



199 Figure 1. Temporal variation of growing-season  $CH_4$  and  $N_2O$  fluxes under five levels of nitrogen deposition (0, 20, 40, 80 and  
 200  $160 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) and three water table levels in 2018 and 2019. Error bars represent the SE ( $n=3$ ).

201




Figure 2. Effects of nitrogen deposition levels on cumulative CH<sub>4</sub> and N<sub>2</sub>O emissions at three water table levels during the growing seasons in 2018 and 2019. Error bars represent the SE (n=3). Different letters above the bars indicate statistically significant differences ( $P < 0.05$ ).

Table 3. The individual and interactive effects of water table (WT) and nitrogen (N) deposition levels on cumulative CH<sub>4</sub> and N<sub>2</sub>O emissions in 2018 and 2019 using generalized least square (GLS) models.

	CH <sub>4</sub>		N <sub>2</sub> O	
	F	P	F	P
2018				
WT	24.88	<b>&lt;0.001***</b>	36.68	<b>&lt;0.001***</b>
N	1.37	0.27	239.38	<b>&lt;0.001***</b>
WT×N	15.15	<b>&lt;0.001***</b>	4.28	<b>0.002**</b>
2019				
WT	615.89	<b>&lt;0.001***</b>	351.26	<b>&lt;0.001***</b>
N	5.99	<b>0.001**</b>	989.75	<b>&lt;0.001***</b>
WT×N	18.01	<b>&lt;0.001***</b>	5.23	<b>&lt;0.001***</b>

Bold font denotes significant values. The statistical significance levels used were: \*:  $0.01 < P < 0.05$ ; \*\*:  $0.001 < P < 0.01$ ; \*\*\*:  $P < 0.001$ .




Figure 3. The relationship between cumulative greenhouse gas (CH<sub>4</sub> and N<sub>2</sub>O) emissions and nitrogen deposition levels at three water table levels. A linear model was used to estimate cumulative N<sub>2</sub>O emission at five nitrogen deposition levels, while a generalized additive model (GAM) was used to assess the response of cumulative CH<sub>4</sub> emission to nitrogen deposition levels. The statistical significance levels used were: \* =  $P < 0.05$  and  $> 0.01$ ; \*\* =  $P < 0.01$  and  $> 0.001$ ; \*\*\*  $P < 0.001$ ; NS = not significant ( $P > 0.05$ ). The shaded areas indicate 95% confidence intervals.

### 3.3 Nitrous oxide

The Zoige alpine peatland acted as a net N<sub>2</sub>O source during the growing seasons, the N<sub>2</sub>O fluxes showing clear temporal variability in 2018 and 2019. The N<sub>2</sub>O flux tended to be highest in early September 2018 and in mid-August 2019, while the lowest flux was observed at the start or the end of the growing seasons (Figure 1). The cumulative N<sub>2</sub>O emissions ranged from 0.02 to 0.49 g N<sub>2</sub>O-N m<sup>-2</sup> in 2018 and from 0.05 to 0.50 g N<sub>2</sub>O-N m<sup>-2</sup> in 2019. The cumulative N<sub>2</sub>O emissions were significantly affected by the WT levels, N deposition and their interaction (Table 3). N deposition significantly increased the cumulative N<sub>2</sub>O emission by 28.9% to 1974.6%, most significantly in the N<sub>160</sub> treatment. However, there was no clear effect of WT levels on N<sub>2</sub>O emissions. We observed a significantly positive and linear relationship between the cumulative N<sub>2</sub>O emissions and N application doses, and the slope and intercept of the linear correlation depended on the WT levels (Figure 3). The linear results also showed that the 1 kg N ha<sup>-1</sup> addition increased the cumulative N<sub>2</sub>O emission by 0.0024, 0.0028 and 0.0023 g N<sub>2</sub>O-N m<sup>-2</sup> under WT<sub>-30</sub>, WT<sub>0</sub> and WT<sub>10</sub> conditions, respectively.

## 4. Discussion

### 4.1 Effects of WT and N deposition on CH<sub>4</sub> emission

The cumulative CH<sub>4</sub> emissions from the Zoige alpine peatland in our study ranged from -0.35 to 29.26 g CH<sub>4</sub>-C m<sup>-2</sup> across the two years, which is in the same order of magnitude as the cumulative CH<sub>4</sub> emissions (25.4-29.6 g CH<sub>4</sub>-C m<sup>-2</sup>) from an alpine wetland of the Qinghai-Tibetan Plateau in a previous study (Wang et al., 2017). As expected, WT levels had a significant positive effect on CH<sub>4</sub> emissions, with higher WT levels increased CH<sub>4</sub> emissions. This corresponds with the results of previous studies, due to the manipulative effects of WT levels on the soil redox conditions (Hoyos-Santillan et al., 2019; Wang et al., 2017). With higher WT levels, SWC increased and likely formed more anaerobic conditions conducive to CH<sub>4</sub> production, leading to elevated CH<sub>4</sub> emissions (Evans et al., 2021; Hoyos-Santillan et al., 2019; Zhang et al., 2020).

In the current study, a moderate level of N deposition positively stimulated the CH<sub>4</sub> emissions, but subsequently the positive effect declined with further N addition. However, this non-linear effect of N deposition was only observed at high WT levels, and the close to zero emissions of CH<sub>4</sub> at low WT levels were not affected by N deposition. To our knowledge, the non-linear effects of N deposition on CH<sub>4</sub> fluxes in the upland ecosystems has been reported in numerous studies (Li et al., 2012; Lafuente et al., 2020; Qu et al., 2021), however, there have been very few attempts in alpine peatlands. Song et al. (2013) reported that the N addition (0-240 kg N ha<sup>-1</sup> yr<sup>-1</sup>) showed non-linear positive effects on CH<sub>4</sub> fluxes in a wetland with the highest CH<sub>4</sub> flux at 60 kg ha<sup>-1</sup> yr<sup>-1</sup> N addition, but not reaching a significant level. Wu et al. (2022) conducted a global meta-analysis and found the N addition (0-300 kg N ha<sup>-1</sup> yr<sup>-1</sup>) could consistently increase the CH<sub>4</sub> emissions in the global wetlands, but the highest value occurred at 50-100 kg N ha<sup>-1</sup> yr<sup>-1</sup> N addition. The results of the aforementioned studies were not in agreement with the present study, due to the differences between the threshold values and the inhibitory or promotive effects of further N addition on CH<sub>4</sub> emission. However, in accordance with the previous studies, the relatively low level of N addition probably alleviated N constraints on microbial metabolism in soils and increased CH<sub>4</sub> production (Currey et al., 2009; Deng et al., 2019). With the further N addition, large amounts of available NO<sub>3</sub><sup>-</sup> likely led to negative and inhibitory effects on the methanogenic activity due to the competition of NO<sub>3</sub><sup>-</sup>-reducing bacteria with methanogens (Liu et al., 2020).

The interactive effects of WT levels and N deposition on the cumulative CH<sub>4</sub> emissions were distinct in our study (Table 3 and Figure 3). We found that the WT levels were more likely to determine the direction and magnitude of CH<sub>4</sub> emissions from alpine peatlands than N deposition. This is in line with the previous study (Evans et al., 2020), which indicated that WT was the overriding factor to control GHG emissions from the peatlands at a global scale. The N deposition non-linearly affected the CH<sub>4</sub> emissions, and the scenario for maximum CH<sub>4</sub> emissions was roughly ca. 20 kg N ha<sup>-1</sup> yr<sup>-1</sup>, which could be slightly changed by the WT levels. As far as we know, this is the first mesocosm experiments in an alpine peatland comparing the interactive effects of multi-level N deposition and WT levels on CH<sub>4</sub> emissions, which makes it challenging to put in context the scope of these results, and we have not found any theory in the previous studies which could best explain the underlying mechanism. We speculate that the N deposition

262 supplied N substrate and WT levels were associated with N utilization by microorganisms. Precisely, the higher WT levels promoted  
263 diffusion of the added N in the water-filled soil pore, and N thus becoming readily accessible in the microbial process to support  
264 CH<sub>4</sub> production (Wang et al., 2017).

#### 265 4.2 Effects of WT and N deposition on N<sub>2</sub>O emission

266 The mesocosms in the Zoige alpine peatland were consistently acting as N<sub>2</sub>O sources, and the cumulative N<sub>2</sub>O emissions (0.02-  
267 0.49 g m<sup>-2</sup> in 2018 and 0.05-0.50 g m<sup>-2</sup> in 2019) did not show significant differences between the two years. The cumulative N<sub>2</sub>O  
268 emission from the Zoige peatland in our study was relatively higher than that in a previous report (0.08-0.2 g m<sup>-2</sup>), which focused  
269 on the drainage or lower water table level (Cao et al., 2018). Unlike the CH<sub>4</sub> emissions, we did not observe a clear pattern for the  
270 effects of WT levels on N<sub>2</sub>O emissions in our study. This is different from the numerous previous studies, which all confirmed the  
271 positive effects of WT on N<sub>2</sub>O emissions in peatlands, due to its positive effects on anaerobic denitrification and N<sub>2</sub>O production  
272 (Gao et al. 2014; Regina et al., 1999; Regina et al., 1996). However, Wang et al. (2017) reported that elevated WT levels from  
273 drained to inundated conditions had no effects on N<sub>2</sub>O fluxes in an alpine wetland of the Qinghai-Tibetan Plateau, which is in line  
274 with our study. The possible reason for this could be due to the large differences of habitat types, soil properties or precipitations  
275 among the regional heterogeneous peatlands.

276 N deposition stimulated N<sub>2</sub>O emissions across the two years in our study, and same results were also found in the previous  
277 reports (Gao et al., 2014; Deng et al., 2019). The N deposition increased soil TN ( $F = 4.49, P = 0.002$ ) in our study and is likely to  
278 supply more N substrate (NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>) in soil (Zhu et al., 2020). The consequently increased N substrate could potentially activate  
279 the microbial process of N<sub>2</sub>O production and increase N<sub>2</sub>O emissions (Yue et al., 2021). Additionally, contrary to the CH<sub>4</sub> emissions,  
280 the N deposition showed consistently linear effects on N<sub>2</sub>O emissions in the peatland, regardless of the WT levels. This is not in line  
281 with a previous study (Song et al., 2013), which indicated that N addition ranging from 0-120 kg N ha<sup>-1</sup> yr<sup>-1</sup> did not show significant  
282 effects on N<sub>2</sub>O emissions in peatlands, while high level (240 kg N ha<sup>-1</sup> yr<sup>-1</sup>) largely increased the N<sub>2</sub>O emissions. However, a global  
283 meta-analysis showed that N<sub>2</sub>O emissions from wetlands demonstrated significant positive response to N enrichment (Deng et al.,  
284 2019), which agreed with our study. The possible explanation could be that the current level of N deposition in the Zoige peatland  
285 did not meet the N-saturation stage, after which the sensitivity of N<sub>2</sub>O emission from the ecosystem to further N addition would  
286 decline (Gomez-casanovas et al., 2016).

287 The N<sub>2</sub>O emission from the alpine peatland was likely primarily determined by N deposition rather than by WT levels. We  
288 confirmed the occurrence of an interactive effect of WT and N deposition on N<sub>2</sub>O emissions, but it was neither synergistic nor  
289 antagonistic. N deposition had linear positive effects on N<sub>2</sub>O emission, and WT level did not alter this linear relationship but slightly  
290 changed the slope and intercept. The N<sub>2</sub>O emissions could be generated from both the aerobic nitrification and anaerobic  
291 denitrification processes. This is confirmed by the previous study (Bateman et al., 2005), which indicate that the denitrification was  
292 the major contributor to N<sub>2</sub>O emission at water-filled pore space (WFPS) > 60% while nitrification was the major one at WFPS <

60%. The overall production of N<sub>2</sub>O could not be uniquely manipulated by the WT levels, though the WT levels could influence the redox conditions and the microbial processes. Additionally, the Zoige peatland being a N-limited ecosystem (Squeo et al., 2006), we observed the overwhelming effects of N deposition on N<sub>2</sub>O emission in the peatland, rather than WT levels.

The growing-season N<sub>2</sub>O emission in the current study increased by 0.0023-0.0028 g N<sub>2</sub>O-N m<sup>-2</sup> in response to the additional 1 kg NH<sub>4</sub>NO<sub>3</sub>-N ha<sup>-1</sup> yr<sup>-1</sup> deposition. This is slightly lower than the levels from previous studies (Gong et al., 2019), which showed that 1 kg annual N ha<sup>-1</sup> addition led to an increase of ca. 0.0076 g N<sub>2</sub>O-N m<sup>-2</sup> during the growing season in a boreal peatland. This could be attributed to the relatively low air temperature at this particular alpine peatland, which hampered the microbial N<sub>2</sub>O production. Furthermore, IPCC (2013) suggested that the default emission factor (the fraction of nitrogen added that is released as N<sub>2</sub>O) is 1%, indicating that 1 kg annual N ha<sup>-1</sup> addition may increase N<sub>2</sub>O emissions by 0.01 kg N<sub>2</sub>O-N ha<sup>-1</sup> (or 0.001 g N<sub>2</sub>O-N m<sup>-2</sup>). The relatively higher emission factor for N<sub>2</sub>O-N in our study was probably due to the high dose of N addition.

### 4.3 Implications and limitations

To what extent the magnitude of non-CO<sub>2</sub> emissions from alpine peatlands could be upscaled in response to the increasing N deposition, and how this effect interacts with the WT levels is still high uncertain. Our study demonstrated that the CH<sub>4</sub> and N<sub>2</sub>O emissions from the alpine peatland system could reach a new state if the projected increases in N deposition continued, in particular the CH<sub>4</sub> emission could be disproportionally increased. In the Zoige peatland, the current level of N deposition was 1.08-17.81 kg N·ha<sup>-1</sup> yr<sup>-1</sup> (Han et al., 2019), and the N deposition was predicted to be a possible doubling or tripling by the end of the century (Lamarque, 2005). Given the highest CH<sub>4</sub> emissions occurred at the level of ca. 20 kg N ha<sup>-1</sup> yr<sup>-1</sup> deposition and the consistent linear correlation between N<sub>2</sub>O emissions and N deposition levels in our study, we infer that the CH<sub>4</sub> and N<sub>2</sub>O emissions had not yet peaked, and the increasing N deposition could still result in the alpine peatland becoming a potential hotspot of GHG emissions in the future. Moreover, the fluctuation of WT levels resulting from anthropogenic drainage or rewetting could affect the GHG emissions but not suppress the N deposition-induced trend. However, our study is limited to a specific habitat, and thus the upscaling of this conclusion contains large uncertainty. To the best of our knowledge, very few advances heretofore have been achieved to elaborate this general conclusion. We believe that our results are useful for predicting the GHG emissions from alpine peatlands in response to the climate change and anthropogenic activities in the future.

It should be noted that some levels of N deposition (80 or 160 kg N ha<sup>-1</sup> yr<sup>-1</sup>) in our study were much higher than the local N deposition (1.08-17.81 kg N ha<sup>-1</sup> yr<sup>-1</sup>). This should not affect our general conclusion, because the non-linear and linear effects of N deposition on CH<sub>4</sub> and N<sub>2</sub>O emissions (respectively) were primarily dependent on the low levels of N deposition (0-40 kg N ha<sup>-1</sup> yr<sup>-1</sup>), and the higher levels did not alter the relationship pattern. Meanwhile, note must be made that we did not measure the net ecosystem CO<sub>2</sub> exchange and wintertime GHG fluxes, which might hamper estimating the annual carbon budget from GHG emissions and SOC change. However, our study focused on the growing-season non-CO<sub>2</sub> emissions from the peatland at different WT levels under the future scenarios of increasing N deposition, and also the non-growing-season GHG emissions had only a minor

324 contribution to the yearly budget due to the low temperature and microbial activities (Peng et al., 2019). In addition, the low  
325 frequency of GHG sampling in 2018 could cause uncertainties in the temporal variation and cumulative emissions of CH<sub>4</sub> and N<sub>2</sub>O,  
326 and this might result in bias in the present result. The monthly measurements of GHG fluxes from peatlands have already been  
327 found in the previous study (Cao et al., 2018), and we also increased the sampling frequency in 2019 to better support the current  
328 conclusion. However, further monitoring of GHG fluxes from the peatland would still be required to eliminate the uncertainties.

## 329 5. Conclusion

330 This study explored the response of non-CO<sub>2</sub> GHG emissions in the alpine peatland to the increasing N deposition at different  
331 WT levels. We found that CH<sub>4</sub> emissions were determined by N deposition, WT levels and their interactive effects. A modest input  
332 of N deposition and high WT levels both stimulated CH<sub>4</sub> emissions. N<sub>2</sub>O emissions were remarkably sensitive to N deposition,  
333 which consistently and linearly increased the N<sub>2</sub>O emissions, irrespective of WT levels. The projected increasing N deposition  
334 suggested an inevitable risk for higher CH<sub>4</sub> and N<sub>2</sub>O emissions from the alpine peatland in the future.

## 335 Data availability

336 All data are available from the corresponding author by request.

## 337 Author contributions

338 **Wantong Zhang:** Conceptualization, Data curation, Software, Writing - original draft, Writing - review & editing, Validation,  
339 Formal analysis. **Zhengyi Hu:** Conceptualization, Supervision. **Joachim Audet:** Data curation, Supervision, Writing – review &  
340 editing. **Thomas A. Davidson:** Supervision, Data curation, Writing – review & editing. **Enze Kang:** Investigation. **Xiaoming**  
341 **Kang:** Investigation, Project administration. **Yong Li:** Investigation. **Xiaodong Zhang:** Investigation, Project administration.  
342 **Jinzhi Wang:** Investigation, Conceptualization, Project administration, Writing - review & editing.

## 343 Funding

344 This study was financially supported by the National Natural Science Foundation of China (41877421 and 31770511).

## 345 Competing interests

346 The contact author has declared that neither they nor their co-author has any competing interests.

## 347 References

- 348 Audet J., Elsgaard L., Kjaergaard C., Larsen S. E. , Hoffmann C. C.: Greenhouse gas emissions from a Danish riparian wetland  
349 before and after restoration, *Ecol. Eng.*, 57, 170-182. <https://doi.org/10.1016/j.ecoleng.2013.04.021>, 2013.
- 350 Bateman E. J., Baggs E.M.: Contributions of nitrification and denitrification to N<sub>2</sub>O emissions from soils at different water-filled  
351 pore space. *Biology and Fertility of Soils* 41:379-388. <https://doi.org/10.1007/s00374-005-0858-3>, 2005
- 352 Cao R., Xi X., Yang Y., Wei X., Wu X. , Sun S.: The effect of water table decline on soil CO<sub>2</sub> emission of Zoige peatland on eastern

- 353 Tibetan Plateau: A four-year in situ experimental drainage, *Appl. Soil Ecol.*, 120, 55-61.  
 354 <https://doi.org/10.1016/j.apsoil.2017.07.036>, 2017.
- 355 Cao R., Chen Y., Wu X., Zhou Q., Sun S.: The effect of drainage on CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions in the Zoige peatland: a 40-month  
 356 in situ study. *Mires Peat*, 21, 1-15. <https://doi.org/10.19189/MaP.2017.OMB.292>, 2018.
- 357 Chen H., Zhu Q., Peng C., Wu N., Wang Y., Fang X., Gao Y., Zhu D., Yang G., Tian J., Kang X., Piao S., Ouyang H., Xiang W.,  
 358 Luo Z., Jiang H., Song X., Zhang Y., Yu G., Zhao X., Gong P., Yao T., Wu J.: The impacts of climate change and human  
 359 activities on biogeochemical cycles on the Qinghai-Tibetan Plateau, *Global Change Biol.*, 19, 10, 2940-55.  
 360 <https://doi.org/10.1111/gcb.12277>, 2013.
- 361 Chen Q., Wang S., Ni Z., Guo Y., Liu X., Wang G., Li H.: No-linear dynamics of lake ecosystem in responding to changes of  
 362 nutrient regimes and climate factors: Case study on Dianchi and Erhai lakes, China, *Sci. Total Environ.*, 781.  
 363 <https://doi.org/10.1016/j.scitotenv.2021.146761>, 2021.
- 364 Cui L., Kang X., Li W., Hao Y., Zhang Y., Wang J., Yan L., Zhang X., Zhang M., Zhou J., Kardol P.: Rewetting decreases carbon  
 365 emissions from the Zoige alpine peatland on the Tibetan plateau, *Sustainability*, 9, 6. <https://doi.org/10.3390/su9060948>,  
 366 2017.
- 367 Currey P. M., Johnson D., Sheppard L. J., Leith I. D., Toberman H., Van Der Wal R., Dawson L. A., Artz R. R. E.: Turnover of  
 368 labile and recalcitrant soil carbon differ in response to nitrate and ammonium deposition in an ombrotrophic peatland,  
 369 *Global Change Biol.*, 16, 8, 2307-2321. <https://doi.org/10.1111/j.1365-2486.2009.02082.x>, 2009.
- 370 Deng L., Huang C., Kim D. G., Shangguan Z., Wang K., Song X., Peng C.: Soil GHG fluxes are altered by N deposition: New data  
 371 indicate lower N stimulation of the N<sub>2</sub>O flux and greater stimulation of the calculated C pools, *Global Change Biol.*  
 372 <https://doi.org/10.1111/gcb.14970>, 2019.
- 373 Evans C. D., Peacock M., Baird A. J., Artz R. R. E., Burden A., Callaghan N., Chapman P. J., Cooper H. M., Coyle M., Craig E.,  
 374 Cumming A., Dixon S., Gauci V., Grayson R. P., Helfter C., Heppell C. M., Holden J., Jones D. L., Kaduk J., Levy P.,  
 375 Matthews R., McNamara N. P., Misselbrook T., Oakley S., Page S., Rayment M., Ridley L. M., Stanley K. M., Williamson  
 376 J. L., Worrall F., Morrison R.: Overriding water table control on managed peatland greenhouse gas emissions, *Nature.*,  
 377 <https://doi.org/10.1038/s41586-021-03523-1>, 2021.
- 378 Frolking S., Talbot J., Jones M. C., Treat C. C., Kauffman J. B., Tuittila E.-S., Roulet N.: Peatlands in the Earth's 21st century  
 379 climate system, *Environ. Rev.*, 19, 371-396. <https://doi.org/10.1139/a11-014>, 2011.
- 380 Gao Y., Chen H., Zeng X.: Effects of nitrogen and sulfur deposition on CH<sub>4</sub> and N<sub>2</sub>O fluxes in high-altitude peatland soil under  
 381 different water tables in the Tibetan Plateau, *Soil Sci. Plant Nutr.*, 60, 3, 404-410.  
 382 <https://doi.org/10.1080/00380768.2014.893812>, 2014.
- 383 Goldberg S. D., Knorr K.-H., Blodau C., Lischeid G., Gebauer G.: Impact of altering the water table height of an acidic fen on N<sub>2</sub>O  
 384 and NO fluxes and soil concentrations, *Global Change Biol.*, 16, 1, 220-233. <https://doi.org/10.1111/j.1365-2486.2009.02015.x>, 2010.
- 386 Gomez-Casanovas N., Hudiburg T. W., Bernacchi C. J., Parton W. J., Delucia E. H.: Nitrogen deposition and greenhouse gas  
 387 emissions from grasslands: uncertainties and future directions, *Global Change Biol.*, 22, 4, 1348-1360.  
 388 <https://doi.org/10.1111/gcb.13187>, 2016.
- 389 Gong Y., Wu J., Vogt J., Le T. B.: Warming reduces the increase in N<sub>2</sub>O emission under nitrogen fertilization in a boreal peatland,  
 390 *Sci. Total Environ.*, 664, 72-78. <https://doi.org/10.1016/j.scitotenv.2019.02.012>, 2019.
- 391 Han Y., Dong S., Zhao Z., Sha W., Li S., Shen H., Xiao J., Zhang J., Wu X., Jiang X., Zhao J., Liu S., Dong Q., Zhou H., Yeomans  
 392 J. C.: Response of soil nutrients and stoichiometry to elevated nitrogen deposition in alpine grassland on the Qinghai-  
 393 Tibetan Plateau, *Geoderma*, 343, 263-268. <https://doi.org/10.1016/j.geoderma.2018.12.050>, 2019.
- 394 Hoyos-Santillan J., Lomax B. H., Large D., Turner B. L., Lopez O. R., Boom A., Sepulveda-Jauregui A., Sjögersten S.: Evaluation  
 395 of vegetation communities, water table, and peat composition as drivers of greenhouse gas emissions in lowland tropical  
 396 peatlands, *Sci. Total Environ.*, 688, 1193-1204. <https://doi.org/10.1016/j.scitotenv.2019.06.366>, 2019.
- 397 IPCC: *Climate Change: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*  
 398 *Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.
- 399 Järveoja J., Peichl M., Maddison M., Soosaar K., Vellak K., Karofeld E., Teemusk A., Mander Ü.: Impact of water table level on  
 400 annual carbon and greenhouse gas balances of a restored peat extraction area, *Biogeosciences*, 13, 9, 2637-2651.

- 401 <https://doi.org/10.5194/bg-13-2637-2016>, 2016.
- 402 Juutinen S., Moore T. R., Bubier J. L., Arnkil S., Humphreys E., Marinck B., Roy C., Larmola T.: Long-term nutrient addition  
403 increased CH<sub>4</sub> emission from a bog through direct and indirect effects, *Sci. Rep.*, 8, 1, 3838.  
404 <https://doi.org/10.1038/s41598-018-22210-2>, 2018.
- 405 Lafuente A., Recio J., Ochoa-Hueso R., Gallardo A., Perez-Corona M. E., Manrique E., Duran J.: Simulated nitrogen deposition  
406 influences soil greenhouse gas fluxes in a Mediterranean dryland, *Sci. Total Environ.*, 737, 139610.  
407 <https://doi.org/10.1016/j.scitotenv.2020.139610>, 2020.
- 408 Laine A. M., Mehtatalo L., Tolvanen A., Frolking S., Tuittila E. S.: Impacts of drainage, restoration and warming on boreal wetland  
409 greenhouse gas fluxes, *Sci. Total Environ.*, 647, 169-181. <https://doi.org/10.1016/j.scitotenv.2018.07.390>, 2019.
- 410 Lamarque J. F.: Assessing future nitrogen deposition and carbon cycle feedback using a multimodel approach: Analysis of nitrogen  
411 deposition, *J. Geophys. Res.*, 110, D19. <https://doi.org/10.1029/2005jd005825>, 2005.
- 412 Liu Z., Li D., Zhang J., Saleem M., Zhang Y., Ma R., He Y., Yang J., Xiang H., Wei H.: Effect of simulated acid rain on soil CO<sub>2</sub>,  
413 CH<sub>4</sub> and N<sub>2</sub>O emissions and microbial communities in an agricultural soil, *Geoderma*, 366.  
414 <https://doi.org/10.1016/j.geoderma.2020.114222>, 2020.
- 415 Pekár S., Brabec M., Bshary R.: Marginal models via gls: A convenient yet neglected tool for the analysis of correlated data in the  
416 behavioural sciences, *Ethology*, 122, 8, 621-631. <https://doi.org/10.1111/eth.12514>, 2016.
- 417 Peng, H., Guo, Q., Ding, H., Hong, B., Zhu, Y., Hong, Y., et al.: Multi-scale temporal variation in methane emission from an alpine  
418 peatland on the Eastern Qinghai-Tibetan Plateau and associated environmental controls. *Agr. Forest. Meteorol.*, 276-277,  
419 <https://doi.org/10.1016/j.agrformet.2019.107616>, 2019.
- 420 Qu S., Xu R., Yu J., Li F., Wei D., Borjigidai A.: Nitrogen deposition accelerates greenhouse gas emissions at an alpine steppe site  
421 on the Tibetan Plateau, *Sci. Total Environ.*, 765, 144277. <https://doi.org/10.1016/j.scitotenv.2020.144277>, 2021.
- 422 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.  
423 URL <https://www.R-project.org/>, 2017
- 424 Regina K., Nykanen H., Silvola J., Martikainen P. J.: Fluxes of nitrous oxide from boreal peatlands as affected by peatland type,  
425 water table level and nitrification capacity, *Biogeochemistry*, 35:401-418. <https://doi.org/10.1007/bf02183033>, 1996
- 426 Regina K., Silvola J., Martikainen P. J.: Short-term effects of changing water table on N<sub>2</sub>O fluxes from peat monoliths from natural  
427 and drained boreal peatlands. *Global Change Biol.*, 5, 183-189. <https://doi.org/10.1046/j.1365-2486.1999.00217.x>, 1999.
- 428 Saiz E., Sgouridis F., Drijfhout F. P., Peichl M., Nilsson M. B., Ullah S.: Chronic atmospheric reactive nitrogen deposition  
429 suppresses biological nitrogen fixation in peatlands, *Environ. Sci. Technol.*, 55, 2, 1310-1318.  
430 <https://doi.org/10.1021/acs.est.0c04882>, 2021.
- 431 Song C., Wang L., Tian H., Liu D., Lu C., Xu X., Zhang L., Yang G., Wan Z.: Effect of continued nitrogen enrichment on greenhouse  
432 gas emissions from a wetland ecosystem in the Sanjiang Plain, Northeast China: A 5 year nitrogen addition experiment, *J.*  
433 *Geophys. Res.-Biogeo.*, 118, 741-751. <https://doi.org/10.1002/jgrg.20063>, 2013
- 434 Squeo F. a., Warner B. g., aravena R., espinoza D.: Bofedales: high altitude peatlands of the central Andes, *Rev. Chil. Hist. Nat.*,  
435 79, 245-255. <http://dx.doi.org/10.4067/S0716-078X2006000200010>, 2006.
- 436 Tiemeyer B., Albiac Borraz E., Augustin J., Bechtold M., Beetz S., Beyer C., Drosler M., Ebli M., Eickenscheidt T., Fiedler S.,  
437 Forster C., Freibauer A., Giebels M., Glatzel S., Heinichen J., Hoffmann M., Hoper H., Jurasinski G., Leiber-Sauheitl K.,  
438 Peichl-Brak M., Rosskopf N., Sommer M., Zeitz J.: High emissions of greenhouse gases from grasslands on peat and other  
439 organic soils, *Global Change Biol.*, 22, 12, 4134-4149. <https://doi.org/10.1111/gcb.13303>, 2016.
- 440 Wang H., Yu L., Zhang Z., Liu W., Chen L., Cao G., Yue H., Zhou J., Yang Y., Tang Y., He J. S.: Molecular mechanisms of water  
441 table lowering and nitrogen deposition in affecting greenhouse gas emissions from a Tibetan alpine wetland, *Global Change*  
442 *Biol.*, 23, 2, 815-829. <https://doi.org/10.1111/gcb.13467>, 2017.
- 443 Wanyama L., Pelster D. E., Butterbach-Bahl K., Verchot L. V., Martius C., Rufino M. C.: Soil carbon dioxide and methane fluxes  
444 from forests and other land use types in an African tropical montane region, *Biogeochemistry*, 143, 171-190.  
445 <https://doi.org/10.1007/s10533-019-00555-8>, 2019.
- 446 Wilson D., Farrell C. A., Fallon D., Moser G., Muller C., Renou-Wilson F.: Multiyear greenhouse gas balances at a rewetted  
447 temperate peatland, *Global Change Biol.*, 22, 12, 4080-4095. <https://doi.org/10.1111/gcb.13325>, 2016.
- 448 Wood S. N.: *Generalized Additive Models: An Introduction with R* (2nd edition). Chapman and Hall/CRC.

- 449 <https://doi.org/10.1201/9781315370279>, 2017.
- 450 Wu J., Cheng X., Xing W., Liu G.: Soil-atmosphere exchange of CH<sub>4</sub> in response to nitrogen addition in diverse upland and wetland  
451 ecosystems: A meta-analysis, *Soil Biol Biochem*, <https://doi.org/164.10.1016/j.soilbio.2021.108467>, 2022.
- 452 Yang G., Chen H., Wu N., Tian J., Peng C., Zhu Q., Zhu D., He Y., Zheng Q., Zhang C.: Effects of soil warming, rainfall reduction  
453 and water table level on CH<sub>4</sub> emissions from the Zoige peatland in China, *Soil Biol. Biochem.*, 78, 83-89.  
454 <https://doi.org/10.1016/j.soilbio.2014.07.013>, 2014.
- 455 Yang G., Wang M., Chen H., Liu L. F., Wu N., Zhu D., Tian J. Q., Peng C. H., Zhu Q. A., He Y. X.: Responses of CO<sub>2</sub> emission  
456 and pore water DOC concentration to soil warming and water table drawdown in Zoige Peatlands, *Atmos. Environ.*, 152,  
457 323-329. <https://doi.org/10.1016/j.atmosenv.2016.12.051>, 2017.
- 458 Yu Z., Loisel J., Brosseau D. P., Beilman D. W., Hunt S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophys.*  
459 *Res. Lett.*, 37, 13. <https://doi.org/10.1029/2010gl043584>, 2010.
- 460 Yue P., Zuo X., Li K., Cui X., Wang S., Misselbrook T., Liu X.: The driving effect of nitrogen-related functional microorganisms  
461 under water and nitrogen addition on N<sub>2</sub>O emission in a temperate desert, *Sci. Total Environ.*, 772, 145470.  
462 <https://doi.org/10.1016/j.scitotenv.2021.145470>, 2021.
- 463 Zhang L., Hou L., Guo D., Li L., Xu X.: Interactive impacts of nitrogen input and water amendment on growing season fluxes of  
464 CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O in a semiarid grassland, Northern China, *Sci. Total Environ.*, 578, 523-534.  
465 <https://doi.org/10.1016/j.scitotenv.2016.10.219>, 2017.
- 466 Zhang W., Wang J., Hu Z., Li Y., Yan Z., Zhang X., Wu H., Yan L., Zhang K., Kang X.: The Primary Drivers of Greenhouse Gas  
467 Emissions Along the Water Table Gradient in the Zoige Alpine Peatland, *Water Air Soil Poll.*, 231, 5.  
468 <https://doi.org/10.1007/s11270-020-04605-y>, 2020.
- 469 Zhang Y., Wang G., Wang Y.: Changes in alpine wetland ecosystems of the Qinghai-Tibetan plateau from 1967 to 2004, *Environ.*  
470 *Monit. Assess.*, 180, 1-4, 189-99. <https://doi.org/10.1007/s10661-010-1781-0>, 2011.
- 471 Zhu J., Chen Z., Wang Q., Xu L., He N., Jia Y., Zhang Q., Yu G.: Potential transition in the effects of atmospheric nitrogen deposition  
472 in China, *Environ. Pollut.*, 258, 113739. <https://doi.org/10.1016/j.envpol.2019.113739>, 2020.
- 473