

Modeling nitrous oxide emissions from agricultural soil incubation experiments using CoupModel

Jie Zhang¹, Wenxin Zhang², Per-Erik Jansson³, Søren O. Petersen¹

¹Department of Agroecology, iClimate, Aarhus University, Tjele, Denmark

²Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

³Department of Sustainable Development, Environmental Science and Engineering, KTH Royal Institute of Technology, Stockholm, Sweden

Correspondence to: Jie Zhang (jiezhang@agro.au.dk)

Formatted: Swedish (Sweden)

Abstract. Efforts to develop effective climate mitigation strategies for agriculture require methods to estimate nitrous oxide (N₂O) emissions from soil. Process-based biogeochemical models have been often used for such estimations but were mainly tested with field- and large-scale measurements estimates, while the sensitivity and uncertainty of model applications to incubation experiments are less investigated. In this study, results from a short-term (43-day) factorial incubation experiment were used to investigate the ability of a process-oriented model (CoupModel) was used to estimate simulate N₂O and carbon CO₂ fluxes, and soil mineral nitrogen (N) dynamics. This study identified the sensitivities of model parameters when estimating three output variables using a contents in a short-term (43-day) factorial incubation experiment (16 treatments). A global sensitivity analysis (GSA) approach. Our results, “Morris screening”, was applied to quantify parameter sensitivity. The GSA suggested that important a higher number of sensitive parameters was associated with N₂O flux estimates and that inter-treatment variations in parameter sensitivities were distinguished by soil moisture levels or NO₃⁻ content, and residue types. Important parameters regarding N₂O flux estimates were linked to the decomposability of soil organic matter (e.g., organic C pool sizes) and the denitrification process (e.g., Michaelis constant and denitrifier respiratory rates). The After calibration, the model was able to simulate low better captured temporal variations and magnitude of gas fluxes and mineral N in unamended soils than in residue-amended soils. Low-magnitude daily and cumulative N₂O fluxes were well simulated with model errors (MEs) close to zero, but the model tended to underestimate N₂O fluxes as observed daily values increased over 0.1 g N m⁻² day⁻¹. Besides, the response of N₂O, where the major mismatch was due to limited success of the model to describe the high emissions to soil moisture during the first few days after crop residue addition. A larger uncertainty was not well reflected in the model, probably related to the indirect involvement of soil moisture response function in the denitrification process also seen in the magnitude of pulse emissions by the posterior simulations. We also evaluated ancillary variables regarding N cycling, which indicates indicated that more frequent measurements and additional types of observed data such as soil oxygen content and the microbial sources of emitted N₂O are required to further evaluate model performance and biases. The current description of the N cycling process in the model may not consistently represent the temporal scale of nitrification and denitrification processes behind N₂O emissions. The major challenges for calibration are were associated with high sensitivities of denitrification parameters to initial soil moisture abiotic conditions and the

~~instantaneous~~ residue amendment. ~~For the development of process-based models, we suggest there is a need to address soil heterogeneity, and to revisit current subroutines of moisture response functions.~~ Model structure uncertainties and improved modeling practices in the context of incubation experiments were discussed.

1 Introduction

The potent greenhouse gas nitrous oxide (N_2O) has been estimated to be responsible for about 7 % of the overall global radiative forcing by long-lived greenhouse gases (World Meteorological Organization, 2021). N_2O emissions from the agricultural sector account for 60-70 % of the total anthropogenic emissions of this gas (Davidson and Kanter, 2014; Syakila and Kroeze, 2011). ~~To provide a scientific basis for developing achievable climate mitigation strategies. However, it is persistently difficult to reduce the uncertainties of N_2O emission estimates, and one reason is associated with the high spatiotemporal variability of N_2O (Kravchenko et al., 2017). To provide a scientific basis for developing achievable climate mitigation strategies, an~~ improved understanding of N_2O production in agricultural soils and quantification of N_2O emissions are urgently needed.

N_2O emissions from agricultural soils are driven by a suite of microbiological processes among which nitrification and denitrification predominate as sources of N_2O . The factors directly regulating nitrification and denitrification activity are the availability of mineral nitrogen (N), oxygen, and degradable carbon (C) sources used by denitrifying organisms (Wijler and Delwiche, 1954). Indirect controls include soil temperature, moisture, pH, and soil texture. During nitrification, where ammonia (NH_3 , at equilibrium with ammonium NH_4^+) is oxidized to nitrate (NO_3^-), a small proportion of N may be lost as N_2O (Firestone and Davidson, 1989). Nitrification mainly occurs in well-aerated soils with moderate water content (Goreau et al., 1980; Li et al., 1992; Parton et al., 1996). In contrast, denitrification is a microbial process that occurs under anaerobic conditions where NO_3^- is reduced to gaseous N. ~~Soil C substrates are electron donors for denitrification, but they are also a sink for oxygen that leads to anaerobic microsites (Sommer et al., 2004), and finally NO_3^- is used as electron acceptors. Soil C substrates are electron donors for denitrification, but they are also a sink for oxygen that can lead to anaerobic microsites (Parkin, 1987) where NO_3^- is used as an electron acceptor for microbial respiration.~~ Nitrifying and denitrifying bacteria are most active to produce N_2O in environments with abundant N relative to assimilatory demands by other microorganisms or plants (Firestone and Davidson, 1989), as is often the case following ~~input~~ soil amendment of fertilizers, manure, or crop residues ~~to the soil~~.

Farming practices influence the ~~potential for~~ interactions between microbial, physical, and chemical processes in the soil, ~~and the uncertainties associated with predicting N_2O emissions from agricultural soils amended with plant residues such as cover crops or green manure can be particularly high.~~ Incorporation of crop residues can reduce NH_3 losses and enhance degradation compared to leaving residues at the soil surface, but ~~at the increased soil~~ ~~same time the N input, elevated~~ water holding capacity, and ~~high local~~ oxygen demand ~~locally of residues~~ may stimulate the development of anaerobic microsites

and bacterial denitrification activity (Kravchenko et al., 2018; Kuzyakov and Blagodatskaya, 2015). ~~Mechanical disturbances via~~ Also, mechanical disturbances caused by tillage may influence soil properties (e.g., porosity, aggregate size distribution, solute and gas diffusivities) and microbial enzyme activities, ~~with subsequent changes in the magnitude of associated with~~ N₂O emissions (Grandy and Robertson, 2006).

The quantification of N₂O emissions from agroecosystems ~~is constrained by logistical~~ based on field experiments often meets challenges ~~associated with logistics, data gaps, and long-term resource~~ availability support (e.g. ~~analytical, manpower, equipment, and budgets~~ research budget). Process-oriented biogeochemical models, e.g., DNDC (Li et al., 1992), DayCent (Parton et al., 1996), APSIM (Keating et al., 2003), and CoupModel (Jansson and Moon, 2001), ~~have been developed~~ can be a complementary approach to ~~quantify N₂O emission, partly compensate for these limitations. In addressing~~ the application of ~~above-mentioned challenges. When applying~~ process-based models, available *in-situ* measurements can be used to infer model parameters and allow simulation of soil N transformations and N₂O emissions ~~at~~ on temporal and spatial scales beyond the monitoring sites, but accurately simulating the magnitude and temporal variability of N₂O fluxes under contrasting ~~contexts still poses~~ field conditions remains a challenge. ~~Those~~ While models may provide reasonable estimates of ~~annual~~ N₂O emissions ~~from soils in a narrow context, usually at specific sites and at annual time scales, but,~~ they become less successful at finer time resolution (e.g., diurnal time steps) ~~and at sites different from the pre-calibrated ones. This), and this~~ represents a barrier ~~into~~ evaluating the effects of agricultural land use and management on greenhouse gas emissions (Brilli et al., 2017). Such model errors are often attributed to physical and biogeochemical processes being inadequately represented, ~~which calls for the improvement of~~ suggesting there is a need to improve process descriptions beyond parameter optimization (Abdalla et al., 2010; Brilli et al., 2017; Gaillard et al., 2018; Uzoma et al., 2015).

Process-based models attempt to reproduce the most relevant physical and biogeochemical processes through understanding grounded in the best available ~~theory at the time they were developed, after which some new empirical adjustments were~~ gradually added knowledge. CoupModel, used in the current investigation, has a high level of detail on ~~soil physical~~ the interaction between biotic and abiotic ~~components~~ processes and has adopted ~~details of~~ submodules of nitrification, denitrification, and gas ~~fluxes~~ transport from the ~~DNDC-model~~ DNDC (Li et al., 2000; Norman et al., 2008). The description of N₂O emissions, including the links between soil environmental factors and biological reactions, is based on a series of hypotheses and results generated from both field measurements and laboratory incubations studies (Li et al., 2000), and the algorithms and parameterization of microbial growth and ~~death~~ decay dynamics were specifically supported by the latter. ~~While our understanding regarding decomposition and denitrification has advanced in recent decades, the incorporation of state-of-the-art knowledge into process-based models has lagged behind. To test the description of N₂O emissions, it is necessary to apply the model to results from properly~~ Hence, controlled laboratory experiments, where the impact of ill-defined pedo-climatic conditions on model predictability can be minimized (Brilli et al., 2017). ~~This may reveal causal relationships behind gas production and transport in a microcosm representing the ecosystem, and suggest new paths for model development.,~~ may be useful to optimize the model prediction of N₂O emissions following residue amendment.

Formatted: English (United Kingdom)

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: English (United Kingdom)

The application of process-based models has often been challenged by the paucity of prior information and measurements compared to the model's demands, and this is also the case when applying a model to incubation experiments. One widely-used model calibration method to bridge the gap between model requirements and available data, and to quantify parameter uncertainties, is "generalized likelihood uncertainty estimation (GLUE)" (Beven and Binley, 1992). During model calibration, uncertainty analysis can help assess ~~whether how robust~~ the model ~~performance is good enough compared to~~ ~~reasonably describes~~ the ~~requirement of the applied use of the model measurements~~ and ~~to evaluate identify~~ possible ~~biases in~~ ~~simulations~~ ~~reasons to explain the model errors~~ (U.S. Environmental Protection Agency, 2009). This may be facilitated by applying a Global Sensitivity Analysis (GSA), which can rank the sensitivities of parameters so that the model calibration can focus on the relatively more sensitive parameters (Vezzaro et al., 2012), and thereby the model's uncertainties can be more efficiently constrained. While model processes and performance have been extensively documented, in many studies N₂O emissions alone were used to train and test the subroutines of nitrification and denitrification (Chen et al., 2008). Evaluation under controlled conditions and with ancillary measurements is noticeably lacking, which makes it difficult to ~~identify model structure limitations assess the submodules related to C and N processes, for example, decomposition, nitrification, and denitrification~~. Thus, a first step in understanding model performance may be an evaluation using new datasets that contain different variables linked to N cycling based on targeted laboratory experiments. To our knowledge, no previous study has attempted a systematic sensitivity and uncertainty analysis in the prediction of N₂O emissions based on ~~laboratory incubation results soil incubation experiments. The present study focused on the role of reactive C and N for N₂O emissions and used simulations of targeted experiments to identify key drivers. Simulating N₂O dynamics in a short-term laboratory study may be considered to zoom in on a single field operation, in this case the incorporation of crop residues by standard tillage operations.~~

For this work, we ~~selected used~~ CoupModel ~~which to simulate incubation experiments~~. CoupModel has a flexible setting of soil layer thickness down to a scale of mm and is proper to study soil physical processes at the scale of incubation experiments. Besides, CoupModel has integrated ~~options for modules that implement parameter calibration and~~ uncertainty estimation and performance evaluation analysis (Jansson, 2012). ~~It has a flexible setting of soil layer thickness down to a scale of mm, which is proper to study soil physical processes at the scale of incubation experiments. Data~~ The calibration data sets used in the model were obtained from a 43-day laboratory incubation using a factorial-based design with various crop residue practices and abiotic factors (Taghizadeh-Toosi et al., 2021). Specifically, our objectives were (i) to conduct a global sensitivity analysis for parameters in a ~~model~~ CoupModel setup that ~~can simulate represents soil N cycling under different and C processes for incubation treatments~~; (ii) to calibrate the model and ~~quantify the assess model~~ uncertainty in the estimates of N₂O and CO₂ emissions, ~~soil ammonium and nitrate under different residue applications~~; and (iii) to discuss any ~~model limitations identified in the model optimization with short-term incubation experiments~~ and suggest directions for future model ~~improvement development~~. We hypothesized that the model is able to simulate the daily and cumulative N₂O emissions under contrasting environments in incubated soil cores. Furthermore, we hypothesized that ~~it would be difficult to~~

130 ~~constrain the only a few~~ parameters ~~of in~~ a complex model ~~could be constrained~~ to an unambiguous solution with limited laboratory measurements. ~~Our findings can thus help diagnose potential causes of model-measurement discrepancies concerning C and N processes, and specify conditions for which model and data collection need to be improved.~~

2 Materials and methods

2.1 Laboratory incubation experiment

135 In spring 2018, soil used for the experiment was collected from the 0-20 cm tilled layer at the Lönnstorp Field Station, Sweden. Red beets had been grown in the previous year with no cover crop during winter. The soil is sandy loam (61.8 % sand, 22.4 % silt, and 15.8 % clay) with a pH of 6.18, C content of 15 g kg⁻¹, and N content of 1.49 g kg⁻¹. After ~~colleitionsampling~~, the soil was partially dried, stored at -20 °C, and thawed ~~onethe~~ day before ~~sieving and being sieved to < 6 mm for use for in~~ the experiment.


140 Treatments were prepared ~~with under~~ four different soil conditions ~~regarding the including two~~ moisture ~~level levels~~ (i.e., 40 or 60 % WFPS) and ~~two~~ nitrate content ~~levels~~ (i.e., no nitrate addition or addition of KNO₃ to 100 mg NO₃-N kg⁻¹ dry wt. soil). Soil cores were prepared by stepwise packing 1 cm layers of soil to a density of 1.25 g cm⁻³ in cylinders ~~to the with a~~ height of 8 cm, at each step adding deionized water or a KNO₃ solution. ~~The Following moisture and nitrate adjustments, the soil treatments were was~~ pre-incubated for one week at 15 °C. The experiment involved two different crop residues, red clover (RC) and winter wheat (WW). RC residues had a C/N ratio of 17.9, and a moisture level corresponding to 80 % of the fresh weight. The WW residues had a C/N ratio of 90.9, and the moisture content corresponded to 20 % of the fresh weight. WW residues had a higher proportion of lignin and ash (11.7 %) than RC residues (5.1 %). ~~In On Day 0 of~~ the experiment, RC or WW residues were either mixed ~~at a rate in an amount~~ of 0.04 g DM cm⁻² into the soil from 0-4 cm depth and then repacked, or residues were placed as a layer at 4 cm depth; only results from the mixed treatments were used in the present study. Incubations with RC and WW took place sequentially, and therefore each residue treatment had its own set of unamended controls. Thus, in total 16 ~~incubation~~ treatments ~~from the incubation experiment (i.e., 40 % vs. 60 % WFPS; ambient nitrate vs. nitrate addition; RC vs. WW; and soil amended with residues vs. unamended controls)~~ were used for this modeling study, ~~including unamended soils (as controls) and soil residue mixtures from either red clover or winter wheat.~~


155 All cylinders were covered at both ends with perforated plastic caps and incubated at 15 °C for up to 43 days. Gas sampling for N₂O and CO₂ flux measurements took place ten times, i.e., on day 1, 3, 6, 9, 13, 16, 22, 29, 36, and 43. Gas concentrations were determined by gas chromatography. Additionally, nitric oxide (NO) fluxes were quantified in four selected treatments set up separately. Soil mineral N pools in all treatments were measured at four destructive samplings after 1, 6, 22, and 43 days of incubation. Further details about the experimental treatments, preparations, and analytical methods are given by Taghizadeh-Toosi et al. (2021).

160 **2.2 Model description and simulation setup**

2.2.1 CoupModel

This study used CoupModel v6.1, which can be downloaded from <http://coupmodel.com>. A detailed description of CoupModel can be found in Jansson & Karlberg (2010). The main structure of the model is a one-dimensional vertical soil profile with user-defined layer thickness and subdivisions. The current setup of CoupModel includes a number of components, of which the following processes are linked to N₂O emissions (Fig. 1): (i) soil organic matter (SOM) decomposition and mineralization; (ii) nitrification and nitrifier growth; (iii) denitrification and denitrifier growth; and (iv) gas diffusion between soil layers and internal exchange of N trace gases between aerobic and anaerobic micro-sites. In the nitrification subroutine, CoupModel accounts for response functions of soil temperature, soil moisture, mineral N concentration, and pH. For denitrification, each step in the chain of denitrification is explicitly ~~calculated~~ simulated, and denitrifier activity is directly influenced by soil temperature, pH, nitrogen oxides, and anaerobic fraction. The anaerobic soil volume fraction is calculated using the “anaerobic balloon” concept, as implemented in the DNDC model (Li et al., 2000; Norman et al., 2008).





175 **Figure 1: A conceptual diagram of major C and N processes in the current setup of CoupModel. The details of parameters and equations in each C or N process can be found in Table S2.**

2.1.2 Simulation settings

180 The modeled soil profile consisted of a single soil layer with a depth of 4 cm for the control treatments and 4.2 cm for the mixed-treatments with residues, allowing a 2 mm increment owing to residue amendment as observed in the experiment. We only simulated the upper half of the 8 cm soil core to have proper gas boundary conditions since the model only allows external gas exchange at the upper boundary side of the soil profile, although whereas in the experiments both ends of the cylinder were exposed to air and the two halves were identical for control soils. For the water process, it was assumed that there was no evaporation from the surface and no vertical water flow across the lower boundary. Constant, and a constant temperature was set for the upper and lower boundaries, in accordance with incubation conditions. The model was initialized based on the measured soil water content, soil porosity, temperature, pH, total organic C and N, and NO_3^- -N, and NH_4^+ -N of the incubated soil cores. The dynamics of SOM dynamics were simulated with first-order kinetics using three pools (litter1, litter2, and humus). Considering there were no explicit pools designed for crop residue addition, we assigned the rapidly decomposable SOM and metabolically easily metabolized residue materials components (e.g., sugars and proteins) to litter1, the

190 moderately decomposable SOM and structural residue materials (e.g., lignin and other fibers) to litter2, and the resistant SOM to humus. ~~When initializing the organic matter pools, we took the soil pool sizes from published papers and assumed that they were already close to a steady state at the start of incubations (see the description in the following paragraphs), and hence the changes observed during incubation were assumed to be mainly caused by the decomposition of the residues introduced.~~ For simulating gas transport, we selected the steady-state mode where the oxygen content within the soil profile ~~is a trade-off was derived from the balance~~ between soil oxygen consumption and ~~surface air~~ diffusive supply ~~from surface air,~~ and ~~N trace~~ ~~the transient storage term of~~ gases ~~are directly lost to~~ ~~in~~ the ~~ambient soil~~ air ~~from the layer in which they are generated was not considered here.~~

195
200 *Calibration datasets* – Measurements used for model calibration were N₂O flux, CO₂ flux, NO flux, NO₃⁻-N content, and NH₄⁺-N content. As the gas fluxes and mineral N content in the upper part with soil-residue mixtures and the lower part with bulk soil were not analyzed separately, ~~we~~ ~~it~~ ~~was~~ assumed that soil C and N turnover in the lower, unamended part was identical to that of control treatments ~~in order~~ to create datasets for the residue-amended part for modeling. Specifically, the amounts of mineral N and gas fluxes recorded on individual sampling days in the controls were divided by two and subtracted from the values recorded in residue-amended soil.

205 *Initial values* – (1) Mineral N: Since the mineral N content in the unamended control soil changed little during incubation and the mineral N content in crop residues was negligible, the initial NH₄⁺-N and NO₃⁻-N values for the control and residue-amended treatments were taken as the measurement in control soil on day 1. (2) Soil moisture: For the control treatments, the initial volumetric water content was calculated from the water-filled pore space (WFPS) levels of 40 or 60 % ~~according to~~ ~~and~~ the total porosity of 0.53. For the residue treatments, the initial volumetric water content was calculated from the moisture content of soil and crop residues (Taghizadeh-Toosi et al., 2021). (3) Organic matter pools: The partitioning of soil organic C between litter1, litter2, and humus was defined by the ratio 0.02:0.54:0.44 (Gijsman et al., 2002). For crop residues, the ~~metabolic~~-fraction of ~~easily metabolized~~ organic C was calculated from the lignin/N ratio: ~~Metabolic fraction =~~ 0.85 - 0.013 (lignin/N) (Gijsman et al., 2002), and hence the organic C allocation between litter1 and litter2 had a ratio of 0.82:0.18 for RC and 0.55:0.45 for WW. The allocation of organic N ~~into~~ different pools followed the pattern of C and the C/N ratios (Table S5).

215 A summary of calibration data sets can be found in Table S1 ~~in~~ ~~of~~ the supplement, in which cumulative gas emissions were estimated by ~~linearly interpolating~~ ~~linear interpolation~~ between sampling dates and ~~integrating~~ ~~integration of~~ the area under emission curves; ~~and,~~ ~~while~~ average mineral N ~~contents~~ were calculated by dividing the integrated values by the sampling period. The results for nitrate in soil cores with residues were not included due to high uncertainty in the calculations that was probably caused by solute transport between the unamended and amended soil layers, as observed in a related incubation experiment using some of the same soil and residue treatments conducted by ~~Lashermes et al. (2021)~~; ~~Lashermes et al. (2022)~~.

2.3 Model sensitivity and uncertainty analysis

2.3.1 Global sensitivity analysis

Given uncertain prior information, the study used Morris screening (Morris, 1991) for a global sensitivity analysis to identify the most important input parameters and process parameters affecting N_2O fluxes. We included seven input parameters related to the characteristics of soil and crop residues (i.e. soil porosity, residue porosity, soil pH, and sizes of organic C pools), with realistic ranges of uncertainty intervals considered. Besides, we considered 45 process parameters involved in the relevant model processes. These parameters are listed in Table S4 parameters to which estimates of N_2O fluxes were sensitive. We included seven input parameters related to the characteristics of soil and crop residues (i.e., soil porosity, crop residue porosity, soil pH, and organic C pool sizes), and the parameter ranges were defined around the mean values of measurements or estimations (Table S4). For parameters supported by measurements, i.e., soil porosity and pH, the ranges were within 25 % of the mean values to represent realistic micro-scale variations in the laboratory setup. The residue porosity was estimated with a bulk density of 0.18 g cm^{-3} and a dry density of 1.3 g cm^{-3} (Lam et al., 2008; Zhang et al., 2012), and with a wider uncertainty range of 40 % due to compressibility. Bulk soil porosity and crop residue porosity were used to calculate the soil-residue mixture bulk density used in the model input, and this calculation is included in the Supplement. The ranges of organic pool fractions were derived from literature values for cultivated soil where the humus pool size (SOC_h) had an uncertainty range of ca. 40 % and the labile carbon pool size (SOC_l) varied within two orders of magnitude considering the marginal fraction. The range of crop residue organic pool sizes was taken from the constraints of the different estimated fractions of the two crop residues.

Besides the seven input parameters, we considered 45 process parameters involved in the relevant model processes. Part of the parameter ranges (e.g., tO_{10a} , d_{pHrate} , CM_m) adopted in the study were based on the most relevant applications of CoupModel, experimental studies, and other process models as shown in Table S4. Ranges for the remaining model-specific parameters, including those involved in nitrification and denitrification, could not be derived from the existing literature, and in the absence of better prior information, we adopted the default ranges set by the model.

The Morris screening method is a commonly used for sensitivity analysis technique, based on an efficient sampling strategy for performing a randomized calculation of one-factor-at-a-time (OAT) sensitivity analysis. This method can be viewed as a compromise between a simple OAT approach and the more complex GSA methods (e.g., variance-based approaches) as it provides a good approximation to the global sensitivity measure of the parameters at an affordable computational cost. Furthermore, it was considered excessive and unnecessary in the present study to adopt a more detailed analysis given the limited data availability available measurements.

The elementary effect (EE) was estimated by comparing the variation of the model's output y^j , with the variation of a given parameter θ_i , according to Eq. (1). The number of iterations n was set to 50, and the optimal perturbation factor Δ was set to 2/3 by dividing the input space into four levels (Morris, 1991). To allow comparison across outputs, the EE was then

standardized by using the standard deviation of the model factor and the standard deviation of the output (SEE, Eq. (2)). The significance of the impact of parameters was tested by comparing the mean of the SEE of those parameters to twice the standard error (*sem*, Eq. (3)) (Sin et al., 2009). If the input factor lies outside this range, it is said to have a significant effect on the output. The codes used in the analysis were adapted from (Sin et al., 2009).

$$EE_i^j = \frac{y(\theta_1^j, \theta_2^j, \dots, \theta_i^j + \Delta_{OPT}, \dots, \theta_{m-1}^j, \theta_m^j) - y(\theta_i^j)}{\Delta} \quad (1)$$

$$SEE_i^j = EE_i^j \cdot \frac{\sigma_{\theta_i}}{\sigma_y} \quad (2)$$

$$sem_i = \pm \frac{\sigma(SEE_i^j)}{\sqrt{n}} \quad (3)$$

The GSA was performed to the model-evaluation measure root mean square error (RMSE) for three variables: N₂O flux, CO₂ flux, and soil NH₄⁺ which had relatively complete measurement data sets. By applying the sensitivity analysis to the likelihood measure, the main factors that drive the model runs with a good fit to data could be identified (Ratto et al., 2001). The results from sensitivity analyses were further used to identify process parameters for inclusion in the uncertainty analysis due to their contribution to output variability.

2.3.2 Uncertainty analysis

Model calibration was conducted separately for each of the 16 treatments to give more flexibility in model parameterization. ~~The calibration was carried out with reference to five measurement variables, namely N₂O flux, CO₂ flux, NH₄⁺ content, NO₃⁻ content, and NO flux (only four treatments), using the “generalized likelihood uncertainty estimation” (GLUE) technique (Beven and Binley, 1992).~~ The calibration was carried out with reference to five measurement variables, namely N₂O flux, CO₂ flux, NH₄⁺ content, NO₃⁻ content, and NO flux (only four treatments), using the GLUE method. The GLUE method does not seek the single best fit to the measured data but utilizes an ensemble of model simulations that represent equally good results using informal likelihood measures, ~~often mentioned as (also called acceptance criteria) to account for parameter equifinalities.~~ In this study, we first described the entire ensemble of model runs as prior runs and after applying selection criteria the selected ensemble of model runs was analyzed as posterior runs or behavioral runs. Based on the calculated sensitivity indices from Morris screening, a total of 26 process parameters were selected for calibration where the parameters with marginal SEEs were omitted and only one denitrifier growth parameter was kept in each step of the denitrification chain (see Table S4). These parameters were uniformly or log-uniformly distributed within the predefined ranges, from which 20,000 parameter sets were then randomly sampled for model runs. Out of these runs, whether a parameter set was accepted or not was based on the defined criteria, which in this study consisted of coefficient of

determination, R^2 , and mean error, ME. The latter is defined as $ME = E(O_i - S_i)$, where S_i and O_i are the ~~time-series-of-the~~ simulated and observed data- at each measurement time step.

280 The ME acceptance threshold of each variable was set to be around the average of daily measurements taking into account the different magnitudes of each variable (see Table S3). ~~In-The ME values of prior runs ME-values-were-for some measurement variables~~ often showed a skewed to-one-side (above distribution deviating from 0, meaning that the model may have systematically over- or below 0) while underestimated these variables. While setting the same threshold on both sides rejected most of the prior model runs, ~~and hence~~ the ME criterion on one side might be looser than the other. For N_2O emissions with marked peak fluxes, a combination of R^2 and ME was used for the selection of posterior parameters to
285 simulate the dynamics and magnitudes. An ensemble of ca. 50 posterior runs was selected with an acceptance rate of 0.25% based on prior simulations. The uncertainties of model predictions were quantified within the limits and posterior probability distributions of parameters.

290 Finally, to investigate whether the treatment effects ~~concerning of~~ soil moisture and nitrate ~~level~~ levels could be represented by a common parameterization, we ~~attempted to calibrate also calibrated~~ process parameters against measurements combined data ~~sets from multiple treatments where measurements~~ from every four treatments with the same residue application ~~were pooled-including~~ “40 % WFPS, -NO₃”, “40 % WFPS, +NO₃”, “60 % WFPS, -NO₃”, and “60 % WFPS, +NO₃”. The prior parameter ensembles used the same 20,000 parameter sets as the single-treatment calibration. Accordingly, the measurement datasets from the four treatments in each group were pooled and thus a larger data set for model evaluation was obtained.

295 The procedure of selecting behavioral runs followed the aforementioned approach based on ME and R^2 . A diagram that describes the analysis workflow for this study is presented in Fig. 2.




Figure 2: A schematic diagram for performing of the workflow of the study to analyze parameter sensitivity and uncertainty analyses of the CoupModel based on 16 treatments from an incubation experiment (see text for details).

3 Results

3.1 Sensitivity analysis







The results of Morris screening were evaluated by comparing the absolute SEEs concerning N₂O flux, CO₂ flux, and soil NH₄⁺ for individual treatments. Figure 3a-c lists all the parameters ranking in the top five SEEs across the 16 treatments. Parameter ranking was performed based on the absolute mean of SEEs – the higher the absolute value, the more important the parameter is was, as shown by the shade of color in Fig. 3a-c. In general, the a higher number of sensitive parameters identified as most influential for soil respiration (were found for N₂O flux (18) than CO₂ flux (8) and NH₄⁺ content showed robustness across treatments as they differed only slightly in their ranking with, respectively, eight and nine different parameters represented. In contrast(9). Also, more inter-treatment variation was variations in the SEEs of parameters were found for N₂O flux. For CO₂ flux and NH₄⁺, inter-treatment variations were found in the parameter ranking for N₂O

emissions with 18 different parameters represented relevant to soil moisture levels (e.g., $p_{\theta_{low}}$ and θ_{wilt}) or NO_3^- content ($d_{growthNO_3}$), or residue types (e.g., $f_{e,12}$, $f_{e,11}$, and cn_m). For N_2O , the parameters exhibiting relatively high SEE values for most treatments belonged to categories of SOM decomposition and denitrification (Table S2), including d_{effNO} , SOC_h , $d_{growthNO_3}$, cn_m , and d_{hrateN_2O} . The model input, SOC_h , characterizing the partitioning of SOM pools in the simulation, was found to be crucial in 13 out of 16 treatments. d_{effNO} represents the respiration of denitrifying bacteria based on NO, and it showed relatively large elementary effects for almost all treatments by directly regulating the reduction step from NO to N_2O . The parameter $d_{growthNO_3}$ describes the loss of NO_3^- from the anaerobic nitrogen pool due to microbial growth. d_{hrateN_2O} represents the N concentration for half rate in the denitrification process and is also known as the Michaelis constant of the enzyme (see n13, n15, n17, n19, and n20 in Fig. 1).

Parameters that had the greatest impact on CO_2 emissions were concentrated in the following: SOC_h , SOM decomposition rates (k_{12} , k_{11}), and the corresponding efficiencies ($f_{e,12}$, $f_{e,11}$). In addition, the two parameters $p_{\theta_{low}}$ and θ_{wilt} , controlling the lower limit of the soil moisture response function for the decomposition of organic matter (see Eq. (5.86) in Table S2), exhibited distinct influences for the treatments at the lower moisture level.

The main processes influencing the NH_4^+ content of the soil were identified as SOM decomposition and denitrification, and influential parameters included: cn_m , SOC_h , k_{12} , and $f_{e,12}$. The C/N ratio of microbes, cn_m , has an influence on the mineral N content by changing the magnitude and direction of soil mineralization/immobilization of nitrogen (see n1-n7 and n11 in Fig. 1). Soil porosity (θ_{soil}) had significant effects on some treatments, especially under higher moisture conditions. Besides, as a key intermediate of mineral nitrogen turnover, the content of NH_4^+ , was also influenced by denitrification-related parameters, such as ($d_{growthNO_3^-}$).

The average elementary effects (SEEs) across all treatments are shown in Fig. 3d-f. For the N_2O flux, all parameters were located inside the wedge indicating that none of these parameters showed a significant effect across all treatments despite their significance in individual treatments. In contrast, we found that the other two variables, CO_2 flux, and NH_4^+ content, were significantly affected by 15 and 28 parameters respectively. Moreover, all parameters showed non-linear effects on the outputs as revealed by their non-zero standard deviations, which suggested that simulated C and N processes did not solely depend on individual parameters but also on their interactions.






Figure 3: Sensitivity analyses for N_2O flux, CO_2 flux, and soil NH_4^+ content in all treatments. (a-c): Heatmaps that include all parameters ranked in the top five places for each treatment based on absolute values of standardized elementary effects (SEEs).

(d-f): Estimated mean and standard deviation of SEEs averaged across the 16 treatments, where the two lines drawn in each subplot correspond to twice the standard error (sem): $\mu_i \pm 2sem_i$ (see [Sect. Section 2.3.1](#)): if a factor is located inside the wedge, it indicates that its impact on the output is considered insignificant and vice versa.

3.2 Uncertainty analysis

3.2.1 Temporal dynamics of N₂O flux, CO₂ flux, and mineral N

The prior simulations generally showed mean errors for gas emissions and soil mineral N largely deviated from zero, and these biases were reduced in the posterior simulations for most model outputs (Fig. 4a-d). In the experimental treatments with RC amendment, N₂O emission rates were consistently low at 40 % WFPS but were markedly higher and peaked on day 3 at 60 % WFPS (Fig. 5a). The highest measured daily N₂O flux was 1.4 g N m⁻² day⁻¹ in the RC treatment with NO₃⁻ addition at 60 % WFPS. Similar patterns were observed for CO₂ emission rates, with emission peaks at an early stage of incubation (day 1 or day 3) and then followed by a decline. In treatments amended with WW, N₂O evolution rates were generally low compared to those with RC amendment, and showed higher rates at 60 % WFPS and in NO₃⁻ amended soil but treatment effects were generally minor. For CO₂ evolution, higher rates were detected by day 1, but there was also a secondary peak after 1-2 weeks. The control treatments of the WW residue incubations showed less CO₂ and N₂O release compared to the control treatments of the preceding RC incubations.

The prior models generally showed significantly biased mean errors in terms of gas emissions and soil mineral N, and their magnitude was reduced in the posterior models for most model outputs (Fig. 4a-d). For N₂O fluxes, most treatments amended with WW and corresponding controls did not show significant deviations from the observed fluxes. In contrast, in treatments amended with RC and corresponding controls, though the absolute MEs had been reduced, there were still significant deviations generally in the direction of underestimating the observed fluxes. For CO₂ emissions, 13 out of 16 treatments showed reduced mean errors in the behavioral models and half of the treatments showed insignificant deviations from the observed fluxes. For soil NH₄⁺ content, there was a severe overestimation for most prior models, but this was alleviated by posterior models, and seven treatments showed insignificant deviations from θ_{zero} after model calibration. For soil NO₃⁻ content in control treatments, the ME ranges of posterior runs were around zero while negative or positive biases existed, especially the former. The simulated evolution of associated variables is depicted in Fig. 5a-d and results are summarized below.

In the experimental treatments with RC amendment, N₂O emission rates were consistently low at 40 % WFPS but were markedly higher and peaked on day 3 at 60 % WFPS (Fig. 5a). The highest measured daily N₂O flux was 1.4 g N m⁻² day⁻¹ in the RC treatment with NO₃⁻ addition at 60 % WFPS. Similar patterns were observed for CO₂ emission rates, with emission peaks at an early stage of incubation (day 1 or day 3) and then followed by a decline. In treatments amended with WW, N₂O evolution rates were generally low compared to those with RC amendment. Higher rates were observed at 60 % WFPS and in NO₃⁻ amended soil, but treatment effects were generally minor. For CO₂ evolution, higher rates were detected by day 1,

370 ~~but there was also a secondary peak after 1-2 weeks for the WW treatments (Fig. 5b). The control treatments of the WW residue incubations showed less CO₂ and N₂O release compared to the control treatments of the preceding RC incubations. For soil mineral N, there was a consistent increase in NH₄⁺-N content by day 1 after residue application compared to the controls, and both NH₄⁺-N and NO₃⁻-N showed smaller variations than gas fluxes (Fig. 5c-d and Fig. S6c-d). The simulated evolution of associated variables is depicted in Fig. A₂O – The accepted~~
375 ~~5a-d for treatments at 60 % WFPS, and in Fig. S6a-d for treatments at 40 % WFPS, respectively, and the results are summarized below.~~


~~N₂O – The posterior~~ simulations (Fig. 5a and Fig. S6a) were able to represent the scenarios with low daily N₂O emissions (10⁻⁵-10⁻² g N m⁻² day⁻¹), while simulations failed to capture the large emission peaks (e.g., 1.4 g N m⁻² day⁻¹ and 0.13 g N m⁻² day⁻¹ for the two RC treatments with NO₃⁻ amendment), or the emission dynamics were reasonably simulated (e.g., R² > 0.4, see Table S3) but the peak values were lower than observed. The N₂O fluxes obtained from the model tended to increase
380 over time and generally agreed with the observed fluxes in the second part of the experiment. ~~Compared with residue-amended soils, the model better described the magnitude and stable trend of N₂O fluxes in control soils.~~

CO₂ – Overall, the ~~behavioral models~~ ~~posterior simulations~~ mimicked the measured dynamics and magnitude of CO₂ emissions well (Fig. 5b and Fig. S6b). There were overestimations or underestimations by the model, most pronounced in the early stage of incubation. By day 14, a second peak of respiration was observed for the WW treatments ~~that was but not simulated~~ ~~captured~~ by ~~the~~ posterior ~~models~~ ~~simulations~~.
385

NH₄⁺ – For RC residue treatments, ~~an increase of NH₄⁺ was observed within the first day, followed by~~ net N mineralization was observed from the early- to mid-stage of the incubation period, ~~followed by and then a declining trend, whereas decline.~~
390 ~~In contrast,~~ in the posterior ~~models in~~ ~~simulations for~~ three of four treatments, the simulated NH₄⁺ ~~predicted a trend of~~ ~~corresponded to an enhanced~~ net N mineralization throughout the incubation (Fig. 5c and Fig. S6c). Such a ~~continuously increasing trend~~ ~~continuous increase of soil NH₄⁺~~ also existed in the prior ~~run~~ ~~simulations~~, which would not be radically altered in the ~~behavioral models~~ ~~posterior simulations~~ by setting a stricter selection criterion for R². For WW residue and control treatments the measured NH₄⁺ content was at the detection limit, and the magnitude of the simulated NH₄⁺ content was either in line with the measurements, or a bit higher.


NO₃⁻ – The ~~change of~~ simulated daily NO₃⁻ content ~~generally captured well the magnitude of the measurements, and showed a declining trend~~ ~~small variations throughout the incubation period~~ for all control treatments, ~~with modeled values comparable to observed data. (Fig. 5d and Fig. S6d).~~
395 While in most of the control treatments except for the ones with high moisture and NO₃⁻ amendment, the observed NO₃⁻ levels remained stable or slightly increased during incubation ~~(Fig. 5d).~~ For the treatments amended with crop residues, the simulated NO₃⁻ content showed a ~~more clear downward trend~~ ~~noticeable decrease~~ throughout the period, consistent with NO₃⁻ being utilized as a substrate of denitrification in the simulation. Though no explicit measurement of NO₃⁻ within the residue-amended layer in the present experiment, the average NO₃⁻ within the entire soil core in RC treatments showed net consumption followed by a rebound (Taghizadeh-Toosi et al., 2021), consistent with
400


observations for the RC-amended layer in a comparable incubation experiment by Lashermes et al. (2021). ~~In view of this~~2022). Given that, the simulated NO_3^- content in residue treatments exhibiting a continuous decline was probably lower than the actual values.



410

Figure 4: Comparison of the ME ranges between prior simulations and posterior (accepted) simulations for (a) daily N_2O fluxes (a), (b) CO_2 fluxes (b), (c) soil NH_4^+ content (c), and (d) soil NO_3^- content (d). Blue. The blue color shows ME values not significantly different from zero by one-sample t -tests (significance level $\alpha = 0.05$). Treatment indices 1-4 represent treatments of mix RC, 5-8 for control RC, 9-12 for mix WW, and 13-16 for control WW, where treatment conditions are, in order: “40% WFPS, $-\text{NO}_3^-$ ”, “40% WFPS, $+\text{NO}_3^-$ ”, “60% WFPS, $-\text{NO}_3^-$ ”, and “60% WFPS, $+\text{NO}_3^-$ ”. No measured data for nitrate in residue treatments.





415 **Figure 5:** Simulated and measured daily (a) N_2O fluxes (a), (b) CO_2 fluxes (b), (c) soil NH_4^+ content (c), and (d) soil NO_3^- content (d) during the 43-day incubation at 60% WFPS. Scatter points represent measured data; and triangles with dashed lines represent simulated data (error bar: 95% confidence interval). Daily measurements presented were re-calculated from the data provided by Taghizadeh-Toosi et al. (2021).

Formatted: Not Superscript/ Subscript

3.2.2 Cumulative gas fluxes and average mineral N content

420 Model predictions of cumulative N_2O fluxes and CO_2 fluxes for the 16 treatments were significant and strongly correlated with the observed fluxes (Fig. 6 and Table 1). For N_2O , there was a bias towards consistent underestimation of high cumulative N_2O fluxes (slope bias $\beta_1 = 0.17$, $\text{ME} = -0.23$). Estimate of The estimated slope in of the linear regression for cumulative CO_2 flux approached $\beta_1 = 1$, indicating there was no consistent bias. For the average NH_4^+ and NO_3^- content, the estimated slopes were also close to unity, and the deviations between prediction and measurement, signified by the relative RMSEs (rRMSEs), were 46% and 11%, respectively. For the average NH_4^+ content in the low range, simulated values were found to overestimate the measured data (Fig. 6c).

425




Figure 6: Simulated and measured (a) cumulative N_2O fluxes (a), (b) cumulative CO_2 fluxes (b), (c) average NH_4^+ content (c), and (d) average soil NO_3^- content (d) during the 43-day incubation (error bar: 95 % confidence interval). Colors indicate different types of residue amendment and markers indicate different abiotic soil environments. Reference lines with a slope of 1.0 are shown on the graphs.

Formatted: Not Superscript/ Subscript

Table 1: Model evaluation of cumulative gas fluxes and average mineral N content from single-treatment calibration procedure and multi-treatment calibration procedure. The mean values of posterior models were compared with the observed data for 16 treatments. Units are valid for the statistics of ME and RMSE.

Calibration	Cumulative N_2O emission (g N m^{-2})		Cumulative CO_2 emission (g C m^{-2})		Average NH_4^+ content (g N m^{-2})		Average NO_3^- content (g N m^{-2})	
	Single-treatment	Multi-treatment	Single-treatment	Multi-treatment	Single-treatment	Multi-treatment	Single-treatment	Multi-treatment
ME	-0.23	0.10	-3.07	4.66	0.03	0.12	-0.11	-1.10
RMSE	0.82	0.71	17.2	12.1	0.06	0.50	0.41	0.50
rRMSE	275 %	238 %	34 %	24 %	46 %	411 %	11 %	14 %
Slope, β_1	0.17	0.56	0.93 ^a	0.97 ^a	1.15	2.13 ^a	1.07 ^a	1.08 ^a

Formatted: English (United Kingdom)


Intercept, β_0	0.02	0.23	0.64	6.16	0.01	-0.01	-0.36	-0.53
R ²	0.96	0.47	0.84	0.93	0.98	0.50	0.98	0.98


^a Values are not significantly different from one by one-sample *t*-tests (significance level $\alpha = 0.05$).

A regression of simulated cumulative N₂O flux residuals against observed data confirmed that underestimations were strongly ($R^2 = 0.92$) associated with the magnitude of observed N₂O fluxes (Fig. S1a). The negative slope of the regression indicated an underestimate of 0.83 g N₂O-N for every 1 g of observed N₂O-N per square meter. A regression of simulated cumulative N₂O flux residual against the residuals of other variables revealed that underestimations were not strongly associated with the residuals of simulated NH₄⁺ and NO₃⁻ (Fig. S1c and d). However, we observed that clustering of residuals concerning for mineral N existed in which underestimations of cumulative N₂O flux tended to occur when soil NH₄⁺ was overestimated and when soil NO₃⁻ was underestimated. Specifically, residuals for cumulative N₂O flux and soil NO₃⁻ were simultaneously underestimated in 53 % of the posterior runs as revealed by scatter points falling in the third quadrant; and underestimations of N₂O flux were accompanied by overestimations of soil NH₄⁺ in 41 % of the posterior runs by looking at scattering points in the fourth quadrant. ~~When only the subset of control treatments was analyzed, the clustering patterns became even more apparent (Fig. S2c and d).~~

3.2.3 Calibration by multiple treatments

Increasing the number of calibration treatments ~~led to reduced~~ resulted in lower uncertainty but, meanwhile, ~~poor~~ performances reduced the performance of posterior models for some treatments (Fig. 7). Cumulative N₂O fluxes were better simulated for the treatments with higher observed fluxes in each group, especially treatments at 60 % WFPS, but were overestimated for others with low observed fluxes. The regression between the mean simulated and measured N₂O flux only accounted for 47 % of the ~~variation~~ variations in the data, much lower than the level of 96 % in the single-treatment calibration procedure (Table 1). Simulated CO₂ and NO₃⁻ were generally close to the observed data. In the same group, simulated CO₂ fluxes ~~were not different~~ showed no difference between the two levels of soil NO₃⁻ ~~input but depended on~~ not for the level levels of moisture. Simulated soil NH₄⁺ showed a good agreement with the measured data in the low range of NH₄⁺ content, but had large model deviations for the four RC residue treatments; ~~(a log-scale is used in Fig. 6c and Fig. 7c).~~ The statistics rRMSE and RMSE are more sensitive to outliers compared to mean error (ME), and regarding NH₄⁺, their high values for NH₄⁺ (rRMSE: 411 % and RMSE: 0.50 in Table 1) would be reduced to 78 % and 0.01, respectively, after removing the four RC treatments.





465 **Figure 7: Simulated and measured (a) cumulative N₂O fluxes (a), (b) cumulative CO₂ fluxes (b), (c) average NH₄⁺ content (c), and (d) average soil NO₃⁻ content (d) during the 43-day incubation (error bar: 95 % confidence interval). Simulated results were obtained from multi-treatment calibration. Colors indicate different types of residue amendment and markers indicate different abiotic soil environments. Reference lines with a slope of 1.0 are shown on the graphs.**

3.2.4 Simulated oxygen status and N₂O sources

470 Simulated oxygen content in the soil cores was close to that of the ambient air, with the modeled volumetric oxygen content in air-filled pores ranging from 19.5 % to 20 % throughout the incubation period for all treatments independent of treatment (Table S7). Still, according to the model, denitrification-derived N₂O accounted for 76-100 % of the total cumulative emissions on average (Table S7).

475 In simulations, the 0-4 cm soil layer was treated as one uniform compartment, and this could have influenced model predictions. To investigate whether increasing the vertical resolution could improve or change the model performance predicted oxygen availability, the soil profile was uniformly divided into five uniform layers. The results showed that underestimations of high N₂O fluxes still existed after this change in the vertical stratified representation of the

model (Table S6). The simulated soil oxygen profiles were still predominantly aerobic for all treatments, as in the single-layer model, but showed stratification overby depth as depicted in Fig. 8. The oxygen level was the lowest at the beginning of incubation and then showed an increase over the period studied, mirroring the trends ofin CO₂ flux. Despite the overall aerobic conditions in the soil, the large proportion of denitrification-derived N₂O emissions was accompanied by the rapid growth of denitrifier biomass (data not shown).




Figure 8: Simulated O₂ content and volumetric anaerobic fraction by the multi-layer model for the treatment with the greatest soil respiration rate (i.e., RC treatment with NO₃⁻ addition at 60 % WFPS). The mean daily values from posterior runs were used here.

3.2.5 Calibrated parameters


Although the number of parameters used for calibration had been reduced by Morris screening, more than half of the 26 calibrated parameters still exhibited random distributions within the predefined ranges (Fig. S4). The many potential high inter-correlations between posterior parameters may be the reason that these explain why some parameters could not be constrained to an unambiguous solution. But five parameters were showing marked variability between treatments in their posterior distributions between treatments as depicted in Fig. 9, where the prior ranges of these parameters are indicated in the ordinate.

In most treatments, the parameter $d_{hrateN \times O_2}$ representing the half-saturation constant of N concentration for half rate in the denitrification process and also known as the Michaelis constant of the enzyme, was well constrained at the lower range of the parameter boundary within 50 mg N L⁻¹ in contrast to the mean of 250 mg N L⁻¹ in the prior range (Fig. 9). A low

495 $d_{hrateNO_3}$ relative to the physical concentration of NO_3^- resulted in a pronounced response of denitrifying bacteria activity to substrate availability (see Eq. (6.44) in Table S2). In some treatments which had no NO_3^- addition, i.e., treatment 3, 7, 9, and 13, the parameter showed more ~~diffused~~ diffuse distribution and higher medians compared to other treatments (Fig. 9). An enzyme with high $d_{hrateNO_3}$ relative to the concentration of substrate is not normally saturated with substrate and thus the rate of formation of product is substrate-limited.

500 In more than half of the treatments, the posterior distribution of cn_m , the microbial C/N ratio (cn_m) involved in calculations of mineralization and immobilization, was concentrated at around 10 on average. However, for some NO_3^- amended treatments which usually had higher N_2O emission rates, i.e., treatments 4, 6, 10, 12, and 14, the distribution of calibrated microbial C/N ratio was not well constrained but similar to the prior distribution with medians up to 20.

505 The parameter representing the rate coefficient for the decay of the litter carbon pool, k_{li} , generally showed higher values in WW treatments than controls, and its range for the high nitrate RC treatment at 60% WFPS (treatment 4) was markedly higher than other treatments, indicating the faster decomposition of labile organic matter. Besides, the efficiency of NO-based denitrifier respiration, d_{effNO} , showed a low-range distribution for treatment 4. Low d_{effNO} values induced a high respiration rate of denitrifiers ~~for~~ carrying out NO reduction (see Eq. (6.47) in Table S2). This treatment also exhibited a low-range distribution of the efficiency of SOM decomposition, $f_{e,li}$, associated with a high fraction of CO_2 production.



510

Figure 9: Variability in five calibrated parameters among the 16 treatments. The boxplots show the 25 % and 75 % percentiles as the tops and bottoms of the boxes, and the medians as the bold lines. Treatment indices 1-4 represent treatments of mix RC, 5-8 for control RC, 9-12 for mix WW, and 13-16 for control WW, where treatment conditions are, in order: “40_% WFPS, -NO₃”, “40_% WFPS, +NO₃”, “60_% WFPS, -NO₃”, and “60_% WFPS, +NO₃”.

515

4 Discussion

4.1 Sensitivity analysis

4.1 Simulation of decomposition and denitrification processes

The sensitivity analyses, investigating the impact of parameter uncertainties on model predictions, ~~revealed~~ showed the importance of parameter interactions and ~~connections~~ associations between different processes in the model (Fig. 3a-f).

520

Compared with CO₂ emissions and soil NH₄⁺ content, N₂O emissions were controlled by a larger ~~number~~ range of parameters related to decomposition and denitrification processes and had generally lower SEEs. According to Fig. 3a, ~~the sizes~~ reliable information about the size of humus and litter pools and their decomposition rates ~~were~~ will be critical in reducing the

uncertainty of ~~simulating~~ simulated N₂O emissions. ~~These results highlight the importance of reliable information on the initial state of the soil with respect to the composition, and recalcitrance of organic matter pools~~ this could be particularly true where part of the soil organic matter ~~within the soil cores could be~~ fresh input from residues. The results confirm the findings of previous experimental and modeling studies showing the importance of substrate heterogeneity for ~~simulations of~~ decomposition and denitrification processes (Brilli et al., 2017; Eusterhues et al., 2003; Sierra et al., 2011). However, ~~some~~ other studies (Dungait et al., 2012; Schmidt et al., 2011) ~~also~~ indicate that ~~the~~ chemical structure of organic molecules alone may not ~~control~~ ~~determine~~ their stability in soil, ~~instead~~ ~~and~~ environmental and biological controls (e.g., accessibility of the SOM to decomposers, abiotic reactions, and desorption) predominate ~~the SOM turnover~~, especially in the longer term.

In our model simulations, C and N in crop residues were allocated to two labile pools, but the allocation ratio did not greatly influence N₂O emissions, soil respiration, or mineral N. This could be related to the fact that the overall C/N ratio of crop residues was kept constant as the sizes of organic matter pools changed in the sensitivity analysis. The influence of crop residues on N₂O emissions may be better reflected in other residue properties, e.g., the C/N ratio and solubility of individual substrates (Aulakh et al., 1991; Surey et al., 2020). Furthermore, it should be noted that ~~the~~ addition of labile carbon from crop residues does not affect the decomposition of native soil organic matter in the model (i.e., no priming effect), as the decomposition of organic matter in labile and recalcitrant pools are calculated separately in CoupModel, ~~similar~~ as in other process-based models. The omission of a priming effect, the importance of which has been shown in field and laboratory studies (Kuziyakov, 2010), may cause models to underestimate the effects of crop residue composition on ~~the turnover of soil C and N~~ soil C and N turnover, particularly in connection with the initial burst of gas emissions.

~~On~~ ~~the other hand~~ ~~present~~ study, denitrifier growth parameters (e.g., d_{effNO} , $d_{growthNO3}$) showed considerable influence on the release of N₂O in most treatments (Fig. 3a). Our results suggested that the influences of microbial activities on N₂O emissions varied between different denitrification steps, and the denitrifier respiration for NO reduction showed a relatively larger and broader impact across treatments than other steps. The analysis of the two statistical measures σ and μ suggested that, rather than a single factor driving the model to become more 'behavioral' in predicting N₂O emissions, the collective effects of multiple parameters were more important, because one single parameter could exhibit various SEEs as other parameters changed, represented by high variability (σ) compared to the mean (μ) in Fig. 3d. ~~In calibrating~~ ~~For calibration of~~ complex models, several combinations of different parameter values might give the same goodness-of-fit between model outputs and measured variables, which is defined as equifinality (Beven and Freer, 2001). The model sensitivity to such parameters is probably attenuated in the case of high-level equifinalities. Besides, the importance of parameter interaction structure associated with equifinality could hinder the constraint of parameters and hence the reduction of uncertainty in N₂O simulations when limited measurement data are available. For instance, the fraction of C mineralized to CO₂, characterized by $(1-f_{e,li})$, and the decay rate of litter1 (k_{l1}), have a product interaction regarding the production of CO₂ (see Eq. (6.3) and Eq. (6.4) in Table S2). Also, the denitrifier growth rates (e.g., $d_{growthNO3}$) and the ~~Michaelis~~ ~~half-saturation~~ constant

Formatted: English (United Kingdom)

555 ~~characterized by of nitrogen substrates ($d_{hrateN_xO_y}$)~~ influence the loss of N from anaerobic N pools by invoking microbial growth via a quotient interaction (see Eq. (6.41) and Eq. (6.44) in Table S2).

The parameters ~~found to which~~ have the greatest impact on soil respiration and NH_4^+ content were associated with SOM composition (SOC_h) and decomposability (k_{l1} , k_{l2} , $f_{e,l1}$, $f_{e,l2}$), suggesting that model uncertainty for soil respiration and soil NH_4^+ could be greatly reduced if experimental data for either SOM composition or decay rates were available. For ~~simulating simulation of~~ soil NH_4^+ , information about microbial C/N ratio (cn_m) and denitrifier growth parameters (e.g., ~~$d_{growthNO_3}$~~ is ~~are~~ also important, because the availability of soil mineral N is closely associated with decomposition dynamics and its consumption by immobilization, nitrification, and denitrification (Lashermes et al., 2022). The ~~influencees influence~~ of soil ~~porosity and~~ wilting point on CO_2 emissions and soil NH_4^+ content ~~were was~~ larger under, ~~respectively, wet and~~ dry conditions. ~~The results can be explained by the fact that soil, while soil porosity had a greater effect on these two variables under wet conditions.~~ Soil porosity and wilting point are key set points of the soil moisture response function controlling the upper and lower bounds of the function, which implies that the measurement of soil hydraulic properties could reduce model uncertainty under contrasting soil moisture levels.

4.2 Model performance and possible explanations for deviations

570 Overall, the performance of posterior models varied between estimated variables and treatments. ~~The timing and magnitude of peak N_2O emissions were more difficult to predict than those of CO_2 emissions even though parameters had been adjusted for individual treatments, and negative errors relative to observations were seen particularly when simulating high N_2O emissions. Evaluation of model bias with respect to the slope β_1 in linear regression demonstrated a tendency across treatments to increasingly underestimate cumulative N_2O flux as the observed flux increased. The problem of underestimating high N_2O fluxes by process-oriented models has been reported in previous studies. For example, Fang et al. (2015) showed that four different algorithms all underestimated the four highest cumulative N_2O fluxes among eight N fertilizer treatments in an irrigated cornfield. Also, The timing and magnitude of peak N_2O emissions were more difficult to predict than those of CO_2 emissions. The GLUE calibration greatly reduced model uncertainty associated with N_2O gas emissions, with an average reduction of 83 % in the posterior ME range for all treatments (Fig. 4a), whereas the model still had difficulties capturing high emissions observed in the incubation (Fig. 5a). This was probably in part due to model biases in describing abiotic and biotic properties, but also due to uncertainties in the measurement dataset as discussed below (see Section 4.4). The model performance was generally consistent with previous attempts to simulate N_2O emissions from agricultural soils where models have been able to reproduce the cumulative emissions but had the same difficulties in describing the actual dynamics (e.g., Fuchs et al., 2020; He et al., 2016). Evaluation of model bias with respect to the slope β_1 in linear regression of model results vs. observations (Table 1) showed that the model had a better performance in its simulation of CO_2 fluxes and soil mineral contents than N_2O fluxes. The major mismatch between the model and measurements was associated with the underestimation of N_2O flux during the first three days of the experiment.~~

590 Nevertheless, our model showed initial success by better capturing the timing of N_2O for mix RC, particularly for “60 %
WFPS, +NO₃” ($R^2 > 0.4$, Figure 5a) than other treatments. According to Gaillard et al. (2018) evaluated the simulated N_2O
flux from three process-oriented models (DNDC, DayCent, and EPIC) and reported an underestimation of 0.01-0.93 kg N_2O -
N ha⁻¹ for every 1 kg of observed N_2O -N ha⁻¹ across models. who evaluated the simulated N_2O flux from three process-
oriented models (DNDC, DayCent, and EPIC), an underestimation of 0.01-0.93 kg N_2O -N ha⁻¹ for every 1 kg of observed
595 N_2O -N ha⁻¹ across these models were reported. When Fang et al. (2015) applied four different algorithms to an irrigated corn
field, they found that all algorithms underestimated the four highest cumulative N_2O fluxes among eight N fertilizer
treatments and explained this with an underestimation of soil water content. In the present study, parameter-induced biases
may have been an important source of simulation uncertainty as shown in Fig. 5, as some microbial coefficients were loosely
constrained, with more denitrification-related parameters than other process parameters (Fig. 3). So far very few studies have
reported parameterization and calibration of C and N cycling for incubation experiments. Our identified parameter
sensitivities and constrained parameters could provide a reference for future modeling of similar systems. More discussion of
parameter calibration is presented in Section 4.4.

600 Residual analysis revealed that the model had a tendency to simultaneously underestimate NO_3^- and overestimate NH_4^+ when
 N_2O emission was underestimated, and this trend was even more pronounced when looking at control treatments only (Figs.
S1 and S2). This suggests the nitrification rates may have been underestimated by the model and calls for revisiting the
parameterization of the nitrification process. The simulated accumulation of NH_4^+ in the RC residue treatments, in contrast to
the transient NH_4^+ peaks observed (Fig. 5c), indicates that the modeled NH_4^+ release linked to decomposition was greater
605 than the NH_4^+ consumption by microbial immobilization and nitrification simulated. Nylander et al. (2011) showed that a low
nitrification rate simulated by the CoupModel was possibly the reason for the overestimation of the amount of soil NH_4^+ in a
model of an organic cropping system. In our study, however, the weak and insignificant relationship between N_2O flux
residuals and the residuals for mineral N indicates that N_2O underestimation at high flux ranges may be due to other factors.

610 The unamended soil treatments (controls) were well described by the model, indicating that the repacked and pre-incubated
soil was a suitable representation of normal soil conditions. For amended soils, a rapid increase in soil NH_4^+ was observed by
day 1, while the modeled NH_4^+ tended to have a delayed increase for most residue treatments (Fig. 5c). According to
Lashermes et al. (2022), the red clover residues contained water-soluble N (WSN) corresponding to ca. 1 g N m⁻² which can
account for the observed increase. Cutting and mixing residues into the soil probably accelerated the release and
mineralization of N from the WSN pool compared to a real field situation (Angers and Recous, 1997), challenging the model
615 that was calibrated to describe decomposition in natural environments (see Section 4.4 for more discussion). It also stresses
the importance of realistic experimental setups for model parameterization under controlled laboratory conditions. The
accumulating trend of NH_4^+ simulated for the RC residue treatments was in contrast to the transient NH_4^+ peaks observed
(Fig. 5c), which indicates that the modeled NH_4^+ release from residue decomposition was greater than the NH_4^+ consumption
by microbial immobilization and nitrification that was simulated. By setting stricter selection criteria of ME and R^2 regarding

620 ~~NH₄⁺, such a trend could be avoided, but this was accompanied by decreased N₂O emission rates owing to reduced net N mineralization (data not shown). In our study, the relationships between N₂O flux residuals and the residuals for mineral N were weak and not significant, indicating that the N₂O underestimation at high flux ranges may also be due to other factors.~~

Inaccurate estimation of proximal factors such as soil water content and temperature by the pedo-climatic subroutines has been ~~put forward as~~ a main cause of errors in simulating C and N emissions in many process-based models (Brilli et al., 625 2017). ~~In our, but in the present~~ study, the soil water content and temperature were ~~assumed~~-constant during incubation, ~~but heterogeneity in the~~. ~~Instead, a heterogeneous~~ distribution of water ~~within residue-amended soil cores~~ could be a problem ~~when initializing for the description of soil environment~~ water content in the model. ~~Water because water~~ retention capacity in the soil might be altered by the practice of adding crop residues. ~~Lashermes et al. (2021)~~Lashermes et al. (2022) found that adding crop residues to soil increased the average WFPS of this layer from 60 % to 63 %. ~~Kravchenko et al. (2017)~~Also, 630 ~~Kravchenko et al. (2017)~~ found that specific gravimetric moisture of plant residues in soil could vary in the range ~~of 60-220 %~~-% and that residues were characterized by high moisture even at low soil water contents. Hence, the main effect of crop residues on the abiotic soil environment is probably not the marginal change in the average soil moisture content, but more likely the co-occurrence of elevated water content and labile C and N within the soil core. Residue fragments with high water retention capacity could represent microenvironments markedly different from those of the bulk soil and promote N₂O emissions (Kravchenko et al., 2017). ~~Model results indicated that the simulated O₂ content at 0-4 cm depth had only slight changes overall during incubation and was close to the saturation partial pressure in soil air owing to faster diffusion supply compared to soil respiration rates (Fig. 8). In the experiments, most of the oxygen consumption likely occurred in the microenvironment around residue debris. This is supported by observations of O₂ concentration (Kravchenko et al., 2017).~~

~~Model results indicated that the simulated O₂ content of soil air at 0-4 cm depth changed only slightly during incubation and was close to the saturation partial pressure in the atmosphere owing to faster diffusion supply compared to soil respiration rates (Fig. 8). However, most O₂ consumption likely occurred in the microenvironment around residue particles. This is supported by observations of O₂ concentration profiles in soil using O₂ microsensors (Markfoged et al., 2011) and planar optodes (Kravchenko et al., 2017) showing (Kravchenko et al., 2017), which showed the aerated O₂ partial pressure in the soil matrix away from organic hotspots and steep gradients in O₂ between bulk soil and hotspots of manure and residues, respectively. Nevertheless, Model simulations showed found that denitrification was the major N₂O-producing process that produced N₂O in the experiment, accounting for 76-100 % of the total estimated N₂O emissions. Parkin (1987) found that a thin water film even as little as 20 μm, could be enough to deplete air and support denitrification at the surface of decaying litter, and it is thus possible that the observed high N₂O fluxes were produced via denitrification despite an overall high aeration status within the soil. Water absorption by residue fragments from the surrounding soil could create local anoxic environments conducive to denitrification while also enabling the release of produced gases via drained pores (Kravchenko et al., 2018). In existing process-based models the heterogeneity in physical and biochemical processes caused by organic amendments is not included a high average oxygen status within the soil. However, existing process-based models do not~~

640
645
650

Field Code Changed

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

655 describe the heterogeneity in physical and biochemical processes caused by organic amendments, which may limit the ability of these models to reflect the microscale anaerobiosis and SOC availability, and to predict peak N₂O emissions such as those observed in RC treatments (Fig. 5a). Some studies have explored possibilities/methods to incorporate spatial variability into denitrification models, although conceptual frameworks considering heterogeneous environments for effects of heterogeneity on greenhouse gas emissions have only in recent years emerged and gained attention (Sihi et al., 2020). Using a stochastic modeling approach, Parkin (1987) found that the patchy dispersion pattern of high denitrification microsites was a major factor influencing the overall rates of denitrification. Based on a parsimonious numerical model, Sihi et al. (2020) used probability distribution functions to represent soil microsite production and consumption of three greenhouse gases, which explained occasional observations of simultaneous N₂O uptake (reduction) and CH₄ uptake (oxidation) that were not typically captured by other models. We suggest that model development should improve on the description/representation of microscale processes in soil, for example by parameterizing the distribution and extent of heterogeneity in, e.g., organic amendments and clay content, and by establishing the degree of anaerobiosis associated with organic hotspots and bulk soil separately.

670 The simultaneous underestimation of N₂O and NO₃⁻ could be linked in model runs (Fig. 5d and Fig. S1) may have been due to the an incomplete description of nitrate supply in the residue-amended 0-4 cm soil layer, which assumed there was no exchange with the lower 4-8 cm bulk soil layer. In a separate incubation experiment using the same soil type and experimental design and several of the same treatments, Lashermes et al. (2021) Lashermes et al. (2022) found that adding RC residue to the 0-4 cm soil layer induced a decrease in the NO₃⁻ content of the unamended 4-8 cm depth layer, indicating that NO₃⁻ removal in the above upper, amended layer influenced the NO₃⁻ dynamics in the bottom layer presumably during denitrification caused by a diffusive mass transfer between the two layers due to net consumption of NO₃⁻ within the top layer during denitrification. In the current model framework, solute transport is only simulated/possible by convection (driven by water flow) and does not include diffusion driven by concentration gradients. The model was originally designed for Under field conditions, and at this spatial scale infiltration is the main mechanism for solute transport between compartments. However, in the short term, but for a period after organic amendments, diffusive NO₃⁻ supply from the bulk soil can be the most important source of electron acceptor for denitrification, as observed in earlier incubation studies (Nielsen et al., 1996; Petersen et al., 1996). The current solute transport process may thus not be sufficient to properly simulate N₂O production in microbial hotspots on small scales, especially under low flow rates or for short travel distances where diffusive flux becomes increasingly important (Flury and Gimmi, 2002). Microbial turnover could accelerate the recycling of N and increase substrate availability for nitrification and denitrification locally (De Bruijn et al., 2009), but in this and some other process-based models, microbial N is not connected to the mineral N pool or included in the calculation of total N budget, which could be another reason for model discrepancies in mineral N dynamics. (Flury and Gimmi, 2018).

685 Only a few parameters showed distinct probability distribution patterns after calibration while others exhibited uniform distributions as the prior sampling. This result was in accordance with the second hypothesis, which should be related to the

Formatted: Strikethrough

690 ~~limited size of calibration data set in each treatment and the equifinalities between parameters.~~ The differences in the posterior parameter ~~distribution hold~~ distributions provided information about the ~~characteristics~~ simulation of ~~simulated~~ C and N processes between treatments, ~~although such variability may also reflect potential model limitations.~~ For example, the ~~well-constrained~~ microbial biomass C/N ratio (cn_m) ~~was constrained~~ within 10 in most treatments ~~was~~, consistent with observations that, on average, the C/N ratio of the soil microbial biomass varies between 6 and 10 ~~at a global scale~~ (Xu et al., 2013) and does not easily ~~adapt in composition to change with~~ litter quality (Spohn, 2015). Fungal cells typically have a C/N ratio ranging from 10 to 15 while bacteria range from 3.5 to 7 (Paul, 2007). In some treatments associated with extra NO_3^- input and high N_2O emissions, ~~however~~, the microbial C/N ratios in accepted runs exhibited relatively high values, closer to the soil-residue mixture C/N ratio. According to Eq. (6.7) and Eq. (6.8) in Table S2, a relatively high cn_m ~~could lead to a low level~~ coincides with simulations of low humification (i.e., less labile C and N converted to recalcitrant matter) as well as intense N mineralization (i.e., more organic N in litter pools converted to NH_4^+). ~~This could be associated with the underestimation of NO_3^- availability discussed above, especially for the treatments amended with crop residues.~~ Meanwhile, the relatively low values of estimated Michaelis constant $d_{\text{hrateN}_2\text{O}}$ suggested a high microbial affinity for soluble nitrogen oxides, ~~accelerating microbial denitrification.~~ ~~In treatments without NO_3^- addition, respiration via denitrification could be limited by the availability of electron acceptors through the respiratory chain, explaining an increase in the apparent Michaelis constant for N substrate reduction (Khalil et al., 2005).~~ ~~These trends all promote N availability for denitrification, and we propose with reference to Lashermes et al. (2022) that C/N ratios above 10 were related to an inaccurate assumption that there was no diffusive supply of NO_3^- from the 4-8 cm soil layer.~~ Including solute diffusion in the model may be able to change the posterior distributions of both parameters by better mimicking the mineral N supply.

700
705
710
715 Compared to control ~~treatments, the faster decay of organic matter (k_{d1} , k_{d2}) soils, C and higher CO_2 formation rate ($f_{e,11}$, $f_{e,12}$) N dynamics in the crop-residue treatments could reflect the need-amended soil, especially at the high moisture level (Fig. 5a), were difficult to mobilize N for use in nitrification and denitrification processes.~~ ~~Fast~~ simulate. In the experiment, rapid organic matter turnover in the residue-soil mixture was possibly caused by a high concentration of decomposer microorganisms associated with residue fragments. ~~Additionally, in contrast to natural soils, human disturbance of the soil in the laboratory could stimulate indigenous microbial communities resulting in rapid biological phenomena (Calderon et al., 2001; Thiessen et al., 2013).~~ ~~As discussed above, the rapid decay of organic matter (k_{d1} , k_{d2}) and high CO_2 formation rate ($f_{e,11}$, $f_{e,12}$) in the calibrated parameters in these amended treatments reflected the need to mobilize N for nitrification and denitrification processes to simulate observed N_2O emissions from amended soils.~~

It should be noted that the model deviations for N_2O flux were not caused by the spatial resolution of the vertical soil profile, which has been a problem in some studies (e.g. Xing et al., 2011), as the model performance concerning N_2O prediction was not improved in the multi-layer model (Table S6) where the one-layer soil profile had been sub-divided into five layers for simulations. Deviations between modeled results and measured values are more likely to have resulted from limitations in the description of the N processes behind N_2O emissions. For example, increasing the number of layers would not reflect the

Formatted: Strikethrough

720 ~~microscale processes associated with crop residue fragments and soil aggregates, nor would it address the missing description of solute diffusion between interfaces.~~

4.3 Treatment effects

725 ~~We did not investigate~~We also noted that the model deviations for N₂O flux were not caused by the spatial resolution of the vertical soil profile, which has been a problem in some studies (e.g., Xing et al., 2011), as model performance concerning N₂O prediction was not improved in the multi-layer model (Table S6) where the one-layer soil profile had been sub-divided into five layers for simulations. This was not surprising since structural uncertainty was not addressed in this way, for example, increasing the number of layers would not reflect the microscale processes associated with crop residue fragments and soil aggregates, nor would it address the missing description of solute diffusion between interfaces.

4.3 Treatment effects

730 ~~We only investigated~~ how the model responded to the specific change ~~of~~for two soil moisture and NO₃⁻ levellevels, but the results we obtained after calibrating the model against multiple treatments indicated ~~the challenges in predicting that it is difficult to predict~~ N₂O emissions under varying soil environmental conditions using a common model parameterization (Fig. 7). ~~Similar~~7). In the multiple treatment calibration, similar cumulative N₂O fluxes were simulated for treatments with the same NO₃⁻ level regardless of the soil moisture level, which was different from observations. ~~In~~For instance, in the experiment, ~~in with~~ RC treatments, higher N₂O fluxes were associated with the higher WFPS level (60 %) rather than with the higher NO₃⁻ level, ~~although there was but with~~ a strong interaction between the two factors (Taghizadeh-Toosi et al., 2021). ~~The problem to describe~~However, single treatment calibration showed more success to simulate the effects of incubation studies by process-based models was discussed in different soil moisture levels on N₂O fluxes. In a recent study by Grosz et al. (2021) ~~who also found that it difficult to describe treatments in incubation studies with~~ three N₂O models (DNDC, CoupModel, and DeNi) ~~that~~ responded to controlling factors in the same direction as measurements with frequencies from only 19 % to 67 %. ~~Different from their study, in which no systematic calibration of model parameters was performed, the~~In CoupModel, soil moisture effects have indirect effects on denitrification through decomposition, nitrification, and gas diffusion processes in which the soil moisture response function and its association with anaerobic conditions were accounted for. Hence, soil moisture may have had less or delayed effects on the N₂O emissions in model applications than in soil incubations with non-uniform distribution of moisture and C and N sources. The model deviations in our study, obtained by calibration of multiple treatments, suggested that potential limitations in model assumptions or the description of mechanisms were more ~~critical reasons~~important for unsatisfactory model responses than parameterization. ~~In CoupModel, while~~It should be noted, however, that only two moisture levels were included in the denitrification subroutine is sensitive to changes in soil temperature, pH, mineral N ~~experimental setup~~, and SOC content, the soil moisture has indirect and average effects on denitrification through decomposition, nitrification, and gas diffusion processes, but the effects of heterogeneity in the distribution of water and microbial activities are not represented. Therefore, ~~soil~~absence of a more

740
745
750

~~detailed moisture may have less effect on the N₂O flux estimation gradient, we were unable to draw more concrete conclusions about the model's performance in model applications than in real soil environments with heterogeneity in the distribution of C and N sources, and simulating soil moisture. This is still one of the most challenging tasks facing soil biogeochemical models. effects.~~

755 On the other hand, ~~our results showed a tendency to better predict with the same type of residue amendment,~~ treatments with higher ~~cumulative N₂O fluxes in the same group – either high soil moisture level or high N input – were better simulated by the model (Fig. 7a).~~ This can be understood from the characteristics of the calibration dataset and selection criteria. The high flux samples represented only a minor fraction of the total samples (i.e., 40 sampling points) in each group but were higher than the rest of them by orders of magnitude (Fig. 5a). The application of the ME criterion mainly constrained model
760 deviations for the high fluxes in ~~one each~~ data set, and less so for minor fluxes. It may be argued that this limitation could be improved by applying more stringent additional criteria such as R². However, this would reduce the acceptance rate or even refuse all posterior runs. Interestingly, Vezzano et al. (2012) obtained similar results in a GLUE context by using the Nash-Sutcliffe-based likelihood and stormwater measurements with large internal variability, and concluded that the choice of selection criteria should be based not only on its mathematical features but also by looking at the characteristics of the
765 available data.

~~We also found that our capacity to evaluate model performance was limited by the data available for model estimation and calibration. Some model parameters were not assessed in the incubation experiment (e.g. soil/residue labile C content and microbial biomass) and their values were either estimated or determined by calibration. The quality and temporal resolution in the measurement of controlling factors such as NO₃⁻ and NH₄⁺ were limited, and improving these aspects may reduce uncertainty in model prediction and facilitate model evaluation. By looking at the patterns of simulated N₂O emissions and ancillary variables, we identified potential problems behind model principles, which should be investigated with experimental studies designed carefully for model use. Previous studies, including global sensitivity analyses (Metzger et al., 2016; Wu et al., 2019) and model evaluations (Grosz et al., 2021), have specific suggestions to this end, such as improving measurement frequencies, evaluating sensitive input variables (e.g. decomposability of labile C), measuring more variables
770 regarding N cycle (e.g. N₂, NO) and using state-of-the-art techniques (e.g. ¹⁵N gas flux methods). We understand that collecting all data types discussed here is not always possible or practical, but encourage modelers to report more model outputs regarding N cycles even in the absence of observations, particularly the denitrification products, soil oxygen content, and anaerobic fraction, which was not done very often in previous studies.~~

4.4 Improving modeling practice and experimental design

780 ~~Some reasons for deviations between model simulations and experimental results were already discussed, but possibly other microbial and physiochemical processes are also not well described by current modeling practices. For example, enzymatic processes accounting for N₂O production and reduction could change progressively during incubation, which is not reflected~~

785 in fixed parameter values, as in this study. Khalil et al. (2005), coupling model and soil batch experiments, demonstrated the importance of an increase in the ability of denitrifiers to reduce N₂O for correct simulation of N₂O dynamics and variations in N₂O/(N₂O+N₂) ratios, especially when anaerobic conditions lasted more than one day. Microbial biomass and enzyme production can also vary under field conditions. Davidson et al. (2012) found that allowing model parameters to vary seasonally was more effective for modeling field data than calibrating parameters fixed at constant values to the entire annual dataset.

790 In view of these reports, we tried to test the sensitivity of simulated N₂O emissions to dynamic parameters using the default setup as a benchmark. Two parameters, d_{effNO} and $d_{\text{effN}_2\text{O}}$, were selected to simulate an increase with time in the ability of denitrifiers to produce and reduce N₂O, respectively, by assigning different values before and after day 3. The results (Fig. S7) show that the time course of N₂O emissions could be manipulated, with an intensification of the N₂O reduction activity over time changing the timing of an emission peak. It indicates that the use of fixed model parameters for short-term incubation experiments could lead to systematic errors if biotic properties such as the activity of enzymes or the affinity for a substrate of soil microbes vary significantly.

800 The rapid N mineralization in RC treatments during the early stage of incubation (Fig. 5c) may have been associated with disturbances in the experimental setup (see Section 4.2), which is beyond the description of the model framework. We conducted a simple sensitivity test with initial NH₄⁺ content on treatment 4 using default parameters and one calibrated parameter combination from the posterior ensemble. We found that by setting the initial NH₄⁺ content to different values between the observations on day 0 (value taken from the corresponding control) and day 1 to account for the rapid mineralization of WSN, simulated N₂O in the early stage could be enhanced (Fig. S8), and the impact was linked to parameter values. When using process-based models to simulate such systems, the measurements taken shortly after experiment initialization can be more suitable model inputs compared to the values from the starting point, and thus frequent measurements in the early stage of experiments are required.

805 We also found that our capacity to evaluate model performance was limited by the data available for model estimation and calibration. Firstly, some model parameters were not assessed in the incubation experiment (e.g., soil/residue labile C content and microbial biomass) and their values were either estimated or determined by calibration. Proximal chemical analysis related to these properties of soil samples could help reduce model uncertainties. Secondly, the quality and temporal resolution of the measurements of controlling factors such as NO₃⁻ and NH₄⁺ were limited. We used mean values of
810 recalculated measurements to compare with model simulations, where any measurement error was included in parameter ranges. Having explicit measurements in each layer (i.e., soil-residue mixture and bulk soil layer) of incubated soil cores, and higher measurement frequencies of mineral N and gas emissions within the first 24 hours of the experiment, are promising tools to improve the reliability of experimental data for modeling. Thirdly, in soil N₂O modeling, measuring more ancillary variables regarding N cycling (e.g., N₂, NO) and using state-of-the-art techniques (e.g., ¹⁵N gas flux methods) could

815 help test the validity of model principles (Grosz et al., 2021). Although collecting all data types discussed here is not always
possible or practical, we consider that reporting ancillary model outputs regarding N cycle processes even in the absence of
observations, particularly the denitrification products, soil oxygen content, and anaerobic fraction, will aid model evaluation
efforts, which was not done very often in previous studies. Accordingly, closer collaboration between modelers and
experimentalists is essential to increase mutual understanding, enforce better modeling practices, and find new knowledge by
820 merging data and modeling.

5 Conclusion

The current setup of CoupModel, when applied to results from an incubation study, indicated that parameters associated with the ~~decomposability~~decomposition of SOM and denitrifier growth were important in regulating soil respiration and mineral N dynamics. A high level of parameter interaction and equifinality issues existed ~~regarding~~in simulating N₂O emissions;
825 ~~hindering the determination of~~ has brought us challenges to quantify parameter sensitivities and improve parameter constraints.

The parameters showing posterior distributions that differed from the prior distributions revealed specific modeled microbial processes between treatments and may be used as references behind observations. For example, in the treatments ~~without~~with NO₃⁻ addition, the ~~availability~~relatively low values of N-substrates to denitrifiers ~~was limited according to~~
830 ~~$d_{hrateNO_3}$~~ in the posterior distribution of Michaelis constants, suggested a high microbial affinity for soluble nitrogen oxides,
accelerating microbial denitrification. More intense SOM decomposition was simulated in residue treatments compared to ~~controls-~~the “control” simulations.

The uncertainty analysis demonstrated ~~that~~ a model bias towards underestimating high-range daily and cumulative N₂O
fluxes, ~~which~~ was likely associated with ~~an inaccurate~~ simplified description of microbial processes in mineral N dynamics.
835 ~~Residual analysis indicated that nitrification rate could be underestimated but did not sufficiently explain the model~~
~~deviations.~~ While the simulated soil respiration response to soil moisture was generally in line with the direction of
measurement, the modeled N₂O emissions were not as sensitive to the WFPS as the measured data, ~~probably because~~but
further assessment of the ~~indirect effect of soil~~ this requires experimental data with more moisture response function on the
~~denitrification process.~~ ~~Discussing levels. We discussed~~ potential limitations in model principles related to structure
840 uncertainties based on current model outcomes (i.e., the prediction bias, we described several oxygen profiles, N₂O sources,
and moisture effects) and the emerging knowledge in recent experimental studies. Several suggestions for model
improvement ~~were described,~~ were described, including the use of new parameters and equations to represent microscale heterogeneity, and
a re-examination of the effects of soil moisture on denitrification processes: ~~with the assistance of more experimental data,~~
Under the current modeling framework, allowing for dynamic microbial parameters in calibration and careful consideration

845 of initial mineral N conditions may assist in better model representation of microbial and physiochemical processes in the context of incubation experiments as well as N application in field conditions.

Generally, we conclude that modeling N₂O emissions in controlled experiments is useful to identify the need for prior knowledge in both basic (e.g., decomposability of SOM) and elaborate (e.g., denitrifier growth) aspects of the process-based model for reducing the uncertainty of N₂O flux estimates. Moreover, we identified a potential model bias deviations
850 from observations and discussed future steps that may be required to assess its/their sources. We believe there is a need to modify model equations and revisit basic model assumptions with high quality measurement data sets and improve the experimental design that enable/enables more intensive/effective modeling practices, model evaluation/evaluation, and comparisons/comparison.

Code availability

855 The codes and descriptions of the models are available from the authors on request.

Data availability

The laboratory experiment and the modeled gas flux data are available from the authors on request.

Author contribution

860 JZ and SOP conceived the study. JZ designed the modeling experiments and carried them out, with contributions from WZ and SOP. PEJ made a version of CoupModel available model for the study and provided advice for model application. JZ prepared the manuscript and figures with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

865 This study was financially supported by Independent Research Fund Denmark (DRF) (Project acronym: modelN2O). Dr. Arezoo Taghizadeh-Toosi is acknowledged for having provided the experimental data. We further thank Iris Vogeler Cronin for providing insights on the model results. Zhang W. acknowledged the grant from the Swedish Research Council VR 2020-05338.

References

- 870 Abdalla, M., Jones, M., Yeluripati, J., Smith, P., Burke, J. and Williams, M.: Testing DayCent and DNDC model simulations of N₂O fluxes and assessing the impacts of climate change on the gas flux and biomass production from a humid pasture, *Atmos. Environ.*, 44, 2961–2970, doi:10.1016/j.atmosenv.2010.05.018, 2010.
- [Angers, D. A. and Recous, S.: Decomposition of wheat straw and rye residues as affected by particle size, *Plant Soil* 1997 1892, 189, 197–203, doi:10.1023/A:1004207219678, 1997.](#)
- 875 Aulakh, M. S., Walters, D. T., Doran, J. W., Francis, D. D. and Mosier, A. R.: Crop residue type and placement effects on denitrification and mineralization, *Soil Sci. Soc. Am. J.*, 55, 1020–1025, doi:10.2136/SSSAJ1991.03615995005500040022X, 1991.
- Beven, K. and Binley, A.: The future of distributed models: model calibration and uncertainty prediction, *Hydrol. Process.*, 6, 279–298, doi:10.1002/HYP.3360060305, 1992.
- 880 Beven, K. and Freer, J.: Equifinality, data assimilation, and uncertainty estimation in mechanistic modelling of complex environmental systems using the GLUE methodology, *J. Hydrol.*, 249, 11–29, doi:10.1016/S0022-1694(01)00421-8, 2001.
- Brilli, L., Bechini, L., Bindi, M., Carozzi, M., Cavalli, D., Conant, R., Dorich, C. D., Doro, L., Ehrhardt, F., Farina, R., Ferrise, R., Fitton, N., Francaviglia, R., Grace, P., Iocola, I., Klumpp, K., Léonard, J., Martin, R., Massad, R. S., Recous, S., Seddaiu, G., Sharp, J., Smith, P., Smith, W. N., Soussana, J.-F. and Bellocchi, G.: Review and analysis of strengths and 885 weaknesses of agro-ecosystem models for simulating C and N fluxes, *Sci. Total Environ.*, 598, 445–470, doi:10.1016/j.scitotenv.2017.03.208, 2017.
- [De Bruijn, A. M. G., Butterbach-Bahl, K., Blagodatsky, S. and Grote, R.: Model evaluation of different mechanisms driving freeze-thaw N₂O emissions, *Agric. Ecosyst. Environ.*, 133, 196–207, doi:10.1016/j.agee.2009.04.023, 2009.](#)
- Chen, D., Li, Y., Grace, P. and Mosier, A. R.: N₂O emissions from agricultural lands: a synthesis of simulation approaches, 890 *Plant Soil*, 309, 169–189, doi:10.1007/s11104-008-9634-0, 2008.
- Davidson, E. A. and Kanter, D.: Inventories and scenarios of nitrous oxide emissions, *Environ. Res. Lett.*, 9, 105012, doi:10.1088/1748-9326/9/10/105012, 2014.
- [Davidson, E. A., Samanta, S., Caramori, S. S. and Savage, K.: The Dual Arrhenius and Michaelis – Menten kinetics model for decomposition of soil organic matter at hourly to seasonal time scales, *Glob. Chang. Biol.*, 18, 371–384, doi:10.1111/j.1365-2486.2011.02546.x, 2012.](#)
- 895 Dungait, J. A. J., Hopkins, D. W., Gregory, A. S. and Whitmore, A. P.: Soil organic matter turnover is governed by accessibility not recalcitrance, *Glob. Chang. Biol.*, 18, 1781–1796, doi:10.1111/j.1365-2486.2012.02665.x, 2012.

- 900 Eusterhues, K., Rumpel, C., Kleber, M. and Kögel-Knabner, I.: Stabilisation of soil organic matter by interactions with minerals as revealed by mineral dissolution and oxidative degradation, *Org. Geochem.*, 34, 1591–1600, doi:10.1016/j.orggeochem.2003.08.007, 2003.
- Fang, Q. X., Ma, L., Halvorson, A. D., Malone, R. W., Ahuja, L. R., Del Grosso, S. J. and Hatfield, J. L.: Evaluating four nitrous oxide emission algorithms in response to N rate on an irrigated corn field, *Environ. Model. Softw.*, 72, 56–70, doi:10.1016/j.envsoft.2015.06.005, 2015.
- 905 Firestone, M. K. and Davidson, E. A.: Microbiological basis of NO and N₂O production and consumption in soil, *Exch. Trace Gases between Terr. Ecosyst. Atmos.*, 47, 7–21, 1989.
- Flury, M. and Gimmi, T. F.: 6.2 Solute Diffusion, in *Methods of Soil Analysis: Part 4 Physical Methods*, edited by J. H. Dane and C. G. Topp, pp. 1323–1351, Soil Science Society of America, Inc., Madison, U.S., ~~2002~~2018.
- 910 Fuchs, K., Merbold, L., Buchmann, N., Bretscher, D., Brilli, L., Fitton, N., Topp, C. F. E., Klumpp, K., Lieffering, M., Martin, R., Newton, P. C. D., Rees, R. M., Rolinski, S., Smith, P. and Snow, V.: Multimodel evaluation of nitrous oxide emissions from an intensively managed grassland. *J. Geophys. Res. Biogeosciences*, 125, 1–21, doi:10.1029/2019JG005261, 2020.
- Gabrielle, B., Laville, P., Duval, O., Nicoulaud, B., Germon, J. C. and Hénault, C.: Process-based modeling of nitrous oxide emissions from wheat-cropped soils at the subregional scale, *Global Biogeochem. Cycles*, 20, GB4018, doi:10.1029/2006GB002686, 2006.
- 915 Gaillard, R. K., Jones, C. D., Ingraham, P., Collier, S., Izaurrealde, R. C., Jokela, W., Osterholz, W., Salas, W., Vadas, P. and Ruark, M. D.: Underestimation of N₂O emissions in a comparison of the DayCent, DNDC, and EPIC models, *Ecol. Appl.*, 28, 694–708, doi:10.1002/eap.1674, 2018.
- 920 Gijsman, A. J., Hoogenboom, G., Parton, W. J. and Kerridge, P. C.: Modifying DSSAT crop models for low-input agricultural systems using a soil organic matter–residue module from CENTURY, *Agron. J.*, 94, 462–474, doi:10.2134/agronj2002.4620, 2002.
- Goreau, T. J., Kaplan, W. A., Wofsy, S. C., McElroy, M. B., Valois, F. W. and Watson, S. W.: Production of NO₂⁻ and N₂O by nitrifying bacteria at reduced concentrations of oxygen, *Appl. Environ. Microbiol.*, 40, 526–532, doi:10.1128/aem.40.3.526-532.1980, 1980.
- 925 Grandy, A. S. and Robertson, G. P.: Initial cultivation of a temperate-region soil immediately accelerates aggregate turnover and CO₂ and N₂O fluxes, *Glob. Chang. Biol.*, 12, 1507–1520, doi:10.1111/j.1365-2486.2006.01166.x, 2006.
- Grosz, B., Well, R., Dechow, R., Köster, J. R., Khalil, M. I., Merl, S., Rode, A., Ziehmer, B., Matson, A. and He, H.:

Evaluation of denitrification and decomposition from three biogeochemical models using laboratory measurements of N₂, N₂O and CO₂, *Biogeosciences*, 18, 5681–5697, doi:10.5194/bg-18-5681-2021, 2021.

930 [He, H., Jansson, P. E., Svensson, M., Meyer, A., Klemetsson, L. and Kasimir, Å.: Factors controlling Nitrous Oxide emission from a spruce forest ecosystem on drained organic soil, derived using the CoupModel, *Ecol. Modell.*, 321, 46–63, doi:10.1016/j.ecolmodel.2015.10.030, 2016.](#)

Jansson, P.-E.: CoupModel: model use, calibration, and validation, *Trans. ASABE*, 55, 1335–1344, doi:10.13031/2013.42245, 2012.

935 Jansson, P.-E. and Karlberg, L.: Coupled heat and mass transfer model for soil-plant-atmosphere systems, [online] Available from: <https://www.coupmodel.com/documentation> (Accessed 10 February 2022), 2010.

Jansson, P.-E. and Moon, D. S.: A coupled model of water, heat and mass transfer using object orientation to improve flexibility and functionality, *Environ. Model. Softw.*, 16, 37–46, doi:10.1016/S1364-8152(00)00062-1, 2001.

940 Keating, B. ., Carberry, P. ., Hammer, G. ., Probert, M. ., Robertson, M. ., Holzworth, D., Huth, N. ., Hargreaves, J. N. ., Meinke, H., Hochman, Z., McLean, G., Verburg, K., Snow, V., Dimes, J. ., Silburn, M., Wang, E., Brown, S., Bristow, K. ., Asseng, S., Chapman, S., McCown, R. ., Freebairn, D. . and Smith, C. .: An overview of APSIM, a model designed for farming systems simulation, *Eur. J. Agron.*, 18, 267–288, doi:10.1016/S1161-0301(02)00108-9, 2003.

Khalil, K., Renault, P., Guerin, N. and Mary, B.: Modelling denitrification including the dynamics of denitrifiers and their progressive ability to reduce nitrous oxide: comparison with batch experiments, *Eur. J. Soil Sci.*, 56, 491–504, doi:10.1111/j.1365-2389.2004.00681.x, 2005.

945 Kravchenko, A. N., Toosi, E. R., Guber, A. K., Ostrom, N. E., Yu, J., Azeem, K., Rivers, M. L. and Robertson, G. P.: Hotspots of soil N₂O emission enhanced through water absorption by plant residue, *Nat. Geosci.*, 10, 496–500, doi:10.1038/ngeo2963, 2017.

950 Kravchenko, A. N., Fry, J. E. and Guber, A. K.: Water absorption capacity of soil-incorporated plant leaves can affect N₂O emissions and soil inorganic N concentrations, *Soil Biol. Biochem.*, 121, 113–119, doi:10.1016/J.SOILBIO.2018.03.013, 2018.

Kuzyakov, Y.: Priming effects: interactions between living and dead organic matter, *Soil Biol. Biochem.*, 42, 1363–1371, doi:10.1016/j.soilbio.2010.04.003, 2010.

Kuzyakov, Y. and Blagodatskaya, E.: Microbial hotspots and hot moments in soil: concept & review, *Soil Biol. Biochem.*, 83, 184–199, doi:10.1016/j.soilbio.2015.01.025, 2015.

955 [Lam, P. S., Sokhansanj, S., Bi, X., Lim, C. J., Naimi, L. J., Hoque, M., Mani, S., Womac, A. R., Narayan, S. and Ye, X. P.:](#)

[Bulk density of wet and dry wheat straw and switchgrass particles, *Appl. Eng. Agric.*, 24, 351–358, doi:10.13031/2013.24490, 2008.](#)

Lashermes, G., Recous, S., Alavoine, G., Janz, B., Butterbach-Bahl, K., Ernfors, M. and Laville, P.: N₂O emissions from decomposing crop residues are strongly linked to their initial soluble fraction and early C mineralization, *Sci. Total Environ.*, 806, 150883, doi:10.1016/j.scitotenv.2021.150883, 2022.

Li, C., Frolking, S. and Frolking, T. A.: A model of nitrous oxide evolution from soil driven by rainfall events: 1. model structure and sensitivity, *J. Geophys. Res. Atmos.*, 97, 9759–9776, doi:10.1029/92JD00509, 1992.

Li, C., Aber, J., Stange, F., Butterbach-Bahl, K. and Papen, H.: A process-oriented model of N₂O and NO emissions from forest soils: 1. model development, *J. Geophys. Res. Atmos.*, 105, 4369–4384, doi:10.1029/1999JD900949, 2000.

Markfoged, R., Nielsen, L. P., Nyord, T., Ottosen, L. D. M. and Revsbech, N. P.: Transient N₂O accumulation and emission caused by O₂ depletion in soil after liquid manure injection, *Eur. J. Soil Sci.*, 62, 541–550, doi:10.1111/j.1365-2389.2010.01345.x, 2011.

[Metzger, C., Nilsson, M. B., Peichl, M. and Jansson, P. E.: Parameter interactions and sensitivity analysis for modelling carbon heat and water fluxes in a natural peatland, using CoupModel v5, *Geosci. Model Dev.*, 9, 4313–4338, doi:10.5194/gmd-9-4313-2016, 2016.](#)

Morris, M. D.: Factorial sampling plans for preliminary computational experiments, *Technometrics*, 33, 161–174, doi:10.1080/00401706.1991.10484804, 1991.

Nielsen, T. H., Nielsen, L. P. and Revsbech, N. P.: Nitrification and coupled nitrification-denitrification associated with a soil-manure interface, *Soil Sci. Soc. Am. J.*, 60, 1829–1840, doi:10.2136/sssaj1996.03615995006000060031x, 1996.

Norman, J., Jansson, P.-E., Farahbakhshazad, N., Butterbach-Bahl, K., Li, C. and Klemedtsson, L.: Simulation of NO and N₂O emissions from a spruce forest during a freeze/thaw event using an N-flux submodel from the PnET-N-DNDC model integrated to CoupModel, *Ecol. Modell.*, 216, 18–30, doi:10.1016/j.ecolmodel.2008.04.012, 2008.

[Nylinder, J., Stenberg, M., Jansson, P.-E., Klemedtsson, Å. K., Weslien, P. and Klemedtsson, L.: Modelling uncertainty for nitrate leaching and nitrous oxide emissions based on a Swedish field experiment with organic crop rotation, *Agric. Ecosyst. Environ.*, 141, 167–183, doi:10.1016/j.agee.2011.02.027, 2011.](#)

Parkin, T. B.: Soil microsites as a source of denitrification variability, *Soil Sci. Soc. Am. J.*, 51, 1194–1199, doi:10.2136/sssaj1987.03615995005100050019x, 1987.

Parton, W. J., Mosier, A. R., Ojima, D. S., Valentine, D. W., Schimel, D. S., Weier, K. and Kulmala, A. E.: Generalized model for N₂ and N₂O production from nitrification and denitrification, *Global Biogeochem. Cycles*, 10, 401–412,

- 985 doi:10.1029/96GB01455, 1996.
- Paul, E. A., Ed.: Soil microbiology, ecology and biochemistry, 3rd ed., Elsevier, Amsterdam, Netherlands., 2007.
- Petersen, S. O., Nielsen, T. H., Frostegård, Å. and Olesen, T.: O₂ uptake, C metabolism and denitrification associated with manure hot-spots, *Soil Biol. Biochem.*, 28, 341–349, doi:10.1016/0038-0717(95)00150-6, 1996.
- Ratto, M., Tarantola, S. and Saltelli, A.: Sensitivity analysis in model calibration: GSA-GLUE approach, *Comput. Phys. Commun.*, 136, 212–224, doi:10.1016/S0010-4655(01)00159-X, 2001.
- 990 Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S. and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, doi:10.1038/nature10386, 2011.
- Sierra, C. A., Harmon, M. E. and Perakis, S. S.: Decomposition of heterogeneous organic matter and its long-term stabilization in soils, *Ecol. Monogr.*, 81, 619–634, doi:10.1890/11-0811.1, 2011.
- 995 Sihi, D., Davidson, E. A., Savage, K. E. and Liang, D.: Simultaneous numerical representation of soil microsite production and consumption of carbon dioxide, methane, and nitrous oxide using probability distribution functions, *Glob. Chang. Biol.*, 26, 200–218, doi:10.1111/gcb.14855, 2020.
- Sin, G., Gernaey, K. V. and Lantz, A. E.: Good modeling practice for PAT applications: propagation of input uncertainty and sensitivity analysis, *Biotechnol. Prog.*, 25, 1043–1053, doi:10.1021/bp.166, 2009.
- 1000 Sommer, S. G., Petersen, S. O. and Møller, H. B.: Algorithms for calculating methane and nitrous oxide emissions from manure management, *Nutr. Cycl. Agroecosystems*, 69, 143–154, doi:10.1023/B:FRES.0000029678.25083.fa, 2004.
- Spohn, M.: Microbial respiration per unit microbial biomass depends on litter layer carbon-to-nitrogen ratio, *Biogeosciences*, 12, 817–823, doi:10.5194/bg-12-817-2015, 2015.
- 1005 Surey, R., Schimpf, C. M., Sauheitl, L., Mueller, C. W., Rummel, P. S., Dittert, K., Kaiser, K., Böttcher, J. and Mikutta, R.: Potential denitrification stimulated by water-soluble organic carbon from plant residues during initial decomposition, *Soil Biol. Biochem.*, 147, 107841, doi:10.1016/j.soilbio.2020.107841, 2020.
- Syakila, A. and Kroeze, C.: The global nitrous oxide budget revisited, *Greenh. Gas Meas. Manag.*, 1, 17–26, doi:10.3763/ghgmm.2010.0007, 2011.
- 1010 Taghizadeh-Toosi, A., Janz, B., Labouriau, R., Olesen, J. E., Butterbach-Bahl, K. and Petersen, S. O.: Nitrous oxide emissions from red clover and winter wheat residues depend on interacting effects of distribution, soil N availability and moisture level, *Plant Soil*, 466, 121–138, doi:10.1007/s11104-021-05030-8, 2021.

U.S. Environmental Protection Agency: Guidance on the development, evaluation, and application of environmental models (report no. EPA/100/K-09/003), [online] Available from: https://www.epa.gov/sites/default/files/2015-04/documents/cred_guidance_0309.pdf (Accessed 9 February 2022), 2009.

Uzoma, K. C., Smith, W., Grant, B., Desjardins, R. L., Gao, X., Hanis, K., Tenuta, M., Goglio, P. and Li, C.: Assessing the effects of agricultural management on nitrous oxide emissions using flux measurements and the DNDC model, *Agric. Ecosyst. Environ.*, 206, 71–83, doi:10.1016/j.agee.2015.03.014, 2015.

Vezzaro, L. and Mikkelsen, P. S.: Application of global sensitivity analysis and uncertainty quantification in dynamic modelling of micropollutants in stormwater runoff, *Environ. Model. Softw.*, 27–28, 40–51, doi:10.1016/j.envsoft.2011.09.012, 2012.

Vezzaro, L., Eriksson, E., Ledin, A. and Mikkelsen, P. S.: Quantification of uncertainty in modelled partitioning and removal of heavy metals (Cu, Zn) in a stormwater retention pond and a biofilter, *Water Res.*, 46, 6891–6903, doi:10.1016/j.watres.2011.08.047, 2012.

Wijler, J. and Delwiche, C. C.: Investigations on the denitrifying process in soil, *Plant Soil*, 5, 155–169, doi:10.1007/BF01343848, 1954.

World Meteorological Organization: WMO Greenhouse gas bulletin: the state of greenhouse gases in the atmosphere based on global observations through 2020, [online] Available from: https://library.wmo.int/index.php?lvl=notice_display&id=21975#.YgFFTrMJaq (Accessed 7 February 2022), 2021.

~~Wu, M., Ran, Y., Jansson, P. E., Chen, P., Tan, X. and Zhang, W.: Global parameters sensitivity analysis of modeling water, energy and carbon exchange of an arid agricultural ecosystem, *Agric. For. Meteorol.*, 271, 295–306, doi:10.1016/j.agrformet.2019.03.007, 2019.~~

Xing, H., Wang, E., Smith, C. J., Rolston, D. and Yu, Q.: Modelling nitrous oxide and carbon dioxide emission from soil in an incubation experiment, *Geoderma*, 167–168, 328–339, doi:10.1016/j.geoderma.2011.07.003, 2011.

Xu, X., Thornton, P. E. and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems, *Glob. Ecol. Biogeogr.*, 22, 737–749, doi:10.1111/geb.12029, 2013.

~~Zhang, Y., Ghalay, A. E. and Li, B.: Physical properties of wheat straw varieties cultivated under different climatic and soil conditions in three continents, *Am. J. Eng. Appl. Sci.*, 5, 98–106, doi:10.3844/ajeassp.2012.98.106, 2012.~~