



Lateral carbon export has low impact on the net ecosystem carbon balance of a polygonal tundra catchment

Lutz Beckebanze^{1,2}, Benjamin R. K. Runkle^{1,3}, Josefine Walz^{1,4}, Christian Wille⁵, David Holl^{1,2}, Manuel Helbig^{1,6}, Julia Boike^{7,8}, Torsten Sachs⁵, and Lars Kutzbach^{1,2}

¹Institute of Soil Science, Universität Hamburg, Germany

²Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Germany

³Department of Biological and Agricultural Engineering, University of Arkansas, Fayetteville, AR, USA

⁴Climate Impacts Research Centre, Institute for Ecology and Environmental Science, Umeå University, 98107 Abisko

⁵Helmholtz-Zentrum Potsdam – Deutsches Geo Forschungs Zentrum (GFZ), Potsdam, Germany

⁶Department of Physics & Atmospheric Science, Dalhousie University, Halifax, Canada

⁷Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

⁸Department of Geography, Humboldt-Universität zu Berlin, Germany

Correspondence: Lutz Beckebanze (lutz.beckebanze@uni-hamburg.de)

Abstract. Permafrost-affected soils contain large quantities of soil organic carbon (SOC). Changes in the SOC pool of a particular ecosystem can be related to its net ecosystem carbon balance (NECB) in which the balance of carbon (C) influxes and effluxes is expressed. For polygonal tundra landscapes, accounts of ecosystem carbon balances in the literature are often solely based on estimates of vertical carbon fluxes. To fill this gap, we present data regarding the lateral export rates of dissolved inorganic carbon (DIC) and dissolved organic carbon (DOC) from a polygonal tundra site in the North-Siberian Lena River Delta, Russia. We use water discharge observations in combination with concentration measurements of water-borne carbon to derive the lateral carbon fluxes from one season (2 June–8 September 2014 for DOC, 8 June–8 September 2014 for DIC). To put the lateral C fluxes into context, we furthermore present the surface–atmosphere eddy covariance fluxes of carbon dioxide (CO₂) and methane (CH₄) from this study site.

The results show cumulative annual lateral DIC and DOC fluxes of 0.31–0.38 g m⁻² and 0.06–0.08 g m⁻², respectively. Vertical turbulent fluxes of CO₂-C and CH₄-C accumulated to -19.0 ± 1.2 g m⁻² and 1.0 ± 0.02 g m⁻² during the 93-day observation period (8 June–8 September 2014). Thus, the lateral C export represented about 2% of the net ecosystem exchange of CO₂ (NEE). However, the relationship between lateral and surface–atmosphere fluxes changed over the observation period. At the beginning of the growing season (early June), the lateral C flux outpaced the surface-directed net vertical turbulent CO₂ flux, causing the polygonal tundra landscape to be a net carbon source during this time of the year. Later in the growing season, the vertical turbulent CO₂ flux dominated the NECB. Notably, the surface–atmosphere C flux served as a good approximation for the NECB at our study site during the growing season.

Keywords. net ecosystem carbon balance, dissolved inorganic carbon, dissolved organic carbon, polygonal tundra



20

1 Introduction

Permafrost regions have accumulated $1,300 \pm 200$ Pg of soil organic carbon (SOC), of which 472 ± 27 Pg are stored within the top 1 m of soil (Hugelius et al., 2014). In a warming climate, previously frozen SOC can be mobilised and lost from the permafrost-affected ecosystems through vertical and lateral carbon fluxes. Many studies (e.g., Koven et al., 2015; Schuur et al., 2015) focus on vertical gaseous carbon (C) fluxes in the form of the greenhouse gases carbon dioxide (CO_2) and methane (CH_4). However, C loss can also occur laterally as dissolved organic carbon (DOC) and dissolved inorganic carbon (DIC), which is exported through runoff of meltwater and rainwater (e.g., Fouché et al., 2017; Olefeldt and Roulet, 2014) and may be emitted in the form of GHGs to the atmosphere outside of the spatial observation range (such as in coastal regions as shown by Loughheed et al., 2020). Thus, to estimate the net ecosystem carbon balance (NECB), both vertical and lateral C fluxes must be considered (Chapin et al., 2006). Although scholars have identified lateral C transport as an important mechanism of C losses from terrestrial ecosystems in the Arctic (Zhang et al., 2017), little is known about the contribution of lateral C fluxes to the NECB since basic Arctic landscape C balance models focus solely on vertical C fluxes. So far, lateral C fluxes have only been included in NECB estimations in two subarctic boreal catchments in Northern Sweden, which show that lateral C fluxes contribute 6–15% (Chi et al., 2020) and 4–28% to the annual NECB (Öquist et al., 2014). To our knowledge, there has been no attempt yet to combine the lateral and vertical C fluxes in an Arctic ecosystem. Here, we estimate the NECB for a Siberian Arctic tundra ecosystem and present the individual flux contributions during one growing season.

Since 2002, the vertical C fluxes of CO_2 and CH_4 have been observed at our study site (Holl et al., 2019) using the eddy covariance method (Baldocchi, 2003). The study site, located in the North-Siberian Lena River Delta, is characterized by polygonal lowland tundra landscape. In this study, we combine the vertical C fluxes (F_{CO_2} and F_{CH_4}) with the lateral C fluxes (F_{DOC} and F_{DIC}) to derive the NECB for one growing season in 2014. We also compare the temporal dynamics of DIC and DOC concentrations with respect to the water discharge rate to find a potential driver for the concentrations of dissolved carbon. In summary, this study examines two research questions: (1) What is the influence of the lateral waterborne C fluxes on the NECB of the polygonal tundra landscape? (2) How do DOC and DIC concentrations and fluxes develop over the growing season?

2 Methods

2.1 Study site

The study site, Samoylov Island (Fig. 1), which is located in the southern part of the Russian Lena River Delta, consists of two geomorphological units: a modern floodplain in the west ($\sim 1.5 \text{ km}^2$) and a late-Holocene river terrace in the east ($\sim 3 \text{ km}^2$, Boike et al., 2013). The floodplain and river terrace are at elevations of 0–8 m and 8–13 m meters, respectively, above the water level of the Lena River (Boike et al., 2012a). Rain and meltwater from the river terrace drain towards the Lena River




Figure 1. Top: Map of Northern Russia (a) and the Lena River Delta (b). Bottom: Map of Samoylov Island (c) with the maximum (green) and minimum (yellow) catchment size (d). The blue cross in (d) marks the eddy covariance tower's location, while the yellow cross in (c) denotes the location of the meteorological station with the active layer depth observations. Panel (e) shows the locations of outflows A₁, A₂, and B. Map data from © OpenStreetMap contributors 2020, distributed under the Open Data Commons Open Database License (ODbL) v1.0 (a and b) and modified after Boike et al. (2012b; c, d, and e).

and the lower-lying floodplain. Polygonal tundra with a shallow active layer (less than 1 m) and wet to moist tundra vegetation (sedges, mosses, dwarf shrubs) characterizes the river terrace. Following the World Reference Base for Soil Resources (WRB, 2014), the main soil types include Histic Cryosols in polygon centers and Turbic Glacic Cryosols at elevated polygon rims on the river terrace (Pfeiffer and Grigoriev, 2002; Boike et al., 2013; Zubrzycki et al., 2014).



55 2.2 Catchment size

We estimated the catchment size by analyzing a digital elevation model (DEM) of Samoylov Island (Boike et al., 2012a) in ArcMap 10.1 (Environmental Systems Research Institute, ESRI). This DEM has a vertical and horizontal accuracy of less than 1 m. Due to the low elevation gradient in some catchment areas, we also used field observations from 2019 and shoreline measurements during different stages of the spring flood in 2014 to validate our estimations of the catchment size. To distinguish flow paths, we furthermore used the orthomosaic of Samoylov Island produced by Boike et al. (2012b) with an average horizontal resolution of 0.33 m. Based on this methodology, we estimated a larger catchment size than Helbig et al. (2013). This catchment drains through three outflows (A_1 , A_2 , and B). We estimated a catchment size of 0.69–0.84 km² (Fig. 1, panel (d); green denotes the maximum estimate, while yellow denotes minimum estimate).

The polygonal tundra in this catchment is characterized by intact and degraded ice-wedge polygons with water-filled centers as well as polygons with dry centers. Water-filled troughs between the polygons are also present (Kartozia, 2019). A low-lying and largely inundated area stretches from the center of the catchment towards the outflows A_1 and A_2 (Fig 1, panel (d)).

2.3 Water discharge

Between 2 June and 8 September 2014, we measured the water level at outflows A_1 , A_2 , and B using pressure sensors (Mini-Diver, Schlumberger Water Services, Netherlands) placed within 10 cm of the weir wall outside of the zone of fast flowing water. To calculate the water level above the diver, we subtracted the barometric pressure from the diver pressure. We measured the barometric pressure at the eddy covariance tower (CS100, Campbell Scientific, USA). Prior to the water discharge rate calculation, we corrected the water level estimations with a linear relationship with manual water level measurements (obtained using a ruler). The water discharge rates at outflows A_1 and A_2 were observed with V-notch weirs as described by Helbig et al. (2013). To calculate the water discharge rates, we used Kulin and Compton's 1975 method for calculating V-notch weirs:

75 $Q_a = \frac{8}{15} C_{weir} \sqrt{2g} \tan\left(\frac{1}{2}\alpha\right) h^{\frac{3}{2}}$, where Q_a is the discharge rate in Ls⁻¹, h is the water level above the notch determined from the pressure sensor in feet (ft), $C_{weir} = 0.58$ is the dimensionless weir constant, $g = 9.81 \text{ ms}^{-2}$ is the gravitational acceleration, and $\alpha = 60^\circ$ is the angle of the V-notch. After performing the water discharge rate calculations, we corrected the estimated water discharge rates linearly using manual bucket measurements (with a stop watch and a defined bucket volume).

Outflow B was located about 150 meters north of the A-outflows and the discharge rate was measured using a long-throated flume (RBC flume, 13.17.02, Eijkelpamp Agrisearch Equipment, the Netherlands). We calculated Q_b using the manufacturer's equation for this RBC flume: $Q_b = -0.066 + 0.016 h + 0.00063 h^2 + 7 \cdot 10^{-7} h^3$, where h is the water level above the notch in mm and Q_b is the water discharge rate in Ls⁻¹.

2.4 Dissolved inorganic carbon (DIC)

Between 8 June and 8 September 2014, we deployed a CO₂ sensor in the water column at outflow A_1 (CO₂ measurement system with multisensor module MSM-S2, UIT GmbH, Germany). This sensor measured the concentration of dissolved CO₂ (C_{dCO_2}) every 5 minutes (each measurement 15 seconds long).



DIC consists of dissolved CO₂ (as free CO₂ and carbonic acid; [H₂CO₃]), bicarbonate ions (HCO₃⁻), and carbonate ions (CO₃⁻²). In a freshwater system, each component's contribution to the DIC concentration depends on water temperature and pH; the bicarbonate equilibrium describes this relationship. More details on the bicarbonate equilibrium can be found in the book by Dodds and Whiles (2010). We calculated the carbonic acid concentration (C_{HCO₃⁻}) from C_{dCO₂}, water temperature, and pH: $C_{HCO_3^-} = \frac{K_1 \cdot C_{dCO_2}}{a(H^+)}$ where $a(H^+) = 10^{-pH}$. The dimensionless value of K_1 is temperature-dependent, and following Wong and Hsu (1991), is described as $K_1 = 10^{15.11 - 0.034 \cdot T - 3406.12 \cdot T^{-1}}$, where T describes the water temperature in the unit K .

The pH value was frequently, but not continuously measured throughout the 2014 growing season (N=40). To fill the gaps in the pH time series, we applied a running mean. The pH values varied between 6.60 and 6.99; therefore, the contribution of carbonate to C_{DIC} was not relevant. Due to the negligible amount of carbonate, we calculated C_{DIC} as the sum of C_{HCO₃⁻} and C_{dCO₂}.

The C_{dCO₂} sensor failed to record accurate measurements between 18 July and 30 July, and thus we excluded the recorded values during this period. To fill this data gap, we applied an artificial neural network (ANN), targeting C_{dCO₂} and using four input parameters (air temperature, relative air humidity, vertical CO₂ flux, and DOC concentration). We set up the ANN as a multilayer perceptron with 10 hidden neurons in Matlab's Deep Learning Toolbox (MATLAB, 2019b) using Levenberg-Marquardt backpropagation as an optimization algorithm. We divided the data sets into training (70%), validation (15%), and testing (15%) subsets. We assessed an agreement between the observed data and the independent testing subset as $R_{adj}^2 = 0.79$. Based on this agreement, we used the ANN for gap-filling of the C_{dCO₂} data set.

2.5 Dissolved organic carbon (DOC)

We routinely analyzed unfiltered water samples from all three outflows using a portable UV-Vis spectrometer probe (spectrolyzer, s::can Messtechnik GmbH, Austria). The measurements were supported by lab analyses to calibrate the spectrometer probe observations and increase data availability. Water samples for calibration were collected in acid-washed glass bottles, acidified to a pH value of 2, cooled to 4 °C for transport, and filtered (40 μm) prior to analysis. Analysis was conducted using a total organic carbon (TOC) analyzer (TOC-L, Shimadzu, Japan). We estimated C_{DOC} from unfiltered water samples following the workflow presented by Avagyan et al. (2014). This approach is based on the finding that different absorbance bands of a spectrometer probe can be suitable for the description of the DOC concentration, depending on the types of organic compounds in the sample water. We found a good agreement of $R_{adj}^2 = 0.82$ between C_{DOC} from the spectrometer probe and C_{DOC} from the TOC analyzer (Fig. A3). Details of the method can be found in supplement A1.

2.6 DOC and DIC flux

The catchment area-normalized lateral carbon fluxes of DOC (F_{DOC}) and DIC (F_{DIC}) are the product of water discharge rate Q and C_{DOC} and C_{DIC}, respectively, divided by the area of the catchment: $F_{DOC} = Q \cdot C_{DOC} / a$ and $F_{DIC} = Q \cdot C_{DIC} / a$, where a describes either the minimum or maximum estimated catchment size.



2.7 Environmental conditions

120 Precipitation and air temperature were recorded throughout the study period at the meteorological station in the southern
part of the island in one-hour intervals; Boike et al. (2019) published these measurements (data obtained from Boike et al.
(2019)). The growing degree days (GDD_{10}) were calculated as the sum of all positive differences between the daily mean air
temperature and the reference temperature (defined as 10 °C). The thaw depth was measured at a 150 grid point array next to
the meteorological station by pushing a metal rod vertically into the ground (Boike et al., 2019) and was obtained from the
125 GTNP database (GTNP Database, 2019).

2.8 Eddy covariance flux

We estimated the net vertical fluxes of CO₂ (F_{CO_2}), CH₄ (F_{CH_4}), and evapotranspiration (ET) using an eddy covariance
(EC) measurement system. Holl et al. (2019) described raw data processing of CO₂ fluxes; and the gap-filled time series were
obtained from Holl et al. (2018). High-frequency fluctuations in CH₄ concentration were observed with a Licor 7700 gas
130 analyzer (Licor Biosciences, USA). Data processing of CH₄ fluxes followed the same method as described in Holl et al. (2019)
for open-path CO₂ fluxes. Gap-filling of CH₄ fluxes was performed by applying a running mean of 48 hours. ET fluxes were
observed using a Licor 7500A gas analyzer (Licor Biosciences, USA), and the data processing followed Helbig et al. (2013).

2.9 Cumulative fluxes and uncertainty estimation

135 To quantify the impact carbon losses due to lateral transport have on the total carbon balance of Samoylov Island, we calculated
the cumulative carbon fluxes of CO₂, CH₄, DIC, and DOC for the period between 8 June and 8 September 2014 in 30-minute
intervals. Between 2 June and 7 June 2014, the component of C_{DIC} was not obtained yet, therefore, this period is not included
in the sums of carbon fluxes. However, this period is still part of the study period, since the spring flood had a great influence
on the DOC flux dynamics.

140 Flux uncertainties are related to combinations of random and systematic errors. Uncertainty from random errors can occur
for several reasons; for example, it can be caused by turbulence characteristics and the limited precision of measurement
instruments. Systematic errors occur due to systematic, yet unknown bias. Regarding the measurements of the vertical gas
fluxes and lateral carbon fluxes, careful calibrations of gas, DIC, and DOC analyzers as well as discharge estimates should
have minimized systematic errors. However, we could not calibrate our estimation of the catchment size. We treated the extent
145 of the maximum and minimum areas of the catchment as a systematic error. In this study, uncertainties from random errors are
indicated by the \pm symbol, and the ranges of uncertainties from systematic errors are indicated with a hyphen (–). Details of
the uncertainty estimation can be found in Appendix A2.



3 Results

3.1 Environmental conditions

150 To put the observation year of 2014 into perspective, we compared the meteorological conditions at our study site between
8 June and 8 September 2014, to the meteorological conditions during the same 93-day period over 20 years (1998 – 2018,
Fig. A1). With 87 °C at the end of the 93-day period, the growing degree days (GDD, Fig. A1 a) in 2014 were among the
average values, as established by the comparison dataset ($67 \frac{118}{49}$ °C, Median $\frac{75\text{th Percentile}}{25\text{th Percentile}}$). The 2014 thaw depths in the center
and the rim of the polygons studied (Fig. A1 b and c) in 2014 were among the deepest recorded in a 17-year companion dataset
155 (2002–2018).

The vertical water balance shows the precipitation, evapotranspiration, and water runoff rate between 2 June and 8 September
2014 (Fig. A2). The precipitation accumulation of 94 mm is within the average range for 14 of the available years between
1998–2018 ($95 \frac{138}{76}$ mm, Median $\frac{75\text{th Percentile}}{25\text{th Percentile}}$). In the same period, the evapotranspiration accumulated to 161 mm, and the lateral
water runoff accumulated to 23–38 mm. The 2014 spring flood of the Lena River flooded parts of the catchment. Therefore,
160 the lateral water runoff was largely influenced by the out-flowing water of this spring flood, visible at the beginning of the
observation period. The larger overall loss of water (183–189 mm) stands out more than the accumulation of water (94 mm)
during the observation period. However, one component has been neglected in this water balance: the snow accumulation in
winter, which was not observed between 2013 and 2014. However, in the winter of 2008 – 2009, there was a mean snow
accumulation of 65 ± 35 mm on Samoylov Island (snow water equivalent; Boike et al., 2013).

165 3.2 Lateral carbon flux dynamics

To focus more closely on lateral C flux dynamics, we examine the relationship between water discharge and DIC and DOC
concentration (Fig. 2). At outflow A₁, high DIC concentrations were generally associated with low water discharge. With
decreasing water discharge, the DIC concentration rose. A similar effect can be seen with the DOC concentration, which
continuously increases as the water discharge rate decreases data recorded during the river flood are excluded. A comparison
170 of the DIC and DOC concentrations shows that DIC concentrations were $4.31 \frac{6.41}{3.28}$ times higher than DOC concentrations
(Median $\frac{75\text{th Percentile}}{25\text{th Percentile}}$).

3.3 Net ecosystem carbon balance

In this section, we present the NECB for the study period consisting of the lateral (F_{DOC} and F_{DIC}) and vertical carbon
175 fluxes (F_{CO_2} and F_{CH_4}). The cumulative fluxes of all NECB components between 8 June and 8 September are summarized in
Fig. 3 and Tab. A1. Values with dominant systematic errors are expressed as ranges with an en-dash symbol, and values with
dominant random errors are expressed with a plus-minus symbol.

During the 93-day period in 2014, the NECB accumulated to $-17.6 - -17.5 (\pm 1.2)$ g m⁻². The vertical fluxes of F_{CO_2} and




Figure 2. Dissolved inorganic carbon (DIC, triangles) and dissolved organic carbon (DOC, circles) concentrations (note the different scales) against the water discharge rate on a semi-logarithmic scale. DOC concentration is shown on the right y-axis, while data after the spring flood (June 2–8) are shown in transparent circles. The DIC concentration was not observed during the spring flood.

F_{CH_4} contributed -19.0 ± 1.2 and 1.0 ± 0.02 $g\ m^{-2}$ to the NECB, respectively, while the lateral fluxes of F_{DIC} and F_{DOC} contributed $0.31 - 0.38$ and $0.06 - 0.08$ $g\ m^{-2}$ to the NECB, respectively, to the NECB. Thus, within the study period, lateral C fluxes exported 1.95 – 2.42% of the net ecosystem exchange (lateral-C-flux/NEE), i.e. the net C uptake due to the balance of photosynthesis and respiration.

We also split these cumulative fluxes into mean monthly fluxes (Fig. 4, a–e). During the periodic spring flood that occurred in 2014 partially in the first days of June (2–7 June Fig. 4 a), high lateral DOC flux ($13.0 - 15.8$ $mg\ m^{-2}\ d^{-1}$) and CH_4 -C flux (3.6 ± 0.3 $mg\ m^{-2}\ d^{-1}$) outpaced the CO_2 -C uptake (-7.0 ± 21.1 $mg\ m^{-2}\ d^{-1}$) and indicate an ecosystem carbon source (positive NECB). During this period, F_{DIC} was not yet observed. Therefore, the NECB is expected to be a stronger C source than presented in Fig. 4 a.

During the second part of June ($-213 - -211 \pm 11$ $mg\ m^{-2}\ d^{-1}$, Fig. 4 b) and in July ($-541 - -540 \pm 11$ $mg\ m^{-2}\ d^{-1}$, Fig. 4 c), the negative NECB indicates that the ecosystem served as a carbon sink due to high levels of plant CO_2 uptake. In August, the CO_2 sink strength decreased (Fig. 4 d) resulting in a mean daily CO_2 -C flux of only -24 ± 14 $mg\ m^{-2}\ d^{-1}$. At the same time, vertical CH_4 fluxes reached their maximum. Lateral DOC and DIC fluxes declined and the ecosystem acted as a weak carbon sink with an NECB of $-7.3 - -6.8 (\pm 14)$ $mg\ m^{-2}\ d^{-1}$. During the eight September days within the study period





Figure 3. Schematic illustration of all four cumulative flux components of the NECB (DIC, DOC, CH_4-C , and CO_2-C) in $g\ m^{-2}$ during the study period in 2014. The NECB is shown in the top right corner. Uncertainties from systematic errors are denoted with an en dash (–), while uncertainties from random errors are indicated with the plus-minus symbol (\pm). The picture in the background was taken on 30 August 2016 and provided by Jean-Louis Bonne.

(Fig. 4 e), all fluxes acted as carbon sources, and the NECB accumulated to $457 - 458 (\pm 25)\ mg\ m^{-2}\ d^{-1}$. With a relative contribution of 97%, the CO_2-C emission dominated the NECB in September.

195 4 Discussion

4.1 Comparison of DOC and DIC dynamics

We found a negative correlation between the water discharge rate and the DIC concentration (Fig. 2), meaning that higher water discharge rates dilute and decrease the DIC concentration. This result indicates that in other years, the DIC flux would not increase linearly with greater precipitation in other years. The results by Öquist et al. (2014) show precipitation to be an important driver of interannual variability in lateral C flux dynamics. Our study period had comparatively normal summer precipitation rates. In other years with higher precipitation rates, we would expect to find a higher water discharge rate. However,





based on our results, we do not expect a sharp rise in DIC flux to result from a higher water discharge rate. In one study, which focused on a catchment in Northern Sweden, a tripling in water discharge rate increased the annual lateral ^{14}C -export by only 2% (Campeau et al., 2017). In contrast, in another study in Northern Sweden, annual DIC export increased exponentially with rising water discharge rates (Öquist et al., 2014). Our findings suggest that the DOC flux has a low overall impact on the NECB in the studied polygonal tundra ecosystem; thus, an increase in the DOC flux in seasons with higher precipitation would likely affect the NECB only to a minor degree. However, DOC export from polygonal tundra may increase if arctic climate change will lead to accelerated degradation of ice-wedges, which is expected to enhance drainage of the permafrost landscape (Liljedahl et al., 2016).

210

One unexpected finding was the relationship between DOC and DIC concentrations with a C_{DIC}/C_{DOC} ratio of $4.31^{6.41}_{3.28}$ (Median $^{75\text{th Percentile}}_{25\text{th Percentile}}$). This ratio differs from those in other studies, which report a mean ratio of 0.65 from an Alaskan permafrost-affected watershed (Kling et al., 2000), a ratio of 0.24–1.30 in Canadian boreal biomes (Hutchins et al., 2019), and a ratio of 0.28 in a mixed coniferous forest in Northern Sweden (Chi et al., 2020). However, one study reported a C_{DIC}/C_{DOC} ratio of up to 11.6 in an ice-rich permafrost catchment in Northern Alaska (O'Donnell et al., 2019). A previous study at our study site found C_{DIC}/C_{DOC} ratios between 6.6 and 15.5 at the island's northern floodplain outlet in September 2008 (Abnizova et al., 2012), which are higher values than we present in this study, with a ratio of $4.12^{4.42}_{3.82}$ in September 2014 (Median $^{75\text{th Percentile}}_{25\text{th Percentile}}$, not shown in the results). The high C_{DIC}/C_{DOC} ratio hints to effective degradation and mineralization of dissolved organic matter (DOM) in the surface waters of the studied catchment. Such effective degradation of DOM by either photo-oxidation (Cory et al., 2014, 2015) and/or microbial decomposition (e.g., Drake et al., 2015; Mann et al., 2015; Spencer et al., 2015) has been found and intensively studied in other arctic catchments. The studied polygonal tundra catchment is characterised by (1) a low relief and (2) mostly shallow water bodies (depth < 1 m). Both factors enhance decomposition and mineralization of DOM: The low relief leads to long residence times of DOM before export, and the shallowness of the water bodies allows for intense light exposure and photodegradation of DOM, which, in turn, may promote microbial mineralization (Cory et al., 2015).

225

4.2 Net ecosystem carbon balance

We estimated the NECB using the lateral C fluxes (DIC and DOC flux) and the vertical fluxes of CO_2 and CH_4 . Our results indicate that vertical CO_2 uptake dominated the NECB during the study period. The lateral C fluxes exported only 1.95 – 2.42% of the NEE. During the complete study period, we found the lateral C fluxes' contribution to the NECB to be smaller than the uncertainty range of the observed CO_2 uptake. Nevertheless, our results also show that lateral carbon loss can exceed vertical carbon uptake at the beginning of the growing season. This finding shows that lateral C fluxes can play an essential role in the NECB during intensive water runoff periods, as we show in Fig. 4 a.

230

The question remains whether the resulting relationship between lateral C fluxes and the NECB would be similar in the previous and following growing seasons. A previous study at the site of the instant study includes a 15-year record of eddy covariance CO_2 fluxes between 19 July and August 23 of each year (Holl et al., 2019). It shows that CO_2 uptake in 2014 was

235



among the lowest values in the 15-year record. In 12 other years, CO₂ uptake was stronger compared to the 2014 period. According to these data, we assume that the influence of lateral C fluxes on the NECB would have played an even less important role in many other years compared to 2014. In two studies from this site, researchers reported low but varying average CH₄-C fluxes in two summer seasons: in the first study, the CH₄-C fluxes vary between 7.5 mg m⁻² d⁻¹ (28 June–22 July 2004) and 17.3 mg m⁻² d⁻¹ (18 July–25 July 2003; Wille et al., 2008). In the second study, Beckebanze et al. (2022) reported CH₄-C fluxes of 12.55^{16.07}_{9.65} mg m⁻² d⁻¹ (Median^{75th Percentile}_{25th Percentile}, 11 July–10 September 2019) Thus, our CH₄-C flux estimation of 10.6 mg m⁻² d⁻¹ in July 2014 lies within the range of estimates for other years.

We also investigate the question of whether the measured EC flux would be representative of the entire catchment. Instruments at the EC tower were mounted at a height of 4.15 m, and the tower was located approximately 850 m southeast of the A-outflows (see Fig. 1). The normalized mean contributions of four surface classes (based on the classification by Muster et al., 2012) within the footprint of the EC flux amounted to 66% (dry tundra), 18% (wet tundra), 8% (overgrown water), and 7% (water) in 2014 (Holl et al., 2019). Within the entire catchment (maximum estimated extent), these four surface classes amounted to 63% (dry tundra), 16% (wet tundra), 9% (overgrown water), and 11% (open water; Muster et al., 2012). Therefore, the distribution of tundra surface classes within the footprint of the EC flux is similar to the distribution of tundra surface classes within the catchment and the observed EC fluxes can be considered representative of vertical fluxes for the entire catchment.

This study discounts the contributions of particulate organic carbon, since we only found small differences between filtered (average 6.01 mg L⁻¹) and unfiltered water samples (average 6.07 mg L⁻¹) with respect to total carbon content. Thus, we suggest that particulate organic carbon would contribute only very little to the lateral C flux.

Previous studies that estimate the NECB from subarctic ecosystems have reported that the contribution of lateral C fluxes to the NECB can vary substantially from year to year. There are two studies from catchments in Northern Sweden. The first study shows that lateral C fluxes exported 6.4 and 15.4% of the NEE over two years in a forest and mire catchment (Chi et al., 2020). The second study, based on a 13-year period, shows that the lateral C fluxes export 4 – 28% of the annual NEE (Öquist et al., 2014).

5 Conclusions

At the polygonal tundra site in the Arctic Lena River Delta, which we investigated for this study, the net ecosystem carbon balance was periodically dominated by laterally exported dissolved carbon. The relative impact of these water-borne carbon losses on the total net ecosystem carbon balance was particularly high in the early and late growing season. During the Lena River spring flood, the in absolute and relative terms largest amounts of dissolved organic carbon were exported. In the late vegetation period, the relatively high impact of lateral C fluxes can largely be explained by low net ecosystem exchange rates of carbon dioxide due to generally diminished plant activity.

The contribution of lateral C fluxes to the cumulative NECB decreased on Samoylov Island over the growing season and was, in contrast to temperate and boreal ecosystems, negligible compared to cumulative vertical growing season carbon fluxes. We therefore conclude that the annual NECB of a polygonal tundra landscape is sufficiently described when only vertical



270 flux measurements are performed. Only studies which describe short-term tundra C balances should take lateral C export into account, particularly during or immediately following snow melt. Furthermore, in regions with rapid landscape degradation, lateral C fluxes could play a different, more relevant role in an ecosystem's carbon balance.

Data availability. The data will be published at Pangaea (doi will be added as soon as it becomes available).

275 *Author contributions.* Benjamin R. K. Runkle, Christian Wille, and Lars Kutzbach designed the experiments, and Benjamin R. K. Runkle, Christian Wille, David Holl, and Lars Kutzbach carried out the fieldwork. Benjamin R. K. Runkle, Josefine Walz, Lutz Beckebanze, and Lars Kutzbach developed the idea for the analysis, and Christian Wille provided processed eddy covariance data. The formal analysis and data visualization were done by Lutz Beckebanze. David Holl and Lars Kutzbach supervised it. Lars Kutzbach, Julia Boike, and Torsten Sachs provided resources for the instrumentation. Lutz Beckebanze prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

280 *Disclaimer.* This study was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2037 'CLICCS - Climate, Climatic Change, and Society' – Project Number: 390683824, contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg". Longterm measurements of CO₂ and CH₄ fluxes were supported by the projects "CarboPerm" (grant no. 03G0836A) and "KoPF" (grant no. 03F0764A), both funded by the German Federal Ministry of Education and Research (BMBF). Torsten Sachs and Christian Wille were supported by the Helmholtz Association of German Research Centres through a Helmholtz Young Investigators Group grant to Torsten Sachs (grant no. VH-NG-821).

285 *Acknowledgements.* We would like to thank the members of the Russian–German field campaign *LENA 2014*, especially Sandra Petersen (Universität Hamburg) for sampling analysis and the crew of the Russian research station Samoylov for logistical as well as technical support. We are grateful to Tim Eckhardt and Leonardo de Aro Galera for valuable discussions about the data analysis and the manuscript and Sarah Wiesner for being a great support as a Ph.D. advisory panel chair (all at the Universität Hamburg). We also thank Jean-Louis Bonne (Université de Reims Champagne-Ardenne) for allowing us to use the picture of the polygonal tundra.



290 Appendix A: Supplementary Material

A1 DOC concentration from a spectrometer probe

We used a multiple stepwise regression model (MSR; following Draper and Smith (2014)) to estimate C_{DOC} from a spectrometer probe in order to obtain a longer time series of C_{DOC} compared to the time series from the TOC analyzer. We compared the C_{DOC} analyzed in the TOC analyzer with the absorbance bands from a spectrometer probe to find suitable absorbance
295 bands to describe the DOC concentration. The spectrometer probe measured the absorbance (a_λ) of the sampled water probe between the 200 and 740 nm wavelength (λ) in 2.5 nm steps. In this analysis, we focused on the commonly used absorbance values between 250 and 740 nm as well as Ratio 1 (a_{465}/a_{665}) and Ratio 2 (a_{255}/a_{365}). Absorbance values below 250 nm were neglected due to possible interference with inorganic substances, following Avagyan et al. (2014). For the application of the MSR model, we split the data set of C_{DOC} from the TOC analyzer and the absorbance values from the spectrometer probe
300 into training (75%) and validation sets (25%). Details on the application of the MSR can be found in Avagyan et al. (2014). We applied the MSR model in Matlab R2019b using the *stepwisefit* function.

We used the following wavelengths and ratios as predictors for the DOC concentration at the three outflows: 250 nm and ratio 2 at outflow A₁; 250 nm, 300 nm, and 722.5 nm at outflow A₂; and 250 nm, 690 nm, and 712.5 nm at outflow B. In figure A3, the validation set of C_{DOC} from the TOC analyzer and the spectrometer probe are shown.

305 A2 Flux uncertainty estimation

We calibrated the observed water discharge rate Q against manual water bucket and manual height measurements. Therefore, we assumed random errors from both pressure sensors to dominate the uncertainty of Q . According to the manuals, our used Diver pressure sensor has a typical accuracy of 0.05% at full scale (however, the error is not further specified). The atmospheric pressure sensor has an accuracy of ± 1 hPa (one standard deviation). We used the Gaussian error propagation to
310 estimate the resulting uncertainty u_Q following two steps. First, we estimated the resulting error of the height measurement $u_{p_h} = \sqrt{u_{p_d}^2 + u_{p_a}^2}$, where u_{p_h} describes the uncertainty of the water level height measurement in hPa, and u_{p_d} and u_{p_a} describe the error of the Diver and the atmospheric pressure sensor, respectively. We converted u_{p_h} to the unit of mm, u_h , by dividing through the density of water and the earth's gravitational force. Second, we estimated the resulting uncertainty of Q as $u_Q = \frac{\delta Q}{\delta h} \cdot u_h$, where $\frac{\delta Q}{\delta h}$ describes the partial derivative from Q with respect to h .

315 The uncertainty of the DOC concentration results from the limits of the TOC analyzer (TOC-L, Shimadzu, Japan). The manufacturer states a maximum error of 1.5% in repetitive measurements. We used the RMSE between the modeled DOC concentration from the spectrometer and the DOC concentration from the TOC analyzer to estimate DOC concentration's uncertainty, $u_{C_{DOC}}$.

For the estimation of the uncertainty of C_{DIC} (u_{DIC}) we needed to consider the uncertainty of C_{dCO_2} , water temperature, and pH. According to the C_{dCO_2} sensor's manual, the sensor has an accuracy of 5% and, after calibration, an offset of up to 1 mg L⁻¹. The accuracy of the water temperature probe is given as $u_{t_w} = 0.2$ K. We estimated the pH uncertainty from the standard deviation of multiple measurements of the same water sample. The overall uncertainty of C_{DIC} was calculated using Gaussian
320



error propagation as: $u_{DIC} = \sqrt{u_{dCO_2}^2 + u_{HCO_3}^2}$ with $u_{HCO_3} = \sqrt{\left(\frac{\delta C_{HCO_3}}{\delta C_{dCO_2}}\right)^2 \cdot u_{dCO_2}^2 + \left(\frac{\delta C_{HCO_3}}{\delta pH}\right)^2 \cdot u_{pH}^2 + \left(\frac{\delta C_{HCO_3}}{\delta T_w}\right)^2 \cdot u_{T_w}^2}$.

We estimated the systematic and random uncertainty of the lateral C flux separately. Systematic uncertainty, described as
 325 $F_{DOC,sys}$ and $F_{DIC,sys}$, occurs due to systematic error of the catchment size and is estimated as $F_{DOC,sys} = F_{DOC_{a_{max}}} - F_{DOC_{a_{min}}}$,
 where $F_{DOC_{a_{min}}}$ and $F_{DOC_{a_{max}}}$ denote the DOC flux calculated with the largest and smallest assumed catchment size (re-
 sulting in the smallest and the largest DOC flux, respectively).

The random uncertainty of the lateral C flux, $F_{DOC,rand}$ and $F_{DIC,rand}$, resulting from random errors, is estimated as:

$$F_{DOC,rand} = F_{DOC} \sqrt{\left(\frac{u_Q}{Q}\right)^2 + \left(\frac{u_{C_{DOC}}}{C_{DOC}}\right)^2}.$$

330

We estimated the systematic uncertainty range of the cumulative lateral C flux ($\sum F_{DOC,sys}$ and $\sum F_{DIC,sys}$) as the differ-
 ence between the cumulative fluxes with the smallest and the largest assumed catchment size:

$\sum F_{DOC,sys} = \Delta t \sum_{t_1}^{t_n} F_{i_{a_{max}}} - \Delta t \sum_{t_1}^{t_n} F_{i_{a_{min}}}$, where Δt describes the duration of the measurement interval and t_1 and
 t_n denote the first and the last time step of the measurement, respectively.

335 We estimated the random uncertainty of the cumulative lateral C flux ($\sum F_{DOC,rand}$ and $\sum F_{DIC,rand}$) as:

$$\sum F_{DOC,rand} = \Delta t \sqrt{\sum_{t_1}^{t_n} \left(F_{DOC} \sqrt{\left(\frac{u_Q}{Q}\right)^2 + \left(\frac{u_{C_{DOC}}}{C_{DOC}}\right)^2} \right)^2}.$$

For the uncertainty estimation of DIC, we replaced DOC with DIC in the four equations above. In instances in this text
 where only a range of lateral C flux is provided, we ignored the random uncertainty and focused on the dominant systematic
 340 uncertainty.

The uncertainty of the vertical EC fluxes u_{FCO_2} and u_{FCH_4} were estimated in the flux processing software *EddyPro* following
 Finkelstein and Sims (2001). Details on the flux uncertainty estimation of F_{CO_2} can be found in Holl et al. (2019). We estimated
 the uncertainty of the cumulative vertical fluxes $u_{\sum F_{CO_2}}$ using the Gaussian error propagation for random uncertainties,
 resulting in $u_{\sum F_{CO_2}} = \Delta t \sqrt{\sum_{t_1}^{t_n} u_{FCO_2}^2}$ and $u_{\sum F_{CH_4}} = \Delta t \sqrt{\sum_{t_1}^{t_n} u_{FCH_4}^2}$ for F_{CO_2} and F_{CH_4} , respectively.



Table A1. Mean daily flux components (first five rows) and cumulative flux components (last row) of the NECB (CO₂-C, CH₄-C, DIC, and DOC) during the measurement period in 2014. Uncertainties from systematic errors are shown with an en dash (–) and uncertainties from random errors are denoted by the plus-minus symbol (±).

	NECB	vert CO ₂ -C Flux	vert CH ₄ -C Flux	DIC Flux	DOC Flux	Unit
2–7 June	9.6 – 12.3 (± 21)	-7.0 ± 21.1	3.6 ± 0.3	—	13.0 – 15.8	mg m ⁻² d ⁻¹
8–30 June	-213 – -211 (± 11)	-228 ± 11	8.2 ± 0.2	5.44 – 6.58	1.51 – 1.83	mg m ⁻² d ⁻¹
1–31 July	-541 – -540 (± 11)	-556 ± 11	10.6 ± 0.2	3.37 – 4.07	0.42 – 0.51	mg m ⁻² d ⁻¹
1–31 August	-7.3 – -6.8 (± 14)	-24 ± 14	14.4 ± 0.2	2.01 – 2.43	0.31 – 0.37	mg m ⁻² d ⁻¹
1–8 September	457 – 458 (± 25)	444 ± 25	9.9 ± 0.4	2.75 – 3.32	0.80 – 0.97	mg m ⁻² d ⁻¹
Season Sum (8 June– 8 September)	-17.6 – -17.5 (± 1.2)	-19.0 ± 1.2	1.0 ± 0.02	0.31 – 0.38	0.06 – 0.08	g m⁻²




Figure A1. Cumulative growing degree days (GDD; a) and thaw depths in the center and the rim of a polygon (b and c, respectively) between 8 June and 8 September of 2002–2018. The year 2014 is highlighted in orange. All other years are displayed in gray.




Figure A2. Vertical water balance at the study site between 2 June and 8 September 2014 with cumulative precipitation amounts above zero and cumulative evapotranspiration and water runoff rate below zero. Cumulative precipitation and evapotranspiration for the years 1998–2018 and 2007–2018, respectively, are shown in lighter colors (with the exception of 2014, which is shown in darker colors).




Figure A3. Validation set of C_{DOC} from the TOC analyzer and the spectrometer probe with the corresponding linear regression (dashed line). The solid line represents the 1:1 line.



345 References

- Abnizova, A., Siemens, J., Langer, M., and Boike, J.: Small ponds with major impact: The relevance of ponds and lakes in permafrost landscapes to carbon dioxide emissions, *Global Biogeochemical Cycles*, 26, <https://doi.org/10.1029/2011GB004237>, 2012.
- Avagyan, A., Runkle, B. R., and Kutzbach, L.: Application of high-resolution spectral absorbance measurements to determine dissolved organic carbon concentration in remote areas, *Journal of Hydrology*, 517, 435–446, <https://doi.org/10.1016/j.jhydrol.2014.05.060>, 2014.
- 350 Baldocchi, D. D.: Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future, *Global Change Biology*, 9, 479–492, <https://doi.org/10.1046/j.1365-2486.2003.00629.x>, 2003.
- Beckebanze, L., Rehder, Z., Holl, D., Wille, C., Mirbach, C., and Kutzbach, L.: Ignoring carbon emissions from thermokarst ponds results in overestimation of tundra net carbon uptake, *Biogeosciences*, 19, 1225–1244, <https://doi.org/10.5194/bg-19-1225-2022>, 2022.
- Boike, J., Dietrich, R., Günther, F., Langer, M., Piel, K., Scheller, S., Scheritz, M., and Schneider, W.: Digital Elevation Model of Samoylov Island, Lena River Delta, Siberia, Russia, with link to ArcGIS files, <https://doi.org/10.1594/PANGAEA.802255>, 2012a.
- 355 Boike, J., Grüber, M., Langer, M., Piel, K., and Scheritz, M.: Orthomosaic of Samoylov Island, Lena Delta, Siberia, <https://doi.org/10.1594/PANGAEA.786073>, 2012b.
- Boike, J., Kattenstroth, B., Abramova, E., Bornemann, N., Chetverova, A., Fedorova, I., Fröb, K., Grigoriev, M., Grüber, M., Kutzbach, L., et al.: Baseline characteristics of climate, permafrost and land cover from a new permafrost observatory in the Lena River Delta, Siberia (1998–2011), *Biogeosciences*, 10, 2105–2128, <https://doi.org/10.5194/bg-10-2105-2013>, 2013.
- 360 Boike, J., Nitzbon, J., Anders, K., Grigoriev, M., Bolshiyarov, D., Langer, M., Lange, S., Bornemann, N., Morgenstern, A., Schreiber, P., Wille, C., Chadburn, S., Gouttevin, I., Burke, E., and Kutzbach, L.: A 16-year record (2002–2017) of permafrost, active-layer, and meteorological conditions at the Samoylov Island Arctic permafrost research site, Lena River delta, northern Siberia: an opportunity to validate remote-sensing data and land surface, snow, and permafrost models, *Earth System Science Data*, 11, 261–299, <https://doi.org/10.5194/essd-11-261-2019>, 2019.
- 365 Boike, J., Nitzbon, J., Anders, K., Grigoriev, M. N., Bolshiyarov, D. Y., Langer, M., Lange, S., Bornemann, N., Morgenstern, A., Schreiber, P., Wille, C., Chadburn, S., Gouttevin, I., and Kutzbach, L.: Meteorologic data at station Samoylov (2002–2018, level 2, version 201908), link to archive, PANGAEA, <https://doi.org/10.1594/PANGAEA.905232>, in: Boike, J et al. (2019): Measurements in soil and air at Samoylov Station (2002–2018), version 201908. Alfred Wegener Institute - Research Unit Potsdam, PANGAEA, <https://doi.org/10.1594/PANGAEA.905236>, 2019.
- 370 Campeau, A., Bishop, K. H., Billett, M. F., Garnett, M. H., Laudon, H., Leach, J. A., Nilsson, M. B., Öquist, M. G., and Wallin, M. B.: Aquatic export of young dissolved and gaseous carbon from a pristine boreal fen: Implications for peat carbon stock stability, *Global Change Biology*, 23, 5523–5536, <https://doi.org/10.1111/gcb.13815>, 2017.
- Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., Clark, D. A., Harmon, M. E., Schimel, D. S., Valentini, R., et al.: Reconciling carbon-cycle concepts, terminology, and methods, *Ecosystems*, 9, 1041–1050, <https://doi.org/10.1007/s10021-005-0105-7>, 2006.
- 375 Chi, J., Nilsson, M. B., Laudon, H., Lindroth, A., Wallerman, J., Fransson, J. E. S., Kljun, N., Lundmark, T., Ottosson Löfvenius, M., and Peichl, M.: The Net Landscape Carbon Balance—Integrating terrestrial and aquatic carbon fluxes in a managed boreal forest landscape in Sweden, *Global Change Biology*, 26, 2353–2367, <https://doi.org/10.1111/gcb.14983>, 2020.



- 380 Cory, R., Harrold, K., Neilson, B., and Kling, G.: Controls on dissolved organic matter (DOM) degradation in a headwater stream: the influence of photochemical and hydrological conditions in determining light-limitation or substrate-limitation of photo-degradation, *Biogeosciences*, 12, 6669–6685, <https://doi.org/10.5194/bg-12-6669-2015>, 2015.
- Cory, R. M., Ward, C. P., Crump, B. C., and Kling, G. W.: Sunlight controls water column processing of carbon in arctic fresh waters, *Science*, 345, 925–928, <https://doi.org/10.1126/science.1253119>, 2014.
- 385 Dodds, W. K. and Whiles, M. R.: Chapter 13 - Carbon, in: *Freshwater Ecology (Second Edition)*, edited by Dodds, W. K. and Whiles, M. R., *Aquatic Ecology*, pp. 323–343, Academic Press, London, second edition edn., <https://doi.org/10.1016/B978-0-12-374724-2.00013-1>, 2010.
- Drake, T. W., Wickland, K. P., Spencer, R. G., McKnight, D. M., and Striegl, R. G.: Ancient low-molecular-weight organic acids in permafrost fuel rapid carbon dioxide production upon thaw, *Proceedings of the National Academy of Sciences*, 112, 13946–13951, <https://doi.org/10.1073/pnas.1511705112>, 2015.
- 390 Draper, N. R. and Smith, H.: *Applied regression analysis*, vol. 326, John Wiley & Sons, 2014.
- Environmental Systems Research Institute (ESRI), Redlands, C.: *ARC GIS Release 10.1*, 2012.
- Finkelstein, P. L. and Sims, P. F.: Sampling error in eddy correlation flux measurements, *Journal of Geophysical Research: Atmospheres*, 106, 3503–3509, <https://doi.org/10.1029/2000JD900731>, 2001.
- 395 Fouché, J., Lafrenière, M. J., Rutherford, K., and Lamoureux, S.: Seasonal hydrology and permafrost disturbance impacts on dissolved organic matter composition in High Arctic headwater catchments, *Arctic Science*, 3, 378–405, <https://doi.org/10.1139/as-2016-0031>, 2017.
- GTNP Database, 2019: <https://gtnp.arcticportal.org/>, access in February 2019.
- Helbig, M., Boike, J., Langer, M., Schreiber, P., Runkle, B. R. K., and Kutzbach, L.: Spatial and seasonal variability of polygonal tundra water balance: Lena River Delta, northern Siberia (Russia), *Hydrogeology Journal*, 21, 133–147, <https://doi.org/10.1007/s10040-012-0933-4>, 2013.
- Holl, D., Wille, C., Sachs, T., Schreiber, P., Runkle, B. R. K., Beckebanze, L., Langer, M., Boike, J., Pfeiffer, E.-M., Fedorova, I., Bolshiyarov, D. Y., Grigoriev, M. N., and Kutzbach, L.: A long-term (2002 to 2017) record of closed path and open path eddy covariance CO₂ net ecosystem exchange fluxes from the Siberian Arctic, <https://doi.org/10.1594/PANGAEA.892751>, 2018.
- 405 Holl, D., Wille, C., Sachs, T., Schreiber, P., Runkle, B. R. K., Beckebanze, L., Langer, M., Boike, J., Pfeiffer, E.-M., Fedorova, I., Bolshiyarov, D. Y., Grigoriev, M. N., and Kutzbach, L.: A long-term (2002 to 2017) record of closed-path and open-path eddy covariance CO₂ net ecosystem exchange fluxes from the Siberian Arctic, *Earth System Science Data*, 11, 221–240, <https://doi.org/10.5194/essd-11-221-2019>, 2019.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirmermeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- 410 Hutchins, R. H., Prairie, Y. T., and del Giorgio, P. A.: Large-Scale Landscape Drivers of CO₂, CH₄, DOC, and DIC in Boreal River Networks, *Global Biogeochemical Cycles*, 33, 125–142, <https://doi.org/10.1029/2018GB006106>, 2019.
- 415 Kartoziia, A.: Assessment of the Ice Wedge Polygon Current State by Means of UAV Imagery Analysis (Samoylov Island, the Lena Delta), *Remote Sensing*, 11, <https://doi.org/10.3390/rs11131627>, 2019.



- Kling, G. W., Kipphut, G. W., Miller, M. M., and W, J. O.: Integration of lakes and streams in a landscape perspective: the importance of material processing on spatial patterns and temporal coherence, *Freshwater Biology*, 43, 477–497, <https://doi.org/10.1046/j.1365-2427.2000.00515.x>, 2000.
- 420 Koven, C. D., Schuur, E., Schädel, C., Bohn, T., Burke, E., Chen, G., Chen, X., Ciais, P., Grosse, G., Harden, J. W., et al.: A simplified, data-constrained approach to estimate the permafrost carbon–climate feedback, *Philos. Trans. Royal Soc. A*, 373, 20140423, <https://doi.org/10.1098/rsta.2014.0423>, 2015.
- Kulin, G. and Compton, P. R.: A guide to methods and standards for the measurement of water flow, 421, US Dept. of Commerce, National Bureau of Standards, 1975.
- 425 Liljedahl, A. K., Boike, J., Daanen, R. P., Fedorov, A. N., Frost, G. V., Grosse, G., Hinzman, L. D., Iijma, Y., Jorgenson, J. C., Matveyeva, N., et al.: Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology, *Nature Geoscience*, 9, 312–318, 2016.
- Lougheed, V. L., Tweedie, C. E., Andresen, C. G., Armendariz, A. M., Escarzaga, S. M., and Tarin, G.: Patterns and Drivers of Carbon Dioxide Concentrations in Aquatic Ecosystems of the Arctic Coastal Tundra, *Global Biogeochemical Cycles*, 34, 1–13, <https://doi.org/10.1029/2020GB006552>, 2020.
- 430 Mann, P. J., Eglinton, T. I., McIntyre, C. P., Zimov, N., Davydova, A., Vonk, J. E., Holmes, R. M., and Spencer, R. G.: Utilization of ancient permafrost carbon in headwaters of Arctic fluvial networks, *Nature Communications*, 6, 1–7, <https://doi.org/10.1038/ncomms8856>, 2015.
- MATLAB, 2019b: MATLAB Deep Learning Toolbox, the MathWorks, Natick, MA, USA, 2019b.
- Muster, S., Langer, M., Heim, B., Westermann, S., and Boike, J.: Subpixel heterogeneity of ice-wedge polygonal tundra: a multi-scale analysis of land cover and evapotranspiration in the Lena River Delta, Siberia, *Tellus B: Chemical and Physical Meteorology*, 64, 17301, <https://doi.org/10.3402/tellusb.v64i0.17301>, 2012.
- 435 O'Donnell, J. A., Carey, M. P., Koch, J. C., Xu, X., Poulin, B. A., Walker, J., and Zimmerman, C. E.: Permafrost Hydrology Drives the Assimilation of Old Carbon by Stream Food Webs in the Arctic, *Ecosystems*, 23, 435–453, <https://doi.org/10.1007/s10021-019-00413-6>, 2019.
- 440 Olefeldt, D. and Roulet, N. T.: Permafrost conditions in peatlands regulate magnitude, timing, and chemical composition of catchment dissolved organic carbon export, *Global Change Biology*, 20, 3122–3136, <https://doi.org/10.1111/gcb.12607>, 2014.
- Öquist, M. G., Bishop, K., Grelle, A., Klemedtsson, L., Köhler, S. J., Laudon, H., Lindroth, A., Ottosson Löfvenius, M., Wallin, M. B., and Nilsson, M. B.: The Full Annual Carbon Balance of Boreal Forests Is Highly Sensitive to Precipitation, *Environmental Science & Technology Letters*, 1, 315–319, <https://doi.org/10.1021/ez500169j>, 2014.
- 445 Pfeiffer, E.-M. and Grigoriev, M. N.: Russian-German Cooperation SYSTEM LAPTEV-SEA 2000: The Expedition LENA 2001, *Berichte zur Polar-und Meeresforschung (Reports on Polar and Marine Research)*, 426, 2002.
- Schuur, E. A., McGuire, A. D., Schädel, C., Grosse, G., Harden, J., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., and Lawrence, D. M.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- Spencer, R. G., Mann, P. J., Dittmar, T., Eglinton, T. I., McIntyre, C., Holmes, R. M., Zimov, N., and Stubbins, A.: Detecting the signature of permafrost thaw in Arctic rivers, *Geophysical Research Letters*, 42, 2830–2835, <https://doi.org/10.1002/2015GL063498>, 2015.
- 450 Wille, C., Kutzbach, L., Sachs, T., Wagner, D., and Pfeiffer, E.-M.: Methane emission from Siberian arctic polygonal tundra: eddy covariance measurements and modeling, *Global Change Biology*, 14, 1395–1408, <https://doi.org/10.1111/j.1365-2486.2008.01586.x>, 2008.
- Wong, G. T. and Hsu, C.-H.: The first apparent dissociation constant of carbonic acid in marine waters from 0 to 20‰ and 0 to 30 °C, *Estuarine, Coastal and Shelf Science*, 32, 257 – 265, [https://doi.org/10.1016/0272-7714\(91\)90019-8](https://doi.org/10.1016/0272-7714(91)90019-8), 1991.



- 455 WRB, I. W. G.: International soil classification system for naming soils and creating legends for soil maps, *World Soil Resources Reports*, 106, 2014, 2014.
- Zhang, X., Hutchings, J. A., Bianchi, T. S., Liu, Y., Arellano, A. R., and Schuur, E. A.: Importance of lateral flux and its percolation depth on organic carbon export in Arctic tundra soil: Implications from a soil leaching experiment, *Journal of Geophysical Research: Biogeosciences*, 122, 796–810, <https://doi.org/10.1002/2016JG003754>, 2017.
- 460 Zubrzycki, S., Kutzbach, L., and Pfeiffer, E.-M.: Permafrost-affected soils and their carbon pools with a focus on the Russian Arctic, *Solid Earth*, 5, 595–609, <https://doi.org/10.5194/se-5-595-2014>, 2014.