

Technical note: Common ambiguities in plant hydraulics

Yujie Wang¹ and Christian Frankenberg^{1,2}

¹Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California 91125, USA

²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California 91109, USA

Correspondence: Yujie Wang (wyujie@caltech.edu)

Abstract. Plant hydraulics gains increasing interest in plant eco-physiology and vegetation modeling. However, the hydraulic properties and profiles are often improperly represented thus leading to biased results and simulations, e.g., the neglect of gravitational pressure drop results in overestimated water flux. We highlight the commonly seen ambiguities and/or misunderstandings in plant hydraulics, including (1) distinction between water potential and pressure, (2) differences among hydraulic conductance and conductivity, (3) xylem vulnerability curve formulations, (4) [model complexity](#), (5) stomatal model representations, (56) bias from analytic estimations, (67) whole plant vulnerability, and (78) neglected temperature dependencies. We recommend careful thinking before using or modifying existing definitions, methods, and models.

1 Introduction

Plant hydraulics gains increasing interest in understanding plants' responses and acclimation/adaptation to the environment (Santiago et al., 2004; McDowell et al., 2008; McDowell, 2011; Meinzer et al., 2010; Allen et al., 2010; Anderegg et al., 2012, 2016; Gleason et al., 2016; Wang et al., 2021a; Liu et al., 2021) and modeling canopy carbon and water fluxes within vegetation and land surface models (Buckley and Mott, 2013; Manzoni et al., 2013; McDowell et al., 2013; Sperry et al., 2017; Kennedy et al., 2019; Liu et al., 2020; Wang et al., 2020, 2021b; Sabot et al., 2022). However, xylem hydraulic properties and flow pressure profile are often improperly represented, due to the ambiguities and misunderstandings of various plant hydraulic parameters, though the plant hydraulic models used in topical research are already dramatically simplified compared to a complicated hydraulic architecture (Tyree and Ewers, 1991). For instance, distinctions between (a) water potential and pressure, (b) hydraulic conductance and conductivity, and (c) division and derivative are often not recognized. Further, the pursuit of simplicity, analytical solution, and novelty consequently results in modifications of known and well tested functional forms. However, while researchers should be encouraged to try "new" approaches, it is important to keep in mind whether these changes or new methods (a) are correct and (b) need to be tested before moving forward. Any research violating the two principles would be unwarranted, not matter how "reasonable" they appear to be.

For example, regarding the modeling of plant hydraulics, since Wolf et al. (2016) and Sperry et al. (2017) advanced the stomatal optimization theory (Cowan and Farquhar, 1977) by quantifying hydraulic risk under a general gain-risk optimization framework, an increasing number of new models or variants have been developed (e.g., Anderegg et al., 2018; Dewar et al., 2018; Eller et al., 2018; Wang et al., 2020, 2021a; Chen et al., 2022); and many plant hydraulics-based models show predictive skills comparable to statistical methods (Anderegg et al., 2018; Venturas et al., 2018; Eller et al., 2020; Wang et al., 2020;

Sabot et al., 2022). Nevertheless, these tested models are not always replicated correctly as researchers tend to mutate the formulations and sometimes hypotheses, such as the neglect of the rhizosphere component that plays an important role in drought stress conditions (Sperry et al., 1998; Sperry and Love, 2015; Sperry et al., 2016; Wang et al., 2020). Although the modifications often resemble tested models, they are often used without being thoroughly tested. Reasons behind the lack of model testing include (a) there is not yet a well established method or database to conveniently benchmark the new model variants, (b) research that focus on varying the formulations and testing the variants are not encouraged by reviewers due to lack of novelty, and (c) one may not recognize the changes or differences that have been made. Here, we list some common ambiguities and misunderstandings in plant hydraulics, and recommend careful thinking before using or modifying existing definitions, methods, and models.

2 Water potential and pressure

Water ~~potential-and-pressure~~ movement in plants includes mass flow through xylem conduits and diffusion between xylem conduits and capacitance tissues. Water mass flow (from site 1 to 2) in xylem is driven by the net force at the target plane per area (driving pressure, DP), which is $DP = P_{x1} - P_{x2} + \rho gh_1 - \rho gh_2$ as in Figure 1a; and the dissolved ions only play a role in liquid water density. Water diffusion across the cell membrane (from xylem conduit to the cell) is driven by the potential difference, which is $P_x - P_c + \Psi_{mx} - \Psi_{mc}$ as in Figure 1a; and the dissolved ions play a role through the osmotic potential.




Figure 1. Diagram for the difference between potential and pressure in plant hydraulics. (a) Difference among pressure difference (ΔP), potential difference ($\Delta\Psi$), and mass flow driving pressure (DP). Water potential (Ψ) is the sum of pressure (P , P_x for xylem pressure and P_c for cell turgor pressure), osmotic potential (Ψ_m , Ψ_{mx} for xylem and Ψ_{mc} for cell), and gravitational potential related to height (ρgh). (b) Example of why $\Delta\Psi$ cannot be used for mass flow. In scenario 1 (left) where $\Psi_{mx2} = \Psi_{mx1}$ (labeled in blue), the $\Delta\Psi = DP$. In scenario 2 (right) where $\Psi_{mx2} < \Psi_{mx1}$ (labeled in red), $\Delta\Psi > DP$.

As ion concentration in xylem conduit is very low, the osmotic potential in xylem conduit (Ψ_{mx}) is often ignored. Further, as there is not a general name for the term $P_x + \rho gh$, water potential is imperceptibly used in place of mass flow driving pressure

(i.e., DP) because of the gravity term in it. To date, many people use water potential rather than water pressure when modeling and describing mass water flow. This, though may be easier for people to understand, is not correct. For example, in Figure 1b, if the osmotic potential at the bottom and top are the same, the driving pressure and water potential difference are the same. However, if the osmotic potential at the top is more negative than at the bottom (for example, via adding a very thin layer of high concentration salt solution), the driving pressure will be lower than the water potential difference. Thus, using potential difference for water mass flow is technically incorrect, and it is necessary to clarify the terminology to distinguish them.

Water potential and pressure used in plant hydraulics are both defined as a difference from a reference value: water potential (often denoted as Ψ or ψ in literature) is typically defined as the difference from the potential of pure water in the soil, and water pressure (often denoted as P or p) is typically defined as the difference from the environmental air pressure. Water potential gradients drive water flow through permeable media such as xylem conduits, while water pressure difference between xylem water and surrounding air is responsible for air-seeded conduit cavitation (Sperry and Tyree, 1988; Tyree and Sperry, 1989), which occurs when the pressure difference exceeds the capillary pressure at the air-water interface. Thus, using water potential to describe xylem vulnerability curve should be avoided. Although the Δ values of the two are often interchangeable in many scenarios (e.g., when there is no height change or external air pressure), one needs to be cautious to avoid ambiguity. Note that flow rate calculation from hydraulic conductance, or vice-versa, should use water potential (equation 1). For instance, for a 100 m tall tree with no transpiration, leaf water potential is equal to soil water potential; however, leaf xylem water pressure would be approximately 1 MPa more negative than xylem pressure at the tree base; and using pressure drop here to derive flow rate will be incorrect when there is height change. On the other hand, calculation of hydraulic dysfunction caused by xylem cavitation should use water pressure (ΔP between xylem water and surrounding air). In this case, using leaf water potential to estimate xylem conductance loss is incorrect, as there is not a unique conductance for a given water potential.

- Use pressure in xylem cavitation;
- Use pressure in water mass flow;
- Use potential in water diffusion across the cell membrane (e.g., water exchange between xylem and living cell).

The ambiguity in potential and pressure results in inaccurate terminology in the literature, for example. A commonly seen sophistic “mistake” is the use of leaf water potential and vulnerability curve. Leaf water potential is often estimated using the pressure chamber method (Scholander et al., 1964; Boyer, 1967). While it is well known that the pressure chamber method gives the applied pressure at the free meniscus of the cut end. However, the term “potential” is not accurate here, as the pressure chamber method gives the applied pressure at the free meniscus of the cut end. For example, for a tree with no transpiration, gives a decent estimate of the xylem water pressure, people often refer to the measurement as leaf water potential equals soil water potential; however, the measured “potentials” values differ for leaves at different heights because of the gravity. The fact is that the measured “potential” value from pressure chambers does not have a gravity term in it, and it is as (a) xylem conduit water has very low solute content, and (b) gravity term is often negligible compared to the very negative leaf xylem water pressure. However, it is always more accurate to treat it as an equivalent pressure, say leaf water pressure or a balance pressure (at the end of water flow xylem). For example, when the whole plant is under equilibrium, leaf

water potential should be equal everywhere, but the measured leaf xylem pressure would differ for leaves at different height.

Similar logic applies to xylem water potential and xylem water pressure, and so does the thermocouple psychrometers method (Boyer and Knippling, 1965; Boyer, 1968). ~~Given that water potential does not properly represent *in vivo* water pressure, one~~

80 ~~should avoid using the term water potential to describe plant's vulnerability curve~~ It is recommended to refer to the measurement as leaf/xylem water pressure or balance pressure in the future, rather than leaf/xylem water potential.

3 Hydraulic conductance and conductivity

Hydraulic conductance (k) and conductivity (K) are also often confused in the literature (e.g., Kannenberg et al., 2019; Cardoso et al., 2020; Li et al., 2021). Hydraulic conductance (flow rate divided by ~~potential difference~~ driving pressure) and electrical

85 conductance (current divided by voltage difference) are bulk properties, and the two are analogous. However, while electrical conductivity is the intrinsic property of a uniform material, hydraulic conductivity is a bulk property given the nonuniform structure of the tissue (e.g., only the sapwood is hydraulically conductive). The most widely used definitions for conductance and conductivity are: (a) hydraulic conductance (namely k) is the ratio between flow rate through the segment (E) and ~~potential~~

~~difference~~ (ΔP driving pressure ($\Delta P - \rho g \Delta h$)), and segment length (L) and area not accounted for by k , (b) hydraulic conduc-

90 tivity (namely K) is the ratio between flow rate and ~~potential~~ driving pressure gradient (area not accounted for), (c) sapwood area specific hydraulic conductivity (K_S) is the ratio between hydraulic conductivity and xylem sapwood area (A_S), and (d) leaf area specific hydraulic conductivity (K_L) is the ratio between hydraulic conductivity and leaf area the xylem supports

(A_L):

$$k = \frac{E}{\Delta P - \rho g \Delta h}, \quad (1)$$

95 ~~$K = k \cdot L$,~~ $K = k \cdot L, \quad (2)$

$$K_S = \frac{K}{A_S}, \quad (3)$$

$$K_L = \frac{K}{A_L}. \quad (4)$$

Note that only K_S and K_L are per unit conducting area, and thus can be treated as “intrinsic” properties for comparison purpose: K_S for sapwood water permeability and K_L for leaf water supply capability. However, K_L may not best describe leaf

100 water supply capability. For example, if two branches have the same K_S , leaf area, and sapwood area, but only differ in their length, the computed K and K_L would be the same for the two branches even though the actual leaf water supply capabilities differ. In comparison, conductance of the entire branch divided by leaf area of the branch, i.e., ~~leaf area specific hydraulic~~

~~conductance~~ leaf area specific hydraulic conductance (k_L) as inspired by leaf area specific whole plant hydraulic conductance, would be a better measure for leaf water supply. The k_L can be estimated using

105 $k_L = \frac{k_{\text{branch}}}{A_L}, \quad (5)$

where k_{branch} is hydraulic conductance of the ~~entire~~ entire branch (not a stem segment).

4 Xylem vulnerability curve (VC)

Various formulas have been used to represent xylem VC, and the three most common ones are Weibull cumulative probability function (equation 6) (e.g., Sperry et al., 2016; Love et al., 2019), logistic function (equation 7) (e.g., Feng et al., 2018; Huber et al., 2019), and power function (equation 8) (e.g., Eller et al., 2018; Liu et al., 2020):

$$\frac{k}{k_{\max}} = \exp \left[- \left(\frac{-P}{B} \right)^C \right] = 2^{- \left(\frac{P}{P_{50}} \right)^C}, \quad (6)$$

$$\frac{k}{k_{\max}} = 1 - \frac{1}{1 + a \cdot \exp(b \cdot P)} = 1 - \frac{1}{1 + \exp[b \cdot (P - P_{50})]}, \quad (7)$$

$$\frac{k}{k_{\max}} = \frac{1}{1 + m \cdot (-P)^n} = \frac{1}{1 + \left(\frac{P}{P_{50}} \right)^n}, \quad (8)$$

where B , C , a , b , m , and n are vulnerability function parameters, and P_{50} is the water pressure at which the tissue loses 50% of its conductance. Note that there are also more complex VC formulations based on the three, such as dual-Weibull function used in hydraulic fiber bridge (Cai et al., 2014; Pan and Tyree, 2019) and cavitation fatigue (Feng et al., 2015; Zhang et al., 2018).

We should be aware that the logistic VC function (equation 7, or formulation based on it) does not always start from 1 when $P = 0$. This problem is minor for sigmoidal VCs (s-shaped); however, the offset at $P = 0$ could introduce bias if the VC becomes more exponential (r-shaped, see Fig. 5 of Huber et al. (2019) for an example). In this case, fitting VC using equation 7 would result in overestimated k_{\max} and less negative P_{50} . Thus, equation 7 should be rescaled to minimize the bias, and the modified formulation is

$$\frac{k}{k_{\max}} = \frac{a \cdot \exp(b \cdot P)}{1 + a \cdot \exp(b \cdot P)} \cdot \frac{1 + a}{a} = \frac{(1 + a) \cdot \exp(b \cdot P)}{1 + a \cdot \exp(b \cdot P)}, \quad (9)$$

$$P_{50} = - \frac{\log(2 + a)}{b}. \quad (10)$$

125 5 ~~Division and derivative~~ Hydraulic model complexity

~~A Plant hydraulic models have various complexities depending on the various aims of research and difficulties in model parameterization (Tyree and Ewers, 1991; Tyree and Zimmermann, 2002). In terms of flow profiles, the models can be categorized to steady state and non-steady state models. The steady state models use a constant flow rate within roots, stem, and leaves. The non-steady state models employ a changing flow rate within or among different tissues given the water exchange between xylem and capacitance tissues. In terms of the model complexity, the models range from a single element to a xylem network (say multiple roots and multiple canopy layers). Further, hydraulic conductance of an element may change with the growth of plants; for example, the drought legacy, maximum hydraulic conductance, and VC vary with the stack of new tree rings (McCulloh and Sperry, 2005; Cai and Tyree, 2010). Although more complex models may better represent the water flow and pressure profiles within the plants, increasing difficulties in model parameterization makes these more complex models less~~

135 appealing to users. However, inappropriate model selection could result in biased results, for instance, modeling plant hydraulics at steady state for plants with high water capacity and ignoring vessel tapering effect when modeling xylem growth. Thus, it is important to select plant hydraulic models with adequate complexity in topical research. See the section below for a detailed example of how reduced model complexity (ignoring VC segmentation) may bias the modeled hydraulic risk and thus stomatal responses.

140 **6 Stomatal model representation**

Plant hydraulics-based stomatal models are gaining increasing interest in the vegetation and land modeling communities (e.g., Kennedy et al., 2019; Sabot et al., 2020) as they predict stomatal closure at dry environmental conditions without employing an arbitrary tuning factor (often known as the β factor) (Powell et al., 2013). For instance, the recently developed optimality theory-based models propose that plants should balance the gain and risk associated with stomatal functioning (Wolf et al., 2016; Sperry et al., 2017).
 145 When plants open their stomata more, plants gain more photosynthetic carbon, but lose more water and have higher risk in hydraulic failure; therefore, plants are supposed to find a sweet zone to maximize the difference between the gain and risk. These optimality theory models, particularly those weigh the risk based on plant hydraulics, show comparable or better predictive skills compared to the statistical approaches (Anderegg et al., 2018; Eller et al., 2018; Venturas et al., 2018; Wang et al., 2020; Sperry et al., 2017). However, a common mistake when using plant hydraulics-based models is that one does not follow the original model formula or hypothesis. For example, the Sperry et al. (2017) model defines the risk of stomatal opening associated with stomatal functioning (Θ) as

$$\Theta = A_{\max} \cdot \frac{1 - \frac{k_{\text{canopy,ref}} - k_{\text{canopy}}(P_{\text{canopy}})}{k_{\text{canopy,ref}} - k_{\text{crit}}}}{k_{\text{canopy,ref}} - k_{\text{crit}}}, \quad (11)$$

$$k_{\text{canopy}} = \frac{dE}{dP}, \quad (12)$$

where A_{\max} is the maximal achievable photosynthetic rate at the given setting, P_{canopy} is the leaf xylem end water pressure in the canopy, and $k_{\text{canopy,ref}}$ is the maximum k_{canopy} when transpiration rate is 0, and k_{crit} is the k_{canopy} when transpiration rate is maximum. The k_{crit} by definition is 0, as a minimum incremental transpiration rate results in infinity increase in xylem pressure (the $dE/dP = 0$ in Figure 2). Thus, in the subsequent research where Sperry et al. (2017) was tested (Venturas et al., 2018; Wang et al., 2020), the model has been reformulated to

$$\Theta = A_{\max} \cdot \left[1 - \frac{k_{\text{canopy}}(P_{\text{canopy}})}{k_{\text{canopy,ref}}} \right]. \quad (13)$$

160 Note here that k_{canopy} is the derivative of a water supply curve at given soil water potential and canopy water potentialspressure, and $k_{\text{canopy}}/k_{\text{canopy,ref}}$ is different from (a) relative conductance of root, stem, or leaf xylem (i.e., k/k_{ref} , where k_{ref} is the maximum k at a reference xylem pressure), and (b) relative whole-plant hydraulic conductance ($k_{\text{plant}} = E/\Delta\Psi$, $k_{\text{plant}} = E/DP$). However, model descriptions from various sources may be contrasting: Sperry et al. (2017) expressed their risk in derivative form; Eller et al. (2018) expressed their risk calculations in three completely different ways including division (their equation

165 2.3), derivative (their equation 2.6), and point estimation (their equation 2.8); Mencuccini et al. (2019) interpreted the two models based on k_{plant} ; Wang et al. (2020) interpreted the two models based on the derivative forms. See Fig. 2 for how the three quantities differ.




Figure 2. Difference between quantities used in plant hydraulics. E : flow rate through the xylem, P : xylem water pressure, k xylem hydraulic conductance. The simulation is done for a plant with soil water potential of -0.5 MPa, no gravity term, and no drought legacy effect from previous xylem embolism.

For a xylem that does not have height change or VC segmentation, $dE/dP = k(P)$. Otherwise, using $k(P)$ to proxy dE/dP could result in biases, particularly when gravity is not negligible and when tissue VCs differ dramatically (Sperry et al., 2016; McCulloh et al., 2019). Using the parameters of a real plant as an example (data from Wang et al., 2019), it is obvious that none of the root, stem, leaf, or whole-plant hydraulic conductance is a good dE/dP proxy (Fig. 3). Therefore, researchers should test the models that differ from the original forms. To note, the primary reason that Sperry et al. (2017) used dE/dP was to account for the VC segmentation. Using stem VC (easiest to measure; typically more resistant than roots and leaves) to proxy root and leaf VCs in stomatal models would likely result in less sensitive stomatal response to environmental stimuli such as soil moisture (Fig. 3).

7 Analytic solution and estimation

The pursuit of simplicity and analytic solution often leads to biased results, for example, using leaf or stem VC as a proxy for dE/dP (e.g., Fig. 3) and ignoring the impact of gravity (e.g., Fig. 4). As a result, it is important to distinguish true analytic ~~solution from analytic estimation~~ solution from analytic estimation. For example in a xylem water supply curve, when gravitational pressure drop is neglected, flow rate at a given canopy water pressure will be overestimated (Fig. 4). The more the




Figure 3. Comparison of different risk measures of stomatal opening. **(a)–(e)** Values are relative to the maximum when soil water potential is 0 and transpiration rate is 0. **(f)–(j)** Values are relative to the maximal when transpiration rate is 0 at the given soil water potential. Ψ_{soil} : soil water potential, E : transpiration rate of the whole plant, E_{crit} : maximum transpiration rate of the whole plant beyond which the plant desiccate, dE/dP : derivative of a water supply curve (a plot of E vs. top of canopy xylem water pressure, P), k : hydraulic conductance at a given xylem water pressure, k_{max} : maximum k when there is no cavitation. The simulation is done assuming there is no drought legacy effect from previous xylem embolism. For the simulation, the plant has a root:stem:leaf resistance ratio of 2:1:1; root and stem height change are 1 and 10 m, respectively; VCs are represented using a Weibull function; Weibull B in MPa and C are 1.879 and 2.396 for root, 2.238 and 9.380 for stem, and 1.897 and 2.203 for leaf (data from Wang et al., 2019).

height changes, the more E is overestimated (Fig. 4). The absolute value of E overestimation decreases with more negative soil water potential, whereas the relative E overestimation increases with more negative soil water potential (Fig. 4). Thus, given the potentially great biases, it is recommended to verify any analytic or numerical estimations against true numerical solution before using them in research.

185 8 Whole-plant vulnerability “curve”

The ideas of whole-plant conductance and VC largely advance the understanding of how plant traits coordinate as they provide a simple way to correlate different traits (see McCulloh et al. (2019) for an overview). Whole-plant hydraulic conductance (k_{plant}) depends on not only the upstream water potential (namely Ψ_{soil}) but also the downstream water potential (Ψ_{canopy}):

$$k_{\text{plant}} = f(\Psi_{\text{soil}}, \Psi_{\text{canopy}}) = \frac{E}{\Psi_{\text{canopy}} - \Psi_{\text{soil}}} \text{ pressure } (P_{\text{canopy}}): k_{\text{plant}} = f(\Psi_{\text{soil}}, P_{\text{canopy}}) = \frac{E}{DP}$$

190 the hidden assumptions when using the term *whole-plant hydraulic conductance* whole-plant hydraulic conductance (or any similar terms): upstream water potential (soil water potential in this scenario) is the same everywhere, and *downstream water*




Figure 4. Water supply curve at different soil water potential for xylem with different height change. E : xylem flow rate, P : xylem water pressure. (a)–(d) Water supply curves when soil water potential is 0, -0.5 , -1 , and -1.5 MPa, respectively. The simulation is done assuming there is no drought legacy effect from previous xylem embolism.

potential (leaf-water-potential-in-this-scenario) *driving pressure* is the same everywhere, regardless of plant height, canopy light conditions, and root/stem/leaf network. Therefore, in the practice of modeling or research, the two assumptions are barely met. Further, note that k_{plant} is a bulk parameter from root to leaves, and xylem water pressure and xylem hydraulic conductance are profiles rather than being constant along the flow path. Therefore, by definition, there is not a whole-plant vulnerability *curve*; instead, $f(\Psi_{\text{soil}}, \Psi_{\text{canopy}}) - f(\Psi_{\text{soil}}, P_{\text{canopy}})$ is a whole-plant vulnerability *surface* (Fig. 5). It is obvious that none of Ψ_{soil} , P_{canopy} , or a mean pressure can predict a unique k_{plant} (although the change of k_{plant} is relatively smaller for the mean pressure; dotted line in Fig. 5). Further, drought legacy effect from previous non-refillable xylem embolism (Anderegg et al., 2015) would further complicate the scenario as the “surface” changes with drought legacy. Therefore, it is not recommended to use the term *whole-plant vulnerability curve* *whole-plant vulnerability curve* in research.

9 Temperature effects

When modeling plant hydraulics, the temperature effects on viscosity (η), surface tension (γ), and osmotic potential are typically ignored. However, when water temperature decreases from 25 °C to 10 °C (298.15 K to 283.15 K), (1) viscosity of water increases by 43.8% , meaning a $> 40\%$ increase in pressure drop along the flow path for a given flow rate; (2) surface tension of water increases by 3.1% , meaning that capillary force withholding the air-water interface at the pit membrane increases by 3.1% for a given curvature radius (xylem becomes more resistant to cavitation) and that soil metric potential becomes 3.1% more negative for a given soil water content; (3) soil osmotic water potential would be 5.0% less negative for a given ion concentration. Therefore, a more reasonable way to describe a xylem VC (e.g., using Weibull function) should be:

$$k = k_{\text{max},25} \cdot \frac{\eta_{25}}{\eta} \cdot \exp \left[- \left(\frac{-P}{B_{25}} \cdot \frac{\gamma_{25}}{\gamma} \right)^{C_{25}} \right], \quad (14)$$




Figure 5. Whole-plant vulnerability surface. Whole-plant hydraulic conductance is computed using $k_{\text{plant}} = \frac{E}{\Psi_{\text{soil}} - \Psi_{\text{canopy}}}$ where Ψ_{soil} is soil water potential and $\Psi_{\text{canopy}} = P + \rho gh$, where P is canopy water potential. Ψ_{canopy} differs from canopy xylem water pressure (P) in that P , and h is more negative than Ψ_{canopy} by the gravitational pressure drop height. The dotted line plots the scenario when mean xylem water pressure is -1.6 MPa. The simulation is done assuming there is no drought legacy effect from previous xylem embolism.

210 where the subscript 25 denotes the values are at a reference temperature of 25 °C. In other words, k_{max} needs to be scaled to $k_{\text{max},25} \cdot \eta_{25}/\eta$, and P needs to be scaled to $P \cdot \gamma_{25}/\gamma$.

10 Conclusions

Plant hydraulics is often improperly represented in research, potentially resulting in ambiguities to those who are not familiar with the terminologies. This paper documents differences among commonly seen ambiguous and miscellaneous terms that are often not recognized, and the mistakes and misunderstandings researchers may make when using established methods and models. The mathematics and visualizations of the documented items will help researchers in their research and teaching associated with plant hydraulics.

Author contributions. YW and CF wrote the manuscript.

Competing interests. No competing interests

220 *Acknowledgements.* We gratefully acknowledge the generous support of Eric and Wendy Schmidt (by recommendation of the Schmidt Futures) and the Heising-Simons Foundation. This research has been supported by the National Aeronautics and Space Administration (NASA) Carbon Cycle Science grant 80NSSC21K1712 awarded to Christian Frankenberg. We benefit from the discussion with Dr. Tom Buckley for the proper usages of water potential and water pressure.

References

- 225 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Venetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. T., et al.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *Forest Ecology and Management*, 259, 660–684, 2010.
- Anderegg, W. R. L., Berry, J. A., Smith, D. D., Sperry, J. S., Anderegg, L. D., and Field, C. B.: The roles of hydraulic and carbon stress in a widespread climate-induced forest die-off, *Proceedings of the National Academy of Sciences*, 109, 233–237, 2012.
- 230 Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Shevliakova, E., Williams, A., Wolf, A., Ziaco, E., and Pacala, S.: Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models, *Science*, 349, 528–532, 2015.
- Anderegg, W. R. L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A. F. A., Choat, B., and Jansen, S.: Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe, *Proceedings of the National Academy of Sciences*, 113, 5024–5029, 2016.
- 235 Anderegg, W. R. L., Wolf, A., Arango-Velez, A., Choat, B., Chmura, D. J., Jansen, S., Kolb, T., Li, S., Meinzer, F. C., Pita, P., Resco de Dios, V., Sperry, J. S., Wolfe, B. T., and Pacala, S.: Woody plants optimise stomatal behaviour relative to hydraulic risk, *Ecology Letters*, 21, 968–977, 2018.
- Boyer, J. S.: Leaf water potentials measured with a pressure chamber, *Plant Physiology*, 42, 133–137, 1967.
- 240 Boyer, J. S.: Relationship of water potential to growth of leaves, *Plant physiology*, 43, 1056–1062, 1968.
- Boyer, J. S. and Naylor, E. B.: Isopiestic technique for measuring leaf water potentials with a thermocouple psychrometer, *Proceedings of the National Academy of Sciences of the United States of America*, 54, 1044, 1965.
- Buckley, T. N. and Mott, K. A.: Modelling stomatal conductance in response to environmental factors, *Plant, Cell & Environment*, 36, 1691–1699, 2013.
- 245 Cai, J. and Tyree, M. T.: The impact of vessel size on vulnerability curves: Data and models for within-species variability in saplings of aspen, *Populus tremuloides* Michx., *Plant, Cell & Environment*, 33, 1059–1069, 2010.
- Cai, J., Li, S., Zhang, H., Zhang, S., and Tyree, M. T.: Recalcitrant vulnerability curves: Methods of analysis and the concept of fibre bridges for enhanced cavitation resistance, *Plant, Cell & Environment*, 37, 35–44, 2014.
- Cardoso, A. A., Visel, D., Kane, C. N., Batz, T. A., García Sánchez, C., Kaack, L., Lamarque, L. J., Wagner, Y., King, A., Torres-Ruiz, J. M., 250 et al.: Drought-induced lacuna formation in the stem causes hydraulic conductance to decline before xylem embolism in *Selaginella*, *New Phytologist*, 227, 1804–1817, 2020.
- Chen, C., Riley, W. J., Prentice, I. C., and Keenan, T. F.: CO₂ fertilization of terrestrial photosynthesis inferred from site to global scales, *Proceedings of the National Academy of Sciences*, 119, e2115627 119, 2022.
- Cowan, I. R. and Farquhar, G. D.: Stomatal function in relation to leaf metabolism and environment, *Symposia of the Society for Experimental 255 Biology*, 31, 471–505, 1977.
- Dewar, R., Mauranen, A., Mäkelä, A., Hölttä, T., Medlyn, B., and Vesala, T.: New insights into the covariation of stomatal, mesophyll and hydraulic conductances from optimization models incorporating nonstomatal limitations to photosynthesis, *New Phytologist*, 217, 571–585, 2018.

- 260 Eller, C. B., Rowland, L., Oliveira, R. S., Bittencourt, P. R. L., Barros, F. V., da Costa, A. C. L., Meir, P., Friend, A. D., Mencuccini, M., Sitch, S., and Cox, P.: Modelling tropical forest responses to drought and El Niño with a stomatal optimization model based on xylem hydraulics, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20170315, <https://doi.org/10.1098/rstb.2017.0315>, 2018.
- Eller, C. B., Rowland, L., Mencuccini, M., Rosas, T., Williams, K., Harper, A., Medlyn, B. E., Wagner, Y., Klein, T., Teodoro, G. S., et al.: Stomatal optimization based on xylem hydraulics (SOX) improves land surface model simulation of vegetation responses to climate, *New Phytologist*, 226, 1622–1637, 2020.
- 265 Feng, F., Ding, F., and Tyree, M. T.: Investigations concerning cavitation and frost fatigue in clonal 84K poplar using high-resolution cavitron measurements, *Plant Physiology*, 168, 144–155, 2015.
- Feng, X., Ackerly, D. D., Dawson, T. E., Manzoni, S., Skelton, R. P., Vico, G., and Thompson, S. E.: The ecohydrological context of drought and classification of plant responses, *Ecology letters*, 21, 1723–1736, 2018.
- Gleason, S. M., Westoby, M., Jansen, S., Choat, B., Hacke, U. G., Pratt, R. B., Bhaskar, R., Brodribb, T. J., Bucci, S. J., Cao, K.-F., et al.: 270 Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world’s woody plant species, *New Phytologist*, 209, 123–136, 2016.
- Huber, A. E., Melcher, P. J., Piñeros, M. A., Setter, T. L., and Bauerle, T. L.: Signal coordination before, during and after stomatal closure in response to drought stress, *New Phytologist*, 224, 675–688, 2019.
- Kannenbergh, S. A., Novick, K. A., and Phillips, R. P.: Anisohydric behavior linked to persistent hydraulic damage and delayed drought 275 recovery across seven North American tree species, *New Phytologist*, 222, 1862–1872, 2019.
- Kennedy, D., Swenson, S., Oleson, K. W., Lawrence, D. M., Fisher, R., Lola da Costa, A. C., and Gentine, P.: Implementing plant hydraulics in the community land model, version 5, *Journal of Advances in Modeling Earth Systems*, 11, 485–513, 2019.
- Li, S., Fang, L., Hegelund, J. N., and Liu, F.: Elevated CO₂ modulates plant hydraulic conductance through regulation of PIPs under progressive soil drying in tomato plants, *Frontiers in plant science*, 12, 666066, <https://doi.org/10.3389/fpls.2021.666066>, 2021.
- 280 Liu, H., Ye, Q., Gleason, S. M., He, P., and Yin, D.: Weak tradeoff between xylem hydraulic efficiency and safety: climatic seasonality matters, *New Phytologist*, 229, 1440–1452, 2021.
- Liu, Y., Kumar, M., Katul, G. G., Feng, X., and Konings, A. G.: Plant hydraulics accentuates the effect of atmospheric moisture stress on transpiration, *Nature Climate Change*, 10, 691–695, 2020.
- Love, D. M., Venturas, M. D., Sperry, J. S., Brooks, P. D., Pettit, J. L., Wang, Y., Anderegg, W. R. L., Tai, X., and Mackay, D. S.: Dependence 285 of aspen stands on a subsurface water subsidy: Implications for climate change impacts, *Water Resources Research*, 55, 1833–1848, 2019.
- Manzoni, S., Vico, G., Palmroth, S., Porporato, A., and Katul, G.: Optimization of stomatal conductance for maximum carbon gain under dynamic soil moisture, *Advances in Water Resources*, 62, 90–105, 2013.
- McCulloh, K. A. and Sperry, J. S.: Patterns in hydraulic architecture and their implications for transport efficiency, *Tree Physiology*, 25, 257–267, 2005.
- 290 McCulloh, K. A., Domec, J.-C., Johnson, D. M., Smith, D. D., and Meinzer, F. C.: A dynamic yet vulnerable pipeline: Integration and coordination of hydraulic traits across whole plants, *Plant, Cell & Environment*, 42, 2789–2807, 2019.
- McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D. G., et al.: Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought?, *New phytologist*, 178, 719–739, 2008.
- 295 McDowell, N. G.: Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality, *Plant physiology*, 155, 1051–1059, 2011.

- McDowell, N. G., Fisher, R. A., Xu, C., Domec, J.-C., Hölttä, T., Mackay, D. S., Sperry, J. S., Boutz, A., Dickman, L., Gehres, N., et al.: Evaluating theories of drought-induced vegetation mortality using a multimodel–experiment framework, *New Phytologist*, 200, 304–321, 2013.
- 300 Meinzer, F. C., McCulloh, K. A., Lachenbruch, B., Woodruff, D. R., and Johnson, D. M.: The blind men and the elephant: the impact of context and scale in evaluating conflicts between plant hydraulic safety and efficiency, *Oecologia*, 164, 287–296, 2010.
- Mencuccini, M., Manzoni, S., and Christoffersen, B.: Modelling water fluxes in plants: From tissues to biosphere, *New Phytologist*, 222, 1207–1222, 2019.
- Pan, R. and Tyree, M. T.: How does water flow from vessel to vessel? Further investigation of the tracheid bridge concept, *Tree physiology*, 39, 1019–1031, 2019.
- 305 Powell, T. L., Galbraith, D. R., Christoffersen, B. O., Harper, A., Imbuzeiro, H., Rowland, L., Almeida, S., Brando, P. M., Costa, A. C. L., Costa, M. H., and Moorcroft, P. R.: Confronting model predictions of carbon fluxes with measurements of Amazon forests subjected to experimental drought, *New Phytologist*, 200, 350–365, 2013.
- Sabot, M. E., De Kauwe, M. G., Pitman, A. J., Medlyn, B. E., Verhoef, A., Ukkola, A. M., and Abramowitz, G.: Plant profit maximization improves predictions of European forest responses to drought, *New Phytologist*, 226, 1638–1655, 2020.
- 310 Sabot, M. E., De Kauwe, M. G., Pitman, A. J., Medlyn, B. E., Ellsworth, D. S., Martin-StPaul, N., Wu, J., Choat, B., Limousin, J.-M., Mitchell, P. J., et al.: One Stomatal Model to Rule Them All? Towards Improved Representation of Carbon and Water Exchange in Global Models, *Journal of Advances in Modeling Earth Systems*, p. e2021MS002761, <https://doi.org/10.1029/2021MS002761>, 2022.
- Santiago, L. S., Goldstein, G., Meinzer, F. C., Fisher, J. B., Machado, K., Woodruff, D., and Jones, T.: Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian forest canopy trees, *Oecologia*, 140, 543–550, 2004.
- 315 Scholander, P. F., Hammel, H., Hemmingsen, E., and Bradstreet, E.: Hydrostatic pressure and osmotic potential in leaves of mangroves and some other plants, *Proceedings of the National Academy of Sciences of the United States of America*, 52, 119–125, 1964.
- Sperry, J. S. and Love, D. M.: What plant hydraulics can tell us about responses to climate-change droughts, *New Phytologist*, 207, 14–27, 2015.
- 320 Sperry, J. S. and Tyree, M. T.: Mechanism of water stress-induced xylem embolism, *Plant Physiology*, 88, 581–587, 1988.
- Sperry, J. S., Adler, F. R., Campbell, G. S., and Comstock, J. P.: Limitation of plant water use by rhizosphere and xylem conductance: Results from a model, *Plant, Cell & Environment*, 21, 347–359, 1998.
- Sperry, J. S., Wang, Y., Wolfe, B. T., Mackay, D. S., Anderegg, W. R. L., McDowell, N. G., and Pockman, W. T.: Pragmatic hydraulic theory predicts stomatal responses to climatic water deficits, *New Phytologist*, 212, 577–589, 2016.
- 325 Sperry, J. S., Venturas, M. D., Anderegg, W. R. L., Mencuccini, M., Mackay, D. S., Wang, Y., and Love, D. M.: Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost, *Plant, Cell & Environment*, 40, 816–830, 2017.
- Tyree, M. T. and Ewers, F. W.: The hydraulic architecture of trees and other woody plants, *New Phytologist*, 119, 345–360, 1991.
- Tyree, M. T. and Sperry, J. S.: Vulnerability of xylem to cavitation and embolism, *Annual review of plant biology*, 40, 19–36, 1989.
- Tyree, M. T. and Zimmermann, M. H.: *Xylem structure and the ascent of sap*, Springer Science & Business Media, 2002.
- 330 Venturas, M. D., Sperry, J. S., Love, D. M., Frehner, E. H., Allred, M. G., Wang, Y., and Anderegg, W. R. L.: A stomatal control model based on optimization of carbon gain versus hydraulic risk predicts aspen sapling responses to drought, *New Phytologist*, 220, 836–850, 2018.
- Wang, Y., Sperry, J. S., Venturas, M. D., Trugman, A. T., Love, D. M., and Anderegg, W. R. L.: The stomatal response to rising CO₂ concentration and drought is predicted by a hydraulic trait-based optimization model, *Tree Physiology*, 39, 1416–1427, 2019.

- 335 Wang, Y., Sperry, J. S., Anderegg, W. R. L., Venturas, M. D., and Trugman, A. T.: A theoretical and empirical assessment of stomatal optimization modeling, *New Phytologist*, 227, 311–325, 2020.
- Wang, Y., Anderegg, W. R., Venturas, M. D., Trugman, A. T., Yu, K., and Frankenberg, C.: Optimization theory explains nighttime stomatal responses, *New Phytologist*, 230, 1550–1561, 2021a.
- 340 Wang, Y., Köhler, P., He, L., Doughty, R., Braghieri, R. K., Wood, J. D., and Frankenberg, C.: Testing stomatal models at the stand level in deciduous angiosperm and evergreen gymnosperm forests using CliMA Land (v0.1), *Geoscientific Model Development*, 14, 6741–6763, 2021b.
- Wolf, A., Anderegg, W. R. L., and Pacala, S. W.: Optimal stomatal behavior with competition for water and risk of hydraulic impairment, *Proceedings of the National Academy of Sciences*, 113, E7222–E7230, 2016.
- Zhang, W., Feng, F., and Tyree, M. T.: Seasonality of cavitation and frost fatigue in *Acer mono Maxim.*, *Plant, Cell & Environment*, 41, 1278–1286, 2018.