

Dispersal of bacteria and stimulation of permafrost decomposition by Collembola

Sylvain Monteux^{1,2,3}, Janine Mariën⁴, Eveline J. Krab^{1,3}

¹Department of Soil and Environment, Sveriges Lantbruksuniversitet SLU, Uppsala, 750 07, Sweden

5 ²Department of Environmental Science, Stockholms Universitet, Stockholm, 106 91, Sweden

³Climate Impacts Research Centre, Umeå Universitet, Abisko, 981 07, Sweden

⁴Department of Animal Ecology, Vrije Universiteit Amsterdam, Amsterdam, 1081 HV, The Netherlands


Correspondence to: Sylvain Monteux (monteux@pm.me)

Abstract. Contrary to most soils, permafrost soils have the atypical feature of being almost entirely deprived of soil fauna. Abiotic constraints on the fate of permafrost carbon after thawing are increasingly understood, but biotic constraints remain scarcely investigated. Incubation studies, essential to estimate effects of permafrost thaw on carbon cycling, typically measure the consequences of permafrost thaw in isolation from the topsoil, and thus do not account for the effects of altered biotic interactions because of e.g. colonization by soil fauna. Microarthropods facilitate the dispersal of microorganisms in soil, both on their cuticle (ectozoochory) and through their digestive tract (endozoochory), which may be particularly important in permafrost soils, considering that microbial community composition can strongly constrain permafrost biogeochemical processes.

Here we tested how a model species of microarthropod (the Collembola *Folsomia candida*) affected aerobic CO₂ production of permafrost soil over a 25 days incubation. By using collembola stock cultures grown on permafrost soil or on an arctic topsoil, we aimed to assess the potential for endo- and ectozoochory of soil bacteria, while cultures grown on gypsum and sprayed with soil suspensions would allow to observe only ectozoochory.

The presence of collembola introduced bacterial ASVs absent in the no-collembola control, regardless of their microbiome manipulation, when considering presence-absence metrics (unweighted UniFrac metrics), which resulted in increased species richness. However, these introduced ASVs did not induce changes in bacterial community composition as a whole (accounting for relative abundances, weighted UniFrac), which might only become detectable in the longer-term.

CO₂ production was increased by 25.85% in the presence of collembola, about half of which could be attributed to collembola respiration based on respiration rates measured in the absence of soil. We argue that the rest of the CO₂ being respired can be considered a priming effect of the presence of collembola, i.e. a stimulation of permafrost CO₂ production in the presence of active microarthropod decomposers. Overall, our findings underline the importance of biotic interactions in permafrost biogeochemical processes, and the need to explore the additive or interactive effects of other soil food web groups of which permafrost soils are deprived.



Graphical abstract. Yedoma permafrost was incubated in absence or presence of *Folsomia candida* collembola, and CO₂ production and bacterial community composition were assessed after 25 days. The presence of collembola increased CO₂ production by 26%, of which half was attributed to collembola respiration and the other half to priming effects. Collembola presence further modified bacterial community composition (16S V4V5), although this effect was independent of the different treatments aiming to manipulate the skin and/or gut microbiome of the collembola.

1. Introduction

Carbon fluxes from soils are largely governed by the rate of decomposition of organic matter. Soil fauna is a crucial component in organic matter decomposition (García-Palacios et al., 2013; Griffiths et al., 2021), both directly through mechanical degradation of litter and corpses into smaller pieces, and indirectly through feeding habits controlling the abundances of other decomposer groups such as fungi, microbial eukaryotes or bacteria (Hanlon and Anderson, 1979; Kaneda and Kaneko, 2008; Frouz et al., 2020; Potapov et al., 2020). In arctic soils, the scarcity of macrofaunal decomposers (e.g. earthworms: Blume-Werry et al., 2020) coupled to the high abundance of microbivorous microarthropods such as collembola (Potapov et al., 2022), results in a particularly strong impact of collembola on decomposition through microbial population control (Koltz et al., 2018, Crowther et al., 2012, Seastedt 1984).

Contrary to other important groups of soil fauna, such as earthworms, millipedes (Golovatch and Kime, 2009; Berman et al., 2015) or woodlice (Sfenthourakis and Hornung, 2018), collembola are ubiquitous in Arctic soils where they can reach high
45 densities – up to 130,000 individuals per m² in high arctic Greenland (Sørensen et al., 2006). However, collembola are mostly abundant in the topsoil, and to our best knowledge have never been observed in the perennially-frozen subsoil, the permafrost. Permafrost soils are a prominent feature of Arctic landscapes, and the huge carbon stock they represent and positive feedback to climate change that their thawing will likely induce causes concern (Masson-Delmotte et al., 2021). The
50 frozen conditions over long periods of time have eliminated most fauna and non-microbial life from these environments, and despite the possibility of resuscitating organisms such as plants (Yashina et al., 2012), nematodes (Shatilovich et al., 2018) or rotifers (Shmakova et al., 2021), newly-thawed permafrost soils generally harbour an extremely simplified soil food web entirely deprived of metazoans. Despite this particularity, the absence of collembola and the consequence of their possible introduction on biogeochemical cycling of newly-thawed permafrost has been mostly overlooked.

55 In contrast to collembola, some microorganisms survive and/or thrive in permafrost conditions, and microbial adaptations to frozen conditions have been studied (Mackelprang et al., 2011, 2017; Hultman et al., 2015; Bottos et al., 2018). However, not all microbes survive these conditions, and the combination of environmental constraints exerted over long periods of time (Mackelprang et al., 2017) and strong dispersal limitations (Bottos et al., 2018) result in microbial communities that can be deprived of some functions (Knoblauch et al., 2018; Monteux et al., 2020; Barbato et al., 2022). The re-introduction of
60 such functions can result in drastic changes in permafrost processes, and sizable impacts on greenhouse gas production have been observed in vitro for CH₄ and CO₂ (Knoblauch et al., 2018; Monteux et al., 2020) and confirmed in situ for N₂O (Marushchak et al., 2021). Upon thawing, this re-introduction of missing function, or ecological rescue (Calderón et al., 2017) requires microorganisms to migrate into this newly-available habitat, which could happen for instance laterally through airborne dispersal (Harding et al., 2011) – e.g. for permafrost exposed to the air in abrupt thaw processes (Inglese et al., 2017) – or vertically through percolation in the soil column when the active layer becomes deeper (Monteux et al., 2018;
65 Johnston et al., 2019).

Collembola are another possible vector for the dispersal of microorganisms into newly-thawed permafrost (Buse et al., 2014). With the deepening of the active layer, collembola migration into newly-thawed permafrost is unlikely since they
70 mostly reside in topsoil layers, but it will likely occur in soil mixing events (Väisänen et al., 2020) such as thermokarst, active layer detachment or thaw slump processes where newly-thawed permafrost is exposed to surface conditions. Like all large organisms, collembola host a variety of microorganisms, their microbiome (Agamennone et al., 2015; Leo et al., 2021). Because micro-arthropods, such as collembola, can move across large distances compared to fungi or bacteria, the microbial species in collembolan microbiomes might be among the first to colonize and establish into newly-thawed permafrost when
75 collembola access it. This could occur through a combination of two main processes: ectozoochory, where the microbiome of the cuticle disperses into the new habitat, and endozoochory where microorganisms disperse after transiting through the

gut of the animals. To our best knowledge, whether collembola affect the biogeochemical functioning of newly-thawed permafrost, and whether and how they can serve as a vector for microbial colonization has not been explored yet..

80 We incubated permafrost from the Yedoma domain, which represents a large carbon stock in parts of Siberia and North-
America (Strauss et al., 2017) and has previously been shown to lack certain microbial functions (Monteux et al., 2020), in
the presence or absence of a model species of microarthropod (collembola *Folsomia candida*), and assessed CO₂ production
and bacterial community composition over a 25 days aerobic incubation. We used *Folsomia candida* collembola from a
stock culture, as well as collembola subjected to manipulation of their cuticle microbiome, or cuticle and gut microbiomes,
85 to test the following hypotheses:

1. Collembola presence in permafrost will alter bacterial community composition, through their grazing.
2. Collembola more closely exposed to topsoil bacteria will change permafrost bacterial community composition
further than those exposed less intricately. In other words, we expect a gradual change between clean collembola,
collembola with cuticle microbiome manipulated (ectozoochory) and collembola with both gut and cuticle
90 microbiomes manipulated (endozoochory).
3. Collembola presence in permafrost will increase CO₂ production, through both collembola respiration and a
stimulation of microbial activity ('priming effect').
4. The gradual introduction of distinct bacterial communities hypothesized above (2) will result in increased CO₂
production, owing to the functional limitations of Yedoma permafrost microbial community in terms of CO₂
95 production.

2. Materials and Methods

2.1 Experimental design

2.1.1 Soils

The Yedoma sediment used to assess the impacts of collembola on permafrost CO₂ production and bacterial community
100 composition originated from the Cold Region Research Engineering Laboratory (CRREL) Permafrost Tunnel (Fox, Alaska,
USA). The sediment was sampled from the upper silt unit and is an upper Pleistocene silty deposit, previously described in
details (Shur et al., 2004; Mackelprang et al., 2011, 2017; Monteux et al., 2020). This sediment was chosen due to its
microbial communities being vulnerable to invasions and exhibiting functional limitations, allowing to discern impacts of
introduced microorganisms on broad proxies such as CO₂ production (Monteux et al., 2020). Approximately 35 g (fresh
105 weight) homogenized sediment was set in 200ml glass jars, sealed with parafilm to allow for gas but not moisture or
microorganism exchange, and pre-incubated at 10 °C for 11 days before inoculation.

To manipulate the collembola microbiome and make it more similar to that found in natural settings, we collected a topsoil (0-15 cm depth) from a subarctic meadow (Kärkevage, 30 km west of Abisko, northern Sweden, 68°24'23.8"N, 18°18'51.6"E) in September 2019 and kept it frozen until the cultivation of collembola. We could not obtain topsoil from the same location as the Yedoma permafrost due to practical constraints during the Covid-19 pandemic.

2.1.2 Collembola

A strain of the collembola species *Folsomia candida* was obtained from Vrije Universiteit Amsterdam and cultured 6 months prior to the onset of the experiment. *Folsomia candida* is a parthenogenetic ground-dwelling collembola, which has been routinely used as a model organism in soil ecology. Stock cultures were maintained on a gypsum and coal medium and fed baker's yeast, traces of mould were removed and fresh yeast and water were added once to twice a week, and fresh stock cultures were started monthly.

2.1.3 Collembola inoculation treatments

Two months prior to the onset of the experiment, separate stock cultures were established on gypsum and coal medium supplemented with a 2-3 cm layer of topsoil, to obtain collembola which both skin and gut microbiome were colonized with topsoil microorganisms ('topsoil stock culture'). In parallel, similar stock cultures using permafrost sediment were established as an additional control ('permafrost stock culture'). These stock cultures with soil or sediment were supplemented with yeast and water alike the gypsum stock culture, to maintain high adult population densities prior to the experiment. One day before inoculating the incubation jars, soil suspensions were made from the topsoil and permafrost sediment (5 and 10 g, respectively, in 100 ml ddH₂O), shaken at 150 rpm for 1 hour and filtered (Ahlstrom-Munksjö grade 006, 1.5 µm pore size) to manipulate the skin microbiome of collembola in the ectozoochory treatments.

All incubation jars containing permafrost were randomly assigned a treatment on the day of inoculation among the following, replicating each treatment across 6 jars (Figure 1). The treatments consisted in adding to the jars with permafrost:

- No collembola (hereafter, 'No-collembola control');
- Collembola grown on gypsum stock culture ('Collembola');
- Collembola grown on gypsum stock culture and sprayed with topsoil suspension ('Ectozoochory');
- Collembola grown on gypsum stock culture and sprayed with permafrost suspension ('Ectozoochory control');
- Collembola grown on topsoil stock culture ('Endozoochory');
- Collembola grown on permafrost stock culture ('Endozoochory control').




Figure 1: Summary of the experimental design. *Folsomia candida* collembola were grown in a stock culture on gypsum and fed with yeast, then subjected to different treatments before being inoculated into jars filled with Yedoma permafrost sediment (30-80 individuals per jar). Ectozoochory was assessed by spraying the collembola with soil suspensions, while endozoochory was assessed by growing collembola in stock cultures supplemented with soil. The jars were incubated in the dark for 25 days at 10 °C, CO₂ production was measured throughout the incubation period and bacterial community composition was determined at the end of the incubation.

135

To obtain a baseline allowing to quantify the effect of collembola on permafrost CO₂ production and bacterial community composition, Yedoma permafrost was incubated in absence of collembola ('no-collembola control'). For all other treatments, the permafrost was supplemented with adult collembola, isolated from the stock cultures using a hand-held vacuum cleaner mounted with a 10 ml pipet tip. The collembola were transferred into a black plastic tray, allowing to spread them and pick out the adults, excluding juveniles as much as possible. From this tray, a similar amount of collembola (30 – 80 individuals, in the range of values used in the literature e.g. by Kaneda and Kaneko (2008)) was sampled with the vacuum cleaner, then inoculated into each jar by pouring them into a plastic funnel, with the collembola provenance or manipulation depending on the treatment. Prior to transferring them into incubation jars, collembola used in the ectozoochory treatment and its control

140

were sprayed with the corresponding soil suspensions. To limit cross-contamination, separate funnels, as well as separate 10 ml pipet tips on the vacuum cleaner were used for the different treatments. A picture of the inside of each jar was taken to count the exact number of collembola, then the jars were closed with rubber septa, flushed with moisturized CO₂-free air and incubated. To estimate collembola respiration per individual, collembola were also incubated in 6 jars under identical conditions, except that the soil was replaced with a few drops of autoclaved ddH₂O to prevent dehydration (hereafter, no-soil calibration).

150 2.1.4. Incubation

We dark-incubated all flasks for 25 days under aerobic conditions at 10 °C. This incubation temperature is similar to summer active-layer temperatures in permafrost affected areas and within the thermal tolerance range of psychrophilic microorganisms (D'Amico et al., 2006). We used a short (25 days) incubation period to ensure a relatively stable collembola population level, by limiting uncertain numbers of newly-hatched collembola individuals, since the eggs of *Folsomia candida* take 18-20 days to hatch at 16 °C (Marshall & Kevan, 1962) and presumably longer than 25 days at 10 °C.

2.2. Measurements

2.2.1. CO₂ production

Headspace air was sampled with a syringe to measure CO₂ concentrations (EGM-5 IRGA, PP Systems, Amesbury, Massachusetts, USA) at intervals ensuring CO₂ concentrations remained below 20,000 ppm to prevent a toxic CO₂ build-up (i.e. after 3, 7, 14 and 25 days). After each measurement, the jars were flushed with 0.45 µm-filtered CO₂-free air moisturized by bubbling it through two 5l bottles of ddH₂O, for 3 min at 1 to 2 l min⁻¹, i.e. with at least 15 times the volume of the jar. CO₂ concentrations were adjusted for changes in temperature and atmospheric pressure to calculate CO₂ production rates (τ) as follows:

$$\tau_{(i)} = \frac{[CO_2]_i \times (P_i V / RT_i)}{(\Delta t)_i}$$

165 where (Δ_t)_i is the time interval between measurement (i) and previous flushing, P_i is atmospheric pressure at measurement time, V the headspace volume, R the ideal gas constant and T_i the temperature. To calculate cumulative CO₂ production over the entire incubation, we summed up the quantity of CO₂ present in the headspace at each sampling, within each jar.

2.2.2. Bacterial community

170 Microcentrifuge tubes (1.5 ml) were filled with soil and snap-frozen in dry ice to analyse microbial communities from the jars harvested after one day and at the end of the incubation. Initial soils were sampled likewise, after thawing and homogenizing but before pre-incubation for the permafrost soils and before preparing soil suspensions for active layer soils. The frozen tubes were kept at -20 °C for up to 4 months before freeze-drying, then homogenized by bead-beating (Precellys

CK-68 15 ml tubes, 2×30 s at 4500 rpm). DNA was extracted from 183 to 285 mg of homogenized freeze-dried soil using DNEasy PowerSoil Pro Kit (Qiagen) according to the manufacturer's instructions, and DNA concentrations in the extracts were measured on a Qubit 1.0 fluorometer.

The V4-V5 region of the 16S ribosomal RNA gene was targeted in PCR amplification using primers 515F (5' GTGYCAGCMGCCGCGGTAA 3') and 926R (5' CCGYCAATTYMTTTRAGTTT 3') with Illumina sequencing adapters, using 12.5 μ l Phusion Taq Green PCR mastermix (Thermo Scientific), 0.25 μ M of each primer, 2 μ l of DNA extract diluted to 5 ng μ l⁻¹ and nuclease-free water to 25 μ l reaction volume. PCR conditions were as follows: initial denaturation (98 °C, 3 min), 25 cycles of denaturation (98 °C, 15 s), annealing (50 °C, 30 s) and elongation (72 °C, 40 s), and a final elongation (72 °C, 10 min), after which PCR products were checked by electrophoresis on 1% agarose SB gel. 20 μ l of PCR products were cleaned and their DNA concentrations normalized using a SequalPrep Normalization Plate kit (Invitrogen), according to manufacturer's instructions. Three DNA extraction blanks and two PCR blanks were included as negative controls, as well as a mock community as positive control (ZymoBIOMICS, diluted to 5 and 0.5 ng μ l⁻¹ in two replicates each).

A second PCR step was performed to add Nextera dual-indexing barcodes, using 30 μ l reaction volume, 1 μ M of each primer and 5 μ l cleaned PCR product. PCR conditions were as follows: initial denaturation (98 °C, 3 min), 8 cycles of denaturation (98 °C, 30 s), annealing (55 °C, 30 s) and elongation (72 °C, 40 s), and a final elongation (72 °C, 10 min), after which PCR products were checked by electrophoresis on 1% agarose SB gel. 25 μ l of PCR products were cleaned and their DNA concentrations normalized using a SequalPrep Normalization Plate kit (Invitrogen). Serial elution across columns was used to increase concentration of the pooled products, i.e. using only 8×20 μ l elution buffer instead of 96×20 μ l. The eluted DNA was pooled, its concentration was measured on a Qubit fluorometer and the size distribution of the amplicons was measured by automated electrophoresis (Agilent 2100 Bioanalyzer). The library was then sent for sequencing on an Illumina MiSeq with V3 chemistry (2×300 bp, 15% PhiX spike-in) at the SNP&SEQ Technology Platform in Uppsala. Demultiplexing was performed by the sequencing facility, and data deposited at ENA with accession number PRJEB51992.

2.3. Data analysis

2.3.1. Bioinformatics

All bioinformatics and statistics were performed in R v4.1.3 (R Core Team, 2019), unless specified otherwise. The whole analysis pipeline is found at https://git.bolin.su.se/bolin/Collembola_vector and the processed data and figure-generating script at Zenodo (Monteux et al., 2022). In short, amplicon sequence variants (ASVs) were created with DADA2 1.18.0 (pseudo-pooling; Callahan et al., 2016) after removing primers and adapters with cutadapt (v3.10; Martin, 2011). Taxonomy was assigned to ASVs with the RDP naïve Bayesian classifier (v1.8; Wang et al., 2007), and ASVs resolved to the genus rank were further assigned a species rank by the exact string-matching algorithm implemented in DADA2 (*assignSpecies*),

205 using SILVA v138.1 reference data (Quast et al., 2013). Putative contaminant ASVs were manually selected from those identified *in silico* using the *decontam* algorithm (Davis et al., 2018) with combined prevalence- and frequency-based methods using the default threshold of 0.1, and separate prevalence- and frequency-based methods with a threshold of 0.05. Eight contaminant ASVs amounting up to 0.024% of the total reads were removed. Appropriateness of the bioinformatics analysis parameters was judged by visually assessing the composition of mock communities (microbial community DNA
210 standard, ZymoBIOMICS) at the genus level, leading to ASVs amounting up to less than 10 reads and/or present in fewer than 3 samples being removed from the dataset. Sequencing depth did not associate with experimental treatments (ANOVA $F_{5,30}=0.701$, $P=0.627$), therefore read numbers were converted to proportional abundances within samples to normalize sample sizes.

2.3.2. Diversity analyses

215 The effect of the different treatments on bacterial communities was visualized using principal coordinates analysis (PCoA) and tested with permutational multivariate analyses of variance (PerMANOVA, *adonis* function in *vegan* package) after verifying homoscedasticity (*betadisper*). Two distance matrixes were computed for that purpose, using weighted- and non-weighted UniFrac distances (Lozupone et al., 2011), to distinguish between compositional effects accounting for bacterial relative abundances and for only presence-absence, respectively. Pairwise contrasts were subsequently computed using the
220 wrapper provided in the *pairwiseAdonis* R package (Arbizu, 2021).

In addition, the same effect was tested with the more robust and sensitive *manyGLM* approach (Wang et al., 2012), which avoids certain pitfalls of distance-based methods (Warton et al., 2012). ManyGLM were fitted with negative binomial distribution, after visually checking that assumptions were met, on the non-normalized ASV count data using the default PIT-trap resampling with 1999 bootstrap permutations, and a likelihood-ratio testing method. An analysis of deviance was
225 carried out (*anova.manyglm*), subsequently using the provided wrapper for pairwise comparisons with free step-down P value adjustment.

2.3.3. CO₂ production analysis

A repeated-measures ANOVA was carried out, using Greenhouse-Geisser epsilon correction to degrees of freedom to account for the violation of the sphericity assumption, to assess the interactive effect of the collembola manipulation
230 treatments over time. Since this interactive effect was not statistically significant ($P=0.56$, Appendix A1), we removed the time dimension and used cumulative CO₂ production at the end of the incubation period below.

To assess our hypothesis that collembola presence would increase CO₂ production we used two-sample t-tests with unequal variances to compare the cumulative CO₂ production at the end of the incubation between the no-collembola control and all other jars.

235 To explore differences between the collembola microbiome manipulations, we used a one-way ANOVA followed by treatment contrasts to assess the difference from the control (i.e. all treatment and controls compared to the no-collembola

control) and selected orthogonal contrasts to assess the effects of each treatment (i.e. each treatment compared to its own control).

2.3.4. Estimated collembola respiration

240 Based on photographs taken upon inoculating the incubation jars with collembola and three days after inoculating, we counted the exact number of live collembola (i.e. ignoring collembola which had apparently not moved between the two pictures were taken). Only a few animals did not survive the transfer, and no dead animals were observed at the end of the incubation. The amount of collembola per jar varied across treatments ($F_{4,25} = 3.19$, $P = 0.030$), although no pairs of treated jars significantly differed from each other (when including the no-soil calibration jars, $F_{5,30} = 4.048$, $P = 0.006$; Appendix B),
245 we therefore needed to account for differing collembola numbers. We averaged the respiration per individual in the no-soil calibration set for each of the CO_2 concentration measurement times, and multiplied this amount by the number of collembola present in each jar to estimate the amount of CO_2 produced by collembola basal respiration in treatment jars.

2.3.5. Response-ratio calculations

A response-ratio of CO_2 production in collembola treatments was calculated by dividing cumulative CO_2 production by the average of the no-collembola control. This was done both for the net CO_2 production after subtracting the estimated collembola basal respiration (RR_{soil}), and for the gross CO_2 production (RR_{gross}). This allows to partition the difference in altered CO_2 production between what was respired by the collembola and a putative priming effect on soil CO_2 production.

3. Results

3.1. Bacterial communities




Figure 2: Response of permafrost bacterial communities to the addition of collembola undergoing different microbiome manipulation. a: Principal coordinates analysis (PCoA) of normalized abundance of bacterial ASVs (Amplicon Sequence Variants) based on weighted UniFrac phylogenetic distance (i.e. accounting for relative abundances); **b:** PCoA of normalized abundance of bacterial ASVs based on non-weighted UniFrac phylogenetic distance (i.e. presence-absence only). Ellipses indicate 80% confidence interval around treatment centroids, and axes are scaled to the percentage of explained variance. The asterisk indicates that the No-collembola control (dark brown) in **b** significantly differs from all other treatments (pairwise PerMANOVA $p < 0.05$) and collembola treatments did not differ from the stock-collembola treatment. Pairwise comparisons are summarized in Appendix C1.

255

260

265

Bacterial community composition was overall largely unaffected by the presence of collembola. Taking relative abundances into account, no effect of the different treatments was identified with PERMANOVA ($P > 0.05$, Figure 2a), but a significant treatment effect was observed using the presence-absence distance metrics (PERMANOVA $F_{5,30}=2.91$, $P = 0.001$, Figure 2b). More specifically, treatments containing collembola differed from the no-collembola control (Figure 2b) and to some extent from each other (Appendix C1). Using manyGLM, a significant treatment effect was observed (manyGLM analysis of deviance $P < 0.001$; Table 1). The ectozoochory treatment and its control (collembola sprayed with topsoil or permafrost soil suspension, respectively) differed significantly from the no-collembola control in the post-hoc test ($0.01 < P < 0.05$), while the other treatments including collembola tended to differ from the no-collembola control ($0.05 < P < 0.1$, Table 1). Similarly to the presence-absence analysis, treatments that included collembola did not significantly differ from each other in the manyGLM analysis. Overall, soil with collembola exhibited higher alpha-diversity metrics (richness estimators Chao1, ACE, number of observed ASVs, as well as Shannon and Fisher diversity indexes, Figure 3, Appendix C2) than the no-collembola control but those did not differ among the collembola treatments.

Table 1: Effect of addition of collembola on Yedoma sediment bacterial communities.
 Analysis of deviance test on a manyGLM model fitted on 1206 ASVs assuming negative-binomial distribution, using 1999 iterations of PIT-trap resampling.
P values below 0.05 are denoted in bold; ***: $P < 0.001$; *: $0.01 < P < 0.05$; : $0.05 < P < 0.1$.

			<i>Multivariate test</i>	Res.Df	Df.diff	Dev	<i>P</i>	
			Treatment	30	5	11742	<0.001	***
<i>Multivariate test contrasts</i>					Observed statistic	Adjusted <i>P</i>		
	Control	vs Collembola			3124	0.082	.	
	Control	vs Ectozoochory control			4085	0.013	*	
	Control	vs Ectozoochory			4274	0.011	*	
	Control	vs Endozoochory control			3276	0.064	.	
	Control	vs Endozoochory			3117	0.082	.	
	Collembola	vs Ectozoochory control			1533	0.669		
	Collembola	vs Ectozoochory			1580	0.669		
	Collembola	vs Endozoochory control			1842	0.658		
	Collembola	vs Endozoochory			1720	0.669		
	Ectozoochory control	vs Ectozoochory			1755	0.669		
	Ectozoochory control	vs Endozoochory control			2335	0.378		
	Ectozoochory control	vs Endozoochory			1752	0.669		
	Ectozoochory	vs Endozoochory control			2375	0.378		
	Ectozoochory	vs Endozoochory			2031	0.577		
	Endozoochory control	vs Endozoochory			1976	0.586		

270




Figure 3: Response of permafrost bacterial alpha diversity to the addition of collembola undergoing different microbiome manipulation. a: Number of observed ASVs (Amplicon Sequence Variants) per sample; **b:** Abundance-based Coverage Estimator ACE. Coloured symbols and error-bars are means and standard-errors within a treatment (n=6), small grey symbols are individual samples. Error-bars are only shown when exceeding the symbols size. Asterisks denote that the no-collembola control differs from all other treatments (estimated marginal means pairwise comparison, $0.001 < \text{Holm-adjusted } P < 0.01$).

3.2. CO₂ production

Overall, the collembola addition resulted in higher CO₂ production than the no-collembola control, but the ectozoochory and endozoochory treatments did not result in higher CO₂ production than their respective controls (using gross CO₂ production, as in RR_{gross}, Appendixes A2, A3). When accounting for collembola basal respiration (as in RR_{soil}), only the ectozoochory treatment (p=0.005) and to a lesser extent its control (p=0.069) differed from the no-collembola control (Appendixes A2, A3).

We hypothesized an increased CO₂ production in the zoochory treatments through an effect on bacterial communities, however that effect was absent, we therefore tested for the overall effect of collembola presence across all treatments compared to the no-collembola control (Figure 4). Using a t-test with unequal variances, we observed a 25.85% increase in CO₂ production in presence of collembola (RR_{gross}; Figure 4a). When subtracting estimated collembola basal respiration, this increase was on average 13.22%, (RR_{soil}; Figure 4b), thus roughly half of the observed increase in CO₂ production could be attributed to collembola basal respiration.




Figure 4: Response-ratios of permafrost cumulative CO₂ production rates to addition of collembola. a: RR_{gross}: measured CO₂ production rates; **b:** RR_{soil}: measured CO₂ production rates with estimated collembola basal respiration subtracted. All values are divided by the mean of the no-collembola control, small grey circles are individual values, large symbols are means, error-bars are standard errors of the mean (n=6 and 30 for brown and black symbols, respectively), and violin plots denote the shape of the distributions with maximum width proportional to the number of observations. Statistics are derived from two-sample t-tests with unequal variances.

285

4. Discussion

We showed that collembola stimulated CO₂ production from permafrost sediment through ‘priming’, while subtly altering the presence of certain bacterial ASVs, irrespective of the microbiome manipulation imposed on the collembola.

Colonization of permafrost by collembola therefore has the potential to result in bacterial community changes and to
290 increase C turnover from tundra soils.

4.1. Introduction of bacteria by collembola

Collembola introduced new bacteria in thawed permafrost soil, resulting in an increase in alpha-diversity and significant
differences in bacterial community composition, on a presence-absence basis. This pattern was consistent in all treatments
where collembola were added, but did not depend on the microbiome manipulation treatment assigned to the collembola
295 (Figure 2b, Figure 3). Despite their effects on the presence of certain bacteria, collembola did not impose broader community
changes, as bacterial communities were not affected when considering relative abundances as we had hypothesized (Figure
2a, Table 1). However, given the subtle effect of collembola on bacteria during this short incubation, this suggests that on
longer time scales collembola may modify bacterial communities more dramatically.

4.2. Differences between ecto- and endo-zoochory treatments

300 Although collembola introduced new bacteria into permafrost soil, our results did not support that they did so differently
through ecto- and/or endo-zoochory. This suggests that collembola presence mostly introduced bacteria that were part of the
core collembola microbiome, or that the way in which their microbiome changes was not well represented by our treatments.
For example, spraying may not have altered the cuticle microbiome as collembola cuticle is particularly hydrophobic
(Hensel et al., 2013a, b; Nickerl et al., 2013). This should however not have been an issue for the Endozoochory treatment,
305 in which collembola were in contact with the topsoil for a prolonged (>1 month) period of time. Despite this prolonged
exposure, no additional effect of Endozoochory on bacterial community composition was observed, compared to
Ectozoochory or Collembola-only treatments. Collembola typically harbor a gut microbiome reflecting their diet and direct
environment (Xiang et al., 2019; Leo et al., 2021), and observing soil-coloured feces in the topsoil and permafrost stock
cultures suggests that soil bacteria were present in the gut of collembola used in the Endozoochory treatments. In contrast
310 with the permafrost stock cultures, where the yeast was necessary as an additional food source to maintain stable
populations, collembola populations remained stable in topsoil stock cultures similar to those used in the Endozoochory
treatment even in absence of yeast, further indicating their consumption of soil particles and associated microorganisms.
Therefore, the absence of effect of the Ectozoochory treatment on permafrost bacterial communities may indicate that
collembola cuticle does not serve as a vector for bacterial dispersal, possibly due to its omniphobic structure. Further, the
315 lack of difference between the effects of Collembola and Endozoochory treatments suggests that few to none of the bacteria
present in this topsoil are able to establish in- or survive transit through collembolan gut. This suggests against generalist
'hitch-hiking' bacteria using microfauna guts as a means of dispersal, emphasizing instead that such hitch-hiking bacteria are
similar regardless of the feeding context of the collembola.

320 It is not unlikely that the observed limited response of bacterial community to zoochory treatments was transient, and that
collembola presence would have eventually affected community composition also in terms of relative abundances. A longer
incubation period may have resulted in stronger effects on bacterial community composition, as observed by Coulibaly et al.
(2019), but this would have been at the expense of controlling the number of collembola in the jars, and thus of being able to
325 way to detect such transient effects. It should also be noted that our semi-quantitative approach to bacterial community
composition did not allow us to assess putative changes in absolute abundances, although such changes are not necessarily
observed in similar studies (Kaneda and Kaneko, 2008).

Collembola effects on fungal communities may have indirectly affected bacterial communities, and thus represent a potential
330 additional mechanism by which collembola have affected the presence of certain bacteria. It is not unlikely that fungi would
have responded more strongly to the presence of mostly fungivorous *Folsomia candida*. Using fungi as focus organism in
this study, however, may have had its difficulties. For example, the use of *Candida albicans* as food source for the
collembola stock cultures may have strongly distorted fungal community composition, thus requiring questionable
bioinformatic workarounds. Further, assessing the effects on fungal communities may not allow to distinguish between the
335 consequences of collembola as a vector for microbial migration and as a grazing consumer. Although assessing responses of
the whole soil food web may have given better insights in potential mechanisms by which collembolan affect thawed
permafrost soil, we here chose to focus on their effects on bacteria given their preponderant role on realizing soil functions in
this system (Monteux et al., 2020).

4.3. Collembola effects on CO₂ production

340 Our hypothesis stating that soil CO₂ production would be increased in presence of collembola was supported by our results.
However, since no differences in bacterial community composition was detected between the different collembola
treatments, the rationale for our follow-up hypothesis (attributing an increased influence on CO₂ production to the treatments
exhibiting stronger changes in bacterial community) was not met. We therefore did not deem relevant to formally evaluate
this hypothesis. Our results suggest that it would be rejected (Appendix A3), but we could not say whether that is because
345 the rationale is wrong or because the manipulation of bacterial communities did not yield the expected effect.

Overall, collembola presence increased respiration by 25%, half of which was attributed to respiration of collembola
themselves. We interpret the remaining half (13.22%) as a “priming effect”, i.e. a stimulation of SOM-derived CO₂
production induced by the presence of collembola. The mechanism behind this priming effect remain unclear, and could for
350 instance relate to stimulation of microbial activity by the input of more labile substrates by the collembola (digestion by-
products). Overall, our results add to a list of contrasting findings on the direction and magnitude of priming of SOM
decomposition by collembola presence in soils in general. Earlier studies that compared CO₂ production in presence or

absence of collembola or other faunal groups have mostly focused on litter (e.g. Hanlon and Anderson, 1979). The scarce studies on soil CO₂ production including collembola reported contrasted findings, with some studies observing increases in soil CO₂ emissions in presence of collembola by up to +400% (Addison and Parkinson, 1978; Bakonyi, 1989; Kaneda and Kaneko, 2008; Wang et al., 2017), while others found no significant changes (Theenhaus et al., 1999; Lubbers et al., 2020; Lucas et al., 2020). Collembola effect on CO₂ production in litter can be species-specific (Hanlon and Anderson, 1979), and our results could support this as we found a similar increase as in three studies where *Folsomia candida* was used (Bakonyi, 1989; Kaneda and Kaneko, 2008; Wang et al., 2017). Kaneda and Kaneko (2008) reported a density-dependent increase in CO₂ production with the addition of *F. candida*, which showed no effect at the density used in our study but an increase with 5-10 times higher density (i.e. 400 individuals in 30g soil compared to our 30-80). Accordingly, we did not find any trend supporting a density-dependent response in our data. Nevertheless, we suggest that permafrost soils colonized by collembola in particular, and soil fauna in general, will likely see an increase in their rates of decomposition, and CO₂ production in particular, due to both priming and the basal respiration of the newly-established food web.

Our experimental setup did not allow us to distinguish further between soil and collembola respiration. We estimated basal respiration, though this may not perfectly reflect reality, considering that collembola in jars without soil may not have behaved similarly to those in jars with soil. Faced with a scarcity of food sources, they may have slowed their metabolism to endure that stress, which could result in decreased basal respiration, thus leading us to overestimate the magnitude of “priming”. We could not find reference values in the literature for *Folsomia candida* respiration rates, but Addison & Parkinson (1978) estimated the CO₂ production of two high-Arctic collembolan species to 182.6 and 250.6 µl CO₂ per g live weight per hour at 10 °C, for *Hypogastrura tullbergi* and *Folsomia regularis*, respectively. Assuming a CO₂ density of 1.977 kg / m³ and 170 µg per adult individual of *Folsomia candida*, our values range between 47 and 220 µl CO₂ per live weight per hour. Addison and Parkinson (1978) refer to Mitchell (1973) for a description of the method used for estimating respiration rates, where the measurements appear to be carried out over a period of 20 hours. When considering only the first measurement date in our data (3 days after inoculation), the rates are closer to the values reported by Addison and Parkinson (1978) with 157.5 ± 18.5 µl CO₂ per live weight per hour (mean ± SE, n=6). In future studies of this mechanism, it would be interesting to use stable isotope methods to partition the collembola- vs soil organic matter-derived CO₂ production more accurately.

Beyond the CO₂ production increase due to priming by soil fauna, our collembola addition treatments may have induced a “microbial community priming”, in the form of the release of microbial functional limitations. The Yedoma sediment used in this study indeed lacks certain microbial taxa and functions, and the onset of an exotic microbial community can result in large increases in CO₂ production (Monteux et al., 2020). Considering the limited to non-significant (depending on the distance metrics) effect of collembola addition on bacterial communities, we do not think this mechanism explains our observed results, although we do not rule out this explanation.

5. Conclusions

Collembola presence modified bacterial communities in newly-thawed permafrost and primed its CO₂ production, highlighting the importance of non-microbial decomposers for the fate of soil organic matter in thawing permafrost. An emerging theme in permafrost research is the missing functions in permafrost microbial communities due to a lack of certain microbial groups, hampering the production of CO₂, CH₄ and N₂O (Knoblauch et al., 2018; Monteux et al., 2020; Marushchak et al., 2021). It is also becoming clearer that the functionality of permafrost microbes can vary across space (Barbato et al., 2022), therefore the modalities of microbial dispersal into newly-thawed permafrost likely affect the fate of its organic matter and the rate of release of greenhouse gases. Our findings suggest that collembola may accelerate the release of greenhouse gases, possibly in part through the introduction of microorganisms, although those did not seem to depend strongly on the preceding collembola environment. In nature, thawing permafrost mostly occurs at the bottom of the seasonally-thawing active layer, often below the water-table and at depths far from the topsoil where most collembola reside. Different soil fauna, such as earthworms, may access such newly-thawed permafrost but their current geographical range does not overlap much with the permafrost region. However, rapid thaw events such as hillslope thermokarst, active layer detachments, or retrogressive thaw slumps expose newly-thawed permafrost to surface conditions, thus making it susceptible to colonization by collembola. Fauna-induced dispersal of microorganisms is most likely to take place in these environments, and should be further investigated in realistic field conditions.

It remains unclear whether the increase in CO₂ production we attribute to priming stems from increased functionality of the microbial community due to bridging of functional limitations, or from other mechanisms. Mechanical breakdown of organic material and digestion processes might provide microorganisms with more easily available substrates, while disruption of the soil pore structure may also result in increased microbial decomposition activity. Several studies indicate increases in CO₂ production with the presence of soil fauna, however to our best knowledge no studies have specifically partitioned to which extent such increases were due to faunal respiration or to priming effects. We argue that elucidating this question would be an important next step towards opening the ‘black box’ that soil systems still often represent, thus helping to mechanistically address the effects of global changes.

6. Appendices

Appendix A1: Effect of collembola addition on daily CO₂ production over time, repeated measures ANOVA.

P [GG] indicates Repeated Measures-ANOVA *P* using Greenhouse-Geisser correction on degrees of freedom to account for violation of the assumption of sphericity (Mauchly's test).

P values below 0.05 are denoted in bold; ***: *P* < 0.001; *: 0.01 < *P* < 0.05.

<i>Repeated Measures ANOVA</i>	DFnum	DFden	F	<i>P</i>	ges	
Treatment	5	30	3.499	0.013	0.259	*
Date	3	90	65.263	<0.001	0.465	***
Treatment:Date	15	90	0.854	0.616	0.054	

<i>Mauchly's sphericity test</i>	W	<i>P</i>	
Date	0.165	<0.001	***
Treatment:Date	0.165	<0.001	***

<i>Sphericity corrections</i>	GGe	p[GG]	
Date	0.544	<0.001	***
Treatment:Date	0.544	0.562	

Appendix A2: Effects of collembola additions on response ratios of cumulative CO₂ production at the end of the incubation, excluding (RR_{soil}) or including (RR_{gross}) estimated collembola respiration. Holm adjustment for *P* values of non-orthogonal contrasts.

P values below 0.05 are denoted in bold; ***: $P < 0.001$; **: $0.001 < P < 0.01$; *: $0.01 < P < 0.05$; .: $0.05 < P < 0.1$.

RR _{soil}	ANOVA	Df	Df.Res	Sum Sq	Mean Sq	F value	<i>P</i>
	Treatment		5	30	0.003	0.001	3.728


<i>Contrast vs no-collembola Control</i>	Estimate	SE	Df	t.ratio	Adjusted <i>P</i>
Collembola	0.012	0.008	30	1.573	0.379
Ectozoochory control	0.020	0.008	30	2.525	0.068 .
Ectozoochory	0.028	0.008	30	3.626	0.005 **
Endozoochory control	0.003	0.008	30	0.443	0.764
Endozoochory	0.007	0.008	30	0.887	0.764

<i>Orthogonal contrasts</i>	Estimate	Std. Error	t value	<i>P</i>
Collembola vs Control	0.006	0.004	1.573	0.126
Ectozoochory vs Ectozoochory control	0.002	0.004	0.444	0.660
Endozoochory vs Endozoochory Control	0.004	0.004	1.101	0.280

RR _{gross}	ANOVA	Df	Df.Res	Sum Sq	Mean Sq	F value	<i>P</i>
	Treatment		5	30	0.005	0.001	4.184


<i>Contrast vs no-collembola Control</i>	Estimate	SE	Df	t.ratio	Adjusted <i>P</i>
Collembola	0.024	0.009	30	2.754	0.030 *
Ectozoochory control	0.030	0.009	30	3.362	0.009 **
Ectozoochory	0.038	0.009	30	4.331	0.001 ***
Endozoochory control	0.020	0.009	30	2.270	0.031 *
Endozoochory	0.023	0.009	30	2.629	0.030 *

<i>Orthogonal contrasts</i>	Estimate	Std. Error	t value	<i>P</i>
Collembola vs Control	0.012	0.004	2.754	0.010 **
Ectozoochory vs Ectozoochory control	0.002	0.004	0.359	0.722
Endozoochory vs Endozoochory Control	0.004	0.004	0.969	0.340



Appendix A3: Response-ratios of permafrost cumulative CO₂ production rates to addition of collembola.
a: RR_{gross}: measured CO₂ production rates; **b:** RR_{soil}: measured CO₂ production rates with estimated collembola basal respiration subtracted. All values are divided by the mean of the no-collembola control, small grey circles are individual jar values, large symbols are means, error-bars are standard errors of the mean (n=6). Black symbols denote significant difference from the control, different symbols denote different statistical significance (EMmeans treatment contrasts with Holm adjustment for multiple comparisons, Supplementary Table 2, .: 0.05 < P < 0.1; *: 0.01 < P < 0.05; **: 0.001 < P < 0.01; ***: P < 0.001).

415



Appendix B: Number of adult collembola individuals per incubation jar. Small grey circles are individual jar values, large symbols are means, error-bars are standard errors of the mean (n=6), and violin plots denote the shape of the distributions with maximum width proportional to the number of observations. Different black letters denote statistically significant pairwise differences (EMmeans pairwise contrasts with Holm adjustment for multiple comparisons, 0.05 < P < 0.1).

Appendix C1: Effects of collembola additions on bacterial community composition (permutational multivariate ANOVAs and pairwise contrasts), using distance matrixes based on relative abundances (weighed UniFrac) or presence-absence (non-weighted UniFrac). Holm adjustment for *P* values of pairwise contrasts.

P values below 0.05 are denoted in bold; ***: $P < 0.001$; **: $0.001 < P < 0.01$; *: $0.01 < P < 0.05$; .: $0.05 < P < 0.1$.

<i>PerMANOVA</i>	Df	Df.Res	F.Model	R2	P
Weighted UniFrac (relative abundance)	5	30	1.261	0.174	0.101
Non-weighted UniFrac (presence-absence)	5	30	3.018	0.335	< 0.001

<i>Pairwise contrast</i>	Df	F.Model	R2	Adjusted P	
Control vs Collembola	1	3.236	0.244	0.025	*
Control vs Endozoochory control	1	4.356	0.303	0.025	*
Control vs Ectozoochory	1	6.777	0.404	0.022	*
Control vs Ectozoochory control	1	5.658	0.361	0.025	*
Control vs Endozoochory	1	4.777	0.323	0.012	*
Collembola vs Endozoochory control	1	1.767	0.150	0.102	
Collembola vs Ectozoochory	1	1.751	0.149	0.115	
Collembola vs Ectozoochory control	1	1.348	0.119	0.148	
Collembola vs Endozoochory	1	1.434	0.125	0.148	
Endozoochory control vs Ectozoochory	1	3.068	0.235	0.025	*
Endozoochory control vs Ectozoochory control	1	2.711	0.213	0.025	*
Endozoochory control vs Endozoochory	1	2.156	0.177	0.025	*
Ectozoochory vs Ectozoochory control	1	1.497	0.130	0.113	
Ectozoochory vs Endozoochory	1	1.958	0.164	0.025	*
Ectozoochory control vs Endozoochory	1	1.820	0.154	0.025	*

Appendix C2: Effects of collembola additions on bacterial alpha diversity (ANOVAs and pairwise contrasts) , with total number of observed ASVs, Abundance-based Coverage Estimator (ACE), Shannon's and Fisher's diversity indexes. Holm adjustment for *P* values of pairwise contrasts.

P values below 0.05 are denoted in bold; ***: $P < 0.001$; **: $0.001 < P < 0.01$; *: $0.01 < P < 0.05$; .: $0.05 < P < 0.1$.

<i>ANOVA</i>	Df	Df.Res	Sum Sq	Mean Sq	F value	<i>P</i>	
Observed ASVs	5	30	94389	18878	9.313	<0.001	***
ACE	5	30	98265	19653	8.517	<0.001	***
Shannon	5	30	0.582	0.116	4.667	0.003	**
Fisher	5	30	2141	428	12.644	<0.001	***

<i>Treatment contrast (vs Control)</i>		Estimate	SE	df	t.ratio	<i>P</i>	
Observed ASVs	Collembola	113.500	25.994	30	4.366	0.001	**
	Ectozoochory control	141.667	25.994	30	5.450	<0.001	***
	Ectozoochory	162.333	25.994	30	6.245	<0.001	***
	Endozoochory control	106.000	25.994	30	4.078	0.001	**
	Endozoochory	106.333	25.994	30	4.091	0.001	**
ACE	Collembola	115.2	27.7	30	4.155	0.001	**
	Ectozoochory control	145.4	27.7	30	5.245	<0.001	***
	Ectozoochory	165.2	27.7	30	5.956	<0.001	***
	Endozoochory control	106.0	27.7	30	3.821	0.003	**
	Endozoochory	106.5	27.7	30	3.840	0.003	**
Shannon	Collembola	0.230	0.091	30	2.521	0.070	.
	Ectozoochory control	0.326	0.091	30	3.571	0.006	**
	Ectozoochory	0.378	0.091	30	4.148	0.001	**
	Endozoochory control	0.198	0.091	30	2.171	0.144	
	Endozoochory	0.347	0.091	30	3.800	0.003	**
Fisher	Collembola	16.806	3.360	30	5.002	<0.001	***
	Ectozoochory control	21.731	3.360	30	6.468	<0.001	***
	Ectozoochory	24.006	3.360	30	7.145	<0.001	***
	Endozoochory control	14.598	3.360	30	4.345	0.001	**
	Endozoochory	17.301	3.360	30	5.149	<0.001	***

420 **7. Code availability**

All code used to process the raw DNA data and produce the figures and tables presented in text is found at https://git.bolin.su.se/bolin/Collembola_vector

The code used to generate figures from the processed DNA data is also found at <https://dx.doi.org/10.5281/zenodo.6461323>

8. Data availability

425 All 16S sequencing data is found at ENA under project accession number PRJEB51992.

All processed data used to generate the findings presented in text are found at <https://dx.doi.org/10.5281/zenodo.6461323> and https://git.bolin.su.se/bolin/Collembola_vector

9. Sample availability

430 Due to practical constraints, the exact permafrost material and collembola cultures used throughout the manuscript are no longer available. Permafrost from the same location may be obtained through contacting the USA Army CRREL, while *Folsomia candida* strains may be obtained by contacting JM. Frozen permafrost samples from the end of the incubation period, as well as aliquots of DNA extracts used for this study may be obtained from the corresponding author upon reasonable request.

10. Author contributions

435 SM, JM and EK designed the experiment. JM provided *Folsomia candida* strains and guidance for culturing. EK provided topsoil from Kärkevagge. SM performed the experiment, data analysis, and wrote the manuscript with input from all co-authors.

11. Competing interests

The authors declare that they have no conflict of interest.

440 **12. Acknowledgements**

This study was funded by a grant from Formas (Dnr 2017-01182) awarded to E.J.K. and Kempestiftelserna awarded to S.M. We thank the Department of Forest Mycology and Plant Pathology, SLU, for hosting the molecular work, T. H. Douglas from the US Army Cold Regions Research and Engineering Laboratory's Permafrost Tunnel (Alaska) for assistance and

permission to sample. We also thank F. Keuper from the French National Research Institute for Agriculture, Food and
445 Environment and E. Dorrepaal from Umeå University for providing the permafrost sample.

Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The facility is part of the National Genomics
Infrastructure (NGI) Sweden and Science for Life Laboratory. The SNP&SEQ Platform is also supported by the Swedish
Research Council and the Knut and Alice Wallenberg Foundation.

13. References

450 Addison, J. A. and Parkinson, D.: Influence of Collembolan Feeding Activities on Soil Metabolism at a High Arctic Site,
Oikos, 30, 529–538, <https://doi.org/10.2307/3543348>, 1978.

Agamennone, V., Jakupovic, D., Weedon, J. T., Suring, W. J., van Straalen, N. M., Roelofs, D., and Roling, W. F. M.: The
microbiome of *Folsomia candida*: an assessment of bacterial diversity in a *Wolbachia*-containing animal, *FEMS Microbiol*
Ecol, 91, UNSP fiv128, <https://doi.org/10.1093/femsec/fiv128>, 2015.

455 Arbizu, P. M.: pairwiseAdonis, R, 2021.

Bakonyi, G.: Effects of *Folsomia candida* (Collembola) on the microbial biomass in a grassland soil, *Biol Fert Soils*, 7, 138–
141, <https://doi.org/10.1007/BF00292572>, 1989.

Barbato, R. A., Jones, R. M., Douglas, T. A., Doherty, S. J., Messan, K., Foley, K. L., Perkins, E. J., Thurston, A. K., and
Garcia-Reyero, N.: Not all permafrost microbiomes are created equal: Influence of permafrost thaw on the soil microbiome
460 in a laboratory incubation study, *Soil Biology and Biochemistry*, 108605, <https://doi.org/10.1016/j.soilbio.2022.108605>,
2022.

Berman, D. I., Meshcheryakova, E. N., and Mikhajlova, E. V.: Cold Hardiness and Range of the Myriapod *Angarozonium*
Amurense (Polyzoniidae, Diplopoda, Arthropoda) in Permafrost Environments, *Cryoletters*, 36, 237–242, 2015.

Bottos, E. M., Kennedy, D. W., Romero, E. B., Fansler, S. J., Brown, J. M., Bramer, L. M., Chu, R. K., Tfaily, M. M.,
465 Jansson, J. K., and Stegen, J. C.: Dispersal limitation and thermodynamic constraints govern spatial structure of permafrost
microbial communities, *FEMS Microbiol Ecol*, 94, fiy110, <https://doi.org/10.1093/femsec/fiy110>, 2018.

Buse, T., Ruess, L., and Filser, J.: Collembola gut passage shapes microbial communities in faecal pellets but not viability of
dietary algal cells, *Chemoecology*, 24, 79–84, <https://doi.org/10.1007/s00049-013-0145-y>, 2014.

Calderón, K., Spor, A., Breuil, M.-C., Bru, D., Bizouard, F., Violle, C., Barnard, R. L., and Philippot, L.: Effectiveness of
470 ecological rescue for altered soil microbial communities and functions, *The ISME Journal*, 11, 272–283,
<https://doi.org/10.1038/ismej.2016.86>, 2017.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2: High-resolution
sample inference from Illumina amplicon data, *Nat Methods*, 13, 581–583, <https://doi.org/10.1038/nmeth.3869>, 2016.

Coulibaly, S. F. M., Winck, B. R., Akpa-Vinceslas, M., Mignot, L., Legras, M., Forey, E., and Chauvat, M.: Functional
475 Assemblages of Collembola Determine Soil Microbial Communities and Associated Functions, *Front. Environ. Sci.*, 7,
<https://doi.org/10.3389/fenvs.2019.00052>, 2019.

- D'Amico, S., Collins, T., Marx, J.-C., Feller, G., Gerday, C., and Gerday, C.: Psychrophilic microorganisms: challenges for life, *EMBO reports*, 7, 385–389, <https://doi.org/10.1038/sj.embor.7400662>, 2006.
- 480 Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., and Callahan, B. J.: Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data, *Microbiome*, 6, 226, <https://doi.org/10.1186/s40168-018-0605-2>, 2018.
- Frouz, J., Novotná, K., Čermáková, L., and Pivokonský, M.: Soil fauna reduce soil respiration by supporting N leaching from litter, *Applied Soil Ecology*, 153, 103585, <https://doi.org/10.1016/j.apsoil.2020.103585>, 2020.
- 485 García-Palacios, P., Maestre, F. T., Kattge, J., and Wall, D. H.: Climate and litter quality differently modulate the effects of soil fauna on litter decomposition across biomes, *Ecology Letters*, 16, 1045–1053, <https://doi.org/10.1111/ele.12137>, 2013.
- Golovatch, S. and Kime, R.: Millipede (Diplopoda) distributions: a review, *Soil Organisms*, 81, 2009.
- Griffiths, H. M., Ashton, L. A., Parr, C. L., and Eggleton, P.: The impact of invertebrate decomposers on plants and soil, *New Phytologist*, 231, 2142–2149, <https://doi.org/10.1111/nph.17553>, 2021.
- 490 Hanlon, R. D. G. and Anderson, J. M.: The effects of collembola grazing on microbial activity in decomposing leaf litter, *Oecologia*, 38, 93–99, <https://doi.org/10.1007/BF00347827>, 1979.
- Harding, T., Jungblut, A. D., Lovejoy, C., and Vincent, W. F.: Microbes in High Arctic Snow and Implications for the Cold Biosphere, *Appl. Environ. Microbiol.*, 77, 3234–3243, <https://doi.org/10.1128/AEM.02611-10>, 2011.
- Hensel, R., Helbig, R., Aland, S., Voigt, A., Neinhuis, C., and Werner, C.: Tunable nano-replication to explore the omniphobic characteristics of springtail skin, *NPG Asia Mater*, 5, e37–e37, <https://doi.org/10.1038/am.2012.66>, 2013a.
- 495 Hensel, R., Helbig, R., Aland, S., Braun, H.-G., Voigt, A., Neinhuis, C., and Werner, C.: Wetting Resistance at Its Topographical Limit: The Benefit of Mushroom and Serif T Structures, *Langmuir*, 29, 1100–1112, <https://doi.org/10.1021/la304179b>, 2013b.
- Hultman, J., Waldrop, M. P., Mackelprang, R., David, M. M., McFarland, J., Blazewicz, S. J., Harden, J., Turetsky, M. R., McGuire, A. D., Shah, M. B., VerBerkmoes, N. C., Lee, L. H., Mavrommatis, K., and Jansson, J. K.: Multi-omics of permafrost, active layer and thermokarst bog soil microbiomes, *Nature*, 521, 208–212, <https://doi.org/10.1038/nature14238>, 2015.
- 500 Inglese, C. N., Christiansen, C. T., Lamhonwah, D., Moniz, K., Montross, S. N., Lamoureux, S., Lafrenière, M., Grogan, P., and Walker, V. K.: Examination of Soil Microbial Communities After Permafrost Thaw Subsequent to an Active Layer Detachment in the High Arctic, Arctic, Antarctic, and Alpine Research, 49, 455–472, <https://doi.org/10.1657/AAAR0016-066>, 2017.
- 505 Johnston, E. R., Hatt, J. K., He, Z., Wu, L., Guo, X., Luo, Y., Schuur, E. A. G., Tiedje, J. M., Zhou, J., and Konstantinidis, K. T.: Responses of tundra soil microbial communities to half a decade of experimental warming at two critical depths, *PNAS*, 116, 15096–15105, <https://doi.org/10.1073/pnas.1901307116>, 2019.
- Kaneda, S. and Kaneko, N.: Collembolans feeding on soil affect carbon and nitrogen mineralization by their influence on microbial and nematode activities, *Biol Fertil Soils*, 44, 435–442, <https://doi.org/10.1007/s00374-007-0222-x>, 2008.
- 510 Knoblauch, C., Beer, C., Liebner, S., Grigoriev, M. N., and Pfeiffer, E.-M.: Methane production as key to the greenhouse gas budget of thawing permafrost, *Nature Climate Change*, 8, 309–312, <https://doi.org/10.1038/s41558-018-0095-z>, 2018.

- 515 Leo, C., Nardi, F., Cucini, C., Frati, F., Convey, P., Weedon, J. T., Roelofs, D., and Carapelli, A.: Evidence for strong environmental control on bacterial microbiomes of Antarctic springtails, *Sci Rep*, 11, 2973, <https://doi.org/10.1038/s41598-021-82379-x>, 2021.
- Lozupone, C., Lladser, M. E., Knights, D., Stombaugh, J., and Knight, R.: UniFrac: an effective distance metric for microbial community comparison, *ISME J*, 5, 169–172, <https://doi.org/10.1038/ismej.2010.133>, 2011.
- 520 Lubbers, I. M., Berg, M. P., De Deyn, G. B., van der Putten, W. H., and van Groenigen, J. W.: Soil fauna diversity increases CO₂ but suppresses N₂O emissions from soil, *Global Change Biology*, 26, 1886–1898, <https://doi.org/10.1111/gcb.14860>, 2020.
- Lucas, J. M., McBride, S. G., and Strickland, M. S.: Trophic level mediates soil microbial community composition and function, *Soil Biology and Biochemistry*, 143, 107756, <https://doi.org/10.1016/j.soilbio.2020.107756>, 2020.
- 525 Mackelprang, R., Waldrop, M. P., DeAngelis, K. M., David, M. M., Chavarria, K. L., Blazewicz, S. J., Rubin, E. M., and Jansson, J. K.: Metagenomic analysis of a permafrost microbial community reveals a rapid response to thaw, *Nature*, 480, 368–371, <https://doi.org/10.1038/nature10576>, 2011.
- Mackelprang, R., Burkert, A., Haw, M., Mahendrarajah, T., Conaway, C. H., Douglas, T. A., and Waldrop, M. P.: Microbial survival strategies in ancient permafrost: insights from metagenomics, *The ISME Journal*, 11, 2305–2318, <https://doi.org/10.1038/ismej.2017.93>, 2017.
- 530 Martin, M.: Cutadapt removes adapter sequences from high-throughput sequencing reads, *EMBnet.journal*, 17, 10–12, <https://doi.org/10.14806/ej.17.1.200>, 2011.
- Marushchak, M. E., Kerttula, J., Diáková, K., Faguet, A., Gil, J., Grosse, G., Knoblauch, C., Lashchinskiy, N., Martikainen, P. J., Morgenstern, A., Nykamb, M., Ronkainen, J. G., Siljanen, H. M. P., van Delden, L., Voigt, C., Zimov, N., Zimov, S., and Biasi, C.: Thawing Yedoma permafrost is a neglected nitrous oxide source, *Nat Commun*, 12, 7107, <https://doi.org/10.1038/s41467-021-27386-2>, 2021.
- 535 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, Ö., Yu, R., and Zhou, B. (Eds.): *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2021.
- 540 Mitchell, M. J.: An improved method for microrespirometry using gas chromatography, *Soil Biology and Biochemistry*, 5, 271–274, [https://doi.org/10.1016/0038-0717\(73\)90011-4](https://doi.org/10.1016/0038-0717(73)90011-4), 1973.
- Monteux, S., Weedon, J. T., Blume-Werry, G., Gavazov, K., Jassey, V. E. J., Johansson, M., Keuper, F., Olid, C., and Dorrepaal, E.: Long-term in situ permafrost thaw effects on bacterial communities and potential aerobic respiration, *The ISME Journal*, 12, 2129–2141, <https://doi.org/10.1038/s41396-018-0176-z>, 2018.
- 545 Monteux, S., Keuper, F., Fontaine, S., Gavazov, K., Hallin, S., Juhanson, J., Krab, E. J., Revaillet, S., Verbruggen, E., Walz, J., Weedon, J. T., and Dorrepaal, E.: Carbon and nitrogen cycling in Yedoma permafrost controlled by microbial functional limitations, *Nature Geoscience*, 13, 794–798, <https://doi.org/10.1038/s41561-020-00662-4>, 2020.
- Nickerl, J., Helbig, R., Schulz, H.-J., Werner, C., and Neinhuis, C.: Diversity and potential correlations to the function of Collembola cuticle structures, *Zoomorphology*, 132, 183–195, <https://doi.org/10.1007/s00435-012-0181-0>, 2013.

- 550 Potapov, A., Bellini, B. C., Chown, S. L., Deharveng, L., Janssens, F., Kováč, L., Kuznetsova, N., Ponge, J.-F., Potapov, M., Querner, P., Russell, D., Sun, X., Zhang, F., and Berg, M. P.: Towards a global synthesis of Collembola knowledge: challenges and potential solutions, *soil organisms*, 92, 161–188, <https://doi.org/10.25674/so92iss3pp161>, 2020.
- Quast, C., Priesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic Acids Res*, 41, D590–596, <https://doi.org/10.1093/nar/gks1219>, 2013.
- 555 R Core Team: R: A Language and Environment for Statistical Computing, 2019.
- Sfenthourakis, S. and Hornung, E.: Isopod distribution and climate change, *Zookeys*, 25–61, <https://doi.org/10.3897/zookeys.801.23533>, 2018.
- 560 Shatilovich, A. V., Tchesunov, A. V., Neretina, T. V., Grabarnik, I. P., Gubin, S. V., Vishnivetskaya, T. A., Onstott, T. C., and Rivkina, E. M.: Viable Nematodes from Late Pleistocene Permafrost of the Kolyma River Lowland, *Dokl Biol Sci*, 480, 100–102, <https://doi.org/10.1134/S0012496618030079>, 2018.
- Shmakova, L., Malavin, S., Iakovenko, N., Vishnivetskaya, T., Shain, D., Plewka, M., and Rivkina, E.: A living bdelloid rotifer from 24,000-year-old Arctic permafrost, *Current Biology*, 31, R712–R713, <https://doi.org/10.1016/j.cub.2021.04.077>, 2021.
- 565 Shur, Y., French, H. M., Bray, M. T., and Anderson, D. A.: Syngenetic permafrost growth: cryostratigraphic observations from the CRREL tunnel near Fairbanks, Alaska, *Permafrost and Periglacial Processes*, 15, 339–347, <https://doi.org/10.1002/ppp.486>, 2004.
- Sørensen, L. I., Holmstrup, M., Maraldo, K., Christensen, S., and Christensen, B.: Soil fauna communities and microbial respiration in high Arctic tundra soils at Zackenberg, Northeast Greenland, *Polar Biol*, 29, 189–195, <https://doi.org/10.1007/s00300-005-0038-9>, 2006.
- 570 Strauss, J., Schirrmeister, L., Grosse, G., Fortier, D., Hugelius, G., Knoblauch, C., Romanovsky, V., Schädel, C., Schneider von Deimling, T., Schuur, E. A. G., Shmelev, D., Ulrich, M., and Veremeeva, A.: Deep Yedoma permafrost: A synthesis of depositional characteristics and carbon vulnerability, *Earth-Science Reviews*, 172, 75–86, <https://doi.org/10.1016/j.earscirev.2017.07.007>, 2017.
- 575 Theenhaus, A., Scheu, S., and Schaefer, M.: Contramensal interactions between two collembolan species: effects on population development and on soil processes, *Functional Ecology*, 13, 238–246, <https://doi.org/10.1046/j.1365-2435.1999.00303.x>, 1999.
- Väisänen, M., Krab, E. J., Monteux, S., Teuber, L. M., Gavazov, K., Weedon, J. T., Keuper, F., and Dorrepaal, E.: Meshes in mesocosms control solute and biota exchange in soils: A step towards disentangling (a)biotic impacts on the fate of thawing permafrost, *Applied Soil Ecology*, 151, 103537, <https://doi.org/10.1016/j.apsoil.2020.103537>, 2020.
- 580 Wang, M., Zhang, W., Xia, H., Huang, J., Wu, Z., and Xu, G.: Effect of Collembola on mineralization of litter and soil organic matter, *Biol Fertil Soils*, 53, 563–571, <https://doi.org/10.1007/s00374-017-1200-6>, 2017.
- Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R.: Naive Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy, *Applied and Environmental Microbiology*, 73, 5261–5267, <https://doi.org/10.1128/AEM.00062-07>, 2007.

- 585 Wang, Y., Naumann, U., Wright, S. T., and Warton, D. I.: mvabund– an R package for model-based analysis of multivariate abundance data, *Methods in Ecology and Evolution*, 3, 471–474, <https://doi.org/10.1111/j.2041-210X.2012.00190.x>, 2012.
- Warton, D. I., Wright, S. T., and Wang, Y.: Distance-based multivariate analyses confound location and dispersion effects, *Methods in Ecology and Evolution*, 3, 89–101, <https://doi.org/10.1111/j.2041-210X.2011.00127.x>, 2012.
- 590 Xiang, Q., Zhu, D., Chen, Q.-L., Delgado-Baquerizo, M., Su, J.-Q., Qiao, M., Yang, X.-R., and Zhu, Y.-G.: Effects of diet on gut microbiota of soil collembolans, *Science of The Total Environment*, 676, 197–205, <https://doi.org/10.1016/j.scitotenv.2019.04.104>, 2019.
- Yashina, S., Gubin, S., Maksimovich, S., Yashina, A., Gakhova, E., and Gilichinsky, D.: Regeneration of whole fertile plants from 30,000-y-old fruit tissue buried in Siberian permafrost, *PNAS*, 109, 4008–4013, <https://doi.org/10.1073/pnas.1118386109>, 2012.

595