



Large contribution of soil N₂O emission to the global warming potential of a large-scale oil palm plantation despite changing from conventional to reduced management practices

Guantao Chen¹, Edzo Veldkamp¹, Muhammad Damris², Bambang Irawan³, Aiyen Tjoa⁴, Marife D. Corre¹

5 ¹Soil Science of Tropical and Subtropical Ecosystems, Faculty of Forest Sciences and Forest Ecology, University of Goettingen, Göttingen 37077, Germany

²Faculty of Science and Technology, University of Jambi, Jl. Raya Jambi-Ma. Bulian km. 15, Mendalo Darat, Muaro, Jambi 36361, Indonesia

³Forestry Faculty, University of Jambi, Campus Pinang Masak Mendalo, Jambi 36361, Indonesia

10 ⁴Faculty of Agriculture, Tadulako University, Jl. Soekarno Hatta, km 09 Tondo, Palu 94118, Indonesia

Correspondence to: Guantao Chen (gchen1@gwdg.de)

Abstract. Conventional management of oil palm plantations, involving high fertilization rate and herbicide application, result in high yield but with large soil greenhouse gas (GHG) emissions. This study aimed to assess a practical alternative to conventional management, namely reduced fertilization with mechanical weeding, to decrease soil GHG emissions without sacrificing production. We established a full factorial experiment with two fertilization rates (conventional and reduced fertilization, equal to nutrients exported via fruit harvest) and two weeding methods (herbicide and mechanical), each with four replicate plots, since 2016 in a ≥ 15 -year old, large-scale oil palm plantation in Indonesia. Soil CO₂, N₂O, and CH₄ fluxes were measured during 2019 – 2020 and yield was measured during 2017 – 2020. Fresh fruit yield (30 ± 1 Mg ha⁻¹ yr⁻¹) and soil GHG fluxes did not differ among treatments ($P \geq 0.11$), implying legacy effects of over a decade of conventional management prior to the start of experiment. Annual soil GHG fluxes were 5.5 ± 0.2 Mg CO₂-C ha⁻¹ yr⁻¹, 3.6 ± 0.7 kg N₂O-N ha⁻¹ yr⁻¹, and -1.5 ± 0.1 kg CH₄-C ha⁻¹ yr⁻¹ across treatments. The palm circle, where fertilizers are commonly applied, covered 18% of the plantation area but accounted 79% of soil N₂O emission. The net primary production of this oil palm plantation was 17150 ± 260 kg C ha⁻¹ yr⁻¹ but 62% of this was removed by fruit harvest. The global warming potential of this plantation was 3010 ± 750 kg CO₂-eq ha⁻¹ yr⁻¹ of which 55% was contributed by soil N₂O emission and only < 2% offset by soil CH₄ sink.

25 1 Introduction

With increasing demand for vegetable oil, oil palm as a productive woody oil crop is widely planted in the tropics (Descals et al., 2021). Globally, the oil palm-planted area rapidly increased from 4 million ha in 1980 to 28 million ha in 2019 (FAO, 2021), and oil palm plantations are expected to continue to expand to meet the increasing demand of a growing world population (OECD, 2022). Oil palm expansion drives tropical deforestation (Vijay et al., 2016) and is accompanied by serious reductions in multiple ecosystem functions, e.g. decreases in C storage (Kotowska et al., 2015; van Straaten et al., 2015), nutrient cycling and retention (Allen et al., 2015; Kurniawan et al., 2018), and biodiversity losses (Clough et al., 2016). Despite these losses of ecosystem multifunctionality, profit gains increase under oil palm plantations (Grass et al., 2020), which increase farm and non-farm households' income and improve their livelihood (Bou Dib et al., 2018). Moreover, the high yield of oil palm plays an invaluable role in meeting human demand for vegetable oil (Thomas et al., 2015; Rochmyaningsih, 2019). However, there



35 is a need for a balance between economic gains and maintaining or avoiding further degradation of ecosystem functions (Bessou
et al., 2017). Conventional management practices of large-scale oil palm plantations (> 50 ha planted area, can be up to 20000
ha and owned by corporations), particularly high fertilization rates and herbicide application, are agents of these decreases in
ecosystem functions (Tao et al., 2016; Ashton-Butt et al., 2018; Rahman et al., 2019). Thus, there is a need for practical
40 solutions that can easily be implemented in order to reduce these negative impacts on ecosystem functions without sacrificing
productivity and profit.

Soil greenhouse gas (GHG) emissions are a concern in oil palm plantations (Kaupper et al., 2020; Skiba et al., 2020). Compared
to forests, the reductions in plant biomass production, soil organic carbon (SOC) and soil microbial biomass as well as C
removal from the field via fruit harvest and increase in soil bulk density in oil palm plantations largely decrease the latter's
GHG abatement capability (Kotowska et al., 2015; van Straaten et al., 2015; Clough et al., 2016). This capability is influenced
45 by agricultural management practices, especially fertilization rates (Sakata et al., 2015; Hassler et al., 2017; Rahman et al.,
2019). N fertilization in oil palm plantations during the wet season can increase soil N₂O emissions (Aini et al., 2015; Hassler
et al., 2017), a potent GHG and agent of ozone depletion (Davidson et al., 2000). Soil N₂O emissions from tropical agriculture
is largely controlled by soil mineral N availability, which in turn is influenced by N fertilization rate, and soil moisture, as N₂O-
production processes of denitrification and nitrification in the soil are favored under high soil mineral N and moisture levels
50 (Khalil et al., 2002; Liyanage et al., 2020; Quiñones et al., 2022). In Indonesia, a large-scale oil palm plantation with commonly
high N fertilization rate has higher soil N₂O emissions than smallholder oil palm plantations (< 5 ha of planted area per
household) with low N fertilization rates (Hassler et al., 2017).

Soil CO₂ emissions originate from heterotrophic and root respiration (Bond-Lamberty et al., 2004). The temporal pattern of
soil CO₂ emission follows the seasonal dynamics of soil moisture and/or soil temperature, exhibiting low soil CO₂ emissions
55 at low soil moisture content, increases toward an optimum soil moisture, and decrease towards water saturation when oxygen
availability and gas diffusion limit soil CO₂ emissions (Sotta et al., 2007; van Straaten et al., 2011). The spatial pattern of soil
CO₂ emissions in tropical ecosystems is influenced by the spatial variation in soil organic matter, bulk density, root biomass
and available N and P levels in the soil (Adachi et al., 2006; Hassler et al., 2015; Cusack et al., 2019; Tchiofo Lontsi et al.,
2020). Another important GHG that is influenced by management practices in tropical plantations or croplands is CH₄ (Hassler
60 et al., 2015; Quiñones et al., 2022). Soil surface CH₄ flux is the net effect of CH₄ production by methanogenic archaea and CH₄
oxidation by methanotrophic bacteria (Hanson and Hanson, 1996). In well-drained tropical soils, CH₄ oxidation usually is more
dominant than CH₄ production, resulting in a net soil CH₄ uptake or negative CH₄ flux (Veldkamp et al., 2013). Seasonal
pattern of soil surface CH₄ flux in smallholder oil palm plantations in Indonesia reflects the seasonal variation of soil moisture,
with lower CH₄ uptake during the wet season than the dry season (Hassler et al., 2015). In well-drained tropical soils with low
65 N availability, soil CH₄ uptake or methanotrophic activity is enhanced with increase in soil mineral N content (Veldkamp et
al., 2013; Hassler et al., 2015; Tchiofo Lontsi et al., 2020). However, in tropical agricultural soils with high soil mineral N
levels (NH₄⁺ and NO₃⁻) from high N fertilization, competition of NH₄⁺ against CH₄ for the active site of mono-oxygenase
enzyme can reduce soil CH₄ uptake (Hanson and Hanson, 1996). Veldkamp et al. (2001) observed a temporary inhibition of
CH₄ uptake for approximately three weeks following NH₄⁺-based fertilizer application in tropical pasture soils. In contrast,
70 high soil NO₃⁻ level (e.g. resulting from nitrification of applied N fertilizer) can inhibit CH₄ production in the soil since NO₃⁻
is preferred over bicarbonate as an electron acceptor (Martinson et al., 2021; Quiñones et al., 2022). Nonetheless, across forests,



smallholder rubber and oil palm plantations in Indonesia, the overriding pattern is the increase in soil CH₄ uptake with increase in soil mineral N, suggesting the prevailing control of N availability on methanotrophic activity in the soil (Hassler et al., 2015). The spatial patterns of soil surface CH₄ fluxes depict the spatial variations in soil properties that affects soil moisture content and gas diffusivity, such as soil texture (Veldkamp et al., 2013; Tchifo Lontsi et al., 2020), soil bulk density and organic matter (e.g. at a plot or landscape scale; Tchifo Lontsi et al., 2020).

In large-scale oil palm plantations, the typical management practices of fertilization, weeding, and pruning of senesced fronds result in three distinctive spatial management zones: palm circle (weeded and fertilizers are applied), inter-row (weeded but not fertilized) and frond-stacked area (where pruned fronds are piled) (Fig. 1; Formaglio et al., 2021). In the palm circle and inter-row, frequent management activities (weeding, pruning and harvesting) result in soil compaction by foot traffic (increased soil bulk density) and the low litter input in these zones exhibits low SOC and microbial biomass, and low soil N cycling rate (Formaglio et al., 2021). Additionally, root biomass is high in the palm circle (Dassou et al., 2021). In the frond-stacked area, decomposition of fronds results in large SOC (with decreased soil bulk density), large microbial and fine root biomass, and high soil N cycling rate (Moradi et al., 2014; Rüegg et al., 2019; Formaglio et al., 2020; Dassou et al., 2021). Overall, the differences in soil properties and root biomass among these spatially distinct management zones (Formaglio et al., 2021) potentially drive the spatial variation of soil GHG fluxes from oil palm plantations (Hassler et al., 2015, 2017; Aini et al., 2020). Thus, estimating soil GHG emissions from oil palm plantations should take into account the spatial variability among management zones within a site or plot.

This study aimed to (1) assess differences in soil GHG fluxes from conventional high fertilization rates with herbicide application compared to alternative management of reduced fertilization rates (equal to nutrient exported via fruit harvest) with mechanical weeding; and (2) determine the controlling factors of the soil GHG fluxes from a large-scale oil palm plantation. A 2 × 2 factorial field experiment with conventional and reduced fertilization rates as well as herbicide and mechanical weed control was established in a ≥ 15-year old, large-scale oil palm plantation in Jambi, Indonesia starting in November 2016. The earlier studies during the first 1.5 years of this oil palm management experiment show comparable gross rates of soil N cycling, microbial biomass (Formaglio et al., 2021), and root biomass (Ryadin et al., 2022) among treatments. Thus, we hypothesized that during 2.5–3.5 years of this management experiment, the reduced fertilization with mechanical weeding will have comparable soil CO₂ and CH₄ fluxes but lower soil N₂O emissions than the conventional fertilization with herbicide weeding. Moreover, we hypothesized that the fertilized palm circle that has high soil bulk density and root biomass but low SOC, soil microbial biomass and N cycling rate (Dassou et al., 2021; Formaglio et al., 2021) will have large soil CO₂ and N₂O emissions but small soil CH₄ uptake. The unfertilized inter-row that has high soil bulk density but low SOC, microbial biomass and N cycling rate (Formaglio et al., 2021) will have small soil CO₂, N₂O emissions and CH₄ uptake. The frond-stacked area (i.e. unfertilized but piled with pruned fronds) that has large SOC, microbial biomass and N cycling rate but low soil bulk density (Formaglio et al., 2021) will have large soil CO₂ emissions and CH₄ uptake but small soil N₂O emissions. In this study, we assessed the soil GHG footprint and the global warming potential (GWP) of a typical large-scale oil palm plantation in order to evaluate reduced management (i.e. reduced fertilization rate with mechanical weeding) against the commonly employed conventional management (i.e. high fertilization rate with herbicide application).



2 Materials and methods

2.1 Site description and experimental design

This study was conducted in a large-scale oil palm plantation in Jambi, Indonesia ($1^{\circ}43'8''$ S, $103^{\circ}23'53''$ E, 73 m above sea level). The plantation was 2025 ha and established between 1998 and 2002, and thus was ≥ 18 years old during our measurement of soil GHG fluxes from July 2019 to June 2020. The oil palms were planted in a triangular pattern with 8 m spacing between palms and the planting density was 142 palms ha^{-1} . Mean annual (2010–2020) air temperature is 26.9 ± 0.2 °C and mean annual precipitation is 2078 ± 155 mm. The soil is Acrisol with a sandy clay loam texture (Table S1). More than 18 years of management induced three distinct zones within this oil palm plantation: palm circle, inter-row, and frond-stacked area (Fig. 1) (Formaglio et al., 2020). The palm circle (a 2 m radius from the palm base) is the zone where fertilizers and lime are applied (in April and October of each year) and is weeded every three months; this represents 18% of the plantation area. The inter-row is unfertilized but weeded every six months; this represents 67% of the plantation area. The frond-stacked area is where senesced fronds are piled and is neither fertilized nor weeded; this represents 15% of the plantation area. As consequences of these management activities, SOC and total N stocks are higher whereas soil bulk density is lower in the frond-stacked area than the palm circle and inter-row. The effective cation exchange capacity and pH, influenced by the applied lime as well as from decomposed leaf litter, are higher in the palm circle and frond-stacked area than the inter-row (Table S1) (Formaglio et al., 2020).

This oil palm management experiment had started in November 2016 – a 2×2 factorial design of two fertilization rates and two weeding methods: conventional fertilization – herbicide weeding (ch), conventional fertilization – mechanical weeding (cw), reduced fertilization – herbicide weeding (rh), and reduced fertilization – mechanical weeding (rw). These four treatments were randomly assigned to four plots (50 m \times 50 m each) in a block and there were four replicate blocks (Fig. 1). Within each plot, we selected two subplots in the inner 30 m \times 30 m area and each subplot included the three management zones where all measurements were carried out (Fig. 1).

Conventional fertilization rates were 260 kg N, 50 kg P, and 220 kg K $\text{ha}^{-1} \text{yr}^{-1}$, commonly practiced in large-scale oil palm plantations in Jambi, Indonesia (Formaglio et al., 2020). Reduced fertilization rates were 136 kg N, 17 kg P, and 187 kg K $\text{ha}^{-1} \text{yr}^{-1}$, equal to the nutrients exported by fruit harvest (detail calculation given by Formaglio et al., 2021). The fertilizer sources were urea, triple superphosphate, muriate of potash or NPK-complete. For herbicide weed control, glyphosate was used at a rate of 1.5 L $\text{ha}^{-1} \text{yr}^{-1}$ in the palm circle (split into four applications yr^{-1}) and 0.75 L $\text{ha}^{-1} \text{yr}^{-1}$ in the inter-row (split into two applications yr^{-1}). Mechanical weeding used a brush cutter with the same weeding frequencies as the herbicide applications. All treatments received the same rates of lime (426 kg dolomite $\text{ha}^{-1} \text{yr}^{-1}$) and micronutrients (142 kg micro-mag $\text{ha}^{-1} \text{yr}^{-1}$ with 0.5% B_2O_3 , 0.5% CuO, 0.25% Fe_2O_3 , 0.15% ZnO, 0.1% MnO and 18% MgO), applied only in the palm circle.

2.2 Soil greenhouse gas fluxes

Soil CO_2 , N_2O , and CH_4 fluxes were measured monthly from July 2019 to June 2020, using vented static chambers. Measurement schedules were random among plots (i.e. 16 plots, each with two subplots that each encompassed three management zones) such that we covered the temporal variability of soil GHG fluxes (e.g. peak of soil N_2O emissions during two weeks following fertilization), as observed in our earlier study (Hassler et al., 2017). During the year-round measurements,



chamber bases (0.04 m² area) were installed permanently at each management zone (i.e. palm circle, inter-row, and frond-stacked area) in all subplots of 16 plots (Fig. 1), totaling to 96 chambers, by inserting these into the soil at approximately 0.02 m depth. On a measurement day, the chamber bases were covered for 28 minutes with polyethylene covers (11 L total volume) that were equipped with a Luer-lock sampling port. Four gas samples (23 mL each) were taken using syringes at 1, 10, 19, 28 minutes following chamber closure and injected into pre-evacuated 12 mL glass vials (Labco Exetainers, Labco Limited, Lampeter, UK) with rubber septa. On each monthly measurement, 384 gas samples were taken (i.e. 16 plots × 2 subplots × 3 management zones × 4 chamber headspace-sampling intervals). As a check for possible leakage, we also stored standard gases into pre-evacuated 12 mL glass vials in the same period as the field gas samples. All gas samples were transported to the Goettingen University, Germany for analysis.

Gas samples were analyzed using a gas chromatograph (SRI 8610C, SRI Instruments Europe GmbH, Bad Honnef, Germany) equipped with a flame ionization detector to measure CH₄ and CO₂ concentrations (with a methanizer) as well as an electron capture detector for N₂O analysis (with a make-up gas of 5% CO₂–95% N₂). Soil CO₂, N₂O, and CH₄ fluxes were calculated from the linear change in concentrations with chamber closure time, adjusted with the measured air temperature and atmospheric pressure during sampling. We found that the concentrations of all standard gases stored in the same duration as the field gas samples stayed the same as those in the standard gases at our laboratory. The quality check for each flux measurement was based on the linear increase of CO₂ concentrations with chamber closure time ($R^2 \geq 0.9$). For soil CH₄ and N₂O, all flux measurements (including zero and negative fluxes) were included in the data analysis. For an overall value of soil CO₂, N₂O, and CH₄ fluxes in a plot, fluxes were weighted by the areal coverages of the three management zones (see above). Area-weighted annual soil CO₂, N₂O, and CH₄ fluxes were estimated based on trapezoidal extrapolations between measured fluxes and sampling day intervals.

2.3 Soil variables

Concurrent with soil GHG flux measurement, soil temperature, mineral N, and moisture content in top 5 cm depth were determined. Soil temperature was recorded using a portable thermometer (Greisinger GMH 3210, Greisinger Messtechnik GmbH, Regenstauf, Germany). At about 1 m away from each chamber, soil samples were collected and pooled from the two subplots for each management zone. Part of each soil sample was added to a prepared bottle containing 150 mL 0.5 M K₂SO₄ for immediate mineral N extraction. Upon arrival at the field laboratory, the bottles were shaken for 1 hour, filtered and the extracts were immediately frozen. The remaining soil sample was oven-dried at 105 °C for 24 h to determine the gravimetric moisture content, which was used to calculate the dry mass of the soil extracted for mineral N. The moisture content was the expressed as water-filled pore space (WFPS), using the mineral soil particle density of 2.65 g cm⁻³ and the average soil bulk density in the top 5 cm (1.23 g cm⁻³ in the palm circle, 1.20 g cm⁻³ in the inter-row, and 0.52 g cm⁻³ in the frond-stacked area). In April and May 2020, we were unable to conduct WFPS and mineral N measurements due to restrictions from COVID-19 pandemic. The frozen extracts were transported by air to Goettingen University and analyzed for NH₄⁺ and NO₃⁻ concentrations using continuous flow injection colorimetry (SEAL Analytical AA3, SEAL Analytical GmbH, Norderstedt, Germany).

2.4 Global warming potential estimation

The GWP of this ≥ 18-year old, large-scale oil palm plantation was estimated based on Malhi et al. (1999), as also used in our



earlier work in agricultural land use (Quiñones et al., 2022). First, the net primary production (NPP) was estimated as the sum of aboveground biomass C production, fruit biomass C production, frond litter biomass C input, root biomass C production, and root litter biomass C input. Within the inner 30 m × 30 m area per plot, stem height, harvested fruit weight, and the number of pruned fronds per palm were recorded during 2017–2020 (Iddris et al., 2023). Aboveground biomass per palm was calculated using the allometric growth equation (Asari et al., 2013): $\text{kg biomass palm}^{-1} = 71.797 \times \text{palm stem height} - 7.0872$; and biomass production per palm was the difference in biomass between two consecutive years. Annual aboveground biomass C production ($\text{g C m}^{-2} \text{ yr}^{-1}$) = annual biomass production per palm (kg palm^{-1} , 2019–2020) × planting density (142 palms ha^{-1}) × tissue C concentration (0.41 g C g^{-1}) × 10^{-1} (for unit conversion). Annual fruit biomass C production ($\text{g C m}^{-2} \text{ yr}^{-1}$) = annual fruit harvest per palm (kg palm^{-1} , mean of 2019–2020) × planting density × tissue C concentration (0.63 g C g^{-1}) × 10^{-1} . Frond litter biomass C input ($\text{g C m}^{-2} \text{ yr}^{-1}$) = annual litter production per palm (kg palm^{-1} , mean of 2019–2020) × planting density × tissue C concentration (0.47 g C g^{-1}) × 10^{-1} . Data of root biomass C production ($140 \text{ g C m}^{-2} \text{ yr}^{-1}$) and root litter biomass C input ($45 \text{ g C m}^{-2} \text{ yr}^{-1}$) were taken from Kotowska et al. (2015).

Second, the net ecosystem productivity (NEP) was calculated as $\text{NEP (g C m}^{-2} \text{ yr}^{-1}) = \text{heterotrophic respiration} - (\text{NPP} - \text{fruit biomass C})$ (Malhi et al., 1999; Quiñones et al., 2022). Our measured soil CO_2 fluxes included both autotrophic and heterotrophic respirations. We assumed 70% heterotrophic contribution to soil CO_2 flux, based on a long-term quantification in a forest in Sulawesi, Indonesia (van Straaten et al., 2011). As the frond litter also contributes to heterotrophic respiration upon decomposition, we assumed this fraction to be 80% of frond litter biomass C, based on the frond-litter decomposition rate in the same plantation (Iddris et al., 2023). Third, the GWP ($\text{g CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$) = $(\text{NEP} \times 3.67) + (\text{soil N}_2\text{O fluxes} \times 298) + (\text{soil CH}_4 \text{ fluxes} \times 25)$, of which 3.67 is C-to- CO_2 conversion, and 298 and 25 are CO_2 -equivalents of N_2O and CH_4 , respectively, for a 100-year time horizon (IPCC, 2006). Negative and positive symbols indicate the direction of the flux: (–) for C uptake and (+) for C export or emission from the plantation.

2.5 Statistical analysis

The mean value of two subplots for the soil GHG fluxes and soil temperature were used to represent each plot and management zone on each sampling day. The normality of distribution and equality of variance were first tested using Shapiro-Wilk's test and Levene's test, respectively. Linear mixed-effects (LME) models with Tukey's HSD test were used to assess the differences in soil GHG fluxes and soil variables (WFPS, soil temperature, and mineral N content) among treatments (Crawley, 2013). In the LME models, management (2×2 factorial of fertilization rates, weed control and their interaction) was considered as the fixed effect whereas plot and sampling day were taken as random effects, and statistical analysis were conducted for each management zone. As there were no significant differences among treatments (Table 1), we tested differences among the three management zone across treatments; for the latter, management zone was the fixed effect in the LME model and plot and sampling day were random effects. We also assessed if there were seasonal differences, and the year-round measurements were categorized into dry (precipitation $\leq 80 \text{ mm month}^{-1}$) and wet seasons; this was conducted for each management zone, and season was the fixed effect in the LME model and plot and sampling day were random effects. For all the above analyses, the LME models further included a variance function that allows variance heteroscedasticity of the fixed effect, and/or a first-order temporal autoregressive process that assumes decreasing auto-correlation between sampling days with increasing time difference, if these improve the model performance based on Akaike information criterion. The model residual was checked



using diagnostic plots and finally soil CO₂ and N₂O fluxes, and soil NH₄⁺ and NO₃⁻ concentrations were re-analyzed after log-transformation as the model residual distributions approximated the normal distribution.

215 The relationships between soil GHG fluxes and soil variables were determined by Spearman's Rank correlation test. Correlations tests were conducted on the means of the four replicate plots on each measurement day for each management zone ($n = 144$ for soil temperature, from 4 treatments \times 3 management zones \times 12 monthly measurements; $n = 120$ for WFPS and mineral N content, from 4 treatments \times 3 management zones \times 10 monthly measurements). All data analyses were performed using the R version 4.0.5 (R core Team, 2021). The statistical significance for all the tests was set at $P \leq 0.05$.

220 3 Results

3.1 Soil greenhouse gas fluxes and global warming potential estimate

Soil CO₂ emissions from the palm circle and frond-stacked area were higher in the wet season than in the dry season ($P \leq 0.03$; Fig. S1). Soil N₂O emissions from the palm circle sharply increased after fertilizer application and returned to background levels after two months (Fig. S2). Excluding the direct effects after fertilization, the palm circle had higher soil N₂O emissions
225 in the wet season than the dry season ($P = 0.03$; Fig. S2). Soil CH₄ uptake was higher in the dry season than the wet season in all three management zones ($P \leq 0.01$; Fig. S3). The frond-stacked area showed consistent CH₄ uptake throughout the measurement period whereas 33% and 17% of measured soil CH₄ fluxes in the palm circle and inter-row, respectively, were net CH₄ emissions (Fig. S3).

Reduced and conventional management had comparable soil CO₂ emissions (fertilization, weeding and interaction: $P \geq 0.13$),
230 N₂O emissions (fertilization, weeding and their interaction: $P \geq 0.14$), and CH₄ fluxes (fertilization, weeding and their interaction: $P \geq 0.26$) in each management zone (Table 1). However, there were clear differences in soil GHG fluxes among the three management zones. The palm circle had the highest soil N₂O emissions and lowest soil CH₄ uptake ($P \leq 0.01$; Table 1); the inter-row had the lowest soil CO₂ and N₂O emissions ($P \leq 0.01$; Table 1); the frond-stacked area had the highest soil CO₂ emissions and CH₄ uptake ($P \leq 0.01$; Table 1). The palm circle accounted for 25%, the inter-row for 45%, and the frond-stacked area for 30% of the annual soil CO₂ emissions (Fig. 2). The palm circle comprised 79% of the annual soil N₂O emissions although it only accounted for 18% of the plantation area (Fig. 2). The frond-stacked area with 15% areal coverage contributed to 41% of the annual soil CH₄ uptake and the palm circle with 18% of plantation area contributed 5% of the annual soil CH₄ uptake (Fig. 2).

We calculated the GWP across 16 plots (Fig. 3) as there were no significant differences in soil GHG fluxes among treatments
240 (Table 1). Additionally, the reduced and conventional management had also comparable fruit yield during four years (2017–2020) of treatments (fertilization, weeding and their interaction: $P \geq 0.07$; Table S2). The NPP was larger than the soil heterotrophic respiration (which was assumed to be 70% of the measured soil respiration + 80% C emissions from decomposition of frond litter; see Methods), but 62% of this NPP was removed from the field via fruit harvest (Fig. 3). Thus, this oil palm plantation turned into a net C source (i.e. positive NEP value; Fig. 3). Summing the NEP, soil N₂O emissions, and
245 soil CH₄ uptake in terms of CO₂ equivalent (100-year time horizon; see Methods), the GWP of this ≥ 18 -year old, large-scale oil palm plantation was contributed by 55% soil N₂O emissions and only counterbalanced by $< 2\%$ soil CH₄ sink (Fig. 3).



3.2 Soil variables

Fertilization and weeding treatments did not affect soil temperature (fertilization, weeding and their interaction: $P \geq 0.21$) and WFPS (fertilization, weeding and interaction: $P \geq 0.24$) in each management zone (Table 2). Soil NO_3^- concentration was lower in reduced than conventional fertilization, particularly in the frond-stacked area ($P \leq 0.01$; Table 2). There was an interaction effect of fertilization and weeding on soil NH_4^+ concentration in the frond-stacked area ($P = 0.02$; Table 2); however, neither fertilization nor weeding solely affect soil NH_4^+ concentration in any of the management zones ($P \geq 0.08$; Table 2). Across treatment plots, the three management zones showed comparable soil temperature while the palm circle and inter-row had higher WFPS than the frond-stacked area ($P \leq 0.01$; Table 2). The soil NH_4^+ and NO_3^- concentrations in the palm circle and frond-stacked area were larger than the inter-row ($P \leq 0.01$; Table 2).

Soil CO_2 emissions were positively correlated with WFPS in the palm circle ($\rho = 0.37$, $P = 0.02$) and frond-stacked area ($\rho = 0.72$, $P \leq 0.01$; Fig. 4a) and were positively correlated with soil temperature in the frond-stacked area ($\rho = 0.60$, $P \leq 0.01$) and inter-row ($\rho = 0.29$, $P = 0.05$; Fig. 4b). Soil N_2O emissions were positively correlated with soil mineral N in the palm circle ($\rho = 0.58$, $P \leq 0.01$) and inter-row ($\rho = 0.32$, $P = 0.04$; Fig. 4f). Soil CH_4 uptake decreased with increase in WFPS in all three management zones ($\rho = 0.44\text{--}0.81$, $P \leq 0.01$; Fig. 4g). In the frond-stacked area, soil CH_4 uptake decreased with increase in soil temperature ($\rho = 0.54$, $P \leq 0.01$; Fig. 4h) but increased with increase in soil mineral N ($\rho = -0.40$, $P = 0.01$; Fig. 4i); also in the frond-stacked area, a positive relationship between soil temperature and WFPS was observed ($\rho = 0.37$, $P = 0.02$; Fig. 4j). We did not find any other significant correlations between soil GHG fluxes and the measured soil variables.

4 Discussion

4.1 Soil CO_2 emissions

Area-weighted soil CO_2 emissions (Table 1) were only about one-third of the soil CO_2 emissions from forests in the same study area ($187\text{--}196 \text{ mg C m}^{-2} \text{ h}^{-1}$; Hassler et al., 2015) but within the range reported for oil palm plantations on mineral soils in Southeast Asia ($45\text{--}195 \text{ mg C m}^{-2} \text{ h}^{-1}$; Hassler et al., 2015; Sakata et al., 2015; Aini et al., 2020; Drewer et al., 2021b). Specifically, soil CO_2 emissions from the inter-row were in the lower end of this range and soil CO_2 fluxes from the frond-stacked area were in the middle of this range. These earlier studies deployed different spatial sampling designs for measuring soil GHG fluxes from oil palm plantations. Sakata et al. (2015) measured soil CO_2 fluxes at 1 m away from the palm base and Hassler et al. (2015) measured soil CO_2 fluxes at 1.8–5 m away from the palm base. Aini et al. (2020) measured soil CO_2 fluxes from the fertilized area (within 1 m from the palm base) and unfertilized area whereas no information on sampling location was given by Drewer et al. (2021b). Measurement locations to represent spatial management zones should be stated when reporting soil GHG fluxes from oil palm plantations in order to facilitate comparisons as well as to warrant spatial extrapolation.

The three management zones differed in soil CO_2 fluxes caused by their differences in SOC (Table S1), microbial biomass (Fig. S4) as drivers of heterotrophic respiration, and root biomass (Nelson et al., 2014) that influences autotrophic respiration. In this mature large-scale oil palm plantation, senesced fronds have been piled on the frond-stacked area for more than a decade. This results in 40% larger SOC stocks (Table S1) and 3–5 times larger microbial biomass (Fig. S4) in the frond-stacked area than in the palm circle and inter-row (Formaglio et al., 2020, 2021). The positive correlation of microbial biomass C (MBC) to soil



CO₂ fluxes (Fig. S4) supported our second hypothesis whereby differences in soil CO₂ emissions among management zones are driven in part by microbial biomass size and available organic C for heterotrophic respiration. Substantial heterotrophic respiration as well as presence of roots in the frond-stacked area (Rüegg et al., 2019; Dassou et al., 2021) explained its highest soil CO₂ emissions (Table 1). On the other hand, the palm circle has higher root biomass than the inter-row (Nelson et al., 2014), and higher soil CO₂ emissions from the palm circle than the inter-row (Table 1) may be caused by their disparate autotrophic respiration as their soil SOC stocks (Table S1) and MBC did not differ (Formaglio et al., 2021). However, in the same oil palm plantation, the different fertilization and weeding treatments, analyzed across the three management zones, did not influence soil MBC (Formaglio et al., 2021) as well as the fine root biomass in the top 10 cm depth (Ryadin et al., 2022) after one year of this management experiment. These findings support our first hypothesis whereby there was no short-term differences between reduced and conventional managements on soil CO₂ emissions. Nonetheless, we emphasize that the lower soil respiration in oil palm plantations compared to the forests (Hassler et al., 2015) is supported by its decreases in SOC (van Straaten et al., 2015; Allen et al., 2016), root and litter production (Kotowska et al., 2015) and microbial biomass (Allen et al., 2015; Formaglio et al., 2021). Also, the higher soil ¹⁵N natural abundance and lower soil C:N ratio in oil palm plantations than the forests (Hassler et al., 2015; van Straaten et al., 2015) signify a highly decomposed organic matter which, combined with reduced SOC, suggest reduced available C for microbial biomass and heterotrophic activity (Allen et al., 2015; Formaglio et al., 2021).

The seasonal pattern of soil CO₂ emissions from land uses in Indonesia is commonly influenced by soil moisture (Hassler et al., 2015; van Straaten et al., 2011). In our present study, the positive correlation between soil CO₂ emissions and soil moisture (ranging from 10%–55% WFPS; Fig. 4a) depicted a reduced soil respiration during the dry season, particularly in the palm circle and frond-stacked area, suggesting diminished autotrophic and heterotrophic respiration in these management zones when soil moisture was low (Fig. 4a; Fig. S1). Previous studies show that autotrophic and heterotrophic respiration increase toward an optimum WFPS (e.g. WFPS between 50%–55%; Sotta et al., 2007; van Straaten et al., 2011). Unlike the previous study conducted in 2013 in the same area (Hassler et al., 2015), our measured WFPS did not reach beyond 55%, as the annual rainfall during our study year (2019–2020) was lower than in 2013 and the sandy clay loam texture of our present soil may facilitate well-drained conditions. Thus, we did not observe a parabolic relationship of soil CO₂ emissions with WFPS beyond 55% as observed by Hassler et al. (2015). Although we observed a positive relationship between soil CO₂ emissions and soil temperature (Fig. 4b), largely in the frond-stacked area, this maybe confounded by WFPS as the soil temperature and WFPS were auto-correlated (Fig. 4j). Thus, soil temperature was not a dominant controlling factor for the seasonal pattern of soil CO₂ emissions in this oil palm plantation where soil temperature also only varied narrowly during our measurement period (25–28 °C; Fig. 4b).

4.2 Soil N₂O emissions

Area-weighted soil N₂O emissions (Table 1) were within the range reported for oil palm plantations on mineral soils (8–117 μg N m⁻² h⁻¹; Aini et al., 2015; Sakata et al., 2015; Hassler et al., 2017; Rahman et al., 2019; Drewer et al., 2021b). Specifically, the soil N₂O emissions from the unfertilized inter-row and frond-stacked areas at our site were close to the lower end of this range whereas those from the fertilized palm circle were larger than the upper end of this range (Fig. S2). This pattern supported our second hypothesis, whereby soil N₂O emission was primarily influenced by soil N availability (i.e. mineral N; Table 2; Fig.



4f). These pulses of N₂O emissions from the palm circle peaked at around two weeks following N fertilization and went down to the background emissions after at most eight weeks (Fig. S2) (Aini et al., 2015; Hassler et al., 2017; Rahman et al., 2019). Although both the inter-row and frond-stacked areas had no direct N fertilizer application, litter decomposition in the frond-stacked area resulted in higher gross rates of N mineralization and nitrification, indicating higher soil N availability, than the inter-row (Formaglio et al., 2021). This explained the higher soil N₂O emissions from the frond-stacked area than the inter-row (Table 1). However, these internal soil-N cycling processes provide slow release of mineral N as opposed to the pulse release of mineral N level from N fertilization, and hence the soil N₂O emissions and mineral N levels were larger in the palm circle than the frond-stacked area (Tables 1 and 2). These findings signified the main control of soil N availability on soil N₂O emissions (Fig. 4f) (Davidson et al., 2000). We did not observe a correlation of soil N₂O emissions with WFPS (Fig. 4d) possibly because our sandy clay loam soil was relatively dry to moist in the top 5 cm depth ($\leq 55\%$ WFPS) during our measurement period. In sum, the direct N fertilizer application on the palm circle (although covering only 18% of the plantation area) caused the extremely high soil N₂O emissions (Fig. S2), accounting 79% of the annual soil N₂O emission at the plantation level (Fig. 2). Our findings highlight that the palm circle was a hot spot of soil N₂O emissions, and such management-induced spatial heterogeneity must be accounted for in accurately quantifying soil N₂O emissions from large-scale oil palm plantations.

The large and comparable soil N₂O emissions between the conventional and reduced fertilization treatments (Table 1; Fig. 2) were contrary to our first hypothesis. However, this finding was consistent with the soil N availability, i.e. mineral N (Table 2) as well as gross and net rates of soil N mineralization and nitrification, which did not differ among treatments during 1–4 years of this management experiment (Formaglio et al., 2021; unpublished data of net N mineralization and nitrification rates). These results implied a substantial legacy effect of the past decade of conventional management (high fertilization rate and herbicide application) prior to the start of this management experiment. It is important to note that our reduced fertilization treatment was still 1.5–3 times higher than the N fertilization rates in smallholder oil palm plantations, and these reduce fertilization treatment displayed 2–4 times larger soil N₂O emissions than the smallholder plantations (Hassler et al., 2017). Also, the soil mineral N levels in this large-scale oil palm plantation were larger in any of the management treatments (Table 2) compared to smallholder oil palm plantations (Hassler et al., 2017). In the reduced fertilization treatment, soil mineral N was possibly not the limiting factor for N₂O production, since the peaks of soil N₂O emissions following fertilization in both reduced and conventional fertilization treatments were comparable (Fig. S2). This supports the conclusion of Formaglio et al. (2020) that the decadal over-fertilization of this large-scale oil palm plantation causes large stocks of mineral N, leached below the root zone, and despite four years of reduced fertilization, mineral N stored at deeper depths can contribute to microbial production of N₂O.

These findings imply the need to adjust fertilization rates with age of oil palm plantation to maintain good yield while reducing the environmental impact. Apparently, years of over fertilization can have lasting effects well beyond the period when fertilization management changes. As the palm circle is a hotspot of N₂O emissions, improved nutrient management in this zone may have the potential to minimize fertilizer-induced N₂O emissions, e.g. through application of slow-release N fertilizers, use of nitrification inhibitors, adjusting N application rate with age of the plantation, and understory vegetation to take up and recycle excess mineral N (Sakata et al., 2015; Ashton-Butt et al., 2018; Cassman et al., 2019). Moreover, return of organic residues (empty fruit bunches or mill effluent) should be encouraged to improve nutrient retention and recycling, and to reduce dependency on chemical fertilizers (Bakar et al., 2011; Formaglio et al., 2021).



4.3 Soil CH₄ uptake

Area-weighted soil CH₄ uptake (Table 1) was comparable to CH₄ uptake reported for oil palm plantations ($-15 \pm 3 \mu\text{g C m}^{-2} \text{ h}^{-1}$) but lower than soil CH₄ uptake in forests on similar loam Acrisol soils in Jambi, Indonesia ($-29 \pm 12 \mu\text{g C m}^{-2} \text{ h}^{-1}$) (Hassler et al., 2015). The soil CH₄ uptake at our site was larger than that reported by Drewer et al., (2021a) ($-3 \pm 1 \mu\text{g C m}^{-2} \text{ h}^{-1}$) from ≤ 12 years old oil palm plantations on clay Acrisol soil in Malaysian Borneo, which they attributed to very high CH₄ emissions from a plot adjacent to a riparian area. Also, the soil CH₄ uptake at our site was larger than that reported by Aini et al. (2020) (ranging from -1 to $13 \mu\text{g C m}^{-2} \text{ h}^{-1}$) for an oil palm plantation on sandy clay loam Cambisol soil in Jambi, Indonesia, which they explained by the high WFPS ($> 80\%$) during their measurement period. Clay content or soil texture is the main site factor that correlate positively to soil CH₄ fluxes (Veldkamp et al., 2013), indicating that the higher the clay content the lower is the soil CH₄ uptake. High clay content soils have a low proportion of coarse pores (Hillel, 2003), which are important for gas diffusive transport. Furthermore, soils with high clay content have high water-holding capacity, which hinders gas diffusion from the atmosphere to the soil and limits CH₄ availability to methanotrophic activity in the soil (Keller and Reiners, 1994; Veldkamp et al., 2013). Thus, the disparity of soil CH₄ uptake between our present site and these above-mentioned studies was attributed to their differences in soil texture or drainage status as well as rainfall conditions or WFPS during the measurement periods, which all influence CH₄ diffusion from the atmosphere to the soil and thereby its uptake.

The positive correlation between soil CH₄ fluxes and WFPS (Fig. 4g), which reflected the same spatial pattern and positive correlation between soil CH₄ fluxes and soil bulk densities (Fig. S5), was attributed to the reduction of CH₄ diffusion from the atmosphere to the soil with increases in WFPS and soil bulk density (Veldkamp et al., 2013; Martinson et al., 2021). Differences in soil bulk density that result from management practices in oil palm plantations (Table S1) (Formaglio et al., 2021) affect soil total porosity, WFPS and gas diffusivity (Keller and Reiners, 1994; Hassler et al., 2015). The frond-stacked area has large SOC (Table S1) and low soil bulk density (Fig. S5) or high porosity (Formaglio et al., 2021), resulting in low WFPS (Table 2). Thus, with the high soil porosity in the frond-stacked area, gas diffusion may not limit CH₄ availability to methanotrophic activity in the soil, resulting in the highest soil CH₄ uptake among the management zones (Table 1). Where gas diffusion was favorable in the frond-stacked area, the increase in soil mineral N had increased CH₄ uptake (Fig. 4i), suggesting that mineral N availability enhanced CH₄ uptake once gas diffusion is not limiting (Veldkamp et al., 2013; Hassler et al., 2015). Conversely, the palm circle and inter-row have small SOC (Table S1) and high soil bulk density (Fig. S5) or low porosity (Formaglio et al., 2021), resulting in high WFPS (Table 2), which may have limited gas diffusion and possibly created anaerobic microsites, and thereby the occasional soil CH₄ emissions during the wet season (Fig. S3). Thus, the observation that soil mineral N did not influence soil CH₄ fluxes from the palm circle and inter-row (Fig. 4i) was possibly because gas diffusion limitation was the overriding factor controlling soil CH₄ uptake. Aside from improving the soil biochemical properties with the decadal piling of senesced fronds on the frond-stacked area (Table S1), which favor for increases in soil microbial biomass (Fig. S4) and N cycling rate (Formaglio et al., 2021), stimulating soil CH₄ sink or methanotrophic activity is one proof of the multiple benefits of conserving soil organic matter, e.g. its role on soil GHG abatement (Veldkamp et al., 2020). Foot traffic from management practices in the palm circle and inter-row as well as reduced litter inputs had increased soil bulk density and decreased SOC (Table S1), accompanied by reductions in microbial biomass and its activity (e.g. low soil N cycling rate; Formaglio et al., 2021), including reduced methanotrophic activity (Table 1). At the plantation level, the overall comparable soil CH₄ uptake between the reduced and conventional management treatments indicated that changes in fertilization rates and weeding methods



390 did not yet affect the drivers of soil CH₄ uptake (e.g. comparable soil mineral N and WFPS among treatments; Table 2) at least during the first four years of this experiment. Instead, the spatial differences in soil CH₄ uptake suggest that restoring the function of soil as CH₄ sink should be geared towards increasing soil organic matter, e.g. alternating locations of piled fronds with unused inter-rows, returning empty fruit bunches and other processing by-products, and avoiding plant biomass burning in establishing the next generation oil palm plantation (Bakar et al., 2011; Carron et al., 2015; Bessou et al., 2017).

395 4.4 Global warming potential

The GWP of this ≥ 18 -year old, large-scale oil palm plantation (Fig. 3) was in the lower end of the estimate from another part of this plantation near a peat soil (GWP of 686 ± 353 g CO₂-eq m⁻² yr⁻¹; Meijide et al., 2020). The slight difference between our estimated GWP and this previous study, aside from the latter's proximity to a peat soil, can be due to plantation age, different climatic conditions during separate study years, and different employed methods. First, as to plantation age, oil palm acts as a net C source one year after forest conversion with a net ecosystem exchange (NEE) of 1012 ± 51 g C m⁻² yr⁻¹ and becomes a net C sink at 12 years old (NEE of -754 ± 38 g C m⁻² yr⁻¹; Meijide et al., 2020). However, C removed from the field via fruit harvest turns this plantation into a net C source (NEP of 146 ± 94 g C m⁻² yr⁻¹; Meijide et al., 2020). At our study plots, the average annual yield during 2017–2020 across treatments (Fig. 3; Table S2) was higher than that of Meijide et al. (2020) (900 ± 49 g C m⁻² yr⁻¹). Our estimated NEP (i.e. heterotrophic respiration – (NPP – fruit biomass C); Fig. 3) (Quiñones et al., 2022; Iddris et al., 2023) was in the lower end of Meijide et al.'s (2020) NEP estimate. We attributed this small NEP to be due to large biomass and yield production as oil palm trees aged and also possibly due to reduced heterotrophic respiration as SOC had already decreased and reported to attain a steady-state low level after ~ 15 years from forest conversion (van Straaten et al., 2015). Secondly, the different climate conditions during our study year compared to the study by Meijide et al. (2020) may also have contributed to the differences in biomass and yield production as well as heterotrophic respiration. Our measurement period (2019–2020) had annual rainfall within the 10-year average, whereas Meijide et al.'s (2020) study years (2014–2016) included a severe drought in 2015 caused by a strong El Niño Southern Oscillation that induced prolonged smog events in Jambi, Indonesia (Field et al., 2016). Drought combined with smoke haze reduce the productivity and CO₂ uptake in this oil palm plantation (Stiegler et al., 2019), supporting the low fruit yield measured by Meijide et al. (2020). Thirdly, our different estimation methods may have contributed to the small disparity between our GWP estimate and that by Meijide et al. (2020), who measured NEE by eddy covariance technique. Our method and theirs both have advantages and disadvantages; ours based on measurements of NPP components and assumption of heterotrophic respiration contribution to measured soil CO₂ emissions (van Straaten et al., 2011) was spatially replicated, inexpensive and practicable to deploy without a need for electricity source in the field (Baldocchi, 2014), e.g. for plot-scale experiment on different fertilization regimes and weeding practices.

420 As soil N₂O emissions contributed substantially (55%) to the GWP of this large-scale oil palm plantation while soil CH₄ sink had only minor offset ($< 2\%$) (Fig. 3), reducing its GHG footprint could be achieved by decreasing soil N₂O emissions and increasing soil CH₄ uptake (see above). Finally, the large NPP of this ≥ 18 -year old, large-scale oil palm plantation and reduced heterotrophic respiration (due to reduced SOC after ~ 15 years from forest conversion; van Straaten et al., 2015) contributed to its small NEP that accounted 46% of the GWP. In the perspective of long-term oil palm management, extending the rotation period from 25 years to 30 years to prolong accumulation of plant biomass C (Meijide et al., 2020), avoiding large biomass

425



loss during establishment of the next generation oil palms (e.g. not burning but leaving cut palm trees on the field), and enhancing SOC stocks will reduce the GHG footprint of oil palm plantations.

5 Conclusions

430 During the first four years of this management experiment, soil GHG fluxes, GWP, and yield in reduced fertilization with mechanical weeding remained similar to conventional fertilization with herbicide application, signifying the strong legacy effect of over a decade of high fertilization regime prior to the start of our experiment in this mature oil palm plantation. Reducing soil N₂O emissions is the key to reducing GHG footprint in this oil palm plantation as soil N₂O emissions contributed substantially to the GWP of this oil palm plantation. The palm circle and frond-stacked area both showed high mineral N availability, but the palm circle was driven by the high fertilization rates and rendering it as a hotspot of soil N₂O emissions. In 435 contrast, the frond-stacked area support high root and microbial biomass, low soil N₂O emissions and high soil CH₄ uptake. Thus, reasonably expanding the frond-stacked area and returning organic residues from oil palm fruit processing in order to reduce dependency on chemical fertilizers will help reduce the GHG footprint while maintaining high production in oil palm plantations.

Acknowledgements

440 This study was funded by the Deutsche Forschungsgemeinschaft (DFG) project number 192626868—SFB 990 /2-3) in the framework of the Collaborative Research Center 990 EForTS as part of project A05. Guantao Chen was supported by China Scholarship Council. The PT Perkebunan Nusantara VI company provided no funding and did not have any influence on the study design, data collection, analysis, or interpretation. We thank PTPN VI for allowing us to conduct research in their plantation. We are especially thankful to our Indonesian field and laboratory assistants, Fajar Sidik, Mohammed Fatoni and the 445 project Z01 field personnel, for managing the field implementation of this experiment. We thank Andrea Bauer, Dirk Böttger, Kerstin Langs, and Natalia Schröder for their assistance in the laboratory analysis. This study was conducted under the research permit 148/E5/E5.4/SIP/2019.

Author contributions

EV and MDC conceptualized the study, implemented the field-plot design and measurement methodologies. AT, BI and MD 450 facilitated field access, logistical support, collaborator agreements and material exports. GC conducted the field works and data analysis, and wrote the first draft of the manuscript. MDC and EV revised extensively the manuscript, and all co-authors approved the final manuscript.

Funding

This study was funded by the Deutsche Forschungsgemeinschaft (DFG) project ID 192626868 (SFB 990/2-3) as part of project 455 A05.



Data availability

All data of this study are deposited on the GRO Göttingen Research Online data repository (<https://data.goettingen-research-online.de>) and under the following DOIs: 10.25625/NR8SRD, which is accessible to all members of the Collaborative Research Center (CRC) 990. Based on the data sharing agreement within the CRC 990, these data are currently not publicly accessible
460 but will be made available through a written request to the senior author.

Declaration

Conflict of interest At least one of the (co-)authors is a member of the editorial board of Biogeosciences.

References

- 465 Adachi, M., Bekku, Y. S., Rashidah, W., Okuda, T., and Koizumi, H.: Differences in soil respiration between different tropical ecosystems, *Appl Soil Ecol*, 34, 258–265, <https://doi.org/10.1016/j.apsoil.2006.01.006>, 2006.
- Aini, F. K., Hergoualc’h, K., Smith, J. U., and Verchot, L.: Nitrous oxide emissions along a gradient of tropical forest disturbance on mineral soils in Sumatra, *Agric Ecosyst Environ*, 214, 107–117, <https://doi.org/10.1016/j.agee.2015.08.022>, 2015.
- 470 Aini, F. K., Hergoualc’h, K., Smith, J. U., Verchot, L., and Martius, C.: How does replacing natural forests with rubber and oil palm plantations affect soil respiration and methane fluxes? *Ecosphere*, 11, e03284, <https://doi.org/10.1002/ecs2.3284>, 2020.
- Allen, K., Corre, M. D., Tjoa, A., and Veldkamp, E.: Soil Nitrogen-Cycling Responses to Conversion of Lowland Forests to Oil Palm and Rubber Plantations in Sumatra, Indonesia, *PLoS One*, 10, e0133325, <https://doi.org/10.1371/journal.pone.0133325>, 2015.
- 475 Allen, K., Corre, M. D., Kurniawan, S., Utami, S. R., and Veldkamp, E.: Spatial variability surpasses land-use change effects on soil biochemical properties of converted lowland landscapes in Sumatra, Indonesia, *Geoderma*, 284, 42–50, <https://doi.org/10.1016/j.geoderma.2016.08.010>, 2016.
- Asari, N., Suratman, M. N., Jaafar, J., and Khalid, M. Md.: Estimation of Above Ground Biomass for Oil Palm Plantations Using Allometric Equations, *Int Proc Chem Biol Environ Eng*, 32, 12–16, <https://doi.org/10.7763/IPCBE.2013.V58.22>, 2013.
- 480 Ashton-Butt, A., Aryawan, A. A. K., Hood, A., Naim, M., Purnomo, D., Suhardi, Wahyuningsi, R., Willcock, S., Poppy, G. M., Caliman, J.-P., Turner, E. C., Foster, W. A., Peh, K. S.-H., and Snaddon, J. L.: Understorey vegetation in oil palm plantations benefits soil biodiversity and decomposition rates, *Front For Glob Chang*, 1, 10, <https://doi.org/10.3389/ffgc.2018.00010>, 2018.
- 485 Bakar, R. A., Darus, S. Z., Kulaseharan, S., and Jamaluddin, N.: Effects of ten year application of empty fruit bunches in an oil palm plantation on soil chemical properties, *Nutr Cycl Agroecosyst*, 89, 341–349, <https://doi.org/10.1007/s10705-010-9398-9>, 2011.
- Baldocchi, D.: Measuring fluxes of trace gases and energy between ecosystems and the atmosphere - the state and future of the eddy covariance method, *Glob Chang Biol*, 20, 3600–3609, <https://doi.org/10.1111/gcb.12649>, 2014.
- 490 Bessou, C., Verwilghen, A., Beaudoin-Ollivier, L., Marichal, R., Ollivier, J., Baron, V., Bonneau, X., Carron, M. P., Snoeck, D., Naim, M., Aryawan, A. A. K., Raoul, F., Giraudoux, P., Surya, E., Sihombing, E., and Caliman, J. P.: Agroecological



- practices in oil palm plantations: Examples from the field, *OCL - Oilseeds and fats, Crops and Lipids*, 24, D305, <https://doi.org/10.1051/ocl/2017024>, 2017.
- 495 Bond-Lamberty, B., Wang, C., and Gower, S. T.: A global relationship between the heterotrophic and autotrophic components of soil respiration? *Glob Chang Biol*, 10, 1756–1766, <https://doi.org/10.1111/j.1365-2486.2004.00816.x>, 2004.
- Bou Dib, J., Alamsyah, Z., and Qaim, M.: Land-use change and income inequality in rural Indonesia, *For Policy Econ*, 94, 55–66, <https://doi.org/10.1016/j.forpol.2018.06.010>, 2018.
- 500 Carron, M. P., Pierrat, M., Snoeck, D., Villenave, C., Ribeyre, F., Suhardi, Marichal, R., and Caliman, J. P.: Temporal variability in soil quality after organic residue application in mature oil palm plantations, *Soil Res*, 53, 205–215, <https://doi.org/10.1071/SR14249>, 2015.
- Cassman, N. A., Soares, J. R., Pijl, A., Lourenço, K. S., van Veen, J. A., Cantarella, H., and Kuramae, E. E.: Nitrification inhibitors effectively target N₂O-producing *Nitrosospora* spp. in tropical soil, *Environ Microbiol*, 21, 1241–1254, <https://doi.org/10.1111/1462-2920.14557>, 2019.
- 505 Clough, Y., Krishna, V. V., Corre, M. D., Darras, K., Denmead, L. H., Mejjide, A., Moser, S., Musshoff, O., Steinebach, S., Veldkamp, E., Allen, K., Barnes, A. D., Breidenbach, N., Brose, U., Buchori, D., Daniel, R., Finkeldey, R., Harahap, I., Hertel, D., Holtkamp, A. M., Hörandl, E., Irawan, B., Jaya, I. N. S., Jochum, M., Klarner, B., Knohl, A., Kotowska, M. M., Krashevskaya, V., Kreft, H., Kurniawan, S., Leuschner, C., Maraun, M., Melati, D. N., Opfermann, N., Pérez-Cruzado, C., Prabowo, W. E., Rembold, K., Rizali, A., Rubiana, R., Schneider, D., Tjitrosoedirdjo, S. S., Tjoa, A., Tscharnkte, T., and Scheu, S.: Land-use choices follow profitability at the expense of ecological functions in Indonesian smallholder landscapes, *Nat Commun*, 7, 13137, <https://doi.org/10.1038/ncomms13137>, 2016.
- 510 Crawley, M. J.: *The R book second edition*, John Wiley & Sons Ltd, Chichester, UK, 2013.
- Cusack, D. F., Ashdown, D., Dietterich, L. H., Neupane, A., Ciochina, M., and Turner, B. L.: Seasonal changes in soil respiration linked to soil moisture and phosphorus availability along a tropical rainfall gradient, *Biogeochemistry*, 145, 235–254, <https://doi.org/10.1007/s10533-019-00602-4>, 2019.
- 515 Dassou, O., Nodichao, L., Aholoukpè, H., Cakpo, Y., and Jourdan, C.: Improving the methodology for root biomass estimation in monocotyledonous tree plantations: case of oil palm (*Elaeis guineensis*. Jacq) in West Africa, *Plant Soil*, 465, 593–611, <https://doi.org/10.1007/s11104-021-04939-4>, 2021.
- Davidson, E. A., Keller, M., Erickson, H. E., Verchot, L. V., and Veldkamp, E.: Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides, *Bioscience*, 50, 667–680, 2000.
- 520 Descals, A., Wich, S., Meijaard, E., Gaveau, D. L. A., Peedell, S., and Szantoi, Z.: High-resolution global map of smallholder and industrial closed-canopy oil palm plantations, *Earth Syst Sci Data*, 13, 1211–1231, <https://doi.org/10.5194/essd-13-1211-2021>, 2021.
- Drewer, J., Kuling, H. J., Cowan, N. J., Majalap, N., Sentian, J., and Skiba, U.: Comparing Soil Nitrous Oxide and Methane Fluxes From Oil Palm Plantations and Adjacent Riparian Forests in Malaysian Borneo, *Front For Glob Chang*, 4, 145, <https://doi.org/10.3389/ffgc.2021.738303>, 2021a.
- 525 Drewer, J., Leduning, M. M., Griffiths, R. I., Goodall, T., Levy, P. E., Cowan, N., Comynn-Platt, E., Hayman, G., Sentian, J., Majalap, N., and Skiba, U. M.: Comparison of greenhouse gas fluxes from tropical forests and oil palm plantations on mineral soil, *Biogeosciences*, 18, 1559–1575, <https://doi.org/10.5194/bg-18-1559-2021>, 2021b.
- FAO: Food and Agriculture Organization, FAOSTAT. <https://www.fao.org/faostat/en/#data/QCL>, 2021
- 530 Field, R. D., Van Der Werf, G. R., Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., Levy, R., Livesey, N. J., Luo, M., Torres, O., and Worden, H. M.: Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought, *Proc Natl Acad Sci U S A*, 113, 9204–9209, <https://doi.org/10.1073/pnas.1524888113>, 2016.



- Formaglio, G., Veldkamp, E., Duan, X., Tjoa, A., and Corre, M. D.: Herbicide weed control increases nutrient leaching compared to mechanical weeding in a large-scale oil palm plantation, *Biogeosciences*, 17, 5243–5262, <https://doi.org/10.5194/bg-17-5243-2020>, 2020.
- 535 Formaglio, G., Veldkamp, E., Damris, M., Tjoa, A., and Corre, M. D.: Mulching with pruned fronds promotes the internal soil N cycling and soil fertility in a large-scale oil palm plantation, *Biogeochemistry*, 154, 63–80, <https://doi.org/10.1007/s10533-021-00798-4>, 2021.
- Grass, I., Kubitzka, C., Krishna, V. V., Corre, M. D., Mußhoff, O., Pütz, P., Drescher, J., Rembold, K., Ariyanti, E. S., Barnes, A. D., Brinkmann, N., Brose, U., Brümmer, B., Buchori, D., Daniel, R., Darras, K. F. A., Faust, H., Fehrmann, L., Hein, J., Hennings, N., Hidayat, P., Hölscher, D., Jochum, M., Knohl, A., Kotowska, M. M., Krashevskaya, V., Kreft, H., Leuschner, C., Lobite, N. J. S., Panjaitan, R., Polle, A., Potapov, A. M., Purnama, E., Qaim, M., Röhl, A., Scheu, S., Schneider, D., Tjoa, A., Tschardtke, T., Veldkamp, E., and Wollni, M.: Trade-offs between multifunctionality and profit in tropical smallholder landscapes, *Nat Commun*, 11, 1186, <https://doi.org/10.1038/s41467-020-15013-5>, 2020.
- 540 Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiol Rev*, 60, 439–471, <https://doi.org/10.1128/membr.60.2.439-471.1996>, 1996.
- Hassler, E., Corre, M. D., Tjoa, A., Damris, M., Utami, S. R., and Veldkamp, E.: Soil fertility controls soil-atmosphere carbon dioxide and methane fluxes in a tropical landscape converted from lowland forest to rubber and oil palm plantations, *Biogeosciences*, 12, 5831–5852, <https://doi.org/10.5194/bg-12-5831-2015>, 2015.
- 550 Hassler, E., Corre, M. D., Kurniawan, S., and Veldkamp, E.: Soil nitrogen oxide fluxes from lowland forests converted to smallholder rubber and oil palm plantations in Sumatra, Indonesia, *Biogeosciences*, 14, 2781–2798, <https://doi.org/10.5194/bg-14-2781-2017>, 2017.
- Hillel, D.: *Introduction to Environmental Soil Physics*, Elsevier, <https://doi.org/10.1016/B978-0-12-348655-4.X5000-X>, 2003.
- 555 Iddris, N. A., Formaglio, G., Paul, C., von Groß, V., Chen, G., Angulo-Rubiano, A., Berkelmann, D., Brambach, F., Darras, K. F. A., Krashevskaya, V., Potapov, A., Wenzel, A., Irawan, B., Damris, M., Daniel, R., Grass, I., Kreft, H., Scheu, S., Tschardtke, T., Tjoa, A., Veldkamp, E., and Corre, M. D.: Mechanical weeding enhances ecosystem multifunctionality and profit in industrial oil palm, *Nat Sustain*. <https://doi.org/10.1038/s41893-023-01076-x>, 2023.
- 560 IPCC: *Guidelines for national greenhouse gas inventories chapter 11: N₂O emissions from managed soils, and CO₂ emissions from lime and urea application*, 2006.
- Kaupper, T., Hetz, S., Kolb, S., Yoon, S., Horn, M. A., and Ho, A.: Deforestation for oil palm: impact on microbially mediated methane and nitrous oxide emissions, and soil bacterial communities, *Biol Fertil Soils*, 56, 287–298, <https://doi.org/10.1007/s00374-019-01421-3>, 2020.
- 565 Keller, M. and Reiners, W. A.: Soil-atmosphere exchange of nitrous oxide, nitric oxide, and methane under secondary succession of pasture to forest in the Atlantic lowlands of Costa Rica, *Global Biogeochem Cycles*, 8, 399–409, <https://doi.org/10.1029/94GB01660>, 1994.
- Khalil, M. I., Rosenani, A. B., Van Cleemput, O., Fauziah, C. I., and Shamshuddin, J.: Nitrous Oxide Emissions from an Ultisol of the Humid Tropics under Maize-Groundnut Rotation, *J Environ Qual*, 31, 1071–1078, <https://doi.org/10.2134/jeq2002.1071>, 2002.
- 570 Kotowska, M. M., Leuschner, C., Triadiati, T., Meriem, S., and Hertel, D.: Quantifying above- and belowground biomass carbon loss with forest conversion in tropical lowlands of Sumatra (Indonesia), *Glob Chang Biol*, 21, 3620–3634, <https://doi.org/10.1111/gcb.12979>, 2015.
- 575 Kurniawan, S., Corre, M.D., Matson, A.L., Schulte-Bisping, H., Utami, S.R., van Straaten, O. and Veldkamp, E.: Conversion of tropical forests to smallholder rubber and oil palm plantations impacts nutrient leaching losses and nutrient retention efficiency in highly weathered soils. *Biogeosciences*, 15, <https://doi.org/10.5194/bg-15-5131-2018>, 5131–5154, 2018.



- Liyanage, A., Grace, P. R., Scheer, C., de Rosa, D., Ranwala, S., and Rowlings, D. W.: Carbon limits non-linear response of nitrous oxide (N₂O) to increasing N inputs in a highly-weathered tropical soil in Sri Lanka, *Agric Ecosyst Environ*, 292, 106808, <https://doi.org/10.1016/j.agee.2019.106808>, 2020.
- 580 Malhi, Y., Baldocchi, D. D., and Jarvis, P. G.: The carbon balance of tropical, temperate and boreal forests, *Plant Cell Environ*, 22, 715–740, <https://doi.org/10.1046/j.1365-3040.1999.00453.x>, 1999.
- Martinson, G. O., Müller, A. K., Matson, A. L., Corre, M. D., and Veldkamp, E.: Nitrogen and phosphorus control soil methane uptake in tropical montane forests, *J Geophys Res Biogeosci*, 126, e2020JG005970, <https://doi.org/10.1029/2020JG005970>, 2021.
- 585 Meijide, A., de la Rua, C., Guillaume, T., Röhl, A., Hassler, E., Stiegler, C., Tjoa, A., June, T., Corre, M. D., Veldkamp, E., and Knohl, A.: Measured greenhouse gas budgets challenge emission savings from palm-oil biodiesel, *Nat Commun*, 11, 1089, <https://doi.org/10.1038/s41467-020-14852-6>, 2020.
- Moradi, A., Teh, C. B. S., Goh, K. J., Husni, M. H. A., and Ishak, C. F.: Decomposition and nutrient release temporal pattern of oil palm residues, *Ann Appl Biol*, 164, 208–219, <https://doi.org/10.1111/aab.12094>, 2014.
- 590 Nelson, P. N., Webb, M. J., Banabas, M., Nake, S., Goodrick, I., Gordon, J., O’Grady, D., and Dubos, B.: Methods to account for tree-scale variability in soil- and plant-related parameters in oil palm plantations, *Plant Soil*, 374, 459–471, <https://doi.org/10.1007/s11104-013-1894-7>, 2014.
- OECD: Oilseeds and oilseed products, in: *OECD-FAO Agricultural Outlook 2022-2031*, OECD Publishing, Paris, <https://doi.org/10.1787/5778f78d-en>, 2022.
- 595 Quiñones, C. M. O., Veldkamp, E., Lina, S. B., Bande, M. J. M., Arribado, A. O., and Corre, M. D.: Soil greenhouse gas fluxes from tropical vegetable farms, using forest as a reference, *Nutr Cycl Agroecosyst*, 124, 59–79, <https://doi.org/10.1007/s10705-022-10222-4>, 2022.
- R Core Team: A language and environment for statistical computing. Foundation for Statistical Computing, Vienna, Austria, 2021.
- 600 Rahman, N., Bruun, T. B., Giller, K. E., Magid, J., van de Ven, G. W. J. J., and de Neergaard, A.: Soil greenhouse gas emissions from inorganic fertilizers and recycled oil palm waste products from Indonesian oil palm plantations, *GCB Bioenergy*, 11, 1056–1074, <https://doi.org/10.1111/gcbb.12618>, 2019.
- Rochmyaningsih, D.: Making peace with oil palm, *Science*, 365, 112–115, <https://doi.org/10.1126/science.365.6449.112>, 2019.
- 605 Rüegg, J., Quezada, J. C., Santonja, M., Ghazoul, J., Kuzyakov, Y., Buttler, A., and Guillaume, T.: Drivers of soil carbon stabilization in oil palm plantations, *Land Degrad Dev*, 30, 1904–1915, <https://doi.org/10.1002/ldr.3380>, 2019.
- Ryadin, A. R., Janz, D., Schneider, D., Tjoa, A., Irawan, B., Daniel, R., and Polle, A.: Early effects of fertilizer and herbicide reduction on root-associated biota in oil palm plantations, *Agronomy*, 12, 199, <https://doi.org/10.3390/agronomy12010199>, 2022.
- 610 Sakata, R., Shimada, S., Arai, H., Yoshioka, N., Yoshioka, R., Aoki, H., Kimoto, N., Sakamoto, A., Melling, L., and Inubushi, K.: Effect of soil types and nitrogen fertilizer on nitrous oxide and carbon dioxide emissions in oil palm plantations, *Soil Sci Plant Nutr*, 61, 48–60, <https://doi.org/10.1080/00380768.2014.960355>, 2015.
- Skiba, U., Hergoualc’h, K., Drewer, J., Meijide, A., and Knohl, A.: Oil palm plantations are large sources of nitrous oxide, but where are the data to quantify the impact on global warming? *Curr Opin Environ Sustain*, 47, 81–88, <https://doi.org/10.1016/j.cosust.2020.08.019>, 2020.
- 615 Sotta, E. D., Veldkamp, E., Schwendenmann, L., Guimarães, B. R., Paixão, R. K., Ruivo, M. de L. P., Lola da Costa, A. C., and Meir, P.: Effects of an induced drought on soil carbon dioxide (CO₂) efflux and soil CO₂ production in an Eastern Amazonian rainforest, Brazil, *Glob Chang Biol*, 13, 2218–2229, <https://doi.org/10.1111/j.1365-2486.2007.01416.x>, 2007.



- 620 Stiegler, C., Meijide, A., Fan, Y., Ali, A. A., June, T., and Knohl, A.: El Niño-Southern Oscillation (ENSO) event reduces CO₂ uptake of an Indonesian oil palm plantation, *Biogeosciences*, 16, 2873–2890, <https://doi.org/10.5194/bg-16-2873-2019>, 2019.
- Tao, H. H., Slade, E. M., Willis, K. J., Caliman, J. P., and Snaddon, J. L.: Effects of soil management practices on soil fauna feeding activity in an Indonesian oil palm plantation, *Agric Ecosyst Environ*, 218, 133–140, <https://doi.org/10.1016/j.agee.2015.11.012>, 2016.
- 625 Tchifo Lontsi, R., Corre, M. D., Iddris, N. A., and Veldkamp, E.: Soil greenhouse gas fluxes following conventional selective and reduced-impact logging in a Congo Basin rainforest, *Biogeochemistry*, 151, 153–170, <https://doi.org/10.1007/s10533-020-00718-y>, 2020.
- Thomas, M., McLaughlin, D., Grubba, D., Buchanan, J., McLaughlin, D., Grubba, D., and Buchanan, J.: Sustainable sourcing guide for palm oil users: A practical handbook for US consumer goods and retail companies, Conservation International and WWF US, 83, 2015.
- 630 van Straaten, O., Veldkamp, E., and Corre, M. D.: Simulated drought reduces soil CO₂ efflux and production in a tropical forest in Sulawesi, Indonesia, *Ecosphere*, 2, art119, <https://doi.org/10.1890/ES11-00079.1>, 2011.
- van Straaten, O., Corre, M. D., Wolf, K., Tchienkoua, M., Cuellar, E., Matthews, R. B., and Veldkamp, E.: Conversion of lowland tropical forests to tree cash crop plantations loses up to one-half of stored soil organic carbon, *Proceedings of the National Academy of Sciences*, 112, 9956–9960, <https://doi.org/10.1073/pnas.1504628112>, 2015.
- 635 Veldkamp, E., Weitz, A. M., and Keller, M.: Management effects on methane fluxes in humid tropical pasture soils, *Soil Biol Biochem*, 33, 1493–1499, [https://doi.org/10.1016/S0038-0717\(01\)00060-8](https://doi.org/10.1016/S0038-0717(01)00060-8), 2001.
- Veldkamp, E., Koehler, B., and Corre, M. D.: Indications of nitrogen-limited methane uptake in tropical forest soils, *Biogeosciences*, 10, 5367–5379, <https://doi.org/10.5194/bg-10-5367-2013>, 2013.
- 640 Veldkamp, E., Schmidt, M., Powers, J. S., and Corre, M. D.: Deforestation and reforestation impacts on soils in the tropics, *Nat Rev Earth Environ*, 1, 590–605, <https://doi.org/10.1038/s43017-020-0091-5>, 2020.
- Vijay, V., Pimm, S. L., Jenkins, C. N., and Smith, S. J.: The impacts of oil palm on recent deforestation and biodiversity loss, *PLoS One*, 11, e0159668, <https://doi.org/10.1371/journal.pone.0159668>, 2016.



Table 1. Soil CO₂, N₂O, and CH₄ fluxes (means ± SE, n = 4 plots) from different fertilization and weeding treatments in an ≥ 18-year old, large-scale oil palm plantation, Jambi, Indonesia, measured monthly from July 2019 to June 2020

Soil greenhouse gas	Management zones	Treatments				across treatments	LME model <i>P</i> -values		
		ch	cw	rh	rw		numDF = 1, denDF = 12	Fertilization	Weeding
CO ₂ flux (mg C m ⁻² h ⁻¹)	Palm circle	82.06 ± 9.95	84.27 ± 8.70	92.32 ± 12.69	93.51 ± 5.70	88.04 ± 4.48 b	0.95	0.94	0.89
	Inter-row	38.54 ± 4.73	41.46 ± 3.41	50.34 ± 4.14	39.86 ± 5.31	42.55 ± 2.32 c	0.31	0.47	0.13
	Frond-stacked area	126.21 ± 6.23	132.95 ± 11.17	120.23 ± 11.15	117.95 ± 6.57	124.33 ± 4.34 a	0.21	0.89	0.65
	area-weighted	59.52 ± 3.49	62.60 ± 2.05	68.38 ± 6.37	61.23 ± 4.45	62.93 ± 2.14	0.55	0.66	0.22
N ₂ O flux (μg N m ⁻² h ⁻¹)	Palm circle	198.96 ± 53.70	219.50 ± 131.74	301.90 ± 121.53 (205.17 ± 85.71) ^a	91.47 ± 7.55	202.96 ± 46.14a	0.38	0.16	0.31
	Inter-row	10.14 ± 3.94	7.40 ± 1.56	13.67 ± 5.56	4.48 ± 1.13	8.92 ± 1.81 c	0.90	0.14	0.26
	Frond-stacked area	12.82 ± 0.98	16.12 ± 4.26	37.50 ± 26.14 (17.73 ± 6.98) ^a	9.74 ± 2.91	19.04 ± 6.59 b	0.38	0.47	0.62
	area-weighted	44.53 ± 8.18	46.56 ± 24.31	69.13 ± 28.67	20.93 ± 1.87	45.29 ± 9.67	0.53	0.20	0.07
CH ₄ flux (μg C m ⁻² h ⁻¹)	Palm circle	-2.89 ± 1.19	-3.19 ± 1.63	-3.74 ± 2.46	-5.97 ± 1.70	-3.95 ± 0.87 a	0.26	0.66	0.29
	Inter-row	-16.94 ± 5.39	-14.06 ± 1.69	-14.13 ± 6.27	-14.46 ± 2.98	-14.90 ± 2.03 b	0.89	0.66	0.84
	Frond-stacked area	-42.53 ± 1.93	-38.89 ± 3.36	-36.09 ± 6.59	-44.66 ± 6.44	-40.54 ± 2.39 c	0.88	0.73	0.28
	area-weighted	-18.25 ± 3.74	-15.89 ± 1.42	-15.55 ± 4.75	-17.46 ± 2.97	-16.79 ± 1.57	0.98	0.73	0.58

For each soil greenhouse gas, different letters within each column indicate significant differences among management zones across treatments (2² factorial ANOVA with linear mixed-effects models and Tukey HSD test at p ≤ 0.05); numDF and denDF are numerator and denominator degrees of freedom, respectively. ch: conventional fertilization – herbicide weeding, cw: conventional fertilization – mechanical weeding, rh: reduced fertilization – herbicide weeding, rw: reduced fertilization – mechanical weeding


^a For soil N₂O fluxes, values in parenthesis excluded two extreme outliers in the palm circle (5311 and 4325 μg N m⁻² h⁻¹) and one extreme outlier in the frond-stacked area (1934 μg N m⁻² h⁻¹)



Table 2. Soil temperature, water content, and mineral N concentrations (means ± SE, $n = 4$ plots) measured in the top 5 cm from different fertilization and weeding treatments in an ≥ 18-year old, large-scale oil palm plantation, Jambi, Indonesia, measured monthly from July 2019 to June 2020

Soil factors	Management zones	Treatments				across treatments			LME model P -values		
		ch	cw	rh	rw	treatments			Fertilization	Weeding	Interaction
soil temperature (°C)	Palm circle	26.2 ± 0.2	26.2 ± 0.2	26.3 ± 0.2	26.2 ± 0.1	26.2 ± 0.1 a			0.44	0.57	0.55
	Inter-row	26.1 ± 0.2	26.2 ± 0.2	26.2 ± 0.1	26.0 ± 0.1	26.1 ± 0.1 a			0.89	0.91	0.21
	FronD-stacked area	26.1 ± 0.1	26.2 ± 0.1	26.2 ± 0.1	26.1 ± 0.1	26.2 ± 0.1 a			0.66	0.93	0.21
Water-filled pore space (%)	Palm circle	38.8 ± 2.5	35.6 ± 2.3	34.9 ± 2.9	40.4 ± 2.1	37.5 ± 1.2 a			0.89	0.71	0.32
	Inter-row	34.4 ± 3.0	35.4 ± 1.4	36.5 ± 1.5	36.9 ± 1.9	35.8 ± 1.0 a			0.53	0.71	0.74
	FronD-stacked area	25.2 ± 2.4	26.7 ± 3.0	27.9 ± 3.0	23.6 ± 1.1	25.8 ± 1.2 b			0.86	0.53	0.24
NH ₄ ⁺ (µg N g ⁻¹)	Palm circle	6.3 ± 5.1	15.2 ± 10.9	1.7 ± 0.8	13.7 ± 8.3	9.3 ± 3.6 a			0.42	0.08	0.64
	Inter-row	0.7 ± 0.1	0.8 ± 0.1	0.8 ± 0.1	0.8 ± 0.1	0.7 ± 0.1 c			0.24	0.84	0.26
	FronD-stacked area	1.8 ± 0.1	4.1 ± 1.9	2.3 ± 0.1	1.8 ± 0.1	2.50 ± 0.5 b			0.52	0.27	0.02
NO ₃ ⁻ (µg N g ⁻¹)	Palm circle	5.2 ± 2.3	14.6 ± 6.1	4.5 ± 1.9	10.0 ± 4.7	8.6 ± 2.1 a			0.42	0.29	0.88
	Inter-row	0.4 ± 0.1	0.6 ± 0.2	0.6 ± 0.3	0.3 ± 0.1	0.5 ± 0.1 c			0.54	0.39	0.14
	FronD-stacked area	10.1 ± 2.5	14.1 ± 1.4	5.3 ± 1.0	3.6 ± 0.4	8.3 ± 1.3 b			<0.01	0.53	0.16


For each soil factor, different letters within each column indicate significant differences among management zones across treatments (2^2 factorial ANOVA with linear mixed-effects models and Tukey HSD test at $p \leq 0.05$); numDF and denDF are numerator and denominator degrees of freedom, respectively. ch: conventional fertilization – herbicide weeding, cw: conventional fertilization – mechanical weeding, rh: reduced fertilization – herbicide weeding, rw: reduced fertilization – mechanical weeding



655

Figure 1: Experimental set-up. A 2×2 factorial experiment design with four blocks (OM1–4) within which are the four treatments (each plot was $50 \text{ m} \times 50 \text{ m}$; ch: conventional fertilization – herbicide weeding, cw: conventional fertilization – mechanical weeding, rh: reduced fertilization – herbicide weeding, rw: reduced fertilization – mechanical weeding) (a). Two subplots were selected in the central $30 \text{ m} \times 30 \text{ m}$ area in each plot (b). In each subplot, soil GHG flux measurements were conducted at each management zone (palm circle, inter-row, and frond-stacked area) using vented static chambers (c)

660



665 **Figure 2: Annual soil CO₂, N₂O, and CH₄ fluxes (means ± SE, *n* = 4 plots), weighted by the areal coverages of**
the palm circle (18%), inter-row (67%), and frond-stacked area (15%) under different fertilization and
weeding treatments in an ≥ 18-year old, large-scale oil palm plantation, Jambi, Indonesia. ch: conventional
fertilization – herbicide weeding, cw: conventional fertilization – mechanical weeding, rh: reduced fertilization –
herbicide weeding, rw: reduced fertilization – mechanical weeding. Annual soil N₂O emissions from rh were
670 calculated excluding three extreme outliers: 5311 and 4325 μg N m⁻² h⁻¹ in the palm circle and 1934 μg N m⁻² h⁻¹ in
the frond-stacked area. Annual soil CO₂, N₂O, and CH₄ fluxes were not statistically tested since these are trapezoidal
extrapolations between measurement periods




Figure 3: Ecosystem net global warming potential (GWP) from an ≥ 18 -year old, large-scale oil palm plantation, Jambi, Indonesia (means \pm SE, $n = 16$ plots). Negative and positive symbols indicate the direction of the flux: (-) for C uptake and (+) for C export or emission from the plantation. Net primary productivity (NPP, $\text{g C m}^{-2} \text{ yr}^{-1}$) = aboveground biomass C production + fruit biomass C + frond litter biomass C + belowground biomass C production + root litter biomass C. Aboveground biomass C production = annual biomass production per palm (2019–2020) \times planting density \times tissue C concentration. Annual fruit biomass C = annual fruit production per palm (mean of 2019–2020) \times planting density \times tissue C concentration. Frond litter biomass C = annual litter production per palm (mean of 2019–2020) \times planting density \times tissue C concentration. Belowground biomass C production and root litter C were taken from oil palm sites in the same area (Kotowska et al., 2015). Net ecosystem productivity (NEP, $\text{g C m}^{-2} \text{ yr}^{-1}$) = heterotrophic respiration – (NPP – fruit biomass C) (Malhi et al., 1999; Quiñones et al., 2022). Heterotrophic respiration was assumed to be 70% of soil respiration (van Straaten et al., 2011) + 80% from decomposition of frond litter (Iddris et al., 2023). GWP ($\text{g CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$) = NEP_in $\text{CO}_2 + \text{N}_2\text{O_CO}_2\text{-eq} + \text{CH}_4\text{_CO}_2\text{-eq}$, whereby the CO_2 -equivalents for N_2O and CH_4 are their annual fluxes multiplied by 298 and 25, respectively, for a 100-year time frame (IPCC 2006)




Figure 4: Relationships among soil CO₂, N₂O, and CH₄ fluxes and soil factors. Spearman rank coefficients (rho) marked by * indicates $p \leq 0.05$. Each data point was the average of four replicate plots per treatment on each measurement period ($n = 48$ (i.e. 4 treatments \times 12 months) for soil temperature; $n = 40$ (i.e. 4 treatments \times 10 months) for soil water-filled pore space (WFPS) and total mineral N). P – palm circle, IR – inter-row, FS – frond-stacked area

690