

Lawns and meadows in urban green space – A comparison from perspectives of greenhouse gas, drought resilience and plant functional types

Justine Trémeau^{1,2,3,4}, Beñat Olascoaga^{4,5,6}, Leif Backman¹, Esko Karvinen¹, Henriikka Vekuri¹ and Liisa Kulmala^{1,7}

¹ Finnish Meteorological Institute, Helsinki, Finland

² AgroParisTech – Institut des sciences et industries du vivant et de l'environnement, Palaiseau, France

³ Université Paris-Saclay, Orsay, France

⁴ Institute for Atmospheric and Earth System Research (INAR), Department of Physics, Faculty of Science, University of Helsinki, Helsinki, Finland

⁵ School of Resource Wisdom, Department of Biological and Environmental Science, Faculty of Mathematics and Science, University of Jyväskylä, Jyväskylä, Finland

⁶ Helsinki Institute of Sustainability Science (HELSUS), Department of Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, Helsinki, Finland

⁷ Institute for Atmospheric and Earth System Research (INAR), Forest Sciences, University of Helsinki, Helsinki, Finland

Correspondence: Justine Trémeau (justine.tremeau@fmi.fi) and Liisa Kulmala (liisa.kulmala@fmi.fi)

Abstract. Today, city planners design urban futures by considering environmental degradation and climate change and biodiversity loss mitigation. Here, we studied the greenhouse gas fluxes of urban lawns and meadows and linked the observations with plant functional types and soil properties. In eight lawns and eight meadows in the Helsinki Metropolitan Area, Finland, carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) fluxes were measured using manual chambers and plant functional types were recorded. Four of these sites i.e., an irrigated lawn, an old mesic meadow, a non-irrigated lawn, and a young dry meadow, were more intensively studied in 2021-2022. The process-based ecosystem model JSBACH was utilized together with the momentary observations collected approximately every second week on CO₂ exchange to quantify the annual carbon (C) balance of these sites. On the remaining sites, we studied the initial dynamics of conversion from lawns to meadows by transforming part of lawns to meadows in late 2020 and conducting measurements from 2020 to 2022. We found that without considering the impact of management and unstabilized soil, lawns are clear sinks of carbon (NEE = -336 ± 187 and -157 ± 139 g CO₂ m⁻² yr⁻¹) compared with meadows (NEE = -151 ± 198 and -63 ± 134 g CO₂ m⁻² yr⁻¹), and the mean photosynthetic production (GPP) of the irrigated lawn and mesic meadow was the highest in this study, whereas the dry meadow had the lowest GPP. The studied lawns were stronger C sinks, compared to the meadows. However, the net exchange values were uncertain as the soils were not in equilibrium with the vegetation at all sites, which is common for urban habitats, and modeling the heterotrophic emissions was therefore challenging. The conversion from a lawn to a meadow did not affect the fluxes of CH₄ and N₂O. Moreover, the mesic meadow was more resistant to drought events than the non-irrigated lawn. Lastly, the proportion of herbaceous flowering plants (forbs, including legumes) other than grasses was found to be higher in meadows than in lawns. Even though social and economic aspects also steer urban development, these results can guide planning when considering biodiversity environmentally friendlier green spaces and carbon-smartness.

Keywords. Carbon balance, photosynthesis, respiration, CH₄, N₂O, JSBACH, plant functional types, urban nature

1 Introduction

The impacts of climate change and biodiversity crises environmental degradation are increasingly evident. The These crises have feedbacks are interlinked with each other by feedbacks and as climate change threatens biodiversity, species and

~~ecosystems~~, the loss of biodiversity and ~~possible~~ degradation of ecosystems ~~also~~ affects the climate system. Therefore, ~~it is crucial to understand~~ understanding the role ~~that of~~ various ecosystems in ~~the land-use sector play in~~ regulating climate and supporting biodiversity, ~~and to find~~ within the land-use sector is crucial. It is also important to identify land management practices that can effectively mitigate the climate and ~~biodiversity~~ other environmental crises simultaneously. Well-known management practices for the land use sector include, for example, conservation and restoration, and integrated land-use planning taking into account the relationships between different land uses (Niemelä et al., 2010; Pörtner et al., 2021).

In the urban context, green spaces not only sequester atmospheric carbon (C) but ~~also~~ provide vital ecosystem services, such as cooling, ~~recreation~~ recreational values, purification of air and water, and ~~reducing the risk~~ reduction ~~for of~~ flooding (Niemelä et al., 2010; Belmeziti et al., 2018; Lampinen et al., 2021; Shen et al., 2023). Cities actively seek optimal green ~~areas~~ space planning and management practices to mitigate and adapt to climate change, and there are ~~also~~ numerous initiatives aimed at promoting urban green spaces and ~~green~~ infrastructure (Haaland and van den Bosch, 2015). One of ~~these~~ these is Article 6 ~~in of~~ the Nature Restoration ~~proposal~~ proposals launched by the European Commission (European Commission, 2022), which sets targets for increasing ~~urban~~ green ~~urban~~ spaces in cities, towns and suburbs.

Within urban landscapes, lawns ~~constitute~~ are one of the most common features of green space (Hedblom et al., 2017; Ignatieva et al., 2020), and are usually subjected to frequent and intense management to fulfil social, aesthetic and recreational purposes (Ignatieva et al., 2015, 2017; Zobec et al., 2020; Paudel and States, 2023). Although many types of lawns exist, most of the urban green spaces worldwide are dominated by turfgrass lawns. These lawns typically contain certain selected species of grasses, ~~such as~~ in Southern Finland ~~typically~~ *Agrostis capillaris*, *Alopecurus pratensis*, *Dactylis glomerata*, *Festuca ovina*, *F. pratensis*, *F. rubra* and *Poa annua* (Tonteri and Haila, 1990), ~~yet they~~. They can also harbor other species of ~~forbs and~~ grasses ~~that are and additional plant functional types such as forbs, all~~ spontaneously established, ~~and~~ thus giving lawns the ability to behave like semi-natural grasslands (Thompson et al., 2004; Fischer et al., 2013). Reducing the conventional turfgrass lawn management has been proved to enhance the abundance, ~~species~~ richness and diversity of ~~vegetation~~ plants and arthropods (Venn and Kotze, 2014; Watson et al., 2020), ~~thus~~ these passively created urban grasslands can have a positive effect on biodiversity. An alternative approach to creating more environmentally friendly and ~~biodiverse~~ heterogeneous urban grasslands consists of substituting or modifying part of the lawnscape from the short, monocultural, homogeneous setting of grass species into meadows, a flowering setting with extensive management and possible active incorporation of forbs (Southon et al., 2017; Lane et al., 2019; Norton et al., 2019; Bretzel et al., 2020; Marshall et al., 2023). This practice is becoming increasingly common as more cities and other communities look for ways to create sustainable and low-maintenance green space (Smith et al., 2015; Unterweger et al., 2017; Norton et al., 2019; Ignatieva et al., 2020). Within the context of climate mitigation and adaptation, the capacity of urban lawns to assimilate carbon varies widely due to climatic, edaphic and biological factors. Furthermore, management factors (e.g., mowing and irrigation) can make lawns behave either as carbon sinks or sources (Selhorst and Lal, 2013; Kong et al., 2014; Wang et al., 2022). For instance, soil organic carbon content in this type of urban green space tends to be larger than in natural grasslands (Kaye et al., 2005; Pouyat et al., 2006). Yet, carbon emissions from urban lawn soils have also been shown to be higher than those from natural and extensively managed urban grasslands (Kaye et al., 2005; Upadhyay et al., 2021). Unlike lawns and other types of grasslands, meadows do not only comprise grasses but also additional plant functional types such legumes and other forbs, and the relative proportions of different plant functional types may influence the overall carbon dynamics and ~~greenhouse gas~~ (GHG) exchange. However, the relative impact of urban lawns and meadows on soil and vegetation dynamics, carbon balance and ~~greenhouse gas~~ (GHG) exchange, remains poorly understood.

It is known that natural ecosystems with ~~higher~~ more heterogeneous vegetation ~~species richness~~ generally show higher photosynthetic productivity ~~than less diverse ones~~, due to a more efficient use of resources by a more ~~diversified~~ varying vegetation contributing to carbon assimilation (Fornara and Tilman, 2008; Lange et al., 2015; Yang et al., 2019; Chen et al., 2020). Additionally, ~~they~~ this could lead to a more resistant functioning of the ecosystem ~~in~~ during extreme weather events, as

85 different grasses and forbs can respond differently to changes in environmental conditions (De Keersmaecker et al., 2016; Hossain et al., 2022). ~~This means that, for~~For example, during a drought event some plant species may be more resistant, making the functioning of a diverse green space less vulnerable ~~than~~compared to a ~~less-diverse~~more homogeneous setting. However, the relationship between ~~species-richness~~vegetation, productivity and resistance is not straightforward but influenced by many factors, such as management, vegetation community and environmental conditions (Vogel et al., 2012; De
90 Keersmaecker et al., 2016; Jung et al., 2020).

The aim of this study ~~is~~was to determine the climatic impact of transforming urban lawns into urban meadows in northern Europe. In practice, we had four specific research questions:

1. Does the carbon balance differ between urban lawns and urban meadows?
- 95 2. Are urban meadows more tolerant to extreme weather events than urban lawns?
3. Does the transformation from an urban lawn to an urban meadow increase GHG emissions?
4. Do plant functional types ~~and their proportions~~ affect GHG fluxes in urban grasslands?

In addition, we were interested to see if we could detect any connections between the different plant functional types and ~~carbon~~C and nitrogen (N) pools. To answer these questions, we measured GHG fluxes in urban lawns, in recently created
100 urban meadows and in an older urban meadow in the Helsinki Metropolitan Area in Finland. Observations of the sites were used to set up a land-surface model, which was used to simulate annual ~~carbon~~C balances. The empirical data set included intensive sites with high temporal coverage of CO₂ exchange to estimate the ~~carbon~~C balance and satellite sites with high spatial coverage to study the transformation process.

2 Materials and method

105 2.1 Site description

2.1.1 Study region

The measurements were collected within the Helsinki Metropolitan Area (60°10' N, 24°57' E, Fig. 1), which is situated on the northern coast of the Gulf of Finland. In Helsinki, the mean annual precipitation and temperature were 653 mm and 6.5 °C, respectively, for the reference period of 1991–2020, with monthly mean temperatures being above 10 °C from May to
110 September (Jokinen et al., 2021). According to the Köppen climate classification, the climate is humid continental (Dfb) (Kottek et al., 2006). The data collection took place on intensive and satellite sites which ~~are~~ all ~~are~~ considered urban ~~locations~~.

2.1.2 Intensive sites

In order to study the CO₂ exchange with high temporal coverage and to compare the ~~carbon~~C balances of urban grasslands, two pairs of urban lawns and urban meadows, situated 4 kilometers apart, were selected for the intensive measurement sites
115 (Fig. 1). In Kumpula (KMP) neighborhood, an old, mesic and mesotrophic urban meadow, which could be associated to the “~~Mesi~~mesic perennial anthropogenic herbaceous vegetation (V39)” in the EUNIS classification (European Environment Agency, 2023) and hereinafter referred to as “mesic meadow”, was paired with a highly managed urban lawn, hereinafter referred to as “irrigated lawn”. ~~The~~During the campaign, the meadow ~~is~~was an old agricultural field on which farming practices were abandoned about 40 years ago. Currently, *Aegopodium podagraria*, *Lupinus polyphyllus* (non-native invasive), *Dactylis glomerata*, *Anthriscus sylvestris*, *Elymus repens*, *Lamium album* and *Urtica dioica* are the dominant ~~plant~~ species. It ~~is~~was cut once a year in autumn and most of the vegetation clippings ~~are~~were taken away. It ~~is~~was neither irrigated nor fertilized. The lawn, mainly covered by *Poa pratensis*, was installed about 15 years ago. ~~It is currently~~During the campaign, it was mowed automatically by a mowing robot that ~~operates~~operated daily between 6 PM and 11 AM, and the grass clippings ~~are~~were

125 pounded and left on the site. It ~~is~~was irrigated during the summer when needed and was last fertilized in spring 2021. The mesic meadow and the irrigated lawn were situated 150 meters apart from each other.

In Viikki (VKI) neighborhood, the urban lawn and urban meadow – hereinafter referred to as “non-irrigated lawn” and “dry meadow” – ~~are~~were situated inside a public park 60 meters apart from each other. The urban lawn of *Festuca sp.* was sown in 2005 and ~~is~~managed as a “utility lawn” (Viherympäristöliitto, 2023). ~~It is~~During the campaign, it was mown regularly so that the grass height ~~is~~was 4–12 cm, and the clippings ~~are~~were mostly left on the site. There ~~is~~was no irrigation, and the lawn ~~has~~had not been fertilized for several years. In 2020, a section of the park’s lawns was transformed into a dry and nutrient-poor urban meadow by replacing the topsoil with a layer of recycled sand and sowing seeds (Riikonen and Karilas, 2021). This young dry urban meadow tends toward a “~~Dry~~dry perennial anthropogenic herbaceous vegetation (V38)” in the EUNIS classification (European Environment Agency, 2023). In 2022, the dominant species were *Trifolium repens*, *Tripleurospermum inodorum*, ~~Leucanthemum~~*Leucanthemum vulgare*, *Centaurea jacea*, *Phleum pratense* and *Plantago major*, and some rare
135 species such as *Dianthus deltoides*, *Campanula rotundifolia* and *Galium verum* were also observed. It was fertilized in spring 2021 and mown for the first time in autumn 2022. After mowing, most of the clippings were removed from the site.

2.1.3 Satellite sites

To study the transformation process from an urban lawn into an urban meadow, six other locations belonging to local student housing associations were selected for high spatial coverage GHG measurements in the urban green space. Six urban lawns
140 over 20 years old, each located near the developing urban meadows and all subject to regular lawn management, were selected and studied as control cases. All of the six lawns were predominantly covered by grasses (mostly *Poa pratensis*, *Festuca rubra* and *Lolium ~~perenne~~perenne*). In each of the locations, 50 m² of the already established lawn was transformed into an urban meadow site by volunteering and community participation. Site transformations were conducted during late autumn 2020 by manually turning over the existent soil up to a depth of ca. 35 cm, followed by the sowing of 14 pollinator-friendly forbs at a
145 density of 0.06 g m⁻². The commercial seed mixture (Suomen niittysiemen Oy, Koski Tl, Finland) also included four grass species at a density up to 0.6 g m⁻². Consequently, the hemiparasitic forb *Rhinanthus minor*, which is known to have negative effects on grass cover (Chaudron et al., 2021), was sown at a 5 g m⁻² density. Every year since the lawn transformation, each of the developing urban meadows ~~were~~was managed by manually removing on-site tree seedlings (mainly *Acer platanoides*) before a one-time mowing event in autumn by hand scythe. Lastly, all mowing clippings were removed from the sites.

150




Figure 1. Locations of the intensive (yellow) and satellite (green) measurement sites around the Helsinki metropolitan area. Background maps built with topographic database by the National Land Survey of Finland (2023) and global administrative borders by GADM (2023).

155

Table 1. Site characteristics at the intensive and satellite sites. **The sown** mixture indicates seeds of pollinator-friendly forbs, grass species and *Rhinanthus minor* which were sown in late 2020 and veg. refers to vegetation. (± standard deviation), calculated for at least four replicates. At each VKI site there was only one replicate for each variable. Due to Finnish classification, approximations were made for sand (0.06–2.0 mm), silt (0.002–0.06 mm) and clay (<0.002 mm).

Experiment	Location	Management	Soil texture	Vegetation's characteristic	Irrigation	Sand (%)	Silt (%)	Clay (%)	pH	SOC (%)	SON (%)	CN ratio
Intensive	KMP	irrigated lawn	sandy loam	Grasses (Poa)	Yes	69.5	22.1	8.4	5.6	3.8 (± 1.2)	0.25 (± 0.06)	14.8 (± 2.8)
		meadow	sandy loam	Mesic veg.	No	59.1	31.8	9.1	6.5	5.6 (± 1.8)	0.46 (± 0.16)	12.3 (± 0.7)
	VKI	lawn	sandy loam	Grasses (Fescue)	No	72.2	25.6	2.2	6.3	2.8	0.19	14.5
		meadow	sandy loam	Xerophilic veg.	No	72.4	17.1	10.5	6.1	2.5	0.17	14.9
Satellite	JMT-3	lawn	silt loam	Grasses	No	5.2	68.9	26.0	5.3 (± 0.1)	5.0 (± 0.4)	0.46 (± 0.04)	10.8 (± 0.1)
		meadow	loamy sand	Sown mixture	No	73.7	23.5	2.8	6.5 (± 0.2)	4.6 (± 0.7)	0.18 (± 0.03)	25.7 (± 0.2)
	JMT-7	lawn	loamy sand	Grasses	No	82.7	14.9	2.4	6.5 (± 0.1)	4.0 (± 0.5)	0.26 (± 0.04)	15.0 (± 0.5)
		meadow	loamy sand	Sown mixture	No	81.4	16.0	2.6	6.5 (± 0.2)	3.2 (± 0.3)	0.21 (± 0.02)	15.3 (± 0.6)
	KO-4	lawn	sandy loam	Grasses	No	66.2	29.4	4.4	6.4 (± 0.3)	4.4 (± 1.2)	0.29 (± 0.08)	15.4 (± 0.6)
		meadow	sandy loam	Sown mixture	No	63.3	32.7	4.0	6.1 (± 0.4)	4.0 (± 0.7)	0.19 (± 0.04)	20.9 (± 1.7)
	PK-3	lawn	sandy loam	Grasses	No	68.8	27.6	3.6	6.2 (± 0.1)	4.0 (± 0.7)	0.19 (± 0.03)	21.1 (± 0.4)
		meadow	sandy loam	Sown mixture	No	68.4	27.1	4.5	6.1 (± 0.1)	5.4 (± 0.3)	0.32 (± 0.04)	17.2 (± 1.8)
	SK-1	lawn	sandy loam	Grasses	No	57.3	33.9	8.8	5.6 (± 0.1)	4.5 (± 0.3)	0.37 (± 0.03)	12.3 (± 0.5)
		meadow	sandy loam	Sown mixture	No	54.1	36.5	9.4	5.4 (± 0.0)	4.2 (± 0.1)	0.35 (± 0.01)	12.0 (± 0.3)
SMT-12	lawn	loam	Grasses	No	47.7	42.7	9.7	5.8 (± 0.1)	3.6 (± 0.3)	0.29 (± 0.05)	12.8 (± 1.3)	
	meadow	loam	Sown mixture	No	51.8	39.6	8.6	6.1 (± 0.2)	4.4 (± 0.7)	0.31 (± 0.05)	14.2 (± 0.4)	

160

2.1.4 Sampling design

In this study, the lawn and the meadow within a pair were paired considering the short distance between each other and the two sites were mostly visited on the same day. GHG flux measurements were conducted at each site along a transect. All the data was collected at four fixed quadrats of approximately 1 m² distributed along a single transect. At the intensive sites, most

165 of the quadrats were about 10 meters apart, but to fit the study setup to the landscape design, there were some variations mainly
due to passing walking paths: at the non-irrigated lawn, the quadrats were separated by 10 m, 5 m and 10 m; at the dry meadow,
the quadrats were separated by 17 m, 10 m and 10 m; at the mesic meadow, the quadrats were located exactly ~~located~~ 10
meters apart; and last, the quadrats on the irrigated lawn were located in two separate transects 15 meters away from each
other, with the two quadrats on each transect being 3 meters apart. At the satellite sites, from the centroid of the site, a ~~10m~~10
170 m long transect was established along the length of the site, with four quadrats of 1-m² area each uniformly distributed.
~~The soil~~Soil temperature, soil moisture, soil samples and plant ~~inventories~~functional type cover percentages were collected
close to the GHG measurement quadrats except for the irrigated lawn ~~where, in which~~ we used ~~the~~ soil samples collected from
the same plot by Ahongshangbam et al. (2023).

2.2 Flux measurements

175 2.2.1 Measurement protocol

The GHG flux measurements were conducted with manual chambers using different setups at the intensive and satellite sites.
At the intensive sites, the carbon dioxide (CO₂) exchange measurements were conducted more frequently than at the satellite
sites, and they also included the light response (LR) of net ecosystem exchange (NEE). At the satellite sites, the measurements
also included methane (CH₄) and nitrous oxide (N₂O) exchanges, but measurements were conducted with opaque chambers,
180 thus disabling photosynthesis. The chamber measurements were always performed within the above-mentioned quadrats.

At the intensive sites, the light response of NEE was measured fortnightly between June and September 2021–2022. Additional
measurements of NEE, but mainly of total ecosystem respiration (TER), were also conducted in May and in October–
November. Over the two years of measurements, KMP lawn was visited 23 times, including 15 days of LR, KMP meadow 23
185 times, including 16 days of LR, VKI lawn and meadow 20 times, including 15 days of LR, and at each site, most of the time
the four collars were measured at a site (Table S2). The measurement setup consisted of a transparent chamber attached to a
CO₂ and H₂O analyzer (Li-840A, LI-COR, Inc., Nebraska, USA), an air temperature and humidity sensor (BME280, Bosch
Sensortec GmbH, Reutlingen, Germany), and a photosynthetically active radiation (PAR) sensor (PQS1, Kipp & Zonen, Delft,
Netherlands). The air was circulated through the analyzer at a flow rate of 1 L min⁻¹, and the chamber was equipped with a
190 small fan to ensure air mixing within the chamber headspace. We used two differently sized chambers, either 0.072 m³ or
0.288 m³, and three collars of different heights (18.3 cm, 20.8 cm and 50 cm) to individually match the height of the
chamber+collar combination to that of the vegetation we were measuring during each measurement day. The transparent
chamber was placed on the ground while avoiding damaging the plants, and a tight seal was created by using sand-filled cloth
bags as insulation. In meadows, when a collar was used, it was placed on the ground and sealed with the cloth bags, and the
195 chamber was placed on top of the collar. The duration of each chamber closure was at least 2 minutes. To determine the light
response of NEE, we repeated the measurement five times at each collar under different PAR intensities ~~–(i.e., 100%, 50%,
25%, 12.5% and 0% of the prevailing sunlight level–)~~ created by shading the chamber with netted or opaque fabrics. We first
measured without covering the chamber and then covered it with 1–3 layers of mesh cloth. Eventually, we measured the last
recording with an opaque cover. There was a pause of at least one minute in between the single light intensities to ventilate
200 the chamber and to allow the plants some time to acclimatize to the changing light level. While conducting the measurements
on the field, we allowed for ±10% variation in the PAR level during a single closure.

At the satellite sites, GHG measurements were conducted with an opaque chamber monthly, between May–September 2021–
2022. Additional background measurements were conducted in July, August and October 2020, before the lawn
205 transformations took place. In total, the sites were visited thirteen times: three times before transformation, including two times

with CH₄ and N₂O measurements and one time with only CO₂, and ten times after transformation, including six times with CH₄ and N₂O measurements and four times with only CO₂ (Table S2). A single measurement was performed at each of the four selected 1-m² quadrats at each site (Table S2). To allow a good representativity of CO₂ measurements, but also measurements of N₂O and CH₄, two different types of devices were used over the different months (Table S2). The measurement setup in August and October 2020, as well as in May, July and September 2021 and 2022 consisted of a CO₂, H₂O, N₂O and CH₄ analyzer (DX4015, Gaset Technologies Oy, Vantaa, Finland) connected to an opaque chamber equipped with a small fan. This Fourier Transform Infrared-based analyzer was zero-calibrated at the beginning of each measurement day. The chamber volume was either 0.00456 m³ or 0.268 m³ depending on the height of the vegetation measured. An additional collar of 20-cm height was needed to allow enough room for tall vegetation in July 2021 and 2022. The measurements in July 2020 and in June and August 2021 and 2022 were conducted with a setup consisting of an opaque chamber (volume 0.007434 m³) equipped with a CO₂ probe (GMP343, Vaisala Oyj, Vantaa, Finland), a relative humidity and air temperature sensor (HMP75, Vaisala), and a small fan. This chamber is introduced in detail by Ryhti et al. (2021). When measuring only CO₂ (i.e., July 2020, June and August 2021 and 2022), the duration of each chamber closure was 4–5 minutes. We extended the closure time to 15 minutes when measuring also N₂O and CH₄ (i.e., August and October 2020, May, July and September 2021 and 2022).

2.2.2 Flux calculation

The measurements represent the total ecosystem fluxes: ~~total~~ i.e., the sum of inflow and outflow of CH₄ fluxes (μg CH₄ m⁻² s⁻¹), ~~total~~ N₂O fluxes (μg N₂O m⁻² s⁻¹) and CO₂ fluxes (NEE (μg CO₂ m⁻² s⁻¹). NEE includes possible photosynthetic input (GPP, μg CO₂ m⁻² s⁻¹) and the outflow by respiration (TER, μg CO₂ m⁻² s⁻¹), which includes both autotrophic and heterotrophic respiration. This can be mathematically written as:

$$NEE(t) = TER(t) + GPP(PAR(t)), \quad (1)$$

where PAR (μmol m⁻² s⁻¹) is photosynthetically active radiation and t is time. Here we ~~use~~ used meteorological notation i.e., we ~~consider~~ considered the inflow of carbon to be negative (GPP ≤ 0) and therefore NEE ~~can~~ could be either negative (i.e., sink of CO₂), when the absolute value of GPP ~~is~~ was higher than TER, or positive (i.e., source), when TER ~~is~~ was higher than the absolute value of GPP.

NEE or the total flux of other GHG fluxes during each chamber closure was calculated with the following Equation (2):

$$flux = \left(\frac{dC(t)}{dt} \right)_{t=0} \frac{M_{GHG} P V}{R T A}, \quad (2)$$

where $\left(\frac{dC(t)}{dt} \right)_{t=0}$ is the time derivative (ppm s⁻¹ = 10⁻⁶ s⁻¹) of the linear regression, M_{GHG} is the molecular mass of the GHG (44.01 g mol⁻¹ for CO₂, 16.05 g mol⁻¹ for CH₄ and 44.02 g mol⁻¹ for N₂O), P is the air pressure (Pa), V is the system (chamber + possible collar) volume (m³), R is the universal gas constant (8.31446 J mol⁻¹ K⁻¹), T is the temperature inside the chamber headspace during the closure (K) and A is the soil surface i.e. the basal area of the chamber (m²).

TER, CH₄ and N₂O fluxes were obtained from the measurement conducted with the opaque cover (intensive sites) or with the opaque chamber (satellite sites).

To ensure a proper air mixing inside the chamber, at least ten seconds were removed from the beginning of each closure before fitting a linear regression into the GHG concentration values. Otherwise, the quality control and the length of the measurements included in the flux calculations differed between the instruments and GHGs. For the Vaisala equipment, CO₂ fluxes were calculated with the first minute of the recordings with R (version 4.2.2) after visual validation of linearity. Python (version 3.9.7) was used to calculate the fluxes measured by the two other devices. With the Gaset analyzer, the CO₂ fluxes were calculated from the first five minutes of the recordings whereas CH₄ and N₂O fluxes, which were significantly lower fluxes

and closer to the detection limits of the equipment, were calculated with the first seven minutes of the recordings. For the intensive sites, where the LI-COR analyzer was used, all fluxes were calculated from the first minute of the recording. There, possible poor-quality measurements were eliminated from the end of the measurement period if there was at least 1 minute of good quality data. The quality control measures were stability of light conditions inside the chamber during the closure and normalized root mean squared error (NRMSE) of the fit. Yet, it is worth noting that in cases where the flux is nearly zero, NRMSE can be very large even though the measurement is not erroneous. Therefore, we did not discard measurements based on NRMSE if $(\frac{dC(t)}{dt})_{t=0}$ was below 0.1 ppm s⁻¹. Otherwise, measurements were discarded if NRMSE was larger than 0.05. Accordingly, 41 NEE measurements out of 1225 recordings were discarded from the intensive site dataset, but none of them referred to the opaque chamber measurements, i.e., TER. Thus, TER was represented by 23 and 20 days of data at KMP and VKI sites, respectively. Regarding the satellite sites, the filtering process described above was also applied to the Gasset dataset resulting in discarding 23 measurements of TER out of 384 recordings over the three years of measurements, but no CH₄, nor N₂O measurements (384 recordings of each in total). All TER measurements (240 TER recordings in total) measured with the Vaisala device were considered as valid (Table S2).

260

2.2.3 Light response (LR) fitting and daily GPP

In order to fit LR curves to the CO₂ flux data, we first subtracted TER from the other NEE rates to obtain an estimate for GPP (Eq. 1). GPP(PAR=0) was set to zero. The LR curve was expected to be a rectangular hyperbola (Ruimy et al., 1995), as follows:

$$GPP = \frac{PAR \alpha GPmax}{PAR \alpha + GPmax} \quad (3)$$

where PAR ($\mu\text{mol m}^{-2} \text{s}^{-1}$) is photosynthetically active radiation, α ($\text{mg } \mu\text{mol}^{-1}$) is the initial slope between GPP and PAR, and GPmax ($\text{mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) is the light-saturated photosynthesis rate. The parameters α and GPmax were estimated using non-linear least squares minimization. α was allowed to vary between -0.1 and $-1 \cdot 10^{-8} \text{ mg } \mu\text{mol}^{-1}$ and GPmax between -5 and $-1 \cdot 10^{-7} \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$.

270 In order to fit the LR curve for one collar, a minimum of three measurements done at different light intensities was required. After processing the light response calculation, LR curves where α had standard error over $0.0008 \text{ mg } \mu\text{mol}^{-1}$ or GPmax standard error was higher than $10 \text{ mg CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ were discarded from the dataset. In addition, LR was discarded if the highest PAR intensity in the response was under $500 \mu\text{mol m}^{-2} \text{ s}^{-1}$. That resulted in 154 LR curves out of 212, representing 14 days of data at KMP meadow, VKI lawn and meadow and 10 days at KMP lawn, which were used in further analyses.

275

Daily GPP was calculated for each collar and for each measurement day using Eq. 3, fitted parameters for the collar and day, and continuous PAR measurements from SMEAR III station (Institute of Atmospheric Research, 2023). First, GPP was estimated for each half hour with the fitted parameters and the 30-min average PAR. Then, all half-hourly GPPs were summed up to obtain the daily GPP of the collar on the particular day.

280 To compare the mean GPP at the different intensive sites, we selected consecutive weeks where all sites had a high number of high-quality LR curves. We estimated the mean daily GPP for all sites by utilizing the means of the obtained LR parameters at each site and the 30-minute average PAR values over the two weeks. The two-week periods which were included in this comparison started on May 31, June 14 and 28, July 12 and August 9 in 2021 and on May 16, June 13 and 27, and August 8 in 2022.

2.3.1 Soil temperature and soil moisture

Soil temperature and soil moisture were always measured together with the chamber measurements within the 1-m² quadrats at both the intensive and the satellite sites. One replicate of soil temperature was measured within the quadrats at a 10-cm depth with a handheld soil thermometer (HH376, Omega Engineering Inc., Connecticut, USA), and 4–5 replicates of soil moisture at 5 cm depth were measured with a handheld setup (ML3 ThetaProbe and HH2 Moisture Meter, Delta-T Devices Ltd., Cambridge, UK).

2.3.2 Soil characterization

At the intensive sites, overall soil characteristics, soil density and C and N contents were analyzed.

To determine the overall soil characteristics, we collected altogether 1 L of soil up to 25 cm depth at each of the sites. At the irrigated lawn, the pooled sample comprised 16–18 individual samples collected in 2020 around the site with a thin auger (d = 2.3 cm). At the rest of the remaining intensive sites, the samples were collected in 2021 with a slightly larger auger (d = 5.0 cm) to collect 4 individual samples that were then pooled together. The fresh samples were stored in a fridge and sent to the lab within 1–3 days after collection. From these samples, particle size distribution (analyzed according to Elonen (1971)) and the overall soil characteristics were analyzed at a commercial laboratory (Eurofins Viljavuuspalvelu Oy, Mikkeli, Finland).

Soil density samples were collected by digging a small pit and carefully inserting a steel cylinder horizontally into an undisturbed pit wall at 10 cm depth. When the cylinder was fully inserted, it was gently detached by removing the surrounding soil while ensuring that all soil within the cylinder remained in place in order to achieve volumetric accuracy. Five replicates were collected at KMP lawn and three replicates at the other intensive sites. After collection, the samples were dried at 105 °C for 48 h and the dry weights were weighed.

Other soil samples were collected to determine C and N content. At KMP sites, samples for C and N were collected with a soil auger down to a 30 cm depth. Altogether, six individual samples were collected at the lawn and the meadow. At VKI, we used a soil auger to collect eight samples and then pooled them together. The irrigated lawn samples were first sieved with a 2-mm mesh sieve and then dried at 105 °C for 24 h. At the other intensive sites, the samples were first dried at 105 °C for 48 h, and then sieved with a 2-mm mesh sieve. Total soil C and N contents were determined from the dried and milled samples of soil with grain size smaller than 2 mm with an elemental CN analyzer (LECO, Michigan, USA). The soil C and N contents were assumed to be only organic components without carbonates i.e., soil organic carbon (SOC) and soil organic nitrogen (SON).

At the satellite sites, overall soil characteristics, soil C and N contents and soil particle distribution were also determined. All soil samples from the satellite sites were collected during early July 2022, and all samples were extracted within the 1-m² quadrats. A total of four soil cores, dug by means of using a soil auger down to ca. 15 cm depth, were pooled together to create a composite soil sample. In each of the sites, a total of four composite soil samples were collected in ziplock bags and kept on ice during soil collection. All soil samples were then stored at –20 °C, until shipped and analyzed at a commercial lab (Eurofins Viljavuuspalvelu Oy, Mikkeli, Finland). Total soil C and N contents were assumed to be only organic components without carbonates, and SOC and SON were determined from freeze-dried and milled samples with an elemental CN analyzer (LECO, Michigan, USA). For the case of soil particle size distribution, a fraction of each of the soil samples was repeatedly washed with a 30% H₂O₂ solution to remove its organic matter content. The resulting soil samples were then dried out at 105 °C for 48 h, before sieving (mesh size = 0.6 mm) the final mineral soils. The sieved soil samples were then mixed with a sodium pyrophosphate solution (0.05 M), and their sand, silt and clay proportions were calculated by means of laser diffraction particle size analyzer (Bekman Coulter LS230, Beckman Coulter Inc., California, USA).

2.4 Vegetation inventory

Vegetation inventories [to estimate plant functional type covers](#) were conducted in all intensive and satellite sites on June 27th and July 21st 2022, respectively, according to the quadrat sampling method. In each of the quadrats (see Section 2.1.4), the cover of each plant species was estimated and classified in one of the following plant functional type categories: forb, grass, horsetail, legume, moss, sedge or tree, where forb includes all other families of flowering vascular plants, which do not belong to one of the already listed categories. An average cover proportion was calculated from the four quadrats to estimate the overall proportion of each plant functional type at a site.

All collars were also photographed from above on each measurement day to calculate the green cover percentage of the basal area within the chamber using Canopeo (Patrignani and Ochsner, 2015).

2.5 Ecosystem modelling

To derive annual estimates of NEE, TER and GPP, the intensive sites were simulated using the JSBACH (Reick et al., 2013), which is the land component of the Max Planck Institute Earth system model (MPI-ESM) (Giorgetta et al., 2013). JSBACH is a process-based model and calculates the dynamic carbon cycle and key driving factors including seasonal dynamics in leaf area, momentary CO₂ fluxes, evapotranspiration, soil moisture, litter production and soil carbon dynamics.

The model was driven with hourly data (2005-2022) of air temperature, precipitation, shortwave and longwave radiation, relative humidity and wind speed. The driver data were derived from observations from the Kumpula weather station (60°12'14.0" N, 24°57'38.9" E) operated by the Finnish Meteorological Institute (Finnish Meteorological Institute, 2023a). The data was gapfilled by observations from the nearby urban measurement station SMEAR III (Järvi et al., 2009). Hourly ERA5-Land data (Muñoz-Sabater et al., 2021) was used to fill a small number of remaining gaps.

The vegetation in JSBACH is represented by plant functional types (PFT). Here, the model was set up for simulating only one PFT, C3 grass, for each site. The Logistic Growth Phenology (LoGro-P) model (Böttcher et al., 2016) is used to describe the phenology in JSBACH, where the temporal development of the leaf area index (LAI) of grass depends on both temperature and soil moisture. The maximum LAI for each site (Table S3) was set based on Sentinel-2 data (Nevalainen et al., 2022; Nevalainen, 2022), whereas the model simulated the seasonal LAI dynamics driven by temperature and precipitation. In addition, in the simulation of the mesic meadow the shedding of the grass was activated 65 days after the growth started, to better simulate the observed LAI.

According to the soil particle size distribution, the soil texture at all sites was sandy loam. The parameters describing the soil properties follow the recommendations by Hagemann and Stacke (2015). However, the volumetric field capacity and wilting point for each site were adjusted based on the soil moisture measurements (Table S3). The root depths of the lawns were assumed to be shallower than [that](#) of the meadows (Table S3).

The photosynthesis of C3-plants in JSBACH is described by the model by Farquhar et al. (1980). The photosynthesis is calculated once to get the unstressed canopy conductance, which is then scaled based on the soil moisture in the root zone to get the canopy conductance and photosynthesis under water stress. The available water in the root zone depends on the field capacity, root depth and a scaling factor $f = (W - W_{wilt}) / (W_{crit} - W_{wilt})$. The scaling factor is applied for relative soil moisture values between $W_{crit} = f_{crit} * W_{max}$ and $W_{wilt} = f_{wilt} * W_{max}$, where W_{max} is the maximum moisture content in the root zone. No water is available to the vegetation when the soil moisture reaches $f_{wilt} * W_{max}$. The factors f_{crit} and f_{wilt} are given in Table S3. Moreover, the daily GPP values were used to adjust the photosynthetic parameters to meet the observations at each intensive site.

The model was used to derive the annual average GPP, TER and NEE for the period 2005–2022. The simulations were set up to represent habitats where the soil organic matter accumulates over time from the litter of standing vegetation. This was achieved by running a long spin-up period (thousands of years). As it is often the case in urban areas, the soils at the intensive

sites had not been accumulated from the litter of the current vegetation, and therefore the soil carbon pools in the model are not equal to the ones present at the sites. Due to this, the simulations may not reproduce the observed TER, but instead represent a more general situation for these habitats. However, we also performed additional simulations where the soil carbon pools were adjusted to meet the observed TER values in 2021 and 2022. The agreement between the simulated and observed carbon fluxes was evaluated by R^2 and RMSE.

2.6 Drought definition

Since drought can be defined by the deficit of precipitation and the increase in evaporation (Wang et al., 2021), we estimated the drought period at our intensive sites with the Standardized Precipitation-Evapotranspiration Index (SPEI). We calculated the drought period at a 14-day scale with the SPEI R package (Vicente-Serrano et al., 2010; Beguería et al., 2014). The Penman-Monteith equation (Monteith, 1965) was used to calculate the Potential Evapotranspiration (PET) with the FAO-56 method (Allen et al., 1998). The altitude was fixed at 10 m and the latitude at $60^{\circ}11'$ N.

The minimum temperature, maximum temperature, wind speed, cloud amount, dew point temperature, relative humidity and air pressure, all collected at an hourly scale at the Kumpula meteorological station and averaged at a daily level, were downloaded from the Finnish Meteorological Database (Finnish Meteorological Institute, 2023b).

The drought periods were defined weekly with SPEI under -1.5 , corresponding to the highest value describing a severe drought (Wang et al., 2021): a week with at least one daily SPEI value under -1.5 was considered a drought week.

2.7 Statistical analyses

2.7.1 Statistical analyses regarding drought stress

With the collected data, the resilience of the different grassland vegetation types to drought events was estimated. As resilience is the result of both components – resistance and recovery – we studied in this paper only the resistance, which was defined as “the magnitude of disturbance that a system can absorb before shifting from one state to another” (Capdevila et al., 2021).

Thus, to estimate the direct impact of drought on lawns and meadows, we chose to primarily compare the old mesic meadow (KMP meadow) and the non-irrigated lawn (VKI lawn), since both of these vegetation types were older than 15 years and non-irrigated. In addition, we also included the irrigated lawn (KMP lawn) in the comparison to see the effect of irrigation. The yearly summer means of TER, daily GPP, green cover, soil temperature and soil moisture were calculated for each site, by considering that the summer season starts on week number 22 and ends on week number 34 (data collected from May 31st–August 24th 2021 and June 3rd–August 25th 2022). Lastly, we calculated the differences between the measured values of the variable and the summer average of the same variable at the site level. These calculated differences were then normalized at a site and year level by subtracting the annual means of the differences and dividing the whole by the annual standard deviation of the differences.

With the summer data, we defined two categories: data collected during the drought events and data collected outside of the drought events. Finally, with the Mann-Whitney U tests, we compared, pair by pair, VKI lawn, KMP lawn and KMP meadow in the different summer conditions.

2.7.2 Dynamics of GHG fluxes at the satellite sites

Considering the transformation from lawns into meadows at the satellite sites, Shapiro-Wilk tests were used to evaluate the normal distribution of each variable (i.e., CO_2 fluxes, CH_4 fluxes, NO_2 fluxes, soil moisture and soil temperature) by year and treatment (meadow vs. lawn). Then, differences in given variables between lawns and meadows were assessed either by t-tests or Mann-Whitney U tests depending on the normality of the data.

405 2.7.3 Statistical analyses ~~of~~ regarding plant variable contribution functional types

The effects of plant functional types on SOC, SON and GHG fluxes were studied in all intensive and satellite sites with the ~~plant~~vegetation and fluxes data collected in 2022. The fluxes were averaged for the growing season, from May to September. First, Mann-Whitney U tests were used to compare the proportions of grasses, legumes and forbs between lawns and meadows. Then, the effects of vegetation ~~functional types~~ were assessed by applying linear mixed-effects models (LMMs), using the R lme4 package (Bates et al., 2015), where a factor describing the locations of the lawn/meadow pairs (Fig. 1) was included as a random effect. This factor was introduced to consider the spatial variability between the sites, as the environmental conditions of the lawn and meadow within each site were considered to be similar. If predictive variables, i.e., plant functional types, correlated over 0.70, one of the correlated ~~variables~~plant functional types was discarded from the LMM according to the lowest cover proportion and the lowest occurrence in the dataset. Next, the normal distribution of each regression's residuals was visually checked with Quantile-Quantile (Q-Q) plots. In total, 6 mixed-effect multiple regressions including the proportions of the selected plant functional types as predictive variables were run by applying a backward selection. At each iteration, predictive variables with p-values higher than 0.1 were removed and variables with p-values lower than 0.05 were considered ~~to be~~ statistically significant. P-values were adjusted by a Kenward-Roger approach (Kenward and Roger, 1997) from R lmerTest package (Kuznetsova et al., 2017) and a conditional R² was used to assess the quality of the model.


420 3 Results

3.1 Drought occurrences

There was 7% and 12% more precipitation in 2021 and 2022 compared to the 653-mm value of the reference period 1991–2020 (Jokinen et al., 2021), but in both years July was drier than the long-term average, with 17% and 15% less precipitation in 2021 and 2022 respectively (47 mm in 2021 and 49 mm in 2022 compared to 57 mm in 1991–2020). At the same time, 2021 and 2022 were 0.1 and 0.4 °C warmer than the reference which was 6.5 °C. The summers were on average hotter than the reference period with 2.5 °C and 1.8 °C higher temperatures in 2021 and 2022, respectively, compared to 16.6 °C. These particularly hot and dry weeks were observed from June 14th to August 1st, 2021, and from June 6th to July 17th, 2022, (Fig. 2a).

At the intensive sites, the measured soil moisture (Fig. 2b) also showed a drastic decrease at all sites during these two periods, except at the irrigated lawn (KMP lawn), where the soil moisture remained relatively high during the whole summer. The highest soil temperature was also reached during the same period (Fig. 2c). In 2021 and 2022, those hot and dry weeks also corresponded to the decrease phase of LAI, except at the dry meadow (VKI meadow), where LAI increased slowly from spring until August (Fig. 2d). The measured green cover notably followed the same patterns as the satellite LAI data (Fig. 2e).

Using the SPEI, two drought periods were defined: the first one from June 14th to August 8th in 2021, and the second one from June 27th to July 17th in 2022 (Fig. 2). Thus, the defined drought event in 2021 happened early in the summer season and lasted eight weeks, whereas the drought event in 2022 arrived later in the summer and lasted only three weeks. These two drought events co-occurred with the high temperature and low soil moisture periods described above.



440 **Figure 2.** (a) Weekly precipitation (Prec.) and mean air temperature (Tav) recorded by FMI meteorology station, (b) manually
 445 measured soil moisture at a 5 cm depth, (c) manually measured soil temperature at a 10 cm depth, (d) leaf area index from satellite
 Sentinel-2 and (e) manually measured green cover at the intensive study sites with standard deviations calculated for each site at a
 daily scale. The irrigated lawn is KMP lawn, the non-irrigated lawn is VKI lawn, the mesic meadow is KMP meadow, and the dry
 meadow is VKI meadow. Red rectangles indicate the drought periods according to the SPEI and light grey rectangles represent the
 summer season. In panel (a), Tmax and Tmin represent the weekly instantaneous maximum and minimum mean temperatures.

3.2 CO₂ fluxes

3.2.1 Observed seasonal variation

At all intensive sites, the measured TER increased at the beginning of the measurement campaign and declined during the
 drought events at the non-irrigated sites (Fig. 3 and S1). When comparing the differences between the four sites, the mean
 450 momentary TER was the highest at the mesic meadow (KMP meadow, 0.44 and 0.31 mg CO₂ m⁻² s⁻¹ in 2021 and 2022,
 respectively) and the lowest at the dry meadow (0.29 and 0.26 mg CO₂ m⁻² s⁻¹ in 2021 and 2022, respectively). In 2021, the
 highest recorded mean values (0.86, 0.84, 0.59, 0.59 mg CO₂ m⁻² s⁻¹) were measured at the non-irrigated lawn (VKI lawn),

mesic meadow, dry meadow, irrigated lawn, on June 7th, July 2nd, July 19th, August 11th, respectively. In 2022, the highest values (0.71, 0.49, 0.65, 0.55 mg CO₂ m⁻² s⁻¹) were recorded at the dry meadow, irrigated lawn, mesic meadow, non-irrigated lawn on June 23rd, June 30th, July 1st, August 17th, respectively. In general, daily TER were higher at KMP sites (irrigated lawn and mesic meadow) compared with VKI sites (non-irrigated lawn and dry meadow), although the highest values were not reached in a specific order.

There were site-specific variations in the seasonal pattern of GPP (Fig. S1). In the first campaign year, after the first measurements, the daily GPP sink decreased at all sites except the dry meadow, for which it was the first growing season after establishment. In 2022, the daily GPP sinks first increased and then started to decrease around the end of June (Fig. 3). ~~The~~In 2021, the highest recorded daily uptakes ($\text{=(-0.48, =-0.51, =-0.40, =-0.46 mg CO}_2 \text{ m}^{-2} \text{ s}^{-1})$) were measured at the non-irrigated lawn, mesic meadow, dry meadow, irrigated lawn, on June 7th, June 18th, August 5th, August 11th, respectively, ~~in 2021~~. In 2022, the order had also slightly changed as the daily highest uptakes ($\text{=(-0.37, =-0.58, =-0.42 and =-0.47 mg CO}_2 \text{ m}^{-2} \text{ s}^{-1})$) were recorded at the non-irrigated lawn, mesic meadow, irrigated lawn, dry meadow on June 10th, June 13th, June 13th and June 23rd. Thus, the measured GPP values were higher at KMP sites (irrigated lawn and mesic meadow) than at VKI sites (non-irrigated lawn and dry meadow).

The mean daily GPP rates during May-August 2021–2022 were $-24.4, -16.7, -23.9, \text{ and } -8.7 \text{ g CO}_2 \text{ m}^{-2} \text{ day}^{-1}$ at the irrigated lawn, non-irrigated lawn, mesic meadow and dry meadow, respectively, when comparing only the consecutive weeks that included high-quality measurements at all sites. The mean observed respiration rates were 0.39, 0.40, 0.51, and 0.35 g CO₂ m⁻² s⁻¹, respectively, during the same months.

3.2.2 JSBACH performance

In order to represent the average fluxes of the different habitats, the model was primarily initialized to reach a steady state with each vegetation type and therefore, the simulated soil carbon pool represents the result of long-term carbon input of the site itself. The simulated soil moisture (Fig. S2) varied according to irrigation, precipitation and evaporation and showed lower values in summer, particularly in June and July, and also in August in 2022. The modelled seasonal dynamics in soil moisture followed the observations in general; however, some of the observations were considered to represent moisture content above the volumetric field capacity. The maximum soil moisture in the model is limited by the field capacity (Fig. S2). At the same time, the modelled soil temperature between 10 and 15 cm depth agreed with the measured soil temperature at 10 cm depth (Fig. S3). Modelled LAI varied in accordance with the observations especially at the mesic meadow and in the second year at the dry meadow (Fig. S4). The dry meadow was established very recently and was found to be sparse and inhomogeneous with respect to the vegetation, and therefore the drought response was challenging to capture in the model simulation, particularly in 2021. The drought response for the non-irrigated lawn agreed with the observations in 2021, while in 2022 the recovery was too strong in the simulation. At the irrigated lawn, the simulated LAI decreased later in 2022 than the observed LAI (Fig. S4).

Simulated TER followed the dynamics in the observed seasonal cycle and, for example, showed a similar response to the 2022 drought but generally underestimated the overall level of the non-irrigated lawn and the dry meadow. The drought response was stronger in the mesic and dry meadow (Fig. 3e-h, Fig. S5e-h). For the irrigated lawn the benefits of irrigation may have been overestimated, seen as a weaker drought response in the model than in the observations. The irrigation used in the model was estimated as an average over the whole area, while there probably was less irrigation where the measurement equipment was installed. The highest R² values for TER were observed at the irrigated lawn and mesic meadow and the lowest at the dry meadow (Table 2). Adjusting the carbon pools to fit closer to observed TER values in 2021 and 2022 (Fig. S6-7) mainly decreased RMSE but did not notably improve R² values at the different vegetation types in 2021 and 2022 (Table 2).

The model was able to simulate the overall level and the seasonal cycle in GPP at the irrigated lawn, non-irrigated lawn, and mesic meadow (Fig. 3a-c, Fig. S5a-c, Table S5). At the dry meadow, the model overestimated the increase in GPP in the early


495 season during its first growing season in 2021 (Fig. S5d). The vegetation cover in the newly established meadow was very low in the early growth season in 2021, which was not accounted for in the model. The agreement was better in the following year (Fig. 3d, Table S5). The benefit of irrigation was also somewhat overestimated for GPP during the dry periods. Simulated daily mean GPP followed the observations closely for non-irrigated lawn and mesic meadow (R^2 between 0.41 and 0.85, Table 2), while the dry meadow was difficult to simulate especially in 2021 (Table 2, Fig. S5).

500 3.2.3 Annual balances

During 2005–2021, simulated annual GPP and TER were on average the highest at the irrigated lawn and mesic meadow (Table 3). GPP and TER were the lowest in the dry meadow (Table 3) being approximately 40% of those of the irrigated lawn and meadow. The non-irrigated lawn's GPP and TER were approximately 70% of those of the irrigated lawn. The lawns had, on average, more negative NEE i.e., a greater sink of carbon, than meadows, and looking at the ratio between standard deviation and NEE, meadows showed higher year-to-year variations than the lawns (Table 3). The sink of the irrigated lawn was approximately 53% higher than that of the non-irrigated lawn (VKI lawn). Additionally, it was found that 13–16 % of TER occurred between November and March.

505 According to the simulations that were adjusted to meet the TER observations, annual NEE were 328, 2663, 1476 and 2140 g $\text{CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ in 2021 and -195, 1496, 571 and 1205 g $\text{CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ in 2022, at the irrigated lawn, the non-irrigated lawn, the mesic meadow and the dry meadow respectively (Table S4).

510



515 **Figure 3. Seasonal dynamics of mean total ecosystem respiration (TER, abcd) and daily photosynthesis (GPP, efgh) in the four intensive sites in 2022. Grey continuous lines represent JSBACH simulations and triangles and dots represent the mean of manual measurements with standard deviation bars. (a, e) Irrigated lawn = KMP lawn; (b, f) non-irrigated lawn = VKI lawn; (c, g) mesic meadow = KMP meadow; (d, h) dry meadow = VKI meadow. Red rectangles indicate the drought periods according to the SPEI and light grey rectangles represent the summer season. The model simulated the whole year but for clarity, Jan-March and Dec are not visible as there were no measurements during those months.**

520 **Table 2. R² and root mean square error (RMSE, mg CO₂ m⁻² s⁻¹) calculated for total ecosystem respiration (TER) and gross primary production (GPP) in the standard simulation where the soil carbon was stabilized based on the standing vegetation, and for the adjusted simulations (adj soil c) where the soil carbon pool was set so that simulated TER met the observations in 2021 and 2022.**


Year	Site	TER				GPP			
		standard		adj soil c		standard		adj soil c	
		R ²	RMSE						
2021	irrigated lawn	0.86	0.07	0.84	0.07	0.17	0.13	0.15	0.11
	non-irrigated lawn	0.61	0.24	0.47	0.18	0.41	0.10	0.42	0.10
	mesic meadow	0.80	0.23	0.79	0.19	0.85	0.12	0.87	0.10
	dry meadow	0.15	0.20	0.04	0.19	0.02	0.13	0.01	0.13
2022	irrigated lawn	0.62	0.09	0.68	0.08	0.38	0.13	0.39	0.11
	non-irrigated lawn	0.61	0.13	0.53	0.10	0.65	0.08	0.66	0.09
	mesic meadow	0.71	0.12	0.67	0.11	0.85	0.12	0.86	0.10
	dry meadow	0.23	0.20	0.26	0.16	0.49	0.11	0.49	0.11

525 **Table 3. Mean annual (± standard deviations calculated for 2005–2022) of ecosystem respiration (TER), photosynthetic uptake (GPP) and net ecosystem exchange (NEE) modeled with JSBACH at the four intensive sites during the years 2005–2022.**

Sites	Characteristics	TER (g CO ₂ m ⁻² yr ⁻¹)	GPP (g CO ₂ m ⁻² yr ⁻¹)	NEE (g CO ₂ m ⁻² yr ⁻¹)
KMP lawn	Irrigated lawn	4331 (± 177)	-4667 (± 218)	-336 (± 187)
VKI lawn	Non-irrigated lawn	3039 (± 251)	-3196 (± 231)	-157 (± 139)
KMP meadow	Mesic meadow	4127 (± 291)	-4278 (± 233)	-151 (± 198)
VKI meadow	Dry meadow	1765 (± 136)	-1828 (± 161)	-63 (± 134)

3.3 Extreme weather resistance

530 Green cover, TER and daily GPP were significantly decreased at the non-irrigated lawn compared to the records outside the drought and to the mature mesic meadow and the irrigated lawn during the prolonged drought in 2021 (Fig. 4ace) and during the shorter drought in 2022 (Fig. 4bdf). In 2022, TER was significantly higher at the irrigated lawn and the mesic meadow during the drought than outside of the drought period (Fig. 4d).




535

Figure 4. The resistance indices of an irrigated lawn (KMP lawn), a non-irrigated lawn (VKI lawn) and an old mesic urban meadow (KMP meadow) to drought events in 2021 (first column) and 2022 (second column), calculated for the green cover (a, b), the TER (c, d) and the daily GPP converted to positive values (e, f). The values are the differences between the measured values and the summer average, normalized at a yearly level. Summer includes Jun–Aug and droughts were defined in Fig. 2 (June 14th–August 8th, 2021 and June 27th–July 17th, 2022). (□ for p-value ≤ 0.10; * for p-value ≤ 0.05; ** for p-value ≤ 0.01; and *** for p-value ≤ 0.001).

540

3.4 CH₄ and N₂O fluxes

545



550 **Figure 5. Box and whisker plots of measured (a) total ecosystem respiration (TER), (b) CH₄ and (c) N₂O fluxes of lawns and transformed meadows at the satellite sites before (grey background) and after transformation (white background), which happened at the end of 2020 only at the transformed meadows. The environmental conditions were considered to be similar between the two treatments. Statistical differences between both types of green spaces were tested at the year level (α for p-value ≤ 0.10; and * for p-value ≤ 0.05).**

555 The measurements at the control lawn and transformed meadow of each site were conducted consecutively during the same day, and therefore the environmental conditions were ~~considered to be~~ as similar as possible. Indeed, soil moisture and temperature did not differ between the treatments during the measurements (Fig. S9). At the satellite sites, before the transformation in 2020, there were no differences between lawn and meadow sites in terms of TER, CH₄ and N₂O fluxes measured in darkened conditions (Fig. 5). There was some general year-to-year variation in the median and range of observed CO₂ and CH₄ fluxes, but the fluxes did not differ between lawns and transformed meadows in 2021 or 2022 (p > 0.1, Fig. 5ab). The mean N₂O fluxes were slightly lower in meadows than in lawns in 2021 and 2022 (p-value < 0.1, Fig. 5c).

560 3.5 Plant functional type predictor variables for C and N cycles

565 On average, the total vegetated area of the studied meadows (Table 4) did not significantly differ from those on lawns (U = 40.000, p-value = 0.442), but the cover proportions of the main plant functional types shared by both vegetation types differed. Meadows had a significantly lower proportion of grassesgrass cover (i.e., 25%) than lawns (65%, U = 10.000, p-value = 0.002). Most of the meadow area was covered by forbs other than legumes (i.e., 65%), a proportion significantly larger (U = 16.000, p-value = 0.002) than that found in lawns (i.e., 19%). Nevertheless, the proportion of legumeslegume cover was found to be statistically similar (U = 39 000, p-value = 0.505) in meadows (11%) and lawns (6%).

570 Next, we explored possible correlations between cover proportions of common plant functional types and variables related to C and N cycles. First, we discarded sedges from the analysis since it their cover proportion was correlated with that of horsetails (> 0.7, Table S7) and were only present at one site and in a low proportion (Table 3). ~~None~~ The cover proportions of none of the functional plant types are significantly associated to TER, nor SON. Yet, CH₄ flux appeared to be positively associated with moss proportioncover (Table 5), and. Additionally, N₂O flux was significantly negatively associated to with forb proportioncover and positively to associated with horsetail cover proportion (Table 5). SOC was slightly positively explained by forb cover proportion (Table 5). The C/N ratio was significantly negatively associated with legume, grass and forb cover proportions (Table 5).

575

580

Table 4. Cover proportions (%) of the different plant functional types: grasses (*Poaceae*), legumes (*Fabaceae*), forbs (other families of flowering vascular plants, which do not belong to one of the listed categories), trees, sedges (*Carex*), horsetails (*Equisetum*), and mosses (*Bryophyta*), inventoried on June 27th and 21st July 2022. The “total” column is the sum of all the plant functional type cover, a fully covered quadrat with only one layer of vegetation should have a 100% cover, >100% indicates layered vegetation (short and tall grassland plants) and <100% indicates the presence of bare soil.

Experiment	Location	Management	Grasses	Legumes	Forbs	Trees	Sedges	Horsetails	Mosses	Total
Intensive	KMP	lawn	70.5	6.0	21.5	0.0	0.0	0.0	0.0	98.0
		meadow	30.0	16.5	70.3	0.0	0.0	0.0	0.0	116.8
	VKI	lawn	92.3	0.3	7.5	0.0	0.0	0.0	0.0	100.0
		meadow	6.3	38.3	23.3	0.0	0.0	0.0	0.0	67.8
Satellite	JMT-3	lawn	78.8	6.6	10.1	0.0	0.0	0.0	0.0	95.4
		meadow	6.5	9.1	42.7	0.4	0.0	0.0	0.0	58.8
	JMT-7	lawn	21.3	20.6	37.5	0.0	0.0	0.0	0.0	79.4
		meadow	31.9	13.0	68.3	0.0	0.0	0.0	0.0	113.1
	KO-4	lawn	78.8	3.9	15.1	0.0	0.0	0.3	0.0	98.0
		meadow	10.0	0.4	53.0	0.0	0.0	0.0	0.0	63.4
	PK-3	lawn	39.8	0.8	10.2	0.2	0.0	0.0	38.6	89.5
		meadow	23.8	7.0	104.8	0.0	0.5	30.0	0.0	166.0
	SK-1	lawn	65.0	9.7	24.3	0.0	0.0	0.0	0.0	99.0
		meadow	81.3	2.0	52.8	0.0	0.0	0.0	0.0	136.0
	SMT-12	lawn	66.3	2.3	27.8	1.3	0.0	0.0	0.0	97.7
		meadow	15.8	1.8	106.0	0.3	0.0	0.0	0.0	123.8

585

Table 5. The connection between the cover proportions (%) of different plant functional types and total ecosystem respiration of CO₂ (TER), CH₄ fluxes, N₂O fluxes, Soil C content (SOC), Soil N content (SON) and C/N ratio. The plant functional types are grasses (*Poaceae*), legumes (*Fabaceae*), forbs (other families of flowering vascular plants, which do not belong to one of the listed categories), trees, sedges (*Carex*), horsetails (*Equisetum*), and mosses (*Bryophyta*). (NS for p-value > 0.10; □ for p-value ≤ 0.10; * for p-value ≤ 0.05; ** for p-value ≤ 0.01; and *** for p-value ≤ 0.001). ^{sat.} indicates values that were only measured at satellite sites.

Response variables	Predictive variables						R ²
	Grasses	Legumes	Forbs	Trees	Horsetails	Mosses	
TER (mg CO ₂ m ⁻² s ⁻¹)	NS	NS	NS	NS	NS	NS	/
CH ₄ fluxes (ng CH ₄ m ⁻² s ⁻¹) ^{sat.}	NS	NS	NS	NS	NS	2.698 □	0.63
N ₂ O fluxes (ng N ₂ O m ⁻² s ⁻¹) ^{sat.}	NS	NS	-0.279 **	NS	1.092 **	NS	0.98
SOC (% , < 2 mm)	NS	NS	0.011 □	NS	NS	NS	0.56
SON (% , < 2 mm)	NS	NS	NS	NS	NS	NS	/
C/N ratio	-0.159 ***	-0.281 **	-0.079 *	NS	NS	NS	0.69

590 4 Discussion

It is well known that converting lawns into meadows can increase biodiversity the presence of different plants and arthropods in cities (Venn and Kotze, 2014; Wastian et al., 2016; Chollet et al., 2018). In this study, we wanted to understand better how transforming lawns into meadows in Nordic urban areas could affect their GHG fluxes and how varying cover proportions of their different plant functional types could influence the C and N cycles of these green spaces. The studied sites, an irrigated lawn, a non-irrigated lawn, an old mesic spontaneous meadow and a young dry meadow, were mainly C sinks. Moreover, with a focus on the transformation dynamic from lawns into meadows, no significant differences in measured GHG fluxes (i.e., CO₂, CH₄ and N₂O) were found between the six control lawns and six transformed meadows in the two following years after the conversion. However, the studied mesic meadow appeared to be more resistant to drought stress than an unirrigated lawn. This resistance could notably be explained by plant diversity a shift in vegetation functional types and a less homogeneous vegetation setting, as discussed later. Furthermore, as expected, the forbs cover proportion of forbs was found to be higher in

600

meadows than in lawns and seems to be negatively associated with N₂O fluxes and C/N ratio, and positively associated to SOC of the green space.

4.1 Fluxes in urban grasslands

Carbon neutrality, or even a net positive scenario, is nowadays one of the major goals for cities and states in mitigating climate change (IPCC, 2022). Our aim was to understand how CO₂ sequestration differs between lawns and meadows by studying contrasting vegetation types, a mesic mesotrophic meadow and a dry nutrient-poor meadow together with irrigated and non-irrigated lawns, to determine the full range of their GHG exchange. We found that the photosynthetic production (GPP) of an irrigated lawn and a mesic meadow were roughly equal on a mean daily and an annual scale whereas the GPP of a non-irrigated lawn was notably lower. At an annual level, irrigation increased the GPP of a lawn by over 40% and the sink (NEE) by more than 100%. It has already been found that water input improves carbon uptakes (Thienelt and Anderson, 2021) and demonstrated by Zirkle et al. (2011) that irrigated lawns store up to 10 g C m⁻² yr⁻¹ more carbon in soil than non-irrigated ones. Also, our analysis ~~on~~of annual NEE indicated that an irrigated lawn is a stronger sink, approximately 50 g C m⁻² yr⁻¹ higher than a non-irrigated lawn.

Here, the model estimates for NEE ~~had some uncertainties were uncertain~~ as the model was unable to simulate some of the observed momentary total ecosystem respiration (TER) values during summertime (Fig. 3)), which caused the full carbon balance estimate to be more uncertain than the estimated GPP. The uncertainty of the annual TER and NEE estimate is further highlighted by the mean observed TER between May and August, where the order of the sites differed from the ones in simulated annual TER values. However, the momentary TER varies based especially on soil moisture, temperature, and autotrophic activity (Ryan and Law, 2005; Subke et al., 2006; Mäki et al. 2022) and therefore, making a mean without standardizing the environmental conditions might be inaccurate, as the momentary measurements at the satellite sites were collected during different times i.e. environmental conditions.

In this experiment, we measured the collars at various times of day ~~due to~~for practical reasons, even though Pavelka et al. (2018) suggest measuring ecosystem respiration in the morning around 9–10 to get the average level of daily respiration. Moreover, with manual chamber measurements, we were able to measure only once or twice a month and most of the time under ~~a~~sunny weather, whereas with automatic chambers or eddy covariance (Hiller et al., 2011; Thienelt and Anderson, 2021), it would have been possible to estimate the daily and monthly variations at a finer scale and to reduce the bias caused by hot and dry days. Therefore, especially during the summertime, the measured values most probably are overestimations of the daily average whereas in the early spring and in autumn, when the diurnal cycle in soil temperature and plant activity are smaller, the observations fit the model estimate closely. Decina et al. (2016) found that during the growing season, TER was about 0.198 ± 0.006 mg CO₂ m⁻² s⁻¹ at urban lawns in Boston, USA, which is lower than the values measured at our sites. Last, it has been shown that during cold days, under snow and/or frozen ground, grasslands are a source of CO₂ (Hiller et al., 2011; Jasek-Kamińska et al., 2020). The model was also run for winter months with the heterotrophic soil respiration decreasing at the same time as the soil temperature until reaching a minimum in late winter. Although the highest ecosystem activities and emissions take place in the warm summer months, to fully validate the annual balance, it would be useful to have some additional wintertime observations. Nevertheless, we can be quite confident in the simulated photosynthetic production, since GPP principally occurs during the snow-free seasons.

Furthermore, soil respiration is highly related to soil carbon quality and quantity (Davidson and Janssens, 2006). Here, it seems evident that the studied soil was not stabilized but, rather, that growing media with unknown properties was either brought to the site or some earlier vegetation type built the soil carbon storage. This is most evidentdistinct in the young sites (less than 15 years old) and most managed sites i.e., the lawns and the dry meadow. It highlights unpredictable features of urban soils which are characterized by high anthropogenic disturbances, unknown origins and changes in land use. Thus, the quality and quantity of organic matter cannot be connected to a linear history even at a city scale (Pouyat et al., 2006; Setälä et al., 2016;

Ivashchenko et al., 2019; Sushko et al., 2019; Cambou et al., 2021). Here, we chose to estimate the carbon balance over an extended time in stabilized conditions and all our sites were estimated to be small sinks of carbon varying between 63–336 g CO₂ m⁻² yr⁻¹. However, adjusting the model parameters to reproduce values closer to the observed TER values turned most of the sites ~~into~~ sources of atmospheric CO₂ during the measurement years 2021 and 2022. Even if those values represent the current situation at the study sites, those represent also an unstabilized state. ~~Whereas~~ whereas the extended runs stand for long-term carbon balances of these vegetation types.

Many have reported urban grasslands to be C sources at least in certain conditions (Allaire et al., 2008; Hiller et al., 2011; Bezyk et al., 2018), but some have also reported sinks of about 20–180 g CO₂ m⁻² yr⁻¹ (Thienelt and Anderson, 2021). However, most of the studies focus on the respiration rate, omitting GPP (Kaye et al., 2005; Decina et al., 2016; Lerman and Contosta, 2019; Sushko et al., 2019; Upadhyay et al., 2021). Jasek et al. (2020) found that the annual mean of TER for a mix of different types of urban grassland is 424 ± 43 g C m⁻² yr⁻¹ in Krakow, Poland, and Kaye et al. (2005) measured that irrigated lawns in Northern Colorado, USA, emits around 2777 ± 273 g C m⁻² yr⁻¹, where our annual TER values vary between 1181 ± 48 g C m⁻² yr⁻¹ at the irrigated lawn and 481 ± 37 g C m⁻² yr⁻¹ at the dry meadow. In the end, we found that lawns were stronger sinks than the meadows which is in line with Peoplau et al. (2016) who found that soil C was higher in lawns than in meadows, which was mainly attributed to the clippings left at the site and fertilization.

It is noteworthy that lawns in Finland are well maintained to support soil fertility (Viherympäristöliitto, 2023), whereas meadows grow on various types of soil ranging from poor and dry to mesic and fertile. We found that the mesic meadow, with high (max. 1.3 m) and dense vegetation in summer, had higher photosynthetic production and annual sink than the dry meadow sown in late 2020. However, during the campaign that took place 0–2 years after the establishment, the dry meadow was still in an initial phase i.e., its carbon sequestration potential may not have been fully realized, with about six-times lower CO₂ uptake than the mesic meadow. In addition, in the same meadow, there were dissimilar features among the four sampled quadrats: one of them was scarce in plants, another one had a tall and dense vegetation, and the two others had a short dense vegetation, which contributed to making the ecosystem model simulations and the analysis even more challenging. For these reasons, it would be useful in the future to compare homogeneous meadows with varying fertility – from mesic mesotrophic to dry nutrient-poor, but comparable age.

Recent research has demonstrated that urban grasslands can act as both sources and sinks for CH₄ and N₂O, with the magnitude and direction of fluxes being dependent on factors such as soil temperature, moisture, and management such as irrigation or fertilization (Livesley et al., 2010; Law et al. 2021; Künnemann et al. 2023; Zhan et al., 2023). Bezyk et al. (2018) measured urban herbaceous areas and found the sink to be higher (about -5 ng CH₄ m⁻² s⁻¹) than our mean values (-2.28 ± 36.51 ng CH₄ m⁻² s⁻¹) where the variation between collars and dates was high. The measured N₂O fluxes in this study were low and the mean was even slightly negative (-0.60 ± 12.43 ng N₂O m⁻² s⁻¹). Chapuis-Lardy et al. (2007) reported also small N₂O sinks for temperate grasslands varying between -0.2 ng N₂O m⁻² s⁻¹ in a grass mixture in Canada to -109 ng N₂O m⁻² s⁻¹ in an artificial grass-clover mixture in Switzerland. However, most of the reported values from urban grasslands are positive, the mean ranging mainly between 2.9 to 14.1 ng N m⁻² s⁻¹ (Livesley et al., 2010; Gillette et al. 2016; Law et al. 2021; Künnemann et al. 2023). Zhan et al. (2023) reported that on average, biogenic N₂O emissions from soils in various urban environments were 9.5 ng N m⁻² s⁻¹ (median: 5.7 ng N m⁻² s⁻¹), rates which are more than double compared with non-urban environments.

With the satellite sites, we focused on the transformation process and found no negative climate impacts in terms of greenhouse gases during the campaign which covered two growing seasons after the transformation. ~~However, it is possible that we missed some~~ Zhan et al. (2023) synthesized that the change from a natural grassland to a lawn notably increases annual N₂O emissions and decreases CH₄ uptake, yet there was just one study available on CH₄ uptake on such land use change. The opposite change in land management in this study decreased N₂O emissions (p<0.1) but did not notably affect CH₄ fluxes. Those are the balance between methanogenic and methanotrophic activity by microorganisms (Conrad, 1996) and therefore, soil community plays a key role in the CH₄ balance. Yet, possible changes in the community due to the conversion from lawns to meadows were not

reflected in CH₄ fluxes, at least not during the two years studied after conversion. However, it is possible that we missed some momentary sinks or peaks of emissions as the measuring frequency was just once a month. Nonetheless, it is quite safe to conclude that even if such unmonitored peaks occurred, their significance on an annual level would be minor as there was no indication of any differences between the treatments during the measurements. On the other hand, it is evident that destroying vegetation during the transformation process decreases photosynthetic input at least during the following autumn and spring when lawns continue photosynthetic activities but when the new meadow has not developed yet. Therefore, it would have been interesting to measure GPP as well and conduct the experiment over a longer period of time, as it has been shown that even grassland restoration requires many more years (Muller et al., 1998; Waldén and Lindborg, 2016; Kose et al., 2021). In addition, the CO₂, H₂O, N₂O and CH₄ analyzer used at the satellite sites has a low sensitivity for small fluxes, at least in certain conditions (Kohl et al., 2019), in comparison with more advanced analyzers. However, we were only interested in the possible difference between the two treatments and not the actual rates nor annual balances and therefore, we find the choice to be acceptable. In any case, it would be important to study the CH₄ exchange, N cycle and particularly N₂O fluxes in northern urban grasslands in more detail. In addition, the applied transformation process was the same at all the satellite sites and it would be interesting to also study the impact of different transformation processes on GHG fluxes.

Finally, even though the spectra of our four intensive sites was wide, the four sites were unique in their environmental conditions, vegetation, and management. In order to provide a stronger categorization, it would have been beneficial to study more replicates intensively. Moreover, mowing frequency was not studied, nor its carbon footprint, although studies have found notable impacts on source and sink due to managementsmanagement (Allaire et al., 2008; Poeplau et al., 2016; Lerman and Contosta, 2019; Thienelt and Anderson, 2021) leaving room for improvements for further studies.

4.2 CO₂ fluxes impacted by drought events

Extreme climateweather events may occur more often in the coming years and during the two years of measurements in this study, dry summers impacted the ecosystems. Here, we focus only on the resistance component, i.e. the capacity of the system to absorb the drought disturbance during the event (Vogel et al., 2012; Capdevila et al., 2021). According to the comparison between drought and the outside-of-drought periods, the unirrigated lawn reached a more desiccated stage during the period of the summer, whereas the meadows seemed to endure the deficit of water and the hot weather as well as the irrigated lawn. In unirrigated lawns, such a suppression is a physiological reaction to drought stress reducing photosynthesis (GPP) and respiration (TER), as also found in other studies (Allaire et al., 2008; Hiller et al., 2011; Vogel et al., 2012). Yet, it has been also demonstrated in several studies that species-rich grasslands better endure extreme weather events (Vogel et al., 2012; De Keersmaecker et al., 2016).

The greater resistance of meadows compared to non-irrigated lawns could be explained by several factors, such as plant complementarity, microclimate and management (Vogel et al., 2012; De Keersmaecker et al., 2016; Bernath-Plaisted et al., 2023). (1) Lawns are usually sown, weeded and managed in order to maintain the desired low species composition and aesthetic appeal; on the other hand, plant diversity is usually much richer in meadows. Yet, the *sampling effect* theory (Tilman et al., 1997; Loreau and Hector, 2001) argues that in a more diverse grassland, there is more chance to find at least one species, which is well adapted to drier habitats. For this reason, there is a higher chance for at least one plant species to survive during drought events in meadows than in lawns. (2) The second theory is *niche complementarity*, which results from niche differentiations and the benefits of interspecific interactions between species (Tilman et al., 1997), and induces a better individual species performance (Tilman et al., 1997; Loreau and Hector, 2001). Such a high diversity of plants and niches, as we can find in meadows, leads to a better nutrient, light and water-use efficiency (De Boeck et al., 2006; Walde et al., 2021), and helps to face water shortage. In relation to (1) and (2), meadows usually contain vegetation with deeper root systems, which allow plant individuals to supply themselves with water and reduce competition for the available water in the upper soil layers during drought periods. (3) Moreover, this complementarity in meadows is reinforced by microclimate anomalies

(Bernath-Plaisted et al., 2023). The tall and dense vegetation of meadows creates a buffer to face climate extreme events by offering shade to the lower layers and keeping the humidity beneath the meadow canopy. (4) Finally, management, and notably mowing frequency, is also described as one of the factors impacting drought resistance. Regrowth of vegetation is more sensitive to extreme weather stress than vegetation at a later stage of its growth dynamics (Vogel et al., 2012). Hence, lawns are more affected, since they are typically mown every two to three weeks during the growing season, whereas meadows are cut one to two times a year. Thus, vegetated urban meadows are able to function almost as well as irrigated lawns and as well as during regular summertime weather since, e.g., during and outside drought periods, the green cover and the photosynthetic uptake are comparable.

In this study, we did not have a reference year without a drought, as, according to daily SPEI calculation, both years of measurements were affected by extreme drought stress. Further, it would be important to understand how these ecosystems function during a normal climate year, especially to compare with a year similar as 2021 when the drought period covered 8 weeks out of 13 weeks of summer, even though the pre-drought reference and post-drought stable stages used in this study would better estimate the resistance itself (Capdevila et al., 2021).

4.3 Effect of plant functional types on C and N status

It is known that biodiversity, including its component associated with vegetation, increases drastically. Previous studies have already reported more heterogeneous and floristically rich green space settings after conversions of lawns into meadows (Venn and Kotze, 2014; Chollet et al., 2018; Norton et al., 2019). Yet, in our study we wanted to understand how the composition of vegetation, defined here in terms of cover proportions of its different plant functional types, could affect the SOC and the GHG fluxes at our sites. It has previously been found that settings in which vegetation comprises a larger number of plant species could explain a higher rate of TER (Dias et al., 2010) and could further enhance N and C storage in the soils (Fornara and Tilman, 2008; Oelmann et al., 2011; Mueller et al., 2013; Cong et al., 2014; Lange et al., 2015). Nevertheless, Wei et al. (2017) demonstrated that under scenarios in which vegetation was more diverse in its plant functional group types, the nitrate pool and the N mineralization rate were lowered. Additionally, Mueller et al. (2013) also found that higher diversity enhances N transformation and NH_4^+ pool, but a too high plant diversity could also have the opposite effect. Thus, in our study, we explored the possible connections that the different plant functional types, i.e. grasses, legumes, forbs, trees, sedges, horsetails and mosses, could have on the C and N cycles in our study sites. According to our analysis, CH_4 fluxes were connected to the cover proportion of hygrophytes such as mosses, even though the relation between soil moisture and CH_4 fluxes in urban grasslands seems to be hardly predictable (Groffman and Pouyat, 2009; Costa and Groffman, 2013; Bezyk et al., 2023). Although some of the fluxes were linked to the cover percentages of both horsetails and mosses, it would be beneficial to validate our results with a broader dataset since some plant functional types were scarcely represented: mosses and sedges were present at only one site, horsetails at two sites and tree saplings were surveyed at four sites out of the sixteen sites included in the study (Table 4). Nevertheless, our analysis shows that SOC increases with forb cover percentage, of which the proportion is higher in meadows. Hence, meadows could store more carbon than lawns because: (1) some SOC could have been incorporated into the soil due to the site transformation; (2) meadows could have produced more litter that enriched the soil in carbon; and (3) with a deeper root system, forbs could have allocated more carbon into the soil. Indeed, it has been shown that sites with woody plants, which also have deeper root systems, have a higher stock of carbon than lawns (Setälä et al., 2016; Lindén et al., 2020). However, such an increase in soil carbon was not supported by the flux analysis. One of the shortfalls in our study regarding C and N cycles and the proportion of plant functional types could be that it was studied with only a one-year dataset in order to include both the satellite and the intensive sites. We utilized SOC and SON pools in our analyses, even though their accumulation rate would have been a more interesting metric, as the standing pools also

770 reflect the historical land use of the sites. However, a three-year period would not have been long enough to observe changes in the pools due to their slow changing rate of less than 2% per year (Fornara and Tilman, 2008).

Some ~~other~~previous studies have ~~also~~ tried to estimate ~~the correlation~~correlations between plant functional ~~typetypes~~ and soil characteristics, by studying C3 and C4 grasses and legumes (Fornara and Tilman, 2008; Yang et al., 2019). Fornara and Tilman (2008) estimated that C4 grasses may increase soil carbon accumulation by 193%, and legumes by 522%. ~~Moreover, even though~~Although legumes did not appear to be significant in our SOC model, and negatively correlated to C/N ratio, legumes are known to improve the soil N availability and the litter quality by reducing the C/N ratio (Fornara and Tilman, 2008; Yang et al., 2019). Thus, ~~different~~ plant functional types undoubtedly impact the C and N cycles; however, the low representativity of some plant functional types and the low number of replicates limit any further interpretation of the robustness of our analyses. ~~To draw a~~Drawing stronger ~~conclusion~~conclusions on the impact ~~of that~~ the ~~cover of varying plant~~ functional types ~~have~~ on carbon and nitrogen dynamics, ~~it would be necessary~~ ~~quire~~ to study a larger number of sites and also compare chosen mixtures of plants ~~with known plant functional type cover proportions~~ on soils with the same properties.

5 Conclusions

~~In order to slow down biodiversity loss, cities~~Cities around the world are considering converting ~~their~~urban lawns into meadows. ~~Another priority is to~~ ~~to curb environmental degradation~~, mitigate climate change and achieve carbon neutrality. In this study, we focused on urban grasslands from a climate and vegetation perspective by studying maintained lawns in comparison with mature and newly transformed meadows in northern Europe at the border between the boreal and temperate zones. The ~~transformation~~transformation from lawns ~~to~~into meadows did not ~~imply~~cause any additional negative climate effects in terms of ecosystem respiration, CH₄ and N₂O fluxes. However, while meadows are known to ~~have higher species richness~~be more heterogenous in their vegetation and ~~diversity of~~ plant functional types ~~when compared to lawns~~, we found that meadows ~~did~~ not increase carbon sequestration compared with lawns on an annual scale, and ~~even that NEE is the sink~~ was usually ~~more negative~~greater in lawns than in meadows. Nevertheless, ~~in from~~ a ~~climate warming~~ perspective, ~~with of warming climate and~~ more frequent extreme events, ~~meadows resist~~the ability of urban grasslands to tolerate dry periods is an important feature. Meadows resisted drought events better than lawns and, ~~were~~ at least in the short term, ~~are~~ better able to ~~adapt cities to tolerate future~~ extreme drought events ~~induced by climate change and aid city-level adaptation~~. Nonetheless, it would be necessary to check this result with a larger number of sites and investigate the overall resilience with ~~a~~ longer time series dataset, to help stakeholders and city planners ~~to~~ make better decisions to optimize land-use. Thus, considering ~~biodiversity~~, climate warming and carbon neutrality, it is necessary to find the right trade-off between lawns and meadows in cities. Regarding ~~the~~ vegetation, forbs proportion was found to be higher in meadows and some plant functional type proportions, including ~~those of~~ grasses, legumes, forbs, horsetails and mosses, ~~are~~were correlated to specific C and N status. Nevertheless, as this analysis ~~was done with an only~~ ~~comprised~~ one-year ~~dataset, it, a link between functional plant types and~~ C and N cycles would be necessary to ~~investigate~~be investigated with ~~more~~a larger number of urban grasslands, ~~on the link between plants and C and N cycle~~. Moreover, it would be beneficial to further study the carbon storage process and its longevity, as well as ~~the biodiversity~~vegetation dynamics ~~during and the shifting of plant functional type proportions during~~ the transformation from lawns to meadows or drought resilience processes, and to consider the social aspects of the conversion of lawns to meadows.

805 Data availability

The data used in the manuscript can be downloaded from Finnish Meteorological Institute's data storage: <https://doi.org/10.23728/fmi-b2share.920c1e5f08a74a6d9dfcb3a08cfc6734> (Trémeau et al., 2023).

Code availability

The scripts used to calculate LI-COR and Gasmeter fluxes can be found at: <https://github.com/hvekuri/Chamber-codes/> (Vekuri, 810 2023).

Supplement link

Author contributions

LK, BO, EK and JT designed the research; BO, EK and JT participated in the data collection; HV, LK, EK and JT participated in the flux calculations; JT and BO conducted the statistical data analyses; LB performed the JSBACH simulations; all authors 815 participated in writing the paper.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This research takes part in the CO-CARBON project supported by the Strategic Research Council working under Academy of 820 Finland (grant no. 335204), in the CarboCity project supported by the Academy of Finland (grant no. 325549), in the ACCC Flagship program of the Academy of Finland (grant no. 337552), in the Lawns into Meadows project supported by Koneen säätiö (grant no. 201902880), and was also supported by the Maj and Tor Nessling foundation (grant no. 202000391).

We warmly thank Olivia Kuuri-Riutta, Pinja Rauhamäki and Elisa Vainio for their help with field data, Suvi Orttenvuori for her help with the preliminary JSBACH simulations, Elina Nieminen for supporting the research in its early phase, Olli 825 Nevalainen for the Sentinel-2 data, Joséphine Couet, Helena Rautakoski and Thomas Merrien for support in the statistical analyses, Mika Korkiakoski, Stéphanie Gerin and Helena Rautakoski for their help with the chamber measurement analysis, Jack Chapman [and Aarni Koiso-Kanttila](#) for the proofreading and valuable comments and finally, we also thank the reviewers for their supports. We also thank the Stara company, represented by Inkeri Salo, the City of Helsinki, represented by Tuuli Ylikotila, and the Finnish Museum of Natural History, represented by Mikael Lindholm, for providing crucial information on 830 the landscape management. Finally, we express gratitude to the City of Helsinki, the Finnish Museum of Natural History, AYY and HOAS for allowing us to transform and/or conduct our measurements on their properties.

References

Ahongshangbam, J., Kulmala, L., Soininen, J., Frühauf, Y., Karvinen, E., Salmon, Y., Lintunen, A., Karvonen, A., and Järvi, L.: Sap flow and leaf gas exchange response to a drought and heatwave in urban green spaces in a Nordic city, *Biogeosciences*, 20, 4455–4475, <https://doi.org/10.5194/bg-20-4455-2023>, 2023. 835

Allaire, S. E., Dufour-L'Arrivée, C., Lafond, J. A., Lalancette, R., and Brodeur, J.: Carbon dioxide emissions by urban turfgrass areas, *Can. J. Soil. Sci.*, 88, 529–532, <https://doi.org/10.4141/CJSS07043>, 2008.

Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration: guidelines for computing crop water requirements, edited by: Food and Agriculture Organization of the United Nations, Food and Agriculture Organization of the United Nations, Rome, 300 pp., 1998. 840

Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models Using lme4, *Journal of Statistical Software*, 67, 1–48, <https://doi.org/10.18637/jss.v067.i01>, 2015.

- 845 Beguería, S., Vicente-Serrano, S. M., Reig, F., and Latorre, B.: Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring, *International Journal of Climatology*, 34, 3001–3023, <https://doi.org/10.1002/joc.3887>, 2014.
- Belmeziti, A., Cherqui, F., and Kaufmann, B.: Improving the multi-functionality of urban green spaces: Relations between components of green spaces and urban services, *Sustainable Cities and Society*, 43, 1–10, <https://doi.org/10.1016/j.scs.2018.07.014>, 2018.
- 850 Bernath-Plaisted, J. S., Ribic, C. A., Hills, W. B., Townsend, P. A., and Zuckerberg, B.: Microclimate complexity in temperate grasslands: implications for conservation and management under climate change, *Environ. Res. Lett.*, 18, 064023, <https://doi.org/10.1088/1748-9326/acd4d3>, 2023.
- Bezyk, Y., Dorodnikov, M., Grzelka, A., and Nych, A.: Characteristics of temporal variability of urban ecosystem-atmosphere CO₂, CH₄, and N₂O fluxes, *E3S Web Conf.*, 44, 00013, <https://doi.org/10.1051/e3sconf/20184400013>, 2018.
- 855 Bezyk, Y., Dorodnikov, M., Górka, M., Sówka, I., and Sawiński, T.: Temperature and soil moisture control CO₂ flux and CH₄ oxidation in urban ecosystems, *Geochemistry*, 125989, <https://doi.org/10.1016/j.chemer.2023.125989>, 2023.
- Böttcher, K., Markkanen, T., Thum, T., Aalto, T., Aurela, M., Reick, C. H., Kolari, P., Arslan, A. N., and Pulliainen, J.: Evaluating Biosphere Model Estimates of the Start of the Vegetation Active Season in Boreal Forests by Satellite Observations, *Remote Sensing*, 8, 580, <https://doi.org/10.3390/rs8070580>, 2016.
- 860 Bretzel, F., Gaetani, M., Vannucchi, F., Caudai, C., Grossi, N., Magni, S., Caturegli, L., and Volterrani, M.: A multifunctional alternative lawn where warm-season grass and cold-season flowers coexist, *Landscape Ecol Eng*, 16, 307–317, <https://doi.org/10.1007/s11355-020-00423-w>, 2020.
- Cambou, A., Saby, N. P. A., Hunault, G., Nold, F., Cannavo, P., Schwartz, C., and Vidal-Beaudet, L.: Impact of city historical management on soil organic carbon stocks in Paris (France), *J Soils Sediments*, 21, 1038–1052, <https://doi.org/10.1007/s11368-020-02869-9>, 2021.
- 865 Capdevila, P., Stott, I., Oliveras Menor, I., Stouffer, D. B., Raimundo, R. L. G., White, H., Barbour, M., and Salguero-Gómez, R.: Reconciling resilience across ecological systems, species and subdisciplines, *Journal of Ecology*, 109, 3102–3113, <https://doi.org/10.1111/1365-2745.13775>, 2021.
- [Chapui-Lardy, L., Wrage, N., Metay, A., Chotte, J.-L., and Bernoux, M.: Soils, a sink for N₂O? A review, *Global Change Biology*, 13, 1–17, <https://doi.org/10.1111/j.1365-2486.2006.01280.x>, 2007.](https://doi.org/10.1111/j.1365-2486.2006.01280.x)
- 870 Chaudron, C., Mazalová, M., Kuras, T., Malenovský, I., and Mládek, J.: Introducing ecosystem engineers for grassland biodiversity conservation: A review of the effects of hemiparasitic *Rhinanthus* species on plant and animal communities at multiple trophic levels, *Perspectives in Plant Ecology, Evolution and Systematics*, 52, 125633, <https://doi.org/10.1016/j.ppees.2021.125633>, 2021.
- 875 Chen, X., Chen, H. Y. H., Chen, C., Ma, Z., Searle, E. B., Yu, Z., and Huang, Z.: Effects of plant diversity on soil carbon in diverse ecosystems: a global meta-analysis, *Biological Reviews*, 95, 167–183, <https://doi.org/10.1111/brv.12554>, 2020.
- Chollet, S., Brabant, C., Tessier, S., and Jung, V.: From urban lawns to urban meadows: Reduction of mowing frequency increases plant taxonomic, functional and phylogenetic diversity, *Landscape and Urban Planning*, 180, 121–124, <https://doi.org/10.1016/j.landurbplan.2018.08.009>, 2018.
- 880 Cong, W.-F., van Ruijven, J., Mommer, L., De Deyn, G. B., Berendse, F., and Hoffland, E.: Plant species richness promotes soil carbon and nitrogen stocks in grasslands without legumes, *Journal of Ecology*, 102, 1163–1170, <https://doi.org/10.1111/1365-2745.12280>, 2014.
- Costa, K. H. and Groffman, P. M.: Factors Regulating Net Methane Flux by Soils in Urban Forests And Grasslands, *Soil Science Society of America Journal*, 77, 850–855, <https://doi.org/10.2136/sssaj2012.0268n>, 2013.
- 885 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- De Boeck, H. J., Lemmens, C. M. H. M., Bossuyt, H., Malchair, S., Carnol, M., Merckx, R., Nijs, I., and Ceulemans, R.: How do climate warming and plant species richness affect water use in experimental grasslands?, *Plant Soil*, 288, 249–261, <https://doi.org/10.1007/s11104-006-9112-5>, 2006.

- 890 De Keersmaecker, W., van Rooijen, N., Lhermitte, S., Tits, L., Schaminée, J., Coppin, P., Honnay, O., and Somers, B.: Species-rich semi-natural grasslands have a higher resistance but a lower resilience than intensively managed agricultural grasslands in response to climate anomalies, *Journal of Applied Ecology*, 53, 430–439, <https://doi.org/10.1111/1365-2664.12595>, 2016.
- Decina, S. M., Hutyra, L. R., Gately, C. K., Getson, J. M., Reinmann, A. B., Short Gianotti, A. G., and Templer, P. H.: Soil respiration contributes substantially to urban carbon fluxes in the greater Boston area, *Environmental Pollution*, 212, 433–439, <https://doi.org/10.1016/j.envpol.2016.01.012>, 2016.
- 895 Dias, A. T. C., van Ruijven, J., and Berendse, F.: Plant species richness regulates soil respiration through changes in productivity, *Oecologia*, 163, 805–813, <https://doi.org/10.1007/s00442-010-1569-5>, 2010.
- Elonen, P.: Particle-size analysis of soil, *Suomen maataloustieteellinen seura*, 122, 1971.
- European Commission: Proposal for a regulation of the European Parliament and the Council on nature restoration, <https://doi.org/10.5281/ZENODO.5657041>, 2022.
- 900 European Environment Agency. EUNIS habitat types hierarchical view - revised groups: https://eunis.eea.europa.eu/habitats-code-browser-revised.jsp?expand=#level_23466, last access: 26 September 2023.
- Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species, *Planta*, 149, 78–90, <https://doi.org/10.1007/BF00386231>, 1980.
- Finnish Meteorological Institute: SmartMet Server, <https://github.com/fmidev/smartmet-plugin-timeseries>, last access: 30 March 2023a.
- 905 Finnish Meteorological Institute. Download observations: <https://en.ilmatieteenlaitos.fi/download-observations>, last access: 21 April 2023b.
- Fischer, L. K., Lippe, M. von der, Rillig, M. C., and Kowarik, I.: Creating novel urban grasslands by reintroducing native species in wasteland vegetation, *Biological Conservation*, 159, 119–126, <https://doi.org/10.1016/j.biocon.2012.11.028>, 2013.
- 910 Fornara, D. A. and Tilman, D.: Plant functional composition influences rates of soil carbon and nitrogen accumulation, *Journal of Ecology*, 96, 314–322, <https://doi.org/10.1111/j.1365-2745.2007.01345.x>, 2008.
- GADM. Maps and data. Ver 4.1: <https://gadm.org/index.html>, last access: 21 April 2023.
- 915 [Gillette, K.L., Qian, Y., Follett, R.F., and Del Grosso, S. Nitrous oxide emissions from a golf course fairway and rough after application of different nitrogen fertilizers, *Journal of Environmental Quality*, 45: 1788–1795. https://doi.org/10.2134/jeq2016.02.0047, 2016.](https://doi.org/10.2134/jeq2016.02.0047)
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornbluh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, <https://doi.org/10.1002/jame.20038>, 2013.
- 920 Groffman, P. M. and Pouyat, R. V.: Methane Uptake in Urban Forests and Lawns, *Environ. Sci. Technol.*, 43, 5229–5235, <https://doi.org/10.1021/es803720h>, 2009.
- 925 Haaland, C. and van den Bosch, C. K.: Challenges and strategies for urban green-space planning in cities undergoing densification: A review, *Urban Forestry & Urban Greening*, 14, 760–771, <https://doi.org/10.1016/j.ufug.2015.07.009>, 2015.
- Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Clim Dyn*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- 930 Hedblom, M., Lindberg, F., Vogel, E., Wissman, J., and Ahrné, K.: Estimating urban lawn cover in space and time: Case studies in three Swedish cities, *Urban Ecosyst*, 20, 1109–1119, <https://doi.org/10.1007/s11252-017-0658-1>, 2017.
- Hiller, R. V., McFadden, J. P., and Kljun, N.: Interpreting CO₂ Fluxes Over a Suburban Lawn: The Influence of Traffic Emissions, *Boundary-Layer Meteorol*, 138, 215–230, <https://doi.org/10.1007/s10546-010-9558-0>, 2011.

- Hossain, M. L., Li, J., Hoffmann, S., and Beierkuhnlein, C.: Biodiversity showed positive effects on resistance but mixed effects on resilience to climatic extremes in a long-term grassland experiment, *Science of The Total Environment*, 827, 154322, <https://doi.org/10.1016/j.scitotenv.2022.154322>, 2022.
- 935
- Ignatieva, M., Ahrné, K., Wissman, J., Eriksson, T., Tidåker, P., Hedblom, M., Kätterer, T., Marstorp, H., Berg, P., Eriksson, T., and Bengtsson, J.: Lawn as a cultural and ecological phenomenon: A conceptual framework for transdisciplinary research, *Urban Forestry & Urban Greening*, 14, 383–387, <https://doi.org/10.1016/j.ufug.2015.04.003>, 2015.
- Ignatieva, M., Eriksson, F., Eriksson, T., Berg, P., and Hedblom, M.: The lawn as a social and cultural phenomenon in Sweden, *Urban Forestry & Urban Greening*, 21, 213–223, <https://doi.org/10.1016/j.ufug.2016.12.006>, 2017.
- 940
- Ignatieva, M., Haase, D., Dushkova, D., and Haase, A.: Lawns in Cities: From a Globalised Urban Green Space Phenomenon to Sustainable Nature-Based Solutions, *Land*, 9, 73, <https://doi.org/10.3390/land9030073>, 2020.
- Institute of Atmospheric Research. SmartSMEAR - Download: <https://smear.avaa.csc.fi/download>, last access: 21 April 2023.
- IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, IPCC, 2022.
- 945
- Ivashchenko, K., Ananyeva, N., Vasenev, V., Sushko, S., Seleznyova, A., and Kudryarov, V.: Microbial C-availability and organic matter decomposition in urban soils of megapolis depend on functional zoning, *Soil & Environment*, 38, 31–41, <https://doi.org/10.25252/SE/19/61524>, 2019.
- Järvi, L., Hannuniemi, H., Hussein, T., Junninen, H., Aalto, P., Hillamo, R., Mäkelä, T., Keronen, P., Siivola, E., Vesala, T., and Kulmala, M.: The urban measurement station SMEAR III: Continuous monitoring of air pollution and surface–atmosphere interactions in Helsinki, Finland, *Atmosphere*, 14, 2009.
- 950
- Jasek-Kamińska, A., Zimnoch, M., Wachniew, P., and Rózański, K.: Urban CO₂ Budget: Spatial and Seasonal Variability of CO₂ Emissions in Krakow, Poland, *Atmosphere*, 11, 629, <https://doi.org/10.3390/atmos11060629>, 2020.
- Jokinen, P., Pirinen, P., Kaukoranta, J.-P., Kangas, A., Alenius, P., Eriksson, P., Johansson, M., and Wilkman, S.: Climatological and oceanographic statistics of Finland 1991–2020, Finnish Meteorological Institute, <https://doi.org/10.35614/isbn.9789523361485>, 2021.
- 955
- Jung, E.-Y., Gaviria, J., Sun, S., and Engelbrecht, B. M. J.: Comparative drought resistance of temperate grassland species: testing performance trade-offs and the relation to distribution, *Oecologia*, 192, 1023–1036, <https://doi.org/10.1007/s00442-020-04625-9>, 2020.
- 960
- Kaye, J. P., McCulley, R. L., and Burke, I. C.: Carbon fluxes, nitrogen cycling, and soil microbial communities in adjacent urban, native and agricultural ecosystems, *Global Change Biology*, 11, 575–587, <https://doi.org/10.1111/j.1365-2486.2005.00921.x>, 2005.
- Kenward, M. G. and Roger, J. H.: Small Sample Inference for Fixed Effects from Restricted Maximum Likelihood, *Biometrics*, 53, 983–997, <https://doi.org/10.2307/2533558>, 1997.
- 965
- Kohl, L., Koskinen, M., Rissanen, K., Haikarainen, I., Polvinen, T., Hellén, H., and Pihlatie, M.: Technical note: Interferences of volatile organic compounds (VOCs) on methane concentration measurements, *Biogeosciences*, 16, 3319–3332, <https://doi.org/10.5194/bg-16-3319-2019>, 2019.
- Kong, L., Shi, Z., and Chu, L. M.: Carbon emission and sequestration of urban turfgrass systems in Hong Kong, *Science of The Total Environment*, 473–474, 132–138, <https://doi.org/10.1016/j.scitotenv.2013.12.012>, 2014.
- 970
- Kose, M., Heinsoo, K., Kaljund, K., and Tali, K.: Twenty years of Baltic Boreal coastal meadow restoration: has it been long enough?, *Restoration Ecology*, 29, e13266, <https://doi.org/10.1111/rec.13266>, 2021.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World Map of the Köppen-Geiger climate classification updated, *International Journal of Climatology*, 26, 259–263, <https://doi.org/10.1127/0941-2948/2006/0130>, 2006.
- 975
- [Künnemann, T., Cannavo, P., Guérin, V., and Guénon, R.: Soil CO₂, CH₄ and N₂O fluxes in open lawns, treed lawns and urban woodlands in Angers, France, *Urban Ecosystems*, 26, 1659–1672 <https://doi.org/10.1007/s11252-023-01407-y>, 2023.](https://doi.org/10.1007/s11252-023-01407-y)
- Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: lmerTest Package: Tests in Linear Mixed Effects Models | Journal of Statistical Software, *Journal of Statistical Software*, 82, <https://doi.org/10.18637/jss.v082.i13>, 2017.

- Lampinen, J., Tuomi, M., Fischer, L. K., Neuenkamp, L., Alday, J. G., Bucharova, A., Cancellieri, L., Casado-Arzuaga, I., Čeplová, N., Cerveró, L., Deák, B., Eriksson, O., Fellowes, M. D. E., de Manuel, B. F., Filibeck, G., González-Guzmán, A., Hinojosa, M. B., Kowarik, I., Lumbierres, B., Miguel, A., Pardo, R., Pons, X., Rodríguez-García, E., Schröder, R., Sperandii, M. G., Unterweger, P., Valkó, O., Vázquez, V., and Klaus, V. H.: Acceptance of near-natural greenspace management relates to ecological and socio-cultural assigned values among European urbanites, *Basic and Applied Ecology*, 50, 119–131, <https://doi.org/10.1016/j.baae.2020.10.006>, 2021.
- 980
- [Law, Q.D., Trappe, J.M., Braun, R.C., and Patton, A.J.: Greenhouse gas fluxes from turfgrass systems: species, growth rate, clipping management, and environmental effects, *Journal of Environmental Quality* 50, 547–557, <https://doi.org/10.1002/jeq2.20222>, 2021.](https://doi.org/10.1002/jeq2.20222)
- 985
- Lane, I. G., Wolfen, J., Watkins, E., and Spivak, M.: Testing the Establishment of Eight Forbs in Mowed Lawns of Hard Fescue (*Festuca brevipila*) for Use in Pollinator Conservation in: *HortScience* Volume 54 Issue 12 (2019), 2019.
- Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-Vázquez, P. G., Malik, A. A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C., Trumbore, S. E., and Gleixner, G.: Plant diversity increases soil microbial activity and soil carbon storage, *Nat Commun*, 6, 6707, <https://doi.org/10.1038/ncomms7707>, 2015.
- 990
- Lerman, S. B. and Contosta, A. R.: Lawn mowing frequency and its effects on biogenic and anthropogenic carbon dioxide emissions, *Landscape and Urban Planning*, 182, 114–123, <https://doi.org/10.1016/j.landurbplan.2018.10.016>, 2019.
- Lindén, L., Riikonen, A., Setälä, H., and Yli-Pelkonen, V.: Quantifying carbon stocks in urban parks under cold climate conditions, *Urban Forestry & Urban Greening*, 49, 126633, <https://doi.org/10.1016/j.ufug.2020.126633>, 2020.
- 995
- [Livesley, S.J., Dougherty, B.J., Smith, A.J., Navaud, D., Wylie, L.J., and Arndt, S.K.: Soil-atmosphere exchange of carbon dioxide, methane and nitrous oxide in urban garden systems: impact of irrigation, fertiliser and mulch, *Urban Ecosystems*, 13, 273–293 <https://doi.org/10.1007/s11252-009-0119-6>, 2010.](https://doi.org/10.1007/s11252-009-0119-6)
- Loreau, M. and Hector, A.: Partitioning selection and complementarity in biodiversity experiments, *Nature*, 412, 72–76, <https://doi.org/10.1038/35083573>, 2001.
- 1000
- [Mäki, M., Ryhti, K., Fer, I., Ľupek, B., Vestin, P., Roland, M., Lehner, I., Köster, E., Lehtonen, A., Bäck, J., Heinonsalo, J., Pumpanen, J., and Kulmala, L.: Heterotrophic and rhizospheric respiration in coniferous forest soils along a latitudinal gradient, *Agricultural and Forest Meteorology* 317, 108876, <https://doi.org/10.1016/j.agrformet.2022.108876>, 2022.](https://doi.org/10.1016/j.agrformet.2022.108876)
- Marshall, C. A. M., Wilkinson, M. T., Hadfield, P. M., Rogers, S. M., Shanklin, J. D., Eversham, B. C., Healey, R., Kranse, O. P., Preston, C. D., Coghill, S. J., McGonigle, K. L., Moggridge, G. D., Pilbeam, P. G., Marza, A. C., Szigeccsan, D., Mitchell, J., Hicks, M. A., Wallis, S. M., Xu, Z., Toccaceli, F., McLennan, C. M., and Eves-van den Akker, S.: Urban wildflower meadow planting for biodiversity, climate and society: An evaluation at King’s College, Cambridge, *Ecological Solutions and Evidence*, 4, e12243, <https://doi.org/10.1002/2688-8319.12243>, 2023.
- 1005
- Metzger, M. J., Shkaruba, A. D., Jongman, R. H. G., and Brunce, R. H. G.: *Description of the European Environmental Zones and Strata*, Wageningen, Alterra, 2012.
- 1010
- Monteith, J. L.: Evaporation and environment, *Symposia of the Society for Experimental Biology*, 19, 205–234, 1965.
- Mueller, K. E., Hobbie, S. E., Tilman, D., and Reich, P. B.: Effects of plant diversity, N fertilization, and elevated carbon dioxide on grassland soil N cycling in a long-term experiment, *Global Change Biology*, 19, 1249–1261, <https://doi.org/10.1111/gcb.12096>, 2013.
- 1015
- Muller, S., Dutoit, T., Alard, D., and Gréville, F.: Restoration and Rehabilitation of Species-Rich Grassland Ecosystems in France: a Review, *Restoration Ecology*, 6, 94–101, <https://doi.org/10.1046/j.1526-100x.1998.06112.x>, 1998.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth System Science Data*, 13, 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- 1020
- National Land Survey of Finland. Topographic Database: <http://www.maanmittauslaitos.fi/en/maps-and-spatial-data/expert-users/product-descriptions/topographic-database>, last access: 21 April 2023.
- Nevalainen, O.: *satellitertools*, <https://github.com/ollinevalainen/satellitertools>, 2022.

- 1025 Nevalainen, O., Niemitalo, O., Fer, I., Juntunen, A., Mattila, T., Koskela, O., Kukkamäki, J., Höckerstedt, L., Mäkelä, L., Jarva, P., Heimsch, L., Vekuri, H., Kulmala, L., Stam, Å., Kuusela, O., Gerin, S., Viskari, T., Vira, J., Hyväluoma, J., Tuovinen, J.-P., Lohila, A., Laurila, T., Heinonsalo, J., Aalto, T., Kunttu, I., and Liski, J.: Towards agricultural soil carbon monitoring, reporting, and verification through the Field Observatory Network (FiON), *Geoscientific Instrumentation, Methods and Data Systems*, 11, 93–109, <https://doi.org/10.5194/gi-11-93-2022>, 2022.
- 1030 Niemelä, J., Saarela, S.-R., Söderman, T., Kopperoinen, L., Yli-Pelkonen, V., Väre, S., and Kotze, D. J.: Using the ecosystem services approach for better planning and conservation of urban green spaces: a Finland case study, *Biodivers Conserv*, 19, 3225–3243, <https://doi.org/10.1007/s10531-010-9888-8>, 2010.
- 1035 Norton, B. A., Bending, G. D., Clark, R., Corstanje, R., Dunnett, N., Evans, K. L., Grafius, D. R., Gravestock, E., Grice, S. M., Harris, J. A., Hilton, S., Hoyle, H., Lim, E., Mercer, T. G., Pawlett, M., Pescott, O. L., Richards, J. P., Southon, G. E., and Warren, P. H.: Urban meadows as an alternative to short mown grassland: effects of composition and height on biodiversity, *Ecological Applications*, 29, e01946, <https://doi.org/10.1002/eap.1946>, 2019.
- Oelmann, Y., Buchmann, N., Gleixner, G., Habekost, M., Roscher, C., Rosenkranz, S., Schulze, E.-D., Steinbeiss, S., Temperton, V. M., Weigelt, A., Weisser, W. W., and Wilcke, W.: Plant diversity effects on aboveground and belowground N pools in temperate grassland ecosystems: Development in the first 5 years after establishment, *Global Biogeochemical Cycles*, 25, <https://doi.org/10.1029/2010GB003869>, 2011.
- 1040 Patrignani, A. and Ochsner, T. E.: Canopeo: A Powerful New Tool for Measuring Fractional Green Canopy Cover, *Agronomy Journal*, 107, 2312–2320, <https://doi.org/10.2134/agronj15.0150>, 2015.
- Paudel, S. and States, S. L.: Urban green spaces and sustainability: Exploring the ecosystem services and disservices of grassy lawns versus floral meadows, *Urban Forestry & Urban Greening*, 84, 127932, <https://doi.org/10.1016/j.ufug.2023.127932>, 2023.
- 1045 Pavelka, M., Acosta, M., Kiese, R., Altimir, N., Brümmer, C., Crill, P., Darenova, E., Fuß, R., Gielen, B., Graf, A., Klemedtsson, L., Lohila, A., Longdoz, B., Lindroth, A., Nilsson, M., Jiménez, S. M., Merbold, L., Montagnani, L., Peichl, M., Pihlatie, M., Pumpanen, J., Ortiz, P. S., Silvennoinen, H., Skiba, U., Vestin, P., Weslien, P., Janous, D., and Kutsch, W.: Standardisation of chamber technique for CO₂, N₂O and CH₄ fluxes measurements from terrestrial ecosystems, *International Agrophysics*, 32, 569–587, <https://doi.org/10.1515/intag-2017-0045>, 2018.
- 1050 Poeplau, C., Marstorp, H., Thored, K., and Kätterer, T.: Effect of grassland cutting frequency on soil carbon storage – a case study on public lawns in three Swedish cities, *SOIL*, 2, 175–184, <https://doi.org/10.5194/soil-2-175-2016>, 2016.
- 1055 Pörtner, H.-O., Scholes, R. J., Agard, J., Archer, E., Arneth, A., Bai, X., Barnes, D., Burrows, M., Chan, L., Cheung, W. L. (William), Diamond, S., Donatti, C., Duarte, C., Eisenhauer, N., Foden, W., Gasalla, M. A., Handa, C., Kätterer, T., Hoegh-Guldberg, O., Ichii, K., Jacob, U., Insarov, G., Kiessling, W., Leadley, P., Leemans, R., Levin, L., Lim, M., Maharaj, S., Managi, S., Marquet, P. A., McElwee, P., Midgley, G., Oberdorff, T., Obura, D., Osman Elasha, B., Pandit, R., Pascual, U., Pires, A. P. F., Popp, A., Reyes-García, V., Sankaran, M., Settele, J., Shin, Y.-J., Sintayehu, D. W., Smith, P., Steiner, N., Strassburg, B., Sukumar, R., Trisos, C., Val, A. L., Wu, J., Aldrian, E., Parmesan, C., Pichs-Madruga, R., Roberts, D. C., Rogers, A. D., Díaz, S., Fischer, M., Hashimoto, S., Lavorel, S., Wu, N., and Ngo, H.: Scientific outcome of the IPBES-IPCC co-sponsored workshop on biodiversity and climate change, *Zenodo*, <https://doi.org/10.5281/ZENODO.4659158>, 2021.
- 1060 Pouyat, R. V., Yesilonis, I. D., and Nowak, D. J.: Carbon Storage by Urban Soils in the United States, *Journal of Environmental Quality*, 35, 1566–1575, <https://doi.org/10.2134/jeq2005.0215>, 2006.
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459–482, <https://doi.org/10.1002/jame.20022>, 2013.
- 1065 Riikonen, A. and Karilas, A.: Maaherranpuiston OmaStadi-niitty, Prosessin kuvaus ja niityn kehittymisen seuranta - VÄLIRAPORTTI 2021 [Maaherranpuisto's OmaStadi meadow, Description of the process and monitoring of the development of the meadow - INTERIM REPORT 2021, in Finnish], Sitowise, 2021.
- Ruimy, A., Jarvis, P. G., Baldocchi, D. D., and Saugier, B.: CO₂ Fluxes over Plant Canopies and Solar Radiation: A Review, in: *Advances in Ecological Research*, vol. 26, edited by: Begon, M. and Fitter, A. H., Academic Press, 1–68, [https://doi.org/10.1016/S0065-2504\(08\)60063-X](https://doi.org/10.1016/S0065-2504(08)60063-X), 1995.
- 1070 [Ryan, M.G. and Law, B.E.: Interpreting, measuring, and modeling soil respiration, *Biogeochemistry*, 73, 3-27, https://doi.org/10.1007/s10533-004-5167-7, 2005.](https://doi.org/10.1007/s10533-004-5167-7)

- 1075 Ryhti, K., Kulmala, L., Pumpanen, J., Isotalo, J., Pihlatie, M., Helmisaari, H.-S., Leppälampi-Kujansuu, J., Kieloaho, A.-J., Bäck, J., and Heinonsalo, J.: Partitioning of forest floor CO₂ emissions reveals the belowground interactions between different plant groups in a Scots pine stand in southern Finland, *Agricultural and Forest Meteorology*, 297, 108266, <https://doi.org/10.1016/j.agrformet.2020.108266>, 2021.
- Selhorst, A. and Lal, R.: Net Carbon Sequestration Potential and Emissions in Home Lawn Turfgrasses of the United States, *Environmental Management*, 51, 198–208, <https://doi.org/10.1007/s00267-012-9967-6>, 2013.
- 1080 Setälä, H. M., Francini, G., Allen, J. A., Hui, N., Jumpponen, A., and Kotze, D. J.: Vegetation Type and Age Drive Changes in Soil Properties, Nitrogen, and Carbon Sequestration in Urban Parks under Cold Climate, *Frontiers in Ecology and Evolution*, 4, 2016.
- Shen, J., Peng, Z., and Wang, Y.: From GI, UGI to UAGI: Ecosystem service types and indicators of green infrastructure in response to ecological risks and human needs in global metropolitan areas, *Cities*, 134, 104176, <https://doi.org/10.1016/j.cities.2022.104176>, 2023.
- 1085 Smith, L. S., Broyles, M. E. J., Larzleer, H. K., and Fellowes, M. D. E.: Adding ecological value to the urban lawnscape. Insect abundance and diversity in grass-free lawns, *Biodivers Conserv*, 24, 47–62, <https://doi.org/10.1007/s10531-014-0788-1>, 2015.
- Southon, G. E., Jorgensen, A., Dunnett, N., Hoyle, H., and Evans, K. L.: Biodiverse perennial meadows have aesthetic value and increase residents' perceptions of site quality in urban green-space, *Landscape and Urban Planning*, 158, 105–118, <https://doi.org/10.1016/j.landurbplan.2016.08.003>, 2017.
- 1090 Subke, J.A., Inghima, I., and Cotrufo M.F.: Trends and methodological impacts in soil CO₂ efflux partitioning: a meta-analytical review, *Global Change Biology*, 12, 921-943, <https://doi.org/10.1111/j.1365-2486.2006.01117.x>, 2006.
- Sushko, S., Ananyeva, N., Ivashchenko, K., Vasenev, V., and Kudryarov, V.: Soil CO₂ emission, microbial biomass, and microbial respiration of woody and grassy areas in Moscow (Russia), *J Soils Sediments*, 19, 3217–3225, <https://doi.org/10.1007/s11368-018-2151-8>, 2019.
- 1095 Thienelt, T. S. and Anderson, D. E.: Estimates of energy partitioning, evapotranspiration, and net ecosystem exchange of CO₂ for an urban lawn and a tallgrass prairie in the Denver metropolitan area under contrasting conditions, *Urban Ecosyst*, 24, 1201–1220, <https://doi.org/10.1007/s11252-021-01108-4>, 2021.
- Thompson, K., Hodgson, J. G., Smith, R. M., Warren, P. H., and Gaston, K. J.: Urban domestic gardens (III): Composition and diversity of lawn floras, *Journal of Vegetation Science*, 15, 373–378, <https://doi.org/10.1111/j.1654-1103.2004.tb02274.x>, 2004.
- 1100 Tilman, D., Lehman, C. L., and Thomson, K. T.: Plant diversity and ecosystem productivity: Theoretical considerations, *Proceedings of the National Academy of Sciences*, 94, 1857–1861, <https://doi.org/10.1073/pnas.94.5.1857>, 1997.
- Tonteri, T. and Haila, Y.: Plants in a boreal city: Ecological characteristics of vegetation in Helsinki and its surroundings, southern Finland, *Annales Botanici Fennici*, 27, 337–352, 1990.
- 1105 Trémeau, J., Karvinen, E., and Olascoaga, B.: Fluxes and plant diversity data in urban grasslands, <https://doi.org/10.23728/FMI-B2SHARE.920C1E5F08A74A6D9DFCB3A08CFC6734>, 2023.
- Unterweger, P. A., Schrode, N., and Betz, O.: Urban Nature: Perception and Acceptance of Alternative Green Space Management and the Change of Awareness after Provision of Environmental Information. A Chance for Biodiversity Protection, *Urban Science*, 1, 24, <https://doi.org/10.3390/urbansci1030024>, 2017.
- 1110 Upadhyay, S., Singh, R., Verma, P., and Raghubanshi, A. S.: Spatio-temporal variability in soil CO₂ efflux and regulatory physicochemical parameters from the tropical urban natural and anthropogenic land use classes, *Journal of Environmental Management*, 295, 113141, <https://doi.org/10.1016/j.jenvman.2021.113141>, 2021.
- Vekuri, H.: Chamber-codes, <https://github.com/hvekuri/Chamber-codes>, 2023.
- 1115 Venn, S. and Kotze, J.: Benign neglect enhances urban habitat heterogeneity: Responses of vegetation and carabid beetles (Coleoptera: Carabidae) to the cessation of mowing of park lawns, *European Journal of Entomology*, 111, 703–714, <https://doi.org/10.14411/eje.2014.089>, 2014.

- Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718, <https://doi.org/10.1175/2009JCLI2909.1>, 2010.
- 1120 Viherympäristöliitto. ABC-Vastaavuus: <https://www.vyl.fi/ohjeet/kunnossapitoluokitus/vastaavuus/>, last access: 21 April 2023.
- Vogel, A., Scherer-Lorenzen, M., and Weigelt, A.: Grassland Resistance and Resilience after Drought Depends on Management Intensity and Species Richness, *PLOS ONE*, 7, e36992, <https://doi.org/10.1371/journal.pone.0036992>, 2012.
- 1125 Walde, M., Allan, E., Cappelli, S. L., Didion-Gency, M., Gessler, A., Lehmann, M. M., Pichon, N. A., and Grossiord, C.: Both diversity and functional composition affect productivity and water use efficiency in experimental temperate grasslands, *Journal of Ecology*, 109, 3877–3891, <https://doi.org/10.1111/1365-2745.13765>, 2021.
- Waldén, E. and Lindborg, R.: Long Term Positive Effect of Grassland Restoration on Plant Diversity - Success or Not?, *PLOS ONE*, 11, e0155836, <https://doi.org/10.1371/journal.pone.0155836>, 2016.
- 1130 Wang, Q., Zeng, J., Qi, J., Zhang, X., Zeng, Y., Shui, W., Xu, Z., Zhang, R., Wu, X., and Cong, J.: A multi-scale daily SPEI dataset for drought characterization at observation stations over mainland China from 1961 to 2018, *Earth System Science Data*, 13, 331–341, <https://doi.org/10.5194/essd-13-331-2021>, 2021.
- Wang, R., Mattox, C. M., Phillips, C. L., and Kowalewski, A. R.: Carbon Sequestration in Turfgrass–Soil Systems, *Plants*, 11, 2478, <https://doi.org/10.3390/plants11192478>, 2022.
- Wastian, L., Unterweger, P. A., and Betz, O.: Influence of the reduction of urban lawn mowing on wild bee diversity (Hymenoptera, Apoidea), *JHR*, 49, 51–63, <https://doi.org/10.3897/JHR.49.7929>, 2016.
- 1135 Watson, C. J., Carignan-Guillemette, L., Turcotte, C., Maire, V., and Proulx, R.: Ecological and economic benefits of low-intensity urban lawn management, *Journal of Applied Ecology*, 57, 436–446, <https://doi.org/10.1111/1365-2664.13542>, 2020.
- Wei, X., Reich, P. B., Hobbie, S. E., and Kazanski, C. E.: Disentangling species and functional group richness effects on soil N cycling in a grassland ecosystem, *Global Change Biology*, 23, 4717–4727, <https://doi.org/10.1111/gcb.13757>, 2017.
- 1140 Yang, Y., Tilman, D., Furey, G., and Lehman, C.: Soil carbon sequestration accelerated by restoration of grassland biodiversity, *Nat Commun*, 10, 718, <https://doi.org/10.1038/s41467-019-08636-w>, 2019.
- [Zhan Y., Yao, Z., Groffman, P.M., Xie, J., Wang, Y., Li G., Zheng, X., and Butterbach-Bahl, K.: Urbanization can accelerate climate change by increasing soil N2O emission while reducing CH4 uptake, *Global Change Biology*, 29, 3489–3502, https://doi.org/10.1111/gcb.16652, 2023.](https://doi.org/10.1111/gcb.16652)
- 1145 Zirkle, G., Lal, R., and Augustin, B.: Modeling Carbon Sequestration in Home Lawns, *HortScience*, 46, 808–814, <https://doi.org/10.21273/HORTSCI.46.5.808>, 2011.
- Zobec, M., Betz, O., and Unterweger, P. A.: Perception of Urban Green Areas Associated with Sociodemographic Affiliation, Structural Elements, and Acceptance Stripes, *Urban Science*, 4, 9, <https://doi.org/10.3390/urbansci4010009>, 2020.