

1      **Soil priming effects and involved microbial community along salt gradients**

2  
3      Haoli Zhang<sup>ab#</sup>, Doudou Chang<sup>a#</sup>, Zhifeng Zhu<sup>c</sup>, Chunmei Meng<sup>b</sup>, Kaiyong Wang<sup>a\*</sup>

4

5      <sup>a</sup> Agricultural College, Shihezi University, Shihezi 832003, China

6      <sup>b</sup> College of Land Science and Technology, China Agriculture University, Yuanmingyuan West Road,

7      Beijing 100193, China

8      <sup>c</sup>China National Seed Group Co.Ltd, Yazhou Science and Technology City, Sanya 572000, China

9

10

11     All correspondence to: Kaiyong Wang

12     Email: wky20@163.com

13     <sup>#</sup> Both equally contribute to this work

14

15     5 Figures

16     3 Tables

17     4 Supplementary Figures

18     27 text pages

19     **Abstract**

20         Soil salinity mediates microorganisms and soil process, like soil organic carbon (SOC) cycling.  
21         Yet, how soil salinity affects SOC mineralization via shaping bacterial communities diversity and  
22         composition remains elusive. Therefore, soils were sampled along a salt gradient (salinity at 0.25%,  
23         0.58%, 0.75%, 1.00% and 2.64%) and incubated for 90 days to investigate i) SOC mineralization (i.e.  
24         soil priming effects induced by cottonseed meal, as substrate) and ii) responsible bacteria community,  
25         by using high throughput sequencing and natural abundance  $^{13}\text{C}$  isotopes (to partition cottonseed meal  
26         derived CO<sub>2</sub> and soil derived CO<sub>2</sub>). We observed negative priming effect during first 28 days of  
27         incubation but turned to positive priming effect after day 56. Negative priming at the early stage might  
28         be due to the preferential utilization of cottonseed meal. The followed positive priming decreased with  
29         the increase of salinity, which might be caused by the decreased alpha diversity of microbial  
30         community in soil with high salinity. Specifically, soil pH and EC along salinity gradient were the  
31         dominant variables modulating the structure of microbial community and consequently SOC priming  
32         (estimated by distance-based multivariate analysis and path analysis). By adopting O2PLS, priming  
33         effects were linked with specific microbial taxa, e.g., Proteobacteria (*Luteimonas*, *Hoeflea* and  
34         *Stenotrophomonas*) were the core microbial genus that attributed to the substrate induced priming  
35         effects. Here, we highlight that the increase of salinity reduced the diversity of microbial community  
36         and shifted dominant microorganisms(*Actinobacteria* and *Proteobacteria* (*Luteimonas*, *Hoeflea* and  
37         *Stenotrophomonas*)) that determined SOC priming effects, which provides a theoretical basis for  
38         understanding of SOC dynamics and microbial drivers under salinity gradient.

39

40         Keywords: *Salt gradient, priming effects, bacterial community, core microorganisms*

41     **1. Introduction**

42       Soil salinization is an increasing environmental problem caused by natural and  
43       human activities in the arid and semi-arid area (Wichern et al., 2006). Salinization is  
44       often a major threat to crop productivity in agricultural land. Soil microorganisms suffer  
45       from osmotic stress. Soil salinity often cause microbial death or dormant. It was widely  
46       reported that the increased salinity decrease microbial biomass, enzymatic activity, and  
47       alpha diversity of microbial community (Laura, 1974; Pathak and Rao, 1998; Rietz and  
48       Haynes, 2003). Soil salinity is reported to the major determinants of composition,  
49       activity of microbial community (Kamble et al., 2014). Although salinity is reported to  
50       be a vital factor in influencing microorganisms in the arid and semi-arid area, limited  
51       studies investigated C processes (e.g. priming effect) driven by microbial community  
52       in salinity soils (Sardinha et al., 2003).

53       Soil organic carbon (SOC) is the largest pool (1500 Pg C) in the terrestrial carbon  
54       (C) cycle, and contains twice as much C as the atmosphere (Filley and Boutton, 2006;  
55       Wiesmeier et al., 2019). The input of substrate C can influence the output (i.e., CO<sub>2</sub>  
56       release) through a phenomenon called priming effect, which was firstly discovered by  
57       LÖhnis (1926). Substrate additions accelerate or decrease soil organic C mineralization,  
58       referred to positive or negative priming effects (Kuzyakov et al., 2000). The intensity  
59       of the priming effect affects the turnover of SOC and thus storage pool (Sullivan and  
60       Hart, 2013). Soil priming effects are affected by many biotic and abiotic factors  
61       (Lavelle, 1997; Martin W, 2019), to investigate abiotic and biotic mechanisms  
62       underlying SOC priming enhance strong understanding of the SOC cycling.

63       Soil priming effects is affected by soil fauna animals (Scheu and Parkinson, 1994),  
64       activities, diversity and composition of microbial community (Di Lonardo et al., 2017;  
65       Fontaine et al., 2011). The microbial decomposers are the major player in the  
66       decomposition process of added C sources. The addition of substrate, such as composts  
67       (Xun et al., 2016), animal sludges (Hartmann et al., 2015), sewage sludges (Su et al.,  
68       2017; Wagner and Raquel, 2011) and plant residues (Dai et al., 2017), generally

69 increases soil microbial biomass C and stimulates the microbial activities thus enhanced  
70 the loss of SOC (positive priming effects) (Fontaine et al., 2003; Bird et al., 2011; Li et  
71 al., 2018; Ali et al., 2019).

72 Concerning abiotic factors, the priming effect can be controlled by climate  
73 variables (Hagemann, 2008), and soil properties, like pH, EC, TN, etc (Blagodatskaya  
74 and Kuzyakov 2008; Luo et al., 2017). To understand how environmental and edaphic  
75 factors affect the processes of SOC mineralization, is important to estimate terrestrial  
76 C pool (Lehmann and Kleber, 2015). Although many studies have tested the effects of  
77 soil pH, SOC content, and other edaphic variables on soil priming effect, few study  
78 investigated soil priming effects in salinity soil (Asghar et al., 2012), especially linked  
79 with soil microbial community structure and their functions in C decomposition (Soina  
80 et al., 2018).

81 Thus, we sampled the soils along natural salinity gradients (0.25%, 0.58%, 0.75%,  
82 1.00%, 2.64% apart from total water-soluble salt). Based on these soils, we conducted  
83 a 90 days of indoor incubation applying C3 substrate of cottonseed meal ( $\delta^{13}\text{C} =$   
84 23.47‰) to C4 soils with salt gradient ( $\delta^{13}\text{C}$  between -14.21‰ and -16.01‰), to  
85 investigate: 1) mineralization rate of cottonseed meal and induced soil priming effects  
86 along salt gradients; 2) diversity of microbial community in the soils with increased  
87 salinity, and 3) identify the bacteria taxa associated Soil priming. We hypothesized that  
88 i) soil microbial community diversity and composition will be different with the  
89 different in soil variables particularly pH and EC along salinity gradients, and ii) Soil  
90 C processes like priming effects will be regulated mainly by microbial community and  
91 especially the core microbial species. To clarify the priming effects and involved  
92 microbial groups would help us better understanding C sequestration potential and  
93 underlying mechanisms in saline soils.

94

## 95 **2. Materials and methods**

### 96 *2.1. Soil sampling and cottonseed meal production.*

97       The soil type was gray desert soil, which was collected from farmlands (82.90°  
98       longitude, 44.96° latitude) in **Xiao Yinpan town**, Bole City, Bortala, northern Xinjiang  
99       Uygur Autonomous Region, northwest China. The farmlands soil is naturally formed  
100      original saline-salinity soil and with a continuous 30 years planting of maize (C4 crop)  
101      and maize straw returning to soil for 7-8 year. In September 2021, we determining the  
102      sampling area, and use the five-point sampling method to collecting non-rhizosphere  
103      soil. The soil samples were indoor air drying and hand-picked to remove visible other  
104      debris, animal and plant residues and then sieved at field moisture (<2mm) and  
105      subsequently adjusted to 40% of water holding capacity (WHC). Texture was  
106      determined by the pipette method without carbonate in all soil samples. They were then  
107      incubated at 25 °C for 7 days before starting the experiments, to allow any early  
108      sampling and sieving effects to subside.

109       Cottonseed meal is a kind of reddish or yellow granular material obtained by  
110      pressing, leaching and other cottonseed. The cottonseed meal was purchased from the  
111      market and dried at 105 °C for 24 h indoor, then further pulverized by a ball mill and  
112      passed through < 2 mm sieve.

113

## 114      2.2. *Soil and substrate analyses*

115       EC and pH of soil and cottonseed meal were measured at a soil: water ratio of 1:5  
116      (weight/weight) (Bao, 2000). Air-dry soil (5 g, <2 mm) and 25 ml of deionised water  
117      were shaken together for 1 min and left to settle for 30 min, which was repeated once  
118      more before pH was determined with a pH electrode. Soil water-soluble salt was  
119      analyzed by weighted at a soil:water ratio of 1:5 (weight/weight). Air-dry soil (5 g, <1  
120      mm) and 25 ml of deionised water were shaken together for 30 min, filtration to obtain  
121      clear filtrate, using thermostat water bath to evaporate and weigh(Bao, 2000). Soil total  
122      carbon (TC), total nitrogen (TN) are collect soil to be tested was dried and ground  
123      through a 0.15mm screen, and a certain amount of treated soil sample was wrapped in

124 tin foil and placed in an element analyzer for determinatio (air-dried, milled <150 µm)  
125 were determined by dry combustion (LECO CNS 2000, LECO Corporation, Michigan,  
126 USA). Soil microbial biomass C was determined by fumigation extraction (Vance et  
127 al., 1987; Wu et al., 1990). The K<sub>2</sub>SO<sub>4</sub> extractable organic C was determined using an  
128 organic carbon autoanalyser (Shimadzu, Analytical Sciences, Kyoto, Japan). Soil  
129 microbial biomass C (Bc) was calculated from: Bc = 2.22 Ec, where Ec = [(organic C  
130 extracted from fumigated soil) minus (organic C extracted from non-fumigated soil)].  
131 The natural δ<sup>13</sup>C (‰) abundance of the soils (air-dried, milled <200 µm) was  
132 determined using an elemental analyser-isotope ratio mass spectrometer (Sercon Ltd,  
133 Crewe, UK). All measurements are given on an oven-dry weight basis (o.d., 105 °C, 24  
134 h).

135 The δ<sup>13</sup>C (‰) abundance of the cottonseed meal (air-dried, milled <200 µm) was  
136 determined using an elemental analyser-isotope ratio mass spectrometer (Sercon Ltd,  
137 Crewe, UK). The main elemental composition of the substrate was determined using  
138 elemental analysis (Vario EL Cube, Hanau, Germany), with the samples combusted at  
139 1200 °C. Natural δ<sup>13</sup>C (‰) abundance ,the total carbon, total nitrogen contents and C/N  
140 of the cottonseed meal was presented in Table 1.

141

#### 142 2.3. *Experimental design*

143 After pre-incubation, five soils with salinity gradient were thoroughly mixed with  
144 cottonseed meal at 20 mg C g<sup>-1</sup> soil (d.w. basis), and incubated over 90 days following  
145 moisture adjustment to 40% of water-holding capacity (WHC) to investigate the  
146 substrate mineralization and priming effects. Each soil sample (40 g d.w. basis) was  
147 incubated in a 100 ml beaker inside a 1 L brown glass jar. Three jars with only water  
148 and NaOH were set as blank. All the jars were sealed with a rubber bung and incubated  
149 in a randomized block design at 25 °C for the 90 days of incubation. The NaOH vials  
150 were changed after 1, 3, 5, 7, 14, 28, 56 and 90 days for determination of evolved CO<sub>2</sub>

151 and  $^{13}\text{C}$ -CO<sub>2</sub> (%). Meanwhile, soil biomass C, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, pH, EC, TC, TN and DNA  
152 extraction were measured at day 28.

153

154 *2.4. Soil CO<sub>2</sub>-C and its isotopic composition*

155 Soil C evolved as CO<sub>2</sub>-C in jars was measured by trapping CO<sub>2</sub> in 1 M NaOH  
156 (20 ml) during soil incubation. After the NaOH (20 ml) trapping CO<sub>2</sub> at different  
157 periods of soil incubation, 5 ml 1 M NaOH of each sample was mixed with 10 ml  
158 deionised water and titrated with 0.05 M standardised HCl by the TIM840 autotitrator  
159 (Radiometer Analytical, Villeurbanne Cedex, France). Meanwhile, the  $\delta^{13}\text{C}$  (%) of  
160 trapped CO<sub>2</sub>-C was precipitated, with 8 ml of the 1 M NaOH (20 ml) mixed with 8 ml  
161 1.5 M BaCl<sub>2</sub> in vials (Aoyama et al., 2000). The BaCO<sub>3</sub> precipitate was trapped on the  
162 glass fibre the filter, rinsed with deionised water several times, and dried overnight  
163 (80 °C), weighed (0.100-0.200 mg) into tin capsules, and analyzed for  $\delta^{13}\text{C}$  on an  
164 elemental analyzer-isotope ratio mass spectrometer (Sercon Ltd, Crewe, UK).

165

166 *2.5. DNA exaction and sequencing*

167 The total soil DNA was extracted from 0.50 g of moist soil using a FastDNA Spin  
168 Kit (MP Biomedicals, Santa Ana, CA, USA) according to the manufacturer's protocol.  
169 The extracted DNA was dissolved in 50  $\mu\text{l}$  of TE buffer, quantified using a  
170 spectrophotometer and stored at -20 °C until sequencing.

171 V3-V4 hypervariable regions of the bacterial 16S rRNA gene were amplified with  
172 primers 341F (5'-CCTAYGGRBGCASCAG-3') and 806R(5'-  
173 GGACTACHVGGGTWTCTAAT-3'). The PCR reactions were conducted with a  
174 thermocycler PCR system (GeneAmp 9700, ABI, USA) by using the following  
175 programs: 3 min of denaturation at 95 °C; followed by 27 cycles of 30 s at 95 °C, 30 s  
176 at 55 °C, and 45 s at 72 °C; and a final extension at 72 °C for 10 min with a thermocycler  
177 PCR system (GeneAmp9700, ABI, USA). PCR amplicons pooled from the triplicate  
178 reactions were purified using a QIAquick PCR purification kit (Qiagen, Shenzhen,

179 China), and quantified using a NanoDrop ND-1000 spectrophotometer (Thermo  
180 Scientific, Waltham, MA, USA). The PCR products were purified, mixed, and sent to  
181 Majorbio, Inc. (Shanghai, China) for sequencing based on the Illumina MiSeq platform.

182

183 *2.6. Calculations*

184 *2.6.1. CO<sub>2</sub>-δ<sup>13</sup>C emission*

185 The mineralisation of cottonseed meal was separated from SOC mineralisation  
186 according to the change of stable isotopic composition ( $\delta^{13}\text{C}$ ) with time. The standard  
187 equation for determining  $\delta^{13}\text{C}$  (‰) is derived from:

188  $\delta^{13}\text{C} (\text{\textperthousand}) = [(\text{R}_{\text{sample}}/\text{R}_{\text{VPDB}}) - 1] \times 1000,$  Eqn. 1

189 where  $\text{R}_{\text{sample}}$  is the mass ratio of  $^{13}\text{C}$  to  $^{12}\text{C}$  of each sample and  $\text{R}_{\text{VPDB}}$  is the  
190 international PDB([Peedee Belemnite](#)) limestone standard. The labeled  $^{13}\text{C}$  (%) of  
191 cottonseed meal was then estimated from:

192  $\text{CO}_2\text{-}^{13}\text{C} (\%) = (\delta_{\text{treatment}} - \delta\text{C4}) / (\delta\text{C3} - \delta\text{C4}),$  Eqn. 2

193 where  $\text{CO}_2\text{-}^{13}\text{C} (\%)$  is the proportion of evolved  $\text{CO}_2$  from C3 (cottonseed meal)  
194 matter,  $\delta_{\text{treatment}}$  is the  $\delta^{13}\text{C}$  (‰) in treatments of soil with cottonseed meal,  $\delta\text{C4}$  is the  
195  $\delta^{13}\text{C}$  (‰) in control soil and  $\delta\text{C3}$  is the  $\delta^{13}\text{C}$  (‰) from cottonseed meal. Thus, the  $\text{CO}_2\text{-}$   
196 C produced from cottonseed meal during the incubation was calculated from:

197  $\text{CO}_2\text{-}^{13}\text{C} (\mu\text{g g}^{-1} \text{soil}) = \text{CO}_2\text{-}^{13}\text{C} (\%) \times \text{total CO}_2\text{-C} (\mu\text{g g}^{-1} \text{soil})/100,$  Eqn. 3

198  $\text{CO}_2$  from SOC was  $\text{CO}_2\text{-}^{13}\text{C}$  subtracted from total evolved  $\text{CO}_2\text{-C}$ . The absolute  
199 soil priming effect (or primed soil  $\text{CO}_2\text{-C}$ ) with the addition of cottonseed meal was  
200 calculated from:

201  $\text{Primed soil CO}_2\text{-C} (\mu\text{g C g}^{-1} \text{soil}) = \text{CO}_2\text{-C}_{\text{treatment}} - \text{CO}_2\text{-C}_{\text{control}}$  Eqn. 4

202 where  $\text{CO}_2\text{-C}_{\text{treatment}}$  is the non-isotopically labeled  $\text{CO}_2\text{-C}$  evolved from  
203 cottonseed meal amended soil,  $\text{CO}_2\text{-C}_{\text{control}}$  is non-isotopically labeled  $\text{CO}_2\text{-C}$  evolved  
204 from soil without cottonseed meal.

205

206 *2.7. Statistics*

207        The data of  $^{16}\text{S}$  gene sequencing were processed using the Quantitative Insights  
208      Into Microbial Ecology (QIIME) 1.9.0-dev pipeline (Caporaso et al., 2010). In brief,  
209      Reads with less than length 200 bp and ambiguous bases were discarded. The sequences  
210      were then binned into operational taxonomic units (OTUs) by UCLUST (Edgar, 2010)  
211      based on 97% pairwise identity. Chimeric OTUs identified by USEARCH (Edgar et al.,  
212      2011) in QIIME were removed. The most abundant sequence from each OTU was  
213      selected to represent that OTU. Taxonomy was assigned to 16S OTUs against a subset  
214      of the Silva 104 database. The representative OTU sequences were aligned using  
215      PyNAST (Caporaso et al., 2010). We obtained between 64,425 and 89,989 clean\_reads  
216      per sample for all experimental samples.

217        To avoid potential bias caused by sequencing depth, all sample datasets were  
218      rarefied for the bacteria  $\alpha$ -diversity and  $\beta$ -diversity analyses. Faith's phylogenetic  
219      diversity was calculated to provide an integrated index of the phylogenetic breadth  
220      across taxonomic levels (Faith, 1992). To compare  $\beta$ -diversity between samples,  
221      principal coordinate analyses based on the unweighted and weighted UniFrac  
222      (Lozupone et al., 2007a) distances were calculated using the function 'pcoa' in the R  
223      package 'Ape'. Additionally, permutational multivariate analysis of variance  
224      (PERMANOVA) was carried out using the function 'adonis' in the R 'vegan' to  
225      measure effect size and significance on  $\beta$ -diversity. The variable influence projection  
226      (VIP) value was processed using the way of O2PLS analysis by the SIMCAP 14  
227      (Version 14.1.0.2047) (Wang et al., 2016). The y-matrix was defined as the  
228      environmental factors datasets and the x-matrix was defined as the microbial  
229      community on genus level dataset.

230        Data were logarithmically transformed and analyzed by ANOVA. All analyses  
231      were performed using SPSS software (13<sup>th</sup> edition). Pearson's correlation analyses were  
232      performed to assess the linear correlation among soil physio-chemical properties and  
233      microbial community. MULTIVARIATE analysis were operated to investigate  
234      interaction of salinity treatments on bacteria community parameters.

235

236 **3. Results**

237 **3.1. Soil physicochemical properties along salt gradients**

238       The major soil physicochemical properties along salt gradients were presented  
239 (Table 1) and all of soil physicochemical properties has significant difference ( $P <$   
240  $0.05$ ). The total soluble salinity content in the soils ranged from 0.25% to 2.64% of  
241 salinity soils, soil salt gradients increasing gradually from salinity 1 samples to salinity  
242 5 samples. The pH and EC in soils ranged from 8.45 to 8.85 and from  $1.06 \text{ ms cm}^{-1}$  to  
243  $7.75 \text{ ms cm}^{-1}$ . Soil total C and N were increased with salinity, ranging from 3.16% to  
244 3.57%, and from 0.18% to 0.26%. The  $\delta^{13}\text{C}$  value for soils are between -14.21‰ and -  
245 16.01‰, which were relatively enriched compared to cottonseed meal (-23.47‰). This  
246 allowed separation of soil derived  $\text{CO}_2$  from total evolved  $\text{CO}_2$ , according to the classic  
247 mixed modeling.

248

249 **3.2. Total  $\text{CO}_2$  evolution**

250       During the whole 90 days of incubation, the cumulative  $\text{CO}_2$  evolved had  
251 similar trends, which the amount of  $\text{CO}_2$  increased with the incubation times (Fig. S1).  
252 The cumulative  $\text{CO}_2$  evolved increased more rapidly with the addition of cottonseed  
253 meal before 14 days, compared to non-amended soils. At 90 days of incubation. The  
254 cumulative  $\text{CO}_2$  evolved in the soil with the lowest salinity (Salinity 1) gave the lowest  
255  $\text{CO}_2$  emission ( $597 \mu\text{g C g}^{-1}$ ) in the non-amended soils (Fig. S1,  $P < 0.001$ ).

256

257 **3.3. Cottonseed derived  $^{13}\text{CO}_2$  and soil priming effects**

258       The total cumulative  $\text{CO}_2\text{-C}$  was divided three parts based the  $\delta^{13}\text{C}$  value,  
259 including basal soil-derived  $\text{CO}_2$ , cottonseed meal-derived  $\text{CO}_2$  and primed soil  $\text{CO}_2$   
260 (Fig.1). The cottonseed meal-derived  $\text{CO}_2$  had a significant contribution to the total  $\text{CO}_2$   
261 evolved during the early incubation period. The cottonseed meal-derived  $\text{CO}_2$  was  
262 significantly higher in Salinity 1, Salinity 2 and Salinity 3 than in Salinity 4 and Salinity

263 5 before 28 days incubation. Meanwhile, the soil priming effects was negative in all  
264 amended soil treatments before 28 days incubation and the direction of priming effect  
265 in most of soil samples turned into positive after 28 days. During the whole 90 days  
266 incubation, there was a negative correlation between cottonseed meal-derived CO<sub>2</sub> and  
267 primed soil CO<sub>2</sub> (Fig. 2).

268

### 269 3.4. Bacterial diversity and community structure

270 The number of sequences ranged from 64,425 to 91,261 for per sample (average  
271 valve of 80,602). About 27,990 OTUs in total were obtained under different five  
272 treatments. Bacterial community diversity was measured by a series of OTU-based  
273 analyses of alpha diversity including chao1 estimator, and observed\_species in the  
274 QIIME pipeline (Fig. 3). Chao1 diversity estimator and observed\_species was  
275 significantly different in treatments, being the highest in Salinity 1, followed by Salinity  
276 3, Salinity 2, Salinity 4 and Salinity 5 ( $P < 0.01$ ). In general, bacterial community  
277 diversity decreased with increasing salinity (Fig. 3).

278 The most abundant phylum in the soils and their correlation with salinity were  
279 shown in Fig. 4. Among them, Actinobacteria was the dominant taxa in all soils, with  
280 the abundance ranging from 50.07 % (Salinity 3) to 68.99 % (Salinity 4). The relative  
281 abundance of Bacteroidetes, Firmicutes, and Deinococcus-Thermus increased with  
282 the salinity, while Acidobacteria decreased with salinity degree.

283 Based on OTUs of five gradient salt treatments, the PCA analysis showed that  
284 treatments from Salinity 2 and Salinity 4 clustered together. Meanwhile, soil samples  
285 of Salinity 1, Salinity 3 and Salinity 5 distributed in the first, fourth and three quadrant,  
286 which indicated that these treatments had large environmental heterogeneity (Fig. S4).

287 In order to visualize the relationship between environmental factors and microbial  
288 community, *Canonical Correspondence Analysis* (CCA) was conducted, showing that  
289 NO<sub>3</sub><sup>-</sup>-N, EC and TC had a more obvious impact than other factors for microbial  
290 community (Fig. 3). Soil EC were positively correlated with pH, NH<sub>4</sub><sup>+</sup>-N, and

291 negatively correlated with TN, TC and MBC. Mantel test and Distance-based  
292 multivariate analysis showed the contribution rate of different environmental factors  
293 account for 78% of the variability of microbial communities (Table 2). The value of pH  
294 (31%) and EC (12%) had a strong influence on microbial community.

295

296 **3.5. Relation between soil microbial community and C dynamics**

297 Based on the O2PLS analysis, the variable influence projection (VIP) values of  
298 bacterial genus more than 1.00% were showed their contributions to C decomposition  
299 of cottonseed meal-derived C, basal soil-derived C, and primed soil C (Table 3). There  
300 were many microbial taxa positively correlating to soil primed CO<sub>2</sub>, for insatnce, genera  
301 of *Actinomarinales*, *Luteimonas*, *Nocardiooides*, *Hoeflea*, *Intrasporangium*,  
302 *Nitrolancea*, *Pseudarthrobacter* and *Stenotrophomonas* had a positive correlation with  
303 primed CO<sub>2</sub>. In order to further to evaluate the relationship between soil properties, soil  
304 bacterial communities and C decomposition, we used the structural equation modeling  
305 (SEM) to suggest the direct and indirect impacts of salinity and microbial community  
306 on soil C decomposition (Fig. 7). The result showed that soil pH and EC had negative  
307 contribution to bacterial diversity, while bacterial diversity had a strong positive  
308 influence on the primed soil C (Fig. 5). For instance, salinity properties of EC had a  
309 directly negative influence on the bacterial diversity but positive influence on the  
310 primed soil C. Meanwhile, pH were negatively correlated with bacterial diversity and  
311 positively correlated with substrate derived C.

312

313 **4. Discussion**

314 **4.1. Soil priming effects along salty gradients**

315 Understanding soil C dynamics along salinity gradients is crucial to predict C  
316 sequestration in salty soils. In the early stage of the incubation, we observed that the  
317 cumulative substrate derived CO<sub>2</sub> in the soils with lower salinity was significantly  
318 higher than soils with higher salinity (Fig. 1), which can be possibly explained by that

319 high salinity inhibited microbial activity. Many studies have reported the influence of  
320 soil salinity on organic matter decomposition, mostly, the decomposition of organic  
321 matter are decreased by salinity (Wichern et al., 2006; Ghollarata and Raiesi, 2007;  
322 Tripathi et al., 2007; Setia et al., 2012). Yet, the response of microbial community to  
323 the increasing levels of salinity and consequent effects on soil priming effects remains  
324 largely unknown.

325 Here, we found soil priming effects was gradually changed from negative to  
326 positive priming effect (Fig. 1). The early pattern of the dynamics of the priming effect  
327 in this study was similar to other studies showing preferential utilization of labile C  
328 substance. The first phase of negative priming effects was likely to be caused by  
329 microbial assimilation of substrate. The soil microbes turned to use the new added  
330 substrate and thus used less of the original SOC. This was attributed to “preferential  
331 substrate utilization” (Perelo et al., 2005).

332 Soil microbial biomass-related growth predominating in the first phase were most  
333 likely to utilize SOC, leading to a positive priming effects after substrate was largely  
334 vanished. The magnitude of priming effects depends on soil microbial biomass size  
335 (Schneckenberger et al., 2008). It was found that the amount of added easily available  
336 organic C is beyond 50% of microbial biomass C (Blagodatskaya and Kuzyakov, 2008).  
337 Namely, the second phase of positive PEs probably was due to increased biomass size  
338 and enhanced demand on SOC. Secondly, C that was assimilated into microbial  
339 biomass in the first stage may also be mineralized in the second stage due to the  
340 turnover of microbial biomass (Shahbaz et al., 2017; Perelo et al., 2005).

341

#### 342 4.2. Microbial community along salt gradients

343 Previous studies concerning the impact of salinity on soil microbial community  
344 used different soils with a range of salt levels. In the present study we investigated the  
345 influence of soil salinity on microbial communities in soils from the closed area  
346 covering a range of salt content. Similarly, Rousk et al. (2011) also used agricultural

347 soils from the same area representing a range of soil salinity. Here, we found microbial  
348 diversity (alpha diversity) decreased with increasing salinity (Fig. 3). The negative  
349 impact on microbial diversity can be explained by that the accumulation of large  
350 amounts of salt in the soil raised the extracellular osmotic concentration (Rath and  
351 Rousk, 2015; Oren, 2011). The high osmotic pressures made it difficult for many  
352 microorganisms to adapt to and thus reduce their biological activity. The changes of  
353 soil microbial community structure were also explained by salinity (Herlemann et al.,  
354 2011; Campbell and Kirchman, 2013). We found that Bacteroidetes, Firmicutes,  
355 Acidobacteria and Deinococcus-Thermus were dominant in these soils (Fig. 4). These  
356 results are supported by previous findings that Firmicutes possess the high salinity  
357 resistance. Other studies also found that Bacteroidetes is dominant taxa in alkaline  
358 saline soil because of its resistant to salt (Valenzuela-Encinas et al., 2009; Keshri et al.,  
359 2013). Other study shows that the dominant phyla are Bacteroidetes and followed by  
360 Proteobacteria in the haloalkaline soil (Keshri et al., 2013). These results are consistent  
361 with the esuarine or marine environments, despite some studies suggest that soil salinity  
362 is not found to be a decisive factor for bacterial community and their growth (Rousk et  
363 al., 2011).

364 The difference of microbial community structure is affected by many soil variables,  
365 and pH and EC were the most important ones (Fig. 3; Table 2). Our results showed that  
366 the value of soil pH and EC would significantly affect the microbial community  
367 structure and the combined contribution rate of these two variables to microbial  
368 community was 43% (Table 2). At high levels of salt and alkaline arid condition, soil  
369 pH has been also shown to have a very powerful influence on the soil bacterial  
370 community structures (Bååth and Anderson, 2003; Fierer and Jackson, 2006; Rousk et  
371 al., 2010). Meanwhile, it is consequently unlikely that soil pH differences between the  
372 studied soils obscured the influence of salt (Rousk et al., 2011). Salinity has been  
373 identified as one of the most potent environmental factors that determine assembly of  
374 microbiome. Salinity has been regarded to play the vital role in shaping microbial

375 community in different ecosystem. This, despite the clear evidence from aquatic  
376 microbial ecology (Lozupone and Knight, 2007b), show a potential for salt to affect  
377 soil microbial communities apart from that of pH (Rath and Rousk, 2015).

378

379 4.3. The core microbial taxa regulating C decomposition along salinity gradient

380 The correlation of microbial taxa and SOC decomposition (priming) were found  
381 according to the results of O2PLS and SEM (Table 3; Fig. 5). Here we showed that  
382 *Streptomyces* (Actinobacteria), *Glycomyces* (branch of Actinobacteria), *Agromyces*  
383 (branch of Actinobacteria), and *Sphingomonas* (branch of Proteobacteria) at the genus  
384 level were significantly correlated with the C process particularly primed soil-driven C.  
385 Most of these functional taxa belonged to Actinobacteria and Proteobacteria. In a recent  
386 study, Ren et al. (2018) found that Actinobacteria had negative impact on SOC  
387 mineralization across land-use change (Fierer et al., 2007; Goldfarb et al., 2011) and  
388 Proteobacteria drove the positive soil respiration (He et al., 2012; Stevenson et al.,  
389 2004), indicating the balance of soil C dynamics were largely regulated by these two  
390 phyla. We found similar result that *Streptomyces* (branch of Actinobacteria) had a  
391 negative correlation with primed soil CO<sub>2</sub>. Actinobacteria are able to grow  
392 preferentially on the C-rich refractory materials and relatively easily decompose the  
393 cellulose, lignocellulose (Khodadad et al., 2011), indicating these microorganisms  
394 preferentially use the C source that is used partially by others.

395 Although some studies suggest soil salinity may not be a vital factor for C  
396 decomposers (Rousk et al., 2011), the composition of microbial community are  
397 considered to play a decisive role in determining C dynamic processes in response to  
398 salt stress (Ramsey et al., 2005; Schimel et al., 2007; Nottingham et al., 2009). Here,  
399 SEM analysis showed that soil pH and EC in salted soils reduced microbial diversity  
400 and thus limited the utilization of SOC by microbial community, It was reported that  
401 high pH and salinity are the major determinants of soil microbial activity and  
402 community structure (Kamble et al., 2014).

403

404 **5. Conclusion**

405 Cotton meal is a kind of organic material with high nitrogen content, adding cotton  
406 meal in salinised soil can stimulate and promote the release of soil nutrients. The  
407 microorganisms mainly use the organic matter in the cotton meal in the pre-culture  
408 period, so the soil carbon excitation is negative excitation, Soil priming effect turned  
409 from negative to positive at the later stage of incubation (day 28), because  
410 microorganisms turned to decompose SOC from the labile substrate. With the increase  
411 of salinity, the diversity of microbial community decreased. Soil microbial community  
412 was mainly controled by soil pH and EC. By O2PLS, we found Actinobacteria and  
413 Proteobacteria (*Luteimonas*, *Hoeflea* and *Stenotrophomonas*) dominant in these soils  
414 were the core microbial taxa that affecting the process of organic C mineralization,  
415 particularly soil primed CO<sub>2</sub>.

416

417 **Acknowledgements**

418 This study was supported by the Special Fund for Key Science & Technology  
419 Program in Xinjiang Province of China (No. 2022B02021-3-1) .

420

421 **Data availability**

422 The datasets used and analysed during the current study available from the  
423 corresponding author on reasonable request.

424

425 **Author contributions**

426 K.W. conceptualized and conducted the experiment. H.Z. and D.C. conducted the  
427 data analysis and wrote the manuscript, conducted the indoor experiment. C.M. and  
428 Z.Z. assisted in conducting the experiment. All authors reviewed the manuscript. All  
429 authors contributed to the manuscript and approved the submitted version.

430

431    **Competing interests**

432        The authors declare no competing interests.

433

434    **Reference**

- 435    Anderson, C.R., Condron, L.M., Clough, T.J., Fiers, M., Stewart, A., Hill, R.A.,  
436        Sherlock, R.R., 2011. Biochar induced soil microbial community change:  
437        implications for biogeochemical cycling of carbon, nitrogen and  
438        phosphorus. *Pedobiologia* 54, 0-320.
- 439    Asghar, H. N., Setia, R., Marschner, P., 2012. Community composition and activity of  
440        microbes from saline soils and non-saline soils respond similarly to changes in  
441        salinity. *Soil Biology and Biochemistry* 47, 175-178.
- 442    Ali N., Khan S., Li Y., Zheng N., Yao H., 2019. Influence of biochars on the  
443        accessibility of organochlorine pesticides and microbial community in  
444        contaminated soils. *Science of The Total Environment* 551-560.
- 445    Bååth, E., Anderson, T.H., 2003. Comparison of soil fungal/bacterial ratios in a pH  
446        gradient using physiological and PLFA-based techniques. *Soil Biology and*  
447        *Biochemistry* 35, 955-963.
- 448    Bao, S.D., 2000. *Soil and Agricultural Chemistry Analysis*, third ed. China Agriculture  
449        Press, Beijing.
- 450    Blagodatskaya, E., Kuzyakov, Y., 2008. Mechanisms of real and apparent priming  
451        effects and their dependence on soil microbial biomass and community structure:  
452        critical review. *Biology and Fertility of Soils* 45, 115-131.
- 453    Bird, J.A., Herman, D.J., Firestone, M.K., 2011. Rhizosphere priming of soil organic  
454        matter by bacterial groups in a grassland soil. *Soil Biology Biochemistry*. 43,  
455        718 – 725.
- 456    Campbell, B.J., Kirchman, D.L., 2013. Bacterial diversity, community structure and  
457        potential growth rates along an estuarine salinity gradient. *The ISME Journal* 7(1),  
458        210-220.

- 459 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,  
460 E.K., Fierer, N., P?a, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley,  
461 S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D.,  
462 Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters,  
463 W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows  
464 analysis of high-throughput community sequencing data. *Nature Methods* 7(5),  
465 335-336.
- 466 Dai, H., Chen, Y., Yang, X., Cui, J., Sui, P., 2017. The effect of different organic  
467 materials amendment on soil bacteria communities in barren sandy loam  
468 soil. *Environmental Science Pollution Research* 24(14), 1-10.
- 469 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST.  
470 *Bioinformatics* 26, (19), 2460-2461.
- 471 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME  
472 improves sensitivity and speed of chimaera detection. *Bioinformatics*. 27(16), 194-  
473 2200.
- 474 Faith D.P., 1992. Conservation evaluation and phylogenetic diversity. *Biological  
475 Conservation* 1-10.
- 476 Filley, T.R., Boutton, T.W., 2006. Ecosystems in flux: molecular and stable isotope  
477 assessments of soil organic matter storage and dynamics. *Soil Biology and  
478 Biochemistry* 38, 3181-3183.
- 479 Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial  
480 communities. *Proceedings of the National Academy of Sciences of the USA* 103,  
481 626-631.
- 482 Fierer, N., Bradford, M., Jackson, R., 2007. Toward an ecological classification of soil  
483 bacteria. *Ecology* 88, 1354–1364.
- 484 Fontaine, S., Mariotti, A., Abbadie, L., 2003. The priming effect of organic matter: A  
485 question of microbial competition?. *Soil Biology and Biochemistry* 35(6), 837-  
486 843.

- 487 Fontaine, S., Henault, C., Aamor, A., Bdioui, N., Bloor, J.M.G., Maire, V., Mary, B.,  
488 Revaillet, S., Maron, P.A., 2011. Fungi mediate long term sequestration of carbon  
489 and nitrogen in soil through their priming effect. *Soil Biology and Biochemistry*  
490 43(1), 86-96.
- 491 Goldfarb, K., Karaoz, U., Hanson, C., Santee, C., Bradford, M., Treseder, K.,  
492 Wallenstein, M., Brodie, E., 2011. Differential growth responses of soil bacterial  
493 taxa to carbon substrates of varying chemical recalcitrance. *Frontiers in*  
494 *Microbiology*. 2, 94-97.
- 495 Ghollarata, M., Raiesi, F., 2007. The adverse effects of soil salinization on the growth  
496 of trifolium alexandrinum l. and associated microbial and biochemical properties  
497 in a soil from iran. *Soil Biology and Biochemistry* 39(7), 1699-1702.
- 498 Hagemann, S., 2008. Vulnerability of permafrost carbon to climate change:  
499 Implications for the global carbon cycle. *Bioscience* 701-714.
- 500 Hartmann, M., Frey, B., Mayer, J., Widmer, F., 2015. Distinct soil microbial diversity  
501 under long-term organic and conventional farming. *The ISME Journal* 9(5), 1177-  
502 1194.
- 503 He, Z., Piceno, Y., Deng, Y., Xu, M., Lu, Z., DeSantis, T., Andersen, G., Hobbie, S.E.,  
504 Reich, P.B., Zhou, J., 2012. The phylogenetic composition and structure of soil  
505 microbial communities shifts in response to elevated carbon dioxide. *The ISME*  
506 *journal*. 6(2), 259-272.
- 507 Herlemann, D. P., Labrenz, M., Jürgens, Klaus, Bertilsson, S., Waniek, J. J., Andersson,  
508 A. F., 2011. Transitions in bacterial communities along the 2000 km salinity  
509 gradient of the baltic sea. *The ISME Journal* 5(10), 1571-1579.
- 510 Kuzyakov, Y., Friedel, J.K., Stahr, K., 2000. Review of mechanisms and quantification  
511 of priming effects. *Soil Biology and Biochemistry* 32, 1485-1498.
- 512 Khodadad, C.L.M., Zimmerman, A.R., Green, S.J., Uthandi, S., Foster, J.S., 2011.  
513 Taxa-specific changes in soil microbial community composition induced by  
514 pyrogenic carbon amendments. *Soil Biology and Biochemistry* 43(2), 385-392.

- 515 Keshri, J., Mody, K., Jha, B.. 2013. Bacterial community structure in a semi-arid  
516 haloalkaline soil using culture independent method. *Geomicrobiology Journal* 30(6), 517-529.
- 518 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature*  
519 528, 60-68.
- 520 LÖhnis, F., 1926. Nitrogen availability of green manures. *Soil Science* 22: 253 – 290.
- 521 Sullivan, B. W., and Hart, S. C., 2013. Evaluation of mechanisms controlling the  
522 priming of soil carbon along a substrate age gradient. *Soil Biology &*  
523 *Biochemistry* 58(2), 293-301.
- 524 Lozupone, C.A., Hamady, M., Kelley, S.T., Knight, R., 2007a. Quantitative and  
525 qualitative beta diversity measures lead to different insights into factors that  
526 structure microbial communities. *Applied and Environmental Microbiology* 73(5),  
527 1576-1585.
- 528 Lozupone, C.A., Knight, R., 2007b. Global patterns in bacterial diversity. *Proceeding*  
529 *of the National Academy of Science* 104(27), 11436-11440.
- 530 Lavelle, P., et al., 1997. Soil function in a changing world: the role of invertebrate  
531 ecosystem engineers. *Eur. J. Soil Biol* 33 (4), 159–193.
- 532 Luo, Y., Zang, H., Yu, Z., Chen, Z., Gunina, A., Kuzyakov, Y., Xu, J., Zhang, K.,  
533 Brookes, P.C., 2017. Priming effects in biochar enriched soils using a three-  
534 source-partitioning approach:  $^{14}\text{C}$  labelling and  $^{13}\text{C}$  natural abundance. *Soil*  
535 *Biology and Biochemistry* 28-35.
- 536 Li X., Chen Q.L., He C., Shi Q., Chen S.C., Reid B.J., Zhu Y.G., Sun G.X., 2018.  
537 Organic Carbon Amendments Affect the Chemodiversity of Soil Dissolved  
538 Organic Matter and Its Associations with Soil Microbial Communities.  
539 *Environmental Science and Technology* 53(1), 50-59.
- 540 Laura, R.D., 1974. Effects of neutral salts on carbon and nitrogen mineralisation of  
541 organic matter in soil. *Plant and Soil* 41, 113–127.
- 542 Nottingham, A.T., Griffiths, H., Chamberlain, P.M., Stott, A.W., Tanner, E.V.J., 2009.

- 543        Soil priming by sugar and leaf-litter substrates: a link to microbial groups. Applied.  
544        Soil Ecology 42, 183–190.
- 545        Oren, A., 2011. Thermodynamic limits to microbial life at high salt concentrations.  
546        Environmental Microbiology 13(8), 1908-1923.
- 547        Perelo, L.W., Jimenez, M., Munch, J.C., 2005. Microbial immobilisation and turnover  
548        of  $^{15}\text{N}$  labelled substrates in two arable soils under field and laboratory conditions.  
549        Soil Biology and Biochemistry 38(5), 912-922.
- 550        Pathak, H., Rao, D.L.N., 1998. Carbon and nitrogen mineralization from added organic  
551        matter in saline and alkaline soils. Soil Biology and Biochemistry. 30, 695–702.
- 552        Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight,  
553        R., Fierer, N., 2010. Soil bacterial and fungal communities across a pH gradient  
554        in an arable soil. The ISME Journal 4, 1340-1351.
- 555        Rousk, J., Elyaagubi, F.K., Jones, D.L., Godbold, D.L., 2011. Bacterial salt tolerance  
556        is unrelated to soil salinity across an arid agroecosystem salinity gradient. Soil  
557        Biology and Biochemistry 43, 1881-1887.
- 558        Ramsey, P.W., Rillig, M.C., Feris, K.P., Gordon, N.S., Moore, J.N., Holben, W.E.,  
559        Gannon, J.E., 2005. Relationship between communities and processes; new  
560        insights from a field study of a contaminated ecosystem. Ecology Letter 8, 1201-  
561        1210.
- 562        Ren, C., Wang, T., Xu, Y., Deng, J., Zhao, F., Yang, G., Han, X., Feng, Y., Ren, G.,  
563        2018. Differential soil microbial community responses to the linkage of soil  
564        organic carbon fractions with respiration across land-use changes. Forest Ecology  
565        and Management 12, 170-178.
- 566        Rath, K.M., Rousk, J., 2015. Salt effects on the soil microbial decomposer community  
567        and their role in organic carbon cycling: A review. Soil Biology and Biochemistry  
568        108-123.
- 569        Shahbaz, M., Kuzyakov, Y., Sanaullah, M., Heitkamp, F., Zelenev, V., Kumar, A.,  
570        Blagodatskaya, E., 2017. Microbial decomposition of soil organic matter is

- 571 mediated by quality and quantity of crop residues: mechanisms and thresholds.  
572 *Biology and Fertility of Soils* 53(3), 287–301.
- 573 Schneckenberger, K., Demin, D., Stahr, K. and Kuzyakov, Y. 2008. Microbial  
574 utilization and mineralization of  $^{14}\text{C}$  glucose added in six orders of concentration  
575 to soil. *Soil Biology and Biochemistry* 40: 1981–1988.
- 576 Schimel, J., Balser, T.C., Wallenstein, M., 2007. Microbial stress–response physiology  
577 and its implications for ecosystem function. *Ecology* 88, 1386–1394.
- 578 Stevenson, B.S., Eichorst, S.A., Wertz, J.T., Schmidt, T.M., Breznak, J.A., 2004. New  
579 strategies for cultivation and detection of previously uncultured microbes. *Applied  
580 and Environmental Microbiology* 70(8), 4748-4755.
- 581 Rietz, D.N., Haynes, R.J., 2003. Effects of irrigation-induced salinity and sodicity on  
582 soil microbial activity. *Soil Biology Biochemistry*. 35, 845–854.
- 583 Scheu, S., Parkinson, D., 1994. Effects of earthworms on nutrient dynamics, carbon  
584 turnover and microorganisms in soils from cool temperate forests of the Canadian  
585 Rocky Mountains - laboratory studies. *Applied Soil Ecology*, 5, 23-27
- 586 Su, J. Q., An, X. L., Li, B., Chen, Q. L., Gillings, M. R., Chen, H., 2017. Metagenomics  
587 of urban sewage identifies an extensively shared antibiotic resistome in  
588 china. *Microbiome* 5, 84-85.
- 589 Soina V.S., Mergelov N.S., Kudinova A.G., Lysak L.V., Demkina E.V., Vorobyova  
590 E.A., Dolgikh A.V., Shorkunov I.G., 2018. Microbial Communities of Soils and  
591 Soil-like Bodies in Extreme Conditions of East Antarctica. *Paleontological  
592 Journal* 52(10), 1186-1195.
- 593 Setia, R., Setia, D., Marschner, P., 2012. Short-term car- bon mineralization in saline-  
594 sodic soils. *Biology and Fertility of Soils* 48: 475–479.
- 595 Sardinha, M., Müller, T., Schmeisky, H., Joergensen, R.G., 2003. Microbial  
596 performance in soils along a salinity gradient under acidic conditions. *Applied Soil  
597 Ecology* 23, 237–244.

- 598 Tripathi, S., Chakraborty, A., Chakrabarti, K., Bandyopadhyay, B.K., 2007. Enzyme  
599 activities and microbial biomass in coastal soils of India. *Soil Biology and*  
600 *Biochemistry* 39: 2840–2848.
- 601 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for  
602 measuring soil microbial biomass C. *Soil Biology and Biochemistry* 19, 703-707.
- 603 Valenzuela-Encinas, C., Neria-González, I., Alcántara-Hernández, R.J., Estrada-  
604 Alvarado, I., Dendooven, L., Marsch, R., 2009. Changes in the bacterial  
605 populations of the highly alkaline saline soil of the former lake Texcoco (Mexico)  
606 following flooding. *Extremophiles* 13 (4), 609 – 621.
- 607 Wang, Z.M., Lu, Z.M., Shi, J.S., Xu, Z.H., 2016. Exploring flavour-producing core  
608 microbiota in multispecies solid-state fermentation of traditional Chinese vinegar.  
609 *Scientific Report* 6, 26818.
- 610 Wiesmeier, Martin., Urbanski, Livia., Hobley, Eleanor., Lang, Birgit., von Ltzow,  
611 Margit., Marin-Spiotta, Erika., van Wesemael, Bas., Rabot, Eva., Lie, Mareike.,  
612 Garcia-Franco, Noelia., Wollschlger, Ute., Vogel, Hans-Jrg., Kgel-Knabner,  
613 Ingrid., 2019. Soil organic carbon storage as a key function of soils - A review of  
614 drivers and indicators at various scales. *Geoderma* 149-162.
- 615 Wagner, B., Raquel, G., 2011. Impacts of sewage sludge in tropical soil: a case study  
616 in brazil. *Applied and Environmental Soil Science* 1-11.
- 617 Wichern, J., Wichern, F., Joergensen, R.G., 2006. Impact of salinity on soil microbial  
618 communities and the decomposition of maize in acidic soils. *Geoderma* 137, 100-  
619 108.
- 620 Wu, J., Joergensen, R.G., Pommerening, B., Chaussod, R., Brookes, P.C., 1990.  
621 Measurement of soil microbial biomass C by fumigation-extraction: an automated  
622 procedure. *Soil Biology and Biochemistry* 22, 1167-1169.
- 623 Xun, W., Zhao, J., Xue, C., Zhang, G., Ran, W., Wang, B., Shen, Q., Zhang, R., 2016.  
624 Significant alteration of soil bacterial communities and organic carbon  
625 decomposition by different longterm fertilization management conditions of

626 extremely low productivity arable soil in South China. Environmental  
627 Microbiology 18, 1907–1917.  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666

**Table 1.** Soil samples and Cottonseed meal properties

	<b>Salinity 1</b>	<b>Salinity 2</b>	<b>Salinity 3</b>	<b>Salinity 4</b>	<b>Salinity 5</b>	<b>Cottonseed meal</b>
<b>Total C (%)</b>	3.38b	3.18c	3.16c	3.57a	3.35b	42.98
<b>Total N (%)</b>	0.18d	0.19d	0.20c	0.22b	0.26a	5.84
<b>C/N ratio</b>	18.32a	16.56b	15.71c	16.54b	12.94d	7.38
<b><math>\delta^{13}\text{C}</math> value (‰)</b>	-14.21a	-14.79c	-14.60b	-14.55b	-16.01d	-23.47
<b>pH (H<sub>2</sub>O)</b>	8.85a	8.45c	8.58b	8.59b	8.55b	7.63
<b>EC (dS m<sup>-1</sup>)</b>	1.06e	1.96c	1.28d	2.64b	7.75a	2.56
<b>Salinity (%)</b>	0.25e	0.58d	0.75c	1.00b	2.64a	ND

668

669

670 **Table 2.** Mantel test and Distance-based multivariate analysis relevance and  
671 contribution rate between soil properties and bacterial community compositions.

	pH	EC	NO <sub>3</sub> <sup>-</sup> -N	NH <sub>4</sub> <sup>+</sup> -N	MBC	TN	TC
Correlation	0.74**	0.56**	0.36**	0.68**	0.31**	0.11	0.27
Contribution	0.31**	0.12**	0.05	0.04	0.16	0.03	0.07**

672 Note: \* p < 0.05, \*\* p < 0.01

673

674

675 **Table 3.** The variable influence projection (VIP) value and Spearman's correlation  
 676 between the relative abundances of genera and C dynamic.

Phylum-Genus	VIP	Cottonseed meal CO <sub>2</sub> -C(µg g <sup>-1</sup> )	Primed soil CO <sub>2</sub> -C(µg g <sup>-1</sup> )	Basal soil CO <sub>2</sub> -C(µg g <sup>-1</sup> )
Actinobacteria-Actinomarinales	1.36		0.63**	
Proteobacteria-Luteimonas	1.31		0.80**	
Actinobacteria-Nocardioides	1.30		0.54*	
Proteobacteria-Hoeflea	1.29		0.73**	
Actinobacteria-Streptomyces	1.27		-0.84**	
Actinobacteria-Glycomyces	1.26	0.63**		
Actinobacteria-Marmoricola	1.26	-0.52		
Proteobacteria-Nitrosospira	1.23		0.59	
Actinobacteria-Intrasporangium	1.22		0.60*	
Actinobacteria-Agromyces	1.19			0.58*
Proteobacteria-Sphingomonas	1.18			0.65**
Actinobacteria-Myceligerans	1.16			
Chloroflexi-Nitrolancea	1.15		0.65**	
Actinobacteria-Pseudarthrobacter	1.06		0.62**	
Proteobacteria-Stenotrophomonas	1.00	-0.50	0.72**	

677 Note: \* p < 0.05, \*\* p < 0.01

678


679

680

681

682

683



684


685 **Fig. 1.** Partitioning of CO<sub>2</sub> evolution after addition of cottonseed meal in different five  
 686 salinity soils. Cumulative CO<sub>2</sub> evolved from salinity soil of 0.25 % (a) , 0.58 % (b) ,  
 687 0.75 % (c) ,1.00% (d) and 2.64%(e) . Error bars represent standard errors of the means  
 688 (n = 3).

689

690

691

692



693

694 **Fig. 2.** Correlation between primed soil mineralisation and cottonseed meal  
 695 mineralisation following different five salinity soils during 90 days incubation

696

697

698

699

700

701

702

703

704

705

706

707


708 **Fig. 3.** Microbial community alpha diversity (Chao1) observed\_species and beta  
 709 diversity. Within each panel, boxplot data refer to maximum date (top line), 99% (the  
 710 second line), mean (the third line), 1% (the fourth line) and minimum date (bottom line)  
 711 of the different treatments, with statistical significance ( $P < 0.05$ ).  
 712


713

714

715

716






717

718 **Fig. 4.** The top 10 of phylums and genes in bacterial community in soils with a gradient  
 719 of salinity

720



$$\chi^2 = 0.85, P = 0.65, \text{GFI} = 0.98, \text{RMSEA} < 0.001$$

721

722 **Fig. 5.** Path analysis detecting the underlying causal relationships between soil salinity  
 723 physicochemical factors and microbial community composition of carbon dynamics in  
 724 the soilt system. Red lines indicate positive relationships, while blue lines indicate  
 725 negative relationships. The width of arrows indicates the strength of significant  
 726 standardized path coefficients ( $P < 0.05$ ). Paths with non-significant coefficients are  
 727 presented as gray lines. \*\*\* $P < 0.001$ ; \*\* $P < 0.01$ ; \* $P < 0.05$

728