

1 **Soil priming effects and involved microbial community along salt gradients**

2
3 Haoli Zhang^{ab#}, Doudou Chang^{a#}, Zhifeng Zhu^c, Chunmei Meng^b, Kaiyong Wang^{a*}

4
5 ^a Agricultural College, Shihezi University, Shihezi 832003, China

6 ^b College of Land Science and Technology, China Agriculture University, Yuanmingyuan West Road,
7 Beijing 100193, China

8 ^cChina National Seed Group Co.Ltd, Yazhou Science and Technology City, Sanya 572000, China

9
10
11 *All correspondence to: Kaiyong Wang*

12 Email: wky20@163.com

13 [#] Both equally contribute to this work

14
15 5 Figures

16 3 Tables

17 4 Supplementary Figures

18 27 text pages

19 **Abstract**

20 Soil salinity mediates microorganisms and soil process, like soil organic carbon (SOC) cycling.
21 Yet, how soil salinity affects SOC mineralization via shaping bacterial communities diversity and
22 composition remains elusive. Therefore, soils were sampled along a salt gradient (salinity at 0.25%,
23 0.58%, 0.75%, 1.00% and 2.64%) and incubated for 90 days to investigate i) SOC mineralization (i.e.,
24 soil priming effects induced by cottonseed meal, as substrate) and ii) responsible bacteria community,
25 by using high throughput sequencing and natural abundance ¹³C isotopes (to partition cottonseed meal
26 derived CO₂ and soil derived CO₂). We observed negative priming effect during first 28 days of
27 incubation and turned to positive priming effect after day 56. Negative priming at the early stage might
28 be due to the preferential utilization of cottonseed meal. The followed positive priming decreased with
29 the increase of salinity, which might be caused by the decreased alpha diversity of microbial
30 community in soil with high salinity. Specifically, soil pH and EC along salinity gradient were the
31 dominant variables modulating the structure of microbial community and consequently SOC priming
32 (estimated by distance-based multivariate analysis and path analysis). By adopting O2PLS, priming
33 effects were linked with specific microbial taxa, e.g., *Proteobacteria* (*Luteimonas*, *Hoeflea* and
34 *Stenotrophomonas*) were the core microbial genus that attributed to the substrate induced priming
35 effects. Here, we highlight that the increase of salinity reduced the diversity of microbial community
36 and shifted dominant microorganisms(*Actinobacteria* and *Proteobacteria* (*Luteimonas*, *Hoeflea* and
37 *Stenotrophomonas*)) that determined SOC priming effects, which provides a theoretical basis for
38 understanding of SOC dynamics and microbial drivers under salinity gradient.

39

40 **Keywords:** *Salt gradient, priming effects, bacterial community, core microorganisms*

41 **1. Introduction**

42 Soil salinization is an increasing environmental problem caused by natural and
43 human activities in the arid and semi-arid area (Wichern et al., 2006). Salinization is
44 often a major threat to crop productivity in agricultural land. Soil microorganisms suffer
45 from osmotic stress. Soil salinity often cause microbial death or dormant. It was widely
46 reported that the increased salinity decrease microbial biomass, enzymatic activity, and
47 alpha diversity of microbial community (Laura, 1974; Pathak et al., 1998; Rietz et al.,
48 2003). Soil salinity is reported to the major determinants of composition, activity of
49 microbial community (Kamble et al., 2014). Although salinity is reported to be a vital
50 factor in influencing microorganisms in the arid and semi-arid area, limited studies
51 investigated C processes (e.g., priming effect) driven by microbial community in
52 salinity soils (Sardinha et al., 2003).

53 Soil organic carbon (SOC) is the largest pool (1500 Pg C) in the terrestrial carbon
54 (C) cycle, and contains twice as much C as the atmosphere (Filley et al., 2006;
55 Wiesmeier et al., 2019). The input of substrate C can influence the output (i.e., CO₂
56 release) through a phenomenon called priming effect, which was firstly discovered by
57 LÖhnis (1926). Substrate additions accelerate or decrease soil organic C mineralization,
58 referred to positive or negative priming effects (Kuzyakov et al., 2000). The intensity
59 of the priming effect affects the turnover of SOC and thus storage pool (Sullivan et al.,
60 2013). Soil priming effects are affected by many biotic and abiotic factors (Lavelle, et
61 al.;; Martin et al., 2019), to investigate abiotic and biotic mechanisms underlying SOC
62 priming enhance strong understanding of the SOC cycling.

63 Soil priming effects is affected by soil fauna animals (Scheu et al., 1994), activities,
64 diversity and composition of microbial community (; Fontaine et al., 2011). The
65 microbial decomposers are the major player in the decomposition process of added C
66 sources. The addition of substrate, such as composts (Xun et al., 2016), animal sludges
67 (Hartmann et al., 2015), sewage sludges (Su et al., 2017; Wagner et al., 2011) and plant
68 residues (Dai et al., 2017), generally increases soil microbial biomass C and stimulates

69 the microbial activities thus enhanced the loss of SOC (positive priming effects)
70 (Fontaine et al., 2003; Bird et al., 2011; Li et al., 2018; Ali et al., 2019).

71 Concerning abiotic factors, the priming effect can be controlled by climate
72 variables (Hagemann, 2008), and soil properties, like pH, EC, TN, etc (Blagodatskaya
73 et al., 2008; Luo et al., 2017). To understand how environmental and edaphic factors
74 affect the processes of SOC mineralization, is important to estimate terrestrial C pool
75 (Lehmann et al., 2015). Although many studies have tested the effects of soil pH, SOC
76 content, and other edaphic variables on soil priming effect, few study investigated soil
77 priming effects in salinity soil (Asghar et al., 2012), especially linked with soil
78 microbial community structure and their functions in C decomposition (Soina et al.,
79 2018).

80 Thus, we sampled the soils along natural salinity gradients (0.25%, 0.58%, 0.75%,
81 1.00%, 2.64% apart from total water-soluble salt). Based on these soils, we conducted
82 a 90 days of indoor incubation applying C3 substrate of cottonseed meal ($\delta^{13}\text{C}=-$
83 23.47‰) to C4 soils with salt gradient ($\delta^{13}\text{C}$ between -14.21‰ and -16.01‰), to
84 investigate: 1) mineralization rate of cottonseed meal and induced soil priming effects
85 along salt gradients; 2) diversity of microbial community in the soils with increased
86 salinity, and 3) identify the bacteria taxa associated Soil priming. We hypothesized that
87 i) soil microbial community diversity and composition will be different with the
88 different in soil variables particularly pH and EC along salinity gradients, and ii) Soil
89 C processes like priming effects will be regulated mainly by microbial community and
90 especially the core microbial species. To clarify the priming effects and involved
91 microbial groups would help us better understanding C sequestration potential and
92 underlying mechanisms in saline soils.

93

94

95 **2. Materials and methods**

96 **2.1. Soil sampling and cottonseed meal production.**

97 The soil type was gray desert soil, which was collected from farmlands (82.90° E,
98 44.96° N) in Xiao Yinpan town , Bole City, Bortala, northern Xinjiang Uygur
99 Autonomous Region, northwest China. The farmlands soil is naturally formed original
100 saline-salinity soil and with a continuous 30 years planting of maize (C4 crop) and
101 maize straw returning to soil for 7-8 year. In September 2021, we determining the
102 sampling area, and use the five-point sampling method to collecting non-rhizosphere
103 soil. The soil samples were indoor air drying and hand-picked to remove visible other
104 debris, animal and plant residues and then sieved at field moisture (<2mm) and
105 subsequently adjusted to 40% of water holding capacity (WHC). Determination of
106 five salinity gradients at 0.25%, 0.58%, 0.75%, 1.00% and 2.64% through soil salinity
107 measurements. Texture was determined by the pipette method without carbonate in all
108 soil samples. They were then incubated at 25 °C for 7 days before starting the
109 experiments, to allow any early sampling and sieving effects to subside.

110 Cottonseed meal is a kind of reddish or yellow granular material obtained by
111 pressing, leaching and other cottonseed. The cottonseed meal was purchased from the
112 market and dried at 105 °C for 24 h indoor, then further pulverized by a ball mill and
113 passed through < 2 mm sieve.

114

115 2.2. Soil and substrate analyses

116 EC and pH of soil and cottonseed meal were measured at a soil: water ratio of 1:5
117 (weight/weight) (Bao, 2000). Air-dry soil (5 g, <2 mm) and 25 ml of deionised water
118 were shaken together for 1 min and left to settle for 30 min, which was repeated once
119 more before pH was determined with a pH electrode. Soil water-soluble salt was
120 analyzed by weighted at a soil:water ratio of 1:5 (weight/weight). Air-dry soil (5 g, <1
121 mm) and 25 ml of deionised water were shaken together for 30 min, filtration to obtain
122 clear filtrate, using thermostat water bath to evaporate and weigh(Bao, 2000). Soil total
123 carbon (TC), total nitrogen (TN) are collect soil to be tested was dried and ground

124 through a 0.15mm screen, and a certain amount of treated soil sample was wrapped in
125 tin foil and placed in an element analyzer for determinatio (air-dried, milled <150 μm)
126 were determined by dry combustion (LECO CNS 2000, LECO Corporation, Michigan,
127 USA). Soil microbial biomass C was determined by fumigation extraction (Vance et
128 al., 1987; Wu et al., 1990). The K_2SO_4 extractable organic C was determined using an
129 organic carbon autoanalyser (Shimadzu, Analytical Sciences, Kyoto, Japan). Soil
130 microbial biomass C (B_c) was calculated from: $B_c = 2.22 E_c$, where $E_c = [(\text{organic C}$
131 $\text{extracted from fumigated soil}) \text{ minus } (\text{organic C extracted from non-fumigated soil})]$.
132 The natural $\delta^{13}\text{C}$ (‰) abundance of the soils (air-dried, milled <200 μm) was
133 determined using an elemental analyser-isotope ratio mass spectrometer (Sercon Ltd,
134 Crewe, UK). All measurements are given on an oven-dry weight basis (o.d., 105 °C, 24
135 h).

136 The $\delta^{13}\text{C}$ (‰) abundance of the cottonseed meal (air-dried, milled <200 μm) was
137 determined using an elemental analyser-isotope ratio mass spectrometer (Sercon Ltd,
138 Crewe, UK). The main elemental composition of the substrate was determined using
139 elemental analysis (Vario EL Cube, Hanau, Germany), with the samples combusted at
140 1200 °C. Natural $\delta^{13}\text{C}$ (‰) abundance ,the total carbon, total nitrogen contents and C/N
141 of the cottonseed meal was presented in Table 1.

142

143 2.3. Experimental design

144 After pre-incubation, five soils with salinity gradient(salinity at 0.25%, 0.58%,
145 0.75%, 1.00% and 2.64%) were thoroughly mixed with cottonseed meal at 20 mg C
146 g^{-1} soil (d.w. basis), and incubated over 90 days following moisture adjustment to 40%
147 of water-holding capacity (WHC) to investigate the substrate mineralization and
148 priming effects. Each soil sample (40 g d.w. basis) was incubated in a 100 ml beaker
149 inside a 1 L brown glass jar. Three jars with only water and NaOH were set as blank.
150 All the jars were sealed with a rubber bung and incubated in a randomized block design

151 at 25 °C for the 90 days of incubation. The NaOH vials were changed after 1, 3, 5, 7,
152 14, 28, 56 and 90 days for determination of evolved CO₂ and ¹³C–CO₂ (%). Meanwhile,
153 soil biomass C, NH₄⁺, NO₃⁻, pH, EC, TC, TN and DNA extraction were measured at
154 day 28.

155

156 2.4. Soil CO₂-C and its isotopic composition

157 Soil C evolved as CO₂-C in jars was measured by trapping CO₂ in 1 M NaOH
158 (20 ml) during soil incubation. After the NaOH (20 ml) trapping CO₂ at different
159 periods of soil incubation, 5 ml 1 M NaOH of each sample was mixed with 10 ml
160 deionised water and titrated with 0.05 M standardised HCl by the TIM840 autotitrator
161 (Radiometer Analytical, Villeurbanne Cedex, France). Meanwhile, the δ¹³C (‰) of
162 trapped CO₂-C was precipitated, with 8 ml of the 1 M NaOH (20 ml) mixed with 8 ml
163 1.5 M BaCl₂ in vials (Aoyama et al., 2000). The BaCO₃ precipitate was trapped on the
164 glass fibre the filter, rinsed with deionised water several times, and dried overnight
165 (80 °C), weighed (0.100-0.200 mg) into tin capsules, and analyzed for δ¹³C on an
166 elemental analyzer-isotope ratio mass spectrometer (Sercon Ltd, Crewe, UK).

167

168 2.5. DNA exaction and sequencing

169 The total soil DNA was extracted from 0.50 g of moist soil using a FastDNA Spin
170 Kit (MP Biomedicals, Santa Ana, CA, USA) according to the manufacturer's protocol.
171 The extracted DNA was dissolved in 50 µl of TE buffer, quantified using a
172 spectrophotometer and stored at -20 °C until sequencing.

173 V3-V4 hypervariable regions of the bacterial 16S rRNA gene were amplified with
174 primers 341F (5'-CCTAYGGRBGCASCAG-3') and 806R(5'-
175 GGACTACHVGGGTWTCTAAT-3'). The PCR reactions were conducted with a
176 thermocycler PCR system (GeneAmp 9700, ABI, USA) by using the following
177 programs: 3 min of denaturation at 95 °C; followed by 27 cycles of 30 s at 95 °C, 30 s
178 at 55 °C, and 45 s at 72 °C; and a final extension at 72 °C for 10 min with a thermocycler

179 PCR system (GeneAmp9700, ABI, USA). PCR amplicons pooled from the triplicate
 180 reactions were purified using a QIAquick PCR purification kit (Qiagen, Shenzhen,
 181 China), and quantified using a NanoDrop ND-1000 spectrophotometer (Thermo
 182 Scientific, Waltham, MA, USA). The PCR products were purified, mixed, and sent to
 183 Majorbio, Inc. (Shanghai, China) for sequencing based on the Illumina MiSeq platform.
 184

185 2.6. Calculations

186 2.6.1. CO₂-δ¹³C emission

187 The mineralisation of cottonseed meal was separated from SOC mineralisation
 188 according to the change of stable isotopic composition ($\delta^{13}C$) with time. The standard
 189 equation for determining $\delta^{13}C$ (‰) is derived from:

$$190 \delta^{13}C (\text{‰}) = [(R_{\text{sample}}/R_{\text{VPDB}}) - 1] \times 1000, \quad \text{Eqn. 1}$$

191 where R_{sample} is the mass ratio of ¹³C to ¹²C of each sample and R_{VPDB} is the
 192 international PDB(Peedee Belemnite) limestone standard. The labeled ¹³C (%) of
 193 cottonseed meal was then estimated from:

$$194 \text{CO}_2\text{-}^{13}\text{C} (\%) = (\delta_{\text{treatment}} - \delta\text{C4}) / (\delta\text{C3} - \delta\text{C4}), \quad \text{Eqn. 2}$$

195 where $\text{CO}_2\text{-}^{13}\text{C}$ (%) is the proportion of evolved CO₂ from C3 (cottonseed meal)
 196 matter, $\delta_{\text{treatment}}$ is the $\delta^{13}C$ (‰) in treatments of soil with cottonseed meal, δC4 is the
 197 $\delta^{13}C$ (‰) in control soil and δC3 is the $\delta^{13}C$ (‰) from cottonseed meal. Thus, the CO₂-
 198 C produced from cottonseed meal during the incubation was calculated from:

$$199 \text{CO}_2\text{-}^{13}\text{C} (\mu\text{g g}^{-1} \text{soil}) = \text{CO}_2\text{-}^{13}\text{C} (\%) \times \text{total CO}_2\text{-C} (\mu\text{g g}^{-1} \text{soil})/100, \quad \text{Eqn. 3}$$

200 CO₂ from SOC was $\text{CO}_2\text{-}^{13}\text{C}$ subtracted from total evolved CO₂-C. The absolute
 201 soil priming effect (or primed soil CO₂-C) with the addition of cottonseed meal was
 202 calculated from:

$$203 \text{Primed soil CO}_2\text{-C} (\mu\text{g C g}^{-1} \text{soil}) = \text{CO}_2\text{-C}_{\text{treatment}} - \text{CO}_2\text{-C}_{\text{control}} \quad \text{Eqn. 4}$$

204 where $\text{CO}_2\text{-C}_{\text{treatment}}$ is the non-isotopically labeled CO₂-C evolved from
 205 cottonseed meal amended soil, $\text{CO}_2\text{-C}_{\text{control}}$ is non-isotopically labeled CO₂-C evolved
 206 from soil without cottonseed meal.

207

208 2.7. Statistics

209 The data of ¹⁶S gene sequencing were processed using the Quantitative Insights
210 Into Microbial Ecology (QIIME) 1.9.0-dev pipeline (Caporaso et al., 2010). In brief,
211 Reads with less than length 200 bp and ambiguous bases were discarded. The sequences
212 were then binned into operational taxonomic units (OTUs) by UCLUST (Edgar, 2010)
213 based on 97% pairwise identity. Chimeric OTUs identified by USEARCH (Edgar et al.,
214 2011) in QIIME were removed. The most abundant sequence from each OTU was
215 selected to represent that OTU. Taxonomy was assigned to 16S OTUs against a subset
216 of the Silva 104 database. The representative OTU sequences were aligned using
217 PyNAST (Caporaso et al., 2010). We obtained between 64,425 and 89,989 clean_reads
218 per sample for all experimental samples.

219 To avoid potential bias caused by sequencing depth, all sample datasets were
220 rarefied for the bacteria α -diversity and β -diversity analyses. Faith's phylogenetic
221 diversity was calculated to provide an integrated index of the phylogenetic breadth
222 across taxonomic levels (Faith, 1992). To compare β -diversity between samples,
223 principal coordinate analyses based on the unweighted and weighted UniFrac
224 (Lozupone et al., 2007a) distances were calculated using the function 'pcoa' in the R
225 package 'Ape'. Additionally, permutational multivariate analysis of variance
226 (PERMANOVA) was carried out using the function 'adonis' in the R 'vegan' to
227 measure effect size and significance on β -diversity. The variable influence projection
228 (VIP) value was processed using the way of O2PLS analysis by the SIMCAP 14
229 (Version 14.1.0.2047) (Wang et al., 2016). The y-matrix was defined as the
230 environmental factors datasets and the x-matrix was defined as the microbial
231 community on genus level dataset.

232 Data were logarithmically transformed and analyzed by ANOVA. All analyses
233 were performed using SPSS software (13th edition). Pearson's correlation analyses were
234 performed to assess the linear correlation among soil physio-chemical properties and

235 microbial community. MULTIVARIATE analysis were operated to investigate
236 interaction of salinity treatments on bacteria community parameters.

237

238 **3. Results**

239 3.1. Soil physicochemical properties along salt gradients

240 The major soil physicochemical properties along salt gradients were presented
241 (Table 1) and all of soil physicochemical properties has significant difference ($P <$
242 0.05). The total soluble salinity content in the soils ranged from 0.25% to 2.64% of
243 salinity soils, soil salt gradients increasing gradually from salinity 1 samples to salinity
244 5 samples. The pH and EC in soils ranged from 8.45 to 8.85 and from 1.06 ms cm^{-1} to
245 7.75 ms cm^{-1} . Soil total C and N were increased with salinity, ranging from 3.16% to
246 3.57%, and from 0.18% to 0.26%. The $\delta^{13}\text{C}$ value for soils are between -14.21‰ and -
247 16.01‰, which were relatively enriched compared to cottonseed meal (-23.47‰). This
248 allowed separation of soil derived CO_2 from total evolved CO_2 , according to the classic
249 mixed modeling.

250

251 3.2. Total CO_2 evolution

252 During the whole 90 days of incubation, the cumulative CO_2 evolved had
253 similar trends, which the amount of CO_2 increased with the incubation times (Fig. S1).
254 The cumulative CO_2 evolved increased more rapidly with the addition of cottonseed
255 meal before 14 days, compared to non-amended soils. At 90 days of incubation. The
256 cumulative CO_2 evolved in the soil with the lowest salinity (Salinity 1) gave the lowest
257 CO_2 emission (597 $\mu\text{g C g}^{-1}$) in the non-amended soils (Fig. S1, $P < 0.001$).

258

259 3.3. Cottonseed derived $^{13}\text{CO}_2$ and soil priming effects

260 The total cumulative $\text{CO}_2\text{-C}$ was divided three parts based the $\delta^{13}\text{C}$ value,
261 including basal soil-derived CO_2 , cottonseed meal-derived CO_2 and primed soil CO_2
262 (Fig.1). The cottonseed meal-derived CO_2 had a significant contribution to the total CO_2

263 evolved during the early incubation period. The cottonseed meal-derived CO₂ was
264 significantly higher in Salinity 1, Salinity 2 and Salinity 3 than in Salinity 4 and Salinity
265 5 before 28 days incubation. Meanwhile, the soil priming effects was negative in all
266 amended soil treatments before 28 days incubation and the direction of priming effect
267 in most of soil samples turned into positive after 28 days. During the whole 90 days
268 incubation, there was a negative correlation between cottonseed meal-derived CO₂ and
269 primed soil CO₂ (Fig. 2).

270

271 3.4. Bacterial diversity and community structure

272 The number of sequences ranged from 64,425 to 91,261 for per sample (average
273 value of 80,602). About 27,990 OTUs in total were obtained under different five
274 treatments. Bacterial community diversity was measured by a series of OTU-based
275 analyses of alpha diversity including chao1 estimator, and observed_species in the
276 QIIME pipeline (Fig. 3). Chao1 diversity estimator and observed_species was
277 significantly different in treatments, being the highest in Salinity 1, followed by Salinity
278 3, Salinity 2, Salinity 4 and Salinity 5 ($P < 0.01$). In general, bacterial community
279 diversity decreased with increasing salinity (Fig. 3).

280 The most abundant phylum in the soils and their correlation with salinity were
281 shown in Fig. 4. Among them, *Actinobacteria* was the dominant taxa in all soils, with
282 the abundance ranging from 50.07 % (Salinity 3) to 68.99 % (Salinity 4). The relative
283 abundance of *Bacteroidetes*, *Firmicutes*, and *Deinococcus-Thermus* increased with
284 the salinity, while *Acidobacteria* decreased with salinity degree.

285 Based on OTUs of five gradient salt treatments, the PCA analysis showed that
286 treatments from Salinity 2 and Salinity 4 clustered together. Meanwhile, soil samples
287 of Salinity 1, Salinity 3 and Salinity 5 distributed in the first, fourth and three quadrant,
288 which indicated that these treatments had large environmental heterogeneity (Fig. S4).

289 In order to visualize the relationship between environmental factors and microbial
290 community, *Canonical Correspondence Analysis* (CCA) was conducted, showing that

291 NO₃⁻-N, EC and TC had a more obvious impact than other factors for microbial
292 community (Fig. 3). Soil EC were positively correlated with pH, NH₄⁺-N, and
293 negatively correlated with TN, TC and MBC. Mantel test and Distance-based
294 multivariate analysis showed the contribution rate of different environmental factors
295 account for 78% of the variability of microbial communities (Table 2). The value of pH
296 (31%) and EC (12%) had a strong influence on microbial community.

297

298 3.5. Relation between soil microbial community and C dynamics

299 Based on the O2PLS analysis, the variable influence projection (VIP) values of
300 bacterial genus more than 1.00% were showed their contributions to C decomposition
301 of cottonseed meal-derived C, basal soil-derived C, and primed soil C (Table 3). There
302 were many microbial taxa positively correlating to soil primed CO₂, for insatnce, genera
303 of *Actinomarinales*, *Luteimonas*, *Nocardioides*, *Hoeflea*, *Intrasporangium*,
304 *Nitrolancea*, *Pseudarthrobacter* and *Stenotrophomonas* had a positive correlation with
305 primed CO₂. In order to further to evaluate the relationship between soil properties, soil
306 bacterial communities and C decomposition, we used the structural equation modeling
307 (SEM) to suggest the direct and indirect impacts of salinity and microbial community
308 on soil C decomposition (Fig. 7). The result showed that soil pH and EC had negative
309 contribution to bacterial diversity, while bacterial diversity had a strong positive
310 influence on the primed soil C (Fig. 5). For instance, salinity properties of EC had a
311 directly negative influence on the bacterial diversity but positive influence on the
312 primed soil C. Meanwhile, pH were negatively correlated with bacterial diversity and
313 positively correlated with substrate derived C.

314

315 4. Discussion

316 4.1. Soil priming effects along salty gradients

317 Understanding soil C dynamics along salinity gradients is crucial to predict C
318 sequestration in salty soils. In the early stage of the incubation, we observed that the

319 cumulative substrate derived CO₂ in the soils with lower salinity was significantly
320 higher than soils with higher salinity (Fig. 1), which can be possibly explained by that
321 high salinity inhibited microbial activity. Many studies have reported the influence of
322 soil salinity on organic matter decomposition, mostly, the decomposition of organic
323 matter are decreased by salinity (Wichern et al., 2006; Ghollarata et al., 2007; Tripathi
324 et al., 2007; Setia et al., 2012). Yet, the response of microbial community to the
325 increasing levels of salinity and consequent effects on soil priming effects remains
326 largely unknown.

327 Here, we found soil priming effects was gradually changed from negative to
328 positive priming effect (Fig. 1). The early pattern of the dynamics of the priming effect
329 in this study was similar to other studies showing preferential utilization of labile C
330 substance. The first phase of negative priming effects was likely to be caused by
331 microbial assimilation of substrate. The soil microbes turned to use the new added
332 substrate and thus used less of the original SOC. This was attributed to “preferential
333 substrate utilization” (Perelo et al., 2005).

334 Soil microbial biomass-related growth predominating in the first phase were most
335 likely to utilize SOC, leading to a positive priming effects after substrate was largely
336 vanished. The magnitude of priming effects depends on soil microbial biomass size
337 (Schneckenberger et al., 2008). It was found that the amount of added easily available
338 organic C is beyond 50% of microbial biomass C (Blagodatskaya et al., 2008). Namely,
339 the second phase of positive PEs probably was due to increased biomass size and
340 enhanced demand on SOC. Secondly, C that was assimilated into microbial biomass in
341 the first stage may also be mineralized in the second stage due to the turnover of
342 microbial biomass (Shahbaz et al., 2017; Perelo et al., 2005).

343

344 4.2. Microbial community along salt gradients

345 Previous studies concerning the impact of salinity on soil microbial community
346 used different soils with a range of salt levels. In the present study we investigated the

347 influence of soil salinity on microbial communities in soils from the closed area
348 covering a range of salt content. Similarly, Rousk et al. (2011) also used agricultural
349 soils from the same area representing a range of soil salinity. Here, we found microbial
350 diversity (alpha diversity) decreased with increasing salinity (Fig. 3). The negative
351 impact on microbial diversity can be explained by that the accumulation of large
352 amounts of salt in the soil raised the extracellular osmotic concentration (Rath et al.,
353 2015; Oren, 2011). The high osmotic pressures made it difficult for many
354 microorganisms to adapt to and thus reduce their biological activity. The changes of
355 soil microbial community structure were also explained by salinity (Herlemann et al.,
356 2011; Campbell et al., 2013). We found that *Bacteroidetes*, *Firmicutes*, *Acidobacteria*
357 and *Deinococcus-Thermus* were dominant in these soils (Fig. 4). These results are
358 supported by previous findings that *Firmicutes* possess the high salinity resistance.
359 Other studies also found that *Bacteroidetes* is dominant taxa in alkaline saline soil
360 because of its resistant to salt (Valenzuela-Encinas et al., 2009; Keshri et al., 2013).
361 Other study shows that the dominant phyla are *Bacteroidetes* and followed by
362 *Proteobacteria* in the haloalkaline soil (Keshri et al., 2013). These results are consistent
363 with the esuarine or marine environments, despite some studies suggest that soil salinity
364 is not found to be a decisive factor for bacterial community and their growth (Rousk et
365 al., 2011).

366 The difference of microbial community structure is affected by many soil variables,
367 and pH and EC were the most important ones (Fig. 3; Table 2). Our results showed that
368 the value of soil pH and EC would significantly affect the microbial community
369 structure and the combined contribution rate of these two variables to microbial
370 community was 43% (Table 2). At high levels of salt and alkaline arid condition, soil
371 pH has been also shown to have a very powerful influence on the soil bacterial
372 community structures (Bååth et al., 2003; Fierer et al., 2006; Rousk et al., 2010).
373 Meanwhile, it is consequently unlikely that soil pH differences between the studied
374 soils obscured the influence of salt (Rousk et al., 2011). Salinity has been identified as

375 one of the most potent environmental factors that determine assembly of microbiome.
376 Salinity has been regarded to play the vital role in shaping microbial community in
377 different ecosystem. This, despite the clear evidence from aquatic microbial ecology
378 (Lozupone et al., 2007b), show a potential for salt to affect soil microbial communities
379 apart from that of pH (Rath et al., 2015).

380

381 4.3. The core microbial taxa regulating C decomposition along salinity gradient

382 The correlation of microbial taxa and SOC decomposition (priming) were found
383 according to the results of O2PLS and SEM (Table 3; Fig. 5). Here we showed that
384 *Streptomyces* (*Actinobacteria*), *Glycomyces* (*branch of Actinobacteria*), *Agromyces*
385 (*branch of Actinobacteria*), and *Sphingomonas* (*branch of Proteobacteria*) at the genus
386 level were significantly correlated with the C process particularly primed soil-driven C.
387 Most of these functional taxa belonged to *Actinobacteria* and *Proteobacteria*. In a
388 recent study, Ren et al. (2018) found that *Actinobacteria* had negative impact on SOC
389 mineralization across land-use change (Fierer et al., 2007; Goldfarb et al., 2011) and
390 *Proteobacteria* drove the positive soil respiration (He et al., 2012; Stevenson et al.,
391 2004), indicating the balance of soil C dynamics were largely regulated by these two
392 phyla. We found similar result that *Streptomyces* (*branch of Actinobacteria*) had a
393 negative correlation with primed soil CO₂. *Actinobacteria* are able to grow
394 preferentially on the C-rich refractory materials and relatively easily decompose the
395 cellulose, lignocellulose (Khodadad et al., 2011), indicating these microorganisms
396 preferentially use the C source that is used partially by others.

397 Although some studies suggest soil salinity may not be a vital factor for C
398 decomposers (Rousk et al., 2011), the composition of microbial community are
399 considered to play a decisive role in determining C dynamic processes in response to
400 salt stress (Ramsey et al., 2005; Schimel et al., 2007; Nottingham et al., 2009). Here,
401 SEM analysis showed that soil pH and EC in salted soils reduced microbial diversity
402 and thus limited the utilization of SOC by microbial community, It was reported that

403 high pH and salinity are the major determinants of soil microbial activity and
404 community structure (Kamble et al., 2014).

405

406 **5. Conclusion**

407 Cotton meal is a kind of organic material with high nitrogen content, adding cotton
408 meal in salinised soil can stimulate and promote the release of soil nutrients. The
409 microorganisms mainly use the organic matter in the cotton meal in the pre-culture
410 period, so the soil carbon excitation is negative excitation, Soil priming effect turned
411 from negative to positive at the later stage of incubation (day 28), because
412 microorganisms turned to decompose SOC from the labile substrate. With the increase
413 of salinity, the diversity of microbial community decreased. Soil microbial community
414 was mainly controlled by soil pH and EC. By O2PLS, we found *Actinobacteria* and
415 *Proteobacteria (Luteimonas, Hoeflea and Stenotrophomonas)* dominant in these soils
416 were the core microbial taxa that affecting the process of organic C mineralization,
417 particularly soil primed CO₂.

418

419 **Acknowledgements**

420 This study was supported by the Special Fund for Key Science & Technology
421 Program in Xinjiang Province of China (No. 2022B02021-3-1) .

422

423 **Data availability**

424 The datasets used and analysed during the current study available from the
425 corresponding author on reasonable request.

426

427 **Author contributions**

428 K.W. conceptualized and conducted the experiment. H.Z. and D.C. conducted the
429 data analysis and wrote the manuscript, conducted the indoor experiment. C.M. and

430 Z.Z. assisted in conducting the experiment. All authors reviewed the manuscript. All
431 authors contributed to the manuscript and approved the submitted version.

432

433 **Competing interests**

434 The authors declare no competing interests.

435

436 **Reference**

437 Anderson, C.R., Condon, L.M., Clough, T.J., Fiers, M., Stewart, A., Hill, R.A.,

438 Sherlock, R.R., 2011. Biochar induced soil microbial community change:
439 implications for biogeochemical cycling of carbon, nitrogen and
440 phosphorus. *Pedobiologia*, 54, 309-320.

441 Asghar, H. N., Setia, R., Marschner, P., 2012. Community composition and activity of
442 microbes from saline soils and non-saline soils respond similarly to changes in
443 salinity. *Soil Biology and Biochemistry*, 47, 175-178.

444 Ali N., Khan S., Li Y., Zheng N., Yao H., 2019. Influence of biochars on the
445 accessibility of organochlorine pesticides and microbial community in
446 contaminated soils. *Science of The Total Environment*, 647, 551-560.

447 Bååth, E., Anderson, T.H., 2003. Comparison of soil fungal/bacterial ratios in a pH
448 gradient using physiological and PLFA-based techniques. *Soil Biology and*
449 *Biochemistry*, 35, 955-963.

450 Bao, S.D., 2000. *Soil and Agricultural Chemistry Analysis*, third ed. China Agriculture
451 Press, Beijing.

452 Blagodatskaya, E., Kuzyakov, Y., 2008. Mechanisms of real and apparent priming
453 effects and their dependence on soil microbial biomass and community structure:
454 critical review. *Biology and Fertility of Soils*, 45, 115-131.

455 Bird, J.A., Herman, D.J., Firestone, M.K., 2011. Rhizosphere priming of soil organic
456 matter by bacterial groups in a grassland soil. *Soil Biology and Biochemistry*, 43,
457 718-725.

458 Campbell, B.J., Kirchman, D.L., 2013. Bacterial diversity, community structure and
459 potential growth rates along an estuarine salinity gradient. *The ISME Journal*, 7,
460 210-220.

461 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,
462 E.K., Fierer, N., P?a, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley,
463 S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D.,
464 Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters,
465 W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows
466 analysis of high-throughput community sequencing data. *Nature Methods*, 7, 335-
467 336.

468 Dai, H., Chen, Y., Yang, X., Cui, J., Sui, P., 2017. The effect of different organic
469 materials amendment on soil bacteria communities in barren sandy loam
470 soil. *Environmental Science Pollution Research*, 24, 1-10.

471 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST.
472 *Bioinformatics*, 26, , 2460-2461.

473 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME
474 improves sensitivity and speed of chimera detection. *Bioinformatics*, 27, 194-220.

475 Faith D.P., 1992. Conservation evaluation and phylogenetic diversity. *Biological*
476 *Conservation*, 61, 1-10.

477 Filley, T.R., Boutton, T.W., 2006. Ecosystems in flux: molecular and stable isotope
478 assessments of soil organic matter storage and dynamics. *Soil Biology and*
479 *Biochemistry*, 38, 3181-3183.

480 Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial
481 communities. *Proceedings of the National Academy of Sciences of the USA*, 103,
482 626-631.

483 Fierer, N., Bradford, M., Jackson, R., 2007. Toward an ecological classification of soil
484 bacteria. *Ecology*, 88, 1354-1364.

485 Fontaine, S., Mariotti, A., Abbadie, L., 2003. The priming effect of organic matter: A
486 question of microbial competition?. *Soil Biology and Biochemistry*, 35, 837-843.

487 Fontaine, S., Henault, C., Aamor, A., Bdioui, N., Bloor, J.M.G., Maire, V., Mary, B.,
488 Revaillet, S., Maron, P.A., 2011. Fungi mediate long term sequestration of carbon
489 and nitrogen in soil through their priming effect. *Soil Biology and Biochemistry*,
490 4, 86-96.

491 Goldfarb, K., Karaoz, U., Hanson, C., Santee, C., Bradford, M., Treseder, K.,
492 Wallenstein, M., Brodie, E., 2011. Differential growth responses of soil bacterial
493 taxa to carbon substrates of varying chemical recalcitrance. *Frontiers in*
494 *Microbiology*, 2, 94-97.

495 Ghollarata, M., Raiesi, F., 2007. The adverse effects of soil salinization on the growth
496 of trifolium alexandrinum l. and associated microbial and biochemical properties
497 in a soil from iran. *Soil Biology and Biochemistry*, 39, 1699-1702.

498 Hagemann, S., 2008. Vulnerability of permafrost carbon to climate change:
499 Implications for the global carbon cycle. *Bioscience*, 701-714.

500 Hartmann, M., Frey, B., Mayer, J., Widmer, F., 2015. Distinct soil microbial diversity
501 under long-term organic and conventional farming. *The ISME Journal*, 9, 1177-
502 1194.

503 He, Z., Piceno, Y., Deng, Y., Xu, M., Lu, Z., DeSantis, T., Andersen, G., Hobbie, S.E.,
504 Reich, P.B., Zhou, J., 2012. The phylogenetic composition and structure of soil
505 microbial communities shifts in response to elevated carbon dioxide. *The ISME*
506 *journal*, , 259-272.

507 Herlemann, D. P., Labrenz, M., Jürgens, Klaus, Bertilsson, S., Waniek, J. J., Andersson,
508 A. F., 2011. Transitions in bacterial communities along the 2000 km salinity
509 gradient of the baltic sea. *The ISME Journal*, , 1571-1579.

510 Kuzyakov, Y., Friedel, J.K., Stahr, K., 2000. Review of mechanisms and quantification
511 of priming effects. *Soil Biology and Biochemistry*, 32, 1485-1498.

512 Khodadad, C.L.M., Zimmerman, A.R., Green, S.J., Uthandi, S., Foster, J.S., 2011.

513 Taxa-specific changes in soil microbial community composition induced by
514 pyrogenic carbon amendments. *Soil Biology and Biochemistry*, 4, 385-392.

515 Keshri, J., Mody, K., Jha, B.. 2013. Bacterial community structure in a semi-arid
516 haloalkaline soil using culture independent method. *Geomicrobiology Journal*, 3,
517 517-529.

518 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature*,
519 528, 60-68.

520 Löhnis, F., 1926. Nitrogen availability of green manures. *Soil Science*, 22, 253-290.

521 Sullivan, B. W., Hart, S. C., 2013. Evaluation of mechanisms controlling the priming
522 of soil carbon along a substrate age gradient. *Soil Biology and Biochemistry*, 5,
523 293-301.

524 Lozupone, C.A., Hamady, M., Kelley, S.T., Knight, R., 2007a. Quantitative and
525 qualitative beta diversity measures lead to different insights into factors that
526 structure microbial communities. *Applied and Environmental Microbiology*, 73,
527 1576-1585.

528 Lozupone, C.A., Knight, R., 2007b. Global patterns in bacterial diversity. *Proceeding*
529 *of the National Academy of Science*, 104, 11436-11440.

530 Lavelle, P., Bignell, D., Lepage, M., Wolters, V., Roger, P., Ineson, P., Heal, O.W.,
531 Dhillon, S., 1997. Soil function in a changing world: the role of invertebrate
532 ecosystem engineers. *European Journal of Soil Biology*, 33, 159-193.

533 Luo, Y., Zang, H., Yu, Z., Chen, Z., Gunina, A., Kuzyakov, Y., Xu, J., Zhang, K.,
534 Brookes, P.C., 2017. Priming effects in biochar enriched soils using a three-
535 source-partitioning approach: ¹⁴C labelling and ¹³C natural abundance. *Soil*
536 *Biology and Biochemistry*, 106, 28-35.

537 Li X., Chen Q.L., He C., Shi Q., Chen S.C., Reid B.J., Zhu Y.G., Sun G.X., 2018.
538 Organic Carbon Amendments Affect the Chemodiversity of Soil Dissolved
539 Organic Matter and Its Associations with Soil Microbial Communities.
540 *Environmental Science and Technology*, 53, 50-59.

541 Laura, R.D., 1974. Effects of neutral salts on carbon and nitrogen mineralisation of
542 organic matter in soil. *Plant and Soil*, 41, 113-127.

543 Nottingham, A.T., Griffiths, H., Chamberlain, P.M., Stott, A.W., Tanner, E.V.J., 2009.
544 Soil priming by sugar and leaf-litter substrates: a link to microbial groups. *Applied*
545 *Soil Ecology*, 42, 183-190.

546 Oren, A., 2011. Thermodynamic limits to microbial life at high salt concentrations.
547 *Environmental Microbiology*, 13, 1908-1923.

548 Perelo, L.W., Jimenez, M., Munch, J.C., 2005. Microbial immobilisation and turnover
549 of ¹⁵N labelled substrates in two arable soils under field and laboratory conditions.
550 *Soil Biology and Biochemistry*, 38, 912-922.

551 Pathak, H., Rao, D.L.N., 1998. Carbon and nitrogen mineralization from added organic
552 matter in saline and alkaline soils. *Soil Biology and Biochemistry*, 30, 695-702.

553 Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight,
554 R., Fierer, N., 2010. Soil bacterial and fungal communities across a pH gradient
555 in an arable soil. *The ISME Journal*, 4, 1340-1351.

556 Rousk, J., Elyaagubi, F.K., Jones, D.L., Godbold, D.L., 2011. Bacterial salt tolerance
557 is unrelated to soil salinity across an arid agroecosystem salinity gradient. *Soil*
558 *Biology and Biochemistry*, 43, 1881-1887.

559 Ramsey, P.W., Rillig, M.C., Feris, K.P., Gordon, N.S., Moore, J.N., Holben, W.E.,
560 Gannon, J.E., 2005. Relationship between communities and processes; new
561 insights from a field study of a contaminated ecosystem. *Ecology Letter*, 8, 1201-
562 1210.

563 Ren, C., Wang, T., Xu, Y., Deng, J., Zhao, F., Yang, G., Han, X., Feng, Y., Ren, G.,
564 2018. Differential soil microbial community responses to the linkage of soil
565 organic carbon fractions with respiration across land-use changes. *Forest Ecology*
566 *and Management*, 12, 170-178.

567 Rath, K.M., Rousk, J., 2015. Salt effects on the soil microbial decomposer community
568 and their role in organic carbon cycling: A review. *Soil Biology and*
569 *Biochemistry*,81, 108-123.

570 Shahbaz, M., Kuzyakov, Y., Sanauallah, M., Heitkamp, F., Zelenev, V., Kumar, A.,
571 Blagodatskaya, E., 2017. Microbial decomposition of soil organic matter is
572 mediated by quality and quantity of crop residues: mechanisms and thresholds.
573 *Biology and Fertility of Soils*, 53, 287-301.

574 Schneckenberger, K., Demin, D., Stahr, K. and Kuzyakov, Y. 2008. Microbial
575 utilization and mineralization of ¹⁴C glucose added in six orders of concentration
576 to soil. *Soil Biology and Biochemistry*, 40,1981-1988.

577 Schimel, J., Balser, T.C., Wallenstein, M., 2007. Microbial stress–response physiology
578 and its implications for ecosystem function. *Ecology*, 88, 1386-1394.

579 Stevenson, B.S., Eichorst, S.A., Wertz, J.T., Schmidt, T.M., Breznak, J.A., 2004. New
580 strategies for cultivation and detection of previously uncultured microbes. *Applied*
581 *and Environmental Microbiology*, 70(8), 4748-4755.

582 Rietz, D.N., Haynes, R.J., 2003. Effects of irrigation-induced salinity and sodicity on
583 soil microbial activity. *Soil Biology Biochemistry*, 35, 845-854.

584 Scheu, S., Parkinson, D., 1994. Effects of earthworms on nutrient dynamics, carbon
585 turnover and microorganisms in soils from cool temperate forests of the Canadian
586 Rocky Mountains - laboratory studies. *Applied Soil Ecology*, 5, 23-27.

587 Su, J. Q., An, X. L., Li, B., Chen, Q. L., Gillings, M. R., Chen, H., 2017. Metagenomics
588 of urban sewage identifies an extensively shared antibiotic resistome in
589 china. *Microbiome*, 5, 84-85.

590 Soina V.S., Mergelov N.S., Kudinova A.G., Lysak L.V., Demkina E.V., Vorobyova
591 E.A., Dolgikh A.V., Shorkunov I.G., 2018. Microbial Communities of Soils and
592 Soil-like Bodies in Extreme Conditions of East Antarctica. *Paleontological*
593 *Journal*, 52, 1186-1195.

594 Setia, R., Setia, D., Marschner, P., 2012. Short-term carbon mineralization in saline-
595 sodic soils. *Biology and Fertility of Soils*, 48, 475-479.

596 Sardinha, M., Müller, T., Schmeisky, H., Joergensen, R.G., 2003. Microbial
597 performance in soils along a salinity gradient under acidic conditions. *Applied Soil*
598 *Ecology*, 23, 237-244.

599 Tripathi, S., Chakraborty, A., Chakrabarti, K., Bandyopadhyay, B.K., 2007. Enzyme
600 activities and microbial biomass in coastal soils of India. *Soil Biology and*
601 *Biochemistry*, 39, 2840-2848.

602 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for
603 measuring soil microbial biomass C. *Soil Biology and Biochemistry*, 19, 703-707.

604 Valenzuela-Encinas, C., Neria-González, I., Alcántara-Hernández, R.J., Estrada-
605 Alvarado, I., Dendooven, L., Marsch, R., 2009. Changes in the bacterial
606 populations of the highly alkaline saline soil of the former lake Texcoco (Mexico)
607 following flooding. *Extremophiles*, 13, 609-621.

608 Wang, Z.M., Lu, Z.M., Shi, J.S., Xu, Z.H., 2016. Exploring flavour-producing core
609 microbiota in multispecies solid-state fermentation of traditional Chinese vinegar.
610 *Scientific Report*, 31, 6, 26818.

611 Martin, W., Livia, U., Hobley, E., Lang, B., von Ltzow, M., Marin-Spiotta, E., Van
612 Wesemael, B., Rabot, E., Lie, M., Garcia-Franco, N., Wollschlger, U., Vogel, H.J.,
613 Kgel-Knabner, I., 2019. Soil organic carbon storage as a key function of soils - A
614 review of drivers and indicators at various scales. *Geoderma*, 149-162..

615 Wagner, B., Raquel, G., 2011. Impacts of sewage sludge in tropical soil: a case study
616 in brazil. *Applied and Environmental Soil Science*, 2011, 1-11.

617 Wichern, J., Wichern, F., Joergensen, R.G., 2006. Impact of salinity on soil microbial
618 communities and the decomposition of maize in acidic soils. *Geoderma*, 137, 100-
619 108.

620 Wu, J., Joergensen, R.G., Pommerening, B., Chaussod, R., Brookes, P.C., 1990.
621 Measurement of soil microbial biomass C by fumigation-extraction: an automated
622 procedure. *Soil Biology and Biochemistry*, 22, 1167-1169.

623 Xun, W., Zhao, J., Xue, C., Zhang, G., Ran, W., Wang, B., Shen, Q., Zhang, R., 2016.
624 Significant alteration of soil bacterial communities and organic carbon
625 decomposition by different longterm fertilization management conditions of
626 extremely lowproductivity arable soil in South China. *Environmental*
627 *Microbiology*, 18, 1907-1917.

628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657

658
659
660
661
662
663
664
665
666
667

Table 1. Soil samples and Cottonseed meal properties

	Salinity 1	Salinity 2	Salinity 3	Salinity 4	Salinity 5	Cottonseed meal
Total C (%)	3.38b	3.18c	3.16c	3.57a	3.35b	42.98
Total N (%)	0.18d	0.19d	0.20c	0.22b	0.26a	5.84
C/N ratio	18.32a	16.56b	15.71c	16.54b	12.94d	7.38
$\delta^{13}\text{C}$ value (‰)	-14.21a	-14.79c	-14.60b	-14.55b	-16.01d	-23.47
pH (H₂O)	8.85a	8.45c	8.58b	8.59b	8.55b	7.63
EC (dS m⁻¹)	1.06e	1.96c	1.28d	2.64b	7.75a	2.56
Salinity (%)	0.25e	0.58d	0.75c	1.00b	2.64a	ND

668
669

670 **Table 2.** Mantel test and Distance-based multivariate analysis relevance and
 671 contribution rate between soil properties and bacterial community compositions.

	pH	EC	NO ₃ ⁻ -N	NH ₄ ⁺ -N	MBC	TN	TC
Correlation	0.74**	0.56**	0.36**	0.68**	0.31**	0.11	0.27
Contribution	0.31**	0.12**	0.05	0.04	0.16	0.03	0.07**

672 Note: * $P < 0.05$, ** $P < 0.01$

673

674

675 **Table 3.** The variable influence projection (VIP) value and Spearman's correlation
 676 between the relative abundances of genera and C dynamic.

Phylum-Genus	VIP	Cottonseed meal CO ₂ -C(μg g ⁻¹)	Primed soil CO ₂ -C(μg g ⁻¹)	Basal soil CO ₂ -C(μg g ⁻¹)
<i>Actinobacteria-Actinomarinales</i>	1.36		0.63**	
<i>Proteobacteria-Luteimonas</i>	1.31		0.80**	
<i>Actinobacteria-Nocardioiodes</i>	1.30		0.54*	
<i>Proteobacteria-Hoeflea</i>	1.29		0.73**	
<i>Actinobacteria-Streptomyces</i>	1.27		-0.84**	
<i>Actinobacteria-Glycomyces</i>	1.26	0.63**		
<i>Actinobacteria-Marmoricola</i>	1.26	-0.52		
<i>Proteobacteria-Nitrospira</i>	1.23		0.59	
<i>Actinobacteria-Intrasporangium</i>	1.22		0.60*	
<i>Actinobacteria-Agromyces</i>	1.19			0.58*
<i>Proteobacteria-Sphingomonas</i>	1.18			0.65**
<i>Actinobacteria-Myceligenans</i>	1.16			
<i>Chloroflexi-Nitrolancea</i>	1.15		0.65**	
<i>Actinobacteria-Pseudarthrobacter</i>	1.06		0.62**	
<i>Proteobacteria-Stenotrophomonas</i>	1.00	-0.50	0.72**	

677 Note:* $P < 0.05$, ** $P < 0.01$

678


679

680

681

682

683



684

685

Fig. 1. Partitioning of CO₂ evolution after addition of cottonseed meal in different five salinity soils. Cumulative CO₂ evolved from salinity soil of 0.25 % (a) , 0.58 % (b) , 0.75 % (c) , 1.00% (d) and 2.64% (e) . Error bars represent standard errors of the means (n = 3).


688

689

690


691

692



693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706

Fig. 2. Correlation between primed soil mineralisation and cottonseed meal mineralisation following different five salinity soils during 90 days incubation



707

708 **Fig. 3.** Microbial community alpha diversity (Chao1) observed_species and beta
 709 diversity. Within each panel, boxplot data refer to maximum date (top line), 99% (the
 710 second line), mean (the third line), 1% (the fourth line) and minimum date (bottom line)
 711 of the different treatments, with statistical significance ($P < 0.05$).


712

713

714


715

716



717
718
719
720

Fig. 4. The top 10 of phylums and genes in bacterial community in soils with a gradient of salinity



$\chi^2 = 0.85, P = 0.65, GFI = 0.98, RMSEA < 0.001$

721

722 **Fig. 5.** Path analysis detecting the underlying causal relationships between soil salinity
 723 physicochemical factors and microbial community composition of carbon dynamics in
 724 the soil system. Red lines indicate positive relationships, while blue lines indicate
 725 negative relationships. The width of arrows indicates the strength of significant
 726 standardized path coefficients ($P < 0.05$). Paths with non-significant coefficients are
 727 presented as gray lines. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$

728