

REVIEWER 1

Excellent manuscript in isotopic applications in archaeology because clearly, the authors have considered the assumptions related to the analytical techniques applied, animal physiology, and all the caveats and careful considerations that are necessary for properly using SI in archaeology. The authors successfully extracted as much information as they could from ungulate tooth enamel using carbon and oxygen isotopes adding more valuable environmental and ecological information in northern and northeastern Spain, regions where such records are growing, providing detailed scenarios for the region during the MP-UP transition. This manuscript is well-written and well-structured. Data are presented well in tables and figures. Besides a few points that need clarification, the article should be accepted for publication.

Line 95: the word “approach” is too general, be more specific.

Line 98: Refrain from using word correlate unless speaking about statistical correlations.

Fig. 2- why is only one circle around 40,000 years ago green and all the others black, needs to be explained in the caption.

Dear reviewer,

We greatly appreciate your positive assessment of our contribution. Regarding the proposed suggestions, we have implemented the following corrections:

1) We have changed “approach” to “characterise”

2) We have changed “correlate” to “relate”

3) We have include a clarification in the figure’s caption, indicating that the green color in the figure represents the site in the Mediterranean/northeastern area (Canyars). The color code corresponds to the chronology of the technocomplexes as indicated in the lower margin.

Thanks for your suggestions.

(Citation: <https://doi.org/10.5194/bg-2023-128-AC1>)

REVIEWER 2

This is a remarkable manuscript that deserves publication after major changes.

From my point of view there are some overinterpretations on the one hand, and some oversimplifications on the other hand. An important problem of paleoenvironmental reconstructions based only on archaeological remains, like this manuscript, is that the obtained data are not in an objective paleoclimatic and paleoenvironmental frame, in this case (northern) Iberia. This study based the paleoenvironmental reconstructions mainly on the archaeological data instead of comparing them with regional continuous paleoclimatic and paleoenvironmental records. I mean, authors suggested that this type of paleoenvironmental studies is key to understand past climate and human interactions. See for example abstract lines 18-19 or the introduction. Authors must keep in mind that the paleoenvironmental reconstruction that they have performed in these archaeological sites are “discontinuous points” in the paleoclimatic record of the Iberian Peninsula (see for example the chronologies in Fig.2). I mean, the most accurate climatic records for the studied periods are marine, lacustrine and speleothem records, where one can observe the “objective” fluctuations of past climate. Authors should acknowledge this point in the manuscript as well as compare/discuss with these records (there are many for the IP, for example Martrat et al., 2004; 2007; Pérez Mejías et al., 2019; Moreno et al., 2012; González-Sampériz et al., 2020; Camuera et al., 2019; 2022, among others). Some of them also discuss about vegetation changes in NE Iberia, close to the study areas, which would help authors contextualize and discuss their interpretations about the animal diets. Similarly, the speleothem records from the north of Iberia would help constrain temperature/precipitation patterns; there are many of these records in the Vasco-Cantabrian area. Other records from the Iberian Peninsula, the Mediterranean coast or the Iberian margin would show the general climatic patterns for this period. Authors should discuss their data (agreement or disagreement) according to these continuous records in order to have a big picture of the paleoclimate and paleoenvironments in the studied period. Otherwise, authors are only comparing the archaeological sites from their two study areas and the work would only be of local interest. In fig. 2 authors included the $\delta^{18}O$ record of Greenland, but they did not discuss their data according to this record. As I mentioned before, they should compare and discuss the Iberian records instead, since there can be some temporal offsets between some of the events of the NGRIP curve and the Iberian/mediterranean records. Taking all this into account, objective 3 (line 96) is not totally achieved. In any case, my main concerns about this work come from the methodological approach.

Dear reviewer,

We highly appreciate all the suggestions and reflections provided for the manuscript, as they will significantly enhance the final publication. We have addressed them point by point as follows.

Regarding the discussion or data in connection with other regional proxies, we will include further discussion involving regional continuous paleoclimatic and paleoenvironmental terrestrial records of Iberia, as recommended (Martrat et al., 2004; 2007; Pérez Mejías et al., 2019; Moreno et al., 2012; González-Sampériz et al., 2020; Camuera et al., 2019; 2022), as well as marine ones (Naughton et al., 2007; Roucoux et al., 2001; Sánchez-Goñi et al., 1999; Sánchez-Goñi et al., 2008). We addressed this in our recent paper (Fernández-García et al., 2023). However, it's worth noting at a regional scale that limited continuous paleoclimatic records can be validated today. Nevertheless, we would like to emphasise that our primary focus is based on reconstructing climate that can be directly or indirectly linked to human presence. This is the main reason archaeological data takes priority in this paper, even though we know that they represent “discontinuous points” within the MIS3 stadial-interstadial fluctuations. We will make this perspective more straightforward in the introduction and incorporate further discussion regarding the regional environmental records.

Chronology: This a very important part of the study and should be presented in a subsection in section 2 or in the methodology (section 3). Please, explain deeply the absolute chronology of the levels where the ungluted remains were collected. Please, specify the dating method: ESR, OSL or 14C in Fig 2. Regarding 14C samples: it is not clear what radiocarbon curve was used to calibrate the 14C samples. Please, re-calibrate all the radiocarbon data with the latest 14C curve from 2020; there are important age shifts after the Holocene comparing the intCal2020 curve and previous calibration curves. Taking into account the confidence intervals of the ages (I suppose 2 sigma for 14C)->Line 98-99: "The chronological resolution in the study areas for this period allows us to correlate regional paleoenvironmental changes with global records": this would be only true for the two sites younger than 30 ka, since the dates of the other sites might overlap stadial and interstadials (and their probability distributions are very large). What are the grey bands in Fig. 2, stadials? Please explain. In any case the chronology of the samples, according to Fig. 2, would only allow to differentiate between stadials/interstadials in the two youngest samples. What is the meaning of the green colour in one date at around 40 ka in Fig. 2 (Canyars)? What is the meaning of the dates (dots and bars) in Fig 2: Do they represent a single dating event, or a sum of distribution of various dates?, please explain this in the caption and in the main text. If it is the sum of distribution, did you use any statistic approach to obtain it (such as the ones that can be obtained from OxCal software?) Do they represent the ages of the whole archeological sites or of the levels where the ungluted teeth were taken? This is very important to specify in the main text since the age of the remains could vary. Therefore, the caption of Fig. 2 should explain these details and there might be a (sub)section in the main text explaining the chronology of the levels where the remains were taken. This is crucial to validate the discussion of the manuscript

We have comprehensively described all sites in Appendix A, opting to conserve space within the main manuscript. However, we acknowledge the reviewer's suggestion to include a synthesis in Section 2 ("Archaeological sites and sampled material"), providing more details on chronology, cultural attributions, and dating methods. It is important to note that our focus does not directly address specific dates in this work; hence, they are not discussed in the methods section. Date calibrations were done using the OxCal software with the IntCal2020 curve, as outlined in previous publications by our team (e.g., Vidal-Cordasco et al., 2022; Fernández-García et al., 2023). Dates provided represent an approximate age for each level, incorporating either multiple dates or a single date. Further details on dating methods and selected dates for each level are in Appendix B ("B1-Dates").

Additionally, detailed information on date calibration have been included in Appendix B (B1-Dates): "Single radiocarbon dates for sites were calibrated using OxCal4.4 software (Ramsey, 2009), considering the INTCAL20 calibration curve (Reimer et al., 2020). For sites with multiple dates, Bayesian chronology modeling was performed using OxCal4.4. Dates were organised into a 'Sequence,' and chronological information for each level was grouped into a single 'Phase' with start and end 'boundaries' to bracket each archaeological level."

Considering the error chronological margins and the limitation into a straightforward correlation with a single climatic stadial (GS) or interstadial (GI) we have modified the sentence "The chronological resolution in the study areas for this period allows us to correlate regional paleoenvironmental changes with global records".

Concerning formal corrections for Figure 2, we have specified that grey bands indicate Greenland Stadials (GS), and, as previously addressed with reviewer 1, the green colour denotes the Mediterranean area. Furthermore, it will be emphasized that detailed chronological information shown in this figure is available in Appendix B for each archaeological level. In the figure, dots represent the median value of the radiocarbon dates of each level that can be calibrated with the IntCal 20 by using the OxCal version 4.4, while bars indicate the minimum and maximum values of the calibrated dates per level, considering in most of the cases the 95% interval of the calibrated dated. In those old dates, where calibration was not possible due to the high standard deviation, such as Axlor levels III, IV and

VI and El Castillo level 21, the dot represents the date uncalibrated and the bars represent the maximum and minimum of the standard deviation. This clarification was included in the figure caption.

Methodology.

1. Authors mentioned that they did not carry out any pre-treatment to remove secondary carbonates, but did authors check the potential presence of secondary carbonates or the preservation of carbonates? This is very important since all the results are based on these values (there is no data of d18O in phosphates). The physical cleaning that was carried out would not remove all the potential secondary carbonates. Secondary carbonates are very common in archeological contexts such as karstic caves (like the ones studied here), and would modify the isotopic composition of carbonates if they are not eliminated. This must be double checked before stating the sample preparation. Authors did not mention methods to double checked that the isotopic signal was the pristine one. They only mentioned in line 503-504: “The carbonate content in our samples, ranging from 3.9% to 8.9%, is similar to the proportion found in modern tooth enamel, suggesting no immediate indication of diagenetic alteration”. However, there is no explanation about the methodology used to calculate this percentage of carbonate. Authors should explain this in the methodology section and add the % of carbonate in a table (e.g. Table 2).

Based on our experience, we have taken into account that the potential side effects resulting from pretreatment and the removal of carbonates can be more problematic than the risks associated with the presence of secondary carbonates. Calcium carbonate content, as briefly stated in the original manuscript, does not indicate a substantial presence of secondary carbonates, and we used this as a diagenetic test in line with recently published recommendations (Chesson et al. 2021). Additionally, phosphate data has previously been generated for a subset of samples at the Axlór site; these data show d18Ocarb-d18Ophos offsets within the expected range for all samples (Pederzani et al. 2023). We, therefore, consider the risk posed by secondary carbonates to be low. And while additional analyses such as FTIR may give more clarity on this, we believe it essential to acknowledge that no pretreatment method can guarantee their complete removal and that all removal methods, in turn, have the potential to introduce significant undesirable side effects, primarily on d18O (Pellegrini and Snoeck, 2016; Snoeck and Pellegrini, 2015). In addition, phosphate analyses were already in progress during the submission of this paper to double-check this issue and the results will be discussed in future works.

We recognised that secondary carbonates and diagenesis tests, to inform the choice of pretreatment methods, were not discussed as clearly as they could have been. In the methods section we will add a more explicit discussion of these points to justify better the methodological approach taken in the study. The carbonate content (%) was already included in Appendix B (B3. Samples raw), but the mean value will be included in Table 2 in the manuscript, as suggested. In agreement with the explanation of Iso-Analytical laboratory, “the equivalent calcium carbonate content values (%) were derived by comparing the total ion beam data for the samples against the pure calcium carbonate references”. We have also included this explanation in Appendix B (B3-Samples-Raw).

2. Regarding the potential treatment to remove the organic matter, authors said: Lines 145-151: “For this reason, in this work, most of the samples were not pretreated, except for the equid samples from Labeko Koba and Aitzbitarte III, and the cervids and equids from El Otero that were sampled and pretreated in the context of the initial project. Pretreatment followed was established by Balasse et al. (2002), where around 7 mg of powdered enamel was prepared and pretreated with 3% of sodium hypochlorite (NaOCl) at room temperature for 24 h (0.1 ml/mg sample), and thoroughly rinsed with deionised water, before a reaction with 0.1M acetic acid for 4 h (0.1 ml/mg sample) (equivalent protocol in Jones et al., 2019).” And afterwards in lines 453-456: “In the case of equid samples from the Vasco-Cantabrian region, it should be considered that they have been pretreated with a combination of NaClO

and acetic acid, which could potentially affect the isotopic values. Samples after organic removal pretreatment can potentially show either higher or lower $\delta^{13}\text{C}$ values and higher $\delta^{18}\text{O}$ values based on previous experiments (Pellegrini and Snoeck, 2016; Snoeck and Pellegrini, 2015)". So, my doubt is: why did authors treat whole batches of samples to evaluate these "side effects" instead of applying both protocols (with pretreatment and without pretreatment) to aliquots of the same samples? Although they finally ended up that "the influence of the pre-treatment on our samples is deemed to be limited.", this was a risk, and now they cannot be 100% sure about this potential influence. Was there any reason to measure the samples where the organic matter was removed in a different IRMS. If yes, please, explain.

The samples presented in this work have been collected over extensive research conducted by the EvoAdapta group. In 2013 we established a collaboration with the University of Cambridge, where pretreatment and IRMS measurements on the horse teeth from Cantabrian sites were achieved. At that time, pretreatment methods were not as methodologically questioned as today.

Some bovine tooth enamel powder samples were also collected in the respective museums during the initial research phase. However, the availability of these bovine teeth leftovers was limited then. Within the scope of the new ERC SUBSILIENCE project, given the new body of knowledge, it was determined that pretreatment initially done at Cambridge might significantly compromise the final results and thus, we decided not to pretreat the samples.

3. Line 182-196 and throughout the calculations and the manuscript: Authors referred to E^* as the fractionation factor. The symbol E is traditionally the enrichment factor, not the fractionation factor (α). $E = (\alpha - 1) \times 1000\text{‰}$. So, what factor have authors applied eventually: fractionation or enrichment factors? Are these factors mixed in the text and in the calculations? This has to be clear, and if the factors are wrongly applied (enrichment instead of fractionation factor), correct the calculations. Although both factor are related ($E = (\alpha - 1) \times 1000\text{‰}$.) the obtained results would differ, and thus, the derived potential interpretations.

This confusion is strictly terminological between fractionation and enrichment factors. As the reviewer suggests and in agreement with the specified references (Merceron et al., 2021; Passey et al., 2005; Tejada-Lara et al., 2018), " E^* " should refer to isotope enrichment or enrichment factor, hence not causing any issue in interpretation or calculations. This term will has been rectified throughout the manuscript.

4. Lecuyer et al. (2021) performed the calculations to correct the effect of atmospheric CO_2 (difference of 1‰ and CO_2 concentration) for the LGM; so, these specific CO_2 corrections can only be applied for the LGM, but in the present manuscript there are no samples for the LGM (23-19 ka). In addition, the correction for the isotopic composition of atmospheric CO_2 should be done specifically for each age of the studied samples, instead of using a general average -7‰: as authors mentioned, a variation of a ca. 1‰ would imply a change in 150mm of precipitation. Check for example the isotopic composition of atmospheric CO_2 reconstructed from ice records for the Late Pleistocene that ranges from -7 to -6.5 ‰ (Eggleston et al., 2016: Paleoceanography, among others). So, please, apply age-specific CO_2 corrections. I mean, when you are quantifying climatic variables, you should reduce the potential error sources. These errors increase by applying a general unspecific correction for all the data (the same affirmation could be applied to all corrections of the isotopic data in the manuscript).

We agree with the reviewer that atmospheric CO_2 levels varied throughout the Late Pleistocene, and ideally, adjustments should be more specific for each age. In our study, the primary aim of this correction is to mitigate the bias introduced by modern atmospheric CO_2 levels in the correlation. We are not

entirely certain that applying the same average correction to all samples implies a higher risk than using a specific value based on chronology. This is because inaccuracies in these calculations could potentially introduce more unpredictable effects into the final interpretation, especially when dealing with imprecise chronological information. In Table 1 from Leunberger et al. (1992) or in Eggleston et al. (2016) it can be appreciated that fluctuations are less than 0.5‰. However, as the reviewer argues, 0.5‰ variations in CO₂ levels will suppose less than -100mm.

We should assume, as Lécuyer et al. (2021) expressed, “that only broad estimates may be obtained” or “our estimates of MAP from d13C-data must be seen as a first approximation that needs to be confronted with estimates from other approaches”. Uncertainties associated with the estimated MAP values are on the order of ±100-200 mm, and this associated error will be included.

In any case, our intention is not to propose an absolute MAP value, but to obtain comparative values on an understandable scale for assessing variations along a sequence or between deposits. It is important to acknowledge that uncertainties, as suggested by the reviewer, must be considered in the final value. We will introduce these appreciations in the text and the associated error.

5. Authors proposed the above-mentioned (oversimplified) correction for the change in the isotopic composition atmospheric CO₂, but they did not apply any correction related to the change of the isotopic composition of the sea-water during the Late Pleistocene, which is the main moisture source for rainfall (I would not mention the moisture and precipitation due to inland evaporation/recycling in the Iberian Peninsula, even during cold periods (Krklec and Dominguez-Villar, 2014), in order to simplify this interpretation). The global isotopic composition of the rain during colder/warmer periods (glacial/interglacial, stadial/interstadial) differs, not only due to the isotopic fractionation caused by temperatures, but also due to the accumulation/release of the lighter water isotopes in the ice sheets/glaciers during cold/warmer periods (Dansgaard, 1964), among others factors affecting the global isotopic composition of sea waters. Therefore, in order to obtain a reliable isotopic data related to precipitation, the obtained dO18 values has to be corrected to remove this effect. See for example, Niedermeyer et al. (2010) or Garcia-Alix et al. (2021) approach to correct past hydrogen isotopes from vegetation, or even Fernández-García et al (2020) for fossil mammals in the studied period of this paper.

We are aware of this possible bias derived from variations in d18O, mainly between glacial and interglacial periods. However, as previously stated, considering the high chronological uncertainty within the dates, we considered it riskier to correct the results in one or another direction randomly. Nevertheless, as this can be a potential source of uncertainty, this explanation will be incorporated in the manuscript at the end of the section “3.4 Oxygen stable isotope compositions as environmental tracers”. From our point of view, ideally, maintaining a consistent value for both corrections would be preferable to address the significant modern bias from the Holocene period. Therefore, we concur with the reviewer for a correction. However, considering the chronological uncertainties, we find it more consistent to apply a general correction specifically for the MIS3 period of -0.6‰ from the obtained d18Omw, as previously proposed by Fernández-García et al. (2019). This has affected tables 3 and 4, as well as figures 4 and 5.

6. Where did these obligate drinkers drink (water source)? Directly from the rain? Ponds? Lakes? Unless they directly drink precipitation waters (oversimplification), this would imply more isotopic fractionation and would also mask the temperature signal. This is even more important in the studied glacial period, and especially in the stadials? Apart from the potential enhanced rain evaporation due to low atmospheric moisture in glacial times (Dansgaard 1964), an increasing evaporation in lakes, ponds (and even in vegetation->enhanced evapotranspiration) during dry periods have been

demonstrated by different isotopic studies in carbonates from freshwater gastropods, bivalves, or ostracods and even from leaf wax isotopes of freshwater and terrestrial plants. Please clarify this issue, and explain this constrain in the methodology since it would affect the reconstructed temperatures.

Animals will likely acquire water from various types of water sources, with some sources, such as small streams, better reflecting seasonal isotopic oscillations than others, like large lakes or evaporating ponds. Therefore, we agree with the reviewer that certain types of drinking behaviours can impact $\delta^{18}\text{O}$ of the animals, and we have considered these concerns in our interpretation. We will provide a more detailed explanation in the methods and discussion sections, particularly about the state-of-the-art insights outlined by Pederzani and Britton (2019).

In our study, we argue that, in animal teeth where the obtained profile is sinusoidal and has a sufficiently large range ($<1.5-2$), predominant seasonal water information is recorded, related to the predominant consumption of water sources that reflects seasonal precipitation reasonably well. However, it is evident that systematic consumption of certain highly buffered water sources can significantly attenuate the final signal recorded. We will discuss this possibility in more depth in the paper.

Regarding enhanced evaporation during more arid stadials, we consider this effect relatively small regarding oxygen isotope data from large bovids sampled in this study. These animals heavily rely on consuming large amounts of liquid drinking water, and the relative contribution of water from plants, which, as the reviewer mentions, is most susceptible to isotopic enrichment through evapotranspiration, is negligible (Hoppe, 2006; Maloiy, 1973). Other water sources strongly affected by evaporative enrichment are large rivers and lakes with long water residence times where a systematic consumption of such waters would lead to an absence of a seasonal sinusoidal $\delta^{18}\text{O}$ signal. However, this is not systematically observed in conjunction with the higher $\delta^{18}\text{O}$ that would be expected in such evaporated water bodies.

Finally, if aridity effects significantly drove $\delta^{18}\text{O}$, we would expect a positive correlation between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$, which also responds to aridity effects, but we do not see such correlation in our data. However, we agree that non-temperature effects on $\delta^{18}\text{O}$ are critical, so we will add the points raised here to the paper's discussion.

7. Precipitation source (North Atlantic Oscillation modes and Mediterranean dynamics). In the Iberian Peninsula, the isotopic composition of precipitation is highly affected by the moisture source in the present - and in the past - (Araguás y Díaz Teijeiro, 2005, Celle-Jeanton et al., 2001; Domínguez-Villar et al., 2013; Krklec and Domínguez-Villar, 2014; Moreno et al., 2010; 2012; 2014; 2021, Toney et al., 2020; García-Alix et al., 2021; Schirrmacher et al., 2020, among others). Therefore, temperature-isotope fractionation equations would not work that well to reconstruct past temperatures. When proper analysis of the isotopic signal of precipitation are performed in N Iberia, there are sampling stations (precipitation, temperatures, isotopes, and moisture sources) where the isotopes from precipitation do not correlate well with temperatures (Moreno et al., 2021). This thoroughly study of the isotopic composition of precipitation in northern Spain ended up with "although important, air temperature only explains part of the observed $\delta^{18}\text{O}_p$ variability and is therefore not the only control." This issue is even more important in coastal areas, as the ones studied in this paper, and even more in the Mediterranean coast (at present there is some influence of amount effect in the Mediterranean areas of Catalonia; Moreno et al., 2021). Therefore, in the best oversimplified case-scenario (not considering previous comments 5 and 6, and admitting large errors due to these potential source and amount effects) if we would want to calculate temperatures from the isotopic values of precipitation we would need an equation for the Vasco-Cantabrian region, and another one for NE-Mediterranean Iberia (different precipitation pattern and forcing). And even more, since atmospheric patterns, and therefore, moisture sources, are not the same during the warm and cold seasons (see the above-mentioned studies),

specific equations for cold and warm seasons should be applied to reconstruct “summer” and “winter” temperatures.

We appreciate the reviewer's concern about differences in the slope of the d18Oprecip-temperature relationship, which is known to vary geographically, seasonally and through time due to differences in moisture source and atmospheric circulation patterns (Dansgaard, 1964; Gat, 1980; Rozanski et al., 1993).

There are essentially two approaches to account for these differences. The first approach, as described by the reviewer, attempts to specify as accurately as possible the particular d18Oprecip-temperature relationship for the study area. This approach usually yields more precise estimates. The second approach instead posits that creating a precisely accurate d18Oprecip-temperature equation for a specific location is extremely difficult as data is not directly available for the areas in question, but often for rather distant measurement stations and because we are additionally projecting into the past where atmospheric circulation patterns were not identical to what they are now.

In this second approach, the inherent variability and uncertainty of the d18Oprecip-temperature relationship is instead represented statistically, and the resulting uncertainty is incorporated into the error range of the temperature estimate by including a large number of data from different measurement stations into the d18Oprecip-temperature calibration data set. This approach yields less precise data but reduces the risk of systematically inaccurate results. In this study, we consider that due to the projection of equations into the past, the second approach is preferable. Still, we acknowledge that the decision between different methods remains debated in the field, with some researchers choosing the closest measurement station, a mixture of stations from one country, or even globally combined data sets (see e.g. Skrzypek et al. 2011; Pryor et al. 2014; Tütken et al. 2007; Lécuyer et al. 2021). We will justify this decision in more detail concerning the literature cited here in the methods section of our paper.

However, in response to the reviewer's suggestion regarding the use of specific equations for cold and warm seasons to reconstruct "summer" and "winter" temperatures, we have carefully considered this recommendation after online reviews. We agree that this approach could significantly enhance the resolution of our results. Consequently, we have opted to apply the linear regression models proposed by Pederzani et al. (2021), which differentiate between mean, summer, and winter temperatures. These correlations are established using monthly climatic records from Western, Southern, and Central Europe stations within the Global Network of Isotopes in Precipitation (IAEA/WMO, 2020). Further details are provided in the Methods section.

8. “MAT was calculated from the d18O mean value between summer and winter in each tooth before modeling to reduce associated error”; However in caption Table 4: “For some profiles with an unclear seasonal shape, MATs were deduced from the original average of teeth without a seasonal profile”. So, what is the correct methodology?, In any case, according to the methodology section, MAT was calculated before modeling to reduce associated error, but summer and winter temperatures after the inverse modelling?. This reasoning is not clear to me: Is there no associated error in the transformation for summer and winter temperatures? This is why there are some odd values, for example, sample AXL60 MAT 12°C, ST 20.4°C and WT 10.8°C (only 1.2°C colder than the MAT).

We appreciate that the reason for treating MAT differently from summer and winter temperature estimation was not as clearly explained as it could have been. Due to the influence of the prolonged enamel mineralisation process, the seasonal amplitude of d18O derived from sequentially sampled teeth necessarily has a lower measured amplitude than the real d18O seasonal variation in the drinking water consumed by the animal. This effect is corrected as much as possible by the inverse modelling procedure. Therefore, it is imperative that summer and winter temperature estimations can be made on the output values of the inverse model rather than directly on measure d18O data, to avoid systematic

underestimation of the seasonal temperature change. However, the same is not necessarily true for mean annual values, as the inverse model will essentially symmetrically expand the d18O seasonal curve, introducing no substantial changes to the data annual means. A systematic comparison of d18O annual means derived from unmodeled vs modelled curves in Pederzani et al. 2021 showed that annual means do not change between these different ways of obtaining d18O annual means for input into a temperature estimation. In this case, it is preferable to derive mean annual temperature estimation directly from d18O enamel annual means, because the inverse model introduces significant uncertainty, which would also impact any derived temperature estimation. We therefore consider the approach of using unmodeled annual means, but modeled summer and winter d18O as the input for temperature estimation to be the more robust method. We have clarified this difference in the text of the paper with reference to Pederzani et al. 2021 where the two approaches are compared in detail.

Regarding the use of different averaging methods for obtaining d18O annual averages, an in-depth comparison in Pederzani et al. 2021 also showed that the results of either using a summer-winter two-point average or an average of all data points across an entire year yield indistinguishable results. Using a two-point average commonly has the advantage that it allows the calculation of an annual mean for teeth that do not record complete annual cycles but do record both a summer peak and a winter trough in d18O. This method, therefore, allows us to maximize the number of data points that can be obtained for MAT estimation. Still, it is not possible to apply this method to specimens that do not show a clear sinusoidal d18O signal, so we chose to average all data points for those specimens instead. This last option, however, may be less reliable, and we have reviewed the text to clarify these uncertainties.

9. The reconstructed meteoric waters are different depending on the species, even in the same level, and therefore, reconstructed temperatures also differ. I'm aware of the different ecological behaviors of the different species, but the MAT should be, at least close. There are also some discrepancies in specimens of the same species in the same archaeological levels.

The interspecies variability in temperature reconstructions is not significantly higher than the intraspecies variability, as demonstrated in Table 4, where estimations are within error for all species. Multiple factors could account for differences for both within the same species and between species. Variability in d18O, and consequently in derived temperature estimations, is expected even within the same species or among individuals from the same population. This variability is influenced not only by minor differences in foraging and drinking behavior but also by slight metabolic and physiological variations. The relationship between d18O of drunked water and d18O of body water and enamel is affected by factors such as body size, metabolic rate, breathing rate, moisture content of food, and feces, among others (Kohn et al. 1996; Magozzi et al. 2019). It is important to note that appreciable variations in d18O have been observed even within small groups of animals living in the same constrained habitat (e.g., Magozzi et al. 2020; Hoppe et al. 2004). Additionally, it should be considered that archaeological levels represent palimpsests of occupations from different periods, which further increases the expected inter-individual variability observed in d18O. To provide more context for interpreting the slight differences observed among other species and individuals, we will include a note in the manuscript to convey the sentiments expressed in this response.

10. The general comparison between the isotopic composition of the faunas of the archaeological sites of both areas is not objective since in the Vasco-Cantabrian area there are remains from 80 ka to 18 ka approx. (14 sections from 5 archaeological sites), but in the Catalanian area there is only one site at around 40 ka (it could have been coeval to the HS4, a period especially cold and dry). Thus, the general comparisons that appeared in some parts of the manuscript between two areas (not the ones specific at ca. 40 ka) are not balanced.

We agree with the reviewer that there is a disbalance between both northern Iberian regions. The lack of data in the Mediterranean region produces this difference. So far, it is the only site with this type of carbonate enamel analysis in the contemporaneous archaeological sites of the Mediterranean. Nevertheless, we considered it interesting to include the Canyars results as it is a well-dated archaeological site in a precise climatic event such as the Heinrich 4. For this reason, we make this evaluation to contrast with the wide-studied Cantabrian region.

11. What is the error associated with the different equations Eq. 1-9? Authors have to take into account the different errors (not only the standard deviations) that are being accumulated in each equation and also plot and mention them in the tables and in the text. For example, a MAT of 15°C +/- 0.5 °C would be accurate, but if the error was +/- 4°C the interpretation would be more open. This is even more important in this work since the oxygen isotopes were measured in carbonates, not in phosphates, and two more equations were used to convert the $\delta^{18}\text{O}$ data of the carbonates to $\delta^{18}\text{O}$ of phosphates. Authors said: lines 277-278: "In these estimations, the associated error from converting $\delta^{18}\text{O}_{\text{phos}}$ to MAT is enlarged by the uncertainty derived from the transformation of $\delta^{18}\text{O}_{\text{carb}}$ (VPDB) to $\delta^{18}\text{O}_{\text{phos}}$ (VSMOW)". Therefore, all the errors should be calculated (for both O and C), included in the plots, tables and when describing the results.

The authors fully agree with the reviewer and this issue will be solved. Associated errors will be calculated following the methodology outlined by Pryor et al. (2014), as done by one of us in Pederzani et al. (2021). We have added this information to the manuscript with a short explanation of the error calculation method, which integrates uncertainties from the consecutive conversion steps involved in the temperature estimation from $\delta^{18}\text{O}_{\text{phos}}$. Concerning the uncertainties arising from the conversion of $\delta^{18}\text{O}_{\text{carb}}$ (VPDB) to $\delta^{18}\text{O}_{\text{phos}}$ (VSMOW), similar considerations have been addressed in previous studies (Pryor et al., 2014; Pederzani et al., 2023). These works have concluded that the impact of this conversion is negligible compared to the error propagation in subsequent calibrations used for temperature estimations from $\delta^{18}\text{O}_{\text{phos}}$. The spreadsheet for error calculation from Pryor et al. (2014) adapted to the correlation on this study can be accessed at GitHub (https://github.com/ERC-Subsiliencia/Ungulate_enamel-carbonate). The associated errors have been placed in Appendix B ("B6-Temperature-estimations"), explained in methods section and mention in Table 4 caption.

In any case, we want to emphasize once again that the main objective of this article was not the reconstruction of temperatures. We are aware that it would be more appropriate to conduct this based on analysis through phosphate fractions, which is currently underway. We consider this climatic approach valuable for its comparative value, but we remain cautious in the interpretation of absolute values.

12. Inverse Modelling. I think that the correct reference for the inverse modeling (main text and appendix D) is Passey et al. (2005) but this one "Passey, B.H., Cerling, T.E., Schuster, G.T., Robinson, T.F., Roeder, B.L., Krueger, S.K., 2005. Inverse methods for estimating primary input signals from time-averaged isotope profiles. Geochim. Cosmochim. Acta 69, 4101–4116." Reviewing this paper, I noted that the reconstructed profiles showed mostly the same trends/changes as the original isotopic data. However, in the case of the reviewed manuscript sometimes these reconstructed profiles exhibited opposite patterns/trends from the original isotopic values. I'm not familiar with these kinds of transformations, but: is this common? This is an extra transformation for the data that would be used to calculate absolute climatic parameters, and thus, and extra potential error source. Could authors double check these calculations?. In addition, in the original paper from Passey et al. (2005), this inverse modeling was also applied to the carbon isotopes. Why authors did not apply this correction also to the carbon isotopes of the sequential sampling?

Firstly, the reviewer is correct about the reference; it was mistakenly listed in the final reference section. It will be edited in the manuscript. Secondly, some of the teeth included in the manuscript, as elaborated within the methods and also discussed in the section “5.2 Seasonality, mobility, and water acquisition: oxygen ratios and intratooth profiles,” exhibit flat and less sinusoidal profiles, which are less suitable for the application of this model as it presupposes an approximately sinusoidal shape of the seasonal d18O curve. This may result in non-straightforward interpretation in the model outcomes for non-sinusoidal curves. Additionally, small but abrupt ‘noise’ spikes in the original d18O curves tend to be amplified in the model output. They can sometimes represent the highest or lowest values globally in the modelled d18O curves. For this reason, we consistently selected peaks and troughs as summer and winter modelled d18O based on where these seasonal regions are located in the original profile. It should be noted that the model reproduces the lag between d18O change in the input to the animal compared to the expression in tooth enamel. Therefore, the modelled d18O curve of the model is constantly shifted in the x-direction relative to the d18Oenamel input time series. This can sometimes lead to seemingly opposing isotopic changes being located in the same x-axis region, but this does not represent an actual reversal of the isotopic trends between the model and the original. Regarding the uncertainty associated with these models, this is represented as a 95% confidence interval, which is presented in each figure in Appendix D as shaded grey areas, and they can also be found in Appendix B (“B6-Model extrema”). Additionally, as previously discussed concerning error calculation, the accumulated errors for winter and summer will be expressed in Table 4.

Finally, regarding the question about applying the model to carbon values, we considered this to add little valuable insight in our particular case as the d13C time series of our specimens do not show appreciable seasonal change and do not conform to an approximately sinusoidal shape. This is due to the absence of a marked seasonal change in diet, or at least one that cannot be detected through carbon isotopic values, which is common in Eurasian individuals consuming C3 plants. In the absence of such seasonal d13C change, modelling of amplitude damping using the Passey et al. (2005) model would not be genuinely appropriate and would likely lead to issues with model convergence, so we did not attempt this. We will add a note to the manuscript to justify this decision (Appendix D).

Minor comments:

The first paragraph in the introduction and the first lines in the abstract deal with the importance of these kinds of studies to understand the human evolution in this region, but eventually, this is not discussed in the manuscript according to the obtained data.

When we referred to understand human evolution, we meant to reconstruct the climatic and environmental conditions that late Neanderthals and early modern human groups had to face in the regions of study during the Middle to Upper Paleolithic. We aimed with this study to propose the local conditions to later evaluate their adaptation strategies and resilience. We will clarify this aspect.

First sentences of the introduction (lines 38-43): please add references. There are interesting papers dealing with this issue in the Iberian Peninsula and in Europe: Neanderthal-AMH-climate change.

Relevant works such as D’Errico and Sánchez Goñi (2003), Finlayson and Carrión (2007), Sepulchre et al. (2007), and Staubwasser et al. (2018) will be cited but they will be updated with recent reviews such as Sánchez Goñi (2020), Timmerman (2020), Klein et al. (2023) or Vidal-Cordasco et al. (2022; 2023).

Lines 49-50. It is ok, but authors are not including some information (and significant climate-related references) in the area dealing with continuous paleoclimatic records, and they only focused on the data obtained from archeological sites (whose paleoenvironmental record is not that continuous); so, line 75 is not summarizing the multiproxy studies in this area.

The regional non-archaeological studies mentioned by the reviewer, as well as references to other non-archaeological regional records, will be incorporated into the introduction, as previously discussed.

Lines 80-87: add references

The references of Passey et al. (2002), Kohn et al. (2004), and Zazzo et al. (2005) will be included to support the discussion regarding isotopic signal preservation and enamel mineralization. On the other hand, references Pederzani & Britton (2019), Ambrose and Norr (1993), Luz et al. (1984), and Pryor et al. (2014) will be used as examples to justify the general interpretation of carbon and oxygen signals.

Regarding the fossil sites that authors call mediterranean". Since the rest of the fossil sites are in "northeastern Iberia", the term "Mediterranean area" is very open and do not specifically identify the studied site, I would say NE Iberia?

The authors have made the adjustment, replacing the term "Mediterranean area" with "northeastern Iberia" as suggested.

There are some issues with the chronology/dates. For example, line 21 abstract: 80 to 15,000 cal BP. Do you mean 80 ka or ky, right? Taking into account the study period, and the accuracy of the dates, I would not use "yr", I would use ky or ka. In addition, authors should round the dates in the text to the nearest hundreds (eg, line 601: 41,136 to 38,570 cal yr BP: 41.1 ka to 38.6 ka). This accuracy does not make sense in the studied period due to the uncertainty of the measurements.

To prevent any potential confusion, as recommended by the reviewer, all dates throughout the paper have been revised for consistency and expressed in thousands of years ago (ka BP and ka cal BP).

Results: please, add the ages (in ka and the different technocomplexes) to the subsection headings of the different archaeological sites, otherwise one has to check Fig. 2 for each site.

Ages have been incorporated into each site subsection to establish a general chronological framework and facilitate reader comprehension: Axlor (ca. 80-50 cal ka BP), El Castillo (ca. 75-47 cal ka BP), Labeko Koba (ca. 42-37 ka cal BP), Canyars (ca. 40 ka cal BP), Aitzbitarte III interior (ca. 27 ka cal BP), and, El Otero (ca. 19 ka cal BP). Nevertheless, these time periods are approximate and serve as a general reference and it is advisable for readers to refer to Figure 2 for detailed information on each level.

I understand the structure of the result description, but the mixed description of the isotopes from different levels of the same archaeological sites, which sometimes have 10 ka of difference between them, is rare to me.

We believe that this is the most efficient way to present the results, as it takes into consideration the specific characteristics of each individual site, which at times can be influenced by baseline isotopic values.

Line 325: MATAs=1-8/-2.1°C? do you mean 1.8?

Yes, we mean 1.8°C. I has been corrected.

Eq 10: P value <?

Yes, it was a mistake. We have corrected it to "<".

All figures with the data plotted according to the chronology. El Castillo 21A appears after Axlor III; but according to the chronology this site is previous (Fig. 2). The same would happen between some levels of Labeko Kova and Canyars-I. Is this correct? In this case, arrange the sites with the real chronological order even though they belong to different archaeological sections.

We understand that this assessment primarily pertains to Figure 5, where our priority was to arrange the sequences in accordance with an internal diachronic order. While we acknowledge the importance of chronological coherence, for a better understanding, initially, we thought to maintain the separation by sites. However, after reviewing online feedback, we have reconsidered the suggested modification and agree with the reviewer's perspective on organizing levels diachronically. Consequently, we have rearranged the levels in Figure 5 in chronological order. (*Note to editor: This review has changed from the initial online response.)

When speaking about range of values (temperatures, isotopes, precipitation), sometimes the lowest values are mentioned before and other times the other way around. Be consistent and cite the lowest values first (Eg, line 404 MATAs).

Thanks for the comments. We have review it, to be consistent in the final version.

Table 3: I suppose that the different group of data (rows) are related to the three groups of specimens. Right? Please, specify the taxon groups in the tables.

As the reviewer has pointed out, there is an error, and it is necessary to designate each taxon in the table, from top to bottom, including total, bovines, and equids.

I do not understand this sentence. Line 510-511: "Based on these arguments, it is suggested that the non-sinusoidal $\delta^{18}O$ signal observed in some individuals is likely attributed to the preservation of the original isotopic signature from water input." This sentence would suggest a bad preservation of the original water composition when a sinusoidal pattern is present?. However, authors explained afterwards some reasons related to the ecology of the specimens. Please, clarify this.

We argue that these non-sinusoidal profiles do not indicate poor preservation in the studied specimens. Instead, we propose that they may be linked to somehow ethological factors of the individuals, capturing an isotopic signature originating from non-seasonal water sources. Further elaboration on this point has been provided in response to previous questions 6 and 9.

Section 5.4. Regarding the $d_{18}O$ of meteoric waters. Authors compare (as a w

hole) their reconstructed $d_{18}O_{mw}$ with the current values in the area and they ended up with similar ranges. Keep in mind that the reconstructed values correspond to the last 80 ka under glacial conditions (different temperatures, precipitation amount, and in some cases, moisture source)

Our comparisons usually concentrate on assessing the range of values without drawing broader inferences. We agree with this consideration, and as discussed in our previous response to the reviewer, we will incorporate these points into the text.

Sometimes the different $d_{13}C$ and $d_{18}O$ have subscripts with the meaning of the isotopic values (eg, $\delta_{13}C_{carb}$, $\delta_{18}O_{carb}$), but another times this is not indicated. This is confusing. Please, add always the subscripts explaining the meaning of the isotopic values.

We have correct the text to ensure that " $d_{13}C$ " and " $d_{18}O$ " always include subscripts to maintain consistency. However, in a few cases we have retained " $d_{13}C$ " and " $d_{18}O$ " because we felt it was not necessary to specify further because they referred to general issues.

I miss a figure summarizing the chronological evolution of reconstructed temperatures and precipitation (with the associated accumulated errors) and comparing them with other continuous records of precipitation or temperatures or vegetation from the Iberian Peninsula, Mediterranean coast or Iberian margin.

We agree that such a figure would be valuable for this study. However, we are presently in the process of refining and expanding the dataset for a more comprehensive regional comparison with phosphate analysis. As suggested in the paper and, as previously noted by the reviewer, the estimation of paleotemperatures was approached tentatively taking into account the uncertainties arising from starting with the carbonate fraction. For this reason, more reliable correlations between our snapshots and continuous regional records will be provided in future works.

Title: ecological evolution of what? This is very general: I would say ecological evolution of ungulate fauna....

Considering suggestion of the reviewer, we propose: "Ecological evolution of ungulates in northern Iberia during the Late Pleistocene through isotopic analysis on teeth"

Please, use always the symbol delta instead of d, there are some "ds" throughout the manuscript.

Thanks for the edits. We have identified this error occurring on three occasions and it has been addressed.

Cited Literature:

Araguas-Araguas, L.J., Diaz Teijeiro, M.F., 2005. Isotope composition of precipitation and water vapour in the Iberian Peninsula. First results of the Spanish Network of Isotopes in Precipitation, in: *Isotopic Composition of Precipitation in the Mediterranean Basin in Relation to Air Circulation Patterns and Climate*. IAEA-TECDOC-1453, Vienna, pp. 173–190.

Camuera, J., Jiménez-Moreno, G., Ramos-Román, M.J., García-Alix, A., Toney, J.L., Anderson, R.S., Jiménez-Espejo, F., Bright, J., Webster, C., Yanes, Y., Carrión, J.S., 2019. Vegetation and climate changes during the last two glacial-interglacial cycles in the western Mediterranean: A new long pollen record from Padul (southern Iberian Peninsula). *Quat Sci Rev* 205, 86–105.

Camuera, J., Ramos-Román, M.J., Jiménez-Moreno, G., García-Alix, A., Ilvonen, L., Ruha, L., Gil-Romera, G., González-Sampériz, P., Seppä, H., 2022. Past 200 kyr hydroclimate variability in the western Mediterranean and its connection to the African Humid Periods. *Sci Rep* 12, 9050.

Celle-Jeanton, H., Travi, Y., Blavoux, B., 2001. Isotopic typology of the precipitation in the Western Mediterranean Region at three different time scales. *Geophys Res Lett* 28, 1215–1218.

Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus* 16, 436–468.

Domínguez-Villar, D., Carrasco, R.M., Pedraza, J., Cheng, H., Edwards, R.L., Willenbring, J.K., 2013. Early maximum extent of paleoglaciers from Mediterranean mountains during the last glaciation. *Sci Rep* 3, 2034.

Eggleston, S., Schmitt, J., Bereiter, B., Schneider, R., Fischer, H., 2016. Evolution of the stable carbon isotope composition of atmospheric CO₂ over the last glacial cycle. *Paleoceanography* 31, 434–452. <https://doi.org/https://doi.org/10.1002/2015PA002874>

Fernández-García, M., López-García, J. M., Royer, A., Lécuyer, C., Allué, E., Burjachs, F., Chacón, M. G., Saladié, P., Vallverdú, J., and Carbonell, E., 2020. Combined palaeoecological methods using small-mammal assemblages to decipher environmental context of a long-term Neanderthal settlement in northeastern Iberia. *Quat. Sci. Rev.*, 228, 106072

García-Alix, A., Camuera, J., Ramos-Román, M.J., Toney, J.L., Sachse, D., Schefuß, E., Jiménez-Moreno, G., Jiménez-Espejo, F.J., López-Avilés, A., Anderson, R.S., Yanes, Y., 2021. Paleohydrological dynamics in the Western Mediterranean during the last glacial cycle. *Glob Planet Change* 202, 103527.

González-Sampériz, P., Gil-Romera, G., García-Prieto, E., Aranbarri, J., Moreno, A., Morellón, M., Sevilla-Callejo, M., Leunda, M., Santos, L., Franco-Múgica, F., Andrade, A., Carrión, J.S., Valero-Garcés, B.L., 2020. Strong continentality and effective moisture drove unforeseen vegetation dynamics since the last interglacial at inland Mediterranean areas: The Villarquemado sequence in NE Iberia. *Quat. Sci. Rev.* 242, 106425.

Krklec, K., Domínguez-Villar, D., 2014. Quantification of the impact of moisture source regions on the oxygen isotope composition of precipitation over Eagle Cave, central Spain. *Geochim Cosmochim Acta* 134, 39–54.

Lécuyer, C., Hillaire-Marcel, C., Burke, A., Julien, M.-A., and Hélie, J.-F., 2021. Temperature and precipitation regime in LGM human refugia of southwestern Europe inferred from $\delta^{13}C$ and $\delta^{18}O$ of large mammal remains. *Quat. Sci. Rev.*, 255, 106796.

Martrat, B., Grimalt, J.O., Lopez-Martinez, C., Cacho, I., Sierro, F.J., Flores, J.A., Zahn, R., Canals, M., Curtis, J.H., Hodell, D.A., 2004. Abrupt Temperature Changes in the Western Mediterranean over the past 250,000 Years. *Science* (1979) 306, 1762–1765.

Martrat, B., Grimalt, J.O., Shackleton, N.J., Abreu Lucia de, Hutterli, M.A., Stocker, T.F., 2007. Four Climate Cycles of Recurring Deep and Surface Water Destabilizations on the Iberian Margin. *Science* (1979) 317, 502–507.

Moreno, A., Stoll, H., Jiménez-Sánchez, M., Cacho, I., Valero-Garcés, B., Ito, E., Edwards, R.L., 2010. A speleothem record of glacial (25–11.6kyr BP) rapid climatic changes from northern Iberian Peninsula. *Global and Planetary Change* 71, 218–231.

Moreno, A., González-Sampériz, P., Morellón, M., Valero-Garcés, B.L., Fletcher, W.J., 2012. Northern Iberian abrupt climate change dynamics during the last glacial cycle: A view from lacustrine sediments. *Quat Sci Rev* 36, 139–153.

Moreno, A., Sancho, C., Bartolomé, M., Oliva-Urcia, B., Delgado-Huertas, A., Estrela, M.J., Corell, D., López-Moreno, J.I., Cacho, I., 2014. Climate controls on rainfall isotopes and their effects on cave drip water and speleothem growth: the case of Molinos cave (Teruel, NE Spain). *Clim Dyn* 43, 221–241.

Moreno, A., Iglesias, M., Azorin-Molina, C., Pérez-Mejías, C., Bartolomé, M., Sancho, C., Stoll, H., Cacho, I., Frigola, J., Osácar, C., Muñoz, A., Delgado-Huertas, A., Bladé, I., Vimeux, F., 2021. Measurement report: Spatial variability of northern Iberian rainfall stable isotope values – investigating atmospheric controls on daily and monthly timescales. *Atmos. Chem. Phys.* 21, 10159–10177.

Niedermeyer, E.M., Schefuß, E., Sessions, A.L., Mulitza, S., Mollenhauer, G., Schulz, M., Wefer, G., 2010. Orbital- and millennial-scale changes in the hydrologic cycle and vegetation in the western African Sahel: insights from individual plant wax δD and $\delta^{13}C$. *Quat Sci Rev* 29, 2996–3005.

Pérez-Mejías, C., Moreno, A., Sancho, C., Martín-García, R., Spötl, C., Cacho, I., Cheng, H., Edwards, R.L., 2019. Orbital-to-millennial scale climate variability during Marine Isotope Stages 5 to 3 in northeast Iberia. *Quat. Sci. Rev.* 224, 105946.

Schirrmacher, J., Andersen, N., Schneider, R.R., Weinelt, M., 2020. Fossil leaf wax hydrogen isotopes reveal variability of Atlantic and Mediterranean climate forcing on the southeast Iberian Peninsula between 6000 to 3000 cal. BP. *PLoS One* 15, e0243662.

Toney, J.L., García-Alix, A., Jiménez-Moreno, G., Anderson, R.S., Moossen, H., Seki, O., 2020. New insights into Holocene hydrology and temperature from lipid biomarkers in western Mediterranean alpine wetlands. *Quat Sci Rev* 240.

References for reviews (different from those indicated by the reviewer):

- Ambrose, S.H., Norr, L., 1993. Experimental Evidence for the Relationship of the Carbon Isotope Ratios of Whole Diet and Dietary Protein to Those of Bone Collagen and Carbonate, in: *Prehistoric Human Bone*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1–37. https://doi.org/10.1007/978-3-662-02894-0_1
- Chesson, L.A., Beasley, M.M., Bartelink, E.J., Jans, M.M.E., Berg, G.E., 2021. Using bone bioapatite yield for quality control in stable isotope analysis applications. *Journal of Archaeological Science: Reports* 35, 102749. <https://doi.org/10.1016/j.jasrep.2020.102749>
- D'Errico, F., Sánchez Gofí, M.F., 2003. Neandertal extinction and the millennial scale climatic variability of OIS 3. *Quaternary Science Reviews* 22, 769–788. [https://doi.org/10.1016/S0277-3791\(03\)00009-X](https://doi.org/10.1016/S0277-3791(03)00009-X)
- Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus XVI*, 436–468.
- Gat, J.R., 1980. The relationship between surface and subsurface waters: water quality aspects in areas of low precipitation. *Hydrological Sciences-Butletin des Sciences Hydrologiques* 25, 257–267.
- Fernández-García, M., Vidal-Cordasco, M., Jones, J.R., Marín-Arroyo, A.B., 2023. Reassessing palaeoenvironmental conditions during the Middle to Upper Palaeolithic transition in the Cantabrian region (Southwestern Europe). *Quaternary Science Reviews* 301, 107928. <https://doi.org/10.1016/j.quascirev.2022.107928>
- Finlayson, C., Carrión, J.S., 2007. Rapid ecological turnover and its impact on Neanderthal and other human populations. *Trends in Ecology and Evolution* 22, 213–222. <https://doi.org/10.1016/j.tree.2007.02.001>
- Hoppe, K.A., Amundson, R., Vavra, M., McClaran, M.P., Anderson, D.L., 2004. Isotopic analysis of tooth enamel carbonate from modern North American feral horses: implications for paleoenvironmental reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology* 203, 299–311. [https://doi.org/10.1016/S0031-0182\(03\)00688-6](https://doi.org/10.1016/S0031-0182(03)00688-6)
- Hoppe, K.A., 2006. Correlation between the oxygen isotope ratio of North American bison teeth and local waters: Implication for paleoclimatic reconstructions. *Earth and Planetary Science Letters* 244, 408–417. <https://doi.org/10.1016/j.epsl.2006.01.062>
- Kohn, M.J., 2004. Comment: tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series. In: Passey, B.H., Cerling, T.E. (Eds.), (2002). *Geochim. Cosmochim. Acta*. vol. 68. pp. 403–405. [https://doi.org/10.1016/S0016-7037\(03\)00443-5](https://doi.org/10.1016/S0016-7037(03)00443-5).
- Kohn, M.J., 1996. Predicting animal $\delta^{18}O$: Accounting for diet and physiological adaptation. *Geochimica et Cosmochimica Acta* 60, 4811–4829. [https://doi.org/10.1016/S0016-7037\(96\)00240-2](https://doi.org/10.1016/S0016-7037(96)00240-2)
- Klein, K., Weniger, G., Ludwig, P., Stepanek, C., Zhang, X., Wegener, C., Shao, Y. 2023. Assessing climatic impact on transition from Neanderthal to anatomically modern human population on Iberian Peninsula: a macroscopic perspective. *Science Bulletin*, 68 (11), 1176–1186.
- Lécuyer, C., Hillaire-Marcel, C., Burke, A., Julien, M.-A., Hélie, J.-F., 2021. Temperature and precipitation regime in LGM human refugia of southwestern Europe inferred from $\delta^{13}C$ and $\delta^{18}O$ of large mammal remains. *Quaternary Science Reviews* 255, 106796. <https://doi.org/10.1016/j.quascirev.2021.106796>
- Luz, B., Kolodny, Y., Horowitz, M., 1984. Fractionation of oxygen isotopes between mammalian. *Geochimica et Cosmochimica Acta* 48, 1689–1693.
- Magozzi, S., Vander Zanden, H.B., Wunder, M.B., Bowen, G.J., 2019. Mechanistic model predicts tissue–environment relationships and trophic shifts in animal hydrogen and oxygen isotope ratios. *Oecologia* 191, 777–789. <https://doi.org/10.1007/s00442-019-04532-8>
- Magozzi, S., Vander Zanden, H.B., Wunder, M.B., Trueman, C.N., Pinney, K., Peers, D., Dennison, P.E., Horns, J.J., Şekercioğlu, Ç.H., Bowen, G.J., 2020. Combining Models of Environment, Behavior, and Physiology to Predict Tissue Hydrogen and Oxygen Isotope Variance Among Individual Terrestrial Animals. *Frontiers in Ecology and Evolution* 8. <https://doi.org/10.3389/fevo.2020.536109>
- Maloiy, G.M.O., 1973. Water metabolism of East African ruminants in arid and semi-arid regions 1. *Zeitschrift für Tierzucht und Zuchtungsbiologie* 90, 219–228. <https://doi.org/10.1111/j.1439-0388.1973.tb01443.x>
- Merceron, G., Berlioz, E., Vonhof, H., Green, D., Garel, M., Tütken, T., 2021. Tooth tales told by dental diet proxies: An alpine community of sympatric ruminants as a model to decipher the ecology of fossil fauna.

- Palaeogeography, Palaeoclimatology, Palaeoecology 562, 110077.
<https://doi.org/10.1016/j.palaeo.2020.110077>
- Naughton, F., Sánchez-Goñi, M.F., Desprat, S., Turon, J.-L., Duprat, J., 2007. Present-day and past (last 25 000 years) marine pollen signal off western Iberia. *Marine micropaleontology* 62, 91–114. <https://doi.org/10.1016/j.marmicro.2006.07.006>
- Passey, B.H., Robinson, T.F., Ayliffe, L.K., Cerling, T.E., Sponheimer, M., Dearing, M.D., Roeder, B.L., Ehleringer, J.R., 2005. Carbon isotope fractionation between diet, breath CO₂, and bioapatite in different mammals. *Journal of Archaeological Science* 32, 1459–1470. <https://doi.org/10.1016/j.jas.2005.03.015>
- Passey, B.H., Cerling, T.E., 2002. Tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series. *Geochim. Cosmochim. Acta* 66, 3225–3234
- Pellegrini, M., Snoeck, C., 2016. Comparing bioapatite carbonate pre-treatments for isotopic measurements: Part 2 — Impact on carbon and oxygen isotope compositions. *Chemical Geology* 420, 88–96. <https://doi.org/10.1016/j.chemgeo.2015.10.038>
- Pederzani, S., Britton, K., 2019. Oxygen isotopes in bioarchaeology: Principles and applications, challenges and opportunities. *Earth-Science Reviews* 188, 77–107. <https://doi.org/10.1016/j.earscirev.2018.11.005>
- Pederzani, S., Britton, K., Aldeias, V., Bourgon, N., Fewlass, H., Lauer, T., McPherron, S.P., Rezek, Z., Sirakov, N., Smith, G.M., Spasov, R., Tran, N.H., Tsanova, T., Hublin, J.J., 2021. Subarctic climate for the earliest *Homo sapiens* in Europe. *Science Advances* 7, 1–11. <https://doi.org/10.1126/sciadv.abi4642>
- Pederzani, S., Britton, K., Jones, J.R., Agudo Pérez, L., Geiling, J.M., Marin-Arroyo, A.B., 2023. Late Pleistocene Neanderthal exploitation of stable and mosaic ecosystems in northern Iberia shown by multi-isotope evidence. *Quaternary Research* 1–25. <https://doi.org/10.1017/qua.2023.32>
- Pryor, A.J.E., Stevens, R.E., Connell, T.C.O., Lister, J.R., 2014. Quantification and propagation of errors when converting vertebrate biomineral oxygen isotope data to temperature for palaeoclimate reconstruction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 412, 99–107. <https://doi.org/10.1016/j.palaeo.2014.07.003>
- Ramsey, C.B., 2009. Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51, 337–360. <https://doi.org/10.1017/S0033822200033865>
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62, 725–757. <https://doi.org/10.1017/RDC.2020.41>
- Roucoux, K.H., Shackleton, N.J., Abreu, L. De, Schönfeld, J., Tzedakis, P.C., 2001. Combined marine proxy and pollen analyses reveal rapid Iberian vegetation response to North Atlantic millennial-scale climate oscillations. *Quaternary Research* 56, 128–132. <https://doi.org/10.1006/qres.2001.2218>
- Rozanski, K., Araguas-araguas, L., Gonfiantini, R., 1993. Isotopic Patterns in Modern Global Precipitation, in: Swart, P.K., Lohman, K.C., McKenzie, J., Savin, S. (Eds.), *Climate Change in Continental Isotopic Records*. Washington, pp. 1–36. <https://doi.org/10.1029/GM078p0001>
- Sánchez-Goñi, M.F., Eynaud, F., Turon, J.-L., Shackleton, N.J., 1999. High resolution palynological record off the Iberian margin: direct land-sea correlation for the Last Interglacial complex. *Earth and Planetary Science Letters* 171, 123–137.
- Sánchez-Goñi, M.F., Landais, A., Cacho, I., Duprat, J., Rossignol, L., 2009. Contrasting intrainterstadial climatic evolution between high and middle North Atlantic latitudes: A close-up of Greenland Interstadials 8 and 12. *Geochemistry, Geophysics, Geosystems* 10, 1–16. <https://doi.org/10.1029/2008GC002369>
- Sánchez-Goñi, M.F. 2020. Regional impacts of climate change and its relevance to human evolution. *Evolutionary Human Sciences* (2020), 2, e55, page 1 of 27
- Sepulchre, P., Ramstein, G., Kageyama, M., Vanhaeren, M., Krinner, G., Sánchez-Goñi, M.F., d'Errico, F., 2007. H4 abrupt event and late Neanderthal presence in Iberia. *Earth and Planetary Science Letters* 258, 283–292. <https://doi.org/10.1016/j.epsl.2007.03.041>
- Skrzypek, G., Wiśniewski, A., Grierson, P.F., 2011. How cold was it for Neanderthals moving to Central Europe during warm phases of the last glaciation? *Quaternary Science Reviews* 30, 481–487. <https://doi.org/10.1016/j.quascirev.2010.12.018>
- Staubwasser, M., Drăguşin, V., Onac, B.P., Assonov, S., Ersek, V., Hoffmann, D.L., Veres, D., 2018. Impact of climate change on the transition of Neanderthals to modern humans in Europe. *Proceedings of the National Academy of Sciences* 115, 9116–9121. <https://doi.org/10.1073/pnas.1808647115>
- Tejada-Lara, J. V., MacFadden, B.J., Bermudez, L., Rojas, G., Salas-Gismondí, R., Flynn, J.J., 2018. Body mass predicts isotope enrichment in herbivorous mammals. *Proceedings of the Royal Society B: Biological Sciences* 285, 20181020. <https://doi.org/10.1098/rspb.2018.1020>
- Timmermann, A.: Quantifying the potential causes of Neanderthal extinction: Abrupt climate change versus competition and interbreeding, *Quaternary Science Reviews*, 238, 106331, <https://doi.org/10.1016/j.quascirev.2020.106331>, 2020.

- Tütken, T., Furrer, H., Vennemann, T.W., 2007. Stable isotope compositions of mammoth teeth from Niederweningen, Switzerland: Implications for the Late Pleistocene climate, environment, and diet. *Quaternary International* 164–165, 139–150. <https://doi.org/10.1016/j.quaint.2006.09.004>
- Vidal-Cordasco, M., Ocio, D., Hickler, T., Marín-Arroyo, A.B., 2022. Ecosystem productivity affected the spatiotemporal disappearance of Neanderthals in Iberia. *Nature Ecology & Evolution* 6, 1644–1657. <https://doi.org/10.1038/s41559-022-01861-5>
- Vidal-Cordasco, M., Terlato, G., Ocio, D., Marín-Arroyo, A.B. 2023. Neanderthal coexistence with *Homo sapiens* in Europe was affected by herbivore carrying capacity. *Sci. Adv.* 9, eadi4099(2023). DOI:10.1126/sciadv.adi4099

Other comments:

The name of one of the coauthors should be read as follows: Montserrat Sanz. Her edited affiliation is: 4 Grup de Recerca del Quaternari (GRQ-SERP), Department of History and Archaeology, Universitat de Barcelona, C/Montalegre 6-8, 08001 Barcelona, Spain. In addition, the affiliation of the first author (Mónica Fernández-García) has changed from the original submission and has been updated. Also, the affiliation of Sarah Pederzani has been adapted.

During the calculation of errors, it was identified that the correlations utilized for the conversion from $\delta^{18}\text{O}_{\text{phosp}}$ to $\delta^{18}\text{O}_{\text{mw}}$ do not correspond to the most updated version. The equations now chosen are the same as those employed in the Axlor site study (Pederzani et al., 2023), which includes a larger number of specimens and is more comprehensive. This, however, implies the modification of Figures 4 and 5. Numbers have also been updated in Tables 3 and 4, in the text and the Supplementary Information (SI). No significant implications have been detected, and the general interpretation aligns with the previous findings.

Throughout the document, minor modifications have also been made to correct grammatical errors or to make the text more fluid. During the revision process, we detected that the section describing the results of El Castillo, which was included in the original version revised by the authors, was missing from the submitted version. It has been added. After a thorough review of the discussion section, some changes have been made to enhance the overall flow and comprehension of the content. The aesthetics of some figures, such as Figure 1, have been improved. In addition, following the comments from the editor, the reference list and tables have been adjusted according to *Biogeosciences* guidelines.