

Investigating the effect of silicate- and calcium-based ocean alkalinity enhancement on diatom silicification

Aaron Ferderer^{1,2}, Kai G. Schulz³, Ulf Riebesell⁴, Kirralee G. Baker^{1,5}, Zanna Chase¹, Lennart T. Bach¹

5 ¹Institute for Marine and Antarctic Studies, Ecology & Biodiversity, University of Tasmania, Hobart, TAS, Australia.

²National Collections and Marine Infrastructure, Commonwealth Scientific and Industrial Research Organisation, Hobart, Tasmania, Australia

³Faculty of Science and Engineering, Southern Cross University, Lismore, NSW, Australia

10 ⁴Marine Biogeochemistry, Biological Oceanography, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁵The Australian Centre for Excellence in Antarctic Science, University of Tasmania, Hobart, Tasmania 7001, Australia (ACEAS)

15 *Correspondence to:* Aaron Ferderer (aaron.ferderer@utas.edu.au)

Abstract. Gigatonne-scale atmospheric carbon dioxide removal (CDR) will almost certainly be needed to supplement the emission reductions required to keep global warming between 1.5 - 2°C. Ocean alkalinity enhancement (OAE) is an emerging marine CDR method with the addition of pulverized minerals to the surface ocean being one widely considered approach. A concern of this approach is the potential for dissolution products released from minerals to impact phytoplankton communities. We conducted an experiment with 10 pelagic mesocosms (M1 – M10) in Raunefjorden, Bergen, Norway, to assess the implications of simulated silicate- and calcium-based mineral OAE on a coastal plankton community. Five mesocosms (M1, M3, M5, M7 and M9) were enriched with silicate (~75 $\mu\text{mol L}^{-1}$ Na_2SiO_3), alkalinity along a gradient from 0 to ~600 $\mu\text{mol kg}^{-1}$, and magnesium in proportion to alkalinity additions. The other five mesocosms (M2, M4, M6, M8, M10) were enriched with alkalinity along the same gradient and calcium in proportion to alkalinity additions. The experiment explored many components of the plankton community, from microbes to fish larvae, and here we report on the influence of simulated mineral based OAE on diatom silicification. Macronutrients (nitrate and phosphate) limited silicification at the onset of the experiment until nutrient additions on day 26. Silicification was significantly greater in the silicate-based mineral treatments, with all genera except *Cylindrotheca*, displaying an increase in silicification as a result of the increased concentration of dissolved silicate. In contrast to the effect of differences in dissolved silicate concentrations between the two mineral treatments, increases in alkalinity only influenced the silicification of two genera, *Pseudo-nitzschia* and *Nitzschia*. ~~*Pseudo-nitzschia* and *Nitzschia* were the only genera directly affected by alkalinity.~~ The four other genera (*Arcocellulus*, *Cylindrotheca*, *Skeletonema*, and *Thalassiosirra*) investigated here displayed no significant changes in silicification as a result of alkalinity increases between 0 and 600 $\mu\text{mol kg}^{-1}$ above natural levels. In summary, our findings illustrate that the enhancement of alkalinity via simulated silicate- and calcium-based methods has limited genus-specific impacts on the silicification of diatoms. This research underscores the importance of understanding the full breadth of different OAE approaches, their risks, co-benefits, and potential for interactive effects. ~~In summary, our findings suggest limited genus-specific impacts of alkalinity on diatoms, while also highlighting the importance of understanding the full breadth of different OAE approaches, their risks, co-benefits, and potential for interactive effects.~~

1 Introduction

Limiting global average surface temperature rise to 1.5 - 2°C above pre-industrial levels necessitates rapid reductions in global CO_2 (carbon dioxide) emissions as well as sustained atmospheric carbon dioxide removal (CDR) (IPCC, 2021; Van Vuuren et al., 2018). However, prior to considering the implementation of large scale CDR methods it is critical to assess the potential ecological impacts of these methods (Bach et al., 2019; Fuss et al., 2018; Renforth and Henderson, 2017).

Ocean alkalinity enhancement (OAE) is considered to be a promising marine CDR method due to its potential to remove CO_2 at a gigatonne scale (Burt et al., 2021; Feng et al., 2017; He and Tyka, 2023; Ilyina et al., 2013; Keller et al., 2014; Paquay and Zeebe, 2013). There are various approaches to implementing OAE, each with techno-economic and environmental advantages and disadvantages (Renforth and Henderson, 2017). Irrespective of the method, all approaches aim to increase the capacity of the ocean to store atmospheric CO_2 by increasing the alkalinity of seawater through the addition of substances which increase alkalinity, the removal of acid and/or

55 neutralisation of protons in seawater, all of which will increase seawater pH (Eisaman et al., 2023). One widely discussed method involves the addition of pulverized alkaline minerals such as magnesium silicates or calcium hydroxides to the surface ocean (Bach et al., 2019; Kheshgi, 1995; Renforth and Henderson, 2017).

In order for minerals to be suitable they must be alkaline, have relatively rapid dissolution rates, and are ideally inexpensive, readily available, and contain minimal potential contaminants (Hartmann et al., 2013; Renforth and Henderson, 2017). Minerals fitting some or most of these criteria include olivine, a silicate-based naturally occurring mineral, as well as quick and/or hydrated lime which are anthropogenic calcium-based minerals (Renforth and Henderson, 2017). However, the effects of dissolution products derived from these minerals on marine communities is yet to be fully assessed (Bach et al., 2019). Indeed, hotspots of dissolution products from mineral-based OAE will inevitably occur at sites of mineral additions resulting in high concentrations of e.g., 65 Mg^{+} , $Si(OH)_4$, Ca^{+} , increased pH, and trace metals (Hartmann et al., 2013). Dissolution products may act to fertilize some organisms while inhibiting others, potentially leading to shifts in plankton communities (Bach et al., 2019; Guo et al., 2022; Hutchins et al., 2023). For example, calcium-based minerals are hypothesized to benefit pelagic and benthic calcifiers, with some studies supporting this (Albright et al., 2016; Bach et al., 2015; Gore et al., 2019) while others found neutral responses (Gately et al., 2023). In contrast, the dissolution of silicate based 70 minerals is expected to benefit silicifying plankton species including diatoms (Bach et al., 2019; Egge and Aksnes, 1992; Hauck et al., 2016). Thus, we expect mineral based OAE to have some impact on marine communities with these impacts being highly dependent on the source minerals used. However, it is important that the environmental impacts and/or co-benefits resulting from OAE are evaluated against the potential climatic benefits.

This study aims to specifically assess and compare the potential implications of increased alkalinity, silicate and 75 magnesium concentrations associated with silicate- based mineral OAE and increased alkalinity and calcium concentrations associated with calcium-based mineral OAE on a coastal plankton community. In order to capture the potential maximum and minimum acceptable levels of alkalinity enhancement, we increased concentrations of alkalinity in steps of $150 \mu mol kg^{-1}$ by 0 to $\sim 600 \mu mol kg^{-1}$. Such an increase in alkalinity is expected to influence the phytoplankton community as concentrations of CO_2 decrease below previously observed thresholds 80 limiting growth (Chen and Durbin, 1994; Hinga, 2002; Paul and Bach, 2020; Riebesell et al., 1993).

Previous work has identified that increases in alkalinity of $500 \mu mol kg^{-1}$ resulted in a significant decrease in silicate uptake and biogenic silica production (Ferderer et al., 2022). Furthermore, it is well known that silicate has a fertilising effect on diatoms, with concentrations above $2 \mu mol kg^{-1}$ often resulting in their dominance within plankton communities (Egge and Aksnes, 1992; Escaravage and Prins, 2002). The dissolution of silicate-based 85 minerals such as olivine to enhance alkalinity is predicted to significantly increase silicate concentrations at sites of addition and projected to induce diatom blooms (Hauck et al., 2016). The influence of varying silicate concentrations on diatoms is well known, however the interaction between OAE and enhanced silicate concentrations as a result of silicate based mineral dissolution is yet to be fully explored. Thus, in this study we focus on assessing the influence of mineral based OAE, along an increasing gradient, on the incorporation of silica 90 into the frustules of diatoms (silicification). Our primary goal is to elucidate the potential risks and or co-benefits of mineral based alkalinity enhancement on diatoms.

2 Methods

2.1 Mesocosm deployment and maintenance

95 On the 7th of May 2022, ten Kiel Off-Shore Mesocosms for Ocean Simulations (KOSMOS, M1-M10; Riebesell
et al., 2013) were deployed from RV ALKOR in Raunefjorden, Bergen, Norway ~1.5 km from the Espegrand
marine research field station (Fig. 1). Mesocosms consisted of a cylindrical polyurethane bag 20 m in length, (2
m in diameter, $\sim 60.01 \pm 0.01 \text{ m}^3$ volume). Mesocosm bags were fitted within 8 m tall floating frames during
deployment which were slowly lowered into the fjord, allowing the bags to gently fill while minimising
disturbance to the plankton community. Once deployed the mesocosm bags remained open at their base (~20 m)
100 and top (~1 m below the sea surface) allowing water exchange between the mesocosms and fjord. Mesocosms
were closed off to the fjord on the 13th of May when divers attached a 2 m long funnel shaped sediment trap to
each mesocosm, and the top of each mesocosm bag was raised ~1 m above the surface (Fig. 1). A ring with the
same diameter as the mesocosms fitted with a 1 mm mesh was passed through each mesocosm after closing to
remove any large nekton or plankton from the mesocosms. The sealing off of the mesocosms from the fjord
105 marked the beginning of the experiment (Day 0). The volume of each mesocosm was determined on day 2 of the
experiment via the addition of a NaCl brine solution. Water inside mesocosms was first homogenized by bubbling
compressed air up through the mesocosms. Following homogenisation, 50 L of NaCl brine was evenly added to
the mesocosm via a bespoke distribution device called “the spider” (Riebesell et al., 2013). The precise addition
of NaCl enabled us to calculate the volume of each mesocosm following (Czerny et al., 2013). Mesocosm bags
110 were cleaned approximately every week from the inside and outside to minimise any potential biofouling which
may impact the results of the experiment. External cleaning of the mesocosms was conducted by divers and/or
surface attendants in small boats using brushes. Internal cleaning of the mesocosms was conducted using a large
ring with rubber blades that was the same diameter as the mesocosms. This ring was sunk inside mesocosms with
a 30 kg weight attached to its base, removing any growth from the inner walls of the mesocosm bags down to 1
115 m above the sediment trap.




Figure 1. Infographic depicting the relevant information pertaining to the mesocosm design and mooring site approximately 1.5 km from the Espesgrend Marine research field station in Bergen, Norway (maps produced in Rstudio using ©Google maps data).

2.2 Setup of OAE treatments and nutrient fertilisation

Mesocosms were split into two treatment groups; a calcium-based (Ca-OAE) treatment (N =5) and silicate-based (Si-OAE) treatment (N =5) with one mesocosm in each group serving as a control. Alkalinity was enhanced along a gradient in each mineral-based treatment ranging from 0 – 600 $\mu\text{mol kg}^{-1}$ using varying amounts of NaOH (Merck), dissolved in 20 L of Milli-Q®. Simulated differences in the type of OAE were established via the addition of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ in the Ca-OAE treatments and $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ and $\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$ in the Si-OAE treatments, all dissolved in 20 L of Milli-Q®. The simulated enhancements of Mg^{2+} and Ca^{2+} were proportional to the addition of NaOH, i.e., increases by half the alkalinity enhancement. In contrast Na_2SiO_3 was increased by equal concentrations (target of 75 $\mu\text{mol L}^{-1}$) in all mesocosms within the Si-OAE treatment group (including the control), instead of a gradient from 0 - 150 $\mu\text{mol L}^{-1}$, which would be the corresponding concentrations for olivine dissolution. This approach was adopted due to metasilicate solubility restrictions (data not shown), the potential for colloid formation to occur at high concentrations and enable clear distinctions to be made between silicate and TA-alkalinity effects. Finally, the increase in TA-alkalinity from silicate additions in a 2:1 ratio was taken into

account by reducing respective NaOH additions and the addition of HCl in the silicate-based control
135 ($\Delta_{\text{alkalinityTA}} = 0 \mu\text{mol kg}^{-1}$).

At the time of closure, all mesocosms had low concentrations of macronutrients ($0.10 \pm 0.019 \mu\text{mol L}^{-1} \text{NO}_3^-$,
0.03 \pm 0.005 $\mu\text{mol L}^{-1} \text{PO}_4^{3-}$ and $0.16 \pm 0.048 \mu\text{mol L}^{-1} \text{Si(OH)}_4$). After observing communities in a prolonged
phase of oligotrophic conditions, macronutrients were added to the mesocosms on day 26 (final range across
140 mesocosms = $3.59 - 3.8 \mu\text{mol L}^{-1} \text{NO}_3^-$, $0.19 - 0.24 \mu\text{mol L}^{-1} \text{PO}_4^{3-}$ and $0.39 - 1.03 \mu\text{mol L}^{-1} \text{Si(OH)}_4$) to stimulate
the phytoplankton community. Macronutrients were added to the mesocosms as two separate 20 L solutions with
one consisting of NaNO_3 (Merck, > 99.5%) and $\text{Na}_2\text{HPO}_4 \cdot \text{H}_2\text{O}$ (Merck, > 99.5%) and the other consisting of
 $\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$ (Roth > 95%). Inorganic nutrient concentrations were measured the day before nutrient additions
and ~2 hours after the addition of nutrients to quantify the additions and ensure appropriate stoichiometry within
145 mesocosms. After the addition of nutrients, it was noted that the stoichiometry of macronutrients was not even
across mesocosms, later identified to have been the result of a mistake during solution preparation. As such, a
second addition of nitrate was completed on day 28 for those mesocosms below target concentrations. All
solutions were added homogeneously to mesocosms using the spider distribution device.

150 Given the additions of alkalinity on day 7 and macronutrients on day 26 and 28, the experiment was divided into
three distinct phases; phase 0 representing the period prior to alkalinity enhancement (day 0 – 6), phase I
representing conditions prior to the addition of macronutrients (day 7 - 28) and phase II representing the period
after nutrient additions (day 29 – 54).

2.3 Sampling methods

155 Sampling of the mesocosms was conducted every second day from small boats with sediment sampling first (0800
– 1000; here and in the following GMT +2) followed by particulate and dissolved substance sampling (0900 –
1300), zooplankton sampling (1000 – 1300) and finally CTD, FastOcean APD/fluoroprobe (1400 – 1600). With
the exception of particulate and dissolved substance sampling, which was carried out at random, mesocosms were
sampled in order from M1 through to M10 with fjord samples taken directly next to M5. Sample containers were
160 stored in boxes to avoid excess light and heat exposure during sampling and upon return to the research station
(directly after each round of sampling) were transferred to a room at ambient water temperature (8.7 – 15.4 °C)
until further processing. The sampling schedule remained consistent with the exception of day 15 where only
sediment sampling was undertaken due to unsafe weather conditions. Additional samples for the determination of
dissolved inorganic nutrients were taken on day 26 and day 28 to assess stoichiometry post nutrient additions
165 performed earlier the same day.

Sinking particles were collected from the sediment traps of each mesocosm via a silicon tube attached to the base
of the sediment trap at one end and a manual vacuum pump at the surface (Boxhammer et al., 2016). Suspended
particulate matter and dissolved substances were collected using 5 L integrated water samplers (IWSs; Hydro-
170 Bios, Kiel). IWSs were equipped with pressure sensors enabling an even collection of water within a specified
depth, from the surface to the top of the sediment trap (0 – 20 m). Four IWSs were taken within each mesocosm
and fjord which were transferred into 10 L polyethylene carboys. Samples for quantification of changes in

carbonate chemistry were collected from the first IWS taken within each mesocosm and filled directly into 500 ml glass bottles following protocols outlined in SOP 1 from (Dickson et al., (2007)).

175

2.4 Carbonate chemistry and dissolved inorganic nutrients

Samples for total alkalinity (TA), pH and dissolved inorganic nutrients were sterile filtered using a peristaltic pump and 25 mm, 0.2 µm pore size, PES membrane, syringe filters to minimise biological processes and remove particles which can influence respective analyses. Dissolved inorganic nutrient concentrations NO_3^- , NO_2^- , PO_4^{3-} , and Si(OH)_4 were determined spectrophotometrically following methods outlined by (Hansen and Koroleff, (1999)). Dissolved inorganic nutrient samples were measured in triplicate to control for technical variability between measurements across the experiment. TA was determined using a two-step open cell titration following SOP3b outlined by (Dickson et al., (2007)). TA samples were measured in duplicate on a Metrohm 826 Compact Titrosampler coupled with an Aquatrode Plus with PT1000 temperature sensor and calibrated against certified reference material (CRM batch 193) supplied by Prof. Andrew Dickson's laboratory. pH was determined in duplicate via spectrophotometric methods outlined in (Dickson et al., (2007)) (not shown here).

180

185

2.5 Particulate matter analysis

Sediment trap samples were processed immediately upon return of the sampling boat to the research station. Sample weight was first determined gravimetrically before resuspension of particles and homogenisation of the sample for subsampling (e.g., dissolution assays, particle sinking velocity). The remaining sample was enriched with 3M FeCl_3 followed by 3M NaOH to enhance flocculation, coagulation and subsequent sedimentation of particles while maintaining pH (Boxhammer et al., 2016). Approximately 1 hour after settling the supernatant was removed and samples centrifuged in two steps: first for 10 min at 5200 g in a 6-16KS centrifuge (Sigma) and then again for 10 min at 5000 g in a 3K12 centrifuge (Sigma). Following each step any supernatant was removed and the remaining pellet freeze dried to remove residual moisture. Finally, the dried samples were pulverised into a homogenous powder using a cell mill and were transported to GEOMAR, Kiel, Germany for further analysis. Subsamples of the powder used to determine concentrations of biogenic silica (BSi) were placed in 60 ml Nalgene™ polypropylene bottles, filled with 25 ml of 0.1 M NaOH solution, and then placed in a shaking water bath at 85 °C. After 135 min the bottles were removed and cooled before the addition (25 ml) of 0.05 M H_2SO_4 to stop the leaching processes. The concentration of dissolved silicate was then measured spectrophotometrically following (Hansen and Koroleff, (1999)). BSi concentrations of the measured subsamples were then scaled to represent the total sample from sediment traps, normalised to mesocosm volume and are reported as the accumulation of BSi in the sediments over the experimental period.

190

195

200

205

210

Analysis of major elemental pools and phytoplankton pigments in the water column subsamples (pre-filtered through a 200 µm screen) of 0.5 – 1 L were taken from carboys after gentle mixing to homogenise samples. Subsamples for BSi were filtered onto cellulose acetate filters (pore size 0.45 µm) using a vacuum filtration system at ≤ 200 mbar and stored in plastic vials at -20 °C until analysis the following day. Filters were then placed in 60 ml Nalgene™ bottles, digested, and analysed following the same methods for BSi in the sediments described above. Chlorophyll *a* was filtered onto glass fibre filters (GFF, nominal pore size = 0.7 µm) while minimising

light exposure. Immediately after filtration filters were stored in plastic vials at -80 °C until analysis the following day. Samples were extracted with 90% acetone and homogenised using glass beads in a cell mill. After homogenisation samples were centrifuged (10 min 800 g, 4°C), then the supernatant was removed and analysed on a fluorometer (Turner 10-AU) to determine Chl *a* concentrations (Welschmeyer, 1994).

215

2.6 PDMPO labelling and determination of silicification rates via fluorescent microscopy

To investigate differences in diatom silicification, 350 ml samples were collected by means of IWS from each mesocosm every two to four days (variation in sampling schedule occurred due to unforeseen circumstances such as extreme weather events and COVID-19 infections). Due to the low abundance of diatoms within mesocosms, (as observed through microscopy and BSi concentrations) samples were gravimetrically concentrated from 350 ml to 70 ml using a 47 mm filtering apparatus and polycarbonate filters (3 µm pore size). Organisms were resuspended by gentle stirring and inversion of the filtration system before the sample was transferred to three 17 ml polycarbonate tubes. The fluorescent dye 2-(4-pyridyl)-5-((4-(2-dimethylaminoethylaminocarbonyl)methoxy)phenyl)oxazole (PDMPO; LysoSensor yellow/blue DND-160 from ThermoFisher Scientific) was added at a final concentration of 0.125 µM before samples were incubated in a flow-through incubator placed outside of the Espegrend Marine research field station for temperature control (inlet location was ~1.5 km from the mesocosms so that temperature in the incubator was similar to mesocosms). The incubator was screened with shade cloth to ~30% incident irradiance and incubations lasted for ~24 hours. At the conclusion of the incubation, 15 ml of the sample was filtered under gentle vacuum pressure onto 25 mm, 0.8 µm black polycarbonate membrane filters. Filters were mounted onto microscope slides with a drop of Prolong gold antifade followed by a glass coverslip and sealed with clear nail varnish. Prepared slides were then stored in the dark at -20 °C for later analysis via fluorescent microscopy at the University of Tasmania, Australia (within 6 months).

Prepared microscope slides were imaged using the software NIS-elements and a Nikon eclipse Ci microscope equipped with a UV-1A (longpass) filter cube, Nikon DS-Ri2 camera and Mercury lamp (Nikon C-SHG1). Prior to analysing slides, a yellow fluorescence slide (Thorlabs FSK3) was imaged ten times and the average fluorescence subtracted from all images taken that day to account for variation in the mercury lamp across imaging days. The entirety of each filter was systematically scanned at ×200 magnification, when a cell was located, it was imaged at ×400 magnification with all cells that were appropriate for measurements (e.g., not overlapping or partially destroyed cells) included in analysis. This gave final counts for each genus ranging from 1 – 176 cells per mesocosm on any given day. Cells were, when possible, identified to genus level as brightfield imaging was not possible on the black polycarbonate filters and fluorescent images did not provide enough detail for accurate identification beyond this level. However, in instances where differentiation between genera was difficult or impossible to complete with high confidence cells were classified based on significant differences in the shape, size and/or details of the frustule of cells. Each genus/group is therefore comprised of similar cells which show distinct differences in characteristics which influence the fluorescence of cells (Table S1). Images were later analysed by quantifying single cell fluorescence following a custom-made procedure in ImageJ on the original TIFF images. For each image, cell/s were selected so that minimal background area was included before the

250 fluorescence of the selected cell was recorded. Background fluorescence was measured at four locations directly
surrounding the cell where no other cells were present, and the average background fluorescence subtracted from
cell fluorescence. Total cell fluorescence, corrected for background fluorescence, was then normalised to cell area
to give mean cell fluorescence. Due to low abundances of diatoms and thus insufficient counts for meaningful
analyses, only days 7, 11 and 17 were analysed prior to the addition of nutrients on day 26 and 28. After the
255 addition of nutrients all filters were analysed, however due to an outbreak of COVID-19, samples for the
determination of silicification could not be taken for the final eight days of the experiment.

Due to technical issues and the unavailability of specific instrumentation, we were unable to measure total
community fluorescence during the experiment and as such convert fluorescence values to BSi incorporation.
260 However, the aim of this experiment was to identify any relative differences in taxa specific rates of silicification,
something which is not achievable through the measurement of BSi. This was achieved with the use of the
fluorescent dye PDMPO which is incorporated at a rate proportional to BSi incorporation in diatoms (Leblanc and
Hutchins, 2005; Mcnair et al., 2015; Znachor and Nedoma, 2008). PDMPO uptake and subsequent fluorescence
within diatom cells therefore provides an appropriate proxy for the incorporation of silica into newly formed
265 frustules irrespective of the units. As such the PDMPO-fluorescence of cells measured here as the fluorescence
of a given cell normalised to cell area, is referred to as silicification throughout this text.

2.7 Statistical analysis

To explore the effect of the alkalinity source mineral and alkalinity enhancement across mesocosms we first
visualised the dataset using non-metric multidimensional scaling (NMDS) plots. Three separate plots were
270 produced to explore the effects of (a) the treatments over the total extent of the experiment, (b) prior to the addition
of nutrients, and (c) post nutrient addition on silicification.


A linear mixed-effects model was used to quantify the influence of the treatments (total alkalinity and alkalinity
source mineral) on diatom silicification. The model was run with alkalinity source mineral, total alkalinity,
275 experimental phase, and diatom genus as fixed effects, and silicification (square root transformed) as the
dependent variable. To account for temporal pseudo-replication in the models, mesocosm ($N = 10$) was nested
within sampling occasion (day) and fitted as a random effect in all models. Several linear mixed-effects models
were fit, with non-significant interactions removed and Akaike Information Criteria (AIC) used to determine the
best model. All statistical analyses were conducted in Rstudio v 2023.6.1.524 (Posit Team, 2023). Estimated
280 marginal means were calculated using the package *emmeans* (Lenth et al., 2023) to determine the significance of
two and three-way interactions within the model. Estimated marginal means of linear trends were calculated using
emtrends from the package *emmeans*, to assess the effect of total alkalinity within significant interactions in the
linear model (Lenth et al., 2023).


285 Finally linear models were used to assess the influence of total alkalinity and alkalinity source mineral on the
concentration of BSi in the water column and accumulation of BSi in the sediments. Linear models were run for
each phase of the experiment with mean water column BSi and mean accumulated sediment BSi over each phase
fitted as dependent variables and mean total alkalinity and alkalinity source mineral as fixed effects. An additional

linear model was run to assess total accumulated BSi in the sediment trap with the accumulated sediment BSi up until day 53 fitted as the dependent variable and total alkalinity and alkalinity source mineral fitted as fixed effects. All statistical analyses including NMDS plots were conducted in Rstudio v 2023.6.1.524 (Posit Team (2023), 2023).

3 Results

~~Chlorophyll *a* concentrations were relatively low at the beginning of the experiment with $1.01 \pm 0.17 \mu\text{g L}^{-1}$ (mean \pm SD) on day 3 (Fig 2d). Concentrations of NO_3^- were below detection limit, thereby constraining phytoplankton growth during phase I of the experiment (mean NO_3^- day 7 – 25 = $0.004 \pm 0.035 \mu\text{mol L}^{-1}$) (Fig 2a). In contrast there was residual PO_4^{3-} ($0.021 \pm 0.022 \mu\text{mol L}^{-1}$) and Si(OH)_4 (Ca-OAE treatment = $0.202 \pm 0.99 \mu\text{mol L}^{-1}$, Si-OAE treatment = $67.929 \pm 1.04 \mu\text{mol L}^{-1}$) which likely supported the phytoplankton community in utilising remineralised nitrogen until the addition of nutrients on day 26 and 28 (Fig 2b, 2c). Although $\sim 759 \mu\text{mol L}^{-1}$ of Na_2SiO_3 was added to the Si-OAE treatment, there was no discernible depletion of Si(OH)_4 during phase I (Fig 2c). The addition of macronutrients (NO_3^- , PO_4^{3-} , and Si(OH)_4) can be seen on day 26, with a secondary addition on day 28 to correct for unwanted differences in the stoichiometry between mesocosms (Fig 2). Chlorophyll *a* concentrations were relatively low at the beginning of the experiment with $1.01 \pm 0.17 \mu\text{g L}^{-1}$ (mean \pm SD) on day 3 (Fig 2d). During phase II, nutrients steadily declined until the majority was depleted between days 39 and 49 (Fig 2). Chlorophyll *a* remained low until day 33, after which it increased across all mesocosms at rates between $0.03 - 0.68 \mu\text{g L}^{-1}$ per day (Fig 2d). The delayed/slow increase in chlorophyll *a* was likely due to the prolonged nutrient deficit within the mesocosms and subsequent small seed population. There was no discernible relationship between total alkalinity and chlorophyll *a*, NO_3^- or PO_4^{3-} observed across the extent of the experimental period or in a particular phase (Fig 2). However, in the Si-OAE treatments initial concentrations of Si(OH)_4 were lowest in the high alkalinity mesocosm, with a difference of $2.45 \mu\text{mol L}^{-1}$ between the $\Delta 0$ and $\Delta 600 \mu\text{mol kg}^{-1}$ alkalinity treatments (Fig 2c). This trend appeared directly after the addition of the treatments but disappeared once nutrient uptake began (Fig 2c). Chlorophyll *a* peaked between days 39 and 49 for all mesocosms with the Si-OAE treatment exhibiting a marginally higher mean peak in chlorophyll *a* (Si-OAE treatment: $3.78 \pm 1.7 \mu\text{g L}^{-1}$, Ca-OAE treatment: $2.97 \pm 0.3 \mu\text{g L}^{-1}$, Fig. 2d). A similar trend is observed in the uptake of NO_3^- with mesocosms in the Si-OAE treatment, exhibiting marginally faster depletion of NO_3^- ($-0.192 \mu\text{mol L}^{-1}$ per day) in comparison to the Ca-OAE treatment ($-0.158 \mu\text{mol L}^{-1}$ per day). There was no discernible relationship between total alkalinity and chlorophyll *a*, NO_3^- or PO_4^{3-} observed across the extent of the experimental period or in a particular phase (Fig 2). However, in the Si-OAE treatments concentrations of Si(OH)_4 were lower in mesocosms with high alkalinity, with a difference of $2.45 \mu\text{mol L}^{-1}$ between the $\Delta 0$ and $\Delta 600 \mu\text{mol kg}^{-1}$ alkalinity treatments (Fig 2c). This trend appeared directly after the addition of the treatments early in the experiment and was conserved until the end of the study (Fig 2e).~~





325 **Figure 2.** Temporal variation in dissolved inorganic nutrients **(a)** nitrate (NO_3^-), **(b)** phosphate (PO_4^{3-}), **(c)** silicate (Si(OH)_4), **(d)** chlorophyll *a* (Chla) and, **(e)** total alkalinity. Dissolved inorganic nutrient measurements commenced on day 0 while chlorophyll *a* measurements commenced on day 3. Vertical dashed lines represent the respective phases of the experiment, phase 0 (pre-alkalinity enhancement), phase I (pre-nutrient addition), phase II (post-nutrient addition).

330

Nonmetric multidimensional scaling (nMDS) (Fig. 3) revealed distinct distances among treatments, including different alkalinity source minerals and total alkalinity, in relation to silicification of the various diatom genera, ~~although relatively small distances are observed between *Pseudo-nitzschia* and *Nitzschia*, likely due to their morphological similarities and therefore similar silica content~~. The distances between polygons, representing the Si- and Ca-based mineral treatments, indicate differences in silicification between the Si- and Ca-based mineral treatments over the total experimental period and after the addition of nutrients (Fig. 3a,c). In contrast, during phase I (pre-nutrient addition) polygons representing the Si- and Ca-based mineral treatments are overlapping, suggesting a weak relationship between alkalinity source mineral (Ca or Si) and silicification prior to the addition of nutrients (Fig. 3b). In all plots, symbols representing the differing levels of total alkalinity within the Si-based treatment are relatively closer, when compared to the Ca-based treatment which exhibits greater spread between differing levels of total alkalinity.




Figure 3. Nonmetric multidimensional scaling ordination exploring the mean silicification of diatom genera across (a) the complete extent of the mesocosm experiment (stress = 0.0502), (b) pre-nutrient addition, phase I (stress = $3.62e^{-5}$), and (c) post nutrient addition, phase II (stress = 0.0612). Due to the low stress values obtained (<0.20), it is assumed that all configurations accurately represent distinct dissimilarities in silicification among diatom genera.

3.1 Results of linear mixed effects model

Analysis of diatom silicification supported the distances observed within the nMDS plots with alkalinity source mineral (Ca or Si) having a significant influence on the silicification of diatom cells (Table 1, Fig. 4). Cells

exposed to the Si-OAE treatment were more heavily silicified irrespective of changes in total alkalinity (Fig. 4). However, the significant interaction between alkalinity source mineral and genus indicates that the effect of the alkalinity source mineral on silicification varies between genera (Table 1). All genera displayed significant differences between the two alkalinity source mineral treatments with the exception of *Cylindrotheca* which showed no significant difference in silicification between the Si- or Ca-based mineral treatments (Table 2). Total alkalinity had a significant effect on silicification between the two experimental phases (t ratio = -2.16, p = 0.03), with silicification increasing as a function of total alkalinity during phase I of the experiment (emtrend = 0.24, t ratio = 2.57, p = 0.01) (Table 1, Figure 4). However, investigation of the three-way interaction revealed the significant effect of total alkalinity on diatom silicification only to be present in phase I of the experiment in the Si-based OAE treatment (t ratio = 3.22, p = 0.002) (Fig. 4, Table 1). Furthermore, the significant interaction between genus and total alkalinity suggests the influence of total alkalinity on silicification varies between genera. Finally, the significant three-way interaction between alkalinity source mineral, genus and total alkalinity suggests that the effect of alkalinity on silicification in given genera was dependent on the alkalinity mineral source (Table 1). Exploration of this interaction revealed the silicification of cells in the genus *Pseudo-nitzschia* ($N = 3510$) to be significantly influenced by alkalinity in both the Ca and Si-based treatments, with silicification increasing with increasing alkalinity (Table 3). In contrast, the genus *Nitzschia* ($N = 677$) displayed a significant increase in silicification in the Si-based treatment only while no other genera displayed significant differences in silicification between total alkalinity levels (Table 3). There were also significant differences in silicification between genera, however this was reduced from eight significant differences in the Si-based mineral treatment to five in the Ca-based mineral treatment (Table S2).

Table 1. Statistical results of linear mixed-effects model assessing the influence of mineral based OAE on diatom silicification.

Source of variation	<i>df</i>	<i>F-value</i>	<i>P-value</i>
Alkalinity source mineral (Ca or Si)	1,64	146.95	<0.001*
Genus	5,11330	1305.96	<0.001*
Total alkalinity	1,64	1.59	0.21
Phase	1,6	5.26	0.06
Alkalinity source mineral*Genus	5,11330	82.07	<0.001*
Alkalinity source mineral*Total alkalinity	1,64	0.04	0.84
Total alkalinity*Genus	5,11330	57.65	<0.001*
Alkalinity source mineral*phase	1,64	1.29	0.26
Total alkalinity*phase	1,64	4.64	0.03*
Alkalinity source mineral*Genus*Total alkalinity	5,11330	5.93	<0.001*
Alkalinity source mineral *Total alkalinity*phase	1,64	4.02	0.05*

Table 2. Results of post-hoc test (*emmeans*) comparing the influence of silicate- and calcium-based mineral treatments on genera specific silicification.

Genus	Contrast	Estimate	Standard error	t-ratio	p-value
<i>Arcocellulus</i>	Ca – Si	-0.69	0.08	-8.64	<0.001*
<i>Cylindrotheca</i>	Ca – Si	0.07	0.11	-0.67	0.51
<i>Nitzschia</i>	Ca – Si	-0.60	0.10	-6.02	<0.001*
<i>Pseudo-nitzschia</i>	Ca – Si	-0.41	0.08	-5.04	<0.001*
<i>Skeletonema</i>	Ca – Si	-1.24	0.09	-14.34	<0.001*
<i>Thalassiosirra</i>	Ca – Si	-0.91	0.09	-10.68	<0.001*

Table 3. Results of post-hoc test (*emtrends*) assessing the influence of total alkalinity on genera specific silicification in across the two alkalinity source mineral treatments.

Genus	Mineral	Total alkalinity mean trend	Standard error	t-ratio	p-value
<i>Arcocellulus</i>	Ca	0.004	0.07	0.06	0.96
<i>Cylindrotheca</i>	Ca	0.10	0.09	1.13	0.26
<i>Nitzschia</i>	Ca	0.08	0.08	1.07	0.29
<i>Pseudo-nitzschia</i>	Ca	0.33	0.07	4.52	<0.001*
<i>Skeletonema</i>	Ca	0.06	0.08	0.79	0.43
<i>Thalassiosirra</i>	Ca	-0.06	0.08	-0.76	0.45
<i>Arcocellulus</i>	Si	0.00	0.07	0.01	0.99
<i>Cylindrotheca</i>	Si	0.06	0.09	0.64	0.52
<i>Nitzschia</i>	Si	0.40	0.08	5.02	<0.001*
<i>Pseudo-nitzschia</i>	Si	0.37	0.07	5.31	<0.001*
<i>Skeletonema</i>	Si	0.03	0.07	0.50	0.62
<i>Thalassiosirra</i>	Si	-0.04	0.07	-0.61	0.55





Figure 4. Single cell silicification of the diatom community depicted as mean fluorescence (PDMPO) normalised to cell surface area and reported in fluorescence units (FU). Data visualised as box plots, with colours representing the different mineral sources (Ca-OAE; blue and Si-OAE; green) and shading indicating the total alkalinity gradient with darker colours indicating higher total alkalinity. Nutrient addition on day 26 and 28 is represented by the dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition).





395 **Figure 5.** Boxplots depicting single cell silicification of different genera. Silicification is shown as mean cell
 fluorescence (PDMPO) normalised to cell area and reported in fluorescence units (FU). Colours representing the
 different mineral sources (Ca-OAE; blue and Si-OAE; green) and shading indicating the total alkalinity gradient
 with darker colours indicating higher total alkalinity. Nutrient addition on day 26 and 28 is represented by the
 dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition), with
 400 genera names in the top right of each plot.

Concentrations of BSi in the water column can be seen decreasing from day 0 (mesocosm closure) and remaining low until day ~33 (Fig 6a). Directly after the addition of Na₂SiO₃ (day 7), BSi in the water column spiked for one day in the silicate-based treatments, while no increase was observed in the calcium-based treatment (Fig 6a). BSi began to increase in all mesocosms between days 33-35, however concentrations in the Ca-OAE treatments remained relatively low (<2 μmol L⁻¹) for the extent of the experiment (Fig 6a). We observed no significant relationship between BSi and alkalinity in either of the alkalinity source mineral treatments (Ca or Si) or any phase of the experiment (Table 3). During phase II, concentrations of BSi in the water column were significantly higher in the Si-OAE treatment when compared to the Ca-OAE treatment (Table 4, Fig 6a). Additionally, we observed significant differences in the accumulation of BSi in the sediments between the two alkalinity source mineral types (Table 4, Fig. 6b). Similar to the water column, there was an initial increase in sedimented BSi in the Si-OAE treatment when compared to the Ca-OAE treatment (Fig 6b). During phase I there was a 0.47 μmol L⁻¹ difference in the amount of BSi accumulated in the sediment trap between the highest (Δ600 μmol kg⁻¹) and lowest (Δ0 μmol kg⁻¹) alkalinity levels in the Si-OAE treatment. However, there was less variability between the Δ150, Δ300 and Δ450 alkalinity treatments, contributing to the non-significant effect of alkalinity on BSi accumulation in the sediment trap (Table 3). Furthermore, irrespective of experimental phase, there was no significant relationship between alkalinity and BSi accumulated in the sediment.


Table 4. Results of linear models exploring the effect of alkalinity source mineral and total alkalinity on average concentrations of BSi in the water column or sediment trap for a given phase.

Phase	Parameter	Source of variation	df	F-value	P-value
Phase 0	BSi water column	Mineral	1	0.70	0.43
		TA	1	0.70	0.43
Phase 0	BSi sediment	Mineral	1	0.20	0.67
		Total alkalinity	1	0.35	0.57
Phase I	BSi water column	Mineral	1	70.88	<0.001*
		TA	1	1.76	0.23
Phase I	BSi sediment	Mineral	1	46.44	<0.001*
		Total alkalinity	1	4.80	0.06
Phase II	BSi water column	Mineral	1	47.15	<0.001*
		TA	1	2.02	0.20
Phase II	BSi sediment	Mineral	1	82.6109	<0.001*
		Total alkalinity	1	0.64	0.45
Day 53	BSi sediment	Mineral	1	45.57	<0.001*
		Total alkalinity	1	0.0005	0.98



Mineral-TA ($\mu\text{mol kg}^{-1}$)

Ca-0	Ca-150	Ca-300	Ca-450	Ca-600
Si-0	Si-150	Si-300	Si-450	Si-600



425 **Figure 6.** Temporal variations of (a) biogenic silica (BSi) in the water column, and (b) accumulation of BSi in the sediment trap across alkalinity source minerals and total alkalinity treatments during the extent of the experimental period. Nutrient addition on day 26 and 28 is represented by the dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition).

4 Discussion

430 Understanding the potential environmental implications of CDR methods such as OAE is a crucial step before decisions are made upon their implementation at large scales. The aim of this mesocosm study was to form part of this research by assessing the potential effects of calcium- and silicate-based mineral OAE on a coastal plankton

community. Here, we specifically discuss the influence of simulated mineral based OAE on diatom community and genus specific silicification. Our results revealed silicate fertilisation associated with silicate-based OAE to significantly increase silicification ~~in of~~ the diatom community and all genera with the exception of *Cylindrotheca*. It is important to note that low initial concentrations of dissolved silicate likely facilitated this observed increase in silicification. Under conditions where dissolved silicate is already replete, further increases due to silicate-based OAE may not result in similar increases in additional silicification. This increase in silicification was primarily a result of the ~~increased difference in the concentration of~~ dissolved silicate concentrations ($\Delta 75 \mu\text{mol kg}^{-1}$) ~~in~~ the between the silicate-based and calcium-based OAE treatments rather than an increase in alkalinity. However, in addition to the influence of silicate fertilisation, *Pseudo-nitzschia* and *Nitzschia* were both significantly affected by changes in alkalinity. *Pseudo-nitzschia* exhibited a significant increase in silicification with increasing alkalinity across both OAE treatments (Si and Ca) while *Nitzschia* only displayed an increase with silicification in the Si-OAE treatment. Overall trends observed in the silicification of the diatom community were confirmed by increased concentrations of BSi in the water column and accumulated in the sediment trap of mesocosms in the Si-OAE treatment. The increase of seawater alkalinity by 0 – 600 $\mu\text{mol kg}^{-1}$ had no effect on the concentration of BSi in the water column, accumulated in the sediment or overall diatom community silicification. In conjunction with published OAE research, our findings highlight the need for research to cover a broad range of environmental conditions, approaches to OAE and marine communities.

4.1 Temporal dynamics of biogenic silica in the water column and sediment

We observed no significant relationship between alkalinity and BSi in the water column or accumulated in the sediments (Fig 6 a,b). In contrast, concentrations of BSi in the water column and accumulated in the sediments were significantly greater in the Si-OAE treatment before and after macronutrient additions. These trends support observations for community- and genera-specific silicification, with diatoms in the Si-OAE being more heavily silicified.

Notably, BSi accumulation in the sediments of the Si-based OAE treatment group was greatest in the highest alkalinity mesocosm ($\Delta 600 \mu\text{mol kg}^{-1}$) and lowest in the control mesocosm ($\Delta 0 \mu\text{mol kg}^{-1}$) during phase I (Fig. 6b). This difference emerged immediately after the addition of Na_2SiO_3 to the Si-based OAE treatment. However, no build-up of chlorophyll *a* or significant build-up of BSi in the water column was observed in the days prior to the emergence of this trend, suggesting the sedimented BSi is an artefact of the Na_2SiO_3 addition to the silicate-based treatments. Our interpretation of these findings are that (1) there were residual precipitates in the Na_2SiO_3 solution added to the mesocosms resulting in increased BSi in the water column, and/or (2) that there was pH dependent, inorganic precipitation of amorphous silicate, with these precipitates sinking out into the sediment trap (Goto, 1956; Okamoto et al., 1957; Owen, 1975). The latter is supported by recent work conducted by Gately et al. (2023), whose abiotic experiments revealed a decrease in dissolved silicate as TA-alkalinity increased, with scanning electron microscopy revealing mineral precipitates formed in high alkalinity seawater to be primarily composed of silicon and oxygen. Precipitation of silica within mesocosms may have been supported by ionic interactions between magnesium (or trace metals e.g. iron, aluminium) and silicate, pressure and relatively low temperatures at depth (Ehlert et al., 2016; Goto, 1956; Spithaki et al., 2018).

4.2 Effect of enhanced silicate concentration on silicification rates

Ca- and Si-based OAE appeared to have a notable influence on silicification of the diatom community. However, this influence is attributed to the mineral treatment type, either silicate or calcium based, with a significant relationship between silicification and alkalinity observed for the *Pseudo-nitzschia* and *Nitzschia* (Si-OAE treatment only). Diatoms in the Si-OAE treatment incorporated considerably more silicate over the 24-hour incubation period resulting in increased silicification. This outcome was expected, especially considering the consistently low (likely limiting) concentrations of Si(OH)_4 observed in the Ca-OAE treatment throughout the majority of the experiment. Si(OH)_4 is the key nutrient in the construction of the silicate-based frustule of diatoms, with low concentrations often becoming a limiting factor for growth (Martin-Jézéquel et al., 2000). It has been shown that before silicate concentrations become growth limiting, diatoms first respond by thinning their frustules (McNair et al., 2018; Paasche, 1975). Whilst this phenomenon has been observed in several studies (McNair et al., 2018; Rocha et al., 2010; Shimada et al., 2009), our mechanistic understanding of how diatoms adjust their silicon quotas is unclear (Milligan et al., 2004). Interestingly, an initial thinning followed by subsequent thickening of diatom frustules has been observed at non-growth limiting silicate concentrations (McNair et al., 2018). This may be an adaptive trade-off by which diatoms respond to lower silicate concentrations by decreasing their silicon quotas in favour of maintaining similar growth rates; a strategy that allows cells to respond to dynamic changes in silicate concentrations while maintaining a similar population size. Such rapid responses have been observed in both culture and field based experiments with diatoms responding to increases in silicate within several hours while responses to nitrate additions, after prolonged nitrate stress, took over 30 hrs (McNair et al., 2018; Rocha et al., 2010). As such, it is possible that the low concentrations of Si(OH)_4 observed in the Ca-OAE treatment, resulted in diatoms prioritising growth over silicate incorporation leading to significantly less silicification when compared to the Si-OAE treatment. Detailed measurements of diatom growth (not assessed here) would be required to confirm this hypothesis. We recommend future experiments consider this and assess diatom growth alongside measures of silicification to enable the exploration of potential trade-offs between growth and silicification.

4.3 Effect of carbonate chemistry manipulations on diatom silicification

We detected no clear relationship between total alkalinity and silicification in this mesocosm study, with the exception of *Pseudo-nitzschia* which was more heavily silicified in higher alkalinity treatments. *Nitzschia* also displayed increases in silicification in higher alkalinity treatments, however this was only observed in the Si-OAE treatment. In addition, we observed a significant effect of total alkalinity on silicification during phase I of the experiment. This result was primarily due to the Si-OAE based treatment, ~~due to the~~ illustrated by the significant three-way interaction between total alkalinity, alkalinity source mineral and experimental phase in the linear mixed effects model. Previous physiological studies have found tight links between diatom silicification and components of the marine carbonate chemistry system (CO_2 and pH) (Gao et al., 2014; Hervé et al., 2012; Li et al., 2019; Petrou et al., 2019; Zepernick et al., 2021). However current research presents some inconsistencies in the relationship between silicification and carbonate chemistry. Petrou et al. (2019) found that silicification

510 decreased at increased pCO₂ and low pH expected as a result of ocean acidification. In contrast, research
conducted by Li et al. (2019) and Zepernick et al. (2021) found the opposite with silicification decreasing at
increasing pH/alkaline conditions. It is important to note that both Li et al. (2019) and Petrou et al. (2019)
simulated ocean acidification, increasing pCO₂ while total alkalinity remained constant. In contrast, Hervé et al.
(2012) and Zepernick et al. (2021) altered the carbonate chemistry of their respective media via additions of NaOH
515 and HCl thereby manipulating the concentrations of carbon species but not altering total DIC values. This allows
for a more direct comparison to be made with the results presented here as OAE in its unequilibrated form results
in changes in carbon species concentrations without significant differences in DIC. Hervé et al. (2012) found silica
incorporation rates to decrease from pH 6.4 to 8.2 and increase from 8.2 to 8.5. Visual inspection of the data
presented by Hervé et al. (2012) showed that the incorporation of silicate into a cell at pH 8.5 was marginally less
520 than that at lower pH values (no statistics were provided for this measurement in the cited article). In support of
this, Zepernick et al. (2021) found no significant difference in the silicification of freshwater diatoms at pH 7.7
and 8.6, only between pH 7.7 and 9.2 was a significant decline in silicification observed. In our study, total
alkalinity values corresponded to a pH range from approximately 8.0 to 8.75 (total scale). Thus, it is possible that
the enhancement of alkalinity in our study and corresponding changes in carbonate chemistry were not extreme
525 enough to result in a significant change in the silicification of the diatom community or specific genera.

The pennate *Pseudo-nitzschia* was the only genus to exhibit a significant increase in silicification with increasing
alkalinity irrespective of the mineral treatment type. In contrast, *Nitzschia* displayed a similar relationship between
silicification and alkalinity however this was only in the Si-OAE based treatment. This relationship is similar to
530 that observed by Petrou et al. (2019) who found several pennate species to exhibit a close relationship between
silicification and carbonate chemistry conditions. Such a finding suggests that changes to silicification as a result
of OAE is likely to be genus or species-specific supporting current knowledge surrounding the large variation in
silicification between species (Martin-Jézéquel et al., 2000; Rousseau et al., 2002; Timmermans et al., 2004).

535 To conclude this section, we would like to highlight that other factors, which were not controlled for in this
experiment, may have also contributed to the lack of a significant difference in silicification observed here.
Silicification is influenced by a range of environmental parameters, such as macronutrient concentrations
(primarily silicate) (Claquin et al., 2002; Shimada et al., 2009), light intensity (Su et al., 2018; Taylor, 1985), and
predation (Liu et al., 2016; Pondaven et al., 2007), which may have masked potential effects of total alkalinity on
540 silicification. It is possible that OAE effects on silicification may be identifiable in other experimental settings
with different boundary conditions and environmental controls. This is supported by the specific scenarios in
which silicification was significantly influenced by total alkalinity here (i.e. during phase I of the experiment in
the Si-OAE based treatment). Our previous OAE study assessing the influence of a ~500 µmol kg⁻¹ alkalinity
increase on coastal Tasmanian plankton communities found significant effects of OAE on silicate dynamics,
545 suggesting changes in diatom community silicification (Ferderer et al., 2022). These differences suggest that
boundary conditions are important and that many studies assessing the effects of OAE on diatom communities
will be needed to extract more robust response patterns across a range of conditions, consistent with conclusions
drawn from a synthesis on diatoms in the context of ocean acidification (Bach and Taucher, 2019).

550 **5 Implications for the implementation of OAE and outlook**

In conclusion, our study underlines that the use of silicate-based minerals for OAE ~~can~~ has the potential to significantly affect silicification of the diatom community and specific genera. This result was expected and consistent with our current understanding of dissolved silicate effects on diatom communities (Baines et al., 2010; Egge and Aksnes, 1992; Hauck et al., 2016; Tréguer et al., 2021). The significant influence of Si(OH)₄ on diatom silicification is not surprising, as it can be a limiting nutrient for diatoms, with previous studies having shown the benefits of increased Si(OH)₄ concentrations as a result of olivine-based mineral dissolution (Baines et al., 2010; Hutchins et al., 2023; Martin-Jézéquel et al., 2000; Wischmeyer et al., 2003). It is important to note that this experiment was designed to specifically assess the potential interactive effects of OAE and magnesium silicate fertilisation associated with silicate based minerals proposed for use in mineral based OAE. Such an interaction was explored as there are several potential minerals proposed for use in mineral based OAE all of which vary in their respective dissolution products. As such we did not control for or assess other ecologically important dissolution products found in silicate based minerals e.g. iron and nickel which have been shown to have significant ecological implications for plankton communities (Xin et al., 2023; Hutchins et al., 2023; Guo et al., 2022; Boyd et al., 2007). Further exploration of potential dissolution products and their effects on plankton communities in the presence and absence of changing carbonate chemistry conditions are encouraged in future research.

In contrast to the clear effects of dissolved silicate fertilisation on silicification, our findings provide ~~we found~~ limited evidence to suggest that the enhancement of seawater alkalinity by 0 – 600 μmol kg⁻¹ affects genera specific silicification. The lack of a clear alkalinity effect on silicification was unexpected, especially in the higher alkalinity treatments, which corresponded to a substantial change in carbonate chemistry conditions. It is important to note that increases in alkalinity above Δ400 μmol kg⁻¹ are considered to be relatively extreme levels of OAE, yet only two genera, *Pseudo-nitzschia* and *Nitzschia*, exhibited significant changes in silicification as a result of the gradient in carbonate chemistry employed here (Schulz et al., 2023). Real world applications are predicted to employ significantly less extreme perturbations of the marine carbonate chemistry system (apart from sites of direct alkalinity addition). As such, one might hypothesize that impacts may be less significant. Furthermore, such perturbations would be relatively short lived in real world applications since dilution of the perturbed water bodies occurs, unlike the sustained changes observed within mesocosms presented here (He and Tyka, 2023; Wang et al., 2023). Irrespective of this, there is substantial empirical evidence suggesting that changes in carbonate chemistry through ocean acidification will influence diatom communities, their growth, various aspects of silicification (e.g. rate, degree of) and subsequently silicate cycling in the ocean (Bach and Taucher, 2019; Gao et al., 2014; Li et al., 2019; Milligan et al., 2004; Petrou et al., 2019; Taucher et al., 2022). Additionally, our previous work has shown that an increase in total alkalinity of ~500 μmol kg⁻¹ has a significant influence on the uptake of dissolved silicate and production of BSi in a coastal phytoplankton community (Ferderer et al., 2022). The mixed outcomes observed here and in the limited OAE studies so far suggest that the responses of diatoms will differ and be dependent on the community and environmental boundary conditions. More community studies, ideally with closely aligned experimental setups, will be needed to discern whether the response of diatoms to OAE forms any robust patterns. Such ecological observations subsequently need mechanistic underpinning, potentially achievable through the intelligent design of physiological experiments (Collins et al.,

590 2022). Ultimately, the goal should be to provide predictive understanding of the role of diatoms and eventually all major functional plankton groups under the differing strategies of OAE.

Data availability

Data are available from the Institute for Marine and Antarctic Studies (IMAS) data catalogue, University of
595 Tasmania (UTAS) <https://doi.org/10.25959/G3FN-HE45>.

Author contributions

UR designed the mesocosm experiment and AF designed the experiment assessing silicification rates with input
from LTB and KGB. AF was responsible for the investigation, data curation, formal analysis, and writing. AF
wrote the manuscript with contributions from LTB, KGS, UR, KGB and ZC.

600 ***Competing interests***

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

We would like to thank all participants of the campaign who assisted in sampling and general scientific discussion
throughout. We would also like to acknowledge Juliane Tammen and Peter Fritzsche for the measurement of
605 dissolved inorganic nutrients, Leila Kittu, Anna Groen, Lucas Krause, Jule Ploschke and Kira Lange for their
measurements of particulate matter, Jana Meyer for the measurement of sedimented material, Julieta Schneider
for the measurement of carbonate chemistry, Andrea Ludwig for her organisation and management of the project
and staff members at the research station for their assistance in daily tasks.

Financial support

610 This study was funded by the OceanNETS project (“Ocean-based Negative Emissions Technologies – analysing
the feasibility, risks and co-benefits of ocean-based negative emission technologies for stabilizing the climate”,
EU Horizon 2020 Research and Innovation Programme Grant Agreement No.: 869357), and the Helmholtz
European Partnering project Ocean-CDR (“Ocean-based carbon dioxide removal strategies”, Project No.: PIE-
0021) with additional support from the AQUACOSM-plus project (EU H2020-INFRAIA Project No.: 871081,
615 “AQUACOSM-plus: Network of Leading European AQUATIC MesoCOSM Facilities Connecting Rivers, Lakes,
Estuaries and Oceans in Europe and beyond”). Additional funding was supplied by a Future Fellowship
(FT200100846) awarded to LTB by the Australian Research Council. This research was also conducted while
Aaron Ferderer was in receipt of an Australian Government Research Training Program (RTP) scholarship.

References

620 Bach, L. T. and Taucher, J.: CO₂ effects on diatoms: A synthesis of more than a decade of ocean acidification
experiments with natural communities, *Ocean Science*, 15, 1159–1175, [https://doi.org/10.5194/OS-15-1159-](https://doi.org/10.5194/OS-15-1159-2019)
2019, 2019.

Baines, S. B., Twining, B. S., Brzezinski, M. A., Nelson, D. M., and Fisher, N. S.: Causes and biogeochemical
implications of regional differences in silicification of marine diatoms, *Global Biogeochemical Cycles*, 24,
625 <https://doi.org/10.1029/2010GB003856>, 2010.

Boxhammer, T., Bach, L. T., Czerny, J., and Riebesell, U.: Technical note: Sampling and processing of
mesocosm sediment trap material for quantitative biogeochemical analysis, *Biogeosciences*, 13, 2849–2858,
<https://doi.org/10.5194/bg-13-2849-2016>, 2016.

- 630 Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale iron enrichment experiments 1993-2005: synthesis and future directions, *Science*, 315, 612–617, <https://doi.org/10.1126/science.1131669>, 2007.
- 635 Chen, C. Y. and Durbin, E. G.: Effects of pH on the growth and carbon uptake of marine phytoplankton, *Marine Ecology Progress Series*, 109, 83–94, <https://doi.org/10.3354/meps109083>, 1994.
- Claquin, P., Martin-Jézéquel, V., Kromkamp, J. C., Veldhuis, M. J. W., and Kraay, G. W.: Uncoupling of Silicon Compared with Carbon and Nitrogen Metabolisms and the Role of the Cell Cycle in Continuous Cultures of *Thalassiosira Pseudonana* (bacillariophyceae) Under Light, Nitrogen, and Phosphorus Control, *Journal of Phycology*, 38, 922–930, <https://doi.org/10.1046/j.1529-8817.2002.t01-1-01220.x>, 2002.
- 640 Collins, S., Whittaker, H., and Thomas, M. K.: The need for unrealistic experiments in global change biology, *Current Opinion in Microbiology*, 68, 102151, <https://doi.org/10.1016/J.MIB.2022.102151>, 2022.
- Czerny, J., Schulz, K. G., Krug, S. A., Ludwig, A., and Riebesell, U.: Technical Note: The determination of enclosed water volume in large flexible-wall mesocosms “KOSMOS,” *Biogeosciences*, 10, 1937–1941, <https://doi.org/10.5194/bg-10-1937-2013>, 2013.
- 645 Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO₂ measurements., *PICES Special Publication 3*, 0–199 pp., 2007.
- Egge, J. and Aksnes, D.: Silicate as regulating nutrient in phytoplankton competition, *Mar. Ecol. Prog. Ser.*, 83, 281–289, <https://doi.org/10.3354/meps083281>, 1992.
- 650 Ehlert, C., Reckhardt, A., Greskowiak, J., Liguori, B. T. P., Böning, P., Paffrath, R., Brumsack, H.-J., and Pahnke, K.: Transformation of silicon in a sandy beach ecosystem: Insights from stable silicon isotopes from fresh and saline groundwaters, *Chemical Geology*, 440, 207–218, <https://doi.org/10.1016/j.chemgeo.2016.07.015>, 2016.
- 655 Escaravage, V. and Prins, T. C.: Silicate availability, vertical mixing and grazing control of phytoplankton blooms in mesocosms, in: *Sustainable Increase of Marine Harvesting: Fundamental Mechanisms and New Concepts*, edited by: Vadstein, O. and Olsen, Y., Springer Netherlands, Dordrecht, 33–48, https://doi.org/10.1007/978-94-017-3190-4_4, 2002.
- Ferderer, A., Chase, Z., Kennedy, F., Schulz, K. G., and Bach, L. T.: Assessing the influence of ocean alkalinity enhancement on a coastal phytoplankton community, *Biogeosciences*, 19, 5375–5399, <https://doi.org/10.5194/BG-19-5375-2022>, 2022.
- 660 Gao, K., Campbell, D. A., Gao, K., and Campbell, D. A.: Photophysiological responses of marine diatoms to elevated CO₂ and decreased pH: a review, *Functional Plant Biol.*, 41, 449–459, <https://doi.org/10.1071/FP13247>, 2014.
- 665 Gately, J. A., Kim, S. M., Jin, B., Brzezinski, M. A., and Iglesias-Rodriguez, M. D.: Coccolithophores and diatoms resilient to ocean alkalinity enhancement: A glimpse of hope?, *Science Advances*, 9, eadg6066, <https://doi.org/10.1126/sciadv.adg6066>, 2023.
- Goto, K.: Effect of pH on Polymerization of Silicic Acid, *J. Phys. Chem.*, 60, 1007–1008, <https://doi.org/10.1021/j150541a046>, 1956.
- 670 Guo, J. A., Strzepek, R., Willis, A., Ferderer, A., and Bach, L. T.: Investigating the effect of nickel concentration on phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement, *Biogeosciences*, 19, 3683–3697, <https://doi.org/10.5194/BG-19-3683-2022>, 2022.
- Hansen, H. P. and Koroleff, F.: Determination of nutrients, in: *Methods of Seawater Analysis*, John Wiley & Sons, Ltd, 159–228, <https://doi.org/10.1002/9783527613984.ch10>, 1999.

- Hauck, J., Köhler, P., Wolf-Gladrow, D., and Völker, C.: Iron fertilisation and century-scale effects of open ocean dissolution of olivine in a simulated CO₂ removal experiment, *Environ. Res. Lett.*, 11, 024007, <https://doi.org/10.1088/1748-9326/11/2/024007>, 2016.
- 675
- He, J. and Tyka, M. D.: Limits and CO₂ equilibration of near-coast alkalinity enhancement, *Biogeosciences Discussion*, <https://doi.org/10.5194/egusphere-2022-683>, 2023.
- Hervé, V., Derr, J., Douady, S., Quinet, M., Moisan, L., and Lopez, P. J.: Multiparametric Analyses Reveal the pH-Dependence of Silicon Biomineralization in Diatoms, *PLOS ONE*, 7, e46722, <https://doi.org/10.1371/JOURNAL.PONE.0046722>, 2012.
- 680
- Hinga, K. R.: Effects of pH on coastal marine phytoplankton, *Marine Ecology Progress Series*, 238, 281–300, <https://doi.org/10.3354/meps238281>, 2002.
- Hutchins, D. A., Fu, F.-X., Yang, S.-C., John, S. G., Romaniello, S. J., Andrews, M. G., and Walworth, N. G.: Responses of keystone phytoplankton groups to olivine dissolution products and implications for carbon dioxide removal via ocean alkalinity enhancement, *bioRxiv*, 2023.04.08.536121, <https://doi.org/10.1101/2023.04.08.536121>, 2023.
- 685
- Leblanc, K. and Hutchins, D. A.: New applications of a biogenic silica deposition fluorophore in the study of oceanic diatoms, *Limnology and Oceanography: Methods*, 3, 462–476, <https://doi.org/10.4319/LOM.2005.3.462>, 2005.
- 690
- Lenth, R. V., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., and Singmann, H.: emmeans: Estimated Marginal Means, aka Least-Squares Means, 2023.
- Li, F., Fan, J., Hu, L., Beardall, J., Xu, J., and Fields, D.: Physiological and biochemical responses of *Thalassiosira weissflogii* (diatom) to seawater acidification and alkalization, *ICES Journal of Marine Science*, 76, 1850–1859, <https://doi.org/10.1093/ICESJMS/FSZ028>, 2019.
- 695
- Liu, H., Chen, M., Zhu, F., and Harrison, P. J.: Effect of Diatom Silica Content on Copepod Grazing, Growth and Reproduction, *Frontiers in Marine Science*, 3, 2016.
- Martin-Jézéquel, V., Hildebrand, M., and Brzezinski, M. A.: Silicon metabolism in diatoms: Implications for growth, *Journal of Phycology*, 36, 821–840, <https://doi.org/10.1046/J.1529-8817.2000.00019.X>, 2000.
- McNair, H. M., Brzezinski, M. A., and Krause, J. W.: Quantifying diatom silicification with the fluorescent dye, PDMPO, *Limnology and Oceanography: Methods*, 13, 587–599, <https://doi.org/10.1002/LOM3.10049>, 2015.
- 700
- McNair, H. M., Brzezinski, M. A., and Krause, J. W.: Diatom populations in an upwelling environment decrease silica content to avoid growth limitation, *Environmental Microbiology*, 20, 4184–4193, <https://doi.org/10.1111/1462-2920.14431>, 2018.
- Milligan, A. J., Varela, D. E., Brzezinski, M. A., and Morel, F. M. M.: Dynamics of silicon metabolism and silicon isotopic discrimination in a marine diatom as a function of pCO₂, *Limnology and Oceanography*, 49, 322–329, <https://doi.org/10.4319/LO.2004.49.2.0322>, 2004.
- 705
- Okamoto, G., Okura, T., and Goto, K.: Properties of silica in water, *Geochimica et Cosmochimica Acta*, 12, 123–132, [https://doi.org/10.1016/0016-7037\(57\)90023-6](https://doi.org/10.1016/0016-7037(57)90023-6), 1957.
- Owen, L.: Precipitation of amorphous silica from high-temperature hypersaline geothermal brines, California Univ., Livermore (USA). Lawrence Livermore Lab., 1975.
- 710
- Paasche, E.: Growth of the plankton diatom *Thalassiosira nordenskioeldii* Cleve at low silicate concentrations, *Journal of Experimental Marine Biology and Ecology*, 18, 173–183, [https://doi.org/10.1016/0022-0981\(75\)90072-6](https://doi.org/10.1016/0022-0981(75)90072-6), 1975.
- Paul, A. J. and Bach, L. T.: Universal response pattern of phytoplankton growth rates to increasing CO₂, *New Phytologist*, 228, 1710–1716, <https://doi.org/10.1111/NPH.16806>, 2020.
- 715

- Petrou, K., Baker, K. G., Nielsen, D. A., Hancock, A. M., Schulz, K. G., and Davidson, A. T.: Acidification diminishes diatom silica production in the Southern Ocean, *Nature Climate Change*, 9, 781–786, <https://doi.org/10.1038/S41558-019-0557-Y>, 2019.
- 720 Pondaven, P., Gallinari, M., Chollet, S., Bucciarelli, E., Sarthou, G., Schultes, S., and Jean, F.: Grazing-induced Changes in Cell Wall Silicification in a Marine Diatom, *Protist*, 158, 21–28, <https://doi.org/10.1016/J.PROTIS.2006.09.002>, 2007.
- Posit Team (2023): RStudio: Integrated Development for R, 2023.
- Riebesell, U., Wolf-Gladrow, D. A., and Smetacek, V.: Carbon dioxide limitation of marine phytoplankton growth rates, *Nature*, 361, 249–251, <https://doi.org/10.1038/361249a0>, 1993.
- 725 Riebesell, U., Czerny, J., von Bröckel, K., Boxhammer, T., Büdenbender, J., Deckelnick, M., Fischer, M., Hoffmann, D., Krug, S. A., Lentz, U., Ludwig, A., Muche, R., and Schulz, K. G.: Technical Note: A mobile sea-going mesocosm system – new opportunities for ocean change research, *Biogeosciences*, 10, 1835–1847, <https://doi.org/10.5194/bg-10-1835-2013>, 2013.
- 730 Rocha, C. L. D. L., Terbrüggen, A., Völker, C., and Hohn, S.: Response to and recovery from nitrogen and silicon starvation in *Thalassiosira weissflogii*: growth rates, nutrient uptake and C, Si and N content per cell, *Marine Ecology Progress Series*, 412, 57–68, <https://doi.org/10.3354/meps08701>, 2010.
- Rousseau, V., Leynaert, A., Daoud, N., and Lancelot, C.: Diatom succession, silicification and silicic acid availability in Belgian coastal waters (Southern North Sea), *Marine Ecology Progress Series*, 236, 61–73, <https://doi.org/10.3354/MEPS236061>, 2002.
- 735 Schulz, K. G., Bach, L. T., and Dickson, A. G.: Seawater carbonate chemistry considerations for ocean alkalinity enhancement research: theory, measurements, and calculations, *State of the Planet*, 2-oae2023, 1–14, <https://doi.org/10.5194/sp-2-oae2023-2-2023>, 2023.
- Shimada, C., Nakamachi, M., Tanaka, Y., Yamasaki, M., and Kuwata, A.: Effects of nutrients on diatom skeletal silicification: Evidence from *Neodenticula seminae* culture experiments and morphometric analysis, *Marine Micropaleontology*, 73, 164–177, <https://doi.org/10.1016/J.MARMICRO.2009.09.001>, 2009.
- 740 Spinhaki, A., Petratos, G., Matheis, J., Hater, W., and Demadis, K. D.: The precipitation of “magnesium silicate” under geothermal stresses. Formation and characterization, *Geothermics*, 74, 172–180, <https://doi.org/10.1016/j.geothermics.2018.03.001>, 2018.
- 745 Su, Y., Lundholm, N., and Ellegaard, M.: The effect of different light regimes on diatom frustule silicon concentration, *Algal Research*, 29, 36–40, <https://doi.org/10.1016/j.algal.2017.11.014>, 2018.
- Taucher, J., Bach, L. T., Prowe, A. E. F., Boxhammer, T., Kvale, K., and Riebesell, U.: Enhanced silica export in a future ocean triggers global diatom decline, *Nature* 2022 605:7911, 605, 696–700, <https://doi.org/10.1038/s41586-022-04687-0>, 2022.
- 750 Taylor, N. J.: Silica incorporation in the diatom *Cosinodiscus granii* as affected by light intensity, *British Phycological Journal*, 20, 365–374, <https://doi.org/10.1080/00071618500650371>, 1985.
- Timmermans, K. R., Van Der Wagt, B., and De Baar, H. J. W.: Growth rates, half-saturation constants, and silicate, nitrate, and phosphate depletion in relation to iron availability of four large, open-ocean diatoms from the Southern Ocean, *Limnology and Oceanography*, 49, 2141–2151, <https://doi.org/10.4319/LO.2004.49.6.2141>, 2004.
- 755 Tréguer, P. J., Sutton, J. N., Brzezinski, M., Charette, M. A., Devries, T., Dutkiewicz, S., Ehlert, C., Hawkings, J., Leynaert, A., Liu, S. M., Monferrer, N. L., López-Acosta, M., Maldonado, M., Rahman, S., Ran, L., and Rouxel, O.: Reviews and syntheses: The biogeochemical cycle of silicon in the modern ocean, *Biogeosciences*, 18, 1269–1289, <https://doi.org/10.5194/BG-18-1269-2021>, 2021.

- 760 Wang, F., Lu, Z., Wang, Y., Yan, R., and Chen, N.: Porewater exchange drives the dissolved silicate export across the wetland-estuarine continuum, *Frontiers in Marine Science*, 10, 812, <https://doi.org/10.3389/FMARS.2023.1206776>, 2023.
- Welschmeyer, N. A.: Fluorometric analysis of chlorophyll a in the presence of chlorophyll b and pheopigments, *Limnology and Oceanography*, 39, 1985–1992, <https://doi.org/10.4319/lo.1994.39.8.1985>, 1994.
- 765 Wischmeyer, A. G., Del Amo, Y., Brzezinski, M., and Wolf-Gladrow, D. A.: Theoretical constraints on the uptake of silicic acid species by marine diatoms, *Marine Chemistry*, 82, 13–29, [https://doi.org/10.1016/S0304-4203\(03\)00033-1](https://doi.org/10.1016/S0304-4203(03)00033-1), 2003.
- Xin, X., Faucher, G., and Riebesell, U.: Phytoplankton Response to Increased Nickel in the Context of Ocean Alkalinity Enhancement, *Biogeosciences Discussions*, 1–15, <https://doi.org/10.5194/bg-2023-130>, 2023.
- 770 Zepernick, B. N., Gann, E. R., Martin, R. M., Pound, H. L., Krausfeldt, L. E., Chaffin, J. D., and Wilhelm, S. W.: Elevated pH Conditions Associated With *Microcystis* spp. Blooms Decrease Viability of the Cultured Diatom *Fragilaria crotonensis* and Natural Diatoms in Lake Erie, *Frontiers in Microbiology*, 12, 2021.
- Znachor, P. and Nedoma, J.: Application of the pdmpo technique in studying silica deposition in natural populations of *fragilaria crotonensis* (bacillariophyceae) at different depths in a eutrophic reservoir1, *Journal of Phycology*, 44, 518–525, <https://doi.org/10.1111/J.1529-8817.2008.00470.X>, 2008.

775