



Continental scientific drilling and microbiology: (extremely) low biomass in crystalline bedrock of central Sweden

George Westmeijer^{1*}, Cristina Escudero^{2,3}, Claudia Bergin⁴, Stephanie Turner¹, Magnus Stähle¹, Maliheh Mehrshad⁵, Prune Leroy⁵, Moritz Buck⁵, Pilar López-Hernández^{5,6}, Jens Kallmeyer⁷, Ricardo Amils²,
5 Stefan Bertilsson⁵ & Mark Dopson¹

¹Centre for Ecology and Evolution in Microbial model Systems (EEMiS), Linnaeus University, Stuvaregatan 4, 39231 Kalmar, Sweden

²Centro de Biología Molecular Severo Ochoa (CSIC-UAM), Universidad Autónoma de Madrid, Cantoblanco, Madrid, Spain

³Geomicrobiology Group, Center for Applied Geoscience, University of Tuebingen, Tuebingen, Germany

10 ⁴Microbial Single Cell Genomics Facility, Science for Life Laboratory, Uppsala, Sweden

⁵Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, P.O. box 7050, 75005 Uppsala, Sweden

⁶Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Sweden

⁷GFZ German Research Centre for Geosciences, Section Geomicrobiology, Potsdam, Germany

15 *Correspondence to:* George Westmeijer (george.westmeijer@lnu.se)

Classification: Microbial ecology

Keywords: COSC-2, deep biosphere, CARD-FISH, whole genome amplification



Abstract

Scientific drilling expeditions offer a unique opportunity to characterize the microbial communities in the subsurface that have been long-term isolated from the surface. With subsurface microbial biomass being low in general, biological contamination from the drilling fluid, sample processing, or molecular work is a major concern. To address this, characterization of the contaminant populations in the drilling fluid and negative extraction controls are essential for assessing and evaluating such sequencing data. Here, crystalline rock cores down to 2250 m depth, groundwater-bearing fractures, and the drilling fluid were sampled for DNA to characterize the microbial communities using a broad genomic approach. However, even after removing potential contaminant populations present in the drilling fluid, notorious contaminants were abundant and mainly affiliated with the bacterial order Burkholderiales. These contaminant microorganisms likely originated from the reagents used for isolating and amplifying DNA despite stringent quality standards during the molecular work. The detection of strictly anaerobic sulfate reducers such as *Candidatus Desulforudis audaxviator* suggested the presence of autochthonous deep biosphere taxa in the sequenced libraries, yet these clades represented only a minor fraction of the sequence counts (< 0.1 %), hindering further ecological interpretations. The described methods and findings emphasize the importance of sequencing extraction controls and can support experimental design for future microbiological studies in conjunction with continental drilling operations.



1 Introduction

40 The concept of the deep biosphere was postulated by Thomas Gold (Gold, 1992) as life in the oceanic and continental subsurface plus sub-glacial lakes, with the continental deep biosphere being estimated to consist of 2 to 6×10^{29} cells (Magnabosco et al., 2018). These cells are active and greatly influence surface life through biogeochemical cycling (Kieft, 2016; Amils et al., 2023) and include archaeal, bacterial, and eukaryotic populations (Borgonie et al., 2015; Bell et al., 2022, 2020) as well as viruses (Rahlff et al., 2021; Holmfeldt et al., 2021; Engelhardt et al., 2014). Together with infiltration from
45 the photosynthesis-driven surface (Osterholz et al., 2022), geogenic hydrogen and carbon dioxide from the deep subsurface are the main energy sources for subsurface life (Pedersen, 2000). Research on the subsurface has been enabled and accelerated by the use of springs (Suzuki et al., 2014), artesian wells (Chapelle et al., 2002), boreholes (Bomberg et al., 2016), mines (Magnabosco et al., 2016), and dedicated underground laboratories (Boylan et al., 2019; Wu et al., 2016). Genomic approaches have greatly increased the understanding of microbial diversity in the continental subsurface and revealed a large proportion
50 of diversity without isolated representatives for which the ecology is not fully understood (Hug et al., 2016).

Subsurface microorganisms inhabit groundwater-filled fractures (Wu et al., 2017), are attached to minerals on the rock surface (Casar et al., 2020), and reside in the pore space of rocks as endoliths (Escudero et al., 2018). The pore space in rocks provides these endolithic communities a surface to colonize, mineral nutrients, and moisture (Golubic et al., 1981). Studies on endolithic communities emphasize the importance of depth (Magnabosco et al., 2018), availability of water and electron donors (Dai et al., 2021), and pore space (Dutta et al., 2018) on microbial biomass and activity. Many of these organisms are chemolithoautotrophs and obtain energy through the oxidation of inorganic compounds, such as sulfide, iron(II), and hydrogen (Lever et al., 2015b). The majority of lithotrophs capture carbon dioxide through the energy-demanding Calvin cycle and this, combined with the often energy- and nutrient-limiting conditions in the subsurface, limits microbial growth (Ghosh and Dam, 2009). Examples of these organisms previously identified in the subsurface are *Thiobacillus denitrificans* that combines the
55 oxidation of sulfides with the reduction of nitrate (Beller et al., 2006; Lopez-Fernandez et al., 2023) or *Acidovorax* sp. that couples the oxidation of ferrous iron to the reduction of nitrate (Pantke et al., 2012; Escudero et al., 2020). Studying the diversity and abundance of these endolithic communities would benefit the understanding of the microbial ecology and biomass in the continental subsurface.

In general, biomass in the continental subsurface is low and hence, contamination during field sampling or sample processing
65 is a major concern, especially when drill cores are employed to retrieve the samples, as allochthonous populations in the drilling fluid are introduced into the subsurface (Smith et al., 2000). With microbial biomass usually being several orders of magnitude lower in the subsurface compared to surface environments (Starnawski et al., 2017), even the infiltration of a small volume of drilling fluid would risk the sample being compromised and unusable for studying microbial diversity. Consequently, contaminant populations introduced along with the drilling fluid need to be identified and the magnitude of any
70 such contamination needs to be assessed prior to downstream ecological interpretations. These contamination control methods often involve the introduction of tracers in the drilling fluid such as fluorescent microspheres or perfluorocarbon tracers that




can be used to assess drilling fluid infiltration into the rock core (Kallmeyer, 2017). Removal of such tracers from the drill core surface through flaming allows one to minimize contamination of the inner core (Lever et al., 2006). Tracer count of the core surface can be combined with cell counts of the drilling fluid to assess microbial contamination originating from the drilling fluid. Furthermore, during molecular analyses, the typically low biomass in the deep continental biosphere increases the risk of detecting notorious reagent contaminants that would cause erroneous conclusions on the nature of the microbial community (Sheik et al., 2018; Salter et al., 2014). Despite the concerns regarding contamination, studies on this vast biome are necessary for understanding the role of the deep subsurface microbiome in carbon and nutrient cycling. However, due to the difficulty of obtaining samples, few investigations of the continental subsurface have been performed and the available data are patchy at best.

Integration of subsurface microbiology into the workings of the International Continental Scientific Drilling Program (ICDP) defined several questions (Mangelsdorf and Kallmeyer, 2010) including understanding subsurface biodiversity; interactions between the deep biosphere and geosphere; and deep life as an analogue for potential life on other planets. Following this workshop, two subsurface scientific drilling projects termed Collisional Orogeny in the Scandinavian Caledonides (COSC-1 and COSC-2), located in western central Sweden, were performed in 2014 and 2020 (Lorenz et al., 2015, 2022). Here, microorganisms present in a hard rock environment were studied down to 2250 m depth by sampling drill cores and groundwater-bearing fractures during the COSC-2 drilling project. The efficacy of the contamination control measures was evaluated by characterizing the microorganisms in the samples of interest combined with those present in the drilling fluid.


2. Results and discussion

2.1 Rock core samples

The aim of this study was to characterize the microbial community in the pore space of subsurface rocks ($n = 50$) and natural fractures ($n = 29$), covering various rock types (Fig. 1). The drilling fluid was sampled ($n = 50 + 29$) simultaneous with the rocks and natural fractures (see section 2.2) to assess the efficiency of the decontamination methods. The drilling fluid contained on average 335 ± 396 cells mL^{-1} (mean \pm sd, $n = 44$). Based upon quenching of the fluorescent beads during sample decontamination (Fig. 2), on average 98.9 ± 3.3 % of these cells were estimated to be destroyed by flame sterilization of the core surface and for 31/44 cores over 99.9 % of the cells were destroyed.



100 **Figure 1 Location and lithology sampling site.** a) The COSC-2 borehole within Sweden (Lat 63.3124° N, Lon 13.5265° E). b) Temperature profile and lithology according to depth. Drill cores were sampled at 50 m intervals from 100 m to 2250 depth ($n = 50$), covering the various lithologies. Temperature increased linearly with depth, ranging from 6.1 °C at the top to 52.3 °C at the bottom of the borehole. Data from Lorenz et al. (2021). Dashed lines in the lithology represents sampled natural fractures ($n = 29$).



105 **Figure 2 Drilling fluid and contamination control.** a) Cell numbers in the drilling fluid over depth, quantified using fluorescence microscopy. b) Quenching of the fluorescent beads using flame sterilization of the core surface. The number of beads was determined before and after surface sterilization using fluorescence microscopy and the higher the fraction of beads quenched, the more successful the sterilization procedure. At 750 m depth, an alternative method of adding the beads was used by immersing the core in a bead bath instead of adding the beads to the drilling fluid due to adhesion of the beads to clay particles. c) The estimated number of remaining beads after sterilization by multiplying the cell number in the drilling fluid (a) with the fraction of unquenched beads (1-b).



110

Despite testing six DNA extraction protocols adapted to low biomass rock samples, none of the extracts from the cores contained detectable DNA ($< 0.05 \text{ ng } \mu\text{l}^{-1}$). This suggested surface sterilization of the rock cores, to remove contaminants from the drilling fluid, was successful as the majority of the cells from the drilling fluid on the core surface were destroyed (Fig. 2) and DNA could be isolated (in detectable quantities) from the drilling fluid itself. Amplification of potentially extracted DNA, targeting different regions of the 16S ribosomal RNA gene using three primer pairs, also did not yield any products. Furthermore, whole genome amplification confirmed the extremely low concentration of DNA as the qPCR of the amplified product revealed that based on similar quantification cycles (Cq), the samples of interest and the negative extraction controls were indistinguishable. Whole genome amplification has previously been shown to amplify DNA from a single cell (Braslavsky et al., 2003) and has been successfully employed on crystalline rock samples (Puente-Sánchez et al., 2018). As previous studies have described microbes to persist within deep crystalline bedrock (Puente-Sánchez et al., 2014, 2018) as well as crystalline bedrock of the Fennoscandian Shield (Purkamo et al., 2020), the lack of microbial cells in the rock samples scrutinized in this study underlines the influence of local lithology on microbial abundance with pore space, the presence of any electron donor, and the availability of water as potential controlling factors. The cause for the observed low biomass was likely a combination of these factors, especially given the variable lithology (Fig. 1). However, the effect of the lithology on the biomass was not feasible to predict prior to field sampling as the expected lithology from the siting differed from the drilled lithology (Lorenz et al., 2022).

115

120

130

135

In addition to DNA extraction and amplification, cell numbers were determined using fluorescence microscopy. DNA staining was observed in the shallowest rock samples (100-150 m depth) and in the fault zone around 1300 m depth (Table 2). However, the presence of microorganisms could not be confirmed by fluorescence *in situ* hybridization with catalyzed reporter deposition (CARD-FISH) as the corresponding fluorescent signals were deemed as unspecific. The lack of a reliable signal could be due to (i) the extremely low abundance of microbial populations; (ii) a low number of ribosomes present in the cells, although this is unlikely given the signal amplification capacity of the CARD method; (iii) insufficient membrane permeabilization despite the use of standard permeabilization methods for gram-positive bacteria and archaea; or (iv) the presence of microbial populations that fail to hybridize with the general bacterial (EUB338 I-III) and archaeal (ARC915) probes even though these probes have been reported to hybridize with a high proportion of microbial diversity (Moter and Göbel, 2000). Given the low detection limit of the method (able to detect single cells; Hoshino et al. 2008) and its widespread application in low-biomass environments (Escudero et al., 2018; Teske, 2005), an extremely low microbial biomass was the most probable cause for the lack of a fluorescent signal.

140

Table 2 Cell counts on selected drill cores. Microbial cells were quantified using fluorescence microscopy combined with the DNA-binding dyes DAPI and Syto-9. However, the presence of microorganisms could not be confirmed by CARD-FISH (described in results).

Depth (m)	Lithology	Cell count (g^{-1})
-----------	-----------	--------------------------------



104		8.6×10^3
112		2.9×10^3
150		3.0×10^3
200	Turbiditic greywackes	0
245		0
299		0
350		0
374		0
800	Alum shale	0
1,298	Ignimbrites + volcanic porphyries	2.1×10^5
1,450		1.7×10^4
1,697	Dolerite	0
1,800		0
2,200	Ignimbrites + volcanic porphyries	0

2.2 Fracture samples

145 DNA was isolated from the groundwater-bearing fractures ($n = 7$) occurring in a fault zone between 1,200 and 1,700 m depth and their respective drilling fluid samples ($n = 6$). Sequencing of the 16S ribosomal RNA gene yielded 2,758 unique amplicon sequence variants (ASVs). Bacterial 16S rRNA copy numbers were in the range of 10^3 - 10^5 copies g^{-1} fracture sample while archaea were hardly detected with copy numbers close to the lower detection limit of the qPCR (Table 1).

150 **Table 1 Details of fracture samples.** Bacterial and archaeal abundances were quantified using qPCR combined with bacterial and archaeal primers targeting the 16S ribosomal RNA gene, respectively. The gene copies are presented as average g^{-1} fracture sample \pm standard deviation ($n = 6$). The fractures at 1,268 and 1,270 m depth shared a drilling fluid sample as these fractures were part of the same drill core. *Number of ASVs after removing the ASVs present in the drilling fluid.

Sampling date	Depth (m)	DNA fracture sample (ng)	DNA drilling fluid (ng)	Bacterial gene copies \pm sd (g^{-1})	Archaeal gene copies \pm sd (g^{-1})	No. ASVs total	No. ASVs drilling fluid	No. ASVs fracture*
29 May '20	1,268	< 0.05	14.3	$2.5 \times 10^4 \pm 102$	91 ± 1	507	351	424
29 May '20	1,270	< 0.05	6.6	$1.2 \times 10^4 \pm 91$	96 ± 1	234	351	133
31 May '20	1,311	< 0.05	< 0.05	$2.9 \times 10^5 \pm 614$	97 ± 1	447	194	342
15 June '20	1,537	< 0.05	< 0.05	$3.5 \times 10^3 \pm 38$	79 ± 0	25	161	22
28 June '20	1,605	< 0.05	11.4	$2.7 \times 10^3 \pm 19$	112 ± 3	324	650	197
28 June '20	1,608	< 0.05	54	$2.5 \times 10^3 \pm 48$	93 ± 2	98	176	65
2 July '20	1,669	< 0.05	15.4	$2.2 \times 10^4 \pm 63$	100 ± 3	105	1,025	101



155 The difference in composition between the microbial communities in the drilling fluid compared to the fractures was small but significant (permutational MANOVA, $R^2 = 0.17$, $p = 0.01$). Removal of ASVs detected in the negative extraction controls ($n = 78$) and the drilling fluid ($n = 1,724$) from the fracture samples reduced the sequence count in the latter by 52 % and 64 %, respectively (Fig. 3). 1,304 ASVs were exclusively detected in the fracture samples, representing 11 % of the sequence counts, and after removing the contaminant populations from the fracture samples for downstream analysis, the difference between the latter and the drilling fluid increased ($R^2 = 0.32$, $p = 0.002$). However, the composition of the microbial community on the
160 fracture surfaces (Fig. 3) strongly suggested contamination due to a high abundance of notorious reagent contaminants such as *Ralstonia*, *Sphingomonas*, *Agrobacterium*, and *Corynebacterium* (Salter et al., 2014; Sheik et al., 2018). Apart from *Corynebacterium*, these genera were also detected in the negative extraction control and together these four genera comprise 46 % of all the sequence counts. Some taxa were more credible autochthonous populations as they are strict anaerobes and have been described in the deep subsurface, such as the sulfate reducers *Pseudodesulfovibrio* sp. or *Candidatus Desulforudis*
165 *audaxviator* (Fig. 4). The latter has been reported in a South African gold mine at 2.8 km depth (Chivian et al., 2008), an aquifer at 2 km depth in Western Siberia (Karnachuk et al., 2019), and a groundwater at 755 m depth in California (Becraft et al., 2021). However, as these taxa represent less than 0.1 % of the sequence counts, ecological interpretations of the data were hindered. In general, these observations emphasize the critical importance of negative extraction controls, in addition to characterization of the drilling fluid, to enable identification of contaminant populations.

170




Figure 3 Microbial community structure fracture samples. a) Differentiation among microbial communities (beta diversity) prior to removing potential contaminants from the drilling fluid and the negative extraction controls. The labels depict the depth of the fracture in m below the surface (elevation field site 320 m). Beta diversity was estimated according to Bray-Curtis dissimilarities and visualized using non-metric dimensional scaling (NMDS). b) Differentiation among microbial communities after removing ASVs in the drilling fluid from



180 the respective fracture samples plus ASVs in the negative extraction controls. As shared ASVs between fracture and drilling fluid samples were excluded, the differentiation among these samples increased and therefore, these samples were further apart. c) Composition of the microbial community, depicting the nine most abundant orders with the remaining orders grouped as "Other". The four most abundant orders are represented by genera described as notorious reagent contaminants such as *Ralstonia* (Burkholderiales), *Sphingomonas* (Sphingomonadales), *Corynebacterium* (Mycobacteriales) and *Agrobacterium* (Rhizobiales). The text below each bar depicts the depth (m) of the fracture. d) Overlap among the microbial communities in the fractures, drilling fluid, and the negative extractions controls. The number of shared ASVs is followed by the abundance of these ASVs. For example, 38 ASVs were detected in both the fracture samples and the controls, and these taxa comprise 52 % of the sequence counts in the fracture samples.

3. Conclusions

185 The goal of this study was to characterize the subsurface microbial community in the pore space of rocks and natural fractures by extracting DNA to be used for constructing sequencing libraries combined with quantification of microbial abundance using CARD-FISH. The difficulty with extracting nucleic acids and the lack of a fluorescent signal in CARD-FISH analysis all indicated an extremely low microbial biomass in the sampled drill cores. In contrast, DNA was isolated from groundwater-bearing fractures, although ecological interpretations of the microbial communities in these fractures were compromised due to a high abundance of notorious contaminant genera. The presence of strict anaerobic sulfate-reducers that have been described as autochthonous deep biosphere microorganisms suggested subsurface microorganisms were present in the fractures. The method to trace contamination by adding fluorescent beads to the drilling fluid was evaluated to be successful. Not only did it allow to assess the success of surface sterilization, but it also allowed fluorescence microscopy on rock fragments taken from the inner core to quickly check if any microspheres were present, indicating contamination. Finally, in addition to the drilling fluid, the source of the drilling fluid (in this case a nearby freshwater lake) could be sampled at the water inlet to better discriminate potential contaminant populations.

4. Methods

4.1 Site description

200 The COSC-2 borehole (Lat 63.3124° N, Lon 13.5265° E) was drilled in an early Paleozoic rock formation with an elevation of 320 m and located in western central Sweden (Fig. 1). The top 780 m of the borehole were turbiditic greywackes that was followed by 45 m of sheared black shales. Underneath this shale zone, a sedimentary basement cover comprised of sandstones and conglomerates extended downwards to 1250 m, followed by ignimbrites and volcanic porphyries down to the total depth of 2276 m, interrupted by dolerite intrusions between 1600 and 1930 m. Temperature increases linearly with depth with an increase of ca. 2 °C every 100 m, starting at 5 °C at the surface and rising to 55 °C at the maximum depth.

205




Figure 4 Phylogenetic tree of selected ASVs in fracture samples. The tree was constructed using maximum likelihood on 16S rRNA gene amplicons (V3-V4 region, approximately 420 base pairs). Numbers on the nodes depict bootstrap values with values of 100 and 70 indicating a robust and moderate support, respectively. The number in the tip label shows the relative abundance with, for example, $2.7e^{-4}$ depicting that this ASV represents 0.027 % of all the sequence counts in the fracture samples, i.e., they were not detected in the drilling fluid nor in the extraction controls.

4.2 Field sampling

The sampling procedure and contamination control were described in detail in Lorenz et al. (2022). Briefly, rock core samples ($n = 50$) were obtained from 100 to 2250 m depth at 50 m intervals. Samples were 20-30 cm in length and either 61 mm in diameter for 100-1550 m depth or 45 mm in diameter for 1600-2250 m depth. The drilling fluid was sampled simultaneously with core recovery to characterize potential contaminants introduced during the drilling operation. The collected drilling fluid was subsampled (1 mL) for cell counts, fixed with 2 % (vol/vol) formaldehyde, stored in the dark at 4 °C, and transported to the home laboratory under identical conditions. Fluorescent microspheres (AFN-09, Radiant Color NV) were added to the drilling fluid in a concentration of 10^7 - 10^8 beads mL^{-1} (stock concentration 10^{18} beads mL^{-1}) that enabled to assess the success of surface sterilization as the bead's fluorescence was quenched when heated above 100 °C (Friese et al., 2017). Samples were collected and transported to the on-site microbiology laboratory where the surface of the rock samples was briefly flame



sterilized to combust contaminants present in the drilling fluid as evidenced by the quenching of the fluorescent microspheres. The number of beads on the core surface was counted using an on-site fluorescence microscope after sampling 2-3 g of surface
225 fragments and compared to the bead concentration in the drilling fluid at the time of sampling. Surface sterilization was repeated until at least 99.9 % of the beads on the core surface could no longer be observed. At 750 m depth, an alternative strategy was utilized by immersing the cores in a bead bath (10^7 - 10^8 beads mL^{-1}) rather than adding the beads to the drilling fluid due to adsorption of the fluorescent microspheres to clay particles. To do so, the bead concentration was quantified by washing the core's surface in a sterile plastic bag with 15 mL sterile, ultrapure water after immersing the core in the bead bath.
230 This liquid was collected, brought on a filter, stained, and analyzed with an on-site fluorescence microscope both before and after surface sterilization of the drill core. The core and the drilling fluid samples were stored on-site at -80°C and transported on dry ice to the home laboratory where they were kept at -80°C until further processing.

In addition to the core samples, natural fractures were sampled ($n = 29$) by collecting material surrounding the fracture using either flame-sterilized forceps or a chisel. 10 mL of sterile, ultrapure water was added to the material before storing the sample
235 at -80°C . Identical to the core samples, the drilling fluid was sampled simultaneously to sampling the fracture. Rock samples (2-3 g) for CARD-FISH were taken from the inner core, fixed in 4 % (vol/vol) formaldehyde in Mackintosh minimal media, and stored on-site at 4°C before being placed at -20°C once back at the home laboratory.

4.3 Cell counts

Fluorescence microscopy was used to quantify the cells in the drilling fluid. To do so, $1 \times$ TE-buffer (pH 8.0) was added to
240 the fixed sample to a final volume of 10 mL and this volume was brought on a polycarbonate black membrane (pore size $0.22 \mu\text{m}$, Whatman) under a pressure of 10 – 15 kPa and placed in a sterile petri dish to dry. Meanwhile, a staining solution was prepared consisting of 30 % 100X SYBR Green I (Invitrogen), 10 % 10 g L^{-1} p-phenylenediamine, and 60 % 1:1 glycerol in PBS. $10 \mu\text{L}$ of the solution was brought on a glass slide and cover slip before adding the filter and let incubate in the dark for 5 min. Finally, 300 cells or ten fields of view (5×5 raster) were counted in the dark at a 1000-fold magnification using an
245 epifluorescence microscope (Olympus BX50) combined with a light source (Olympus) that illuminated the DNA/dye complex at a wavelength of 497 nm.

4.4 CARD-FISH

Rock fragments were aseptically ground to sand grain size with a mortar and pestle. The ground rock was covered with 0.2 % (wt/vol) agarose, dried at 37°C , dehydrated with absolute ethanol, and stored at -20°C . To account for autofluorescence and
250 false positives, each sample was screened with fluorescence microscopy to select suitable fluorophores for avoiding mineral interference. Additionally, controls were carried out using the Non338 probe combined with multiple fluorophores and DNA-binding dyes (DAPI & Syto-9) on sterilized rock fragments by acid wash and overnight incubation at 120°C . More detail on the negative controls is provided in Escudero et al. (2018).



Hybridization, counter-staining, and mounting of the samples were performed as described in Pernthaler et al. (2008) using
255 the probes EUB338 I-III (Amann et al., 1990; Daims et al., 1999), ARC915 (Stahl and Amann, 1991), and Non338 (Wallner
et al., 1993). Endogenous peroxidases were inactivated using the protocol from Ishii et al. (2004). Samples were imaged using
a confocal laser scanning microscope (LSM710, Zeiss) coupled to an inverted platform (Axio Observer, Zeiss) and equipped
with diode (405 nm), argon (458/488/514 nm), helium (543 nm), and neon lasers (633 nm). Images were collected using a 63-
fold/1.4 oil immersion lens. Only the signals that matched the spectrum of the specific fluorophore were considered as positive
260 signals.

4.5 DNA extraction and amplification

Twelve rock samples covering all lithologies (Fig. 1) were subsampled using a flame-sterilized hammer and chisel (ASC
Scientific) on a metal benchtop that was sterilized with both bleach and 70 % ethanol. Subsamples were taken from the inner
core, avoiding the outer 10-20 mm of the core to reduce potential contamination introduced during field sampling. Several
265 subsamples were prepared simultaneously to avoid repeated freeze-thawing of the original sample. The rock fragments were
ground using a flame-sterilized percussion mortar and pestle (ASC Scientific) and stored at -20 °C until DNA extraction. To
verify if the processing of the rock fragments did not hinder downstream work, a positive control was made by adding 10⁹
cells of *Pseudomonas aeruginosa* to sterilized rock fragments prior to grinding the material, while flame-sterilized rock
fragments served as a negative control. After each extraction, the DNA yield was measured using a Qubit 2.0 fluorometer (Life
270 Technologies) with a lower detection limit of 0.05 ng μL⁻¹. As detectable amounts of DNA are not to be expected due to the
low biomass, the DNA extracts subsequently served as template in a PCR with 34 cycles using the primer pair 341F/805R
(Herlemann et al. 2011, product length circa 450 bp). To account for the potential fragmented nature of the DNA, a PCR was
also run using the bacterial primer pair 908F/1075R (Ohkuma and Kudo, 1998), covering a target region of circa 170 bp. The
archaeal primer pair 915F/1059R (Yu et al., 2005) was used to check for the potential presence of Archaea. The three PCRs
275 served as screening for potentially isolated DNA.

Initial attempts to extract DNA used the FastDNA Spin kit for soil (MP Biomedicals), DNeasy PowerSoil kit (Qiagen), and
the DNeasy PowerMax Soil kit (Qiagen). The latter was attempted due to the large amount of input material (10 g). According
to the manufacturer's instructions, the final elution volume was 5 mL and this volume was concentrated to 50 μL using the
DNA/RNA Clean All kit (Norgen Biotek). Due to potential adsorption of DNA to particles, an extraction protocol was designed
280 according to Lever et al. (2015) using 10 g of ground rock. Briefly, this protocol included a pyrophosphate wash (200 mM)
prior to cell lysis (50 °C for 2 h), chemical lysis based on a lysis solution (pH 10) containing 800 mM guanidium-HCl, 30 mM
EDTA, 0.5 % Triton X-100, and 30 mM Tris-HCl, 1 min of bead-beating using the rock fragments as beads, purification using
phenol-chloroform-isoamyl alcohol (pH 8.0) combined with phase-lock tubes (Qiagen), a chloroform-isoamyl alcohol wash
and precipitation overnight at -20 °C using ethanol-NaCl combined with linear polyacrylamide as co-precipitant. The obtained
285 pellet was dried, washed with 70 % ethanol, dissolved in 100 μL nuclease-free water, and purified using the DNA/RNA Clean
All kit (Norgen Biotek). Finally, the nucleic acids were eluted in 35 μL nuclease-free water and used as template for whole



genome amplification (described below). To increase the input material for DNA extraction even further, 100 g of ground rock from two cores was washed while shaking (300 rpm) with 250 mL 1 M phosphate buffer (pH 7) for 1 h at room temperature. The supernatant was decanted and filtered using a membrane filter (0.1 μm pore size, 45 mm in diameter) combined with a vacuum source to capture the cells. The membrane was directly subjected to DNA extraction using the DNeasy PowerWater kit (Qiagen), adjusting the elution volume to 50 μL . The negative extraction control was 250 mL of identical buffer without any rock added and subjected to identical downstream processing. For all methods described, reagents used were molecular grade, all plastic consumables (tips, centrifuge tubes) were PCR grade and placed for 10 min in a UV hood, glassware was baked at 160 $^{\circ}\text{C}$ for 3 h, and the lysis solution was filter sterilized (0.1 μm pore size) prior to autoclaving. The autoclave was cleaned prior to use and filled with fresh deionized water.

Fracture samples ($n = 29$) were extracted using the custom protocol based on Lever et al. (2015a) described above. The paired drilling fluid samples ($n = 28$) were extracted by bringing 30 mL of the fluid on a membrane filter (pore size 0.1 μm) using a vacuum source. The filter served as input for the DNeasy PowerWater kit (Qiagen), following the manufacturer's instructions except for eluting the nucleic acids in 50 μL instead of 100 μL . The DNA concentration of the fracture samples were all below detection limit (0.05 $\text{ng } \mu\text{L}^{-1}$) while the drilling fluid isolates contained 0.1 - 1.1 $\text{ng } \mu\text{L}^{-1}$. PCR amplification (34 cycles, primer pair 341F/805R) resulted in a product for seven out of 29 fracture samples. These amplified products and those from the respective drilling fluid samples were sequenced on an Illumina MiSeq platform at the Science for Life Laboratory, Sweden, generating 2×300 bp paired-end reads. Two negative extraction controls were subjected to identical processing, including amplification, and added to the sequencing project.

305 4.6 Whole genome amplification and PCR screening of amplified DNA

Rock DNA extracts were amplified with the Repli-g Single Cell kit (Qiagen) following the manufacturer's protocol with minor adaptations. DNA amplification was done in 384-well plates to allow for small reaction volumes, and in 96-well plates for larger volumes. The plates were UV-treated with 2 J while covered with an optical sealing and used the same day. Plastic consumables, lysis buffer, neutralization reagent, and nuclease-free water were UV-treated with 2 J directly prior to use and kept on ice. For 10 μL reaction volume, 1 μL of template was lysed with 0.5 μL D1 and neutralized with 0.5 μL N1. For 50 μL reaction volume, 2.5 μL template was lysed with 2.5 μL D1 and neutralized with 5 μL N1. Lysis was done at 95 $^{\circ}\text{C}$ for 15 s followed by 5 min at room temperature before reactions were placed on ice. 0.5 μM SYTO 13 (Invitrogen) was added to the amplification mix as a reporter dye. The amplification was run at 30 $^{\circ}\text{C}$ for 8 h with fluorescence readings every 15 min on a qPCR machine (CFX384 C1000, Bio-Rad) or plate reader (FLUOStar Omega, BMG LABTECH). The reactions were inactivated at 65 $^{\circ}\text{C}$ for 5 min and then stored at -20 $^{\circ}\text{C}$. The amplified DNA was diluted up to 1000-fold and screened with the 341F/805R primers targeting the bacterial 16S rRNA (Herlemann et al., 2011). The PCR was run with LightCycler 480 SYBR Green I Master mix (Roche), primers at a final concentration of 0.25 μM , 2 μl of diluted amplified DNA in a reaction volume of 10 μl on a LightCycler 480 (Roche). Amplification settings were 3 min at 95 $^{\circ}\text{C}$, 40 cycles of 15 s 95 $^{\circ}\text{C}$, 30 s 57 $^{\circ}\text{C}$, and 45 s at 72 $^{\circ}\text{C}$, followed by a melting curve.



320 4.7 Real-time PCR

Microbial abundance was assessed by quantifying the 16S ribosomal RNA gene copies using a real-time PCR (qPCR) on a LightCycler 480 (Roche Diagnostics). The reaction volume (10 μ L) consisted of 5 μ L Platinum SYBR Green qPCR SuperMix-UDG with ROX (Thermo Fisher Scientific), 0.4 μ L 10 μ M primer, 3.2 μ L nuclease-free water, and 1.0 μ L template. Bacterial and archaeal gene fragments were amplified using the primers 908F_mod/1075R (Ohkuma and Kudo, 1998) and 915F/1059R (Yu et al., 2005), respectively. Cycling conditions were 2 min at 95 $^{\circ}$ C, 40 cycles of 15 sec at 95 $^{\circ}$ C and 30 sec at 60 $^{\circ}$ C, followed by a melt curve analysis to assess product specificity. Standard curves were generated with a dilution series of purified PCR product using genomic DNA of pure cultures as template (i.e., *Acidiphilium cryptum* JF-5 for bacteria and *Ferroplasma acidiphilum* BRGM4 for archaea). Standards, samples of interest, and no-template controls were run in triplicates and the former two were 1:10 diluted in nuclease-free water to account for inhibition of the polymerase. The gene copy numbers were reported in gene copies g^{-1} after correcting for the mass of the rock used for DNA extraction and the elution volume. Reaction efficiency for the bacterial and archaeal assays were 95 and 96 %, respectively. A minimum number of three quantification cycles (ΔCq) between samples and no-template controls was maintained as a limit of detection.

4.8 Bioinformatics and reproducibility

Raw sequencing reads from the 16S ribosomal RNA gene amplicons ($n = 13$) were processed using the Ampliseq pipeline (v2.3.0) from the nf-core framework that relied on Nextflow (v21.10.6), Cutadapt (v3.4), FastQC (v0.11.9), DADA2 (v1.22.0), and the SBDI Sativa curated 16S GTDB database (release 207). The Ampliseq pipeline was run with default settings, except for the trimming of the primers whereby reads not containing the primer sequence or containing double copies were discarded from downstream analysis. ASVs present in the drilling fluid were removed from their respective fracture sample. For example, the ASVs detected in the drilling fluid taken at 1605 m depth were removed from the fracture sampled at the same depth (Table 1). By doing so, 62 % of the total sequence counts were removed from the fracture samples. ASVs present in the negative extraction controls ($n = 76$) were also excluded from downstream analysis. Absolute counts were standardized within a sample to relative abundances by dividing an ASVs count by the total number of counts within a sample. The phylogenetic tree was constructed using maximum likelihood and the substitution model (GTR + G + I) was selected based on the Akaike information criterion. The phylogeny was evaluated by bootstrapping ($n = 1000$). Statistics and data visualization were performed in using R (v4.2.1) and a compiled version of the R Markdown document is provided on GitHub at <https://github.com/geweaa/fractures/>.



350 **Data availability:** All sequencing data has been made publicly available at the European Nucleotide Archive under project reference PRJEB65120.

355 **Author contribution:** MD, SB, JK, and RA designed the research; GW, ST, MS, MM, PL, MB, PLH, SB, and MD performed field work; GW, CE and CB performed the molecular work; GW processed and performed ecological interpretations of the sequencing data; GW and MD wrote the manuscript with comments from all authors. MD, SB, and RA provided funding. All authors read and improved the final version.

360 **Competing interests:** The authors declare that they have no conflict of interests.

Acknowledgements

The authors thank Henning Lorenz for coordinating the drilling project and his expertise during field sampling, the COSC-2 drilling team and the on-site scientific personnel, Adrian Martinez and Guillermo Mateos for their assistance with the CARD-
365 FISH experiments, and Remco Hoogervorst, Andreas Arleborg, Ezhilarasan Mani Ezhilan, Egon and Ola Bertilsson for their assistance during field work. The drilling was funded by the International Continental Scientific Drilling Project (ICDP; grant no. 4-2017) with co-funding for scientific operations provided by Swedish Research Council (Vetenskapsrådet; grant nos. 2019-03688 and 2017-00642), the German Science Foundation (Deutsche Forschungsgemeinschaft DFG, grant no. SPP 1006), and the National Science Centre of Poland (Narodowym Centrum Nauki, project no. 2018/29/B/ST10/02315). The scientific
370 drilling operations was performed using the Swedish Research Council funded national research infrastructure for scientific drilling, Riksriggen. The study was supported by The Swedish Research Council (contracts 2018-04311 and 2017-04422). Whole genome amplification and PCR screening were done at the SciLifeLab Microbial Single Cell Genomics Facility at Uppsala University. SB acknowledges financial support from the Swedish Research Council and Science for Life Laboratory. Bioinformatics analyses were carried out utilizing the Uppsala Multidisciplinary Center for Advanced Computational Science
375 (UPPMAX) at Uppsala University (SNIC 2022/22-743 and SNIC 2022/6-242). The computations were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, Uppsala University partially funded by the Swedish Research Council through grant agreements no. 2022-06725 and no. 2018-05973.



380 References

- Amann, R. I., Binder, B. J., Olson, R. J., Chisholm, S. W., Devereux, R., and Stahl, D. A.: Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations, *Appl Environ Microbiol*, 56, 1919–1925, <https://doi.org/10.1128/aem.56.6.1919-1925.1990>, 1990.
- Amils, R., Escudero, C., Oggerin, M., Puente Sánchez, F., Arce Rodríguez, A., Fernández Remolar, D., Rodríguez, N., García Villadangos, M., Sanz, J. L., Briones, C., Sánchez-Román, M., Gómez, F., Leandro, T., Moreno-Paz, M., Prieto-Ballesteros, O., Molina, A., Tornos, F., Sánchez-Andrea, I., Timmis, K., Pieper, D. H., and Parro, V.: Coupled C, H, N, S and Fe biogeochemical cycles operating in the continental deep subsurface of the Iberian Pyrite Belt, *Environ Microbiol*, 25, 428–453, <https://doi.org/10.1111/1462-2920.16291>, 2023.
- 385
- Becraft, E. D., Lau Vetter, M. C. Y., Bezuidt, O. K. I., Brown, J. M., Labonté, J. M., Kauneckaitė-Griguole, K., Salkauskaite, R., Alzbutas, G., Sackett, J. D., Kruger, B. R., Kadnikov, V., van Heerden, E., Moser, D., Ravin, N., Onstott, T., and Stepanauskas, R.: Evolutionary stasis of a deep subsurface microbial lineage, *ISME J*, 15, 2830–2842, <https://doi.org/10.1038/s41396-021-00965-3>, 2021.
- 390
- Bell, E., Lamminmäki, T., Alneberg, J., Andersson, A. F., Qian, C., Xiong, W., Hettich, R. L., Frutschi, M., and Bernier-Latmani, R.: Active sulfur cycling in the terrestrial deep subsurface, *ISME J*, 14, 1260–1272, <https://doi.org/10.1038/s41396-020-0602-x>, 2020.
- 395
- Bell, E., Lamminmäki, T., Alneberg, J., Qian, C., Xiong, W., Hettich, R. L., Frutschi, M., and Bernier-Latmani, R.: Active anaerobic methane oxidation and sulfur disproportionation in the deep terrestrial subsurface, *ISME J*, 16, <https://doi.org/10.1038/s41396-022-01207-w>, 2022.
- Beller, H. R., Chain, P. S. G., Letain, T. E., Chakicherla, A., Larimer, F. W., Richardson, P. M., Coleman, M. A., Wood, A. P., and Kelly, D. P.: The genome sequence of the obligately chemolithoautotrophic, facultatively anaerobic bacterium *Thiobacillus denitrificans*, *J Bacteriol*, 188, <https://doi.org/10.1128/JB.188.4.1473-1488.2006>, 2006.
- 400
- Bomberg, M., Lamminmäki, T., and Itävaara, M.: Microbial communities and their predicted metabolic characteristics in deep fracture groundwaters of the crystalline bedrock at Olkiluoto, Finland, *Biogeosciences*, 13, <https://doi.org/10.5194/bg-13-6031-2016>, 2016.
- 405
- Borgonie, G., Linage-Alvarez, B., Ojo, A. O., Mundle, S. O. C., Freese, L. B., Van Rooyen, C., Kuloyo, O., Albertyn, J., Pohl, C., Cason, E. D., Vermeulen, J., Pienaar, C., Litthauer, D., Van Niekerk, H., Van Eeden, J., Lollar, B. S., Onstott, T. C., and Van Heerden, E.: Eukaryotic opportunists dominate the deep-subsurface biosphere in South Africa, *Nat Commun*, 6, 8952, <https://doi.org/10.1038/ncomms9952>, 2015.
- 410
- Boylan, A. A., Perez-Mon, C., Guillard, L., Burzan, N., Loreggian, L., Maisch, M., Kappler, A., Byrne, J. M., and Bernier-Latmani, R.: H₂-fuelled microbial metabolism in Opalinus Clay, *Appl Clay Sci*, 174, <https://doi.org/10.1016/j.clay.2019.03.020>, 2019.



- Braslavsky, I., Hebert, B., Kartalov, E., and Quake, S. R.: Sequence information can be obtained from single DNA molecules, *Proc Natl Acad Sci U S A*, 100, <https://doi.org/10.1073/pnas.0230489100>, 2003.
- 415 Casar, C. P., Kruger, B. R., Flynn, T. M., Masterson, A. L., Momper, L. M., and Osburn, M. R.: Mineral-hosted biofilm communities in the continental deep subsurface, Deep Mine Microbial Observatory, SD, USA, *Geobiology*, 18, 508–522, <https://doi.org/10.1111/gbi.12391>, 2020.
- Chapelle, F. H., O’Neill, K., Bradley, P. M., Methá, B. A., Ciufu, S. A., Knobel, L. R. L., and Lovley, D. R.: A hydrogen-based subsurface microbial community dominated by methanogens, *Nature*, 415, <https://doi.org/10.1038/415312a>, 2002.
- 420 Chivian, D., Brodie, E. L., Alm, E. J., Culley, D. E., Dehal, P. S., DeSantis, T. Z., Gihring, T. M., Lapidus, A., Lin, L. H., Lowry, S. R., Moser, D. P., Richardson, P. M., Southam, G., Wanger, G., Pratt, L. M., Andersen, G. L., Hazen, T. C., Brockman, F. J., Arkin, A. P., and Onstott, T. C.: Environmental genomics reveals a single-species ecosystem deep within Earth, *Science*, 322, 275–278, <https://doi.org/10.1126/science.1155495>, 2008.
- 425 Dai, X., Wang, Y., Luo, L., Pfiffner, S. M., Li, G., Dong, Z., Xu, Z., Dong, H., and Huang, L.: Detection of the deep biosphere in metamorphic rocks from the Chinese continental scientific drilling, *Geobiology*, 19, <https://doi.org/10.1111/gbi.12430>, 2021.
- Daims, H., Brühl, A., Amann, R., Schleifer, K. H., and Wagner, M.: The domain-specific probe EUB338 is insufficient for the detection of all bacteria: Development and evaluation of a more comprehensive probe set, *Syst Appl Microbiol*, 22, 434–444, [https://doi.org/10.1016/S0723-2020\(99\)80053-8](https://doi.org/10.1016/S0723-2020(99)80053-8), 1999.
- 430 Dutta, A., Dutta Gupta, S., Gupta, A., Sarkar, J., Roy, S., Mukherjee, A., and Sar, P.: Exploration of deep terrestrial subsurface microbiome in Late Cretaceous Deccan traps and underlying Archean basement, India, *Sci Rep*, 8, <https://doi.org/10.1038/s41598-018-35940-0>, 2018.
- Engelhardt, T., Kallmeyer, J., Cypionka, H., and Engelen, B.: High virus-to-cell ratios indicate ongoing production of viruses in deep subsurface sediments, *ISME J*, 8, <https://doi.org/10.1038/ismej.2013.245>, 2014.
- 435 Escudero, C., Vera, M., Oggerin, M., and Amils, R.: Active microbial biofilms in deep poor porous continental subsurface rocks, *Sci Rep*, 8, 1538, <https://doi.org/10.1038/s41598-018-19903-z>, 2018.
- Escudero, C., del Campo, A., Ares, J. R., Sánchez, C., Martínez, J. M., Gómez, F., and Amils, R.: Visualizing microorganism-mineral interaction in the Iberian Pyrite Belt subsurface: the *Acidovorax* case, *Front Microbiol*, 11, <https://doi.org/10.3389/fmicb.2020.572104>, 2020.
- 440 Friese, A., Kallmeyer, J., Kitte, J. A., Martínez, I. M., Bijaksana, S., and Wagner, D.: A simple and inexpensive technique for assessing contamination during drilling operations, *Limnol Oceanogr Methods*, 15, <https://doi.org/10.1002/lom3.10159>, 2017.
- Ghosh, W. and Dam, B.: Biochemistry and molecular biology of lithotrophic sulfur oxidation by taxonomically and ecologically diverse bacteria and archaea, <https://doi.org/10.1111/j.1574-6976.2009.00187.x>, 2009.
- 445 Gold, T.: The deep, hot biosphere, *Proc Natl Acad Sci U S A*, 89, 6045–6049, <https://doi.org/10.1073/pnas.89.13.6045>, 1992.



- Golubic, S., Friedmann, I., and Schneider, J.: The lithobiontic ecological niche, with special reference to microorganisms., *J Sediment Petrol*, 51, <https://doi.org/10.1306/212f7cb6-2b24-11d7-8648000102c1865d>, 1981.
- Herlemann, D. P. R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., and Andersson, A. F.: Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea, *ISME J*, 5, 1571–1579, <https://doi.org/10.1038/ismej.2011.41>, 2011.
- 450 Holmfeldt, K., Nilsson, E., Simone, D., Lopez-Fernandez, M., Wu, X., de Bruijn, I., Lundin, D., Andersson, A. F., Bertilsson, S., and Dopson, M.: The Fennoscandian Shield deep terrestrial virosphere suggests slow motion “boom and burst” cycles, *Commun Biol*, 4, 307, <https://doi.org/10.1038/s42003-021-01810-1>, 2021.
- Hoshino, T., Yilmaz, L. S., Noguera, D. R., Daims, H., and Wagner, M.: Quantification of target molecules needed to detect microorganisms by fluorescence in situ hybridization (FISH) and catalyzed reporter deposition-FISH, *Appl Environ Microbiol*, 74, <https://doi.org/10.1128/AEM.00208-08>, 2008.
- 455 Hug, L. A., Baker, B. J., Anantharaman, K., Brown, C. T., Probst, A. J., Castelle, C. J., Butterfield, C. N., Hermsdorf, A. W., Amano, Y., Ise, K., Suzuki, Y., Dudek, N., Relman, D. A., Finstad, K. M., Amundson, R., Thomas, B. C., and Banfield, J. F.: A new view of the tree of life, *Nat Microbiol*, 1, <https://doi.org/10.1038/nmicrobiol.2016.48>, 2016.
- 460 Ishii, K., Mußmann, M., MacGregor, B. J., and Amann, R.: An improved fluorescence in situ hybridization protocol for the identification of bacteria and archaea in marine sediments, *FEMS Microbiol Ecol*, 50, 203–213, <https://doi.org/10.1016/j.femsec.2004.06.015>, 2004.
- Kallmeyer, J.: Contamination control for scientific drilling operations, *Adv Appl Microbiol*, 98, <https://doi.org/10.1016/bs.aamb.2016.09.003>, 2017.
- 465 Karnachuk, O. V., Frank, Y. A., Lukina, A. P., Kadnikov, V. V., Beletsky, A. V., Mardanov, A. V., and Ravin, N. V.: Domestication of previously uncultivated *Candidatus Desulforudis audaxviator* from a deep aquifer in Siberia sheds light on its physiology and evolution, *ISME J*, 13, <https://doi.org/10.1038/s41396-019-0402-3>, 2019.
- Kieft, T. L.: Their world: a diversity of microbial environments, in: *Microbiology of the deep continental biosphere*, Springer, 225–249, 2016.
- 470 Lever, M. A., Alperin, M., Engelen, B., Inagaki, F., Nakagawa, S., Steinsbu, B. O., and Teske, A.: Trends in basalt and sediment core contamination during IODP expedition 301, *Geomicrobiol J*, 23, <https://doi.org/10.1080/01490450600897245>, 2006.
- Lever, M. A., Torti, A., Eickenbusch, P., Michaud, A. B., Šantl-Temkiv, T., and Jørgensen, B. B.: A modular method for the extraction of DNA and RNA, and the separation of DNA pools from diverse environmental sample types, *Front Microbiol*, 6, 476, <https://doi.org/10.3389/fmicb.2015.00476>, 2015a.
- 475 Lever, M. A., Rogers, K. L., Lloyd, K. G., Overmann, J., Schink, B., Thauer, R. K., Hoehler, T. M., and Jørgensen, B. B.: Life under extreme energy limitation: A synthesis of laboratory- and field-based investigations, *FEMS Microbiol Rev*, 39, 688–728, <https://doi.org/10.1093/femsre/fuv020>, 2015b.



- Lopez-Fernandez, M., Westmeijer, G., Turner, S., Broman, E., Ståhle, M., Bertilsson, S., and Dopson, M.: Thiobacillus as a
480 key player for biofilm formation in oligotrophic groundwaters of the Fennoscandian Shield, NPJ Biofilms
Microbiomes, 9, <https://doi.org/10.1038/s41522-023-00408-1>, 2023.
- Lorenz, H., Rosberg, J. E., Juhlin, C., Bjelm, L., Almqvist, B. S. G., Berthet, T., Conze, R., Gee, D. G., Klonowska, I., Pascal,
C., Pedersen, K., Roberts, N. M. W., and Tsang, C. F.: COSC-1 - drilling of a subduction-related allochthon in the
Palaeozoic Caledonide orogen of Scandinavia, Scientific Drilling, 19, <https://doi.org/10.5194/sd-19-1-2015>, 2015.
- 485 Lorenz, H., Juhlin, C., Rosberg, J.-E., Bazargan, M., Klonowska, I., Kück, J., Lescoutre, R., Rejkjær, S., Westmeijer, G., and
Ziemniak, G.: COSC-2 operational report - Operational data sets, GFZ Data Services,
<https://doi.org/https://doi.org/10.5880/ICDP.5054.003>, 2021.
- Lorenz, H., Rosberg, J. E., Juhlin, C., Klonowska, I., Lescoutre, R., Westmeijer, G., Almqvist, B. S. G., Anderson, M.,
Bertilsson, S., Dopson, M., Kallmeyer, J., Kück, J., Lehnert, O., Menegon, L., Pascal, C., Rejkjær, S., and Roberts,
490 N. N. W.: COSC-2 - Drilling the basal décollement and underlying margin of palaeocontinent Baltica in the Paleozoic
Caledonide Orogen of Scandinavia, Scientific Drilling, 30, 43–57, <https://doi.org/10.5194/sd-30-43-2022>, 2022.
- Magnabosco, C., Ryan, K., Lau, M. C. Y., Kuloyo, O., Sherwood Lollar, B., Kieft, T. L., Van HeerDen, E., and Onstott, T. C.:
A metagenomic window into carbon metabolism at 3 km depth in Precambrian continental crust, ISME J, 10,
<https://doi.org/10.1038/ismej.2015.150>, 2016.
- 495 Magnabosco, C., Lin, L. H., Dong, H., Bomberg, M., Ghiorse, W., Stan-Lotter, H., Pedersen, K., Kieft, T. L., van Heerden,
E., and Onstott, T. C.: The biomass and biodiversity of the continental subsurface, Nat Geosci, 11, 707–717,
<https://doi.org/10.1038/s41561-018-0221-6>, 2018.
- Mangelsdorf, K. and Kallmeyer, J.: Integration of deep biosphere research into the International Continental Scientific Drilling
Program, Scientific Drilling, 10, 46–55, <https://doi.org/10.5194/sd-10-46-2010>, 2010.
- 500 Moter, A. and Göbel, U. B.: Fluorescence in situ hybridization (FISH) for direct visualization of microorganisms,
[https://doi.org/10.1016/S0167-7012\(00\)00152-4](https://doi.org/10.1016/S0167-7012(00)00152-4), 2000.
- Ohkuma, M. and Kudo, T.: Phylogenetic analysis of the symbiotic intestinal microflora of the termite *Cryptotermes*
domesticus, FEMS Microbiol Lett, 164, [https://doi.org/10.1016/S0378-1097\(98\)00244-4](https://doi.org/10.1016/S0378-1097(98)00244-4), 1998.
- Osterholz, H., Turner, S., Alakangas, L. J., Tullborg, E.-L., Dittmar, T., Kalinowski, B. E., and Dopson, M.: Terrigenous
505 dissolved organic matter persists in the energy-limited deep groundwaters of the Fennoscandian Shield, Nat Commun,
13, 4837, <https://doi.org/10.1038/s41467-022-32457-z>, 2022.
- Pantke, C., Obst, M., Benzerara, K., Morin, G., Ona-Nguema, G., Dippon, U., and Kappler, A.: Green rust formation during
Fe(II) oxidation by the nitrate-reducing *Acidovorax* sp. strain BoFeN1, Environ Sci Technol, 46,
<https://doi.org/10.1021/es2016457>, 2012.
- 510 Pedersen, K.: Exploration of deep intraterrestrial microbial life: current perspectives, FEMS Microbiol Lett, 185,
<https://doi.org/10.1111/j.1574-6968.2000.tb09033.x>, 2000.



- Pernthaler, A., Pernthaler, J., and Amann, R.: Sensitive multi-color fluorescence in situ hybridization for the identification of environmental microorganisms, in: *Molecular Microbial Ecology Manual*, 711–726, https://doi.org/10.1007/978-1-4020-2177-0_311, 2008.
- 515 Puente-Sánchez, F., Moreno-Paz, M., Rivas, L. A., Cruz-Gil, P., García-Villadangos, M., Gómez, M. J., Postigo, M., Garrido, P., González-Toril, E., Briones, C., Fernández-Remolar, D., Stoker, C., Amils, R., and Parro, V.: Deep subsurface sulfate reduction and methanogenesis in the Iberian Pyrite Belt revealed through geochemistry and molecular biomarkers, *Geobiology*, 12, <https://doi.org/10.1111/gbi.12065>, 2014.
- Puente-Sánchez, F., Arce-Rodríguez, A., Oggerin, M., García-Villadangos, M., Moreno-Paz, M., Blanco, Y., Rodríguez, N., 520 Bird, L., Lincoln, S. A., Tornos, F., Prieto-Ballesteros, O., Freeman, K. H., Pieper, D. H., Timmis, K. N., Amils, R., and Parro, V.: Viable cyanobacteria in the deep continental subsurface, *Proc Natl Acad Sci U S A*, 115, <https://doi.org/10.1073/pnas.1808176115>, 2018.
- Purkamo, L., Kietäväinen, R., Nuppunen-Puputti, M., Bomberg, M., and Cousins, C.: Ultradeep microbial communities at 4.4 525 km within crystalline bedrock: Implications for habitability in a planetary context, *Life*, 10, <https://doi.org/10.3390/life10010002>, 2020.
- Rahlff, J., Turzynski, V., Esser, S. P., Monsees, I., Bornemann, T. L. V., Figueroa-Gonzalez, P. A., Schulz, F., Woyke, T., Klingl, A., Moraru, C., and Probst, A. J.: Lytic archaeal viruses infect abundant primary producers in Earth's crust, *Nat Commun*, 12, <https://doi.org/10.1038/s41467-021-24803-4>, 2021.
- Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., Turner, P., Parkhill, J., Loman, N. J., and 530 Walker, A. W.: Reagent and laboratory contamination can critically impact sequence-based microbiome analyses, *BMC Biol*, 12, 1–12, <https://doi.org/10.1186/s12915-014-0087-z>, 2014.
- Sheik, C. S., Reese, B. K., Twing, K. I., Sylvan, J. B., Grim, S. L., Schrenk, M. O., Sogin, M. L., and Colwell, F. S.: Identification and removal of contaminant sequences from ribosomal gene databases: Lessons from the Census of Deep Life, *Front Microbiol*, 9, 840, <https://doi.org/10.3389/fmicb.2018.00840>, 2018.
- 535 Smith, D. C., Spivack, A. J., Fisk, M. R., Haveman, S. A., and Staudigel, H.: Tracer-based estimates of drilling-induced microbial contamination of deep sea crust, *Geomicrobiol J*, 17, <https://doi.org/10.1080/01490450050121170>, 2000.
- Stahl, D. A. and Amann, R.: Development and application of nucleic acid probes in bacterial systematics, in: *Sequencing and hybridization techniques in bacterial systematics*, John Wiley & Sons Ltd, Chichester, UK, 205–248, 1991.
- Starnawski, P., Bataillon, T., Ettema, T. J. G., Jochum, L. M., Schreiber, L., Chen, X., Lever, M. A., Polz, M. F., Jørgensen, 540 B. B., Schramm, A., and Kjeldsen, K. U.: Microbial community assembly and evolution in seafloor sediment, *Proc Natl Acad Sci U S A*, 114, <https://doi.org/10.1073/pnas.1614190114>, 2017.
- Suzuki, Y., Konno, U., Fukuda, A., Komatsu, D. D., Hirota, A., Watanabe, K., Togo, Y., Morikawa, N., Hagiwara, H., Aosai, D., Iwatsuki, T., Tsunogai, U., Nagao, S., Ito, K., and Mizuno, T.: Biogeochemical signals from deep microbial life in terrestrial crust, *PLoS One*, 9, <https://doi.org/10.1371/journal.pone.0113063>, 2014.
- 545 Teske, A. P.: The deep subsurface biosphere is alive and well, <https://doi.org/10.1016/j.tim.2005.07.004>, 2005.



- Wallner, G., Amann, R., and Beisker, W.: Optimizing fluorescent in situ hybridization with rRNA-targeted oligonucleotide probes for flow cytometric identification of microorganisms, *Cytometry*, 14, 136–142, <https://doi.org/10.1002/cyto.990140205>, 1993.
- 550 Wu, X., Holmfeldt, K., Hubalek, V., Lundin, D., Åström, M., Bertilsson, S., and Dopson, M.: Microbial metagenomes from three aquifers in the Fennoscandian Shield terrestrial deep biosphere reveal metabolic partitioning among populations, *ISME J*, 10, 1192–1203, <https://doi.org/10.1038/ismej.2015.185>, 2016.
- Wu, X., Pedersen, K., Edlund, J., Eriksson, L., Åström, M., Andersson, A. F., Bertilsson, S., and Dopson, M.: Potential for hydrogen-oxidizing chemolithoautotrophic and diazotrophic populations to initiate biofilm formation in oligotrophic, deep terrestrial subsurface waters, *Microbiome*, 5, 1–13, <https://doi.org/10.1186/S40168-017-0253-Y>, 2017.
- 555 Yu, Y., Lee, C., and Hwang, S.: Analysis of community structures in anaerobic processes using a quantitative real-time PCR method, *Water Science and Technology*, 52, <https://doi.org/10.2166/wst.2005.0502>, 2005.