

Optical and radar Earth observation data for up-scaling methane emissions linked to permafrost degradation in sub-Arctic peatlands in Northern Sweden

Sofie Sjögersten¹, Martha Ledger¹, Matthias Siewert², Betsabé de la Barreda-Bautista^{1,3}, Andrew Sowter⁴, David Gee⁴, Giles
5 Foody³, Doreen S. Boyd³

¹University of Nottingham, School of Biosciences, College Road, Sutton Bonington, Loughborough, LE12 5RD, UK

²University of Umeå, Department of Ecology and Environmental Sciences, KB H4, Linnaeus väg 6, Umeå universitet, 901 87
Umeå, Sweden

10 ³School of Geography, University of Nottingham, University Park, Nottingham, NG7 2RD, UK

⁴Terra Motion Ltd, Ingenuity Centre, Triumph Rd, Nottingham NG7 2TU, UK

Correspondence to: Sofie.Sjogersten@nottingham.ac.uk

Abstract. Permafrost thaw in Arctic regions is increasing methane (CH₄) emissions to the atmosphere, but quantification of such emissions is difficult given the large and remote areas impacted. Hence, Earth Observation (EO) data are critical for
15 assessing both permafrost thaw, associated ecosystem change, and increased CH₄ emissions. Often extrapolation from field measurements using EO is the approach employed. However, there are key challenges to consider. Landscape CH₄ emissions result from a complex local-scale mixture of micro-topographies and vegetation types that support widely differing CH₄ emissions and it is difficult to detect the initial stages of permafrost degradation before vegetation transitions have occurred. This study considers the use of a combination of ultra-high resolution unoccupied aerial vehicle (UAV) data, together with
20 Sentinel-1 and -2 data to extrapolate field measurements of CH₄ emissions from a set of vegetation types which capture the local variation in vegetation on degrading palsas wetlands. We show that the ultra-high resolution UAV data can map spatial variation in vegetation relevant to variation in CH₄ emissions and extrapolate these across the wider landscape. We further show how this can be integrated with Sentinel-1 and Sentinel-2 data. By way of a soft classification, and simple correction of misclassification bias of a hard classification, the output vegetation mapping and subsequent extrapolation of CH₄ emissions
25 matched closely that generated using the UAV data. InSAR assessment of subsidence together with the vegetation classification suggested that high subsidence rates of palas wetland can be used to quantify areas at risk of increased CH₄ emissions. The transition of a 50 ha area currently experiencing subsidence to fen vegetation is estimated to increase emissions from 116 kg CH₄ season⁻¹, to emissions as high as 6500 to 13000 kg CH₄ season⁻¹. The key outcome from this study is that a combination of high and low resolution EO data of different types provides the ability to estimate CH₄ emissions from large
30 geographies covered by a fine mixture of vegetation types and vulnerable to transitioning to CH₄ emitters in the near future. This points to an opportunity to measure and monitor CH₄ emissions from the Arctic over space and time with confidence.

1 Introduction

35 The severe impact of climate heating in the Arctic has raised grave concerns of positive feedbacks from ecosystems that store substantial amounts of soil organic carbon in soils and peat deposits (Biskaborn et al., 2019; Chadburn et al., 2017; Grosse et al., 2016; Hugelius et al., 2020; Mishra et al., 2021; Schneider von Deimling et al., 2012; Sjöberg et al., 2020). Although the overall carbon balance in permafrost and non-permafrost peatlands is similar (Olefeldt et al., 2012), permafrost thaw increases methane (CH₄) emissions across a range of Arctic landscapes (Olefeldt et al., 2013). Any increase in emissions of CH₄ is of particular concern, as it is a powerful greenhouse gas with a global warming potential of 28 that of CO₂ on a 100 year time frame, with Arctic ecosystems acting as strong emitters of CH₄ (Euskirchen et al., 2014; Glagolev et al., 2011; IPCC, 2021; Maksyutov et al., 2010; McGuire et al., 2009; Turetsky et al., 2020).

Palsa are elevated peat mounds with a frozen core formed in peatlands. They are a common landform in the sporadic and discontinuous zone of the permafrost region where palsa peatlands cover substantial areas of the total permafrost region (Ballantyne C. K., 2018). From a climate feedback perspective, these are critical systems as they are carbon (C) rich, with an estimated storage of 100 Gt C in the Arctic region (based on a combination of field and modelled data in: Hugelius et al., 2020; Schuur & Abbott, 2011; Tarnocai et al., 2009). Palsa peatland are dynamic systems that comprise of raised peatland plateaux, which can vary in size from tens of meters to kilometers in size, these plateaux are often surrounded by lower lying waterlogged areas or have borders with thermokarst lakes (Ballantyne C. K., 2018; Zuidhoff & Kolstrup, 2000). As a result of the raised nature of palsa their surface tend to be relatively dry. However, this changes as the permafrost in these peatlands degrade and the peat surface subsides which, over time, results in waterlogging (Olvmo et al., 2020; Sjöberg et al., 2015). Such degradation is already being observed in many areas of the Arctic in response to the greater than average warming occurring in this region (Åkerman & Johansson, 2008; Borge et al., 2017; de la Barreda-Bautista et al., 2022; Luoto & Seppälä, 2003; Olvmo et al., 2020; Sannel et al., 2016; Sannel & Kuhry, 2011). The loss of permafrost and switch to waterlogged conditions represents a profound change in how this ecosystem functions, driving increased CH₄ emissions which have been observed across many Arctic landscapes (Glagolev et al., 2011; Miglovets et al., 2021; Varner et al., 2022; Walter Anthony et al., 2016; Walter et al., 2006). To understand fully the magnitude of these feedbacks and their impacts on the global climate, it is imperative to both quantify current emissions and predict new areas that may become CH₄ emitters across Arctic landscapes undergoing permafrost degradation.

60 Given the size of the areas under threat from warming and their remoteness, remote sensing techniques are necessary to detect when and where permafrost degradation and subsidence is occurring. Coupled with measures of CH₄ flux, this may offer an approach for understanding, estimating, and predicting CH₄ feedbacks at landscape scales across permafrost peatlands of the circumarctic region. Detecting year-to-year degradation of permafrost manifested as subsidence is possible using the Advanced Pixel System using Intermittent Baseline Subset (ASPIS-InSAR) and other InSAR techniques (Alshammari et al., 2018; de la Barreda-Bautista et al., 2022; van Huissteden et al., 2021). Measures of surface subsidence of permafrost affected

areas and the rate of ground motion can give an indication of permafrost degradation extents and rates. Furthermore, the ASPIS-InSAR data is unaffected by clouds, is based on publically available Sentinel-1 data, and can be used to analyse impact at continental scale. This means that ASPIS-InSAR can be used to identify areas that are at risk of increased CH₄ emissions and help to assess the radiative forcing ecosystem feedback resulting from permafrost thaw.

70 Complimentary remote sensing derived vegetation maps can be used to extrapolate plot based CH₄ emission data to the wider landscape (Varner et al., 2022). However, the scale-dependency of ecosystem properties that are non-linearly distributed in space (Siewert, 2018; Siewert & Olofsson, 2020) means that the spatial resolution of the remotely sensed data are important. Coarse spatial resolution remote sensing data, such as that from openly available and high temporal optical satellite-borne sensors, run the risk of missing small areas of those vegetation types with high CH₄ emissions, which may lead to an underestimate of total emissions (Bernhardt et al., 2017). Further, the relatively slow rate of the response of vegetation to ground condition changes resulting from permafrost degradation means mapping vegetation may not fully explain patterns of CH₄ emissions (de la Barreda-Bautista et al., 2022). After subsidence starts to occur, the original vegetation type can persist for many years (>10 years) until the site conditions have changed to a point that the original vegetation is outcompeted (Johansson et al., 2013). This makes it difficult to use vegetation change in isolation as a year-on-year proxy for permafrost degradation and linked potential regulation of CH₄ emissions. Furthermore, permafrost degradation and linked increases in CH₄ production, is driven by several processes across scales. Variation in root inputs of labile carbon or oxygen in the peat profile occur over millimetres and can vary seasonally in response to water levels (Ström & Christensen, 2007; Ström et al., 2003). Lateral erosion and subsidence linked to permafrost degradation occur on meter scales when measured over several years, while changes in the vegetation community from raised palsa vegetation to fen type vegetation occur over tens of meters at decadal time-scales (de la Barreda-Bautista et al., 2022; Varner et al., 2022). Together the different processes associated with permafrost degradation result in a spatially varied landscape with a mixture of functionally distinct geomorphological units with regard to soil moisture content and vegetation productivity, creating a palimpsest of processes and properties determining the effective greenhouse gas feedback of these regions (Siewert et al., 2021). Therefore, remote sensing methodologies which capture fine-scale spatial variation in vegetation composition, in addition to permafrost degradation over seasonal to yearly time scales, are needed to provide appropriate observations of degrading peatlands in the Arctic in order to measure their CH₄ emissions accurately.

With a view to estimating CH₄ emissions resulting from permafrost degradation in peatlands, this paper explores two remote sensing technologies to provide data for extrapolating plot-based methane measurements over sub-Arctic peatlands of northern Sweden. Specific objectives were to (i) carry out detailed field surveys to capture small-sized or local-scale variation in CH₄ emissions linked to vegetation type and permafrost subsidence; (ii) conduct vegetation mapping from Sentinel-2 satellite data, and in particular demonstrate the value of rigorous map validation on the quantification of CH₄ emissions; and (iii) quantify CH₄ emissions from areas experiencing permafrost degradation and those at high risk of subsidence as determined from Sentinel-1 InSAR (processed using ASPIS) data.

2.1 Field site description

Three peatland locations affected by permafrost conditions - the Tourist Station, Storflaket, and Stordalen wetlands (Figure 1) - situated to the south of Lake Torneträsk near Abisko, northern Sweden (68°12'N, 19°03'E, 351 masl), were the focus of study. The study sites are in sub-Arctic climatic zone, the mean annual temperature (MAT) was 0.3 °C and the mean annual precipitation (MAP) was 347.2 mm between 1990-2020 (www.smhi.se). The study sites comprise areas of raised palsas (i.e. relatively dry peat plateaus) and waterlogged areas with fen or birch forest vegetation. All three sites show signs of active permafrost thaw indicated by a lowering of the ground surface (i.e. ground subsidence), high soil moisture levels, and areas of palsa collapse (i.e. complete permafrost degradation and transition from palsa to wetland) (Åkerman & Johansson, 2008). The raised palsa parts of the sites tend to get snow free in April/May but snow can remain in hollows into June.

Previous mapping of the vegetation at these sites using Unoccupied Aerial Vehicles (UAVs) and subsidence estimated using ASPIS-InSAR (de la Barreda-Bautista et al., 2022) show the vegetation on the palsa to be composed of bryophytes (e.g. *Sphagnum fuscum*), lichens and dwarf shrubs (*Empetrum nigrum*, *Andromeda polyfolia*, and *Betula nana*; (Sjögersten et al., 2016)). The most common herbaceous species was *Rubus chamaemorus*. Recently collapsed areas adjacent to and within the plateau areas tended to be vegetated by *Sphagnum* sp. and *Eriophorum vaginatum* and *E. angustifolium*. Extensively collapsed and subsequently flooded areas tended to be vegetated by graminoids, mainly *Carex* and *Eriophorum* species, herbaceous plants mainly *Menyanthes trifoliata*, *Comarum palustre*, and deciduous shrubs, such as *Salix lapponica*, *Salix ssp.* and *Betula nana*. Forested wetland areas tended to be located adjacent to streams and at the perimeters of the wetland areas. The mean peat depth was ca 60, 94, 50, and 40 cm, on palsas, *Sphagnum* sp., sedge, and forested wetland areas, respectively; mean soil C storage in the top 100 cm was 52, 34, 33, 40 kg SOC m⁻² for palsa, *Sphagnum* sp., sedge, and forested wetland, respectively (Siewert, 2018). The depth of the active layer (the seasonally thawed upper part of the soil) varies between drier areas, which have a shallower (ca 30 – 60 cm) active layer, and wetter areas, where the active layer is deeper or where there is no permafrost at all in the upper 1.5 m. In the Torneträsk area, the active layer depths have increased, between 0.2 and 2.0 cm yr⁻¹ over 1978-2006, with a higher rate in more recent years (Åkerman & Johansson, 2008).



Figure 1. Map showing the location of the three permafrost peatland study sites: Tourist Station, Storflaket and Stordalen. Inset: Location of Abisko in northern Sweden. The landcover classes were based on the CORINE land cover classification.

130

2.2 Experimental design

The fieldwork to measure CH_4 emissions at the three study sites was carried out across the dates of 8-15 June, 18-29 of July and 8-12 of September 2021. At each site, we targeted six common vegetation types. Dry lichen, dwarf shrub, moist moss, sphagnum wetland, sedge wetland and willow wetland with the dry lichen, dwarf shrub and moist moss vegetation types occurring on the drier parts on the palsas. The sphagnum wetland, sedge wetland and willow wetland vegetation types occurred in waterlogged fen or thermokarst areas. We also measured CH_4 emissions from subsiding areas with dry lichen, dwarf shrub and moist moss vegetation types, subsiding areas was identified visually based on changes in topography, subsidence and waterlogging. Measurements in each vegetation type were carried out in blocks of ca 10 m diameter with 3-5 sampling points in each block. Sampling blocks were distributed across the sites and for the majority of vegetation types data from at least two spatially separate (i.e. > 100 m apart) blocks were collected. As the dry lichen subsiding vegetation type was not common, this vegetation was sampled less than the other classes. To facilitate the collection of spatially distributed data from the common vegetation types, we selected sites with relatively easy access (e.g. sampling of water logged areas were made in parts of the sites accessible by boardwalks). This approach allowed us to collect CH_4 flux data from a large number of sampling points

135

140

with the aim to generate scalable data from each vegetation type. A total number of 502 measurements were collected across
145 the three sites and the three dates, with all vegetation types measured in several locations at each site. Specifically, on the
raised palsa and subsiding palsa areas, the total number of measurements were 47 and 9 for dry lichen (DL) and dry lichen
subsiding (DLS), respectively, 42 and 68 for dwarf shrub (DS) and dwarf shrub subsiding (DSS), respectively, 50 and 56 moist
moss (MM) and moist moss subsiding (MMS), respectively. In the flooded fen vegetation areas, 102, 78 and 50 measurement
150 were made in *Sphagnum* wetland, sedge wetland and willow wetland, respectively. Detailed description of the vegetation types
are in de la Barreda Bautista et al. (2022). Additional measurements of soil temperature and moisture using a Theta probe
connected to a Theta meter (Delta-T devices, Cambridge, UK) were carried out in each point where methane emissions were
measured. The active layer depth was determined using a graded metal pole in a subset of plots (ca 10 of each vegetation type
in each of the three mires in June, July and Sept 2021 (de la Barreda-Bautista et al., 2022).

155 2.3 CH₄ emissions measurements

Methane flux measurements were conducted using a dynamic closed chamber connected to a Los Gatos ultraportable GHG
analyser (San Jose, California, USA). The Los Gatos analyser uses a cavity ring laser spectroscopy system for detecting CH₄
and affords detection of CH₄ in the field. It also has the advantage of being stable over time and not requiring regular
calibration. Two types of chambers were used, one 15 and one 40 cm tall, the diameter was 40 cm. The chambers were equipped
160 with two push-fit connectors that acted as an inlet and outlet allowing the gas to flow from the chamber to the analyser and
then to flow back into the chamber. After the chamber was placed on the measuring point, it was gently pushed on a 5 cm deep
groove in the peat to create an airtight headspace between the ground surface and the chamber walls. To reduce problems with
bubbles being released into the chamber the chamber was first placed on the ground and then lifted and allowed to vent. The
chambers were then placed back on the ground and the sampling tubes connected. The soil CH₄ concentration (in ppm)
165 increment inside the chamber was measured during 4 to 6 minutes, with data automatically recorded every 20 seconds. If the
change in CH₄ concentrations was not linear, the measurement was stopped, chambers lifted from the soil and a new
measurement was taken after CH₄ readings had stabilized.

Trace gas concentrations, determined by the Los Gatos instrument, were converted to mass units using the ideal gas
law (Eq. 1). Thereafter, using of the slope of the linear change in gas concentration over time and Eq. 2, CH₄ fluxes were
170 calculated. Positive fluxes indicated GHG emissions from the soil into the atmosphere and negative fluxes indicated and uptake
of atmospheric GHG by the soil.

$$n = \frac{PV}{RT} \quad (1)$$

$$175 \quad F = \frac{\text{Slope} \times \text{Volume}_{\text{chamber}}}{\text{Area}_{\text{chamber}}} \quad (2)$$

where, n is the number moles of trace gas (mol L^{-1}), P is the atmospheric pressure (Pa), V is volume of trace gas per litre of air (L L^{-1}), R is the gas law constant ($8314.46 \text{ L}\cdot\text{Pa}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$), and T is the temperature (K). F is the calculated flux for either CH_4 or N_2O ($\text{mol m}^{-2} \text{ hr}^{-1}$), $Slope$ is the slope of the linear regression ($\text{mol L}^{-1} \text{ hr}^{-1}$), $Volume_{chamber}$ is chamber headspace volume (L) and $Area_{chamber}$ is the area of the chamber (m^2). Fluxes were converted to $\mu\text{g m}^{-2} \text{ h}^{-1}$ using the molecular weight of CH_4 (i.e. 16.04 g mol^{-1} , respectively). To avoid excluding very small CH_4 fluxes from the seasonal calculations, fluxes with low r^2 were used in the calculation of cumulative emissions. By contrast, large CH_4 fluxes with $r^2 < 0.7$ were discarded as these fluxes were considered to be either affected by CH_4 ebullition, gas leakage or gas under pressured during transportation of vials. Seasonal fluxes were calculated to the thaw period (i.e. spring-autumn) by scaling the CH_4 flux measurements made in June, July and September to the length of the spring season, summer season and autumn periods, respectively, and then summing these. The length of the three seasons follows a rapid transition due to the high latitude. A typical range was estimated using a combination of seasonal NDVI and air temperature data. The NDVI data from the Torneträsk valley is available from a 6 year UAV time-series (partly unpublished), indicating the typical trajectory of rising and falling values during spring and fall of the snow free season and an NDVI peak plateau (summer) (see (Siewert & Olofsson, 2020). Long term air temperature data from Abisko is monitored by the Swedish meteorological and hydrological service (www.smhi.se).

2.4 UAV-captured data for Digital Elevation Models (DEMs) and vegetation mapping

Multispectral and true colour RGB data ultra high resolution (UHR) data were captured across Storflaket and Stordalen in 2020 and from the Tourist St. site in 2021 from ca 106 and 100 m height using a fixed-wing UAV - Sensefly Ebee fitted with a Parrot Sequoia multispectral sensor. Imagery was collected during peak vegetation season in the last week of July and first week of August approximately at noon. The Parrot Sequoia obtained spectral measurements across four bands: Green ($550\text{nm} \pm 40\text{nm}$), Red ($660\text{nm} \pm 40\text{nm}$), Red Edge ($735\text{nm} \pm 10\text{nm}$) and Near Infrared ($790\text{nm} \pm 40\text{nm}$). The images, ranging between ~1300–2000 per flight, were processed using Pix4D (Berlin, Germany) to produce an orthomosaic for each spectral band. Radiometric calibration was performed using spectral reflectance targets (Mosaic Mill). This resulted in a ground resolution of ~11-13 cm for the multispectral and 2-3 cm for the RGB data. To ensure the orthomosaic and DEM generated from the UAV data were accurately geolocated we collected ca 50 ground control points using a differential GPS (dGPS; Trimble R8s) across all three sites. Further details on the UAV data collection are in de la Barreda Bautista et al. (2022).

The multispectral UAV data was resampled to 0.5m spatial resolution and then classified to produce a vegetation type map for the three study sites using the same nine classes (i.e. the dominant vegetation types found on raised palsa plateaux's, dry lichen, dwarf shrub and moist moss vegetation and areas covered by those vegetation types impacted by subsidence, plus three vegetation types which dominate the flooded areas: *Sphagnum* moss, sedge and willow wetland) as those for which *in situ* CH_4 flux data were measured. A Support Vector Machine classification approach (Foody & Mathur, 2004) was used in ArcMap 10.4 (ESRI) using the Train Support Vector Machine Classifier in the Spatial Analyst package, with simple random sampling of ground data in June 2021 for Storflaket ($n = 258$) and Stordalen ($n = 395$) sites and August 2021 for the Tourist Station ($n = 657$). The ground points from each site were randomly split into training and validation datasets of 70% and 30%

proportions. A cutoff of 70% was used to determine a sufficient accuracy for the vegetation mapping. We decided on this threshold because it is high enough to ensure good overall accuracy, whilst not so high to account for the difficulty of mapping classes with very similar spectral signatures e.g. non-subsiding vegetation classes and their subsiding counterparts. Furthermore, it is also important to note that ground data are never perfect. Indeed, the level of error can be quite large even when arising from authoritative sources. For example, trained aerial photograph interpreters are known to differ in their labelling of forest classes by up to 30% in some studies (Powell et al., 2004).

2.5 Sentinel-2 remotely sensed data for vegetation mapping

We used a neural network classification approach (Xie et al., 2008) within R using the ‘neuralnet’ package (Frtisch et al., 2022) to predict vegetation cover over the three sites of interest using Sentinel-2 data (wavebands B2, B3, B4, B5, B6, B7, B8a, B11 and B12) and elevation, slope and roughness data derived from the ArcticDEM, optical-stereo imagery model at 2 m spatial resolution (Morin et al., 2016). Cloud-free Sentinel-2 reflectance data from 27/07/2019 was used to produce vegetation maps for the sites of interest. Reflectance data was atmospherically corrected using Sen2Cor in ESA SNAP. Only one image was required to cover the wider study domain. To match the 20 m × 20 m resolution of the Sentinel-2 pixels, an average elevation was computed for each pixel. Sentinel-2 wavebands 2, 3 and 4 were re-sampled from 10 m to 20 m resolution using nearest neighbour interpolation to match the resolution of the other wavebands. A Sentinel-2 waveband specification is presented in Supplementary information 1. The UAV-derived vegetation maps were used to train the Sentinel-2 scenes covering the three study sites of the Tourist Station, Storflaket and Stordalen - for this the UAV data was upscaled to 20 m × 20 m using majority classification, this yielded a dataset of 2376 pixels. Training and test datasets were created from this. The testing set used was generated as a random stratified sample comprising 50 pixels per vegetation class, unless there was less than that (e.g., the willow wetland class had only 20 test pixels).

The neural network approach was used to classify the Sentinel-2 data into vegetation classes from which the total areal extents of each vegetation class across the three sites were calculated. We chose a neural network approach for this purpose because of its ability to build both a hard and soft classification and is easily implemented in R with rapid processing. Specifically, a multi-layer perceptron network with backpropagation and weight backtracking was used, therefore the ‘learning rate’ and ‘momentum’ parameters were not required (Xie et al., 2008). Two neural networks were generated: one that produced a standard hard classification (i.e., a single class label is produced for each pixel) and one that produced a soft classification (i.e., grades of class membership is produced for each pixel). The optimal model structure for the majority and the soft classification approaches consisted of twelve principal components, normalised between 0-1, one hidden layer with three neurons (hard classification model) or ten neurons (soft classification model) and an output layer of fifteen units corresponding to the land cover types identified within the three study sites. Both models were developed through trial and error and iteration to arrive at the maximum overall accuracy. The soft classification was undertaken as it enables the estimation of the proportion of different vegetation types within each pixel offering greater understanding of the area covered by each class, in particular to capture and map those smaller patches of vegetation that have a larger contribution to CH₄ flux.

245 Misclassification errors in the maps may substantially bias estimates of class areal extent generated from the maps. An adjustment for misclassification bias in a map can be made if its accuracy has been rigorously evaluated. Here, good practice methods for accuracy and area estimation (Olofsson et al., 2014; Olofsson et al., 2013) were followed to produce misclassification bias-corrected estimates of class area; a bias-correction for the soft classification would ideally require more detailed ground reference data than was available. Thus in total, three vegetation type outputs (maps and class areal extents) 250 from the Sentinel-2 data were produced (soft classification; hard classification and misclassification bias-corrected hard classification). The same 70 % accuracy cutoff as described under the UAV mapping section was applied also to the Sentinel-2 datasets. The areal estimates of the vegetation classes were subsequently multiplied with the mean CH₄ fluxes measured for each vegetation class to produce CH₄ budgets for the entire mapped area.

255

2.6 Surface motion from Sentinel-1 InSAR

We used C-band SAR data from the Sentinel-1 satellites (European Union's Copernicus Programme) with a wavelength and frequency of 5.55 cm and 5.41 GHz respectively to map multi-annual average velocity, subsequently denoted surface motion of the study areas. Details of this approach are in de la Barreda Bautista et al, 2022. Briefly we gathered images from 2017 to 260 2020, focusing on the thaw season between April and October for three separate years. A total of 125 images were acquired from the descending mode and surface motion was obtained using the APSIS method (formerly known as the Intermittent Small Baseline Subset (ISBAS)), which relaxes the requirement for consistent phase stability for all observations using a coherence threshold of 0.45. Following InSAR processing the ground resolution was 20 m x 20 m. Using a stable reference point located in the centre of Abisko village first the multi-annual average velocity (or average surface displacement) between 265 2017 and 2020 (with negative values representing subsidence) was calculated and second, for each season, a time series of surface motion was produced, showing the change in relative height of the surface. Surface motion range for the 2017 and 2020 thaw periods were calculated by subtracting the minimum surface motion value from the maximum surface motion value within respective time series between June and October. The InSAR data were thereafter compared to the CH₄ flux data as described in the statistical analysis section below.

270

2.7 Data analysis

We used residual maximum likelihood (REML) mixed models to test for significant differences in CH₄ fluxes among sites and vegetation and their interactions. For this analysis we used vegetation type as the fixed effect and sampling location (i.e. block), month and site as random effects. The residuals were inspected to ensure the normality assumption of the analysis was met. 275 The standard error of the difference was used to evaluate which vegetation types that differed from each other. We tested for the relationships between the UAV predicted CH₄ budgets per vegetation type and the three different Sentinel-2 based classifications using linear regression analysis. Relationships between ASPIS-InSAR linear motion between the 2017 and 2020 thaw periods and seasonal CH₄ fluxes were explored using REML in which the ground motion categorised into 2.5 mm ranges

were used as the fixed effects and site as the random effect. For this analysis comparison, the CH₄ fluxes extrapolated for each target vegetation type to the three sites using the VHR UAV data, which are able to capture the local-scale variation in the vegetation types, were aggregated to the 20×20 m pixel size of the InSAR data.

We also calculated the net emissions of CH₄ across the subsiding parts of the three wetlands (as detected by InSAR). This was done by summing the CH₄ emissions attributed to each subsiding InSAR pixel for all three sites based on the dominant vegetation type (using the UAV vegetation classification) in that pixel. To explore how CH₄ emissions may change as subsidence progresses and results in palsa collapse, waterlogging and vegetation change we multiplied the currently subsiding wetland areas with the seasonal CH₄ emission that was measured in the lower and higher emitting fen vegetation types, respectively to generate two exemplar future emissions scenarios. This “after permafrost loss and palsa collapse” CH₄ emission scenario used the assumption that all of the subsiding area will be covered with a particular fen vegetation type. We want to highlight that this is not a realistic scenario as the fen vegetation that will replace the current vegetation on the raised palsa will be a mixture of vegetation communities suited to permanently waterlogged conditions. Hence the calculations of future CH₄ emissions should be viewed as illustrations of the ranges of emissions that could occur from the sites following full transition to a fen vegetation type based on their current day emissions.


3 Results

295

3.1 Variation in CH₄ fluxes linked to vegetation type and permafrost degradation


The intact raised palsa vegetation types (DL, DS and MM) had generally low CH₄ fluxes; the dry lichen vegetation type even acted as a weak CH₄ sink in spring (Figure 2a). The areas identified as showing signs of permafrost degradation/subsidence had higher soil moisture content ($F_{8,489} = 167.91$, $P < 0.001$; $SED = 3.6\%$; Figure 3a) and deeper active layer depths ($F_{7,163} = 27$, $P < 0.001$ and $F_{8,91} = 8.15$, $P < 0.001$, for July and September, respectively; Figure 4) than areas that were less affected by degradation. Degrading areas had elevated CH₄ fluxes compared to the corresponding vegetation types in areas that did not display visual signs of degradation ($F_{8,489} = 8.15$, $P < 0.001$; $SED = 2.85 \text{ mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$), with the largest increase in emissions being for subsiding areas of dwarf shrub (Fig 2). The emission from the subsiding palsa areas was highest in summer corresponding to the period of highest air and soil temperatures (Fig 2 b, Fig 3). The highest emissions were from the flooded fen vegetation types, where emissions peaked in autumn (Fig 2 c). Soil temperature peaked in July at the non-flooded sites while at flooded sites temperatures remained relatively constant after the initial thaw period. The coldest soil temperature were measured in DS and DSS areas in September ($F_{8,490} = 9.06$, $P < 0.001$; $SED = 0.7^\circ\text{C}$).

305




310 **Figure 2.** CH₄ fluxes measured in different vegetation types in a) June, b) July, c) September and d) cumulative seasonal CH₄ emissions, “sub.” denotes measurements made in subsiding areas of vegetation types found on the raised palsa plateaus. For September, no data are available from the dry lichen sub. class. Means and SE are shown.

315 Integrated CH₄ fluxes per vegetation type showed highest seasonal fluxes of ca 400 kg CH₄ ha⁻¹ season⁻¹ from willow wetland areas with emissions from *Sphagnum* and sedge wetland vegetation types being ca half those of the willow areas at ca 200 kg CH₄ ha⁻¹ season⁻¹. The seasonal emissions from the subsiding dwarf shrub, moist moss, and dry lichen areas were 50 and 25 kg of CH₄ ha⁻¹ season⁻¹, respectively, which were higher than in un-degraded areas where emission from raised palsa vegetation types were low at less and/or equal to 2 kg of CH₄ ha⁻¹ season⁻¹.



320 **Figure 3. Variation in a) moisture and b) temperature among vegetation types measured in June, July and September, “sub.” denotes measurements made in subsiding areas of vegetation types found on the raised palsa plateaus. For September no data is available from the dry lichen sub. class. Means and SE are shown.**



325 **Figure 4. Variation in active layer depth among vegetation types in a) July and b) September, “sub.” denotes measurements made in subsiding areas of vegetation types found on the raised palsa plateaus. Medians, 25 and 75 percentiles are shown. The whiskers show minimum and maximum values excluding outliers, the black dots represents outliers.**

330 3.2 Scaling CH₄ fluxes via vegetation mapping and calculation of class areal extent

The mapping of the nine vegetation classes found on degrading palsa mires using the VHR UAV data gave an overall classification accuracy of 71% (confusion matrices were generated in base R; Supplementary information 2). The area covered by the target wetland vegetation classes (i.e. excluding rock outcrops, birch forest areas, water bodies and anthropogenic surfaces) across the three sites was ca. 50 ha while the total surveyed area was 83.7 ha (Table 1). Subsequently % estimates of areas are made relative to the target vegetation area. Areas where the visual inspection of the surface topography in the field clearly indicated advanced subsidence equated to 10.0 % of the area covered by the vegetation types. The three subsiding vegetation classes covered 1.5, 2.4 and 1.1 ha, or 3.0, 4.8 and 2.2 % of the target vegetation type area, for dry lichen subsiding, dwarf shrub subsiding and moist moss subsiding, respectively (Table 1). Key here was that the classification was of a high enough accuracy for this output to be used as the “ground” data for the Sentinel-2 classifications. We considered 70% overall

340 accuracy as the threshold for suitable classification accuracy for vegetation classifications. This threshold was decided upon
 because it is high enough to ensure good overall accuracy, whilst not so high to account for the difficulty of mapping classes
 with very similar spectral signatures, for example, vegetation classes compared with their subsiding counterparts.

345 **Table 1. Area of different wetland vegetation types at each of the three study sites and their proportional (Prop.)
 contribution to the area covered by the target vegetation types. (Please note that area of anthropogenic, exposed rock,
 sparse birch forest on mineral soil, forested wetland and water as well as pixels impacted by shadows are not in the
 CH₄ flux study and hence they are not shown in the table).**

Vegetation	Tourist St.		Storflaket		Stordalen		Σ Sites (ha)
	Area (ha)	Prop (%)	Area (ha)	Prop (%)	Area (ha)	Prop (%)	
Dry lichen	0.002	0.1	0.177	1.0	0.245	0.8	0.4
Dry lichen sub	0.004	0.1	1.483	8.3	0.012	0.0	1.5
Dwarf shrub	0.494	15.0	0.897	5.0	1.785	6.1	3.2
Dwarf shrub sub	0.681	20.6	0.450	2.5	1.291	4.4	2.4
Moist moss	0.144	4.4	4.861	27.3	5.612	19.3	10.6
Moist moss sub	0.403	12.2	0.143	0.8	0.571	2.0	1.1
<i>Sphagnum</i>	0.827	25.1	2.250	12.7	7.072	24.3	10.1
Sedge	0.351	10.6	4.835	27.2	5.987	20.6	11.2
Willow	0.392	11.9	2.681	15.1	6.523	22.4	9.6
ΣTarget vegetation	3.3		17.8		29.1		50.2

350

The resultant classification vegetation maps from the Sentinel-2 hard and soft classifications had overall accuracies of 50.2%
 and 83.2%, respectively. The class areal extents from these classification outputs, along with class areal extents obtained when
 hard classification output was corrected for misclassification bias, are presented in Table 2 (which depicts those classes shown
 in Table 1). The hard classification of Sentinel-2 data did not capture the fine-scale vegetation mosaic on the raised part of the
 355 palsa mire (i.e., dry lichen, and dwarf shrub prior to subsidence), instead it mapped the vegetation types in those areas as either
 moist moss or dwarf shrub subsiding vegetation classes, leading to the lower overall classification accuracy than obtained

using the soft classification approach. Once corrected for misclassification bias, the hard classification areal extents for each vegetation class improved considerably in that the estimates are comparable to that mapped using the UAV data and the soft classification of the Sentinel-2 data. The soft classification of the Sentinel-2 data was able to map all nine target vegetation types and followed the same distribution of more or less abundant vegetation types, predicting similar total class areal extents as that of the UAV-derived maps. In both cases (bias corrected hard classification and soft classification), those vegetation types on the raised palsa area missed by the hard classification, have a calculated areal extent.

The soft classification of the Sentinel-2 data produced a higher estimate of the areal extent of the fen vegetation types at 35.4 ha than the UAV classification but succeeded in mapping all the vegetation classes present across the study site. The mapping from the hard classification of Sentinel-2 under-estimated the fen vegetation types slightly at 28.64 and missed some of the other vegetation classes (i.e., Dry lichen, Dry lichen subsiding and Dwarf shrub). After correcting for classification bias, the areal extents of vegetation classes from the hard classification closely matched to those of the UAV mapped areas, with 32.21 ha of the fen vegetation types estimated to occupy the study site.

Based on the UAV-classification vegetation mapping (class areal extents), 30.9 ha (or 61%) of the area covered by the target vegetation types was comprised of the three vegetation classes (*Sphagnum*, sedge and willow wetland – known as fen vegetation types) associated with the highest CH₄ emissions. Using the mapping of areal extents from the UAV data to model CH₄ fluxes shows marked differences in CH₄ fluxes between raised palsas and the wetland types surrounding these (Figure 5; Table 3). Areas of the raised palsa, which are not showing visual signs of subsidence, generally have very low emissions while areas with degrading permafrost and subsidence, as well as those areas with no permafrost emit CH₄. Among the vegetation types that are associated with high CH₄ emissions, the maps highlight the importance of the willow wetland vegetation type as a source of methane from these wetlands. This is because it was both high emissions per area and covers substantial areas in particular at the Stordalen mire site. Although the highest CH₄ emissions are from the areas which have lost their permafrost that are surrounding the remaining palsa, there are indications of increased emissions also within the palsas.

Table 2. Comparing class areal estimates for the target vegetation types using Sentinel-2 (S-2) hard and soft classification and bias-corrected hard classification. For comparison the estimates are provided for the UAV-derived classification which served as the ground data.

	S-2 Hard Classification	S-2 Soft Classification	Bias-corrected S-2 Hard Classification	Ground data (UAV Classification)
Vegetation type	Area (ha)	Area (ha)	Area (ha)	Area (ha)
Dry lichen	0.00	0.4	0.0	0.4
Dry lichen sub	0.00	1.2	0.8	1.5

Dwarf shrub	0.00	3.3	1.1	3.2
Dwarf shrub sub	2.9	2.1	5.3	2.4
Moist moss	13.3	8.1	13.7	10.6
Moist moss sub	0.0	1.3	0.3	1.1
<i>Sphagnum</i>	7.4	10.1	11.2	10.1
Sedge	19.4	13.2	11.7	11.2
Willow	1.8	12.1	9.3	9.6
ΣTarget vegetation	44.8	51.0	52.8	50.2
ΣFen with high CH₄ emissions	28.6	35.4	32.2	30.9

385

Methane emissions from within the palsa areas are commonly associated with subsiding moist moss and subsiding dwarf shrub vegetation, as well as development of *Sphagnum* wetland vegetation in subsiding areas that have become waterlogged either at the edges of the palsas or in their interior following degradation. The subsiding dwarf shrub vegetation is mainly found along the edges of the palsas that are experiencing lateral erosion while the subsiding moist moss vegetation type occurs in the interior parts of the palsa in areas with deeper active layers and the development of a hummocky surface.

390

When CH₄ emissions are integrated across the three wetland areas using the UAV-derived vegetation mapping net emissions are 8519 kg CH₄ season⁻¹ (Table 3). The total emissions are composed of 16.6, 142.6 and 8359.5 kg CH₄ season⁻¹ from raised palsa vegetation (DL, DS, MM), subsiding palsa vegetation (DLS, DSS, MMS) and waterlogged vegetation types (SW, SE WW), respectively. This shows that currently emissions from either intact areas in the early stages of degradation is low compared to the already degraded areas. However, as the areas of raised palsa vegetation and subsiding palsa vegetation are large (ca 20 ha (or 40% of the area covered by target vegetation types) when combined), there are substantial risks of higher CH₄ emissions from these sites associated with continued degradation of the currently subsiding areas and onset of degradation of the remaining palsa areas.

395












Figure 5. (a-c) Detailed vegetation maps separating out wetland vegetation types including vegetation types associated with raised and subsiding palsa areas at the Tourist st., Storflaket and Stordalen mires. The vegetation maps were generated from UAV-captured data. (d-f). Maps of seasonal CH₄ emission based on the seasonal CH₄ emission measured in the nine target vegetation types across the three mire sites. The focus palsa-fen wetland area outlined in purple is the area where ground data on CH₄ fluxes were collected as well as for the vegetation mapping. The full wetland area at each site is indicated by the black boundary line.

Table 3. Seasonal methane emissions for 2021 for the total area of each target vegetation classes within the area covered by the UAV flights. Methane emissions are estimated using three different models using Sentinel-2 data. The CH₄ flux calculated from the UAV data are derived from the area of each vegetation type multiplied with the seasonal CH₄ flux for each vegetation type. Mean and SE are shown.

Vegetation	CH ₄ flux		CH ₄ flux		CH ₄ flux		CH ₄ flux	
	S-2 hard		CH ₄ flux		S-2 bias corrected hard		CH ₄ flux	
	classification		S-2 soft classification		classification		UAV classification	
	(kg CH ₄ season ⁻¹)		(kg CH ₄ season ⁻¹)		(kg CH ₄ season ⁻¹)		(kg CH ₄ season ⁻¹)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Dry lichen	0	0	0.05	0.11	0	0	0.05	0.11
Dry lichen sub	0	0	7.2	3.6	4.7	2.3	8.7	4.3
Dwarf shrub	0	0	7.0	5.5	2.3	1.8	6.7	5.3
Dwarf shrub sub	134.5	41.4	96.7	29.7	248.6	76.4	113.1	34.8
Moist moss	12.3	3.7	7.5	2.2	12.1	3.6	9.8	2.9
Moist moss sub	0.0	0.0	23.4	4.2	4.9	0.9	20.8	3.7
<i>Sphagnum</i>								
wetland	1422.0	400.3	1942.8	547.0	2159.4	607.9	1950.3	549.1
Sedge wetland	4415.3	936.3	2993.5	634.8	2651.8	562.3	2537.7	538.1
Willow wetland	726.2	184.7	4885.9	1242.8	3750.4	953.9	3871.6	984.8
∑ Vegetation types	6710		9964		8834		8519	

Taking the Sentinel-2 sets of estimates of areal extent of vegetation types to estimate the CH₄ emissions of the vegetation types showed both over- and under-estimation of emissions (compared to the UAV-derived CH₄ estimates) - the hard classification under-estimated CH₄ emissions across the site by 21 %; while the soft classification over-estimated emissions by 17% (Table 3). The mismatch between the Sentinel-2 hard classification and the UAV estimates were linked to a misclassification of willow wetland as sedge wetland (Table 2, Supplementary information 2). Correcting the output of the hard classification for


misclassification bias improved the match of CH₄ estimates to that derived from the UAV data, and key was that target vegetation classes were included in the mapping following this post-classification process. For the soft classification, it was the misclassification of sedge and willow wetland area. Correlation analysis between the UAV- and Sentinel-2 classification outputs showed that the bias corrected hard classification was more closely correlated to the UAV predictions, followed closely by the soft classification with $R^2 = 0.99$ and $R^2 = 0.99$ ($P < 0.001$ and $P < 0.001$), respectively. The hard classification was not significantly related to the UAV predicted CH₄ fluxes ($P > 0.05$; Fig 6).



425 **Figure 6. Relationship between UAV estimated methane emissions per vegetation types and the three Sentinel-2 (S-2)**
426 **models. The dashed and solid lines shows the relationships between the soft and bias corrected hard classification and**
427 **the UAV estimated CH₄ emissions, respectively. Note that no regression line was plotted for the hard classification as**
428 **there was no significant relationship.**

430 3.3 Linking InSAR determined ground motion to CH₄ emissions


InSAR detected subsidence across 90 % of the study area, indicating that the majority of the surface of these sites were impacted by mm level rates of subsidence. However, rates of ground motion varied considerably across the study area and among vegetation types. Importantly, across the three sites, areas with relatively low levels of ground motion were associated with the highest mean CH₄ emissions ($F_{8,1135} = 5.63$, $p < 0.001$, $SED = 961.5$; Fig 7) while both uplifting (5 to 7.5 mm) and the most subsiding (-10 to -12.5 mm) areas tended to have lower fluxes. High cumulative CH₄ emissions (summed over all pixels for a given range of ground motions) were found for areas that were either stable or subsiding at an intermediate rate (Fig 8) while the most subsiding areas had low cumulative CH₄ emissions.



440 **Figure 7. Average seasonal CH₄ emissions for different ground motion ranges (categories in 2.5 mm ranges) across the three study sites. Negative values indicate subsidence and positive values indicate uplift. Mean and SE are shown.**

This pattern of areas of high subsidence being linked to areas of low CH₄ emissions was also apparent in the subsidence and CH₄ emissions maps (Fig 9). For example, at the Tourist St. mire areas of high subsidence (comprising mainly subsiding moist moss vegetation Fig 5 a) corresponded to areas with relatively low CH₄ emissions while areas that have already transitioned to *Sphagnum* or willow wetland are associated with bands of high CH₄ emissions along the degrading edge of the palsa plateaux (Fig 5 a and d). Similar patterns were evident on the Storflaket mire, where degrading areas within the palsa plateaux did not display high CH₄ emissions while areas that have transitions to *Sphagnum* or sedge wetland have high CH₄ emissions (Fig 9 b

and e). The Stordalen site showed the largest areas of high subsidence that traversed the boundary between the palsa plateau and the fen areas (Fig 9 c and f). In this part of Stordalen the *Sphagnum* wetland vegetation type was present both in the hollows in the degrading interior of the palsa plateaux and in adjacent areas where the palsas had already collapsed, explaining the high CH₄ emissions (together with the presence of the willow wetland vegetation type) from this area.




455 **Figure 8. Cumulative CH₄ emission (\pm SE) for different ground motions across the thaw seasons of 2017 and 2020 for pixels (i.e. a sum of the CH₄ emissions from all pixel of a given ground motion (categories in 2.5 mm ranges)) of the target vegetation classes. The dry lichen vegetation types did not cover large enough areas to be evaluated on the 20x20 m pixel scale of the ground motion data and is therefore not included in this comparison. Negative values indicate subsidence and positive values indicate uplift.**


460

The net emissions of CH₄ across subsiding parts of the three wetlands (as detected by InSAR) were modest at 116 kg CH₄ season⁻¹. However, when these areas have degraded to such a point that they are fully waterlogged and have transitioned to either *Sphagnum* or sedge vegetation we estimate emissions from these, currently low emitting, areas to be around 6480 kg season⁻¹ or if we use the highest emitting vegetation type as a predictor of future emissions we arrive at 12 960 kg season⁻¹.


465







Tourist st.



Storflaket



Stordalen



470 **Figure 9. Maps of (a-c) ground motion (de la Bautista Barred et al., 2022) and (d-f) CH₄ fluxes at the Toursist st., Storflaket, and Stordalen mire. The black boxes show the areas referred to in the text. The backgrounds to panel a, b and c are orthophotos from the area taken in 2018.**

4 Discussion

475 Methane emissions varied strongly across the palsa peatlands with consistent differences among vegetation types over space and time. The low emissions from the intact dry palsa areas and trends of increased emission following degradation and flooding were comparable with those reported from other part of Scandinavia and Russia e.g. -1.7 to 1.2 kg CH₄ ha⁻¹ season⁻¹ from dry lichen and raised palsa, respectively (Glagolev et al., 2011; Liebner et al., 2015; Miglovets et al., 2021; Nykänen et al., 2003; Olefeldt et al., 2013; Varner et al., 2022). However, there was strong variation in the magnitude of that response depending on the degradation stage, level of flooding and vegetation community present. For example, as long as sites were 480 mainly impacted by increased waterlogging but no vegetation change (i.e. the raised palsa vegetation persisted despite experiencing subsidence), CH₄ emissions only increased modestly (Fig 2). This suggest that waterlogging *per se* was not sufficient to drive high emissions as these were only recorded following flooding and a transition of the vegetation community to *Sphagnum* mosses, sedges and other wetland adapted species.

485 An important finding in our study is the high CH₄ emissions reported from fen and in particular the willow vegetation type. These emissions were consistent across the three study sites and were at the upper range of emissions reported for permafrost fens in the literature e.g. maximum fluxes of ca. 130 to 360 kg CH₄ ha⁻¹ and season⁻¹ (Miglovets et al., 2021; Olefeldt et al., 2013) and in the range of emissions from ponds and thermokarst lakes in degrading permafrost in the literature (Dzyuban, 2002; Glagolev et al., 2011; Walter et al., 2006). Possible explanations for this variation among vegetation types and very high CH₄ emissions are the impact of the vegetation on CH₄ emission both via plant mediated transport, root exudate 490 quality and quantity and the level of rhizosphere oxidation. For example, both *M. trifoliata* and in particular *C. rupestris* which occur in the fen vegetation communities are known to transport CH₄ through their vascular tissues while other species common in our system e.g. *S. lapponica* do not share this trait (Ge, 2022). An additional way in which plants may contribute to high CH₄ production and emissions are by releasing labile carbon in the form of root exudates into the rhizosphere. This carbon flow can represent a substantial part of NPP and can have a large impact on CH₄ emissions (Girkin, 2018; Ström et al., 2003). 495 We speculate that some of the willow vegetation type species found at the study sites, *S. lapponica*, and other *Salix* ssp. and *Comarum palustre*, which were abundant within the willow wetland vegetation type, may drive high emissions via root exudation or those species may reflect specific edaphic conditions that are favourable for CH₄ production. Further, root exudation may be an explanation of high emissions also from the tall sedge vegetation type. However, further studies would be required to establish if the high emissions from the willow wetland vegetation community in our study is consistent across 500 greater spatial ranges as well as the specific drivers of the high emissions. Given the variation in flux rates among waterlogged areas with different vegetation types it is likely that the species composition within the wide vegetation classes often used for

describing permafrost peatlands (raised palsa and fen) directly impact the magnitude of the CH₄ emissions as these peatlands degrade. Hence improved understanding of the plant species specific mechanisms that control CH₄ emissions beyond abiotic drivers (such as water logging, redox, soil and air temperature) (Olefeldt et al., 2013) are important for estimations of landscape level CH₄ emission and how these will be altered as these palsa peatlands continue to degrade and their vegetation composition change.

The fact that small extents and local-scale variation in ground/vegetation conditions strongly impact net emissions (Dias et al., 2010; Ström et al., 2005) is important when it comes to developing spatial models for predicting CH₄ emissions over large areas. Our three models using Sentinel-2 data differed in their ability to detect the local-scale variation in the vegetation. The hard classification model did not succeed in capturing this variability and underestimated the total wetland area. The soft and bias-corrected hard classification was both successful in estimating the overall area of the target vegetation types compared to the UAV estimated areas. When the areal estimates from the three models were used for extrapolating the CH₄ emissions only the bias-corrected hard classification estimated the emissions within a 5% error margin to the UAV estimated fluxes. The hard classification underestimated fluxes by 21 % while the soft classification over estimated emissions with 17%. This means that Sentinel-2 (or other freely available satellite data products e.g. Landsat) have sufficient spatial resolution to be an effective tools for scaling up CH₄ emissions linked to particular vegetation types in the local scale vegetation mosaic that composes degrading peatland. However, among the three Sentinel-2 models we used, the bias-corrected hard classification clearly outperformed the other two models. Both, the uncorrected hard or soft classification, would have included considerable error when used to extrapolate the emissions to regional scale due to scale-dependency leading to inaccuracy in the relative distribution of individual land cover classes at coarser resolutions (Hugelius, 2012; Siewert, 2018). This highlights both the risks of substantially increasing the error of the modelled estimates and the opportunity offered by the bias-corrected hard classification approach for scaling of CH₄ emissions using remotes sensing data (Olofsson et al., 2014; Olofsson et al., 2013). The bias-correction methods used in this study are simple, effective and transferable, however, there is a need for ground data if these approaches are applied to new areas. Spatially distributed high-resolution studies and data collection in remote areas such as the Canadian Arctic and Siberia are in the context of permafrost degradation and CH₄ feedbacks of high value and necessary to calibrate regional to Arctic estimates due to high spatial variability of tundra terrain (Siewert et al., 2021; Siewert & Olofsson, 2020).

The implication of this is that current estimates of changes in GHG emissions following permafrost degradation that often only use the distinction between palsa and fen (e.g. (Miglovetts et al., 2021; Nykänen et al., 2003; Olefeldt et al., 2013; Varner et al., 2022) may result in considerable uncertainty and substantial underestimation of CH₄ emissions linked to degradation. This is because although this simple classification will reflect the major shift from dry to wet conditions and the large increase in CH₄ emissions that are associated with this, hot spots of high emissions will be missed as well as variation in CH₄ emissions among different fen vegetation types. The role of different vegetation types is evident from our CH₄ data from a range of wetland vegetation communities found at the study sites. These shows that variation in CH₄ emissions among common vegetation types within the “fen” vegetation class can result in nearly twice as high emissions on an areal basis. Here

the detailed vegetation mapping capabilities of drones provide an important way forward together with higher resolution aerial photography and satellite data e.g. WorldView, Quickbird, and Planet Ldt's "Super Dove" mission when upscaling plot data to the landscape scale (Siewert et al., 2015; Siewert & Olofsson, 2020).

540 The relatively low CH₄ emissions from areas with high subsidence (Fig 7-9) reflect both the fact that they (i) represent the permafrost degradation front, which is spatially less extensive than already degraded or stable areas, and (ii) that although the subsiding areas of a given vegetation type (Fig 2) release more CH₄ than their intact counter parts, emissions are low compared to areas which are fully degraded and flooded. Further, although CH₄ emissions increased as a specific vegetation types showed signs of permafrost degradation and subsidence (Fig 2) the initial increase in CH₄ emissions at the most subsiding sites is modest compared to the high CH₄ emissions in areas where degradation has progressed to a point where permafrost has
545 thawed all together and subsidence has slowed down (albeit some consolidation of the peat material is still ongoing as suggested by subsidence rates at a lower rate in areas which have lost their permafrost).

Our work suggest that subsidence measured using ASPIS-InSAR can be used to predict increases in CH₄ emissions from areas experiencing permafrost degradation. The InSAR method also offers a highly sensitive early warning system of the initial stages of degradation. This is valuable because early stages of permafrost degradation are difficult to capture using
550 optical data or ground surveys, which tend to rely on changes in the vegetation or increased surface heterogeneity which only become apparent after the degradation has progressed substantially. This is evident in the different areal estimates of the subsiding area using the two different vegetation mapping approaches using UAV and Sentinel 2 data. Mapping for subsiding areas using ground truthing of VHR UAV optical data estimated the subsiding area of the study sites to ca 10% while the InSAR data detected subsidence in 90% of the wetland area. The implication of the InSAR data is that the majority of the palsa
555 peatlands area at our study sites is already experiencing permafrost degradation. Indeed, ongoing large-scale permafrost degradation at the sites is further supported by analysis of historical orthophotos and long term satellite data analysis as well as model predictions that suggest that permafrost in wetland will no longer be supported in northern Scandinavia by 2050 (de la Barreda-Bautista et al., 2022; Fewster et al., 2022; Varner et al., 2022). This raises concerns for the fate of the ca 5500 ha of palsa peatlands in Sweden over the next decades (Backe, 2014). The evidence of large-scale degradation at our study sites
560 and the potential for very high CH₄ emissions from areas that currently have low emissions are a serious risk linked to climate heating in northern Scandinavia and highlights the need for regional estimates of the potential for increased CH₄ emissions associated with degradation of palsa peatlands.

Acknowledgements

565

We are grateful for the field work support from Mattias Dalkvist, Veronica Escubar Ruiz, and staff at Abisko Scientific Research Station. We also acknowledge that the climate data evaluation has been made by data provided by Abisko Scientific Research Station and the Swedish Infrastructure for Ecosystem Science (SITES). We are grateful to Samuel Valman for advice on data processing.

Supplementary information

Supplementary information 1. Sentinel-2 data specification. Wavebands listed with a spatial resolution of 10 m were resampled to 20 m to match other wavebands.

Supplementary information 2. Excel file containing confusion matrixes for the hard and soft classifications.

575

References

- Åkerman, H. J., & Johansson, M. (2008). Thawing permafrost and thicker active layers in sub-arctic Sweden
[\[https://doi.org/10.1002/ppp.626\]](https://doi.org/10.1002/ppp.626). *Permafrost and Periglacial Processes*, 19(3), 279-292.
<https://doi.org/https://doi.org/10.1002/ppp.626>
- 580 Alshammari, L., Large, D. J., Boyd, D. S., Sowter, A., Anderson, R., Andersen, R., & Marsh, S. (2018). Long-Term Peatland
 Condition Assessment via Surface Motion Monitoring Using the ISBAS DInSAR Technique over the Flow
 Country, Scotland. *Remote Sensing*, 10(7). <https://doi.org/10.3390/rs10071103>
- Backe, S. (2014). *Kartering av Sveriges palsmyrar*. Länsstyrelsen.
- 585 Ballantyne C. K. (2018). *Periglacial geomorphology*. John Wiley & Sons.
- Bernhardt, E. S., Blaszcak, J. R., Ficken, C. D., Fork, M. L., Kaiser, K. E., & Seybold, E. C. (2017). Control Points in
 Ecosystems: Moving Beyond the Hot Spot Hot Moment Concept. *Ecosystems*, 20(4), 665-682.
<https://doi.org/10.1007/s10021-016-0103-y>
- Biskaborn, B. K., Smith, S. L., Noetzi, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P., Romanovsky, V. E.,
 590 Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R., Diekmann,
 B., Drozdov, D., Etzelmüller, B., Grosse, G., . . . Lantuit, H. (2019). Permafrost is warming at a global scale. *Nature
 Communications*, 10(1), 264. <https://doi.org/10.1038/s41467-018-08240-4>
- Borge, A. F., Westermann, S., Solheim, I., & Etzelmüller, B. (2017). Strong degradation of palsas and peat plateaus in
 northern Norway during the last 60 years. *The Cryosphere*, 11(1), 1-16. <https://doi.org/10.5194/tc-11-1-2017>
- 595 Chadburn, S. E., Burke, E. J., Cox, P. M., Friedlingstein, P., Hugelius, G., & Westermann, S. (2017). An observation-based
 constraint on permafrost loss as a function of global warming. *Nature Climate Change*, 7(5), 340-344.
<https://doi.org/10.1038/nclimate3262>
- de la Barrera-Bautista, B., Boyd, D. S., Ledger, M., Siewert, M. B., Chandler, C., Bradley, A. V., Gee, D., Large, D. J.,
 Olofsson, J., Sowter, A., & Sjögersten, S. (2022). Towards a Monitoring Approach for Understanding Permafrost
 600 Degradation and Linked Subsidence in Arctic Peatlands. *Remote Sensing*, 14(3).
- Dias, A. T. C., Hoorens, B., Van Logtestijn, R. S. P., Vermaat, J. E., & Aerts, R. (2010). Plant Species Composition Can Be
 Used as a Proxy to Predict Methane Emissions in Peatland Ecosystems After Land-Use Changes. *Ecosystems*,
 13(4), 526-538. <https://doi.org/10.1007/s10021-010-9338-1>
- Dzyuban, A. (2002). Intensity of the microbiological processes of the methane cycle in different types of Baltic lakes.
 605 *Microbiology*, 71(1), 98-104.
- Euskirchen, E. S., Edgar, C. W., Turetsky, M. R., Waldrop, M. P., & Harden, J. W. (2014). Differential response of carbon
 fluxes to climate in three peatland ecosystems that vary in the presence and stability of permafrost. *Journal of
 Geophysical Research: Biogeosciences*, 119(8), 1576-1595. <https://doi.org/https://doi.org/10.1002/2014JG002683>
- 610 Fewster, R. E., Morris, P. J., Ivanovic, R. F., Swindles, G. T., Peregon, A. M., & Smith, C. J. (2022). Imminent loss of
 climate space for permafrost peatlands in Europe and Western Siberia. *Nature Climate Change*, 12(4), 373-379.
<https://doi.org/10.1038/s41558-022-01296-7>
- Foody, G. M., & Mathur, A. (2004). A relative evaluation of multiclass image classification by support vector machines.
IEEE Transactions on Geoscience and Remote Sensing, 42(6), 1335-1343.
<https://doi.org/10.1109/TGRS.2004.827257>

- 615 Ge, M., Koskinen, M., Korrensalo, A., Mäkiranta, P., Lohila, A., and Pihlatie, M. (2022, 23–27 May 2022). Species-specific effects of vascular plants on methane transport in northern peatlands. EGU General Assembly 2022, Vienna, Austria.
- Girkin, N. T. (2018). *Tropical forest greenhouse gas emissions: root regulation of soil processes and fluxes* University of Nottingham].
- 620 Glagolev, M., Kleptsova, I., Filippov, I., Maksyutov, S., & Machida, T. (2011). Regional methane emission from West Siberia mire landscapes. *Environmental Research Letters*, 6(4), 045214.
- Grosse, G., Goetz, S., McGuire, A. D., Romanovsky, V. E., & Schuur, E. A. G. (2016). Changing permafrost in a warming world and feedbacks to the Earth system. *Environmental Research Letters*, 11(4), 040201.
<https://doi.org/10.1088/1748-9326/11/4/040201>
- 625 Hugelius, G. A.-O., Loisel, J. A.-O., Chadburn, S. A.-O. X., Jackson, R. A.-O., Jones, M. A.-O., MacDonald, G., Marushchak, M., Olefeldt, D. A.-O., Packalen, M., Siewert, M. A.-O., Treat, C. A.-O., Turetsky, M., Voigt, C. A.-O., & Yu, Z. A.-O. (2020). Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. (1091-6490 (Electronic)).
- IPCC. (2021). *The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- 630 Johansson, M., Callaghan, T. V., Bosiö, J., Åkerman, H. J., Jackowicz-Korczynski, M., & Christensen, T. R. (2013). Rapid responses of permafrost and vegetation to experimentally increased snow cover in sub-arctic Sweden. *Environmental Research Letters*, 8(3), 035025. <https://doi.org/10.1088/1748-9326/8/3/035025>
- Liebner, S., Ganzert, L., Kiss, A., Yang, S., Wagner, D., & Svenning, M. M. (2015). Shifts in methanogenic community composition and methane fluxes along the degradation of discontinuous permafrost. *Frontiers in Microbiology*, 6, 356.
- 635 Luoto, M., & Seppälä, M. (2003). Thermokarst ponds as indicators of the former distribution of palsas in Finnish Lapland [<https://doi.org/10.1002/ppp.441>]. *Permafrost and Periglacial Processes*, 14(1), 19-27.
<https://doi.org/https://doi.org/10.1002/ppp.441>
- 640 Maksyutov, S., Glagolev, M., Kleptsova, I., Sabrekov, A., Peregon, A., & Machida, T. (2010). Methane emissions from the West Siberian wetlands. AGU Fall Meeting Abstracts,
- McGuire, A. D., Anderson, L. G., Christensen, T. R., Dallimore, S., Guo, L., Hayes, D. J., Heimann, M., Lorenson, T. D., Macdonald, R. W., & Roulet, N. (2009). Sensitivity of the Carbon Cycle in the Arctic to Climate Change. *Ecological Monographs*, 79(4), 523-555. <http://www.jstor.org/stable/40385226>
- 645 Miglovets, M., Zagirova, S., Goncharova, N., & Mikhailov, O. (2021). Methane Emission from Palsa Mires in Northeastern European Russia. *Russian Meteorology and Hydrology*, 46(1), 52-59.
- Mishra, U., Hugelius, G., Shelef, E., Yang, Y., Strauss, J., Lupachev, A., Harden Jennifer, W., Jastrow Julie, D., Ping, C.-L., Riley William, J., Schuur Edward, A. G., Matamala, R., Siewert, M., Nave Lucas, E., Koven Charles, D., Fuchs, M., Palmtag, J., Kuhry, P., Treat Claire, C., . . . Orr, A. (2021). Spatial heterogeneity and environmental predictors of permafrost region soil organic carbon stocks. *Science Advances*, 7(9), eaaz5236.
<https://doi.org/10.1126/sciadv.aaz5236>
- 650 Nykänen, H., Heikkinen, J. E., Pirinen, L., Tiilikainen, K., & Martikainen, P. J. (2003). Annual CO₂ exchange and CH₄ fluxes on a subarctic palsa mire during climatically different years. *Global Biogeochemical Cycles*, 17(1).
- Olefeldt, D., Roulet, N. T., Bergeron, O., Crill, P., Bäckstrand, K., & Christensen, T. R. (2012). Net carbon accumulation of a high-latitude permafrost palsa mire similar to permafrost-free peatlands. *Geophysical Research Letters*, 39(3).
- 655 Olefeldt, D., Turetsky, M. R., Crill, P. M., & McGuire, A. D. (2013). Environmental and physical controls on northern terrestrial methane emissions across permafrost zones. *Global Change Biology*, 19(2), 589-603.
- Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., & Wulder, M. A. (2014). Good practices for estimating area and assessing accuracy of land change. *Remote Sensing of Environment*, 148, 42-57.
<https://doi.org/https://doi.org/10.1016/j.rse.2014.02.015>
- 660 Olofsson, P., Foody, G. M., Stehman, S. V., & Woodcock, C. E. (2013). Making better use of accuracy data in land change studies: Estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sensing of Environment*, 129, 122-131. <https://doi.org/https://doi.org/10.1016/j.rse.2012.10.031>

- 665 Olvmo, M., Holmer, B., Thorsson, S., Reese, H., & Lindberg, F. (2020). Sub-arctic palsa degradation and the role of climatic drivers in the largest coherent palsa mire complex in Sweden (Vissátvuopmi), 1955–2016. *Scientific Reports*, 10(1), 8937. <https://doi.org/10.1038/s41598-020-65719-1>
- Sannel, A. B. K., Hugelius, G., Jansson, P., & Kuhry, P. (2016). Permafrost Warming in a Subarctic Peatland – Which Meteorological Controls are Most Important? [<https://doi.org/10.1002/ppp.1862>]. *Permafrost and Periglacial Processes*, 27(2), 177-188. <https://doi.org/https://doi.org/10.1002/ppp.1862>
- 670 Sannel, A. B. K., & Kuhry, P. (2011). Warming-induced destabilization of peat plateau/thermokarst lake complexes. *Journal of Geophysical Research: Biogeosciences*, 116(G3). <https://doi.org/https://doi.org/10.1029/2010JG001635>
- Schneider von Deimling, T., Meinshausen, M., Levermann, A., Huber, V., Frieler, K., Lawrence, D. M., & Brovkin, V. (2012). Estimating the near-surface permafrost-carbon feedback on global warming. *Biogeosciences*, 9(2), 649-665. <https://doi.org/10.5194/bg-9-649-2012>
- 675 Schuur, E. A. G., & Abbott, B. (2011). High risk of permafrost thaw. *Nature*, 480(7375), 32-33. <https://doi.org/10.1038/480032a>
- Siewert, M. B. (2018). High-resolution digital mapping of soil organic carbon in permafrost terrain using machine learning: a case study in a sub-Arctic peatland environment. *Biogeosciences*, 15(6), 1663-1682. <https://doi.org/10.5194/bg-15-1663-2018>
- 680 Siewert, M. B., Hanisch, J., Weiss, N., Kuhry, P., Maximov, T. C., & Hugelius, G. (2015). Comparing carbon storage of Siberian tundra and taiga permafrost ecosystems at very high spatial resolution. *Journal of Geophysical Research: Biogeosciences*, 120(10), 1973-1994. <https://doi.org/https://doi.org/10.1002/2015JG002999>
- Siewert, M. B., Lantuit, H., Richter, A., & Hugelius, G. (2021). Permafrost Causes Unique Fine-Scale Spatial Variability Across Tundra Soils. *Global Biogeochemical Cycles*, 35(3), e2020GB006659. <https://doi.org/https://doi.org/10.1029/2020GB006659>
- 685 Siewert, M. B., & Olofsson, J. (2020). Scale-dependency of Arctic ecosystem properties revealed by UAV. *Environmental Research Letters*, 15(9), 094030. <https://doi.org/10.1088/1748-9326/aba20b>
- Sjöberg, Y., Marklund, P., Pettersson, R., & Lyon, S. W. (2015). Geophysical mapping of palsa peatland permafrost. *The Cryosphere*, 9(2), 465-478. <https://doi.org/10.5194/tc-9-465-2015>
- 690 Sjöberg, Y., Siewert, M. B., Rudy, A. C. A., Paquette, M., Bouchard, F., Malenfant-Lepage, J., & Fritz, M. (2020). Hot trends and impact in permafrost science. *Permafrost and Periglacial Processes*, 31(4), 461-471. <https://doi.org/https://doi.org/10.1002/ppp.2047>
- Sjögersten, S., Caul, S., Daniell, T. J., Jurd, A. P. S., O'Sullivan, O. S., Stapleton, C. S., & Titman, J. J. (2016). Organic matter chemistry controls greenhouse gas emissions from permafrost peatlands. *Soil Biology and Biochemistry*, 98, 42-53. <https://doi.org/https://doi.org/10.1016/j.soilbio.2016.03.016>
- 695 Ström, L., & Christensen, T. R. (2007). Below ground carbon turnover and greenhouse gas exchanges in a sub-arctic wetland. *Soil Biology and Biochemistry*, 39(7), 1689-1698. <https://doi.org/https://doi.org/10.1016/j.soilbio.2007.01.019>
- 700 Ström, L., Ekberg, A., Mastepanov, M., & Røjle Christensen, T. (2003). The effect of vascular plants on carbon turnover and methane emissions from a tundra wetland. *Global Change Biology*, 9(8), 1185-1192. <https://doi.org/https://doi.org/10.1046/j.1365-2486.2003.00655.x>
- Ström, L., Mastepanov, M., & Christensen, T. R. (2005). Species-specific Effects of Vascular Plants on Carbon Turnover and Methane Emissions from Wetlands. *Biogeochemistry*, 75(1), 65-82. <https://doi.org/10.1007/s10533-004-6124-1>
- 705 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., & Zimov, S. (2009). Soil organic carbon pools in the northern circumpolar permafrost region [<https://doi.org/10.1029/2008GB003327>]. *Global Biogeochemical Cycles*, 23(2). <https://doi.org/https://doi.org/10.1029/2008GB003327>
- Turetsky, M. R., Abbott, B. W., Jones, M. C., Anthony, K. W., Olefeldt, D., Schuur, E. A. G., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D. M., Gibson, C., Sannel, A. B. K., & McGuire, A. D. (2020). Carbon release through abrupt permafrost thaw. *Nature Geoscience*, 13(2), 138-143. <https://doi.org/10.1038/s41561-019-0526-0>
- 710 van Huissteden, J., Teshebaeva, K., Cheung, Y., Magnússon, R. Í., Noorbergen, H., Karsanaev, S. V., Maximov, T. C., & Dolman, A. J. (2021). Geomorphology and InSAR-Tracked Surface Displacements in an Ice-Rich Yedoma Landscape [Original Research]. *Frontiers in Earth Science*, 9(724). <https://doi.org/10.3389/feart.2021.680565>

- 715 Varner, R. K., Crill, P. M., Frolking, S., McCalley, C. K., Burke, S. A., Chanton, J. P., Holmes, M. E., null, n., Saleska, S., & Palace, M. W. (2022). Permafrost thaw driven changes in hydrology and vegetation cover increase trace gas emissions and climate forcing in Stordalen Mire from 1970 to 2014. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 380(2215), 20210022. <https://doi.org/10.1098/rsta.2021.0022>
- 720 Walter Anthony, K., Daanen, R., Anthony, P., Schneider von Deimling, T., Ping, C.-L., Chanton, J. P., & Grosse, G. (2016). Methane emissions proportional to permafrost carbon thawed in Arctic lakes since the 1950s. *Nature Geoscience*, 9(9), 679-682. <https://doi.org/10.1038/ngeo2795>
- Walter, K. M., Zimov, S., Chanton, J. P., Verbyla, D., & Chapin, F. S. (2006). Methane bubbling from Siberian thaw lakes as a positive feedback to climate warming. *Nature*, 443(7107), 71-75.
- Xie, Y., Sha, Z., & Yu, M. (2008). Remote sensing imagery in vegetation mapping: a review. *Journal of Plant Ecology*, 1(1), 9-23. <https://doi.org/10.1093/jpe/rtm005>
- 725 Zuidhoff, F. S., & Kolstrup, E. (2000). Changes in palsa distribution in relation to climate change in Laivadalen, northern Sweden, especially 1960–1997 [[https://doi.org/10.1002/\(SICI\)1099-1530\(200001/03\)11:1<55::AID-PPP338>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1099-1530(200001/03)11:1<55::AID-PPP338>3.0.CO;2-T)]. *Permafrost and Periglacial Processes*, 11(1), 55-69. [https://doi.org/https://doi.org/10.1002/\(SICI\)1099-1530\(200001/03\)11:1<55::AID-PPP338>3.0.CO;2-T](https://doi.org/https://doi.org/10.1002/(SICI)1099-1530(200001/03)11:1<55::AID-PPP338>3.0.CO;2-T)
- 730 Hugelius, G. (2012). Spatial upscaling using thematic maps: An analysis of uncertainties in permafrost soil carbon estimates. *Global Biogeochemical Cycles*, 26(2). <https://doi.org/https://doi.org/10.1029/2011GB004154>
- Powell, R. L., Matzke, N., Souza, C., Clark, M., Numata, I., Hess, L., & Roberts, D. (2004). Sources of error in accuracy assessment of thematic land-cover maps in the Brazilian Amazon. *Remote Sensing of Environment*, 90, 221-234. <https://doi.org/10.1016/j.rse.2003.12.007>