

# Assessing $\text{CO}_2$ Carbon dioxide and $\text{CH}_4$ methane fluxes from mounds of African fungus-growing termites

Matti Räsänen<sup>1</sup>, Risto Vesala<sup>2</sup>, Petri Rönnholm<sup>3</sup>, Laura Arppe<sup>2</sup>, Petra Manninen<sup>1</sup>, Markus Jylhä<sup>1</sup>, Jouko Rikkinen<sup>2,4</sup>, Petri Pellikka<sup>1,5</sup> and Janne Rinne<sup>1,5,6</sup>

<sup>1</sup>Department of Geosciences and Geography, University of Helsinki, P.O. Box 64, 00014, Helsinki, Finland

<sup>2</sup>Finnish Museum of Natural History, University of Helsinki, P.O. Box 64, 00014, Helsinki, Finland

<sup>3</sup>Department of Built Environment, Aalto University, P.O. Box 14100, 00076, Aalto, Finland

<sup>4</sup>Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, P.O. Box 64, 00014, Helsinki, Finland

<sup>5</sup>Natural<sup>5</sup>Wangari Maathai Institute for Environmental and Peace Studies, University of Nairobi, P.O. Box 29053, 00625, Kileleshwa, Kenya

<sup>6</sup>Natural Resources Institute Finland, P.O. Box 2, 00791 Helsinki, Finland

15 Correspondence to: Matti Räsänen (matti.rasanen@helsinki.fi)

**Abstract.** Termites play an essential role in decomposing dead plant material in tropical ecosystems and are thus major sources of gaseous C emissions in many environments. In African savannas, fungus-growing termites are among the ecologically most influential termite species. We studied the gas exchange from mounds of two closely related fungus-growing species (*Macrotermes subhyalinus* and *M. michaelseni*, respectively) in ~~shaded and open~~ two habitats representing different vegetation types (grassland, bushland) together with soil fluxes around the mounds. The fluxes from active termite mounds varied from ~~11.6 to 21.0~~ 11.6 to 21.0 mg  $\text{CO}_2\text{-C m}^{-2}\text{h}^{-1}$  for  $\text{CO}_2$  and from 0.06 to 3.667 mg  $\text{CH}_4\text{-C m}^{-2}\text{h}^{-1}$  for  $\text{CH}_4$  fluxes. Mound  $\text{CO}_2$  fluxes varied seasonally with a 64 % decrease and 35 % increase in the fluxes from dry to wet season at the ~~open~~ grassland and ~~shaded~~ bushland sites, respectively. During the wet season, the  $\text{CO}_2$  fluxes were significantly correlated with termite mound volume.

25 The diurnal measurements from two *M. michaelseni* mounds suggest that the gas fluxes peak during daytime and midnight ~~supporting the previously presented model of thermally driven air circulation inside the mound and possibly enhanced ventilation during the night by the opening of foraging routes, possibly reflecting movement of termites between the mounds and surrounding foraging areas and diurnal changes in mound internal air circulation.~~ The stability of the mound gas fluxes over diurnal and annual scales coincides with the constant nature of the nest internal gas and thermal environment that

30 guarantees continuously favorable conditions for the fungal symbiont. Soil fluxes of both  $\text{CO}_2$  and  $\text{CH}_4$  were enhanced at up to 2 m distance from the mounds compared to the local soil respiration, indicating that, in addition to mound ventilation structures, a small proportion of the produced metabolic gases leave the nest also via surrounding soils.

Formatted: Superscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

5

## 1 Introduction

Termites are major degraders of dead organic plant matter in many tropical and subtropical ecosystems (Bignell and Eggleton 2000; Jouquet et al., 2011), also making them a significant source of atmospheric carbon dioxide and methane (Zimmerman et al., 1983; Khalil et al., 1990). Especially in African dry savannas, the fungus-growing species comprise the most common and ecologically important termite group that may recycle a quarter of total dead plant matter and even 90 % of the woody litter in some arid ecosystems (Collins 1981; Buxton 1981). Fungus-growing termites cultivate symbiotic termite fungi (*Termitomyces* spp., Basidiomycota) in fungus combs, which are specialized structures within underground nest chambers. The combs are built from feces of termite workers and consist of largely undigested plant matter, initially ingested by foraging termites, that the *Termitomyces* further degrades (Rouland-Lefèvre 2000; Vesala et al., 2022a). Eventually, termite hosts get their energy and nutrition from the degraded plant matter and/or the fungal mycelium (Sieber and Leuthold, 1981; Hyodo et al., 2003; Vesala et al., 2019a).

Fungus-cultivation within termites is always associated with a complex caste system including numerous workers, soldiers, and reproductive individuals, and the total number of termites within a colony typically ranges from hundreds of thousands to up to a few million individuals (Darlington, 1986, 1990). Such large insect populations are notable sources of CO<sub>2</sub> already in themselves but, within nests of fungus-growing termites, a major proportion of the nest CO<sub>2</sub> emission is produced by the extensive fungus gardens whose total biomass in mature termite colonies may exceed 20 kilograms (Darlington, 1986; Darlington et al., 1997; Noirot and Darlington, 2000). Due to their methanogenic gut symbionts, termites are also among the major sources of atmospheric CH<sub>4</sub> although estimates about their contribution to the global CH<sub>4</sub> budget vary highly (currently considered to represent 1–4 % of the natural sources) due to uncertainties in methane-related processes and global termite biomass (Sanderson, 1996; Kirschke et al., 2013; Saunio et al., 2020). Methane can also be oxidized by soil methanotrophic microbes occurring within and around termite nests (Chiri et al., 2020). It has been evaluated that methane oxidizing bacteria can mitigate even half of the produced methane in nests of some mound-building termite species (Sugimoto et al., 1998; Nauer et al., 2018).

30

Maintenance of large termite populations and the excessive extensive fungal gardens necessitates effective gas-exchange and stable conditions within the nest interior. Termites of the genera *Macrotermes* and *Odontotermes* build aboveground soil

mounds to stabilize temperature, maintain humidity and enhance gas exchange (Noirot and Darlington, 2000; Korb 2003, 2011). Mound structure can vary a lot depending on termite species and environmental conditions (Korb and Linsenmair 1998; Korb 2003, 2011). Even closely related species may rely on fundamentally different mechanisms of nest ventilation, as in case of Kenyan *Macrotermes michaelseni* and *Macrotermes subhyalinus* that build closed and open mounds, respectively  
5 (Darlington 1984, 1985). The open *M. subhyalinus* mounds have multiple large ventilation shafts at different elevations on the mound surface, that promote wind-induced Venturi effect, resulting in airflow through the shafts that ventilate the underground nest (Weir, 1973; Korb 2011). The closed mounds of *M. michaelseni* lack these openings and, instead, the air is circulated internally within the mound until the gas-exchange between the nest interior and ambient air takes place through the porous mound surface (Noirot and Darlington, 2000; Ocko et al., 2017). -Direct measurements of air velocities and temperatures  
10 within closed mounds have shown that air moves in convective cells following a diurnally oscillating thermal schedule (King et al., 2015; Ocko et al., 2017).

Despite the well-recognized importance of termite mounds as significant sources of greenhouse gases, field studies quantifying gas fluxes from termite mounds are rare and often limited to species that build relatively small mounds. Previous CO<sub>2</sub> flux  
15 measurements from termite mounds have ranged from 40 to ~~22462200~~ mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> (van Asperen et al., 2021; Jamali et al., 2013; Brümmer et al., 2009). For wood-feeding termites in Australia, the seasonal variability in CO<sub>2</sub> and CH<sub>4</sub> was high with larger fluxes occurring during the wet season (Jamali et al., 2013). For soil and wood-feeding termites the relationship between mound CO<sub>2</sub> and CH<sub>4</sub> fluxes has been species-specific (Jamali et al., 2013). With the assistance of *Termitomyces*  
20 symbionts, fungus-growing termites can utilize widely different types of plant matter including grasses, leaf litter, and dead wood (Collins 1981; Lepage 1981a) and the proportion of utilized food sources may vary both geographically and seasonally (Boutton et al., 1983; Lepage et al., 1993; Vesala et al., 2022b). Alternative food sources may also differ in their nutritional value with potential consequences to the gas fluxes- ([Zhou et al. 2023](#)). For example, a lower nitrogen content associated with  
25 grass-based compared to woody plant-based fungus combs found in Vesala et al., (2022b) could necessitate relatively higher CO<sub>2</sub> and CH<sub>4</sub> emissions, as more carbon needs to be eliminated to gain sufficiently protein from the N deficient diet (Higashi et al., 1992; Eggleton and Tayasu, 2001). Whether such factors could affect the mound gas fluxes is currently not known.

In this study, we used static chamber method to investigate CO<sub>2</sub> and CH<sub>4</sub> fluxes originating from mounds of two common species of fungus-growing termites, *M. michaelseni* and *M. subhyalinus*, in southern Kenya- during dry and rainy seasons. As, in the area, both species inhabit a wide range of different environments, termite mounds were measured in two contrasting habitats (open grassland and dense bushland). Distribution of the two species is not completely overlapping and, thus, in our  
30 sampling *M. michaelseni* was dominant at the grassland and *M. subhyalinus* at the bushland site. In Kenyan grasslands, *M. michaelseni* mounds are surrounded by dense networks of foraging tunnels with several openings to soil surface per each m<sup>2</sup> (Lepage 1981a; Darlington 1982) that could also increase soil CO<sub>2</sub> and CH<sub>4</sub> fluxes in mound proximity. -Thus, in addition to direct gas fluxes from the mounds, we also measured soil CO<sub>2</sub> and CH<sub>4</sub> fluxes around the mounds. The study objectives were:

(1) to quantify the gas fluxes from the fungus-growing termite mounds with variable mound volumes and from the soil around the mounds, (2) -study the interactions between the fluxes and different environmental variables and to (3) ~~determine~~observe the diurnal and seasonal variation in mound fluxes.

5

Formatted: Indent: Left: -0,03 cm

Formatted: Indent: Left: 0 cm, Space After: 0 pt

Formatted: Indent: Left: 0 cm

## 2 Materials and methods

### 2.1 Site description

10 The chamber measurements were conducted at the lowland areas surrounding Taita Hills, Kenya (Fig. 1). The mean annual rainfall is 433 mm based on measurements at the Maktau weather station (Räsänen et al., 2020). Termite mounds were measured at bushland and grassland locations that are 15 km apart. The bushland is characterized by over 50 % of thorny shrubs and small *Acacia* spp. and *Commiphora* spp. trees (Wachiye et al., 2020). All termite mounds measured in bushland were located in shaded locations. The grassland site is located at a private game sanctuary for wildlife conservation. The  
15 vegetation is characterized by common savanna grasses such as *Chloris roxburghiana* and *Cenchrus ciliaris* and scattered trees (e.g. *Acacia tortilis*, *Albizia anthelmintica*, *Balanites aegyptica*). Majority of bushland mounds were open mounds build by *M. subhyalinus*, whereas most of the grassland mounds were closed type build by *M. michaelseni* (Table 1, Fig. S1). The grassland ecosystem is assumed to be a result ~~from~~of decades long wildlife conservation, which causes high population of mega-herbivore, namely elephants, which forage with leaves of the trees and bushes, thus destroying the tree cover. The tree  
20 cover and above ground biomass is much higher in the bushland areas, which are not inhabited by elephants as shown in Figure 1B (Amara et al., 2020; Pellikka et al., 2018).~~In true color Sentinel satellite image (Figure 1A), the magenta color indicate grasslands, while brownish colors indicate bushlands. Due to difference in habitat, the termites also forage differently in grasslands and bushlands, and thus may cause different fluxes.~~




Figure 1. Locations of the two measurement sites and weather station in Maktau in Sentinel-2A satellite image, 20 December 2020 (A) (the magenta color indicate grasslands, while brownish colors indicate bushlands) and on above ground biomass map (B) (Amara et al., 2020; Pellikka et al., 2018). Chamber measurement at the grassland (C) and the bushland site (D). Metal collar and wireframe around an open *M. subhyalinus* mound (E). Sentinels Scientific Data HubCE4 (ESA, 2015). Photo credits: Petri Pellikka (C), and Markus Jylhä (D, E).

Formatted: Font: Times New Roman, English (United States)

5 **Table 1. Termite mound characteristics. R is the average mound radius, H is height and  $V_m$  is calculated above ground mound volume (see section 2.2). The status (active or dead) of the colonies was determined after the second measurement campaign by excavation.**

Mound id	Termite species (mound type)	LocationSite location	Type	R (m)	H (m)	$V_m$ (m <sup>3</sup> )	Status in April 2017
MR1	<i>M. michaelseni</i> (closed)	Bushland	closed	0.75	0.50	0.256	active
MR2	<i>M. subhyalinus</i> (open)	Bushland	open	0.65	0.15	0.058	active
MR3	<i>M. subhyalinus</i> (open)	Bushland	open	0.28	0.33	0.023	dead
MR4	<i>M. subhyalinus</i> (open)	Bushland	open	0.75	0.45	0.231	active
MR5	<i>M. subhyalinus</i> (open)	Bushland	open	0.30	0.25	0.020	active
MR6	<i>M. subhyalinus</i> (open)	Bushland	open	0.40	0.30	0.044	active
S1	<i>M. michaelseni</i> (closed)	Grassland	closed	0.75	0.70	0.259 <sup>a</sup>	active
S2	<i>M. michaelseni</i> (closed)	Grassland	closed	0.65	0.35	0.106 <sup>a</sup>	active
S3	<i>M. michaelseni</i> (closed)	Grassland	closed	0.55	0.30	0.217 <sup>a</sup>	dead
S4	<i>M. michaelseni</i> (closed)	Grassland	closed	0.35	0.40	0.074 <sup>a</sup>	dead
S5 <sup>b</sup>	<i>M. subhyalinus</i> (open)	Grassland	open	0.75	0.40	0.193 <sup>a</sup>	active
S6	<i>M. subhyalinus</i> (open)	Grassland	open	0.72	0.15	0.072	dead

<sup>a</sup> These values were estimated using photogrammetric technique.  
<sup>b</sup> The gas fluxes of this mound were not estimated due to nonlinear concentration time series (Fig. S2)

Deleted Cells

Formatted Table

## 2.2 Estimation of mound volume

Volumes of the grassland mounds S1–S5 (Table 1) were measured photogrammetrically by taking a large number of digital images from different positions around the mound (method described in Vesala et al., 2019b). To increase the number of observations, this data was supplemented with five additional comparable-sized mounds measured earlier in a close by area (Vesala et al., 2019b). Linear regression with zero intercept was fitted between equivalent cone volume and photogrammetric mound volume (Fig. 2). This relationship was used to estimate the volume at the bushland site where the photogrammetric method was not feasible due to the surrounding dense vegetation. The equivalent cone volume was calculated as

$$V_{cone} = \frac{1}{3} \pi R^2 \times H, \quad (1)$$

where  $R$  is the average of North-South and West-East measured radii and  $H$  is height. The above ground termite mound volume ( $V_m$ ) was  $0.87 \cdot V_{cone}$  for mounds that were not sampled using the photogrammetric method. The above ground termite mound volumes ranged from 0.3 to 10 % of the total chamber volume.




Figure 2. The relationship between the photogrammetrically measured mound volume and calculated cone volume (Eq. 1) according to equation 1. The data points include grassland mounds (S2 and S4) of this study and five additional mounds measured in Vesala et al., (2019b) (Table S1).

5



### 2.3 Gas flux measurements from termite mounds and soil

CO<sub>2</sub> and CH<sub>4</sub> fluxes were measured using a static chamber method in November 2016 (end of long dry season) and April 2017 (wet season) (Table S2). The chamber collar was installed before each measurement and surrounding sand was applied to insulate collar. The chamber closure time was 15 minutes, and each mound was measured three times with five-minute breaks in between the repetitions. The mean and standard deviation of the flux was calculated from the three repeated measurements. Daytime flux measurements were made between 10 am and 4 pm. To study diurnal variation in gas fluxes, additional nighttime measurements were made at two closed mounds (S1 and S2); the mound S1 and S2 were measured during at least three different time periods from evening until next day with three repeated measurements each time. Mound S1 was measured on 15 November 2016 and mound S2 was measured on 7 November 2016.

The size of the measurement chamber was a compromise between practicality and typical size of termite mounds in the area. The radius of the chamber was 0.75 m and height was 1.5 m, still excluding the vast majority of *Macrotermes* mounds occurring in the area, which are larger on average. The metal collar height was 40 cm. Four upright positioned steel rods and an attaching circular steel frame were used to support the cylindrical shape of the custom made polyvinylchloride sheet. A rubber band was used to seal the polyvinylchloride sheet against the collar. Three people were needed to open and close the chamber. The gas analyzer (LGR Ultra-Portable Greenhouse Gas Analyzer Model 915-0011) was connected to the collar using Swagelok fittings. Chamber air was sampled continuously into the gas analyzer and circulated back to the chamber. The air was sampled inside the chamber using a 2.5 m long teflon tube which had small holes along its length. The sampling line was attach in a spiral shape by supporting it from the middle points of the supporting steel rods and it was connected to the inlet of the gas analyzer at the inside wall of the collar. The air inside the chamber was mixed using three 12 cm fans that were hanging in the middle point of the chamber. The gas analyzer recorded concentration every second. Temperature and relative humidity inside the chamber waswere measured using a Campbell CS215 T/RH sensor attached to the collar. A cellphone timing application (Time meter) was used to record the precise start and end times of each measurement because the gas analyzer produces one continuous measurement file per day. Also, the manual soil moisture and temperature measurements were noted in the timing application. The notes were exported to CSV file which was parsed into numerical form. This procedure enabled to automate flux calculation from raw data and minimize note taking errors.

After termite mound measurement, the inlet lines were connected to a small opaque soil chamber (radius 7 cm, height 22 cm) to measure soil CO<sub>2</sub> and CH<sub>4</sub> fluxes around the termite mound. The soil gas flux measurements were made once for two opposing directions from 2, 4 and 6 m distance from the perimeter of the mound. The soil chamber was placed on patches of bare soil to minimize the effect of vegetation. The sampling time was two minutes. One CH<sub>4</sub> flux measurement around S1

mound at 6 m distance during wet season was discarded because the flux was over thousand times higher than the mean of other measurements around the mound (most likely due to an underground termite nest).

Methane and carbon dioxide fluxes were calculated as

$$F = \frac{PV}{RTA} \frac{dC}{dt}, \quad (2)$$

5

where  $V = V_c - V_m$  for termite mound measurements and  $V = 0.0034 \text{ m}^3$  for the soil measurements,  $V_c = 2.65 \text{ m}^3$  is the termite chamber volume,  $P$  is pressure,  $A$  is the area of the chamber,  $R$  is gas constant ( $8.314 \text{ Jmol}^{-1}\text{K}^{-1}$ ) and  $T$  is air temperature in Kelvin. The slope of concentration increase  $\frac{dC}{dt}$  was determined from a linear regression of the 1 Hz concentration measurements.

#### 10 2.4 Environmental measurements

The soil moisture and temperature were measured manually nearby the mound during each chamber measurement. A handheld soil moisture probe Campbell HydroSense II was used to measure surface soil moisture. Soil temperature was measured using a stainless-steel temperature probe. Termite nest temperature measurements were used from an earlier study (Vesala et al., 2019b) where small temperature sensors (iButton Thermochron DS1922L) were placed next to the first fungus combs that were encountered by digging small hole to the mound. The nest temperature measurements made at the grassland location in 2015 from large open and closed mounds (IDs TS56 and TS19 in Vesala et al., 2019b) were used. These nest temperatures were compared to continuous soil temperature measurements at the Maktou weather station. The weather station soil temperature (Campbell CS650) probe was measuring at 30 cm depth inside a fenced area.

20 The  $\text{CO}_2$  and  $\text{CH}_4$  fluxes measured for each mound were ~~correlated~~ combined with above ground biomass (AGB) data, assessed using airborne laser ~~scanner data~~ scanning and field measurements (Amara et al., 2020; Pellikka et al., 2018) ~~applying~~. This combination aimed to investigate the ~~log~~ working hypothesis that fluxes ~~may vary~~ might be related to the C:N ratios of termite diets that ~~correlate negatively~~ with ~~the amount woody~~ vegetation density in the ~~environment~~ area (Vesala et al., 2022b).


#### 2.5 Statistical analysis

25 Simple linear regression was used to test the statistical relationship between the measured fluxes and explanatory variables. The relationship between fluxes and mound volume was tested separately during the dry season ( $n=11$ ) and wet season ( $n=7$ ). ~~For the~~ The relationship ~~between~~ of the environmental variables ~~and~~ to the fluxes and ~~to~~ the ratio of  $\text{CH}_4$  to  $\text{CO}_2$  flux was tested using all the measurements ( $n=18$ ). The mound fluxes and AGB relationship was tested with simple linear regression during the dry season ( $n=11$ ). Statistical analyses were done using the Scipy Python package (Virtanen et al., 2020).

### 3 Results

#### 3.1 Environmental conditions

Total precipitation before the first measurement campaign was 10 mm from September to early November (Fig. 3), meaning that the short rainy season had not yet started. The precipitation between the campaigns was 153 mm corresponding to a typical dry year amount. The annual mean precipitation during this period between the campaigns was 287 mm with 154 mm standard deviation calculated from six years of precipitation measurements at the weather station. The surface soil moisture was twice as high during the second campaign compared to first campaign (Fig. 3a). The wind speed had a consistent trend during the measurement campaign days showing a minimum around 6 am and a maximum during the early afternoon. Revisiting the data collected in Vesala et al. (2019b) shows that termite mounds in the area have a consistent diurnal temperature rhythm with the maximum nest temperature being always around 8 pm, whereas the soil temperature at 30 cm depth has a maximum around midnight.







Figure 3. (a) Time series of daily rainfall (black) and daily mean soil moisture at 10 cm depth (blue). The gray areas indicate the days of the measurement campaigns. (b) Diurnal mean of the wind speed during the first measurement campaign. (c) Soil temperature at 30 cm depth and termite nest temperatures for open (TS56) and closed (TS19) mounds at the grassland site in 2015 during period of no rainfall (Vesala et al., 2019b). (d) Diurnal mean of the soil and nest temperatures during the same period. The error bars indicate one standard deviation.

### 3.2 Mound CO<sub>2</sub> and CH<sub>4</sub> fluxes

The mean CO<sub>2</sub> fluxes ranged from 35 to 24952100 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> for closed *M. michaelseni* mounds and from 446120 to 809800 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> for open *M. subhyalinus* mounds (Fig. 4). The mean CO<sub>2</sub> flux was 40471000 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> for the grassland site and 380 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> for the bushland site from the active mounds listed in Table 1. The largest CO<sub>2</sub> fluxes were measured during the dry season at the grassland site. The site level mean CO<sub>2</sub> flux decreased 64 % at the grassland and increased 41 % at the bushland from dry to wet season. A comparison of the closed mound S1 and open mound MR4, which have similar volumes (Table 1), shows that the closed S1 mound had higher CO<sub>2</sub> and CH<sub>4</sub> flux than the open MR4 mound during the dry season (Fig. 4). The lowest CO<sub>2</sub> fluxes were registered from the grassland mounds S3 and S4, which turned out to be dead when opened/excavated after the second measurement campaign (Table 1).

The mean CH<sub>4</sub> fluxes ranged from 0.02 to 3.667 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> for closed *M. michaelseni* mounds and 0.06 to 1.452 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> for open *M. subhyalinus* mounds. The ratio of mound CH<sub>4</sub> flux to mound CO<sub>2</sub> flux ranged from 0.4 · 10<sup>-3</sup> to

$2.8 \cdot 10^{-3}$  with no clear difference between termite species or mound volume. The mean ratio was  $1.6 \cdot 10^{-3}$  for grassland and  $1.2 \cdot 10^{-3}$  for bushland sites. The ratio of CH<sub>4</sub> flux to CO<sub>2</sub> flux reduced 36 % at the grassland and 49 % at the bushland site from dry to wet season. The dry season mound CO<sub>2</sub> and CH<sub>4</sub> fluxes were not significantly correlated with the AGB (Fig. S3).






Figure 4. Mean termite mound CO<sub>2</sub> and CH<sub>4</sub> fluxes from grassland and bushland sites during dry and wet seasons: (a-b). (c) Ratio of CH<sub>4</sub> to CO<sub>2</sub> flux. Each mound was sampled once in each season during daytime (10 am to 4 pm) by three repeated measurements. Error bars indicate standard error. Hollow bars indicate open and solid bars closed mounds. The mounds that were dead in April 2017 were only measured during the dry season. The mounds are presented in order of increasing volume for each site.



Based on diurnal measurements in two closed mounds, the largest CO<sub>2</sub> flux was measured around noon at the mound S1 (Fig. 5) coinciding with the maximum diurnal wind speed and the minimum nest temperature (Fig. 3)-5). However, the CO<sub>2</sub> flux increased also during the midnight for both S1 and S2 mounds. For both mounds, the difference between maximum nighttime CO<sub>2</sub> flux compared with daytime maximum was less than 2 standard errors. For S1 mound, the maximum flux value was 1.4 times the daily average for CO<sub>2</sub> and 1.5 times for CH<sub>4</sub>. The nighttime peak in the CO<sub>2</sub> flux coincides with the maximum soil temperature at 30 cm depth at midnight (Fig. 3d). The diurnal cycle of the ratio of CH<sub>4</sub> to CO<sub>2</sub> flux showed one peak around the noon for S1 mound and at 6 pm for S2 which was not sampled during the noon. These peaks are due to the increase in CH<sub>4</sub> flux in proportion to CO<sub>2</sub> flux during the daytime.




Figure 5. Diurnal variation of CO<sub>2</sub> and CH<sub>4</sub> fluxes during 2016 dry season for closed mounds S1 and S2. Mound S1 was measured on 15 November 2016 and mound S2 was measured on 7 November 2016. Error bars indicate standard error calculated from three repeated measurements. The hollow symbols indicate daytime measurements that were done nine days before nighttime measurements for S1 and one day before for S2.

### 3.3 Drivers of mound CO<sub>2</sub> and CH<sub>4</sub> fluxes

The mean mound CO<sub>2</sub> flux was significantly correlated with the termite mound volume during the wet season ( $R^2 = 0.96$ , p-value < 0.01; Fig. 6b), while the relation was scattered during the dry season (Fig. 6a). The CO<sub>2</sub> flux and CH<sub>4</sub> flux were not explained by surface soil moisture or soil temperature during the dry or wet season (Fig. S4). The soil moisture values were low for most mounds due to low precipitation amount during the wet season. The mean CO<sub>2</sub> and CH<sub>4</sub> flux were linearly related (Fig. 6c). The ratio of CH<sub>4</sub> and CO<sub>2</sub> fluxes had no significant correlation with environmental variables and only a weak correlation with the magnitude of CH<sub>4</sub> flux ( $R^2 = 0.34$ , p-value = 0.01). The CH<sub>4</sub> flux was related to the mound volume during the wet season with slightly more scatter than the equivalent CO<sub>2</sub> relation (Fig. 6e). The standard deviation of CO<sub>2</sub> flux was increasing linearly with the mean flux magnitude (Fig. 6f).




Figure 6. The relationship between mound volume and mean CO<sub>2</sub> and CH<sub>4</sub> fluxes during dry and wet seasons (a-b and d-e). (c) The relationship between CH<sub>4</sub> and CO<sub>2</sub> fluxes. **Circles indicate open mounds and triangles indicate closed mounds.** (f) The relationship between standard deviation and mean of the CO<sub>2</sub> flux **calculated from three repeated measurements.**

### 3.4 Soil CO<sub>2</sub> and CH<sub>4</sub> fluxes around termite mounds

The soil CO<sub>2</sub> and CH<sub>4</sub> fluxes around the mounds were sampled to determine the effect of termites on soil fluxes around the mound. The site mean soil CO<sub>2</sub> and CH<sub>4</sub> were always higher at a 2 m distance compared to a 4 and 6 m distance from the perimeter of the mound (Fig. 7 and Table 2). The mean soil CO<sub>2</sub> flux at the bushland was only 29 % of the mean soil flux at the grassland. The dry season means of all soil CO<sub>2</sub> flux measurements were 62 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> and 27 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> at 2 m and 4 to 6 m distance (Table 2). The difference between soil CO<sub>2</sub> flux at 2 m and 4 to 6 m distance from the mound was less pronounced during the wet season. The mean mound CO<sub>2</sub> fluxes were a factor of 25 and 6 higher at the dry and wet seasons in comparison to the 4 to 6 m distance soil CO<sub>2</sub> fluxes (Table 2). The dry season means of all soil CH<sub>4</sub> flux measurements were 0.06507 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> and 0.02002 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> at 2 m and 4 to 6 m distance (Table 2). The dry season mean soil CH<sub>4</sub> flux was positive at the grassland and bushland site, whereas the mean wet season flux was nearly zero with most fluxes being negative (Fig. 7, S5 and S6). The soil CH<sub>4</sub> fluxes at a 2 m distance from the mound are much higher in proportion to fluxes at a 4 m distance at the grassland compared to the bushland (Fig. 7).

**Table 2. Mean CO<sub>2</sub> and CH<sub>4</sub> fluxes of termite mounds and surrounding soils. Data from all active mounds measured at grassland and bushland. The range of measured values for each group is given in the parenthesis.**

	CO <sub>2</sub> flux (mg CO <sub>2</sub> -C m <sup>-2</sup> h <sup>-1</sup> )	CH <sub>4</sub> flux (mg CH <sub>4</sub> -C m <sup>-2</sup> h <sup>-1</sup> )
Dry season		
Termite mound	666 (116–2105)	1.294 (0.159–3.659)
Soil at 2 m distance	62 (10–122)	0.065 (0.001–0.186)
Soil at 4–6 m distance	27 (6–123)	0.020 (–0.003–0.108)
Wet season		
Termite mound	475 (136–881)	0.542 (0.062–1.433)
Soil at 2 m	90 (29–243)	0.006 (–0.034–0.104)
Soil at 4–6 m	83 (15–251)	–0.004 (–0.032–0.065)

Formatted: Not Superscript/ Subscript




Figure 7. (a-b) Mean soil-CO<sub>2</sub> and CH<sub>4</sub> fluxes around the from termite mounds for and surrounding soils during the dry season. Distances 2, 4, and 6 m distances are measured from the perimeter of the mound during the dry season. The mean is to two opposing directions. Mean values and standard errors are calculated from both directions and from all the mounds at the site. Soil around each mound was measured at three different distances and from two different directions. Error bars indicate standard error. (c) The ratio of absolute value of soil-CH<sub>4</sub> and CO<sub>2</sub> flux.

## 4 Discussion

### 4.1 Mound CO<sub>2</sub> and CH<sub>4</sub> fluxes

The measured CO<sub>2</sub> fluxes (from 35 to 24052100 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>) were in similar range as reported in two earlier comparable studies where chamber measurements have been used to study greenhouse gas emissions from termite mounds in Australia (up to 45501600 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>, Jamali et al., 2013) and Amazonia (492-2246190-2200 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup>, van Asperen

et al., 2021). Methane fluxes ranged from 0.02 to 3.667 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> and were 2–20 times lower than those reported in earlier studies (0.4–6 mg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup>, van Asperen et al., 2021; Jamali et al., 2013). ~~As in earlier chamber measurement studies termites have been other than fungus growing species, this result was expected: in fungus growing termites a large proportion of the mound total CO<sub>2</sub> fluxes originate from aerobic fungus gardens which seem not to emit measurable amounts of CH<sub>4</sub> (Darlington et al., 1997).~~ In most fungus-growing species studied so far, workers have exhibited relatively high CH<sub>4</sub> production rates (Rouland et al., 1993; Zhou et al., 2023). Thus, the lower CH<sub>4</sub> emissions, in relation to CO<sub>2</sub>, observed in this study compared to mounds of previously studied termite species (which all have been other than fungus-growing termites) could be attributed to the fact that a large proportion of the total CO<sub>2</sub> in our measurements was originating from the aerobic fungus gardens which do not produce CH<sub>4</sub> (Darlington et al., 1997). Our biased sampling design, with one termite species dominating the bushland and the other the grassland site, does not allow us to reliably compare the two termite species. However, the relatively similar gas fluxes of the two species, or mound types, (Figures 4 and 6) imply that they do not exhibit major differences in terms of CO<sub>2</sub> or CH<sub>4</sub> emissions. This result is also highly expected as the two termite species are closely related (Bagine et al., 1994; Vesala et al., 2017), have very similar population sizes and compositions (Darlington, 1984), and also share their dietary niches (Vesala et al., 2022b). The highest fluxes of both gases were registered from two closed mounds of *M. michaelseni* in the grassland. It is important to notice that we did not obtain any information about the population sizes and, thus, these two *M. michaelseni* mounds may simply have housed larger termite populations with correspondingly larger fungus gardens than the studied *M. subhyalinus* mounds.

~~As found by previous studies, the mound outer dimensions of both *Macrotermes* species studied here, mound outer dimensions,~~ correlate positively with the size of the termite population and the total biomass of fungus combs (Darlington and Dransfield 1987; Darlington 1990). Thus, the fluxes of metabolic gases should logically also show correlation with the mound volume. However, considering the high variation associated with the fluxes, it seems obvious that if the sizes of the studied mounds are too close to each other, the relationships between mound volume and gas fluxes are difficult to detect. In this study, the mound volumes ranged from 0.023 to 0.259 m<sup>3</sup> (Table 1) including notably more size variance than the earlier comparable studies (0.02 to 0.077 m<sup>3</sup>) (Jamali et al., 2013; van Asperen et al., 2021). We found that the measured CO<sub>2</sub> fluxes were linearly correlated with the termite mound volume during the wet season (Fig. 6b) but the relationship was more indistinct during the dry season (Fig 6a). Determining the relationship between mound volume and gas fluxes more reliably would require measurements from still much larger mounds, which however is problematic due to technical limitations. In East Africa, average volumes of *Macrotermes* mounds range from 2 to 3 m<sup>3</sup> depending on the geographical area, and the maximum volumes can exceed 12 m<sup>3</sup> (Pomeroy, 1977). Also in our study area, *M. subhyalinus* mounds with volumes of up to 6 m<sup>3</sup> have been recorded (Vesala et al., 2019b), which is more than 20 times the volume of the largest mound measured here. Thus, the reported average flux values in this study are representative for the smallest *Macrotermes* mounds but when scaling the results to the level of larger *Macrotermes* mounds (or calculating area-based estimates), the effect of mound volumes should be considered to avoid serious underestimations.

Formatted: English (United Kingdom)

Formatted: Font: Not Italic, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

The mounds at the grassland and bushland sites had contrasting seasonality of CO<sub>2</sub> fluxes with 64 % decrease at the grassland and 35 % increase at the bushland site from dry to wet season (Fig. 4). This seasonal change is smaller than what has been measured for grass and wood-feeding termites in Australia, which had a 90 % decrease in CO<sub>2</sub> flux from wet to dry season (Jamali et al., 2013). The relatively small intra-annual changes found between our two measurement campaigns reflects the fact that ~~sterile populations~~ within *Macrotermes* nests the number of termites remain relatively constant throughout the year (Darlington 1986, 1991). However, some intra-annual variation can be caused by ~~biomass changes~~the annually produced brood of alates that, in other nest components. For example, within Kenyan *Macrotermes* *M. michaelseni* nest living biomass can be briefly elevated by the annual brood of alates reaching, reaches its maximum biomass just before swarming in October–November (Darlington 1986). Correspondingly, storages of fungus combs are typically at their largest soon after the long rains in May–June after which they start to decline ~~as food availability decreases and, concurrently, the nutritional resources are increasingly allocated to fat accumulated in developing alates~~ (Lepage 1981b) (Lepage 1981b). Such seasonal changes in biomasses of different nest components could cause intra-annual variation in gas fluxes. Additionally, differences in food quality could potentially affect nest CO<sub>2</sub> emissions. Within our study area, fungus combs of grassland mounds had significantly lower nitrogen contents than those situated in more wooded habitats suggesting that, on average, grass-based diets have lower nutritional value than wood-based diets (Vesala et al., 2021b) (2022b). Such differences in food nutritional quality may reflect to nest CO<sub>2</sub> fluxes as more ~~carbon~~carbohydrates needs to be eliminated to accumulate ~~enough nitrogen~~sufficiently proteins from low-nitrogen food sources compared to those with a higher nitrogen content (Higashi et al, 1992). Most likely, food nutritional quality also changes during the year, with the changes possibly being more dramatic in grasslands having fewer and more competed food sources available for termites than ~~the bushlands with~~ more abundant and diverse ~~bushlands~~food sources. Such local differences in food availability and quality could potentially lead to different seasonality in nest living biomass which might explain the ~~opposite~~contrasting seasonal changes observed in CO<sub>2</sub> fluxes between the two habitats and the higher intra-annual variance in gas fluxes observed in grassland and than in bushland mounds. We note that our seasonal sampling was limited one measurement period during dry and wet season which does not capture the seasonal variability throughout the whole year.

There was a clear decrease in CH<sub>4</sub> to CO<sub>2</sub> flux ratios from dry to wet season measurements; at both study sites. This difference might be partially explained by the smaller biomass of fungus gardens (produce only CO<sub>2</sub>) in relation to that of termites (produce both CO<sub>2</sub> and CH<sub>4</sub>) during dry season than during wet season (Darlington, 1986). Another explanation could be ~~most~~ logically linked to the activity of methanotrophic bacteria within mound structures and/or surrounding soils that have been reported to consume a notable proportion of CH<sub>4</sub> produced in nests of several mound-building termite species (Sugimoto et al., 1998; Nauer et al., 2018). ~~During wet seasons,~~Although the humidity within termite nests remain at a constantly high level throughout the year (Lüscher, 1961; Agarwal, 1980), water content in surrounding soils and mound soil structures have higher water content than can decrease notably during dry periods, which with low precipitation (Jamali et al., 2011; Chen et al., 2019)

Formatted: Subscript

Formatted: Subscript

(Figure 3a). Dry soil conditions during periods with low precipitation most likely ~~enhances the limit~~ microbial activity ~~of the soil microbiome~~ (including the methane-oxidizing bacteria) within termite mound walls and surrounding soils, whereas increases in soil water content after rains could oppositely lead to increased microbial activity and enhanced methanotrophy.

This result is also in congruence with our measurements of nest internal gas concentrations within *M. michaelseni* mounds, which tended to have higher CH<sub>4</sub> to CO<sub>2</sub> concentration ratios during the driest period compared to seasons with higher soil moisture (Vesala et al., ~~in preparation~~ submitted).

Four mounds that were measured during the first campaign were determined to be dead at the end of the second campaign (Table 1). The exceptionally low CO<sub>2</sub> fluxes measured from the mounds S3 and S4, however, strongly suggest that these two mounds were dead, or at least seriously weakened, already during the first measurement campaign. An average lifespan of a *Macrotermes bellicosus* colony is thought to be around four years, whereas the mounds that they build persist typically much longer (Pomeroy, 1976). However, a recent long-term monitoring study in Namibia found that 57% of initially active *M. michaelseni* mounds were still active after 12 years (Wildermuth et al., 2022), demonstrating that some colonies can live much longer. These authors also found that small mounds were more likely to be dead than large mounds after 12 years, showing that large well-established colonies are less vulnerable to disruptions. Hence, dead *Macrotermes* mounds are a common feature in many landscapes and for example, in many places within our study area, their number exceeds that of the active mounds (Vesala et al., 2017). Thus, determination of the proportion of active mounds in a landscape is a necessary precondition for ecosystem-level termite flux estimations for all such termite species that build persistent soil mounds.

#### 4.2 Diurnal variation

Diurnal variation of CO<sub>2</sub> and CH<sub>4</sub> fluxes were ~~studied~~ measured in two closed *M. michaelseni* mounds over a period of one day. The afternoon peaks were on average 46 % and 57 % higher compared to the daily averages for CO<sub>2</sub> and CH<sub>4</sub> fluxes, respectively (Fig. 5). Our observations are in congruence with the nest internal CO<sub>2</sub> concentration measurements by Ocko et al. (2017) who found that CO<sub>2</sub> levels within *M. michaelseni* nests in Namibia peaked in early afternoon, whereas concentrations in night-time and especially soon after sunrise were somewhat lower. A corresponding trend has been documented also from outflow chimneys of Kenyan *Macrotermes jeanneli* that had the highest CO<sub>2</sub> concentrations at 3 pm and the lowest around 3–6 am (Darlington et al., 1997). The daily maximum CO<sub>2</sub> flux in early afternoon found in this study (Fig. 5) agrees well with these earlier data suggesting that the internal CO<sub>2</sub> concentrations within *M. michaelseni* mounds show similar diurnal patterns to the mound CO<sub>2</sub> fluxes. A positive relationship between the nest internal gases and mound fluxes has been demonstrated for a few of mound-building termites (Khalil et al., 1990; Jamali et al., 2013). In agreement with our CH<sub>4</sub> measurements, a similar peak in CH<sub>4</sub> fluxes soon after midday was documented by Jamali et al. (2011) from mounds of three Australian termite species.

The lower CO<sub>2</sub> and CH<sub>4</sub> fluxes during nighttime compared to daytime fluxes are probably mostly associated with the diurnal movements of termites: after sunset a proportion of termite workers and soldiers leave the nest due to foraging activities, thus decreasing the CO<sub>2</sub> and CH<sub>4</sub> production within the nest. During night, termites also open the numerous foraging passages that directly connects the nest interior space to the soil surface near the mound (Lepage 1981a; Darlington 1982). This might lead in enhanced nest ventilation and could additionally affect gas fluxes originating from the mound. In addition to the afternoon flux peaks, we registered another peak in both CO<sub>2</sub> and CH<sub>4</sub> fluxes at midnight, which has not been documented in earlier studies (Fig. 5). Reason for this midnight peak in gas fluxes is not currently understood but might be related to temperature gradient driven diurnal changes in mound internal air currents that slow down and may even shift their direction as the mound surface cools down after sunset (Korb and Linsenmair 2000; Ocko et al., 2017). Temporary decreases in nest internal gas flows in the evening could lead in accumulation of nest internal gas concentrations (Korb, 2011) which, in turn, might trigger slightly enhanced CO<sub>2</sub> and CH<sub>4</sub> fluxes soon after the internal flows are reverted again. Regarding the above discussion, more measurements would be needed to understand the drivers of diurnal variation because we measured only two mounds in one location during the dry season. It would be especially useful to continuously measure the mound internal gas concentrations together with the mound fluxes.

In addition to the daily cycles of CO<sub>2</sub> and CH<sub>4</sub> fluxes, also their mutual relationship between CO<sub>2</sub> and CH<sub>4</sub> showed a clear diurnal pattern. The mound CH<sub>4</sub> to CO<sub>2</sub> flux ratio was higher during the daytime than during the nighttime measurements (Fig 5c). This result also supports the interpretation that termite population within the mound is smaller during nighttime than during daytime due to foraging activity. Termite metabolism Metabolism of termites (together with gut microbiota) produces both CH<sub>4</sub> and CO<sub>2</sub>, but a major proportion of CO<sub>2</sub> originates from the fungal gardens, which are sessile and do not produce CH<sub>4</sub> (Darlington et al., 1997). Thus any changes in termite biomass within the nest should inevitably reflect to the mound CH<sub>4</sub> to CO<sub>2</sub> flux ratios.

After all, the diurnal changes in gas fluxes measured in this study were relatively small. This underlines the fact that plant biomass degradation within mounds of fungus-growing termites is a continuously ongoing process and show only minor diurnal changes. Therefore, also a large proportion of termites stay constantly in the nest taking care of the fungus gardens and thus the diurnal biomass changes within the nest are relatively small. The stable diurnal flux variability is in line with the stable humidity and temperature conditions within mounds of fungus-growing termites. Constant temperature and high humidity are of key importance for the fungal *Termitomyces* symbionts which have relatively narrow range of optimal microclimatic conditions (Lüscher, 1961; Thomas 1981; Wood and Thomas, 1982, 1989).

### 4.3 Soil CO<sub>2</sub> and CH<sub>4</sub> fluxes around the mounds

The soil CO<sub>2</sub> and CH<sub>4</sub> fluxes were always higher close to the mound (at 2 m distance from the mound perimeter) compared to further distances (4 to 6 m from the mound perimeter) (Fig. 7). The mean soil CO<sub>2</sub> flux from 4 to 6 m distance was 27 mg

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)



CO<sub>2</sub>-C m<sup>2</sup> h<sup>-1</sup> during the dry season, which is comparable to the measured soil CO<sub>2</sub> flux seasonal minimum of 20 mg CO<sub>2</sub>-C m<sup>2</sup> h<sup>-1</sup> at the grassland site (Wachiye et al., 2020). Correspondingly, the mean soil CH<sub>4</sub> fluxes at 4 to 6 m distances from the mounds were 0.020 and -0.004 mg CH<sub>4</sub>-C m<sup>2</sup> h<sup>-1</sup> at dry and wet seasons, respectively (Table 2), which is similar to the earlier field study where the soil CH<sub>4</sub> fluxes were slightly negative with no seasonality at the grassland site (Wachiye et al., 2020).

5 Our results thus demonstrate that, although the majority of the gas-exchange between termite nests and ambient air takes place through the mound structures, a small proportion of gases leave the nest also via adjacent soils. at up to 2 m distance. These results differ from the earlier measurements in soil feeding termites in Amazon which had enhanced soil fluxes around the mound only up to 0.5 m from the perimeter of the mound (van Asperen et al., 2021). The difference can be largely explained by the much larger size of the studied/measured mounds in our study but it most likely reflects also some fundamental differences between the mounds of soil-feeding and fungus-growing termites. While within soil-feeding species the mound itself acts as a nest for the colony, mounds built by many fungus-growing species especially *Macrotermes michaelseni* are more or less empty from the insects and the actual nests are largely located underground below the mounds (Noirot and Darlington, 2000; Korb, 2011). Thus, the actual nests of fungus-growing termites may be much larger and extend to a somewhat wider area than their aboveground mounds. *Macrotermes* termites also have an extensive network of foraging tunnels surrounding their nest with the largest radial tunnels having width of 4–7 centimeters (Darlington, 1982). Such structures increase soil porosity around the mounds enhancing water infiltration (Jouquet et al., 2011) and, obviously, also enable some gas-exchange to occur through the soil.

20 Soil CH<sub>4</sub> fluxes around the mounds were systematically lower during wet than during dry season at all three distances (Table 2, Figures S5 and S6). As the CH<sub>4</sub> fluxes from the soils unaffected by termite mounds are shown to be slightly negative in grasslands of the same ecosystem (Wachiye et al., 2020) we are confident that termites were the main source of the soil CH<sub>4</sub> fluxes detected in the current study. Seasonal changes in soil fluxes could thus be caused either by changes in number of termites, their CH<sub>4</sub> production rates, or methane oxidation. During the rainy seasons, termite foraging activity is at the lowest (Lepage, 1981a), which could potentially decrease CH<sub>4</sub> fluxes in mound surroundings as the foragers move from foraging tunnels towards the nests. This should, however, concurrently increase the CH<sub>4</sub> fluxes from the mounds, which was not observed. Instead, we found also the mound CH<sub>4</sub> fluxes to be lower during the rainy than during the dry season. Thus, we suspect that, similarly as discussed above in case of mound fluxes, the soil CH<sub>4</sub> fluxes near the mounds are largely regulated by CH<sub>4</sub> oxidation within the soils surrounding the mounds, which could be enhanced by the increased soil water content after rains (Figure 3a).

30 We found that CH<sub>4</sub> fluxes at 2 m distance from the mounds were more enhanced (compared to further distances) at the grassland than in the bushland (Fig. 7). This might simply reflect the size of the underground nest, which may have extended to a larger area in grassland than in bushland, but might also be affected by the fact that in grasslands termites have more homogeneously distributed food sources (i.e., uniform grass layer) compared to the randomly distributed food resources

available at the bushland site (e.g., fallen branches and logs of dead wood). Optimal exploitation of such differently distributed food sources might lead termites to adjust their foraging networks in different habitats, with grasslands potentially having more densely branched tunnel network with more densely distributed foraging passages in mound surroundings than in bushlands. This, in turn, might lead to a more porous soil structure in the vicinity of grassland mounds. Such interactions are, however, currently poorly known.

## 5 Conclusions

The CO<sub>2</sub> and CH<sub>4</sub> fluxes from fungus-growing termite mounds show diurnal and seasonal variation, albeit lower in magnitude than their wood, grass, and soil-feeding counterparts. The likely reason for this stability is the needed temperature and humidity control for the fungal symbiont. The CO<sub>2</sub> and CH<sub>4</sub> fluxes scale with mound volume over a large volume range during the wet season. The soil flux measurements around the mound also show that the soil CO<sub>2</sub> and CH<sub>4</sub> fluxes are significantly affected by the termites at a 2 m distance from the perimeter of the mound. The fungus-growing termite mounds have lower CH<sub>4</sub> fluxes per unit CO<sub>2</sub> than soil and grass-feeding termites, which is due to the plant matter decomposition by *Termitomyces* fungi that do not emit CH<sub>4</sub>. For upscaling the greenhouse gas fluxes originating from mounds of fungus-growing termites, the work highlights the need to determine include the mound volume in the models, as well as to consider the potentially high proportion of active/inactive termite mounds in the landscape.

*Data availability.* The data used in this study are available online:- <https://doi.org/10.6084/m9.figshare.21739484.v1> (Räsänen et al., 2022).

*Author contributions.* MR, PM, MJ, PP, JR conducted the measurements; MR, RV, PR, LA, PP, JR designed the analysis; All authors contributed to the final version of the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgments.* This work was supported by the Academy of Finland through the projects SMARTLAND (grant number 318645)— Environmental sensing of ecosystem services for developing climate smart landscape to improve food security in East Africa (PI P. Pellikka) and IsoTermes (grant number 333868). We thank Venla Pellikka, Mwadime Mjomba, Muhia Gicheru and Darius Kimuzi from Taita Research Station of the University of Helsinki and Remig Righa Mjomba, Wilson Mkala and Tobius Mwamburi from Taita Hills Wildlife Sanctuary for the support during the field work campaign. Research permit from the National Council for Science and Technology, Kenya, permit N° NCST/RCD/17/012/33 is appreciated.



## References

Agarwal, V. B.: Temperature and relative humidity inside the mound of *Odontotermes obesus* (Rambur) (Isoptera: Termitidae). *Proc. Indian Acad. Sci. (Anim. Sci.)* 89, 91–99, <https://doi.org/10.1007/BF03179148>, 1980.

Amara, E., Adhikari, H., Heiskanen, J., Siljander, M., Munyao, M., Omondi, P., and Pellikka, P.: Aboveground Biomass Distribution in a Multi-Use Savannah Landscape in Southeastern Kenya: Impact of Land Use and Fences, *Land*, 9, 381, <https://doi.org/10.3390/land9100381>, 2020.

Bagine, R., Brandl, R., Kaib, M.: Species Delimitation in *Macrotermes* (Isoptera: Macrotermitidae): Evidence from Epicuticular Hydrocarbons, Morphology, and Ecology, *Annals of the Entomological Society of America*, 87, 498-506, 1994.

Bignell, D. E. and Eggleton, P.: Termites in ecosystems, in *Termites: Evolution, Sociality, Symbioses and Ecology*, edited by T. Abe, D. E. Bignell, and M. Higashi, pp. 363–387, Springer, 2000.

Boutton, T. W., Arshad, M. A. and Tieszen, L. L.: Stable isotope analysis of termite food habits in East African grasslands, *Oecologia*, 59(1), 1–6, doi:10.1007/BF00388065, 1983.

Brümmer, C., Papen, H., Wassmann, R., and Brüggemann, N.: Fluxes of CH<sub>4</sub> and CO<sub>2</sub> from soil and termite mounds in south Sudanian savanna of Burkina Faso (West Africa), *Global Biogeochemical Cycles*, 23(1), <https://doi.org/10.1029/2008GB003237>, 2009.

Buxton, R. D.: Termites and the turnover of dead wood in an arid tropical environment, *Oecologia*, 51(3), 379–384, doi:10.1007/BF00540909, 1981.

Chen, C., Wu, J., Zhu, X., Jiang, X., Liu, W., Zeng, H., and Meng, F.-R.: Hydrological characteristics and functions of termite mounds in areas with clear dry and rainy seasons, *Agriculture, Ecosystems & Environment*, 277, 25–35, <https://doi.org/10.1016/j.agee.2019.03.001>, 2019.

Chiri, E., Greening, C., Lappan, R., Waite, D. W., Jirapanjawat, T., Dong, X., Arndt, S. K. and Nauer, P. A.: Termite mounds contain soil-derived methanotroph communities kinetically adapted to elevated methane concentrations, *ISME J.*, 14(11), 2715–2731, doi:10.1038/s41396-020-0722-3, 2020.

Formatted: English (United Kingdom)

Formatted: Normal

- Collins, N. M.: Populations, Age Structure and Survivorship of Colonies of *Macrotermes bellicosus* (Isoptera: Macrotermitinae), *J. Anim. Ecol.*, 50(1), 293–311, doi:10.2307/4046, 1981. Collins, N. M.: The role of termites in the decomposition of wood and leaf litter in the Southern Guinea savanna of Nigeria, *Oecologia*, 51(3), 389–399, doi:10.1007/BF00540911, 1981.
- 5
- Darlington, J. P. E. C.: Seasonality in mature nests of the termite *Macrotermes michaelseni* in Kenya, *Insectes sociaux*, 33(2), 168–189, 1986.
- 10
- Darlington, J. P. E. C.: Populations in nests of the termite *Macrotermes subhyalinus* in Kenya, *Insectes sociaux*, 37(2), 158–168, doi:10.1007/BF02224028, 1990.
- Darlington, J. P. E. C.: Turnover in the populations within mature nests of the termite *Macrotermes michaelseni* in Kenya, *Insectes sociaux*, 38, 251–262, 1991.
- 15
- Darlington, J. P. E. C. and Dransfield, R. D.: Size relationships in nest populations and mound parameters in the termite *Macrotermes michaelseni* in Kenya, *Insectes sociaux*, 34(3), 165–180, 1987.
- Darlington, J. P. E. C., Zimmerman, P. R., Greenberg, J., Westberg, C. and Bakwin, P.: Production of metabolic gases by nests of the termite *Macrotermes jeanneli* in Kenya, *Journal of Tropical Ecology*, 13(4), 491–510, doi:10.1017/S0266467400010671, 1997.
- 20
- Darlington, J. P. E. C.: [Two types of mound built by the termite \*Macrotermes subhyalinus\* in Kenya, \*International Journal of Tropical Insect Science\*, 5\(06\), 481–492, doi:10.1017/s1742758400004914A](#) method for sampling the populations of large termite nests, *Annals of applied biology*, 104, 427–436, 1984.
- 25
- Darlington, J. P. E. C.: Structure of mature mounds of the termite *Macrotermes michaelseni* in Kenya, *International Journal of Tropical Insect Science*, 6(2), 149–156, doi:https://doi.org/10.1017/S1742758400006536, 1985.
- 30
- Darlington, J. P. E. C.: The underground passages and storage pits used in foraging by a nest of the termite *Macrotermes michaelseni* in Kajiado, Kenya, *Journal of zoology*, 198(2), 237–247, doi:10.1111/j.1469-7998.1982.tb02073.x, 1982.
- Eggleton, P. and Tayasu, I.: Feeding groups, lifestyles and the global ecology of termites, *Ecological research*, 16(5), 941–960, doi:10.1046/j.1440-1703.2001.00444.x, 2001.

Higashi, M., Abe, T. and Burns, T. P.: Carbon-nitrogen balance and termite ecology, *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 249(1326), 303–308, doi:10.1098/rspb.1992.0119, 1992.

- 5 Hyodo, F., Tayasu, I., Inoue, T., Azuma, J. I., Kudo, T. and Abe, T.: Differential role of symbiotic fungi in lignin degradation and food provision for fungus-growing termites (Macrotermitinae: Isoptera), *Functional Ecology*, 17(2), 186–193, doi:10.1046/j.1365-2435.2003.00718.x, 2003.

- 10 Jamali, H., Livesley, S. J., [Dawes, T. Z., Cook, G. D., Hutley, L. B., and Arndt, S. K.: Diurnal and seasonal variations in CH<sub>4</sub> flux from termite mounds in tropical savannas of the Northern Territory, Australia, \*Agricultural and Forest Meteorology\*, 151, 1471–1479, <https://doi.org/10.1016/j.agrformet.2010.06.009>, 2011.](#)

- 15 [Jamali, H., Livesley, S. J., Hutley, L. B., Fest, B. and Arndt, S. K.: The relationships between termite mound CH<sub>4</sub>/CO<sub>2</sub> emissions and internal concentration ratios are species specific, \*Biogeosciences\*, 10\(4\), 2229–2240, doi:10.5194/bg-10-2229-2013](#), 2013.

Jouquet, P., Traoré, S., Choosai, C., Hartmann, C. and Bignell, D.: Influence of termites on ecosystem functioning. Ecosystem services provided by termites, *European Journal of Soil Biology*, 47(4), 215–222, doi:10.1016/j.ejsobi.2011.05.005, 2011.

- 20 Khalil, M. A. K., Rasmussen, R. A., French, J. R. J. and Holt, J. A.: The Influence of Termites on Atmospheric Trace Gases: CH<sub>4</sub>, CO<sub>2</sub>, CHC<sub>13</sub>, N<sub>2</sub>O, CO, H<sub>2</sub>, and Light Hydrocarbons, *Journal of Geophysical Research: Atmospheres*, 95, 3619–3634, 1990.

- 25 King, H., Ocko, S. and Mahadevan, L.: Termite mounds harness diurnal temperature oscillations for ventilation, *Proc. Natl. Acad. Sci. - PNAS*, 112(37), 11589–11593, doi:10.1073/pnas.1423242112, 2015.

- 30 Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G. Dlugokencky, E. J. Bergamaschi, P. Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J. F. Langenfelds, R. L., Le Quere, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., Werf, van der, G. R., V., A., van W., M., Weiss, R. F., Williams, J. E. and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013.

Formatted: English (United Kingdom)

- Korb, J. and Linsenmair, K. E.: The effects of temperature on the architecture and distribution of *Macrotermes bellicosus* (Isoptera, Macrotermitinae) mounds in different habitats of a West African Guinea savanna, *Insectes Soc.*, 45(1), 51–65, doi:10.1007/s000400050068, 1998.
- 5 Korb, J. and Linsenmair, K. E.: Ventilation of termite mounds: new results require a new model, *Behav. Ecol.*, 11(5), 486–494, doi:10.1093/beheco/11.5.486, 2000.
- Korb, J.: Thermoregulation and ventilation of termite mounds, *Naturwissenschaften*, 90(5), 212–219, doi:10.1007/s00114-002-0401-4, 2003.
- 10 Korb, J.: Termite Mound Architecture, from Function to Construction, in *Biology of Termites: A Modern Synthesis*, edited by D. E. Bignell, Y. Roisin, and N. Lo, pp. 349–373, Springer, Dordrecht, 2011.
- Lepage, M. G.: L’impact des populations récoltantes de *Macrotermes michaelsoni* (Sjöstedt) (Isoptera: Macrotermitinae) dans un écosystème semi-aride (Kajiado-Kenya), I - L’activité de récolte et son déterminisme, *Insectes Soc.*, 28(3), 297–308, 1981a.
- 15 Lepage, M. G.: L’impact des populations récoltantes de *Macrotermes michaelsoni* (Sjöstedt) (Isoptera: Macrotermitinae) dans un écosystème semi-aride (Kajiado-Kenya), II - La nourriture récoltée, comparaison avec les grands herbivores, *Insectes Soc.*, 28(4), 309–319, 1981b.
- 20 Lepage, M., Abbadie, L. and Mariotti, A.: Food Habits of Sympatric Termite Species (Isoptera, Macrotermitinae) as Determined by Stable Carbon Isotope Analysis in a Guinean Savanna (Lamto, Cote d’Ivoire), *J. Trop. Ecol.*, 3(3), 303–311, 1993.
- 25 Lüscher, M.: Air-Conditioned Termite Nests, *Sci. Am.*, 205(1), 138–145, doi:10.1038/scientificamerican0761-138, 1961.
- Nauer, P. A., Hutley, L. B. and Arndt, S. K.: Termite mounds mitigate half of termite methane emissions, *Proc. Natl. Acad. Sci. - PNAS*, 115(52), 13306–13311, doi:10.1073/pnas.1809790115, 2018.
- 30 Noirot, C. and Darlington, J. P. E. C.: Termite nests: Architecture, Regulation and Defence, in *Termites: Evolution, Sociality, Symbioses, Ecology*, edited by T. Abe, D. E. Bignell, and M. Higashi, pp. 121–139, Springer., 2000.
- Ocko, S. A., King, H., Andreen, D., Bardunias, P., Turner, J. S., Soar, R. and Mahadevan, L.: Solar-powered ventilation of African termite mounds, *J. Exp. Biol.*, 220(18), 3260–3269, doi:10.1242/jeb.160895, 2017.

Pellikka, P. K. E. V., Heikinheimo, J. V., Hietanen, E. J., Schäfer, M. E., Siljander & J. M., and Heiskanen, 2018. J., Impact of land cover change on aboveground carbon stocks in Afromontane landscape in Kenya, *Applied Geography*, 94, 178–189, <https://doi.org/10.1016/j.apgeog.2018.03.017>, 2018.

5

Pomeroy, D. E.: Studies on a population of large termite mounds in Uganda, *Ecol. Entomol.*, 1(1), 49–61, doi:10.1111/j.1365-2311.1976.tb01204.x, 1976.

10

Pomeroy, D. E.: The Distribution and Abundance of Large Termite Mounds in Uganda, *J. Appl. Ecol.*, 14(2), 465–475, doi:10.2307/2402559, 1977.

Rouland-Lefèvre, C.: Symbiosis with fungi, in *Termites: Evolution, Sociality, Symbioses and Ecology*, pp. 289–306., 2000.

15

Räsänen, M., Merbold, L., Vakkari, V., Aurela, M., Laakso, L., Beukes, J. P., Zyl, P. G. V., Josipovic, M., Feig, G., Pellikka, P., Rinne, J. and Katul, G. G.: Root-zone soil moisture variability across African savannas: From pulsed rainfall to land-cover switches, *Ecohydrology*, 13(5), e2213, doi:10.1002/eco.2213, 2020.

20

Räsänen, M., Vesala, R., Rönholm, P., Arppe, L., Manninen, P., Jylhä, M., Rikkinen, J., Pellikka, P., and Rinne, J.: Dataset for "Influence of termite mound structure and habitat on the mound CO<sub>2</sub> and CH<sub>4</sub> fluxes for fungus-growing termites", figshare [data set], <https://doi.org/10.6084/m9.figshare.21739484.v1>, 2022.

Sanderson, M. S.: Biomass of termites and their emissions of methane and carbon dioxide: A global database, *Global Biogeochem. Cycles*, 10(4), 543–557, doi:10.1029/96GB01893, 1996.

25

Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, G., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentretre, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F.,

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Font: Not Italic

Formatted: Font: Not Italic



- Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q. and Zhuang, Q.: The Global Methane Budget 2000-2017, *Earth Syst. Sci. Data*, 12(3), 1561–1623, doi:10.5194/essd-12-1561-2020, 2020.
- Sieber, R. and Leuthold, R. H.: Behavioural elements and their meaning in incipient laboratory colonies of the fungus-growing Termite *Macrotermes michaelseni* (Isoptera: Macrotermitinae), *Insectes Soc.*, 28(4), 371–382, doi:10.1007/BF02224194, 1981.
- Sugimoto, A., Inoue, T., Kirtibutr, N. and Abe, T.: Methane oxidation by termite mounds estimated by the carbon isotopic composition of methane, *Global Biogeochem. Cycles*, 12(4), 595–605, doi:10.1029/98GB02266, 1998.
- Thomas, R. J.: Ecological studies on the symbiosis of *Termitomyces Heim* with Nigerian Macrotermitinae, University of London., 1981.
- Wachiye, S., Merbold, L., Vesala, T., Rinne, J., Räsänen, M., Leitner, S. and Pellikka, P.: Soil greenhouse gas emissions under different land-use types in savanna ecosystems of Kenya, *Biogeosciences*, 17(8), 2149–2167, doi:[10.5194/bg-17-2149-2020](https://doi.org/10.5194/bg-17-2149-2020), 2020.
- van Asperen, H., Alves-Oliveira, J. R., Warneke, T., Forsberg, B., de Araújo, A. C. and Notholt, J.: The role of termite CH<sub>4</sub> emissions on the ecosystem scale: A case study in the Amazon rainforest, *Biogeosciences*, 18(8), 2609–2625, doi:[10.5194/bg-18-2609-2021](https://doi.org/10.5194/bg-18-2609-2021), 2021.
- Vesala, R., Niskanen, T., Liimatainen, K., Boga, H., Pellikka, P. and Rikkinen, J.: Diversity of fungus-growing termites (*Macrotermes*) and their fungal symbionts (*Termitomyces*) in the semiarid Tsavo Ecosystem, Kenya, *Biotropica*, 49(3), 402–412, doi:[10.1111/btp.12422](https://doi.org/10.1111/btp.12422), 2017.
- Vesala, R., Arppe, L. and Rikkinen, J.: Caste-specific nutritional differences define carbon and nitrogen fluxes within symbiotic food webs in African termite mounds, *Sci. Rep.*, 9(1), 16611–16698, doi:10.1038/s41598-019-53153-x, 2019a.
- Vesala, R., Harjuntausta, A., Hakkarainen, A., Rönholm, P., Pellikka, P. and Rikkinen, J.: Termite mound architecture regulates nest temperature and correlates with species identities of symbiotic fungi, *PeerJ*, 6, e6237, doi:[10.7717/peerj.6237](https://doi.org/10.7717/peerj.6237), 2019b.
- Vesala, R., Arppe, L. and Rikkinen, J.: *Termitomyces* fungus combs - formation, structure and functional aspects, in *Microbial Symbionts: Functions and Molecular Interactions on Host*, edited by D. Dhanasekaran, Academic Press., 2022a.

Vesala, R., Rikkinen, A., Pellikka, P., Rikkinen, J. and Arppe, L.: You eat what you find – local patterns in vegetation structure control diets of African fungus-growing termites, *Ecol. Evol.*, 12, e8566, 2022b.

- 5 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nat. Methods*, 17, 261–272,  
10 <https://doi.org/10.1038/s41592-019-0686-2>, 2020.

Weir, J. S.: Air Flow, Evaporation and Mineral Accumulation in Mounds of *Macrotermes subhyalinus* (Rambur), *J. Anim. Ecol.*, 42(3), 509–520, doi:10.2307/3120, 1973.

- 15 Wildermuth, B., Oldeland, J., Arning, C., Gunter, F., Strohbach, B. and Juergens, N.: Spatial patterns and life histories of *Macrotermes michaelseni* termite mounds reflect intraspecific competition: Insights of a temporal comparison spanning 12 years, *Ecography*, 9, e06306, doi:10.1111/ecog.06306, 2022.

- Wood, T. G. and Thomas, R. J.: The mutualistic association between Macrotermitinae and Termitomyces, in *Insect-fungus interactions*, edited by N. Wilding, N. M. Collins, P. M. Hammond, and J. F. Webber, pp. 69–92, Academic Press., 1989.  
20

[Zhou, Y., Staver, A. C., and Davies, A. B.: Species-level termite methane production rates, \*Ecology\*, 104, e3905, https://doi.org/10.1002/ecy.3905, 2023.](https://doi.org/10.1002/ecy.3905)

- Zimmerman, P. R., Greenberg, J. P., Wandiga, S. O. and Crutzen, P. J.: Termites: A Potentially Large Source of Atmospheric  
25 Methane, Carbon Dioxide, and Molecular Hydrogen, *Science*, 218, 563–565, 1982.