

Distributions of bacteriohopanepolyols in lakes and coastal lagoons of the Azores Archipelago

Nora Richter^{1,2*}, Ellen C. Hopmans¹, Danica Mitrović¹, Pedro M. Raposeiro^{3,4}, Vítor Gonçalves^{3,4}, Ana C. Costa^{3,4}, Linda A. Amaral-Zettler^{1,2,5}, Laura Villanueva^{1,6}, and Darci Rush¹

¹Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, 1790 AB Den Burg, The Netherlands

²Department of Earth, Environmental and Planetary Sciences, Brown University, Providence, USA

³Centro de Investigação em Biodiversidade e Recursos Genéticos, CIBIO, InBIO Laboratório Associado, BIOPOLIS Program in Genomics, Biodiversity and Land Planning – UNESCO Chair – Land Within Sea: Biodiversity & Sustainability in Atlantic Islands, Pólo dos Açores, Portugal

⁴Faculdade de Ciências e Tecnologia da Universidade dos Açores, Ponta Delgada, Açores, Portugal

⁵Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

⁶Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands

Correspondence to: Nora Richter (nora.richter@nioz.nl)

Abstract. Bacteriohopanepolyols (BHPs) are a diverse class of lipids produced by bacteria across a wide range of environments. In this study, we aim to further identify BHPs related to ecological niches and/or specific bacteria by characterizing the distribution of BHPs in suspended particulate matter (SPM) of the water column and in sediments in a range of lakes and coastal lagoons from the Azores Archipelago, as well as in a co-culture enriched for methanotrophs. Sediment samples from Azorean lakes with low oxygen conditions during the summer months (i.e., Azul, Verde, Funda, and Negra) contain relatively high abundances of BHPs that are typically associated with methane-oxidizing (methanotrophic) bacteria (i.e., aminotetrol, aminopentol, and methylcarbamate-aminopentol), as well as the ethenolamine-BHPs (i.e., ethenolamine-BHPentol and ethenolamine-BHhexol) and the N-formylated aminoBHPs. Both ethenolamine-BHPs and N-formylated aminoBHPs were also detected in a methanotroph-methylotroph co-culture that was enriched for methanotrophs obtained from a lake. In the SPM of all water columns, bacteriohopanetetrol (BHT), BHT-cyclitol ether, and aminotriol are the dominant BHPs. In SPM from Lake Funda, nucleoside BHPs (i.e., Me-adenosylhopane_{HG-diMe}, N1-methylinosylhopane, 2Me-N1-inosylhopane, and Me-N1-inosylhopane) are present in low abundance or absent under oxic conditions but increase in concentration near the chemocline, suggesting potential *in situ* production of these nucleoside BHPs rather than an allochthonous origin. In contrast, sediments from shallow, well-mixed lakes (i.e., Empadadas, São Jorge, and Lomba) contain higher abundances of adenosylhopane and N1-methylinosylhopane, which likely originate from bacteria living in nearby soils. Based on our current results we revised the existing R_{soil} index, which was previously used to infer terrestrial inputs to aquatic environments, to exclude any potential nucleosides produced in the lake water column ($R_{\text{soil-lake}}$). In the coastal lagoons, Cubres East and West, methoxylated-BHTs were detected, and higher abundances of ethenolamine-BHT were observed. This study highlights the diversity of BHPs in lakes and coastal lagoons and their potential as taxonomic markers for bacteria associated with certain ecological niches, which can be preserved in sedimentary records.

Formatted: Subscript

Formatted: Subscript

Formatted: Font color: Red

1 Introduction

Bacteriohopanepolyols (BHPs) are pentacyclic triterpenoids found in the cell membrane of many gram-negative and gram-positive bacteria (Rohmer et al., 1984). Hopanoid derivatives of BHPs are considered to be some of the most abundant lipids on Earth and in the geologic record (Ourisson and Albrecht, 1992). BHPs are involved in various cell physiological processes (Welander et al., 2009; Sáenz, 2010; Doughty et al., 2011; Sáenz et al., 2012; Welander and Summons, 2012) and, due to their structural diversity, show potential as chemotaxonomic markers (Kusch and Rush, 2022). Functionalized BHPs were identified in sediment records that span 1.2 Ma (Talbot et al., 2014) as well as in Eocene Cobham Lignite (ca. 56 Ma; Talbot et al., 2016), highlighting their preservation potential as biomarkers in sedimentary records.

Recent studies and advancements in BHP analysis highlight the wide distribution and diversity of BHPs in both marine and terrestrial environments (Talbot and Farrimond, 2007; Pearson et al., 2009; Sáenz et al., 2011; Kusch et al., 2019; Hopmans et al., 2021). Bacteriohopanetetrol (BHT), BHT-cyclitol ether (CE), and aminotriol, for instance, are ubiquitous in environmental samples and culture studies, and are not associated with any specific organisms or environments (e.g., Talbot et al., 2003; Talbot and Farrimond, 2007; Zhu et al., 2011). Nucleoside BHPs, (formerly known as adenosylhopanes) are typically associated with soils, and are used to trace soil inputs to riverine and marine environments using a ratio of nucleosides to the more ubiquitous BHT, known as the R_{soil} index (Taylor and Harvey, 2011; Zhu et al., 2011; De Jonge et al., 2016; Kusch et al., 2019). Recent studies, however, show that certain nucleoside BHPs can also be produced in the marine environment under low-oxygen conditions, complicating interpretations of the R_{soil} index (Kusch et al., 2021b). Hopanoids methylated at the C-2 position are degradation products of 2-methylated-BHPs (e.g., 2Me-BHT), and are often considered diagnostic for cyanobacteria in the geologic record (Summons et al., 1999). 2Me-BHPs are produced by cyanobacteria (Talbot et al., 2008), however, the gene responsible for BHP methylation at the C-2 position was identified in a wide-range of other bacterial phyla (Welander et al., 2010), suggesting multiple biological sources. This highlights the complexity of BHPs in the environment, and the need for more studies to evaluate the potential of BHPs as biomarkers for specific bacterial groups.

Aerobic methane-oxidizing bacteria (MOB) regulate methane emissions in seasonally stratified lakes (Bastviken et al., 2008; Oswald et al., 2015; Guggenheim et al., 2020) and meromictic lakes (Oswald et al., 2016) by oxidizing methane before it reaches the atmosphere. MOB in lacustrine environments typically fall within the two phylogenetic groups in the phylum Proteobacteria: Type I (members of Gammaproteobacteria) and Type II (members of Alphaproteobacteria; Hanson and Hanson, 1996). They are known to produce a wide-range of potentially diagnostic BHPs (Rohmer et al., 1984), including aminotetrol and aminopentol (Neunlist and Rohmer, 1985b, c; Cvejjic et al., 2000; Talbot et al., 2001; van Winden et al., 2012), the recently identified methylcarbamate-aminoBHPs (Rush et al., 2016), and related 3 β -methylated-BHPs (Neunlist and Rohmer, 1985b; Zundel and Rohmer, 1985; Cvejjic et al., 2000). Type I methanotrophs are typically associated with increased production of aminopentol, whereas Type II methanotrophs are distinguished by the production of aminotetrol (Neunlist and Rohmer, 1985b, c; Cvejjic et al., 2000; Talbot et al., 2001; van Winden et al., 2012). However, recent environmental studies, particularly in marine environments, suggest that this distinction is not as clear as previously proposed (Rush et al., 2016; Kusch and Rush, 2022). In addition, m~~However, m~~inor amounts

Field Code Changed

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Dutch (Netherlands)

of these BHPs aminotetrol and aminopentol are also produced by sulfur-reducing bacteria and methylcarbamate-aminotriol was found in cultures of nitrite-oxidizing bacteria (Blumenberg et al., 2006; Elling et al., 2022). Few studies have evaluated the BHP composition of lacustrine methanotrophs nor the utility of these biomarkers for tracing MOB in lakes.

In addition to their potential as taxonomic markers, BHP relative abundance was observed to change under certain environmental conditions. For instance, microcosm and mesocosm experiments that enriched for methanotrophs demonstrate that increasing temperatures led to increased concentrations of aminotriol, aminotetrol, and aminopentol (Osborne et al., 2017; van Winden et al., 2020), whereas a decrease in the relative abundance of unsaturated hopanoids was observed in culture experiments (Bale et al., 2019). Shifts in BHP compositions were also used to interpret changes in salinity (Coolen et al., 2008), redox conditions (Blumenberg et al., 2013; Matys et al., 2017; Rush et al., 2019; Zindorf et al., 2020), and soil inputs (Blumenberg et al., 2013) in sediment records. BHPs and BHP-producers are particularly diverse in lakes (Farrimond et al., 2000; Watson and Farrimond, 2000; Talbot et al., 2003; Talbot and Farrimond, 2007; O’Beirne et al., 2022), yet few studies have associated BHPs with specific lacustrine settings.


The purpose of this study is to characterize the distribution of BHPs in lakes and coastal lagoons with the aims of identifying BHPs that are specific to lacustrine bacteria, with a focus on MOB, and/or specific ecological niches, and of testing whether nucleoside BHPs are produced *in situ* in lakes. We also analyzed BHPs from a methanotrophic-methylotrophic co-culture enriched from a lake water column to identify potential novel BHPs related to lacustrine methanotrophy. We also collected and analyzed sediment and water column suspended particulate matter (SPM) from eight lakes and two coastal lagoons on three different islands in the Azores Archipelago. The Azores Archipelago consists of nine islands with 88 lakes that occur in either topographically depressed areas or volcanic depressions, and range from severely impacted by humans to relatively “pristine” (Pereira et al., 2014). The diversity in lake-types and comparison with marine-influenced sites, i.e. the coastal lagoons, in this study provide an ideal setting to investigate the partitioning of BHPs in the environment and their potential as biomarkers.

2 Materials & Methods

2.1 Sample Collection

Samples were collected in lakes and lagoons on three different islands in the Azores Archipelago in June 2018 (Fig. 1 & Table 1): São Miguel (lakes Azul, Verde, and Empadadas), São Jorge (lake São Jorge and lagoons Cubres East and Cubres West), and Flores (lakes Funda, Lomba, and Negra). June is considered part of the dry season in the Azores, which ranges from April to August. From September to March, the Azores Archipelago receives more rain and is exposed to high winds (Santos et al., 2004; Hernández et al., 2016). During the summer months, deeper lakes such as Azul, Verde, Funda, and Negra stratify and the bottom water becomes suboxic and, in some years, even fully anoxic (Gonçalves, 2008; Gonçalves et al., 2018). In contrast, shallow lakes are typically considered polymictic but might undergo short periods of stagnation during the summer months

Field Code Changed



110 Figure 1: (A) The Azores Archipelago with the location of lakes and coastal sites discussed in this study. Our study sites include (B) lakes Azul, Verde, and Empadadas Norte on São Miguel Island, (C) São Jorge lake and the coastal lagoons (i.e., Cubres East and Cubres West) on São Jorge Island, and (D) lakes Negra, Lomba, and Funda on Flores Island (maps generated in ArcGIS 10.3).

115 **Table 1: Location and characteristics of lakes and lagoons discussed in this study. The annual average water column properties are based on long-term monitoring data from 2003-2017 (University of the Azores monitoring program) where TP = total phosphorous and TN = total nitrogen.**

Site information					Annual Average Water Column properties					
Island	Lake/Lagoon Site	Lat. (°N)	Lon. (°W)	Alt. (mas)	Max. Depth (m)	Trophic State*	Temp. (°C)			
							pH	TP (µg/L)	TN (µg/L)	
São Miguel	Azul	37.87	-25.78	260	25.4	Mesotrophic	16.55	7.51	23.13	0.43
São Miguel	Verde	37.84	-25.79	262	23.5	Eutrophic	15.56	7.74	44.96	0.59
São Miguel	Empadadas Norte	37.82	-25.75	762	3.3	Eutrophic	14.3	6.90	22.67	0.45
São Jorge	São Jorge	38.65	-28.07	900	2.5	-	-	-	-	-
São Jorge	Cubres East	38.64	-27.97	3	2.0	-	-	-	-	-
São Jorge	Cubres West	38.64	-27.97	3	2.0	-	-	-	-	-
Flores	Funda	39.41	-31.22	364	31.9	Eutrophic	14.69	7.72	41.84	0.62
Flores	Lomba	39.43	-31.19	651	15.3	Mesotrophic	14.63	7.01	24.16	0.38
Flores	Negra	39.44	-31.23	540	115.0	Eutrophic	14.66	7.99	98.98	0.60

*University of the Azores monitoring program (2003-2017); Cordeiro et al. (2020)

120 (Gonçalves, 2008). The lakes sampled for this study are either mesotrophic or eutrophic and range in water column depth from 2 to 115 m. During the summer months, lakes Verde and Funda experience large cyanobacterial blooms (Cordeiro et al., 2020). Both lakes Verde and Funda lakes were experiencing a cyanobacterial bloom at the time of sampling.

125 Suspended particulate matter was collected from the surface water, near the oxycline/chemocline at the time of sampling, and from the bottom water of the lake without disturbing the sediment-water interface. Water (5 to 10 L) was immediately filtered through Whatman GF/F 142 mm 0.7 µm pore size filters and frozen in liquid nitrogen. Surface sediment samples (0-1 cm) were also obtained at the same time using a gravity corer (UWITEC-90mm, Austria). In Azul and Verde, surface sediments were collected near the lake shore and in the deepest part of the lake. Replicate samples were collected for sediment cores from lakes Azul (n = 2), Empadadas (n = 4) and Negra (n = 2) and analyzed for reproducibility. All samples were immediately frozen in a liquid nitrogen dewar and transported to the NIOZ Royal NIOZ-Netherlands Institute for Sea Research, where SPM and sediment samples were stored at -80°C and -40°C, respectively, until analysis. At the time of sample collection, water column profiles were obtained using a Hydrolab (Horiba-U50) for dissolved oxygen content (percent saturation and mg/L), temperature, salinity, conductivity, pH, and turbidity (NTU). Additional long-term data was obtained from the Universidad dos Açores long-term monitoring program (data from 2003-2020).

2.2 Cultivation of Methanotroph-Methylotroph co-culture

140 An enrichment co-culture consisting of a methanotroph (43% *Methylobacter* sp.) and a methylotroph (21% *Methylotenera* sp.) was previously isolated from a seasonally stratified and hypereutrophic lake (Lacamas Lake, WA, USA; van Grinsven et al., 2020). This culture was grown in triplicate under oxic conditions in the dark at 15 °C on a nitrate mineral salts (NMS) medium (Whittenbury et al., 1970). For lipid extractions, the cultures were grown on 250 mL NMS media in 580 mL acid-washed and autoclaved glass pressure bottles with butyl rubber stoppers with an addition of methane (16 mL, 99.99% pure) to the headspace. After the methane gas was consumed, the cultures were filtered onto muffled Whatman GF/F 47 mm 0.3 µm pore size filters and frozen at -80 °C. The methylotroph identified in the co-culture is closely related to *Methylotenera versatilis* and *Methylotenera mobilis* (van Grinsven et al., 2020). Therefore, we obtained freeze-dried biomass of *Methylotenera*

Formatted: Font: 7 pt

Formatted Table

Formatted: Font: 7 pt, Bold

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

mobilis (DSM 17540) from the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) culture collection to investigate the presence of BHPs in this methylotroph.

2.3 Lipid extraction

All samples were freeze-dried and extracted using a modified Bligh-Dyer method (Bligh and Dyer, 1959; Bale et al., 2021). Samples were submerged in methanol (MeOH), dichloromethane (DCM), and phosphate buffer (2:1:0.8, v:v:v) and ultrasonically extracted twice. The solvent was collected in a separate flask after each extraction. DCM and phosphate buffer were added to the resulting solvent to obtain a new volume ratio of 1:1:0.9 (v:v:v). The DCM layer was collected and the aqueous layer was washed two more times with DCM. The extraction was repeated using MeOH:DCM:aqueous trichloroacetic acid solution (2:1:0.8, v:v:v) following the same procedure described above. The combined DCM layers were dried under N₂ gas and stored at -20 °C until analysis. Prior to analysis, deuterated diacylglyceryltrimethylhomoserine (DGTS D-9; Avanti® Polar Lipids, USA) was added to the extracts as an internal standard and the samples were re-dissolved in MeOH:DCM (9:1, v:v) before filtering the samples through a 0.45 µm regenerated cellulose syringe filter (4 mm diameter; Grace Alltech, Deerfield, IL).

2.4 UHPLC/HRMS analysis

Samples were analyzed following the methods described in Hopmans et al. (2021). Briefly, samples were analyzed on an Agilent 1290 Infinity I UHPLC coupled to a quadrupole-orbitrap HRMS (Q-Exactive, ThermoFisher Scientific, Waltham, MA) equipped with an Ion Max source and heated ESI probe (HESI) (ThermoFisher Scientific, Waltham, MA). Separation was achieved on an Acquity C18 BEH column (2.1 x 150 mm, 1.7 µm particle; Waters) and pre-column. The solvent system consisted of (A) MeOH:H₂O (85:15) and (B) MeOH:isopropanol (1:1) with both containing 0.12 % (v/v) formic acid and 0.04 % (v/v) aqueous ammonia. Lipids were detected using positive ion monitoring of *m/z* 350-2000 (resolution 70,000 ppm at *m/z* 200) with an inclusion list of calculated exact masses of BHPs: 171 for sediment samples (~~one Lake Verde sediment extract was re-run with 421 exact masses for confirmation~~) and 357 for water column samples. ~~Note, that the difference in the number of exact masses in the inclusion lists reflects the addition of further novel BHPs after their identification. A Lake Verde sediment sample (Verde SS1) that included all the novel BHPs was re-run with an updated inclusion list (421 exact masses) for confirmation of novel BHPs.~~ For untargeted BHP detection and identification, we used data dependent MS² (isolation window 1 *m/z*; resolution 17,500 ppm at *m/z* 200) of the 10 most abundant ions for a total cycle of ca. 1.2 s and dynamic exclusion (6 s) with a 3 ppm mass tolerance. To obtain optimal fragmentation of BHPs, we used a stepped normalized collision energy of 22.5 and 40. Mass calibration was performed every 48 h using a Thermo Scientific Pierce LTQ Velos ESI Positive Ion Calibration Solution.

BHPs were identified based on their retention time, exact mass and fragmentation spectra. Integrations were performed on (summed) mass chromatograms (within 3 ppm mass accuracy) of relevant molecular ions ([M+H]⁺, [M+NH₄]⁺, and [M+Na]⁺). We used an internal standard for normalization between sample runs ~~by calculating the average peak area of the internal standard and correcting the peak areas to the average internal standard.~~ However, we do not have an internal standard for BHP quantification, therefore all BHPs are reported as response

Formatted: Font color: Red

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Red

units (RU). BHPs identified in water column and surface sediment samples are normalized to liters of water filtered (L) and grams of freeze-dried sediment (g), respectively.

2.5 Elemental analysis

Total organic carbon (TOC) and total nitrogen (TN) content was measured for the surface sediments (n=18) as described in (Mitrović et al., (2023). Briefly, the samples were decalcified, powdered, and freeze-dried prior to analysis. All samples were analyzed using an Elementar Vario Isotope Cube ~~directly linked coupled to an~~ Isotope Ratio Mass Spectrometer (IRMS) Elementar Isoprime vision. ~~Samples for total organic carbon (TOC) Prior to TOC analysis, carbonates were removed by pretreating the samples were pretreated~~ with excess 2M hydrochloric acid (HCl) ~~to remove carbonates and were put~~ on a shaker overnight. The resulting samples were neutralized with bidistilled water and freeze-dried. ~~All TOC samples were~~ measured in duplicate with a reproducibility of <0.5 %.

~~$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of bulk organic matter were measured on decalcified samples of surface sediments (n=18) using an Elementar Analyser (Elementar Vario Isotope Cube) coupled to an IRMS (Elementar Isoprime vision). Stable isotope ratios of ^{13}C and ^{15}N are expressed using the δ notation in units per mil, reported relative to Vienna Pee Dee belemnite (V-PDB) and atmospheric N_2 , respectively. The results were normalized to certified standards and standard deviations were <0.05 ‰.~~

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Indent: First line: 1.27 cm

Formatted: Font color: Auto, Superscript

Formatted: Font color: Auto, Superscript

Formatted: Font color: Auto

Formatted: Font color: Red

200

2.6 R_{soil} Index

A soil index (R_{soil}) was developed to trace inputs of terrestrial-derived BHPs into the marine environment (Zhu et al., 2011). This index relies on the observations that BHT is present in higher abundance relative to nucleosides in the marine environment than in the soils (Pearson et al., 2009; Rethemeyer et al., 2010; Zhu et al., 2011). The index is calculated as follows:

$$R_{\text{soil}} = \frac{(\text{soil-marker BHPs})}{(\text{soil-marker BHPs} + \text{BHT})} \quad (1)$$

Previously, “soil-marker BHPs” included only adenosylhopane, N1-methylinosylhopane (i.e., type-2 adenosylhopane after Hopmans et al., 2021), adenosylhopane_{HG-Me} (i.e., type-3 adenosylhopane after Hopmans et al., 2021) and their ring-methylated counterparts. BHT in the R_{soil} equation only refers to the isomer with 17, 21(H), 22R, 32R, 33R, 34S stereochemistry. Previous high latitude studies have revised this index, R'_{soil} , to exclude methylated nucleosides (Doğrul Selver et al., 2012):

$$R'_{\text{soil}} = \frac{(\text{adenosylhopane} + \text{N1-methylinosylhopane} + \text{adenosylhopane}_{\text{HG-Me}})}{(\text{adenosylhopane} + \text{N1-methylinosylhopane} + \text{adenosylhopane}_{\text{HG-Me}} + \text{BHT})} \quad (2)$$

In this study, we tested both of these indices at our sample sites.

2.7 Data visualization and statistical analyses

220 Non-metric multidimensional scaling (NMDS) analysis was conducted using a Bray-Curtis dissimilarity matrix
to visualize the similarities between BHP compositions at our sample sites. All analyses were performed and
visualized in R (version R-4.2.1; R Core Team, 2022) using the vegan package (version 2.5-7; Oksanen et al.,
2020) and ggplot2 (version 3.3.5; Wickham, 2016). NMDS plots were generated for both the water column and
225 sediment samples. To test whether there was a significant difference between our sample sites and sample site-
types (for sediment samples: deep lakes, shallow lakes, and coastal environments; for water column: surface water,
chemocline, and bottom water), we performed analysis of similarities (ANOSIM) tests in R using the vegan
package for the sediment samples and water column samples. A Bray-Curtis dissimilarity matrix was used for
these tests.

3 Results & Discussion


3.1 Diversity of BHPs in lake surface sediments & water columns

230 At our study sites, we identified 83 BHPs in total (Table A1). For the identification of novel BHPs, we discuss
the carbon positions, rings, and modifications to the side-chain of the core BHP structure as shown in Fig. B1.
The novel BHPs are briefly discussed here and described in more detail in Appendix B. In the Azorean lakes,
BHPs were more diverse in surface sediments (average 51 BHPs/sediment sample) than in the water column SPM
(average 11 BHPs/SPM; Fig. 2).

235 Nucleoside BHPs were particularly diverse in the water column SPM and surface sediments in lakes Empadadas,
São Jorge, and Lomba. We identified fourteen nucleoside BHPs, including three inosylhopanes, described by
Hopmans et al. (2021; Appendix B, Fig. B2 & B3). Further, we identified five additional adenosyl-BHP isomers
(Fig. B2), including early eluting isomers (peaks b*, f*, g*, and k*). We also detected four additional inosylhopane
240 isomers in surface sediments from Lake Empadadas with modifications in the position of the methyl group on the
ring structure and/or head group (Fig. B3: peaks b*, c*, f*, and g*).

BHT (17, 21(H), 22R, 32R, 33R, 34S stereochemistry) was predominant in most samples, along with unsaturated
and methylated versions of BHT. In the surface sediment from the coastal lagoons, Cubres East and West, we
245 identified two compounds comparable to BHT, however, in the MS² spectrum we observe the loss a methoxy
moiety (32 Da, CH₃OH), as well as three hydroxyl moieties (Appendix B2, Fig. B4). We tentatively identify these
compounds as methoxylated-BHTs. BHT-cyclitol ether (CE) was also abundant in both water column SPM and
surface sediments (Fig. 2). In addition, we detected MeBHT-CE in surface sediments from Lake Verde, as well
as an isomer where, based on the MS² spectrum, the methylation occurs on the cyclitol ether head group (Appendix
250 B3, Fig. B5). We putatively identify this isomer as BHT-MeCE. BHpentol, BHpentol-CE, BHhexol, and
BHhexol-CE were also identified at our sites but only in the lake surface sediments.

Aminotriol, as well as isomers of aminotriol and unsaturated aminotriol, were abundant in the water column SPM
and surface sediment of the lakes (Fig. 2). Methylated and acylated-versions of aminotriol were also identified,
255 but only in the surface



Commented [NR1]: Changed figure - noticed names for oxazinone and dioxanone were labelled as unknown composites and changed formulated to N-formylated

260 **Figure 2:** (A) Relative abundance of all BHPs found in surface sediment samples labelled with the lake depth (meters) at which they were collected where Emp. = Empadadas, SJ = São Jorge, CE = Cubres East, and CW = Cubres West (all raw data is available in the supplementary material). (B) The response units (RU) of BHPs in the sediment samples normalized to g of dry sediment and (C) the number of BHPs identified in each sample. (D) Relative abundance of BHPs identified in water column samples from the different lakes and lagoons (E) with the total response units/L of water filtered and (F) number of BHPs shown.

265

sediment of the lakes. In contrast to a previous study in a hyper-euxinic and meromictic lake, we do not observe any diunsaturated-aminotriols in the SPM or surface sediments at our study sites (O'Beirne et al., 2022).

270 Aminotetrol and aminopentol, compounds commonly associated with methanotrophic activity (Neunlist and Rohmer, 1985b; Zundel and Rohmer, 1985; Cvejic et al., 2000), were found in almost all of the water column SPM and surface sediments. However, acylated versions of aminopentol (C_{14:0}, C_{15:0}, and C_{16:0}) were only

observed in the surface sediments of Azul and Verde. Two isomers of methylcarbamate (MC)-aminotriol and MC-aminopentol, compounds also associated with methanotrophs, were identified in water column SPM and surface sediment after Rush et al. (2016). The recently described propenolamine-BHT, ethenolamine-BHT, ethenolamine-BHpentol, and ethenolamine-BHhexol (Fig. 3) were identified in our samples after Hopmans et al. (2021). In addition, we putatively identified unsaturated versions of ethenolamine-BHT (Appendix B4, Fig. B6) and ethenolamine-BHhexol (Fig. 3e) with the double bond likely occurring at the Δ^6 and Δ^{11} positions, respectively, based on the retention times. Acylated versions of ethenolamine-BHhexol ($C_{15.0}$, $C_{16.0}$, and $C_{17.0}$) were also observed in lake surface sediments.


In sediment from Lake Verde (Fig. 3, peaks j and m), we identified several unknown composite BHPs previously described in Hopmans et al. (2021) with an assigned elemental composition (AEC) of $C_{36}H_{64}O_4N^+$ (m/z 574.483) and $C_{36}H_{64}O_5N^+$ (m/z 590.483), but no structure was assigned. In addition, we identified the same compound but with an additional hydroxy moiety (Fig. 3, peak o).

~~In Hopmans et al. (2021) several unknown composite BHPs were described with an assigned elemental composition (AEC) of $C_{36}H_{64}O_4N^+$ (m/z 574.483) and $C_{36}H_{64}O_5N^+$ (m/z 590.483), respectively, but no structure was assigned. Here, we detected these novel BHPs in sediment from Lake Verde (Fig. 3, peaks j and m) based on the diagnostic loss of 28 Da (CO) in the MS²-spectra. We also identified the same compound with an additional hydroxy moiety on the side chain in the partial mass chromatogram of m/z 606.473 (elemental composition (EC) $C_{36}H_{64}O_6N^+$; Fig. 3, peak o). We propose that these novel composite BHPs are a series of N-formylated-aminoBHPs: N-formylated-aminotriol (peak j), N-formylated-aminotetrol (peak m) and N-formylated-aminopentol (peak o). The proposed structure for N-formylated-aminopentol is shown in Fig. 3f with the diagnostic fragmentations indicated. In addition, we observe unsaturated versions of N-formylated-aminotriol (Fig. 3b & Appendix B5, Fig. B7), N-formylated-aminotetrol (based on the MS¹ and retention time), and N-formylated-aminopentol with the double bond likely occurring at the Δ^6 position for N-formylated-aminotriol and Δ^{11} position for N-formylated-aminotetrol and N-formylated-aminopentol. In the mass chromatogram for m/z 572.467 (Fig. 3b) from the Lake Verde sediment, we also observe a later-eluting peak ($C_{36}H_{62}O_4N^+$; Fig. 3b and d, peak l). Based on the MS² spectrum, we propose that this is the same compound identified by Elling et al. (2022) in nitrite-oxidizing bacteria. The authors describe an acetylated compound (m/z 656.495; $C_{40}H_{66}O_6N^+$), with the unique loss of 44 Da (CO₂), the loss of two hydroxyl moieties, and an additional loss of 17 Da (NH₃⁺). This compound is likely a cyclized form of N-formylated-aminotriol, and we putatively identify it as oxazinone-aminotriol (Appendix B6).~~

Commented [NR2]: Condense this paragraph on fomrylated-aminoBHPs

In addition to the discussed Δ^6 -ethenolamine-BHT, we observed a later eluting peak at 21.88 mins (Fig. 3a, peak c) in the mass chromatogram of m/z 586.483 ($C_{37}H_{64}O_4N^+$; Δ ppm 0.72). We propose that one of the functionalities is part of a cyclized structure and this BHP is a N-formylated aminotetrol, where the formic acid is part of a cyclic structure and the terminal amino group is methylated. The proposed structure with key fragmentations is shown in Fig. 3c. We tentatively identify this compound as a dioxanone-methylaminotriol (Appendix B7). The MS² spectra is characterized by the distinct loss of 28 Da and a loss of 31 Da (CNH₃). Based on the elemental

composition there appears to be an additional DBE (double bond equivalent) in the structure, but there is no indication that the unsaturation is located in the core structure as the lower mass range of the



Commented [NR3]: Changed figure: removed m/z from losses on structures and changed name from formylated-aminopentol to N-formylated-aminopentol

315

320

325

Figure 3: (A) Partial mass chromatograms of ethenolamine-BHT (peak a), Δ^6 -ethenolamine-BHT (peak b), dioxanone-methylaminotriol (peak c), ethenolamine-BHPentol (peak d), isomers of methylcarbamate aminotriol (peaks e and f), a potential peak for unsaturated ethenolamine-BHPentol (peak g), ethenolamine-BHhexol (peak h), and Δ^{11} -ethenolamine-BHhexol (peak i). (B) Partial mass chromatograms of N-formylated-aminotriol (peak j), Δ^6 -N-formylated-aminotriol (peak k), oxazinone-aminotriol (peak l), N-formylated-aminotetrol (peak m), Δ^{11} -N-formylated-aminotetrol (peak n), N-formylated-aminopentol (peak o), and Δ^{11} -N-formylated-aminopentol (peak p). MS² spectrum of (C) the dioxanone-methylaminotriol in m/z 586.483 (peak c) and (D) the oxazinone-aminotriol composite (peak l) with tentative structures shown. MS² spectrum of (E) [M+H]⁺ Δ^{11} -ethenolamine-BHhexol and (F) [M+H]⁺ N-formylated-aminopentol with tentative structures shown. The chromatograms are from a surface sediment sample from Lake Verde.


fragmentation spectrum is similar to what is expected for BHT. We therefore assume one of the functionalities is part of a cyclized structure and propose this BHP is a formylated aminotetrol, where the formic acid is part of a

eyelic structure and the terminal amino group is methylated. The proposed structure with key fragmentations is shown in Fig. 3c. We tentatively identify this compound as a dioxanone-methylaminotriol (Appendix B7):

3.2 Distribution of BHPs depend on environmental conditions in lakes and coastal lagoons

Based on the NMDS visualization, the more complex BHP distributions in the surface sediments allow these sampling sites to be placed into three lake types (Fig. 4): deep lakes (i.e., Azul, Verde, Funda, and Negra; circles), shallow lakes (i.e., Empadadas, São Jorge, Lomba; squares), and coastal lagoons (i.e., Cubres East and Cubres West; stars). Using ANOSIM we find a significant difference both between the sample sites ($R = 0.89, p < 0.01$) and the sample site-types ($R = 0.72, p < 0.001$). The variability between our sites, particularly in the surface sediments, is reflected in the overall differences in BHP distributions. In Azul and Verde surface sediments collected near the shore and in the deepest part of the lake show comparable BHP distributions. We also observe comparable BHP distributions in sample replicates for lakes Azul, Empadadas, and Negra, thus we will only discuss the average BHP values for the replicate sites. In the water column NMDS plot (Fig. 5) we observe the bottom water samples cluster closer together, whereas the samples from the surface water and chemocline overlap. Using the ANOSIM test, we find ~~there is~~ no significant difference in BHP distributions between the surface water, chemocline, and bottom water of these lakes ($R = 0.04, p = 0.32$) or between the water columns of different sites ($R = 0.06, p = 0.35$).

The deep lakes are monomictic and undergo stratification from June-October, with hypoxic and even anoxic conditions occurring during the summer months in the hypolimnion (Gonçalves, 2008; Raposeiro et al., 2018). These lakes contain a high abundance of BHT, BHT-CE, and aminotriol, particularly in the surface waters and sediment samples (Fig. 2 & Tables A3-A4). Previous studies that identified BHPs in lake sediments also described a high abundance of BHT, BHT-CE, and aminotriol (Farrimond et al., 2000; Talbot and Farrimond, 2007). Similarly, BHP distributions in the water column of a hypereutrophic, meromictic lake, Lake Mahoney (Canada), are dominated by BHT, unsaturated BHT, aminotriol, and BHT-CE in the surface waters and BHT, diunsaturated aminotriol, and aminotriol in the surface sediments (O'Beirne et al., 2022). In the Azores, the deep lakes (circles) are further distinguished by the presence of amino-containing BHPs in the surface sediments (Fig. 2a) that cluster together in the NMDS plot (Fig. 4), including: aminotetrol, aminopentol, MC-aminopentol (green), N-formylated-aminotetrol, N-formylated-aminopentol (gray), ethenolamine-BHpentol, and ethenolamine-BHhexol (gold). This is also reflected in the BHP distributions of the bottom water SPM samples collected in June 2018 (Fig. 6), in which we observe amino-containing BHPs (i.e., aminopentol, ethenolamine-BHhexol, aminopentol, N-formylated-aminopentol) in samples collected below the oxycline. A higher abundance of penta- and hexa-functionalized BHPs, including aminopentol, ~~was~~ previously described in eutrophic lakes relative to lakes with low primary productivity (Farrimond et al., 2000; Talbot and Farrimond, 2007).



Commented [NR5]: Modified figure: changed BHP colors and legend


Figure 5: NMDS visualization of BHP distributions between different water column samples from lakes Azul, Empadadas, Funda, and Negra, as well as Cubres East and West. Note: **f**or Empadadas, Cubres East, and Cubres West we only have surface water column samples since the sampling locations were less than 3.5 m deep.

of BHT, aminotriol, and BHT-CE, and even aminopentol (Talbot and Farrimond, 2007). Similar BHP distributions are observed in the sediments of Empadadas, São Jorge, and Lomba.

385

Finally, Cubres East and West are shallow, coastal lagoons that are influenced by both marine and freshwater inputs. Cubres East (salinity 9.7 ppt) and West (salinity 23.8 ppt) are defined by a high relative abundance of ethenolamine-BHT and propenolamine-BHT, and by the presence of two methoxylated BHPs. In contrast to our freshwater sites, BHT-CE is mainly absent from the coastal lagoons (except for in the surface sediments of Cubres West). Similarly, in land-to-sea transects from the Yangtze (China) and Yenisei (Siberia) Rivers, BHT-CE was more abundant in rivers, estuaries, and the coast relative to the open water (Zhu et al., 2011; De Jonge et al., 2016). Marine samples and fjord samples, appear to predominantly consist


390



Commented [NR6]: Changed figure: changed label from unknown composite to dioxanone-methylaminotriol and from formylated to N-formylated

395 Figure 6: Water column profiles for Lakes Azul (A-E), Funda (F-J), and Negra (K-M). Plots A, F, and K show the relative units/liter (RU/L) of nucleoside BHPs identified in the water column. Plots B, G, and L show the relative response units/liter (RU/L) of BHPs associated with methane-oxidizing bacteria found in the water column. Plots C, H, and M display the response relative units/liter (RU/L) of ethenolamine-BHPs and recently identified composite BHPs (i.e., N-formylated-aminoBHPs and dioxanone-methylaminotriol) identified in the water column. D and I show the turbidity (blue) and dissolved oxygen (pink/black) from the water column at the time of sampling in June 2018. Similarly, E and J display the temperature (blue) and pH (pink/black) profiles of the water column at the time of sampling for both Lakes Azul and Funda.

405



410 **Figure 7: (A) Relative abundance of nucleosides identified in the surface sediment samples for all Azorean lakes (where Emp = Empadadas; SJ = São Jorge, CE = Cubres East; CW = Cubres West) with the (B) R_{soil} (see equation 1), R'_{soil} , and $R_{soil-lake}$ calculated for each sample. (C) The distribution of nucleosides in water column samples and their respective (D) R_{soil} , R'_{soil} , and $R_{soil-lake}$ values are also shown. The red bars indicate the terrestrial end-member values. Note: nucleosides were not detected in the water column SPM of Cubres West.**

415 of BHT and aminotriol (Farrimond et al., 2000; Zhu et al., 2011; Kusch et al., 2021a). In the estuary and coastal samples from the Yangtze River drainage basin, the BHP distributions consist of BHT, 2MeBHT, aminotriol, aminotetrol, aminopentol, nucleoside BHPs, and BHT-CE (Zhu et al., 2011). In general, we observe a similar distribution of BHPs in Cubres East and West.

420 Aminotriol and BHT were the most abundant BHPs at all of our sample sites. The presence of more diverse and abundant amino-BHPs in the deeper and seasonally stratified lakes allow us to distinguish them from the shallower lakes that contain a higher abundance of nucleoside BHPs. Finally, the lower diversity of BHPs and unique presence of methoxylated BHPs in Cubres East and West, separates the marine influenced sites from the rest of the lakes in this study.


425 3.3 Diversity of BHPs in a methanotroph-methylotroph co-culture enriched from a lake water column

Previously, certain BHPs identified in MOB cultures were also detected in lake settings (Talbot et al., 2001; Talbot and Farrimond, 2007; Rush et al., 2016; Neunlist and Rohmer, 1985b, c; Cvejic et al., 2000; O'Beirne et al., 2022). However, neither methanotrophs nor lacustrine settings have been analyzed since ~~the~~ recent methodological advancements have expanded BHP identification (Hopmans et al., 2021).

430 Here, we investigated the BHP composition of a lacustrine methanotroph-methylotroph (*Methylobacter-Methylotenera*) enrichment co-culture isolated from Lacamas Lake, WA, U.S.A. (van Grinsven et al., 2020). BHPs (i.e., aminotriol, 3Me-aminotriol, MC-aminotriol, aminotetrol, 3Me-aminotetrol, MC-aminotetrol, aminopentol, unsaturated-aminopentol, 3Me-aminopentol, MC-aminopentol) have been previously detected in
435 cultures of gammaproteobacterial Type I MOB *Methylobacter* (Osborne, 2015; Rush et al., 2016). Our aim was to potentially identify novel BHPs from a lacustrine enrichment dominated by *Methylobacter*. This enrichment is also abundant in the methylotroph *Methylotenera* (van Grinsven et al., 2020). In order to rule out the synthesis of BHPs by this methylotroph, we investigated its genomic capacity to synthesize BHPs by searching for the key gene responsible for the formation of hopane, and therefore BHP (i.e., squalene hopene cyclase). A protein blast
440 search (NCBI) was performed with the sequence of the squalene hopene cyclase gene of *Bradyrhizobium japonicum* as a query (accession number WP_038942977.1), which did not lead to any hits in the genomes of *Methylotenera* species. In addition, we analyzed the BHP composition of *Methylotenera mobilis*, a closely related strain to the methylotroph also present in the co-culture, which was available in the DSMZ culture collection. Analysis of the *M. mobilis* biomass did not lead to the detection of BHPs. We acknowledge that other bacteria
445 were also present in the enrichment culture (see van Grinsven et al., 2020); however, their BHP contribution is likely minor compared to the more abundant *Methylobacter* sp. present.

~~On the other hand, i~~n the *Methylobacter-Methylotenera* co-culture we identified 22 BHPs (Table A4), with the most abundant being aminotriol (43 %) and aminopentol (43 %) (Fig. 8). Aminotetrol (6 %) was also observed,
450 along with the unsaturated and acylated forms of aminotriol and aminopentol. Although present in low abundance, we identified MC-aminotriol, MC-aminopentol, ethenolamine-BHT, ethenolamine-BHhexol, N-formylated-aminotriol, and N-formylated-aminopentol (Fig. 8, Table A4). Low concentrations of nucleoside BHPs, i.e. adenosylhopane and 2Me-adenosylhopane_{HG-diMe}, were also present. Finally,

Formatted: Font: Italic




Commented [NR7]: Changed: formylated to N-formylated

455 **Figure 8: BHP profile of *Methylobacter-Methylotenera* co-culture (n=3). Note: the BHPs present in low abundance are shown in the inset figure.**

we detected two novel BHPs (m/z 594; $C_{35}H_{64}O_6N^+$; Fig. 9) where the MS^2 spectra show the consecutive loss of five hydroxyl moieties and an additional loss of 29 Da (CH_3N). Based on the MS^2 spectra and the elemental composition, we tentatively identify these compounds as aminohexol BHPs (Appendix B8). Notably, 3 β -methylated-BHPs are absent in the co-culture. Based on past culture studies, aminopentol and aminotretrol are associated with Type I and Type II MOB, respectively (Neunlist and Rohmer, 1985b, c; Cvejic et al., 2000; Talbot et al., 2001; van Winden et al., 2012). The high relative abundance of aminopentol relative to aminotretrol observed in the BHP distribution of the *Methylobacter-Methylotenera* co-culture analyzed in this study is similar to that of other Type I MOB (Neunlist and Rohmer, 1985c; Talbot et al., 2001; van Winden et al., 2012; Rush et al., 2016; Kusch and Rush, 2022). The novel BHPs detected in the co-culture, however, are reported for the first time in this study. We conclude that the suite of BHPs identified in the co-culture, including the novel ethenolamine-BHPs, N-formylated-aminoBHPs, and aminohexols, are attributed to the MOB *Methylobacter* present, and thus can be potentially considered as markers of aerobic methanotrophs in lake settings.

3.4 BHPs as biomarkers for aerobic methanotrophs in lakes

470 Aminotretrol ~~was~~ present in all of the Azorean samples, except for in the surface water of Negra (1 m depth), at 5 m depth in Azul, and in the water column and sediment of Cubres East (Fig. 2). Similarly, aminopentol is observed in the water column and sediment of Azul and Verde (except for at a depth of 5 m in the water column of Azul). In Funda and Negra, aminopentol only occurs in the deepest part of the water column and in the sediment samples. For Empadadas and Cubres East, aminopentol occurs both in the water column and the sediment, but it is only
 475 observed in the sediment sample from Cubres West. Similarly, aminopentol~~it~~ is found in both sediment samples from Lomba and São Jorge. Both aminopentol and aminotretrol ~~are~~ also produced in small



Commented [NR8]: Changed figure: remove *m/z* from losses indicated on the structure

480 **Figure 9: (A)** Partial mass chromatograms of aminohexol I (peak a) and II (peak b) BHPs from the *Methylobacter-Methylotenera* co-culture with. MS² spectra for (B) aminohexol I and (C) aminohexol II and the ~~proposed-tentative~~ structures shown. The asterisk (*) indicates arbitrary placement of the hydroxyl group on the side chain.

amounts by sulfur-reducing bacteria (SRB; Blumenberg et al., 2006). As SRB also synthesize relatively higher amounts of aminotriol, ratios of aminopentol and aminotretol to aminotriol have been used to determine the origin of the penta- and hexa-functionalised amino-BHPs (Blumenberg et al., 2006, 2009, 2012). The absence of anoxic conditions in the lake water column would rule out a major contribution of SRB-produced BHPs during the time of sampling. This is confirmed by the high aminopentol:aminotriol (water column: 0 (aminopentol absent) to 0.37 and sediment: 0.01 to 1.67) and aminotretol:aminotriol (water column: 0 (aminotretol absent) to 1.30 and sediment: 0 (aminotretol absent) to 0.38) ratios in both water column and sediment samples. Recently two species of nitrite-oxidizing bacteria were shown to produce aminopentol in minor amounts under certain growth conditions (Elling et al., 2022), potentially contributing to the aminopentol observed in the surface waters of Lake Azul, Empadadas, and Cubres East. However, in the oxygen-depleted bottom waters of Azul, Funda, and Negra, MOB are likely the primary producers of aminotretol and aminopentol. A relatively higher abundance of aminopentol than aminotretol in the water column and sediment samples would suggest a higher abundance of Type I than Type II MOB, in accordance with previous studies in both dimictic and meromictic lakes (Hanson and Hanson, 1996; Oswald et al., 2015, 2016; Guggenheim et al., 2020).

We observe unsaturated aminopentol (van Winden et al., 2012; Wagner et al., 2014) in the bottom water samples of Lake Funda and Negra (Fig. 6), which could be an adaptation by MOB to colder water temperatures (Bale et al., 2019). We also observe unsaturated aminotriol in the bottom water of Azul (15 m, 14 °C), Funda (29 m, 12 °C), and Negra (100 m; temperature data unavailable). C_{16:0}-N-acyl-aminopentol was reported in trace amounts near a terrestrial methane seep (Hopmans et al., 2021). C_{16:0}-N-acyl-aminopentol, as well as C_{14:0} and C_{15:0}-N-acyl-aminopentol are present in sediment samples from Azul and Verde, whereas in Lomba and Empadadas we only observe C_{16:0}-N-acyl-aminopentol in the sediment (Fig. 2). In the NMDS plot of the surface sediments, the acylated-aminopentols group closely with BHPs related to MOB, aminotetrol and aminopentol (Fig. 4). In addition, C_{14:0} and C_{16:0}-N-acyl-aminopentol are both detected in the *Methylobacter-Methylotenera* co-culture, confirming that MOB could be a potential source for these acyl-amino-BHPs in Lakes Azul, Funda, Lomba, and Empadadas.

MC-aminotriol and MC-aminopentol were produced by the *Methylobacter-Methylotenera* co-culture (Fig. 8), albeit in low abundance. However, this is the first report of ethenolamine-BHT, ethenolamine-BHhexol, N-formylated-aminotriol, and N-formylated-aminopentol in culture. So far, ethenolamine-BHPs and N-formylated-aminoBHPs were only identified near a terrestrial methane seep (Hopmans et al., 2021). Further environmental and culture studies are needed to identify whether these are robust biomarkers for methanotrophs. In all of our samples, except for the water column of Negra at 1 and 7 m depths (Fig. 2), we detected ethenolamine-BHPs, propenolamine, and/or N-formylated-aminoBHPs (Hopmans et al., 2021). Their widespread distribution of these compounds suggests they are derived from multiple producers. Although we cannot identify a direct source for these compounds, the NMDS analysis shows that ethenolamine-BHpentol, ethenolamine-BHhexol, acylated-ethenolamine BHhexols (C_{15:0}, C_{16:0}, C_{17:0}), N-formylated-aminotetrol, and N-formylated-aminopentol clustering with MC-aminopentol, aminopentol, aminotetrol and the acylated-aminopentols near the surface sediments from Azul, Verde, Funda, and Negra (Fig. 4). These aminoBHPs are primarily associated with In addition, ethenolamine-BHhexol and N-formylated-aminopentol are detected in the bottom waters of Lake surface sediments from Azul, Verde, Funda, and Negra and the bottom water from Funda and Negra (Fig. 2 & 6). Azul, Verde, Funda, and Negra all stratify during the summer months and the bottom water in these lakes range from suboxic to anoxic conditions (Gonçalves et al., 2018). This would suggest that ethenolamine-BHpentol, ethenolamine-BHhexol, acylated-ethenolamine BHhexols, N-formylated-aminotetrol, and N-formylated-aminopentol ethenolamines, propenolamine, and/or formylated-amino BHPs are being produced under suboxic to anoxic conditions and/or could be associated with MOB.

BHPs with a 3β-methylation, particularly 3MeBHT and 3Me-aminotriol are often linked to methane oxidation (Neunlist and Rohmer, 1985b; Zundel and Rohmer, 1985; Cvejic et al., 2000). However, 3β-methyl-BHPs are not exclusively produced by MOB (Zundel and Rohmer, 1985; Welander and Summons, 2012). BHPs with a 3β-methylation are also produced by other bacteria in the phyla Alpha-, Beta-, and Gamma-proteobacteria, Cyanobacteria, Acidobacteriota, and Nitrospirae (Zundel and Rohmer, 1985; Sinninghe Damsté et al., 2017; Elling et al., 2022). In addition, 3β-methylation is crucial for bacterial cell survival in the late stationary phase and thus could have a broad taxonomic distribution (Welander and Summons, 2012). We observe 3MeBHT in all

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

sediment samples (except for Cubres East and West; (Fig. 2), however, it was not detected in our water column samples. In contrast, 3Me-aminotriol only occurs in sediment samples from Lomba, Cubres West, São Jorge, Empadadas, Verde, and Azul. The absence of 3Me-BHPs in SPM samples suggests that at the time of sampling
540 the bacteria responsible for 3 β -methylation were not abundant in the water column. Future work will need to evaluate annual changes in both BHP distributions and bacterial communities in the water column to confirm the primary producers of 3Me-BHPs in lakes.

AminoBHPs with a methylcarbamate terminal group (MC-aminoBHPs) were proposed as biomarkers for Type I
545 MOB in marine settings based on culture samples (Rush et al., 2016), but so far, the occurrence of MC-aminoBHPs in lakes has not been evaluated. MC-aminotriol has been observed in nitrite-oxidizing bacteria (Elling et al., 2022). The highest rates of methane-oxidation in seasonally stratified lakes typically occurs at or directly below the chemocline (Rudd et al., 1976; Hanson, 1980; Harrits and Hanson, 1980; Oswald et al., 2015). In the Azorean lakes, MC-aminotriol was found in the oxycline and bottom water of both Lake Azul and Funda (Fig. 6).
550 In Azul and Funda we observe MC-aminotriol near the oxycline, which could also be attributed to nitrite-oxidizing bacteria (Elling et al., 2022). We also observe MC-aminotriol in sediment samples from Verde, Negra, São Jorge, Lomba, Cubres East, and Empadadas. The co-occurrence of MC-aminopentol with 3MeBHT and/or 3Me-aminotriol in sediment samples from Verde, Negra, Lomba, and Empadadas could indicate a common MOB source. Both Lomba and Empadadas are relatively shallow lakes with a fully oxygenated water column, which
555 means the primary site of methane oxidation occurs in the surface sediment and not in the water column (Hanson and Hanson, 1996). The different MC-aminoBHP distributions in the sediment samples could indicate that we were either not sampling during the peak period of methane oxidation in the lake, there was a shift in the MOB community, or methane oxidation was primarily occurring at the sediment surface.

560 Based on the distributions of aminoBHPs and MC-aminoBHPs, MOB are likely present, albeit in low-abundance, at all Azorean sites. In the deeper lakes (i.e., Azul, Funda, and Negra), methane oxidation is likely occurring either at or below the chemocline which would account for the diverse aminoBHPs observed in the water column (Fig. 6). Low $\delta^{13}\text{C}$ values in bulk sediment samples from Lake Funda and primarily reducing conditions associated with the bottom water of the lake, would suggest that methanogenesis is occurring in the bottom water and/or surface
565 sediment of Lake Funda during the summer months (Richter et al., 2022). Similarly, proxy records suggest that eutrophication in Lake Azul contributed to hypoxic conditions in the bottom water, ~~resulting in a shift in the chironomid community~~ (Raposeiro et al., 2017). In both lakes Funda and Azul, an increase in organic matter deposition combined with low oxygen conditions in the bottom water during the summer months would support methane production, and therefore, a community of methane-oxidizing bacteria in the water column of these lakes.
570 A similar distribution of BHPs associated with methane-oxidizing bacteria (i.e. aminopentol, unsaturated aminopentol, and aminotetrol) are found in high abundance in the methane-rich environments of floodplains in the Amazon basin and wetlands in the Congo River basin (Wagner et al., 2014; Spencer-Jones et al., 2015). In contrast, in the shallow lakes and the coastal lagoons methane oxidation might occur both in the water column and in the oxygenated sediment. Further molecular and culture studies, however, are needed to validate these
575 conclusions.

3.4 Autochthonous vs. allochthonous source of nucleoside BHPs

A recent study suggests that some nucleoside BHPs may be *in situ* produced in the chemocline or under anoxic/euxinic conditions in marine water columns (Kusch et al., 2021b). To the best of our knowledge, no studies have investigated nucleoside BHP production in lacustrine environments. Nucleoside BHPs were detected in all water column and sediment samples except the water column of lagoon Cubres West (salinity 23.8 ppt; Fig. 7). The distribution of nucleoside BHPs in the sediment of Lake Empadadas is reflected in the BHPs distribution of its surface water, suggesting that in shallow, well-mixed lakes nucleoside BHPs like adenosylhopane and N1-methylinosylhopane are either derived from production in the water column or surface water run-off (Cooke et al., 2008; Xu et al., 2009; Rethemeyer et al., 2010). In contrast, in Lake Funda we observe an increase in adenosylhopane, adenosylhopane_{HG-diMe}, 2Me-adenosylhopane_{HG-diMe}, Me-adenosylhopane_{HG-diMe}, N1-methylinosylhopane, 2Me-N1-methylinosylhopane, and Me-N1-methylinosylhopane at the chemocline (6 m depth). Of these nucleoside BHPs, only adenosylhopane, adenosylhopane_{HG-diMe}, and 2Me-adenosylhopane_{HG-diMe} are present in the surface water (1 m) of Lake Funda. A pycnocline associated with the chemocline in the Azorean lakes could result in the accumulation of adenosylhopane, adenosylhopane_{HG-diMe}, and 2Me-adenosylhopane_{HG-diMe}, however, the lack of Me-adenosylhopane_{HG-diMe}, N1-methylinosylhopane, 2Me-N1-methylinosylhopane, and Me-N1-methylinosylhopane in the surface water of Lake Funda, could also indicate *in situ* production of these compounds in the water column. Kusch et al. (2021b) found that adenosylhopane, N1-methylinosylhopane, and 2Me-N1-methylinosylhopane were produced within the chemocline and under anoxic/euxinic conditions in marine settings. In addition, adenosylhopane is separate from the rest of the nucleosides in the NMDS plot of only water column samples (Fig. 5), falling in between samples collected near the chemocline and the surface water. N1-methylinosylhopane is also observed in the bottom water of Lake Azul, in addition to adenosylhopane_{HG-Me} and adenosylhopane_{HG-diMe}, suggesting that these nucleosides are likely produced *in situ*. The primary producers of nucleoside BHPs, however, remain unconstrained and are likely associated with a wide range of bacteria. Adenosylhopane, in particular, is produced as an intermediate product in BHP synthesis (Bradley et al., 2010) and has the potential to be sourced from diverse bacteria, including purple non-sulfur bacteria (Neunlist and Rohmer, 1985a; Neunlist et al., 1988; Talbot et al., 2007), nitrifying bacteria (Seemann et al., 1999; Talbot et al., 2007; Elling et al., 2022), and nitrogen-fixing bacteria (Bravo et al., 2001; Talbot et al., 2007).

3.5 Implications for the R_{soil} proxy

The ratio of nucleoside BHPs to BHT (R_{soil} index, Eq. 1; Zhu et al., 2011) and the revised soil index for high latitudes (R'_{soil}, Eq. 2; Doğrul Selver et al., 2012) have previously been used as an indication of soil input to riverine and marine environments (Cooke et al., 2009; Sáenz et al., 2011; Taylor and Harvey, 2011; Zhu et al., 2011; Doğrul Selver et al., 2012, 2015; De Jonge et al., 2016; Kusch et al., 2019). However, the recent indications that nucleoside BHPs may be *in situ* produced in marine (Kusch et al., 2021b) and lacustrine water columns (this study) could affect the application of the R_{soil} proxy. R_{soil} values of marine and lacustrine sediments from the tropics to Arctic range from 0.4-0.9 (Cooke et al., 2008; Pearson et al., 2009; Rethemeyer et al., 2010; Kusch et al., 2021a), and values between 0.5-0.8 have been employed as a tentative terrestrial end-member (Zhu et al., 2011). Here we revise the R_{soil} index to exclude nucleoside BHPs that might be produced in the lake water column (i.e., adenosylhopane, early eluting adenosylhopane_{HG-Me}, Me-adenosylhopane_{HG-diMe}, N1-methylinosylhopane, 2Me-N1-methylinosylhopane, Me-N1-methylinosylhopane). We call this new index, R_{soil-lake}. This ratio is

615 significantly correlated with R_{soil} ($R = 0.94$, $p < 0.0001$) and R'_{soil} ($R = 0.60$, $p < 0.0001$). Further, this ratio is
positively correlated with TOC/TN ($R = 0.81$, $p < 0.005$; Fig. C1), where TOC/TN values between 4 to 10 reflect
organic matter derived from lake algae and values higher than 20 indicate an increased source of vascular land
plants (Meyers, 2003). ~~All of the~~ The low sediment TOC/TN values (values fall between 7 to 11) and isotopically
depleted $\delta^{13}C$ values (-31 to -24 ‰; Fig. C2a) and $\delta^{15}N$, indicating indicate predominantly *in situ* derived organic
620 matter which is in accordance with the high levels of primary productivity observed in all of the lakes (Cordeiro
et al., 2020; Meyers, 2003). Similarly, low $\delta^{15}N$ values (-1 to 2 ‰; Fig. C2b) are commonly associated with high-
levels of nitrogen-fixation in accordance with the cyanobacterial blooms observed in the lakes during the summer
months (Cordeiro et al., 2020; Meyers, 2003)(Meyers, 2003). The slightly more enriched $\delta^{13}C$ values (-19 to -15
625 ‰) in Cubres East and West are typical of marine algae, also suggesting that the organic matter is mainly derived
from *in situ* production (Meyers, 1994).

Formatted: Font: Italic

At all of the sites, we observe lower $R_{soil-lake}$ values in the water column than the sediment. As our water column
SPM samples were collected in the dry season in the Azores (April-August; Hernández et al., 2016), we would
630 anticipate low terrestrial inputs. In contrast, the BHPs detected in the sediments represent an integrated annual
signal, and likely includes a larger input of terrestrial material during the wet season (September-March;
Hernández et al., 2016). The $R_{soil-lake}$ values found in the sediments of the deeper lakes (i.e., Funda, Negra, and
Azul) range from 0.17 to 0.32, and similarly have TOC/TN values between 8.1 and 8.3. The sediments of the
shallow lakes all contain relatively higher $R_{soil-lake}$ values (i.e., Empadadas = 0.57, São Jorge = 0.69, and Lomba
635 = 0.56) and higher TOC/TN values (ranging from 10 to 11.4), which could indicate slightly higher inputs of
terrestrial organic matter, particularly during the wet season. In the coastal lagoons, Cubres East and West, we
observe low $R_{soil-lake}$ values both in the water column (0.04 Cubres East and 0 Cubres West) and sediment samples
(0.21 Cubres East and 0.16 Cubres West). This could reflect fewer terrestrial inputs relative to the other lakes in
this study. Low TOC/TN values (7.08 Cubres East and 7.22 Cubres West) would also indicate a primarily *in situ*
640 source of organic matter (Meyers, 2003), however, further work is needed to verify this.

Our results highlight that caution should be used to ensure that the nucleoside BHPs included in the R_{soil} proxy
are not produced *in situ*. In our current dataset, the $R_{soil-lake}$ index, which excludes nucleoside BHPs that likely
originate from the water column, appears to work for all of our Azorean sites, however additional work is needed
645 to distinguish the primary source of nucleoside BHPs, and to expand the application of $R_{soil-lake}$. More detailed
water column and catchment studies are needed to confirm whether these nucleoside BHPs are actually being
produced within the lakes. Furthermore, similar comparison studies should be performed in marine settings.

4 Conclusions

This study highlights the diversity and complexity of BHP distributions and their interpretations in the lacustrine
and coastal environments. We identified several novel BHPs that are being described for the first time in lacustrine
650 and coastal settings, including unsaturated ethenolamine-BHPs, and N-formylated-aminoBHPs, oxazinone-
aminotriol, and dioxanone-methylaminotriol. Further, we identified several nucleoside BHPs that appear to be
produced in the water column could be produced in the water column of lacustrine settings, however, further work

is needed to verify the source of these BHPs. We propose a revised R_{soil} index, $R_{\text{soil-lakes}}$ to distinguish between
655 ~~terrestrial- and aquatic-derived organic matter in lakes, by excluding any nucleosides that might be produced *in*~~
~~*situ.*~~ Within lakes, aminotetrol, aminopentol, and MC-aminopentol show potential as proxies for MOB. In
addition, the recently identified ethenolamine-BH~~pentol~~BHT, ethenolamine-BHhexol, N-formylated-aminotriol,
and N-formylated-aminopentol are produced by a lacustrine methanotroph, *Methylobacter* sp., and within the
Azorean lakes are associated with low-oxygen conditions. Additional studies focused on describing BHP
660 distributions particularly in lacustrine settings and in culture studies, however, are needed to confirm whether
certain BHPs can be used as biomarkers for biogeochemical cycles or environmental conditions.

Formatted: Subscript

Formatted: Subscript

Formatted: Font: Italic

Appendix A Supplementary Tables

Table A1: BHPs identified in this study with the retention times (tr), assigned elemental composition (AEC), calculated exact mass (M_{calc}), and the Δ ppm ($\Delta \text{ ppm} = ((M_{\text{calc}} - M_{\text{measured}})/M_{\text{calc}}) * 10^6$).

BHP	Sample	tr (min)	AEC	MS ² ion	M _{calc} (m/z)	Δ ppm	Fig. Reference Figure	References
adenosylhopane	São Jorge SS2	21.71	C ₄₀ H ₆₈ O ₃ N ₅	H ⁺	662.500	-0.11	B2, peak a	Hopmans et al. 2010
adenosylhopane _{CHG-Me}	São Jorge SS2	17.42	C ₄₁ H ₆₆ O ₃ N ₅	H ⁺	676.516	0.14	B2, peak b*	this study
2Me-adenosylhopane	São Jorge SS2	21.82	C ₄₁ H ₆₆ O ₃ N ₅	H ⁺	676.516	-0.08	B2, peak c	Hopmans et al. 2010
Me-adenosylhopane, unknown isomer	São Jorge SS2	22.30	C ₄₁ H ₆₆ O ₃ N ₅	H ⁺	676.516	0.04	B2, peak d	Hopmans et al. 2010
adenosylhopane _{CHG-Me}	São Jorge SS2	22.90	C ₄₁ H ₆₆ O ₃ N ₅	H ⁺	676.516	-0.18	B2, peak e	Hopmans et al. 2010
3Me-adenosylhopane	Lomba SS2	23.76	C ₄₁ H ₆₆ O ₃ N ₅	H ⁺	676.516	-1.16		Hopmans et al. 2010
adenosylhopane _{CHG-dMe}	São Jorge SS2	17.60	C ₄₂ H ₆₈ O ₃ N ₅	H ⁺	690.532	-0.81	B2, peak f*	this study
Me-adenosylhopane _{CHG-Me}	São Jorge SS2	17.90	C ₄₂ H ₆₈ O ₃ N ₅	H ⁺	690.532	-0.77	B2, peak g*	this study
2Me-adenosylhopane _{CHG-Me}	São Jorge SS2	22.96	C ₄₂ H ₆₈ O ₃ N ₅	H ⁺	690.532	-0.04	B2, peak h	Hopmans et al. 2010
Me-adenosylhopane _{CHG-Me}	Empadadas SS7	23.62	C ₄₂ H ₆₈ O ₃ N ₅	H ⁺	690.532	-0.81	B2, peak i	Hopmans et al. 2010
adenosylhopane _{CHG-dMe}	São Jorge SS2	25.15	C ₄₂ H ₆₈ O ₃ N ₅	H ⁺	690.532	-0.25	B2, peak j	Hopmans et al. 2010
Me-adenosylhopane _{CHG-dMe}	São Jorge SS2	18.06	C ₄₃ H ₇₀ O ₃ N ₅	H ⁺	704.547	0.82	B2, peak k*	this study
diMe-adenosylhopane _{CHG-Me}	Empadadas SS10	23.68	C ₄₃ H ₇₀ O ₃ N ₅	H ⁺	704.547	-1.38	B2, peak l*	this study
2Me-adenosylhopane _{CHG-dMe}	São Jorge SS2	25.17	C ₄₃ H ₇₀ O ₃ N ₅	H ⁺	704.547	-0.50	B2, peak m	Hopmans et al. 2010
Me-adenosylhopane _{CHG-dMe}	São Jorge SS2	26.09	C ₄₃ H ₇₀ O ₃ N ₅	H ⁺	704.547	-0.40	B2, peak n	Hopmans et al. 2010
2,3-diMe-adenosylhopane _{CHG-dMe}	São Jorge SS2	26.14	C ₄₄ H ₇₂ O ₃ N ₅	H ⁺	718.563	-0.10	B2, peak o	Hopmans et al. 2010
inosylhopane	Empadadas SS8	20.33	C ₄₀ H ₆₅ O ₄ N ₄	H ⁺	663.484	-0.68	B3, peak a	Hopmans et al. 2010
2Me-inosylhopane	Empadadas SS8	20.45	C ₄₁ H ₆₅ O ₄ N ₄	H ⁺	677.500	-0.28	B3, peak b*	this study
Me-inosylhopane, unknown isomer	Empadadas SS8	20.90	C ₄₁ H ₆₅ O ₄ N ₄	H ⁺	677.500	-1.10	B3, peak c*	this study
N1-methylinosylhopane	Empadadas SS8	23.91	C ₄₁ H ₆₅ O ₄ N ₄	H ⁺	677.500	-0.96	B3, peak d	Hopmans et al. 2010
2Me-N1-methylinosylhopane	Empadadas SS8	24.03	C ₄₂ H ₆₇ O ₄ N ₄	H ⁺	691.516	-0.36	B3, peak e	Hopmans et al. 2010
Me-N1-methylinosylhopane, unknown isomer	Empadadas SS8	24.54	C ₄₂ H ₆₇ O ₄ N ₄	H ⁺	691.516	-0.16	B3, peak f*	this study
Me-N1-methylinosylhopane, unknown isomer	Empadadas SS8	24.99	C ₄₂ H ₆₇ O ₄ N ₄	H ⁺	691.516	1.39	B3, peak g*	this study
BHT	São Jorge SS2	20.48	C ₃₃ H ₆₆ O ₄ N	NH ₄ ⁺	564.499	-0.17		
Δ ⁶ -BHT (34S)	São Jorge SS2	18.28	C ₃₃ H ₆₆ O ₄ N	NH ₄ ⁺	562.483	0.34		
Δ ⁶ -BHT (34R)	Empadadas 001	18.43	C ₃₃ H ₆₆ O ₄ N	NH ₄ ⁺	562.483	0.49		
2MeBHT	São Jorge SS2	20.62	C ₃₆ H ₆₈ O ₄ N	NH ₄ ⁺	578.514	-0.14	B4, peak a	
3MeBHT	São Jorge SS2	22.26	C ₃₆ H ₆₈ O ₄ N	NH ₄ ⁺	578.514	0.29		
Δ ⁶ -3MeBHT	Negra SS1	20.15	C ₃₆ H ₆₈ O ₄ N	NH ₄ ⁺	576.499	-0.97		
unsaturated-3MeBHT	São Jorge SS1	22.07	C ₃₆ H ₆₆ O ₄ N	NH ₄ ⁺	576.499	-0.97		
Methoxylated-BHT I	Cubres SS3	21.71	C ₃₆ H ₆₈ O ₄ N	NH ₄ ⁺	578.514	-0.04	B4, peak b	this study
Methoxylated-BHT II	Cubres SS6	22.3	C ₃₆ H ₆₈ O ₄ N	NH ₄ ⁺	578.514	-0.09	B4, peak c	this study
BHT-CE	Verde SS1	17.15	C ₄₁ H ₇₁ O ₃ N	NH ₄ ⁺	708.541	-0.15	B5, peak a	
BHT-Me-CE	Verde SS1	17.25	C ₄₂ H ₇₃ O ₃ N	NH ₄ ⁺	722.557	-1.64	B5, peak b	this study
3MeBHT-CE	Verde SS1	18.63	C ₄₂ H ₇₃ O ₃ N	NH ₄ ⁺	722.557	-1.92	B5, peak c	
Anhydro-BHT	Verde SS1	24.09	C ₃₃ H ₆₆ O ₃ N	NH ₄ ⁺	546.488	-0.90		
BHpentol	Verde SS1	18.59	C ₃₃ H ₆₆ O ₃ N	NH ₄ ⁺	580.494	-1.77		
BHpentol-CE	Verde SS1	15.91	C ₄₁ H ₇₁ O ₂ N	NH ₄ ⁺	724.536	-0.08		

Formatted	... [1]
Formatted	... [2]
Formatted	... [4]
Formatted	... [5]
Formatted Table	... [3]
Formatted	... [6]
Formatted	... [7]
Formatted	... [8]
Formatted	... [9]
Formatted	... [10]
Formatted	... [11]
Formatted	... [12]
Formatted	... [13]
Formatted	... [14]
Formatted	... [15]
Formatted	... [16]
Formatted	... [17]
Formatted	... [18]
Formatted	... [19]
Formatted	... [20]
Formatted	... [21]
Formatted	... [22]
Formatted	... [23]
Formatted	... [24]
Formatted	... [25]
Formatted	... [26]
Formatted	... [27]
Formatted	... [28]
Formatted	... [29]
Formatted	... [30]
Formatted	... [31]
Formatted	... [32]
Formatted	... [33]
Formatted	... [34]
Formatted	... [35]
Formatted	... [36]
Formatted	... [37]
Formatted	... [38]
Formatted	... [39]
Formatted	... [40]
Formatted	... [41]
Formatted	... [42]
Formatted	... [43]
Formatted	... [44]
Formatted	... [45]
Formatted	... [46]
Formatted	... [47]
Formatted	... [48]
Formatted	... [49]
Formatted	... [50]
Formatted	... [51]
Formatted	... [52]
Formatted	... [53]
Formatted	... [54]
Formatted	... [55]
Formatted	... [56]
Formatted	... [57]
Formatted	... [58]
Formatted	... [59]
Formatted	... [60]
Formatted	... [61]
Formatted	... [62]
Formatted	... [63]
Formatted	... [64]
Formatted	... [65]

										Formatted	...	[167]
										Formatted	...	[168]
										Formatted	...	[169]
										Formatted	...	[170]
										Formatted	...	[171]
										Formatted	...	[172]
										Formatted	...	[173]
										Formatted	...	[174]
										Formatted	...	[175]
										Formatted	...	[176]
										Formatted	...	[177]
										Formatted	...	[178]
										Formatted	...	[179]
										Formatted	...	[180]
										Formatted	...	[181]
										Formatted	...	[182]
										Formatted	...	[183]
										Formatted	...	[184]
										Formatted	...	[185]
										Formatted	...	[186]
										Formatted	...	[187]
										Formatted	...	[188]
										Formatted	...	[189]
										Formatted	...	[190]
										Formatted	...	[191]
										Formatted	...	[192]
										Formatted	...	[193]
										Formatted	...	[194]
										Formatted	...	[195]
										Formatted	...	[196]
										Formatted	...	[197]
										Formatted	...	[198]
										Formatted	...	[199]
										Formatted	...	[200]
										Formatted	...	[201]
										Formatted	...	[202]
										Formatted	...	[203]
										Formatted	...	[204]
										Formatted	...	[205]
										Formatted	...	[206]
										Formatted	...	[207]
										Formatted	...	[208]
										Formatted	...	[209]
										Formatted	...	[210]
										Formatted	...	[211]
										Formatted	...	[212]
										Formatted	...	[213]
										Formatted	...	[214]
										Formatted	...	[215]
										Formatted	...	[216]
										Formatted	...	[217]
										Formatted	...	[218]
										Formatted	...	[219]
										Formatted	...	[220]
										Formatted	...	[221]
										Formatted	...	[222]
										Formatted	...	[223]
										Formatted	...	[224]
										Formatted	...	[225]
										Formatted	...	[226]
										Formatted	...	[227]
										Formatted	...	[228]
										Formatted	...	[229]
										Formatted	...	[230]
										Formatted	...	[231]

Δ^6 -N-formylated aminotriol	Verde SS1	20.28	C ₃₆ H ₆₄ O ₄ N	H ⁺	574.483	-0.18	3, peak j	Hopmans et al. 20
Δ^6 -N-formylated aminotriol	Verde SS1	18.03	C ₃₆ H ₆₂ O ₄ N	H ⁺	572.467	-0.86	3, peak k and B9	this study
N-formylated aminotetrol	Verde SS1	18.43	C ₃₆ H ₆₆ O ₅ N	H ⁺	590.478	-1.39	3, peak m	Hopmans et al. 20
Δ^{11} -N-formylated aminotetrol	Verde SS1	16.77	C ₃₆ H ₆₂ O ₅ N	H ⁺	588.462	-1.00	3, peak n	this study
N-formylated aminopentol	Verde SS1	16.44	C ₃₆ H ₆₆ O ₆ N	H ⁺	606.473	-0.70	3, peak o	this study
Δ^{11} -N-formylated aminopentol	Verde SS1	14.95	C ₃₆ H ₆₂ O ₆ N	H ⁺	604.457	-1.45	3, peak p	this study
Δ^6 -oxazinone-aminotriol	Verde SS1	20.72	C ₃₆ H ₆₂ O ₄ N	H ⁺	572.467	-1.07	3, peak l	Elling et al. 2022
aminohexol I	<i>Methylobacter- Methylotenera</i>	13.08	C ₃₅ H ₆₀ O ₆ N	H ⁺	594.473	-0.36	5, peak a	this study
aminohexol II	<i>Methylobacter- Methylotenera</i>	13.94	C ₃₅ H ₆₀ O ₆ N	H ⁺	594.473	-0.08	5, peak b	this study
dioxanone-methylaminotriol	Verde SS1	21.88	C ₃₇ H ₆₆ O ₄ N	H ⁺	586.483	0.72	3, peak c	this study

Formatted	... [351]
Formatted	... [352]
Formatted	... [353]
Formatted	... [354]
Formatted	... [355]
Formatted	... [356]
Formatted	... [357]
Formatted	... [358]
Formatted	... [359]
Formatted	... [360]
Formatted	... [361]
Formatted	... [362]
Formatted	... [363]
Formatted	... [364]
Formatted	... [365]
Formatted	... [366]
Formatted	... [367]
Formatted	... [368]
Formatted	... [369]
Formatted	... [370]
Formatted	... [371]
Formatted	... [372]
Formatted	... [373]
Formatted	... [374]
Formatted	... [375]
Formatted	... [376]
Formatted	... [377]
Formatted	... [378]
Formatted	... [379]
Formatted	... [380]
Formatted	... [381]
Formatted	... [382]
Formatted	... [383]
Formatted	... [384]
Formatted	... [385]
Formatted	... [386]
Formatted	... [387]
Formatted	... [388]
Formatted	... [389]
Formatted	... [390]
Formatted	... [391]
Formatted	... [392]
Formatted	... [393]
Formatted	... [394]
Formatted	... [395]
Formatted	... [396]
Formatted	... [397]
Formatted	... [398]
Formatted	... [399]
Formatted	... [400]
Formatted	... [401]
Formatted	... [402]

Table A2: Water column samples and water column measurements from the time of sampling (June 2018) discussed in this study (where D.O. = dissolved oxygen). The relative abundance of the major BHPs identified in each sample are also shown (where BHT = bacteriohopanetetrol, BHT-CE = BHT-cyclitol ether).

Water Column Samples		June 2018 Water Column						Relative Abundance of Main BHPs		
Island	Lake/Lagoon	Water Depth (m)	Temp. (°C)	pH	Turbidity (NTU)	D.O. (%)	Salinity (ppt)	BHT (%)	BHT-CE (%)	Aminotriol II (%)
São Miguel	Azul	1	20.6	9.9	11.9	97.9	0.1	15.8	60.2	8.7
São Miguel	Azul	5	19.9	8.4	14.1	84.1	0.1	23.0	54.8	9.0
São Miguel	Azul	15	14.2	7.8	8.7	63.3	0.1	22.2	27.4	23.0
São Miguel	Empadadas Norte	1	16.5	8.9	20.9	93.0	0.0	23.7	4.8	17.8
São Jorge	Cubres East	0	22.6	9.9	6.8	95.0	9.7	54.1	0.0	20.9
São Jorge	Cubres West	0.5	24.1	9.9	4.9	-	23.8	47.5	0.0	34.6
Flores	Funda	1	19.7	11.3	29.4	96.4	0.1	29.0	53.9	1.7
Flores	Funda	6	15.2	8.2	5.4	57.3	0.1	23.2	57.6	5.1
Flores	Funda	29	12.3	8.3	4.0	24.1	0.1	6.4	6.8	43.5
Flores	Negra	1	18.7	11.3	57.4	100.6	0.1	8.0	90.8	0.4
Flores	Negra	7	14.7	9.7	7.4	77.6	0.1	13.4	84.5	0.5
Flores	Negra*	100	-	-	-	-	-	22.7	38.5	20.3

*Note: Measurements in Lake Negra are only made up to ~50 m water column depth.

Formatted: Font: 7 pt

Formatted Table

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

675 **Table A3: Location and characteristics of the sediment sample sites discussed in this study (where Z = maximum water depth, Rep. n = replicates, TN = total nitrogen, and TOC = total organic carbon). The relative abundance of the major BHPs identified in each sample are also shown (where Adeno. = adenosylhopane, N1-methylino. = N1-methylinosylhopane, BHT = bacteriohopanetetrol, and BHT-CE = BHT-cyclitol ether).**

Site	Sediment samples				Relative Abundance of Main BHPs										
	Island	Lake/ Lagoon	Depth Z (m)	Rep. n	TN (%)	TOC (%)	$\delta^{15}N$ (‰)	$\delta^{13}C$ (‰)	Adeno. (%)	N1-methylino. (%)	BHT (%)	BHT-CE (%)	Aminotriol II (%)	Aminopentol (%)	Ethenolamine-BH
São Miguel	Azul	5	2	0.59	5.14	2.0	-23.7	3.1	1.2	9.0	10.1	17.2	28.7		
São Miguel	Azul	23	0	0.60	4.43	2.0	-26.0	6.3	2.9	13.0	9.2	13.8	16.1		
São Miguel	Verde	5	0	1.22	8.81	1.1	-26.0	1.4	1.0	12.5	11.2	16.3	11.7		
São Miguel	Verde	22.5	0	0.86	8.18	-1.3	-27.6	1.7	2.0	8.2	19.1	13.8	22.2		
São Miguel	Empadadas Norte	2	4	0.82	8.26	-0.2	-27.4	10.4	13.2	9.5	7.4	11.4	8.7		
São Jorge	São Jorge	2.5	2	0.97	11.00	1.1	-26.5	13.8	15.4	8.9	8.0	7.6	1.3		
São Jorge	Cubres East	2	0	1.95	13.79	1.3	-18.9	1.9	1.1	28.0	0.0	11.3	0.2		
São Jorge	Cubres West	2	0	1.92	13.89	2.1	-15.3	3.6	1.4	23.6	4.0	19.0	2.3		
Flores	Funda	29.7	0	0.89	7.18	1.5	-30.7	2.2	1.1	13.8	19.4	15.5	11.2		
Flores	Lomba	14.3	2	1.94	19.45	1.0	-27.7	11.6	7.6	10.0	9.6	12.4	8.1		
Flores	Negra	114	2	0.83	6.77	2.4	-30.5	5.6	0.8	11.6	19.7	11.2	5.5		

680

Table A4: *Methylobacter-Methylotenera* co-culture (n=3) BHP distributions with BHPs reported as relative abundance and the standard deviation between culture samples. The retention times (tr) and calculated exact masses (M_{calc}) are also shown.

Bacteriohopanepolyol/BHP	tr (min)	M_{calc} (m/z)	Relative Abundance (%)	Standard deviation
adenosylhopane	21.77	662.50	0.174	0.006
2Me-adenosylhopane _{HC-dMe}	25.25	704.547	0.002	0.001
aminotriol II	17.52	546.49	43.308	1.027
Δ^6 -aminotriol	15.47	544.47	0.067	0.032
Δ^{11} -aminotriol	15.88	544.47	3.454	0.133
$C_{14:0}$ -N-acyl-aminotriol	33.91	756.69	0.006	0.002
$C_{16:0}$ -N-acyl-aminotriol	37.41	784.72	0.007	0.004
$C_{16:1}$ -N-acyl-aminotriol	34.44	782.70	0.079	0.043
aminotetrol	16.30	562.48	6.381	0.188
aminopentol	14.69	578.48	43.441	1.242
unsaturated aminopentol	13.51	576.46	2.552	0.117
$C_{14:0}$ -N-acyl-aminopentol	27.49	788.68	0.008	0.001
$C_{15:0}$ -N-acyl-aminopentol	30.82	816.71	0.005	0.001
methylcarbamate-aminotriol II	20.89	604.49	0.126	0.030
methylcarbamate-aminotetrol	18.96	620.49	0.003	0.003
methylcarbamate-aminopentol	16.87	636.48	0.024	0.004
ethenolamine-BHT	20.67	588.50	0.039	0.013
ethenolamine-BHhexol	16.70	620.49	0.009	0.009
N-formylated-aminotriol	20.28	574.48	0.033	0.012
N-formylated-aminopentol	16.44	606.47	0.019	0.012
aminohexol I	13.08	594.47	0.131	0.018
aminohexol II	13.94	594.47	0.117	0.016

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt


Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Appendix B. Identification of BHPs



BHP	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈
BHT	H	H	H	H	OH	OH	OH	OH
2Me-BHT	CH ₃	H	H	H	OH	OH	OH	OH
3Me-BHT	H	CH ₃	H	H	OH	OH	OH	OH
BHpentol	H	H	H	OH	OH	OH	OH	OH
BHhexol	H	H	OH	OH	OH	OH	OH	OH
aminotriol	H	H	H	H	OH	OH	OH	NH ₂
2Me-aminotriol	CH ₃	H	H	H	OH	OH	OH	NH ₂
Me-aminotriol*	H	H	H	H	OH	OH	OH	NH ₂
3Me-aminotriol	H	CH ₃	H	H	OH	OH	OH	NH ₂
aminotetrol	H	H	H	OH	OH	OH	OH	NH ₂
aminopentol	H	H	OH	OH	OH	OH	OH	NH ₂

*Note: The extra methyl group does not occur at the C-2 or C-3 position. The position is unknown.

Figure B1: Core structure of BHPs discussed in the text with the carbon positions and the rings labeled. More complex structures are shown in the figures.

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

Formatted: Font: 7 pt

B1 Novel nucleoside BHPs

We identified nucleoside BHPs based on Hopmans et al. (2021) in surface sediments and water column samples from the Azores Archipelago. Fig. B2 shows the partial mass chromatograms of adenosyl-containing nucleoside BHPs with increasing degrees of methylation, found in surface sediment samples from São Jorge and Empadadas. In addition to the adenosyl-BHPs previously identified by Hopmans et al. (2021; peaks a, c-e, h-j, and m-o), we observe several additional peaks (b*, f*, g*, k*, and l*). In the mass chromatogram m/z 676.516, an early eluting peak (b*) occurs in both samples. The MS² spectrum for peak b* contains one fragment ion related to the head group at m/z 150.077 ($C_6H_8N_5^+$, Δ ppm -0.52), which suggests that the adenosyl head group is methylated. Therefore, we tentatively assign peak b* as early eluting adenosylhopane_{HG-Me} (assigned elemental composition (AEC) = $C_{41}H_{66}O_3N_5$, m/z 676.516, Δ ppm 0.14; Fig. B2d). The mass chromatogram of m/z 690.532, contains two early eluting peaks in São Jorge (peaks f* and g*) and one early eluting peak in Empadadas (peak g*). Peaks f* and g* are putatively identified as Me-adenosylhopane_{HG-Me} and adenosylhopane_{HG-diMe} based on the dominant fragment ions related to the head group at m/z 150.077 ($C_6H_8N_5^+$, Δ ppm -0.52; Fig. B2e) and m/z 164.093 ($C_7H_{10}N_5^+$, Δ ppm -1.09; Fig. B2f), respectively. In the sediment samples from Lake São Jorge, the mass chromatogram for m/z 704.547 contains one early eluting peak (peak k*). The MS² spectrum of peak k* is characterized by a fragment at m/z 164.093 ($C_7H_{10}N_5^+$, Δ ppm -0.78; Fig. B2g), which is associated with a dimethylated head group. Thus, we tentatively identify peak k* as Me-adenosylhopane_{HG-diMe}. In Empadadas, we observe another early eluting peak (peak l*) in the mass chromatogram for m/z 704.547 with an MS² spectrum that contains two main mass fragments associated with the adenosyl head group: m/z 150.077 ($C_6H_8N_5^+$, Δ ppm -0.85) and m/z 164.093 ($C_7H_{10}N_5^+$, Δ ppm -1.27; Fig. B2h). Peak l* is likely another isomer of diMe-adenosylhopane_{HG-Me}, but is co-eluting with Me-adenosylhopane_{HG-Me} (peak i) as suggested by the fragment m/z 150.077 ($C_6H_8N_5^+$).

Inosylhopanes, described in Hopmans et al. (2021), were identified in surface sediments from Lake Empadadas, (peaks a, d, and e; Fig. B3a) as well as several novel isomers of methylated-inosylhopanes (peaks b*, c*, f*, and g*; Fig. B3a). The mass chromatogram of m/z 691.516 contains two early eluting peaks (peak b* and c*). Both mass spectra contain one primary fragment associated with the head group, m/z 137.046 ($C_5H_5ON_4^+$, peak b*, Δ ppm -1.11 and peak c*, Δ ppm -0.82; Fig. B3b), indicating an inosine head group. Thus, we identify these peaks as Me-inosylhopane with the methylation occurring on the core structure. The retention time difference between inosylhopane and peaks b* (0.12 min) and c* (0.57 min), is similar to the retention time difference between BHT and 2MeBHT (0.14 min) and MeBHT with a methylation at an unknown position in the ring system (0.58 min), respectively (Hopmans et al., 2021). Therefore, we tentatively assign peak b* as 2Me-inosylhopane and peak c* as Me-inosylhopane, unknown isomer. For peak f* we observe a fragment of m/z 151.061 ($C_6H_7ON_4^+$, Δ ppm -1.14; Fig. B3c), suggesting a methylation on the inosine head group. We propose that peak f* is an isomer of methyl-N1-methylinosylhopane (elemental composition (EC) = $C_{42}H_{67}O_4N_4$, m/z 691.516). Based on the fragment m/z 151.061 ($C_6H_7ON_4^+$, Δ ppm -1.08; Fig. B3d), peak g* is another isomer of Me-N1-methylinosylhopane with unknown positions of the methyl groups on the ring system and head group. However, we also observe an m/z 164.093 ($C_7H_{10}N_5^+$, Δ ppm -1.15) fragment in the MS² spectra of peak g*. We attribute this to ~~potential~~ a potential co-eluting peak that we are unable distinguish.




Figure B2: Partial mass chromatograms of adenosyl-containing nucleoside BHPs in a sediment sample from (A) Lake São Jorge and (B) Lake Empadadas showing the exact mass and intensity of the highest peak in arbitrary units (AU). Peaks identified as novel nucleoside BHP isomers are marked with a ⁺*. (C) general structure of a nucleoside BHP (D-H) MS² of novel early-eluting nucleoside BHP isomers identified in Azorean lakes and a proposed structure for each nucleobase. Note, the mass spectra of peak l* (panel H) appears to be a mixed spectrum of co-eluting nucleoside BHPs.





Figure B3: (A) Partial mass chromatogram of inosylhopane (peak a), N1-methylinosylhopane (peak d), and 2-methyl-N1-methylinosylhopane (peak e) in a sediment sample from Lake Empadadas. (B) A MS^2 of newly identified peak of putative Me-inosylhopane (peak b^*) with the proposed structure for the nucleobase shown. MS^2 of newly identified 2-methyl-N1-methylinosylhopane isomers for (C) peak f^* and (D) peak g^* with the structures of the proposed nucleobases. Note, the mass spectra for peak g^* appears to be a mixed spectrum.


B2 Methoxylated-bacteriohopanetetrol

In the mass chromatogram of m/z 578.514 ($C_{36}H_{68}O_4N^+$) we identified two peaks that elute after the expected retention times of 2MeBHT (peak a) and 3MeBHT (Fig. B4a). The MS² spectrum of the peak b at 21.71 mins (Fig. B4b) shows a distinct loss of three hydroxyl moieties: m/z 543.477 ($C_{36}H_{63}O_3^+$, Δ ppm -0.68), m/z 525.467 ($C_{36}H_{61}O_2^+$, Δ ppm -0.48), and m/z 507.459 ($C_{36}H_{59}O^+$, Δ ppm -5.37) and a loss of 32 Da (CH_3OH) from m/z 525.466 ($C_{36}H_{61}O_2^+$, Δ ppm -0.48) to m/z 493.441 ($C_{35}H_{57}O^+$, Δ ppm -0.26). The lower mass range of the mass spectrum is comparable to that of BHT (Hopmans et al., 2021). We therefore tentatively identify this BHP as a methoxylated-BHT. Peak c at 22.30 mins shows a similar loss of three hydroxyl moieties and a loss of 32 Da. The later elution time of this peak could result from a different position of the methoxy-group on the side-chain. The position of the methoxy group in both isomers is unknown. We tentatively identify these peaks as methoxylated-BHTs.

A Cubres SS3



B MS² of m/z 578.514, [M+H]⁺ of methoxylated-BHT (peak b)



Commented [NR9]: Changed figure: remove m/z and change to loss

Figure B4: (A) Partial mass chromatogram of 2MeBHT (peak a), methoxylated-BHT I (peak b), and methoxylated-BHT II in a sediment sample from Cubres East. (B) MS² of the methoxylated-BHT I (peak b) and the proposed structure for methoxylated BHT. The asterisk (*) indicates that the position of the methoxy moiety is unknown.

B3 Bacteriohopanetetrol-methyl-cyclitol ether

In the mass chromatogram of m/z 722.555, the calculated exact mass of MeBHT-cyclitol ether (CE), from Verde SS1 we observe two peaks (Fig. B5a): a peak at 17.25 mins (peak b) and 3MeBHT-CE ($C_{42}H_{76}O_8N^+$; peak c) at 18.63 mins. The MS^2 spectrum of peak b (Fig. B5b) shows the consecutive losses of two hydroxyl moieties producing fragment ions at m/z 704.543 ($C_{42}H_{74}O_7N^+$, Δ ppm 3.85) and m/z 686.529 ($C_{42}H_{72}O_6N^+$, Δ ppm 2.95), however, the expected loss of a third hydroxyl moiety (m/z 668.520, $C_{42}H_{70}O_5N^+$) is not observed, likely due to the low intensity. In the lower mass range, the fragments follow the fragmentation pattern for a cyclitol ether head group described in Hopmans et al. (2021), but with an additional mass of 14 Da. The intact head group corresponds to the fragment m/z 194.102 ($C_7H_{16}O_5N^+$, Δ ppm -0.31) followed by the loss of several hydroxyl moieties: m/z 176.091 ($C_7H_{14}O_4N^+$, Δ ppm 1.05), m/z 158.081 ($C_7H_{12}O_3N^+$, Δ ppm -2.47), m/z 140.070 ($C_7H_{10}O_2N^+$, Δ ppm 0.32). We also observe two fragments that correspond to the intact polar head group with two additional carbon atoms, originating from the side chain after fragmentation at the C-33 and C-34 bond: m/z 236.113 ($C_9H_{18}O_6N^+$, Δ ppm 0.06) and m/z 218.102 ($C_9H_{16}O_5N^+$, Δ ppm 0.09). Based on the MS^2 spectrum, we tentatively identify this peak as a BHT cyclitol ether with a methylation on the head group: BHT-Me-CE (AEC = $C_{42}H_{76}O_8N^+$, m/z 722.557, Δ ppm -1.64). Note, that we do not observe a loss of 32 Da indicative of a methoxy moiety (CH_3O^+), therefore we propose that the methyl-group occurs on the ring structure of the cyclitol ether.




Figure B5: (A) Partial mass chromatogram of BHT cyclitol ether (BHT-CE, peak a) and newly proposed BHT-Me-CE (peak b) in a sediment sample from Lake Verde. (B) MS^2 spectrum of $[M+H]^+$ BHT-Me-CE (peak b) and a proposed structure for BHT-Me-CE with the diagnostic fragmentation of the cyclitol ether indicated (Note: the asterisk (*) indicates that the position of the methyl group is unknown, the additional methylation could also occur at the C-39 or C-40 position).

30 B4 Ethenolamine-BHPs

In our samples we identified ethenolamine-BHT (peak a), ethenolamine-BHpentol (peak d), and ethenolamine-BHhexol (peak h) after Hopmans et al. (2021; Fig. 3a). In addition, we identified unsaturated ethenolamine-BHT (peak b) and unsaturated ethenolamine-BHhexol (peak i) in the mass chromatograms of m/z 586.483 ($C_{37}H_{64}O_4N^+$) and m/z 618.474 ($C_{37}H_{64}O_6N^+$), respectively. The fragmentation spectra of both unsaturated ethenolamine-BHPs (Fig. 3e and B6, respectively) showed the characteristic loss of 41 Da (C_2H_3N) and 59 Da (C_2H_5ON) related to the loss of the ethenolamine moiety. In both fragmentation spectra we observe the expected fragment for an unsaturation on the BHP core structure, i.e. m/z 473.413 ($C_{35}H_{53}^+$, Δ ppm 1.54) for an unsaturated BHT and m/z 469.382 ($C_{35}H_{49}^+$, Δ ppm 2.08) for an unsaturated BHhexol. The offset in retention time between ethenolamine-BHT and unsaturated-ethenolamine-BHT (2.33 mins) and ethenolamine-BHhexol and unsaturated-ethenolamine-BHhexol (1.51 mins), respectively, is similar to the offset between BHT and Δ^6 -BHT (2.34 mins) and BHT and Δ^{11} -BHT (1.38 mins), respectively, observed by Hopmans et al. (2021). Based on the retention time offset, we propose that the unsaturation in peaks b and i are at the Δ^6 - and Δ^{11} -position, respectively. In the lower mass range of the MS² spectrum of the proposed Δ^6 -ethenolamine-BHT we observe some deviations from the expected fragments for a Δ^6 -BHT, i.e. a relatively dominant m/z 177 and m/z 201 instead of m/z 203, which is expected for a Δ^6 -BHT core (Hopmans et al., 2021). Perhaps the presence of a conjugation alters the fragmentation pattern of an unsaturated core to a small degree. The MS² spectrum of the proposed Δ^{11} -ethenolamine-BHhexol (peak i) also shows the dominant fragment ion at m/z 177, but does show the expected fragment ion at m/z 203 for an unsaturation at the Δ^{11} -position.

MS² of m/z 586.483, $[M+H]^+$ of Δ^6 -ethenolamine-BHT (peak b)




Figure B6: MS² spectra of $[M+H]^+$ Δ^6 -ethenolamine-BHT from sediment samples in Lake Verde.

50 B5 N-fFormylated-aminoBHPs

We detected two unknown composite BHPs previously described in Hopmans et al. (2021) in the partial mass chromatograms of m/z 574.483 ($C_{36}H_{64}O_4N^+$) and m/z 590.483 ($C_{36}H_{64}O_5N^+$). In addition to several consecutive losses of water representing 3 or 4 hydroxyl moieties on the side chain, the fragmentation spectra of both of these unknown composite BHPs is characterized by an initial loss of 28 Da (CO). Here, we detected these novel BHPs in sediment from Lake Verde (Fig. 3b, peaks j and m). In addition, we identified an additional BHP with m/z 606.473 and an AEC of $C_{36}H_{64}O_6N^+$ (Fig. 3b and f, peak o) with a similar fragmentation: a loss of 28 Da (CO)

from the original mass fragment produces the fragment at m/z 578.484 ($C_{35}H_{64}O_5N^+$, Δ ppm -5.81) and a loss of a hydroxyl moiety (m/z 588.461, $C_{36}H_{62}O_5N^+$, Δ ppm 1.29) followed by a loss of 28 Da (CO) results in the fragment at m/z 560.468 ($C_{35}H_{62}O_4N^+$, Δ ppm -1.76). We also observe five sequential losses of hydroxyl moieties from the parent ion to produce m/z 588.461 ($C_{36}H_{62}O_5N^+$, Δ ppm 2.96), m/z 570.451 ($C_{36}H_{60}O_4N^+$, Δ ppm -0.17), m/z 552.441 ($C_{36}H_{58}O_3N^+$, Δ ppm 2.03), m/z 534.431 ($C_{36}H_{56}O_2N^+$, Δ ppm -1.92), and m/z 516.420 ($C_{36}H_{54}ON^+$, Δ ppm 1.09), respectively. In the fragmentation spectra of peaks j, m, and o we observe losses of 45 Da (CONH₃), e.g. from m/z 534 to m/z 489 for peak o (Fig. 3f). Based on these losses we propose that these novel composite BHPs are a series of N-formylated-aminoBHPs: N-formylated-aminotriol (peak j), N-formylated-aminotetrol (peak m) and N-formylated-aminopentol (peak o). The proposed structure for N-formylated-aminopentol is shown in Fig. 3f with the diagnostic fragmentations indicated.

We also observe unsaturated versions of these N-formylated-aminotriol, N-formylated-aminotetrol, and N-formylated-aminopentol based on the elemental composition $C_{36}H_{62}O_4N^+$ (m/z 572.467, peak k), $C_{36}H_{62}O_5N^+$ (m/z 588.462, peak n), $C_{36}H_{62}O_6N^+$ (m/z 604.458, peak p), respectively. In the lower mass range of the m/z 572.467 spectrum, we observe dominant m/z 163.148 ($C_{12}H_{19}^+$, Δ ppm -0.14), m/z 177.163 ($C_{13}H_{21}^+$, Δ ppm 0.10), m/z 201.164 ($C_{15}H_{21}^+$, Δ ppm 0.53), and m/z 203.179 ($C_{15}H_{23}^+$, Δ ppm -0.46) fragments and a notable absence of a dominant m/z 191.179 ($C_{14}H_{23}^+$). This is similar to the fragmentation patterns observed for unsaturated ethenolamine-BHT (Fig. B7) and unsaturated ethenolamine-BHhexol (Fig. 3e) as discussed in section B4. We also observe the corresponding fragments of the D and E ring with a partially de-hydroxylated side-chain in the spectra: m/z 350.271 ($C_{21}H_{36}O_3N^+$, Δ ppm -3.31) and after two additional hydroxyl losses, m/z 314.249 ($C_{21}H_{32}ON^+$, Δ ppm -5.95), and the loss of the hexane ring on the side-chain to get m/z 231.210 ($C_{17}H_{27}^+$, Δ ppm 0.03). This indicates that the unsaturation is not on the side-chain or the D and E rings. Based on the presence of the m/z 201.164 fragment and a similar offset in retention times between N-formylated-aminotriol and unsaturated N-formylated-aminotriol (2.25 mins; Table A1) to that of ethenolamine-BHT and unsaturated ethenolamine-BHT (2.33 mins; Table A1), we tentatively assign the double bond to the Δ^6 position. In m/z 588.462 ($C_{36}H_{62}O_5N^+$; Fig. 3b), we identify peak n as unsaturated N-formylated-aminotetrol based on the MS¹ spectrum. However, due to a low MS² spectrum we cannot identify the position of the double bond. Based on the retention time, the double bond is likely at the Δ^6 position. The difference in retention time between N-formylated-aminopentol and unsaturated N-formylated-aminopentol (1.49 mins) is similar to that of the ethenolamine-BHhexol and

Commented [NR10]: Changed figure B7: modified name to N-formylated-aminotriol




Figure B7: MS² spectra of [M+H]⁺ Δ^6 -N-formylated aminotriol from sediment samples in Lake Verde.

unsaturated ethenolamine-BHhexol (1.51 mins), therefore the double bond likely occurs in the same place for both compounds. We therefore tentatively identify peak p as Δ^{11} -N-formylated-aminopentol.

B6 Oxazinone-aminotriol

In the mass chromatogram for m/z 572.467, in addition to the unsaturated N-formylated-aminotriol (peak k), we observe a later-eluting peak ($C_{36}H_{62}O_4N^+$; Fig. 3b and d, peak l). The MS² spectrum of this peak contains a loss of 44 Da (CO₂) from the original mass fragment to produce m/z 528.477 ($C_{35}H_{62}O_2N^+$, Δ ppm -1.33) followed by a loss of two hydroxyl moieties (m/z 510.466, $C_{35}H_{60}ON^+$, Δ ppm 1.36 and m/z 492.457, $C_{35}H_{58}N^+$, Δ ppm -2.92). We propose that this is the same compound identified in Elling et al. (2022). The authors describe an acetylated compound (m/z 656.495; $C_{40}H_{66}O_6N^+$), with the unique loss of 44 Da (CO₂), the loss of two hydroxyl moieties, and an additional loss of 17 Da (NH₃⁺). We indicate the loss of 17 Da (NH₃⁺) in the proposed structure; however, we do not observe this loss in our MS² spectrum, which could be attributed to low peak intensity. Based on the proposed structure, this appears to be the cyclized form of the N-formylated-aminotriol. The IUPAC name for the cyclized part of the BHP is 1,3-oxazinone-2-one (National Center for Biotechnology Information, 2023a), therefore we refer to this compound as oxazinone-aminotriol.

100 B7 Dioxanone-methylaminotriol


In the mass chromatogram of m/z 586.483 ($C_{37}H_{64}O_4N^+$; Δ ppm 0.72), an additional peak occurs at 21.88 mins (Fig. 3a, peak c). Similar to the above described N-formylated-aminoBHPs, the MS² spectra is characterized by the distinct loss of 28 Da producing m/z 558.488 ($C_{36}H_{64}O_3N^+$; Δ ppm -0.37; Fig. 3c). In addition, a loss of 31 Da (CNH₅) is observed from m/z 522.467 ($C_{36}H_{60}ON^+$; Δ ppm 0.04) to produce the fragment at m/z 491.424 ($C_{35}H_{55}O^+$; Δ ppm 1.98), indicating a second conjugated moiety. Based on the elemental composition there appears to be an additional DBE (double bond equivalent) in the structure, but there is no indication the unsaturation is located in the core structure as the lower mass range of the fragmentation spectrum is similar to what is expected for BHT. We therefore assume one of the functionalities is part of a cyclized structure and propose this BHP is a type of N-formylated-aminotriol, where the formic acid is part of a cyclic structure and the terminal amino group is methylated. The IUPAC name for the cyclic part of the structure is 1,3-dioxan-2-one (National Center for Biotechnology Information, 2023b), therefore we refer to this compound as a dioxanone-methylaminotriol. The proposed structure with key fragmentations is shown in Fig. 3c.

B8 Aminohexol BHPs

A search for additional BHPs using the m/z 191.179 trace ($C_{14}H_{22}^+$) fragment spectra revealed two early eluting peaks in the mass chromatogram for m/z 594.473 ($C_{35}H_{64}O_6N^+$) in the *Methylobacter-Methylotenera* coculture (Fig. 9a). The MS² spectrum for aminohexol I (Fig. 9b) shows the consecutive loss of five hydroxyl moieties to produce fragments at m/z 594.473 ($C_{35}H_{64}O_6N^+$, Δ ppm -0.36), m/z 576.462 ($C_{35}H_{62}O_5N^+$, Δ ppm 0.59), m/z 558.454 ($C_{35}H_{60}O_4N^+$, Δ ppm -3.55), m/z 540.441 ($C_{35}H_{58}O_3N^+$, Δ ppm 0.83), m/z 522.432 ($C_{35}H_{56}O_2N^+$, Δ ppm -2.30), and m/z 504.418 ($C_{35}H_{54}ON^+$, Δ ppm 3.04). Based on the presence of m/z 369.354 ($C_{27}H_{45}^+$, Δ ppm -6.91) in the MS² spectrum, the additional hydroxyl moiety does not occur on the ring system. The position of the additional hydroxyl moiety on the tail is unknown. Similarly, the MS² spectrum for aminohexol II (Fig. 9c) also shows a loss of five hydroxyl moieties, and an additional loss of 29 Da (CH₃N) from m/z 504.422

(C₃₅H₅₄ON⁺, Δ ppm -3.01) to *m/z* 475.393 (C₃₄H₅₁O⁺, Δ ppm 0.99). However, in the fragmentation spectrum of peak b, we observe several fragments that appear to represent the functionalized tail at *m/z* 180.086 (C₆H₁₄O₅N⁺, Δ ppm -0.12), *m/z* 162.076 (C₆H₁₂O₄N⁺, Δ ppm 0.95), *m/z* 144.0655 (C₆H₁₀O₃N⁺, Δ ppm -0.35), and *m/z* 126.055 (C₆H₈O₂N⁺, Δ ppm -1.55). The complementary ion to the tail fragments is observed at *m/z* 395.368 (C₂₉H₄₇⁺, Δ ppm -0.06), indicating fragmentation at the C22-C30 bond. The presence of the *m/z* 369.354 (C₂₇H₄₅⁺, Δ ppm -0.20) fragment indicates a hopanoid core without the addition of an extra hydroxyl group. We propose that in this isomer, the additional hydroxyl group is located at C-22 on the side-chain, resulting in the initial fragmentation of the side-chain and the progressive loss of the hydroxyl groups from the side-chain fragment. Based on the MS² spectra, we tentatively identify these compounds as aminohexol BHPs.

Appendix C Additional figures



135

Figure C1: Correlation between (A) R_{soil} , (B) R'_{soil} , and (C) $R_{soil-lake}$ with TOC/TN. The different colors represent the different types of samples sites in this study: coastal (Cubres East and West), deep (Azul, Funda, and Negra), and shallow (Empadadas, São Jorge, and Lomba).

140




Figure C2: Distribution of (A) $\delta^{13}C$ (‰, VPDB) and TOC/TN values and (B) $\delta^{15}N$ (‰, Air) and $\delta^{13}C$ (‰, VPDB) values for the surface sediment samples analysed in this study.

Commented [NR11]: Added plots for d15N and d13C values from surface samples

Formatted: Superscript

Data availability

Mass spectra published in this manuscript will be added to the Global Natural Products Social (GNPS) Molecular Networking database (<https://gnps.ucsd.edu/ProteoSAFe/static/gnps-splash.jsp>). All additional data produced in this study ~~this study is available in the will be published in the~~ supplementary material.

150 Author contribution

The study was conceptualized by NR, LAA-Z, and DR. The investigation was carried out by all co-authors. Formal analysis was conducted by NR, ECH, and DM. ECH, PMR, VG, ACC, and LV contributed resources to make this project possible. Funding was acquired by LAA-Z, PMR, LV, and DR. NR wrote the manuscript, and all co-authors contributed to the manuscript.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by the Luso-American Foundation (“Crossing the Atlantic”) (LAA-Z), the National
 155 Funds through FCT - Foundation for Science and Technology under the project UIDB/50027/2020, PMR by (DL57/2016/ ICETA/EEC2018/25) grant, and the NWO gravitation grant for Soehngen Institute of Anaerobic Microbiology (024.002.002). We would like to thank everyone who participated and assisted with ~~the~~ field campaign to the Azores Archipelago in 2018, particularly E. Zettler. Further, we would like to thank Dr. S. van Grinsven for the co-culture and advice. We thank Dr. N. Bale, A. Mets, M. Verweij, D. Dorhout, J. Ossebaar, R.
 160 van Bommel, Dr. M. van der Meer, I. Posthuma, A. Nordeloos, and M. Grego for technical support and advice.

References

- [Bale, N. J., Rijpstra, W. I. C., Sahonero-Canavesi, D. X., Oshkin, I. Y., Belova, S. E., Dedysh, S. N., and
 165 \[Sinninghe Damsté, J. S.: Fatty Acid and Hopanoid Adaption to Cold in the Methanotroph *Methylovulum psychrotolerans*, *Frontiers in Microbiology*, 10, 2019.\]\(#\)](#)
- [Bale, N. J., Ding, S., Hopmans, E. C., Arts, M. G. I., Villanueva, L., Boschman, C., Haas, A. F., Schouten, S.,
 and Sinninghe Damsté, J. S.: Lipidomics of Environmental Microbial Communities. I: Visualization of
 Component Distributions Using Untargeted Analysis of High-Resolution Mass Spectrometry Data, *Frontiers
 in Microbiology*, 12, 2021.](#)

Formatted: Heading 1, Indent: Left: 0 cm, First line: 0 cm, Don't suppress line numbers

Formatted: Font: 10 pt

Formatted: Don't add space between paragraphs of the same style, Line spacing: 1.5 lines

- 170 [Bastviken, D., Cole, J. J., Pace, M. L., and Van de Bogert, M. C.: Fates of methane from different lake habitats: Connecting whole-lake budgets and CH₄ emissions, *Journal of Geophysical Research: Biogeosciences*, 113, <https://doi.org/10.1029/2007JG000608>, 2008.](#)
- [Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, *Canadian Journal of Biochemistry and Physiology*, 37, 911–917, <https://doi.org/10.1139/o59-099>, 1959.](#)
- 175 [Blumenberg, M., Krüger, M., Nauhaus, K., Talbot, H. M., Oppermann, B. I., Seifert, R., Pape, T., and Michaelis, W.: Biosynthesis of hopanoids by sulfate-reducing bacteria \(genus *Desulfovibrio*\), *Environmental Microbiology*, 8, 1220–1227, <https://doi.org/10.1111/j.1462-2920.2006.01014.x>, 2006.](#)
- [Blumenberg, M., Seifert, R., Kasten, S., Bahlmann, E., and Michaelis, W.: Euphotic zone bacterioplankton sources major sedimentary bacteriohopanepolyols in the Holocene Black Sea, *Geochimica et Cosmochimica Acta*, 73, 750–766, <https://doi.org/10.1016/j.gca.2008.11.005>, 2009.](#)
- 180 [Blumenberg, M., Hoppert, M., Krüger, M., Dreier, A., and Thiel, V.: Novel findings on hopanoid occurrences among sulfate reducing bacteria: Is there a direct link to nitrogen fixation?, *Organic Geochemistry*, 49, 1–5, <https://doi.org/10.1016/j.orggeochem.2012.05.003>, 2012.](#)
- [Blumenberg, M., Berndmeyer, C., Moros, M., Muschalla, M., Schmale, O., and Thiel, V.: Bacteriohopanepolyols record stratification, nitrogen fixation and other biogeochemical perturbations in Holocene sediments of the central Baltic Sea, *Biogeosciences*, 10, 2725–2735, <https://doi.org/10.5194/bg-10-2725-2013>, 2013.](#)
- [Bradley, A. S., Pearson, A., Sáenz, J. P., and Marx, C. J.: Adenosylhopane: The first intermediate in hopanoid side chain biosynthesis, *Organic Geochemistry*, 41, 1075–1081, <https://doi.org/10.1016/j.orggeochem.2010.07.003>, 2010.](#)
- 190 [Bravo, J.-M., Perzl, M., Härtner, T., Kannenberg, E. L., and Rohmer, M.: Novel methylated triterpenoids of the gammacerane series from the nitrogen-fixing bacterium *Bradyrhizobium japonicum* USDA 110, *European Journal of Biochemistry*, 268, 1323–1331, <https://doi.org/10.1046/j.1432-1327.2001.01998.x>, 2001.](#)
- [Cooke, M. P., Talbot, H. M., and Farrimond, P.: Bacterial populations recorded in bacteriohopanepolyol distributions in soils from Northern England, *Organic Geochemistry*, 39, 1347–1358, <https://doi.org/10.1016/j.orggeochem.2008.05.003>, 2008.](#)
- 195 [Cooke, M. P., van Dongen, B. E., Talbot, H. M., Semiletov, I., Shakhova, N., Guo, L., and Gustafsson, Ö.: Bacteriohopanepolyol biomarker composition of organic matter exported to the Arctic Ocean by seven of the major Arctic rivers, *Organic Geochemistry*, 40, 1151–1159, <https://doi.org/10.1016/j.orggeochem.2009.07.014>, 2009.](#)
- 200 [Coolen, M. J. L., Talbot, H. M., Abbas, B. A., Ward, C., Schouten, S., Volkman, J. K., and Sinninghe Damsté, J. S.-S.: Sources for sedimentary bacteriohopanepolyols as revealed by 16S rDNA stratigraphy, *Environmental Microbiology*, 10, 1783–1803, <https://doi.org/10.1111/j.1462-2920.2008.01601.x>, 2008.](#)
- [Cordeiro, R., Luz, R., Vilaverde, J., Vasconcelos, V., Fonseca, A., and Gonçalves, V.: Distribution of Toxic Cyanobacteria in Volcanic Lakes of the Azores Islands, *Water*, 12, 3385, <https://doi.org/10.3390/w12123385>, 2020.](#)
- 205 [Cvejić, J. H., Bodrossy, L., Kovács, K. L., and Rohmer, M.: Bacterial triterpenoids of the hopane series from the methanotrophic bacteria *Methylocaldum* spp.: phylogenetic implications and first evidence for an](#)

- unsaturated aminobacteriohopanepolyol, *FEMS Microbiology Letters*, 182, 361–365,
210 <https://doi.org/10.1111/j.1574-6968.2000.tb08922.x>, 2000.
- De Jonge, C., Talbot, H. M., Bischoff, J., Stadnitskaia, A., Cherkashov, G., and Sinninghe Damsté, J. S.: Bacteriohopanepolyol distribution in Yenisei River and Kara Sea suspended particulate matter and sediments traces terrigenous organic matter input, *Geochimica et Cosmochimica Acta*, 174, 85–101, <https://doi.org/10.1016/j.gca.2015.11.008>, 2016.
- 215 Doğrul Selver, A., Talbot, H. M., Gustafsson, Ö., Boulton, S., and van Dongen, B. E.: Soil organic matter transport along an sub-Arctic river–sea transect, *Organic Geochemistry*, 51, 63–72, <https://doi.org/10.1016/j.orggeochem.2012.08.002>, 2012.
- Doğrul Selver, A., Sparkes, R. B., Bischoff, J., Talbot, H. M., Gustafsson, Ö., Semiletov, I. P., Dudarev, O. V., Boulton, S., and van Dongen, B. E.: Distributions of bacterial and archaeal membrane lipids in surface
220 sediments reflect differences in input and loss of terrestrial organic carbon along a cross-shelf Arctic transect, *Organic Geochemistry*, 83–84, 16–26, <https://doi.org/10.1016/j.orggeochem.2015.01.005>, 2015.
- Doughty, D. M., Coleman, M. L., Hunter, R. C., Sessions, A. L., Summons, R. E., and Newman, D. K.: The RND-family transporter, HpnN, is required for hopanoid localization to the outer membrane of
Rhodospseudomonas palustris TIE-1, *Proceedings of the National Academy of Sciences*, 108, E1045–E1051, <https://doi.org/10.1073/pnas.1104209108>, 2011.
- 225 Elling, F. J., Evans, T. W., Nathan, V., Hemingway, J. D., Kharbush, J. J., Bayer, B., Spieck, E., Husain, F., Summons, R. E., and Pearson, A.: Marine and terrestrial nitrifying bacteria are sources of diverse bacteriohopanepolyols, *Geobiology*, 20, 399–420, <https://doi.org/10.1111/gbi.12484>, 2022.
- Farrimond, P., Head, I. M., and Innes, H. E.: Environmental influence on the biohopanoid composition of recent
230 sediments, *Geochimica et Cosmochimica Acta*, 64, 2985–2992, [https://doi.org/10.1016/S0016-7037\(00\)00404-X](https://doi.org/10.1016/S0016-7037(00)00404-X), 2000.
- Gonçalves, V.: Contribuição do estudo das microalgas para a avaliação da qualidade ecológica das lagoas dos Açores: fitoplâncton e diatomáceas bentônicas, University of the Azores, Ponta Delgada, 343 pp., 2008.
- Gonçalves, V., Raposeiro, P.M., Marques, H.S., Vilaverde, J., Balibrea, A., Rosa, F., Sixto, M., and Costa, A.C.:
235 Monitorização das Massas de Água Interiores e de Transição da Região Hidrográfica dos Açores. Relatório Anual do Ano 3 (R5/Ano 3) Ponta Delgada, 2018.
- van Grinsven, S., Sinninghe Damsté, J. S., Harrison, J., and Villanueva, L.: Impact of Electron Acceptor Availability on Methane-Influenced Microorganisms in an Enrichment Culture Obtained From a Stratified
Lake, *Frontiers in Microbiology*, 11, 2020.
- 240 Guggenheim, C., Freimann, R., Mayr, M. J., Beck, K., Wehrli, B., and Bürgmann, H.: Environmental and Microbial Interactions Shape Methane-Oxidizing Bacterial Communities in a Stratified Lake, *Frontiers in Microbiology*, 11, 2020.
- Hanson, R. S.: Ecology and Diversity of Methylotrophic Organisms, in: *Advances in Applied Microbiology*, vol. 26, edited by: Perlman, D., Academic Press, 3–39, [https://doi.org/10.1016/S0065-2164\(08\)70328-9](https://doi.org/10.1016/S0065-2164(08)70328-9),
245 1980.
- Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiological Reviews*, 60, 439–471, <https://doi.org/10.1128/mr.60.2.439-471.1996>, 1996.

- Harrits, S. M. and Hanson, R. S.: Stratification of aerobic methane-oxidizing organisms in Lake Mendota, Madison, Wisconsin, *Limnology and Oceanography*, 25, 412–421, <https://doi.org/10.4319/lo.1980.25.3.0412>, 1980.
- 250 <https://doi.org/10.4319/lo.1980.25.3.0412>, 1980.
- Hernández, A., Kutiel, H., Trigo, R. M., Valente, M. A., Sigró, J., Cropper, T., and Santo, F. E.: New Azores archipelago daily precipitation dataset and its links with large-scale modes of climate variability, *International Journal of Climatology*, 36, 4439–4454, <https://doi.org/10.1002/joc.4642>, 2016.
- Hopmans, E. C., Smit, N. T., Schwartz-Narbonne, R., Sinnighe Damsté, J. S., and Rush, D.: Analysis of non-derivatized bacteriohopanepolyols using UHPLC-HRMS reveals great structural diversity in environmental lipid assemblages, *Organic Geochemistry*, 160, 104285, <https://doi.org/10.1016/j.orggeochem.2021.104285>, 2021.
- 255 <https://doi.org/10.1016/j.orggeochem.2021.104285>, 2021.
- Kusch, S. and Rush, D.: Revisiting the precursors of the most abundant natural products on Earth: A look back at 30+ years of bacteriohopanepolyol (BHP) research and ahead to new frontiers, *Organic Geochemistry*, 172, 104469, <https://doi.org/10.1016/j.orggeochem.2022.104469>, 2022.
- 260 <https://doi.org/10.1016/j.orggeochem.2022.104469>, 2022.
- Kusch, S., Sepúlveda, J., and Wakeham, S. G.: Origin of Sedimentary BHPs Along a Mississippi River–Gulf of Mexico Export Transect: Insights From Spatial and Density Distributions, *Frontiers in Marine Science*, 6, 2019.
- Kusch, S., Wakeham, S. G., Dildar, N., Zhu, C., and Sepúlveda, J.: Bacterial and archaeal lipids trace chemo(auto)trophy along the redoxcline in Vancouver Island fjords, *Geobiology*, 19, 521–541, <https://doi.org/10.1111/gbi.12446>, 2021a.
- 265 <https://doi.org/10.1111/gbi.12446>, 2021a.
- Kusch, S., Wakeham, S. G., and Sepúlveda, J.: Diverse origins of “soil marker” bacteriohopanepolyols in marine oxygen deficient zones, *Organic Geochemistry*, 151, 104150, <https://doi.org/10.1016/j.orggeochem.2020.104150>, 2021b.
- 270 <https://doi.org/10.1016/j.orggeochem.2020.104150>, 2021b.
- Matys, E. D., Sepúlveda, J., Pantoja, S., Lange, C. B., Caniupán, M., Lamy, F., and Summons, R. E.: Bacteriohopanepolyols along redox gradients in the Humboldt Current System off northern Chile, *Geobiology*, 15, 844–857, <https://doi.org/10.1111/gbi.12250>, 2017.
- Meyers, P. A.: Preservation of elemental and isotopic source identification of sedimentary organic matter, *Chemical Geology*, 114, 289–302, [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0), 1994.
- 275 [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0), 1994.
- Meyers, P. A.: Applications of organic geochemistry to paleolimnological reconstructions: a summary of examples from the Laurentian Great Lakes, *Organic Geochemistry*, 34, 261–289, [https://doi.org/10.1016/S0146-6380\(02\)00168-7](https://doi.org/10.1016/S0146-6380(02)00168-7), 2003.
- 280 [https://doi.org/10.1016/S0146-6380\(02\)00168-7](https://doi.org/10.1016/S0146-6380(02)00168-7), 2003.
- Mitrović, D., Hopmans, E. C., Bale, N. J., Richter, N., Amaral-Zettler, L. A., Baxter, A. J., Peterse, F., Miguel Raposeiro, P., Gonçalves, V., Cristina Costa, A., and Schouten, S.: Isoprenoidal GDGTs and GDDs associated with anoxic lacustrine environments, *Organic Geochemistry*, 178, 104582, <https://doi.org/10.1016/j.orggeochem.2023.104582>, 2023.
- 285 <https://doi.org/10.1016/j.orggeochem.2023.104582>, 2023.
- National Center for Biotechnology Information. PubChem Compound Summary for CID 641496, 1,3-Oxazinan-2-one: https://pubchem.ncbi.nlm.nih.gov/compound/1_3-Oxazinan-2-one, last access: 26 January 2023a.
- National Center for Biotechnology Information. PubChem Compound Summary for CID 123834, 1,3-Dioxan-2-one: https://pubchem.ncbi.nlm.nih.gov/compound/1_3-Dioxan-2-one, last access: 26 January 2023b.

Formatted: Don't add space between paragraphs of the same style

Formatted: Don't add space between paragraphs of the same style, Line spacing: 1.5 lines

- Neunlist, S. and Rohmer, M.: A novel hopanoid, 30-(5'-adenosyl)hopane, from the purple non-sulphur bacterium *Rhodopseudomonas acidophila*, with possible DNA interactions., *Biochem J*, 228, 769–771, 1985a.
- 290 Neunlist, S. and Rohmer, M.: Novel hopanoids from the methylotrophic bacteria *Methylococcus capsulatus* and *Methylomonas methanica*. (22S)-35-aminobacteriohopane-30,31,32,33,34-pentol and (22S)-35-amino-3 β -methylbacteriohopane-30,31,32,33,34-pentol, *Biochemical Journal*, 231, 635–639, <https://doi.org/10.1042/bj2310635>, 1985b.
- 295 Neunlist, S. and Rohmer, M.: The Hopanoids of '*Methylosinus trichosporium*': Aminobacteriohopanetriol and Aminobacteriohopanetetrol, *Microbiology*, 131, 1363–1367, <https://doi.org/10.1099/00221287-131-6-1363>, 1985c.
- Neunlist, S., Bisserset, P., and Rohmer, M.: The hopanoids of the purple non-sulfur bacteria *Rhodopseudomonas palustris* and *Rhodopseudomonas acidophila* and the absolute configuration of bacteriohopanetetrol, *European Journal of Biochemistry*, 171, 245–252, <https://doi.org/10.1111/j.1432-1033.1988.tb13783.x>, 1988.
- 300 O'Beirne, M. D., Sparkes, R., Hamilton, T. L., van Dongen, B. E., Gilhooly, W. P., and Werne, J. P.: Characterization of diverse bacteriohopanepolyols in a permanently stratified, hyper-euxinic lake, *Organic Geochemistry*, 168, 104431, <https://doi.org/10.1016/j.orggeochem.2022.104431>, 2022.
- Osborne, K. A.: *Environmental controls on bacteriohopanepolyol signatures in estuarine sediments*, Newcastle University, UK, 2015.
- 305 Osborne, K. A., Gray, N. D., Sherry, A., Leary, P., Mejeha, O., Bischoff, J., Rush, D., Sidgwick, F. R., Birgel, D., Kalyuzhnaya, M. G., and Talbot, H. M.: Methanotroph-derived bacteriohopanepolyol signatures as a function of temperature related growth, survival, cell death and preservation in the geological record, *Environmental Microbiology Reports*, 9, 492–500, <https://doi.org/10.1111/1758-2229.12570>, 2017.
- 310 Oswald, K., Milucka, J., Brand, A., Littmann, S., Wehrli, B., Kuypers, M. M. M., and Schubert, C. J.: Light-Dependent Aerobic Methane Oxidation Reduces Methane Emissions from Seasonally Stratified Lakes, *PLOS ONE*, 10, e0132574, <https://doi.org/10.1371/journal.pone.0132574>, 2015.
- Oswald, K., Milucka, J., Brand, A., Hach, P., Littmann, S., Wehrli, B., Kuypers, M. M. M., and Schubert, C. J.: Aerobic gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters, *Limnology and Oceanography*, 61, S101–S118, <https://doi.org/10.1002/lno.10312>, 2016.
- 315 Ourisson, G. and Albrecht, P.: Hopanoids. I. Geohopanoids: the most abundant natural products on Earth?, *Acc. Chem. Res.*, 25, 398–402, <https://doi.org/10.1021/ar00021a003>, 1992.
- Pearson, A., Leavitt, W. D., Sáenz, J. P., Summons, R. E., Tam, M. C.-M., and Close, H. G.: Diversity of hopanoids and squalene-hopene cyclases across a tropical land-sea gradient, *Environmental Microbiology*, 11, 1208–1223, <https://doi.org/10.1111/j.1462-2920.2008.01817.x>, 2009.
- 320 Pereira, C. L., Raposeiro, P. M., Costa, A. C., Bao, R., Giralt, S., and Gonçalves, V.: Biogeography and lake morphometry drive diatom and chironomid assemblages' composition in lacustrine surface sediments of oceanic islands, *Hydrobiologia*, 730, 93–112, <https://doi.org/10.1007/s10750-014-1824-6>, 2014.
- Raposeiro, P. M., Rubio, M. J., González, A., Hernández, A., Sánchez-López, G., Vázquez-Loureiro, D., Rull, V., Bao, R., Costa, A. C., Gonçalves, V., Sáez, A., and Giralt, S.: Impact of the historical introduction of

Formatted: Don't add space between paragraphs of the same style

Formatted: Don't add space between paragraphs of the same style, Line spacing: 1.5 lines

- 325 [exotic fishes on the chironomid community of Lake Azul \(Azores Islands\), *Palaeogeography, Palaeoclimatology, Palaeoecology*, 466, 77–88, <https://doi.org/10.1016/j.palaeo.2016.11.015>, 2017.](#)
- [Raposeiro, P. M., Saez, A., Giral, S., Costa, A. C., and Gonçalves, V.: Causes of spatial distribution of subfossil diatom and chironomid assemblages in surface sediments of a remote deep island lake, *Hydrobiologia*, 815, 141–163, <https://doi.org/10.1007/s10750-018-3557-4>, 2018.](#)
- 330 [Rethemeyer, J., Schubotz, F., Talbot, H. M., Cooke, M. P., Hinrichs, K.-U., and Mollenhauer, G.: Distribution of polar membrane lipids in permafrost soils and sediments of a small high Arctic catchment, *Organic Geochemistry*, 41, 1130–1145, <https://doi.org/10.1016/j.orggeochem.2010.06.004>, 2010.](#)
- [Richter, N., Russell, J. M., Amaral-Zettler, L., DeGroff, W., Raposeiro, P. M., Gonçalves, V., de Boer, E. J., Pla-Rabes, S., Hernández, A., Benavente, M., Ritter, C., Sáez, A., Bao, R., Trigo, R. M., Prego, R., and Giral, S.: Long-term hydroclimate variability in the sub-tropical North Atlantic and anthropogenic impacts on lake ecosystems: A case study from Flores Island, the Azores, *Quaternary Science Reviews*, 285, 107525, <https://doi.org/10.1016/j.quascirev.2022.107525>, 2022.](#)
- [Rohmer, M., Bouvier-Nave, P., and Ourisson, G. 1984: Distribution of Hopanoid Triterpenes in Prokaryotes, *Microbiology*, 130, 1137–1150, <https://doi.org/10.1099/00221287-130-5-1137>, 1984.](#)
- 340 [Rudd, J. W. M., Furutani, A., Flett, R. J., and Hamilton, R. D.: Factors controlling methane oxidation in shield lakes: The role of nitrogen fixation and oxygen concentration I, *Limnology and Oceanography*, 21, 357–364, <https://doi.org/10.4319/lo.1976.21.3.0357>, 1976.](#)
- [Rush, D., Osborne, K. A., Birgel, D., Kappler, A., Hirayama, H., Peckmann, J., Poulton, S. W., Nickel, J. C., Mangelsdorf, K., Kalyuzhnaya, M., Sidgwick, F. R., and Talbot, H. M.: The Bacteriohopanepolyol Inventory of Novel Aerobic Methane Oxidising Bacteria Reveals New Biomarker Signatures of Aerobic Methanotrophy in Marine Systems, *PLOS ONE*, 11, e0165635, <https://doi.org/10.1371/journal.pone.0165635>, 2016.](#)
- [Rush, D., Talbot, H. M., van der Meer, M. T. J., Hopmans, E. C., Douglas, B., and Sinninghe Damsté, J. S.: Biomarker evidence for the occurrence of anaerobic ammonium oxidation in the eastern Mediterranean Sea during Quaternary and Pliocene sapropel formation, *Biogeosciences*, 16, 2467–2479, <https://doi.org/10.5194/bg-16-2467-2019>, 2019.](#)
- [Sáenz, J. P.: Hopanoid enrichment in a detergent resistant membrane fraction of *Crocospaera watsonii*: Implications for bacterial lipid raft formation, *Organic Geochemistry*, 41, 853–856, <https://doi.org/10.1016/j.orggeochem.2010.05.005>, 2010.](#)
- 355 [Sáenz, J. P., Eglinton, T. I., and Summons, R. E.: Abundance and structural diversity of bacteriohopanepolyols in suspended particulate matter along a river to ocean transect, *Organic Geochemistry*, 42, 774–780, <https://doi.org/10.1016/j.orggeochem.2011.05.006>, 2011.](#)
- [Sáenz, J. P., Sezgin, E., Schwillie, P., and Simons, K.: Functional convergence of hopanoids and sterols in membrane ordering, *Proceedings of the National Academy of Sciences*, 109, 14236–14240, <https://doi.org/10.1073/pnas.1212141109>, 2012.](#)
- 360 [Santos, F. D., Valente, M. A., Miranda, P. M. A., Aguiar, A., Azevedo, E. B., Tomé, A. R., and Coelho, F.: Climate change scenarios in the Azores and Madeira Islands, *World Resources Review*, 16, 19, 2004.](#)
- [Seemann, M., Bisseret, P., Tritz, J.-P., Hooper, A. B., and Rohmer, M.: Novel bacterial triterpenoids of the hopane series from *Nitrosomonas europaea* and their significance for the formation of the C35](#)

- 365 [bacteriohopane skeleton, Tetrahedron Letters, 40, 1681–1684, https://doi.org/10.1016/S0040-4039\(99\)00064-7, 1999.](https://doi.org/10.1016/S0040-4039(99)00064-7)
- [Sinninghe Damsté, J. S., Rijpstra, W. I. C., Dedysh, S. N., Foesel, B. U., and Villanueva, L.: Pheno- and Genotyping of Hopanoid Production in Acidobacteria, *Frontiers in Microbiology*, 8, 2017.](#)
- [Spencer-Jones, C. L., Wagner, T., Dinga, B. J., Schefuß, E., Mann, P. J., Poulsen, J. R., Spencer, R. G. M., Wabakanghanzi, J. N., and Talbot, H. M.: Bacteriohopanepolyols in tropical soils and sediments from the Congo River catchment area, *Organic Geochemistry*, 89–90, 1–13, https://doi.org/10.1016/j.orggeochem.2015.09.003, 2015.](#)
- 370 [Summons, R. E., Jahnke, L. L., Hope, J. M., and Logan, G. A.: 2-Methylhopanoids as biomarkers for cyanobacterial oxygenic photosynthesis, *Nature*, 400, 554–557, https://doi.org/10.1038/23005, 1999.](#)
- 375 [Talbot, H. M. and Farrimond, P.: Bacterial populations recorded in diverse sedimentary biohopanoid distributions, *Organic Geochemistry*, 38, 1212–1225, https://doi.org/10.1016/j.orggeochem.2007.04.006, 2007.](#)
- [Talbot, H. M., Watson, D. F., Murrell, J. C., Carter, J. F., and Farrimond, P.: Analysis of intact bacteriohopanepolyols from methanotrophic bacteria by reversed-phase high-performance liquid chromatography–atmospheric pressure chemical ionisation mass spectrometry, *Journal of Chromatography A*, 921, 175–185, https://doi.org/10.1016/S0021-9673\(01\)00871-8, 2001.](#)
- 380 [Talbot, H. M., Watson, D. F., Pearson, E. J., and Farrimond, P.: Diverse biohopanoid compositions of non-marine sediments, *Organic Geochemistry*, 34, 1353–1371, https://doi.org/10.1016/S0146-6380\(03\)00159-1, 2003.](#)
- 385 [Talbot, H. M., Rohmer, M., and Farrimond, P.: Rapid structural elucidation of composite bacterial hopanoids by atmospheric pressure chemical ionisation liquid chromatography/ion trap mass spectrometry, *Rapid Communications in Mass Spectrometry*, 21, 880–892, https://doi.org/10.1002/rcm.2911, 2007.](#)
- [Talbot, H. M., Summons, R. E., Jahnke, L. L., Cockell, C. S., Rohmer, M., and Farrimond, P.: Cyanobacterial bacteriohopanepolyol signatures from cultures and natural environmental settings, *Organic Geochemistry*, 39, 232–263, https://doi.org/10.1016/j.orggeochem.2007.08.006, 2008.](#)
- 390 [Talbot, H. M., Handley, L., Spencer-Jones, C. L., Dinga, B. J., Schefuß, E., Mann, P. J., Poulsen, J. R., Spencer, R. G. M., Wabakanghanzi, J. N., and Wagner, T.: Variability in aerobic methane oxidation over the past 1.2Myrs recorded in microbial biomarker signatures from Congo fan sediments, *Geochimica et Cosmochimica Acta*, 133, 387–401, https://doi.org/10.1016/j.gca.2014.02.035, 2014.](#)
- 395 [Taylor, K. A. and Harvey, H. R.: Bacterial hopanoids as tracers of organic carbon sources and processing across the western Arctic continental shelf, *Organic Geochemistry*, 42, 487–497, https://doi.org/10.1016/j.orggeochem.2011.03.012, 2011.](#)
- [Wagner, T., Kallweit, W., Talbot, H. M., Mollenhauer, G., Boom, A., and Zabel, M.: Microbial biomarkers support organic carbon transport from methane-rich Amazon wetlands to the shelf and deep sea fan during recent and glacial climate conditions, *Organic Geochemistry*, 67, 85–98, https://doi.org/10.1016/j.orggeochem.2013.12.003, 2014.](#)
- 400 [Watson, D. F. and Farrimond, P.: Novel polyfunctionalised geohopanoids in a recent lacustrine sediment \(Priest Pot, UK\), *Organic Geochemistry*, 31, 1247–1252, https://doi.org/10.1016/S0146-6380\(00\)00148-0, 2000.](#)

- 405 [Welander, P. V. and Summons, R. E.: Discovery, taxonomic distribution, and phenotypic characterization of a gene required for 3-methylhopanoid production, Proceedings of the National Academy of Sciences, 109, 12905–12910, <https://doi.org/10.1073/pnas.1208255109>, 2012.](#)
- [Welander, P. V., Hunter, R. C., Zhang, L., Sessions, A. L., Summons, R. E., and Newman, D. K.: Hopanoids Play a Role in Membrane Integrity and pH Homeostasis in *Rhodopseudomonas palustris* TIE-1, Journal of Bacteriology, 191, 6145–6156, <https://doi.org/10.1128/JB.00460-09>, 2009.](#)
- 410 [Welander, P. V., Coleman, M. L., Sessions, A. L., Summons, R. E., and Newman, D. K.: Identification of a methylase required for 2-methylhopanoid production and implications for the interpretation of sedimentary hopanes, Proc. Natl. Acad. Sci. U.S.A., 107, 8537–8542, <https://doi.org/10.1073/pnas.0912949107>, 2010.](#)
- [Whittenbury, R., Phillips, K. C., and Wilkinson, J. F.: Enrichment, Isolation and Some Properties of Methane-utilizing Bacteria, Journal of General Microbiology, 61, 205–218, <https://doi.org/10.1099/00221287-61-2-205>, 1970.](#)
- 415 [van Winden, J. F., Talbot, H. M., Kip, N., Reichart, G.-J., Pol, A., McNamara, N. P., Jetten, M. S. M., Op den Camp, H. J. M., and Sinninghe Damsté, J. S.: Bacteriohopanepolyol signatures as markers for methanotrophic bacteria in peat moss, Geochimica et Cosmochimica Acta, 77, 52–61, <https://doi.org/10.1016/j.gca.2011.10.026>, 2012.](#)
- 420 [van Winden, J. F., Talbot, H. M., Reichart, G.-J., McNamara, N. P., Benthien, A., and Sinninghe Damsté, J. S.: Influence of temperature on the \$\delta^{13}\text{C}\$ values and distribution of methanotroph-related hopanoids in Sphagnum-dominated peat bogs, Geobiology, 18, 497–507, <https://doi.org/10.1111/gbi.12389>, 2020.](#)
- [Xu, Y., Cooke, M. P., Talbot, H. M., and Simpson, M. J.: Bacteriohopanepolyol signatures of bacterial populations in Western Canadian soils, Organic Geochemistry, 40, 79–86, <https://doi.org/10.1016/j.orggeochem.2008.09.003>, 2009.](#)
- 425 [Zhu, C., Talbot, H. M., Wagner, T., Pan, J.-M., and Pancost, R. D.: Distribution of hopanoids along a land to sea transect: Implications for microbial ecology and the use of hopanoids in environmental studies, Limnology and Oceanography, 56, 1850–1865, <https://doi.org/10.4319/lo.2011.56.5.1850>, 2011.](#)
- [Zindorf, M., Rush, D., Jaeger, J., Mix, A., Penkrot, M. L., Schnetger, B., Sidgwick, F. R., Talbot, H. M., van der Land, C., Wagner, T., Walczak, M., and März, C.: Reconstructing oxygen deficiency in the glacial Gulf of Alaska: Combining biomarkers and trace metals as paleo-redox proxies, Chemical Geology, 558, 119864, <https://doi.org/10.1016/j.chemgeo.2020.119864>, 2020.](#)
- 430 [Zundel, M. and Rohmer, M.: Prokaryotic triterpenoids, European Journal of Biochemistry, 150, 23–27, <https://doi.org/10.1111/j.1432-1033.1985.tb08980.x>, 1985.](#)
- [Bale, N. J., Rijpstra, W. I. C., Sahonero-Canavesi, D. X., Oshkin, I. Y., Belova, S. E., Dedysh, S. N., and Sinninghe Damsté, J. S.: Fatty Acid and Hopanoid Adaption to Cold in the Methanotroph *Methylovulum psychrotolerans*, Frontiers in Microbiology, 10, 2019.](#)
- [Bale, N. J., Ding, S., Hopmans, E. C., Arts, M. G. I., Villanueva, L., Boschman, C., Haas, A. F., Schouten, S., and Sinninghe-Damsté, J. S.: Lipidomics of Environmental Microbial Communities. I: Visualization of Component Distributions Using Untargeted Analysis of High-Resolution Mass Spectrometry Data, Frontiers in Microbiology, 12, 2021.](#)
- 440 [Bastviken, D., Cole, J. J., Pace, M. L., and Van de Bogert, M. C.: Fates of methane from different lake habitats: Connecting whole-lake budgets and CH₄ emissions, Journal of Geophysical Research: Biogeosciences, 113, <https://doi.org/10.1029/2007JG000608>, 2008.](#)

Formatted: Indent: Left: 0 cm, Hanging: 0.5 cm, Don't add space between paragraphs of the same style, Line spacing: 1.5 lines

Formatted: Font: 10 pt, English (United States)

- 445 Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, *Canadian Journal of Biochemistry and Physiology*, 37, 911–917, <https://doi.org/10.1139/o59-099>, 1959.
- Blumenberg, M., Krüger, M., Nauhaus, K., Talbot, H. M., Oppermann, B. I., Seifert, R., Pape, T., and Michaelis, W.: Biosynthesis of hopanoids by sulfate-reducing bacteria (genus *Desulfovibrio*), *Environmental Microbiology*, 8, 1220–1227, <https://doi.org/10.1111/j.1462-2920.2006.01014.x>, 2006.
- 450 Blumenberg, M., Seifert, R., Kasten, S., Bahlmann, E., and Michaelis, W.: Euphotic zone bacterioplankton sources major sedimentary bacteriohopanepolyols in the Holocene Black Sea, *Geochimica et Cosmochimica Acta*, 73, 750–766, <https://doi.org/10.1016/j.gca.2008.11.005>, 2009.
- Blumenberg, M., Hoppert, M., Krüger, M., Dreier, A., and Thiel, V.: Novel findings on hopanoid occurrences among sulfate-reducing bacteria: Is there a direct link to nitrogen fixation?, *Organic Geochemistry*, 49, 1–5, <https://doi.org/10.1016/j.orggeochem.2012.05.003>, 2012.
- 455 Blumenberg, M., Berndmeyer, C., Moros, M., Muschalla, M., Schmale, O., and Thiel, V.: Bacteriohopanepolyols record stratification, nitrogen fixation and other biogeochemical perturbations in Holocene sediments of the central Baltic Sea, *Biogeosciences*, 10, 2725–2735, <https://doi.org/10.5194/bg-10-2725-2013>, 2013.
- Bradley, A. S., Pearson, A., Sáenz, J. P., and Marx, C. J.: Adenosylhopane: The first intermediate in hopanoid side chain-biosynthesis, *Organic Geochemistry*, 41, 1075–1081, <https://doi.org/10.1016/j.orggeochem.2010.07.003>, 2010.
- Bravo, J.-M., Perzl, M., Härtner, T., Kannenberg, E. L., and Rohmer, M.: Novel methylated triterpenoids of the gammacerane series from the nitrogen-fixing bacterium *Bradyrhizobium japonicum* USDA 110, *European Journal of Biochemistry*, 268, 1323–1331, <https://doi.org/10.1046/j.1432-1327.2001.01998.x>, 2001.
- 465 Cooke, M. P., Talbot, H. M., and Farrimond, P.: Bacterial populations recorded in bacteriohopanepolyol distributions in soils from Northern England, *Organic Geochemistry*, 39, 1347–1358, <https://doi.org/10.1016/j.orggeochem.2008.05.003>, 2008.
- Cooke, M. P., van Dongen, B. E., Talbot, H. M., Semiletov, I., Shakhova, N., Guo, L., and Gustafsson, Ö.: Bacteriohopanepolyol biomarker composition of organic matter exported to the Arctic Ocean by seven of the major Arctic rivers, *Organic Geochemistry*, 40, 1151–1159, <https://doi.org/10.1016/j.orggeochem.2009.07.014>, 2009.
- 470 Coolen, M. J. L., Talbot, H. M., Abbas, B. A., Ward, C., Schouten, S., Volkman, J. K., and Damsté, J. S. S.: Sources for sedimentary bacteriohopanepolyols as revealed by 16S rDNA stratigraphy, *Environmental Microbiology*, 10, 1783–1803, <https://doi.org/10.1111/j.1462-2920.2008.01601.x>, 2008.
- 475 Cordeiro, R., Luz, R., Vilaverde, J., Vasconcelos, V., Fonseca, A., and Gonçalves, V.: Distribution of Toxic Cyanobacteria in Volcanic Lakes of the Azores Islands, *Water*, 12, 3385, <https://doi.org/10.3390/w12123385>, 2020.
- Cvejic, J. H., Bodrossy, L., Kovács, K. L., and Rohmer, M.: Bacterial triterpenoids of the hopane series from the methanotrophic bacteria *Methylocaldum* spp.: phylogenetic implications and first evidence for an unsaturated aminobacteriohopanepolyol, *FEMS Microbiology Letters*, 182, 361–365, <https://doi.org/10.1111/j.1574-6968.2000.tb08922.x>, 2000.
- 480 De Jonge, C., Talbot, H. M., Bischoff, J., Stadnitskaia, A., Cherkashov, G., and Sinninghe-Damsté, J. S.: Bacteriohopanepolyol distribution in Yenisei River and Kara Sea suspended particulate matter and sediments

- traces terrigenous organic matter input, *Geochimica et Cosmochimica Acta*, 174, 85–101, <https://doi.org/10.1016/j.gea.2015.11.008>, 2016.
- 485 Doğrul Selver, A., Talbot, H. M., Gustafsson, Ö., Boulton, S., and van Dongen, B. E.: Soil organic matter transport along a sub-Arctic river–sea transect, *Organic Geochemistry*, 51, 63–72, <https://doi.org/10.1016/j.orggeochem.2012.08.002>, 2012.
- 490 Doğrul Selver, A., Sparkes, R. B., Bischoff, J., Talbot, H. M., Gustafsson, Ö., Semiletov, I. P., Dudarev, O. V., Boulton, S., and van Dongen, B. E.: Distributions of bacterial and archaeal membrane lipids in surface sediments reflect differences in input and loss of terrestrial organic carbon along a cross-shelf Arctic transect, *Organic Geochemistry*, 83–84, 16–26, <https://doi.org/10.1016/j.orggeochem.2015.01.005>, 2015.
- Doughty, D. M., Coleman, M. L., Hunter, R. C., Sessions, A. L., Summons, R. E., and Newman, D. K.: The RND-family transporter, HpnN, is required for hopanoid localization to the outer membrane of
- 495 *Rhodopseudomonas palustris* TIE-1, *Proceedings of the National Academy of Sciences*, 108, E1045–E1051, <https://doi.org/10.1073/pnas.1104209108>, 2011.
- Elling, F. J., Evans, T. W., Nathan, V., Hemingway, J. D., Kharbush, J. J., Bayer, B., Spieck, E., Husain, F., Summons, R. E., and Pearson, A.: Marine and terrestrial nitrifying bacteria are sources of diverse bacteriohopanepolyols, *Geobiology*, 20, 399–420, <https://doi.org/10.1111/gbi.12484>, 2022.
- 500 Farrimond, P., Head, I. M., and Innes, H. E.: Environmental influence on the biohopanoid composition of recent sediments, *Geochimica et Cosmochimica Acta*, 64, 2985–2992, [https://doi.org/10.1016/S0016-7037\(00\)00404-X](https://doi.org/10.1016/S0016-7037(00)00404-X), 2000.
- van Grinsven, S., Sinnighe Damsté, J. S., Harrison, J., and Villanueva, L.: Impact of Electron Acceptor Availability on Methane-Influenced Microorganisms in an Enrichment Culture Obtained From a Stratified
- 505 Lake, *Frontiers in Microbiology*, 11, 2020.
- Guggenheim, C., Freimann, R., Mayr, M. J., Beck, K., Wehrli, B., and Bürgmann, H.: Environmental and Microbial Interactions Shape Methane-Oxidizing Bacterial Communities in a Stratified Lake, *Frontiers in Microbiology*, 11, 2020.
- Hanson, R. S.: Ecology and Diversity of Methylotrophic Organisms, in: *Advances in Applied Microbiology*, vol. 26, edited by: Perlman, D., Academic Press, 3–39, [https://doi.org/10.1016/S0065-2164\(08\)70328-9](https://doi.org/10.1016/S0065-2164(08)70328-9), 1980.
- Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiological Reviews*, 60, 439–471, <https://doi.org/10.1128/mr.60.2.439-471.1996>, 1996.
- Harriss, S. M. and Hanson, R. S.: Stratification of aerobic methane-oxidizing organisms in Lake Mendota,
- 515 Madison, Wisconsin, *Limnology and Oceanography*, 25, 412–421, <https://doi.org/10.4319/lo.1980.25.3.0412>, 1980.
- Hernández, A., Kutiel, H., Trigo, R. M., Valente, M. A., Sigró, J., Cropper, T., and Santo, F. E.: New Azores archipelago daily precipitation dataset and its links with large-scale modes of climate variability, *International Journal of Climatology*, 36, 4439–4454, <https://doi.org/10.1002/joc.4642>, 2016.
- 520 Hopmans, E. C., Smit, N. T., Schwartz-Narbonne, R., Sinnighe Damsté, J. S., and Rush, D.: Analysis of non-derivatized bacteriohopanepolyols using UHPLC-HRMS reveals great structural diversity in environmental lipid assemblages, *Organic Geochemistry*, 160, 104285, <https://doi.org/10.1016/j.orggeochem.2021.104285>, 2021.

- Kuseh, S. and Rush, D.: Revisiting the precursors of the most abundant natural products on Earth: A look back at 30+ years of bacteriohopanepolyol (BHP) research and ahead to new frontiers, *Organic Geochemistry*, 172, 104469, <https://doi.org/10.1016/j.orggeochem.2022.104469>, 2022.
- Kuseh, S., Sepúlveda, J., and Wakeham, S. G.: Origin of Sedimentary BHPs Along a Mississippi River–Gulf of Mexico Export Transect: Insights From Spatial and Density Distributions, *Frontiers in Marine Science*, 6, 2019.
- 530 Kuseh, S., Wakeham, S. G., Dildar, N., Zhu, C., and Sepúlveda, J.: Bacterial and archaeal lipids trace chemo(auto)trophy along the redoxcline in Vancouver Island fjords, *Geobiology*, 19, 521–541, <https://doi.org/10.1111/gbi.12446>, 2021a.
- Kuseh, S., Wakeham, S. G., and Sepúlveda, J.: Diverse origins of “soil marker” bacteriohopanepolyols in marine oxygen-deficient zones, *Organic Geochemistry*, 151, 104150, <https://doi.org/10.1016/j.orggeochem.2020.104150>, 2021b.
- 535 Matys, E. D., Sepúlveda, J., Pantoja, S., Lange, C. B., Caniupán, M., Lamy, F., and Summons, R. E.: Bacteriohopanepolyols along redox gradients in the Humboldt Current System off northern Chile, *Geobiology*, 15, 844–857, <https://doi.org/10.1111/gbi.12250>, 2017.
- Meyers, P. A.: Preservation of elemental and isotopic source identification of sedimentary organic matter, *Chemical Geology*, 114, 289–302, [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0), 1994.
- 540 Meyers, P. A.: Applications of organic geochemistry to paleolimnological reconstructions: a summary of examples from the Laurentian Great Lakes, *Organic Geochemistry*, 34, 261–289, [https://doi.org/10.1016/S0146-6380\(02\)00168-7](https://doi.org/10.1016/S0146-6380(02)00168-7), 2003.
- Mitrović, D., Hopmans, E. C., Bale, N. J., Richter, N., Amaral-Zettler, L. A., Baxter, A. J., Peterse, F., Miguel Raposo, P., Gonçalves, V., Cristina Costa, A., and Schouten, S.: Isoprenoidal GDGTs and GDDs associated with anoxic lacustrine environments, *Organic Geochemistry*, 178, 104582, <https://doi.org/10.1016/j.orggeochem.2023.104582>, 2023.
- 545 Neunlist, S. and Rohmer, M.: A novel hopanoid, 30-(5'-adenosyl)hopane, from the purple non-sulphur bacterium *Rhodospseudomonas acidophila*, with possible DNA interactions., *Biochem J*, 228, 769–771, 1985a.
- Neunlist, S. and Rohmer, M.: Novel hopanoids from the methylotrophic bacteria *Methylococcus capsulatus* and *Methylomonas methanica*. (22S)-35-aminobacteriohopane-30,31,32,33,34-pentol and (22S)-35-amino-3β-methylbacteriohopane-30,31,32,33,34-pentol, *Biochemical Journal*, 231, 635–639, <https://doi.org/10.1042/bj2310635>, 1985b.
- 555 Neunlist, S. and Rohmer, M.: The Hopanoids of ‘*Methylosinus trichosporium*’: Aminobacteriohopanetriol and Aminobacteriohopanetetrol, *Microbiology*, 131, 1363–1367, <https://doi.org/10.1099/00221287-131-6-1363>, 1985c.
- Neunlist, S., Bissleret, P., and Rohmer, M.: The hopanoids of the purple non-sulfur bacteria *Rhodospseudomonas palustris* and *Rhodospseudomonas acidophila* and the absolute configuration of bacteriohopanetetrol, *European Journal of Biochemistry*, 171, 245–252, <https://doi.org/10.1111/j.1432-1033.1988.tb13783.x>, 1988.
- 560

Formatted: Font: 10 pt

Formatted: Font: 10 pt, English (United States)

- O'Beirne, M. D., Sparkes, R., Hamilton, T. L., van Dongen, B. E., Gilhooly, W. P., and Werne, J. P.: Characterization of diverse bacteriohopanepolyols in a permanently stratified, hyper-euxinic lake, *Organic Geochemistry*, 168, 104431, <https://doi.org/10.1016/j.orggeochem.2022.104431>, 2022.
- 565 Osborne, K. A.: Environmental controls on bacteriohopanepolyol signatures in estuarine sediments, Newcastle University, UK, 2015.
- Osborne, K. A., Gray, N. D., Sherry, A., Leary, P., Mejeha, O., Bisehoff, J., Rush, D., Sidgwick, F. R., Birgel, D., Kalyuzhnaya, M. G., and Talbot, H. M.: Methanotroph-derived bacteriohopanepolyol signatures as a function of temperature related growth, survival, cell death and preservation in the geological record, *Environmental Microbiology Reports*, 9, 492–500, <https://doi.org/10.1111/1758-2229.12570>, 2017.
- 570 Oswald, K., Milucka, J., Brand, A., Littmann, S., Wehrli, B., Kuypers, M. M. M., and Schubert, C. J.: Light-Dependent Aerobic Methane Oxidation Reduces Methane Emissions from Seasonally Stratified Lakes, *PLOS ONE*, 10, e0132574, <https://doi.org/10.1371/journal.pone.0132574>, 2015.
- Oswald, K., Milucka, J., Brand, A., Haeh, P., Littmann, S., Wehrli, B., Kuypers, M. M. M., and Schubert, C. J.: Aerobic gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters, *Limnology and Oceanography*, 61, S101–S118, <https://doi.org/10.1002/lno.10312>, 2016.
- 575 Ourisson, G. and Albrecht, P.: Hopanoids. I. Geohopanoids: the most abundant natural products on Earth?, *Acc. Chem. Res.*, 25, 398–402, <https://doi.org/10.1021/ar00021a003>, 1992.
- Pearson, A., Leavitt, W. D., Sáenz, J. P., Summons, R. E., Tam, M. C.-M., and Close, H. G.: Diversity of hopanoids and squalene-hopene cyclases across a tropical land-sea gradient, *Environmental Microbiology*, 11, 1208–1223, <https://doi.org/10.1111/j.1462-2920.2008.01817.x>, 2009.
- Pereira, C. L., Raposeiro, P. M., Costa, A. C., Bao, R., Giralt, S., and Gonçalves, V.: Biogeography and lake morphometry drive diatom and chironomid assemblages' composition in lacustrine surface sediments of oceanic islands, *Hydrobiologia*, 730, 93–112, <https://doi.org/10.1007/s10750-014-1824-6>, 2014.
- 585 Raposeiro, P. M., Rubio, M. J., González, A., Hernández, A., Sánchez-López, G., Vázquez Loureiro, D., Rull, V., Bao, R., Costa, A. C., Gonçalves, V., Sáez, A., and Giralt, S.: Impact of the historical introduction of exotic fishes on the chironomid community of Lake Azul (Azores Islands), *Palaeogeography, Palaeoclimatology, Palaeoecology*, 466, 77–88, <https://doi.org/10.1016/j.palaeo.2016.11.015>, 2017.
- Raposeiro, P. M., Saez, A., Giralt, S., Costa, A. C., and Gonçalves, V.: Causes of spatial distribution of subfossil diatom and chironomid assemblages in surface sediments of a remote deep island lake, *Hydrobiologia*, 815, 141–163, <https://doi.org/10.1007/s10750-018-3557-4>, 2018.
- 590 Rethemeyer, J., Schubotz, F., Talbot, H. M., Cooke, M. P., Hinrichs, K.-U., and Mollenhauer, G.: Distribution of polar membrane lipids in permafrost soils and sediments of a small high Arctic catchment, *Organic Geochemistry*, 41, 1130–1145, <https://doi.org/10.1016/j.orggeochem.2010.06.004>, 2010.
- 595 Richter, N., Russell, J. M., Amaral-Zettler, L., DeGroff, W., Raposeiro, P. M., Gonçalves, V., de Boer, E. J., Pla-Rabes, S., Hernández, A., Benavente, M., Ritter, C., Sáez, A., Bao, R., Trigo, R. M., Prego, R., and Giralt, S.: Long-term hydroclimate variability in the sub-tropical North Atlantic and anthropogenic impacts on lake ecosystems: A case study from Flores Island, the Azores, *Quaternary Science Reviews*, 285, 107525, <https://doi.org/10.1016/j.quaseirev.2022.107525>, 2022.
- 600 Rohmer, M., Bouvier-Nave, P., and Ourisson, G.: 1984: Distribution of Hopanoid Triterpenes in Prokaryotes, *Microbiology*, 130, 1137–1150, <https://doi.org/10.1099/00221287-130-5-1137>, 1984.

- Rudd, J. W. M., Furutani, A., Flett, R. J., and Hamilton, R. D.: Factors controlling methane oxidation in shield lakes: The role of nitrogen fixation and oxygen concentration, *Limnology and Oceanography*, 21, 357–364, <https://doi.org/10.4319/lo.1976.21.3.0357>, 1976.
- 605 Rush, D., Osborne, K. A., Birgel, D., Kappler, A., Hirayama, H., Peckmann, J., Poulton, S. W., Nickel, J. C., Mangelsdorf, K., Kalyuzhnaya, M., Sidgwick, F. R., and Talbot, H. M.: The Baeteriohopanepolyol Inventory of Novel Aerobic Methane Oxidising Bacteria Reveals New Biomarker Signatures of Aerobic Methanotrophy in Marine Systems, *PLOS ONE*, 11, e0165635, <https://doi.org/10.1371/journal.pone.0165635>, 2016.
- 610 Rush, D., Talbot, H. M., van der Meer, M. T. J., Hopmans, E. C., Douglas, B., and Sinninghe Damsté, J. S.: Biomarker evidence for the occurrence of anaerobic ammonium oxidation in the eastern Mediterranean Sea during Quaternary and Pliocene sapropel formation, *Biogeosciences*, 16, 2467–2479, <https://doi.org/10.5194/bg-16-2467-2019>, 2019.
- Sáenz, J. P.: Hopanoid enrichment in a detergent resistant membrane fraction of *Crocospaera watsonii*: Implications for bacterial lipid raft formation, *Organic Geochemistry*, 41, 853–856, <https://doi.org/10.1016/j.orggeochem.2010.05.005>, 2010.
- 615 Sáenz, J. P., Eglinton, T. I., and Summons, R. E.: Abundance and structural diversity of bacteriohopanepolyols in suspended particulate matter along a river to ocean transect, *Organic Geochemistry*, 42, 774–780, <https://doi.org/10.1016/j.orggeochem.2011.05.006>, 2011.
- 620 Sáenz, J. P., Sezgin, E., Schwille, P., and Simons, K.: Functional convergence of hopanoids and sterols in membrane ordering, *Proceedings of the National Academy of Sciences*, 109, 14236–14240, <https://doi.org/10.1073/pnas.1212141109>, 2012.
- Santos, F. D., Valente, M. A., Miranda, P. M. A., Aguiar, A., Azevedo, E. B., Tomé, A. R., and Coelho, F.: Climate change scenarios in the Azores and Madeira Islands, *World Resources Review*, 16, 19, 2004.
- 625 Seemann, M., Bisseret, P., Tritz, J. P., Hooper, A. B., and Rohmer, M.: Novel bacterial triterpenoids of the hopane series from *Nitrosomonas europaea* and their significance for the formation of the C35 bacteriohopane skeleton, *Tetrahedron Letters*, 40, 1681–1684, [https://doi.org/10.1016/S0040-4039\(99\)00064-7](https://doi.org/10.1016/S0040-4039(99)00064-7), 1999.
- Sinninghe Damsté, J. S., Rijpstra, W. I. C., Dedys, S. N., Foesel, B. U., and Villanueva, L.: Pheno- and Genotyping of Hopanoid Production in Acidobacteria, *Frontiers in Microbiology*, 8, 2017.
- 630 Spencer-Jones, C. L., Wagner, T., Dinga, B. J., Sehefuß, E., Mann, P. J., Poulsen, J. R., Spencer, R. G. M., Wabakanghanzi, J. N., and Talbot, H. M.: Bacteriohopanepolyols in tropical soils and sediments from the Congo River catchment area, *Organic Geochemistry*, 89–90, 1–13, <https://doi.org/10.1016/j.orggeochem.2015.09.003>, 2015.
- 635 Summons, R. E., Jahnke, L. L., Hope, J. M., and Logan, G. A.: 2-Methylhopanoids as biomarkers for cyanobacterial oxygenic photosynthesis, *Nature*, 400, 554–557, <https://doi.org/10.1038/23005>, 1999.
- Talbot, H. M. and Farrimond, P.: Bacterial populations recorded in diverse sedimentary biohopanoid distributions, *Organic Geochemistry*, 38, 1212–1225, <https://doi.org/10.1016/j.orggeochem.2007.04.006>, 2007.
- 640 Talbot, H. M., Watson, D. F., Murrell, J. C., Carter, J. F., and Farrimond, P.: Analysis of intact bacteriohopanepolyols from methanotrophic bacteria by reversed-phase high-performance liquid

- chromatography–atmospheric-pressure-chemical-ionisation-mass-spectrometry, *Journal of Chromatography A*, 921, 175–185, [https://doi.org/10.1016/S0021-9673\(01\)00871-8](https://doi.org/10.1016/S0021-9673(01)00871-8), 2001.
- 645 Talbot, H. M., Watson, D. F., Pearson, E. J., and Farrimond, P.: Diverse biohopanoid compositions of non-marine sediments, *Organic Geochemistry*, 34, 1353–1371, [https://doi.org/10.1016/S0146-6380\(03\)00159-1](https://doi.org/10.1016/S0146-6380(03)00159-1), 2003.
- Talbot, H. M., Rohmer, M., and Farrimond, P.: Rapid structural elucidation of composite bacterial hopanoids by atmospheric-pressure-chemical-ionisation-liquid-chromatography/ion-trap-mass-spectrometry, *Rapid Communications in Mass Spectrometry*, 21, 880–892, <https://doi.org/10.1002/rem.2911>, 2007.
- 650 Talbot, H. M., Summons, R. E., Jahnke, L. L., Cockell, C. S., Rohmer, M., and Farrimond, P.: Cyanobacterial bacteriohopanepolyol signatures from cultures and natural environmental settings, *Organic Geochemistry*, 39, 232–263, <https://doi.org/10.1016/j.orggeochem.2007.08.006>, 2008.
- Talbot, H. M., Handley, L., Spencer-Jones, C. L., Dinga, B. J., Schefuß, E., Mann, P. J., Poulsen, J. R., Spencer, R. G. M., Wabakanghanzi, J. N., and Wagner, T.: Variability in aerobic methane oxidation over the past 1.2 Myrs recorded in microbial biomarker signatures from Congo fan sediments, *Geochimica et Cosmochimica Acta*, 133, 387–401, <https://doi.org/10.1016/j.gea.2014.02.035>, 2014.
- 655 Talbot, H. M., Bischoff, J., Inglis, G. N., Collinson, M. E., and Pancost, R. D.: Polyfunctionalised bio- and geohopanooids in the Eocene Cobham Lignite, *Organic Geochemistry*, 96, 77–92, <https://doi.org/10.1016/j.orggeochem.2016.03.006>, 2016.
- 660 Taylor, K. A. and Harvey, H. R.: Bacterial hopanoids as tracers of organic carbon sources and processing across the western Arctic continental shelf, *Organic Geochemistry*, 42, 487–497, <https://doi.org/10.1016/j.orggeochem.2011.03.012>, 2011.
- Wagner, T., Kallweit, W., Talbot, H. M., Mollenhauer, G., Boom, A., and Zabel, M.: Microbial biomarkers support organic carbon transport from methane-rich Amazon wetlands to the shelf and deep-sea fan during recent and glacial climate conditions, *Organic Geochemistry*, 67, 85–98, <https://doi.org/10.1016/j.orggeochem.2013.12.003>, 2014.
- 665 Watson, D. F. and Farrimond, P.: Novel polyfunctionalised geohopanooids in a recent lacustrine sediment (Priest Pot, UK), *Organic Geochemistry*, 31, 1247–1252, [https://doi.org/10.1016/S0146-6380\(00\)00148-0](https://doi.org/10.1016/S0146-6380(00)00148-0), 2000.
- Welander, P. V. and Summons, R. E.: Discovery, taxonomic distribution, and phenotypic characterization of a gene required for 3-methylhopanoid production, *Proceedings of the National Academy of Sciences*, 109, 12905–12910, <https://doi.org/10.1073/pnas.1208255109>, 2012.
- 670 Welander, P. V., Hunter, R. C., Zhang, L., Sessions, A. L., Summons, R. E., and Newman, D. K.: Hopanooids Play a Role in Membrane Integrity and pH Homeostasis in *Rhodospseudomonas palustris* TIE-1, *Journal of Bacteriology*, 191, 6145–6156, <https://doi.org/10.1128/JB.00460-09>, 2009.
- 675 Welander, P. V., Coleman, M. L., Sessions, A. L., Summons, R. E., and Newman, D. K.: Identification of a methylase required for 2-methylhopanoid production and implications for the interpretation of sedimentary hopanes, *Proc. Natl. Acad. Sci. U.S.A.*, 107, 8537–8542, <https://doi.org/10.1073/pnas.0912949107>, 2010.
- Whittenbury, R., Phillips, K. C., and Wilkinson, J. F.: Enrichment, Isolation and Some Properties of Methane-utilizing Bacteria, *Journal of General Microbiology*, 61, 205–218, <https://doi.org/10.1099/00221287-61-2-205>, 1970.
- 680

- van Winden, J. F., Talbot, H. M., Kip, N., Reichart, G. J., Pol, A., McNamara, N. P., Jetten, M. S. M., Op den Camp, H. J. M., and Sinninghe Damsté, J. S.: Bacteriohopanepolyol signatures as markers for methanotrophic bacteria in peat moss, *Geochimica et Cosmochimica Acta*, 77, 52–61, <https://doi.org/10.1016/j.gea.2011.10.026>, 2012.
- 685 van Winden, J. F., Talbot, H. M., Reichart, G. J., McNamara, N. P., Benthien, A., and Sinninghe Damsté, J. S.: Influence of temperature on the $\delta^{13}\text{C}$ values and distribution of methanotroph-related hopanoids in Sphagnum-dominated peat bogs, *Geobiology*, 18, 497–507, <https://doi.org/10.1111/gbi.12389>, 2020.
- Xu, Y., Cooke, M. P., Talbot, H. M., and Simpson, M. J.: Bacteriohopanepolyol signatures of bacterial populations in Western Canadian soils, *Organic Geochemistry*, 40, 79–86, <https://doi.org/10.1016/j.orggeochem.2008.09.003>, 2009.
- 690 Zhu, C., Talbot, H. M., Wagner, T., Pan, J.-M., and Pancost, R. D.: Distribution of hopanoids along a land-to-sea transect: Implications for microbial ecology and the use of hopanoids in environmental studies, *Limnology and Oceanography*, 56, 1850–1865, <https://doi.org/10.4319/lo.2011.56.5.1850>, 2011.
- Zindorf, M., Rush, D., Jaeger, J., Mix, A., Penkrot, M. L., Schnetger, B., Sidgwick, F. R., Talbot, H. M., van der Land, C., Wagner, T., Waleczak, M., and März, C.: Reconstructing oxygen deficiency in the glacial Gulf of Alaska: Combining biomarkers and trace metals as paleo-redox proxies, *Chemical Geology*, 558, 119864, <https://doi.org/10.1016/j.chemgeo.2020.119864>, 2020.
- 695 Zundel, M. and Rohmer, M.: Prokaryotic triterpenoids, *European Journal of Biochemistry*, 150, 23–27, <https://doi.org/10.1111/j.1432-1033.1985.tb08980.x>, 1985.

700

Formatted: Font: 10 pt

Formatted: Font: 10 pt, English (United States)

Page 26: [1] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [2] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [3] Formatted Table **Nora Richter** **21-4-2023 10:32:00**

Formatted Table

▲
Page 26: [4] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [5] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [6] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [7] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [8] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [9] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [10] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [11] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [12] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [13] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [14] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [15] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [16] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [17] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [18] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [19] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [20] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [21] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [22] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [23] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [24] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [25] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [26] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [27] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [28] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [29] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [30] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [31] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [32] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [33] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [34] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [35] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲

Page 26: [36] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 26: [37] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [38] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 26: [39] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [40] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 26: [41] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [42] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 26: [43] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [44] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 26: [45] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [46] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 26: [47] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [48] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 26: [49] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [50] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 26: [51] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 26: [52] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 26: [53] Formatted	Nora Richter	21-4-2023 10:31:00

Left

▲
Page 26: [54] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [55] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [56] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [57] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [58] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [59] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [60] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [61] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [62] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [63] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [64] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [65] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [66] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [67] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [68] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [69] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [70] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲

Page 26: [71] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [72] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [73] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [74] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [75] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [76] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [77] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [78] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [79] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [80] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [81] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [82] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [83] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [84] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [85] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [86] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [87] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [88] Formatted **Nora Richter** **21-4-2023 10:32:00**

Page 26: [106] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [107] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [108] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [109] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [110] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [111] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [112] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [113] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [114] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [115] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [116] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [117] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [118] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [119] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [120] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [121] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [122] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [123] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [124] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [125] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [126] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [127] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [128] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [129] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [130] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [131] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [132] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [133] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [134] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [135] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [136] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [137] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [138] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [139] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [140] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲

Page 26: [141] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [142] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [143] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [144] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [145] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [146] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [147] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [148] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [149] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [150] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [151] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [152] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [153] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [154] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [155] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [156] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [157] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [158] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [159] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [160] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [161] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [162] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [163] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 26: [164] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 26: [165] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 26: [166] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [167] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [168] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [169] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [170] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [171] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [172] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [173] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [174] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [175] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲

Page 27: [176] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [177] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [178] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [179] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [180] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [181] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [182] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [183] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [184] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [185] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [186] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [187] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [188] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [189] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [190] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [191] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [192] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [193] Formatted	Nora Richter	21-4-2023 10:32:00

Centered

▲
Page 27: [194] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [195] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [196] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [197] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [198] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [199] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [200] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [201] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [202] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [203] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [204] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [205] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [206] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [207] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [208] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [209] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [210] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲

Page 27: [211] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [212] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [213] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [214] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [215] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [216] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [217] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [218] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [219] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [220] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [221] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [222] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [223] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [224] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [225] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [226] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [227] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [228] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [229] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [230] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [231] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [232] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [233] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [234] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [235] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [236] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [237] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [238] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [239] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [240] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [241] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [242] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [243] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [244] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [245] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲

Page 27: [246] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [247] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [248] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [249] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [250] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [251] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [252] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [253] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [254] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [255] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [256] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [257] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [258] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [259] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [260] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [261] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [262] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [263] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [264] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [265] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [266] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [267] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [268] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [269] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [270] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [271] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [272] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [273] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [274] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [275] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [276] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [277] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [278] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [279] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [280] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲

Page 27: [281] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [282] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [283] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [284] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [285] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [286] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [287] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [288] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [289] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [290] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [291] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [292] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [293] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [294] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [295] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [296] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [297] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [298] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [299] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [300] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [301] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [302] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [303] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [304] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [305] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [306] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [307] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [308] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [309] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [310] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [311] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [312] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [313] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [314] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [315] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲

Page 27: [316] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [317] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [318] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [319] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [320] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [321] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [322] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [323] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [324] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [325] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [326] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [327] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [328] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [329] Formatted	Nora Richter	21-4-2023 10:32:00
Centered		
Page 27: [330] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [331] Formatted	Nora Richter	21-4-2023 10:30:00
Font: 7 pt		
Page 27: [332] Formatted	Nora Richter	21-4-2023 10:31:00
Left		
Page 27: [333] Formatted	Nora Richter	21-4-2023 10:32:00

Centered

▲
Page 27: [334] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [335] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [336] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [337] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [338] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [339] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [340] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [341] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [342] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [343] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [344] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [345] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [346] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [347] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 27: [348] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 27: [349] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 27: [350] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲

Page 28: [351] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [352] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [353] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [354] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [355] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [356] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [357] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [358] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [359] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [360] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [361] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [362] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [363] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [364] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [365] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [366] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [367] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [368] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [369] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [370] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [371] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [372] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [373] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [374] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [375] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [376] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [377] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [378] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [379] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [380] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [381] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [382] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [383] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [384] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [385] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲

Page 28: [386] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [387] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [388] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [389] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [390] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [391] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [392] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [393] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [394] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [395] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [396] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [397] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [398] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [399] Formatted **Nora Richter** **21-4-2023 10:30:00**

Font: 7 pt

▲
Page 28: [400] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲
Page 28: [401] Formatted **Nora Richter** **21-4-2023 10:32:00**

Centered

▲
Page 28: [402] Formatted **Nora Richter** **21-4-2023 10:31:00**

Left

▲