

Supplement – Ntirugulirwa et al

Table S1. Information about the germplasm material and propagation in the nursery prior to plantation at the sites.

Table S2. Stem base diameter (5 cm above soil) at planting in Dec 2017/Jan 2018.

Table S3. Stem height at planting in Dec 2017/Jan 2018.

Table S4. Classification of species into the successional groups (SG) early (ES) and late successional (LS) and the references supporting the classification.

Table S5. P-values and total degrees of freedom (df Tot) of one-way ANOVA for species-specific site effects on D_{base} ; h ; D-RGR_{D10-25}; D-RGR_{D10-25}, H-RGR_{H75-100}; H-RGR_{H250-300}; stems#), and tree mortality.

Table S6. Number of individuals recorded dead at each tri-monthly census of each species and site.

Figure S1. Site map and experimental design at the high elevation site (Sigira).

Figure S2. Site map and experimental design at the mid elevation site (Rubona).

Figure S3. Site map and experimental design at the low elevation site (Makera).

Figure S4. The relation between height and base diameter and the relative growth rate (RGR) for height (H-RGR) and base diameter (D-RGR) in relation to height and base diameter, respectively, for all 20 species measured tri-monthly over two years at all three sites.

Figure S5 a-t. The development of the stem base diameter measured every third month over two years and the relative growth rate (RGR) between measurement intervals for 20 species grown at three sites along an elevation gradient

Figure S6 a-t. The development of the stem height measured every third month over two years and the relative growth rate (RGR) between measurement intervals for 20 species grown at 3 sites along an elevation gradient

Table S1. Information about the germplasm material and propagation in the nursery prior to plantation at the sites. Provenance's location of collection: NYU, Nyungwe tropical montane forest; RUB, Rubona research station; NDI, Ndiza tropical montane forest; RUH, Ruhanda arboretum.

Species:	Scientific name	Code	Germplasm material:					Propagation in Nursery:					
			Provenance	Location of collection	Date of collection	No of mother material trees	Type of material	1000-seed weight ¹ (g)	Moisture content (%)	Start of propagation date	Date of transplantsation	Days in seed-beds	Final no of survivals
<i>Afrocarpus falcatus</i>	<i>Afa</i>	NYU	RUH	2016-06-20	10-15	Seed	5.700	25	2016-08-03	2016-11-08	97	420	100
<i>Albizia gummifera</i>	<i>Agu</i>	NYU	RUB	2017-08-25	1	Seed	80	6	2017-08-31	2017-09-15	15	308	49
<i>Bridelia bridelifolia</i>	<i>Bbr</i>	NYU	NYU	2016-07-20	1-5	Seed	67	23	2016-08-03	2016-09-07	35	692	84
<i>Bridelia micrantha</i>	<i>Bmi</i>	RUB	RUB	2016-09-14	1-5	Seed	50	25	2016-10-03	2016-11-09	37	374	100
<i>Carapa grandiflora</i>	<i>Cgr</i>	NYU	NYU	2016-06-30	10-15	Seed	28.600	32	2016-08-03	2016-11-10	99	528	100
<i>Chrysophyllum gorungosanum</i>	<i>Cgo</i>	NYU	RUH	2016-11-21	5-10	Saplings	600	600	2016-11-21	2016-11-21	378	378	100
<i>Croton megalocarpus</i>	<i>Cme</i>	NYU	RUB	2016-01-10	10-15	Seed	800	8	2016-09-22	2016-10-10	18	562	99
<i>Dombeya tomentosa</i>	<i>Dto</i>	NYU	RUH	2016-11-24	20-30	Cuttings	4	4	2016-11-24	2016-11-24	630	630	100
<i>Entandrophragma excelsum</i>	<i>Eex</i>	NYU	RUH	2016-07-11	5-10	Saplings	634	634	2016-11-07	2016-11-07	410	410	100
<i>Faurea saligna</i>	<i>Fsa</i>	NYU	NYU	2016-04-02	1-5	Seed	4	4	2016-08-03	2016-10-19	77	304	87
<i>Ficus thonningii</i>	<i>Fth</i>	NYU	NYU	2016-09-20	1	Cuttings	11	11	2016-09-20	2016-09-20	1047	97	
<i>Harungana madagascariensis</i>	<i>Hma</i>	RUB	RUB	2016-04-02	10-20	Seed	11	10	2016-08-16	2016-09-07	22	1050	71
<i>Harungana montana</i>	<i>Hmo</i>	NYU	NYU	2016-10-05	1-5	Seed	13	10	2016-08-16	2016-11-10	86	352	90
<i>Macaranga kilimandscharica</i>	<i>Mki</i>	NDI	NDI	2017-05-30	5-10	Seed	60	15	2017-07-01	2017-08-04	34	698	93
<i>Maesa lanceolata</i>	<i>Mla</i>	NYU	NYU	2016-06-20	1-5	Seed	9	6	2016-08-03	2016-10-27	85	454	98
<i>Markhamia lutea</i>	<i>Mlu</i>	NYU	RUB	2016-01-06	1-5	Seed	40	10	2016-08-03	2016-09-14	42	1679	97
<i>Nectandra buchananii</i>	<i>Nbu</i>	NYU	NYU	2016-10-07	1-5	Seed	352	33	2016-09-13	2016-10-30	47	632	65
<i>Polyscias fulva</i>	<i>Pfu</i>	NYU	NYU	2016-07-20	1-5	Seed	11	17	2016-08-03	2016-12-02	121	326	100
<i>Prunus africana</i>	<i>Paf</i>	NYU	NYU	2017-04-28	1	Seed	200	19	2017-04-29	2017-05-17	18	1102	92
<i>Syzygium guineense</i>	<i>Sgu</i>	NYU	NYU	2016-10-18	1-5	Saplings	367	367	2016-10-20	2016-10-20	546	546	85

Table S2. Stem base diameter (5 cm above soil) at planting in Dec 2017/Jan 2018. Mean and standard deviations as well as P-values from one-way ANOVA tests are calculated on plot means (n=18) based on five individuals, i.e. 90 trees per species and site. All individuals within species were randomly selected from the nursery. Full name of species is given in Table S1. Different letters for sites indicate significant differences ($P<0.05$).

Species	HE		ME		LE		P-value	ME (% of HE)	LE (% of HE)
	Mean	SD	Mean	SD	Mean	SD			
Afa	4.6 ± 0.5		4.9 ± 0.4		4.8 ± 0.7		0.31	6	5
Agu	1.0 ± 0.0		1.0 ± 0.1		1.0 ± 0.0		0.77	1	0
Bbr	4.4 ± 0.3 a		5.0 ± 0.4 b		4.3 ± 0.4 a		<0.001	14	-2
Bmi	4.5 ± 0.4 a		4.5 ± 0.4 a		4.2 ± 0.4 b		0.010	0	-8
Cgo	3.0 ± 0.5		2.9 ± 0.6		3.0 ± 0.5		0.92	-2	0
Cgr	7.7 ± 0.9		7.8 ± 0.8		7.5 ± 0.8		0.49	1	-3
Cme	4.0 ± 0.2		3.9 ± 0.3		4.0 ± 0.3		0.71	-2	0
Dto	9.6 ± 1.4		9.0 ± 1.3		8.9 ± 1.7		0.38	-6	-7
Eex	8.5 ± 1.4		8.8 ± 1.6		8.7 ± 1.6		0.78	4	2
Fsa	2.9 ± 0.5		3.1 ± 0.6		3.4 ± 0.6		0.06	7	16
Fth ¹	14.2 ± 3.3		14.8 ± 2.3		13.9 ± 2.3		0.61	4	-2
Hma	4.7 ± 0.5		4.8 ± 0.5		4.5 ± 0.5		0.15	1	-6
Hmo	5.2 ± 0.6		5.1 ± 1.0		5.4 ± 0.4		0.37	-3	4
Mki	1.3 ± 0.2		1.3 ± 0.3		1.3 ± 0.2		0.99	0	-1
Mla	5.4 ± 0.3		5.5 ± 0.5		5.5 ± 0.3		0.76	2	1
Mlu	4.3 ± 0.4		4.3 ± 0.4		4.3 ± 0.3		0.96	-1	-1
Nbu	3.1 ± 0.5		3.1 ± 0.5		3.2 ± 0.5		0.83	1	3
Paf	3.4 ± 0.4		3.3 ± 0.3		3.4 ± 0.4		0.56	-4	-1
Pfu	4.6 ± 0.6		4.6 ± 0.7		4.2 ± 0.4		0.052	0	-9
Sgu	4.3 ± 0.3 a		4.7 ± 0.4 b		4.3 ± 0.6 a		0.018	10	2

¹The diameter measured was of the cutting (see Table S1) and not of the secondary shoot.

Table S3. Stem height at planting in Dec 2017/Jan 2018. Mean and standard deviations as well as P-values from one-way ANOVA are calculated on plot means (n=18) based on 5 individuals, i.e. 90 trees per species and site. All individuals within species were randomly selected from the nursery. Full name of species is given in Table S1. Different letters for sites indicate significant differences (P<0.05).

Species	HE		ME		LE		P-value	ME (% of HE)	LE (% of HE)
	Mean	SD	Mean	SD	Mean	SD			
Afa	40.5 ± 5.6		43.8 ± 5.6		41.8 ± 5.0		0.19	8	3
Agu	5.2 ± 0.5		5.0 ± 0.5		5.0 ± 0.4		0.63	-2	-3
Bbr	54.0 ± 3.9		52.8 ± 2.1		55.1 ± 2.4		0.06	-2	2
Bmi	54.2 ± 4.0		54.0 ± 4.2		54.1 ± 4.1		0.99	0	0
Cgo	16.2 ± 2.1		16.2 ± 2.1		15.4 ± 1.8		0.36	0	-5
Cgr	46.3 ± 4.3	ab	47.8 ± 3.7	a	43.8 ± 5.8	b	0.037	3	-5
Cme	45.4 ± 2.2		45.6 ± 2.8		45.4 ± 3.1		0.96	1	0
Dto	43.3 ± 10.5		43.9 ± 10.3		41.0 ± 10.7		0.67	1	-5
Eex	33.2 ± 3.7		35.0 ± 3.8		35.0 ± 3.7		0.27	5	5
Fsa	26.3 ± 4.5		27.1 ± 5.9		29.3 ± 6.4		0.28	3	11
Fth	51.9 ± 12.2	a	62.0 ± 10.7	b	62.0 ± 10.7	b	0.012	19	19
Hma	53.3 ± 2.9		53.4 ± 2.9		52.1 ± 2.7		0.32	0	-2
Hmo	55.7 ± 4.0		55.4 ± 3.7		56.4 ± 2.9		0.70	-1	1
Mki	7.5 ± 0.8		7.4 ± 1.2		7.7 ± 0.8		0.60	-1	3
Mla	75.9 ± 5.3		76.5 ± 4.4		74.7 ± 4.4		0.50	1	-2
Mlu	22.6 ± 2.9		22.8 ± 2.1		22.7 ± 2.3		0.99	1	0
Nbu	18.6 ± 3.3		18.9 ± 4.5		18.5 ± 3.6		0.95	2	0
Paf	41.4 ± 4.5		38.1 ± 3.6		39.7 ± 3.9		0.057	-8	-4
Pfu	29.6 ± 3.3		30.8 ± 3.0		29.9 ± 2.3		0.41	4	1
Sgu	43.5 ± 3.7		42.3 ± 3.5		43.5 ± 4.3		0.57	-3	0

Table S4. Classification of species into the successional groups (SG) early (ES) and late successional (LS) and the references supporting the classification. For most species, the classification was based on both how it was determined in the literature and observations of the abundance in forests with different degree of disturbances. The country where each study was conducted is given in brackets.

Code	Scientific name	SG	References for successional group
Afa	<i>Afrocarpus falcatus</i>	LS	Hundera et al., 2013a (Ethiopia); Tesfaye et al., 2010 (Ethiopia)
Agu	<i>Albizia gummifera</i>	LS	Chapman & Chapman (Uganda); Eilu and Obua, 2005 (Uganda); Hundera et al., 2013a, b (Ethiopia); Mutiso et al., 2013 (Kenya)
Bbr	<i>Bridelia brideliiifolia</i>	ES	Fisher & Killman, 2008 (Rwanda); African Plant Database, https://africanplantdatabase.ch/
Bmi	<i>Bridelia micrantha</i>	ES	Eilu & Obua, 2005 (Uganda); Fashing et al., 2004 (Kenya); Mutiso et al., 2013 (Kenya)
Cgo	<i>Chrysophyllum gorungosanum</i>	LS	Chapman & Chapman (Uganda); Fisher & Killman, 2008 (Rwanda); Eilu & Obua, 2005 (Uganda)
Cgr	<i>Carapa grandiflora</i>	LS	Fisher & Killman, 2008 (Rwanda); Momo et al., 2016 (Cameroon); Nyirambangutse et al., 2017 (Rwanda)
Cme	<i>Croton megalocarpus</i>	ES	Mutiso et al., 2013 (Kenya); Fashing et al., 2004 (Kenya)
Dto	<i>Dombeya torrida</i>	ES	Eilu & Obua, 2005 (Uganda); Fisher & Killman, 2008 (Rwanda); Tesfaye et al., 2002 (Ethiopia)
Eex	<i>Entandrophragma excelsum</i>	LS	Fisher & Killman, 2008 (Rwanda); Hemp et al., 2017 (Tanzania)
Fsa	<i>Faurea saligna</i>	LS	Eilu & Obua 2005 (Uganda); Fisher & Killman, 2008 (Rwanda); Nyirambangutse et al., 2017 (Rwanda)
Fth	<i>Ficus thonningii</i>	LS ¹	Hundera et al., 2013b (Ethiopia); Kirika et al., 2008 (Kenya)
Hma	<i>Harungana madagascariensis</i>	ES	Mutiso et al., 2013 (Kenya); Fashing et al., 2004 (Kenya)
Hmo	<i>Harungana montana</i>	ES	Fisher & Killman, 2008 (Rwanda); Nyirambangutse et al., 2017 (Rwanda)
Mki	<i>Macaranga kilimandscharica</i>	ES	Eilu & Obua, 2005 (Uganda); Hundera et al., 2013b (Ethiopia); Nyirambangutse et al., 2017 (Rwanda); Ssali et al., 2019, Rutten et al., 2015 (Tanzania)
Mla	<i>Maesa lanceolata</i>	ES	Eilu & Obua 2005 (Uganda); Fashing et al., 2004 (Kenya); Hundera et al., 2013b (Ethiopia); Momo et al., 2016 (Cameroon); Mutiso et al., 2013 (Kenya); Rutten et al., 2015 (Tanzania)
Mlu	<i>Markhamia lutea</i>	ES ²	Chapman & Chapman (Uganda); Mutiso et al 2013; Fashing et al., 2004 (Kenya)
Nbu	<i>Newtonia buchananii</i>	LS	Chapman & Chapman (Uganda); Eilu & Obua, 2005 (Uganda); Fisher & Killman, 2008 (Rwanda)
Paf	<i>Prunus africana</i>	LS	Hundera et al., 2013a, b (Ethiopia); Tesfaye et al., 2010 (Ethiopia)
Pfu	<i>Polyscias fulva</i>	ES	Eilu & Obua, 2005 (Uganda); Fashing et al. 2004 (Kenya); Hundera et al., 2013b (Ethiopia); Mutiso et al., 2013 (Kenya); Nyirambangutse et al., 2017 (Rwanda); Tesfaye et al., 2010 (Ethiopia)
Sgu	<i>Syzygium guineense</i>	LS	Hundera et al., 2013a, b (Ethiopia); Nyirambangutse et al., 2017 (Rwanda); Rutten et al., 2015 (Tanzania); Tesfaye et al., 2010 (Ethiopia); Ssali et al., 2019 (Uganda)

¹Also defined as ES (Hundera et al., 2013b), but here considered as LS species as it mainly occurs in non-disturbed areas, but as a potential epiphyte and strangler the successional strategy is special.

²Mentioned as a possible LS species in Mutiso et al. (2013) and both in disturbed and non-disturbed forests (Chapman & Chapman, 1995) but it has mostly been defined as an ES species.

References to Table S4.

- Chapman, C.A. & Chapman, L.J. (1997) Forest regeneration in logged and unlogged forests of Kibale National Park, Uganda. *Biotropica*, 29, 396-412. <https://doi.org/10.1111/j.1744-7429.1997.tb00035.x>
- Eilu, G. & Obua, J. (2005) Tree condition and natural regeneration in disturbed sites of Bwindi Impenetrable Forest National Park, southwestern Uganda. *Tropical Ecology*, 46, 99-111.
- Fashing, P.J., Forrestel, A., Scully, C. & Cords, M. (2004) Long-term tree population dynamics and their implications for the conservation of the Kakamega Forest, Kenya. *Biodiversity and Conservation* 13, 753-771. <https://doi.org/10.1023/B:BIOC.0000011724.34275.73>
- Fischer E. & Killman D. (2008) Illustrated Field Guide to the Plants of Nyungwe National Park Rwanda. *Koblenz Geographical Colloquia, Series Biogeographical Monographs* 1, 771p.
- Hemp, A., Zimmermann, R., Remmeli, S., Pommer, U., Berauer, B., Hemp, C. & Fischer, M. (2017) Africa's highest mountain harbours Africa's tallest trees. *Biodiversity Conservation*, 26, 103–113. <https://doi.org/10.1007/s10531-016-1226-3>
- Hundera, K., Aerts, R., Fontaine, A., Van Mechelen, M., Gijbels, P., Honnay, O. & Muys, B. (2013a) Effects of coffee management intensity on composition, structure, and regeneration status of Ethiopian moist evergreen Afromontane forests. *Environmental Management*, 51, 801–809. <https://doi.org/10.1007/s00267-012-9976-5>
- Hundera, K., Honnay, O., Aerts, R. & Muys, B. (2013b) The potential of small exclosures in assisting regeneration of coffee shade trees in South-Western Ethiopian coffee forests. *African Journal of Ecology*, 53, 389–397. <https://doi.org/10.1111/aje.12203>
- Kirika, J.M. Bleher, B., Bohning-Gaese, K., Chira, R. & Farwig, N. (2008) Fragmentation and local disturbance of forests reduce frugivore diversity and fruit removal in *Ficus thonningii* trees. *Basic Applied Ecology*, 9, 663-672. <https://doi.org/10.1016/j.baac.2007.07.002>
- Momo, S.M.C., Temgoua, L.F., Ngueguim, J.R. & Nkongmeneck, B-A. (2016) Comparison of plant communities between primary and secondary tropical forests of Mount Oku, Cameroon. *Journal of Ecology and The Natural Environment*, F63DB9364239. <https://doi.org/10.5897/JENE2016.0598>
- Mutiso, F.M., Hitimana, J., Kiyiapi, J.L., Sang, F.K. & Eboh, E. (2013) Recovery of Kakamega tropical rainforest from anthropogenic disturbances. *Journal of Tropical Forest Science* 25, 566–576. <https://www.jstor.org/stable/23616998>
- Nyirambangutse, B., Zibera, E., Uwizeye, F. K., Nsabimana, D., Bizuru, E., Pleijel, H., Uddling, J. & Wallin, G. (2017). Carbon stocks and dynamics at different successional stages in an Afromontane tropical forest. *Biogeosciences*, 14, 1285–1303. <https://doi.org/10.5194/bg-14-1285-2017>
- Rutten, G., Ensslin, R., Hemp, A. & Fischer, M. (2015) Forest structure and composition of previously selectively logged and non-logged montane forests at Mt. Kilimanjaro. *Forest Ecology and Management* 337, 61–66. <http://dx.doi.org/10.1016/j.foreco.2014.10.036>
- Ssali, F., Moe, S.R., Sheil, D. (2019) The differential effects of bracken (*Pteridium aquilinum* (L.) Kuhn) on germination and seedling performance of tree species in the African tropics. *Plant Ecology*, 220, 41-55, <https://doi.org/10.1007/s11258-018-0901-8>
- Tesfaye, G., Teketay, D. & Fetene, M. (2002) Regeneration of fourteen tree species in Harenna forest, southeastern Ethiopia. *Flora*, 197, 461–474. <https://doi.org/10.1078/0367-2530-1210063>
- Tesfaye, G., Teketay, D., Fetene, M. & Beck, E. (2010) Regeneration of seven indigenous tree species in a dry Afromontane forest, southern Ethiopia. *Flora*, 205, 135-143. <https://doi.org/10.1016/j.flora.2008.12.006>

Table S5. P-values and total degrees of freedom (df Tot) of one-way ANOVA for species-specific site effects on tree diameter at base (D_{base}), tree height (H), standardised relative growth rates (RGR) of D_{base} at a D_{base} of 10-25 mm ($D\text{-}RGR_{D10-25}$) and 50-75 mm ($D\text{-}RGR_{D50-75}$), of height at a height of 75-100 cm ($H\text{-}RGR_{H75-100}$) and 250-300 cm ($H\text{-}RGR_{H250-300}$), number of stems per individual (stems#), and tree mortality. Df for site was always 2. The analysis was based on plot averages of each species (maximum 18 plots, 20 species and 3 sites). Species that did not meet the criterion for sufficient number of plot replicates (i.e. at least one individual on ≥ 12 plots per site) were not included in the analysis, indicated by NA (not applicable) in the table. ND, no dead individuals and thus no variance analysis could be conducted.

Species	D_{base}			Height			$D\text{-}RGR_{D10-25}$			$D\text{-}RGR_{D50-75}$			$H\text{-}RGR_{H75-100}$			$H\text{-}RGR_{H250-300}$			Stems#			Mortality		
	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot	P-value	df	Tot
Bbr	<0.001	54	<0.001	54	<0.001	54	0.001	54	<0.001	54	<0.001	53	<0.001	54	0.26	54	ND	54	ND	54	ND	54	ND	54
Hmo	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	0.68	54	0.50	54	0.36	54	0.36	54	0.36	54	0.36	54	0.36	54
Mki	<0.001	54	0.008	54	0.090	54	0.064	51	0.17	47	0.35	53	0.32	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54
Mla	<0.001	54	0.22	54	<0.001	54	<0.001	52	0.94	54	0.70	54	0.004	54	0.057	54	0.057	54	0.057	54	0.057	54	0.057	54
Pfu	0.001	54	<0.001	54	0.014	54	0.018	54	<0.001	54	0.003	53	0.32	54	0.13	54	0.13	54	0.13	54	0.13	54	0.13	54
Agu	<0.001	54	<0.001	54	0.20	50	NA	12	NA	40	NA	24	0.93	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54
Bmi	<0.001	54	<0.001	54	<0.001	54	0.005	53	<0.001	53	0.033	54	0.009	54	0.172	54	0.172	54	0.172	54	0.172	54	0.172	54
Cme	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	0.25	54	0.012	54	0.012	54	0.012	54	0.012	54
Dto	<0.001	54	<0.001	54	<0.001	54	0.024	53	<0.001	53	0.002	54	<0.001	54	ND	54	ND	54	ND	54	ND	54	ND	54
Hma	<0.001	54	<0.001	54	<0.001	54	0.005	54	<0.001	54	0.54	54	0.89	54	0.35	54	0.35	54	0.35	54	0.35	54	0.35	54
Mlu	<0.001	54	<0.001	54	<0.001	54	NA	43	<0.001	44	0.003	47	0.004	54	ND	54	ND	54	ND	54	ND	54	ND	54
Afa	<0.001	54	<0.001	54	0.14	54	0.49	46	0.23	54	0.008	54	0.45	54	0.157	54	0.157	54	0.157	54	0.157	54	0.157	54
Cgr	<0.001	51	<0.001	51	<0.001	50	NA	9	NA	45	NA	12	0.33	54	<0.001	51	<0.001	51	<0.001	51	<0.001	51	<0.001	51
Fsa	<0.001	54	<0.001	54	<0.001	54	0.029	43	<0.001	54	NA	43	0.010	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54
Paf	<0.001	54	<0.001	54	<0.001	54	NA	32	<0.001	53	NA	39	0.002	54	0.013	54	0.013	54	0.013	54	0.013	54	0.013	54
Sgu	0.012	54	0.084	54	0.32	54	0.014	50	0.16	54	0.19	54	0.035	54	0.58	54	0.58	54	0.58	54	0.58	54	0.58	54
Cgo	0.001	54	0.003	54	<0.001	49	NA	0	NA	38	NA	2	0.71	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54
Eex	0.001	54	0.11	54	0.006	54	NA	24	0.13	41	NA	0	0.19	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54
Fth	<0.001	54	<0.001	54	0.002	54	<0.001	52	0.50	52	<0.001	53	0.001	54	0.81	54	0.81	54	0.81	54	0.81	54	0.81	54
Nbu	0.002	54	0.15	54	NA	41	NA	1	0.088	44	NA	15	1.00	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54	<0.001	54

Table S6. Number of individuals recorded dead at each tri-monthly census of each species and site. The graded colour scale from light red to bright red indicate an increasing number of dead individuals. Newly dead trees recorded at each census in percent of remaining trees are presented graphically below the table. Grey marked dates in the table head indicate first census after the annual dry period. However, all trees were irrigated during the entire dry period in 2018, but not in 2019. During the first census after planting (Mar-18) only six trees were recorded dead, but five of them were replaced by new individuals from the nursery and was not included in the mortality data. #, number of dead trees. Full species names are given in Table 2.

Species	High elevation (Sigira)								Mid elevation (Rubona)								Low elevation (Makera)								Species							
	Dec-17				Mar-18				Jul-18				Oct-18				Feb-19				Jun-19				Sep-19							
	1	2	3	4	5	6	7	8	#	%	1	2	3	4	5	6	7	8	#	%	1	2	3	4	5	6	7	8	#	%		
Afa									0	0							1	1	1	1	3	3.3					1	1		1	1	Afa
Agu					1	1	4	5	11	12							2	2	2	2	10	11					11	13	2	1	1	Agu
Bbr						0	0														0	0								0	0	Bbr
Bmi						0	0										1	1			3	3.3								0	0	Bmi
Cgo					2	1	1	1	4	4.4							2	2	5	5	20	22					13	20	11	4	3	Cgo
Cgr					1				1	1.1							1	1	6	2	14	16					14	16	19	11	3	Cgr
Cme						0	0													0	0					1	1	2		4	Cme	
Dto						0	0													0	0								0	0	Dto	
Eex						0	0										1	1	2	1	6	6.7					7	9	4	2	1	Eex
Fsa						1			1	1.1							1				1	1.1					1	9	4	2	2	Fsa
Fth						1			1	2.2							1				2	2.2					1			1	1	Fth
Hma						1	1		2	2.2							1				0	0					1		1	2	2	Hma
Hmo						1	1		2	2.2							1				0	0								1	1	Hmo
Mki						1	4		7	7.8							2		1	1	4	4.4					11	4	8	7	4	Mki
Mla						1			1	1.1							1				0	0								4	1	Mla
Mpl							0	0												0	0								0	0	Mpl	
Nbu					4	2	5	4	21	23							7	6	10	7	25	28					6	21	18	9	3	Nbu
Paf							1		1	1.1							1	7	1	9	10									0	0	Paf
Pfu							0	0											0	0									1	2	2	Pfu
Sgu							1	1.1											2	2.2					1	1	1	1	3	Sgu		
#	0	0	0	6	7	12	12	17	54	3	0	0	9	15	33	12	19	11	99	6	0	1	16	84	84	59	35	17	296	18	#	
%	0	0	0	11	13	22	22	31			0	0	9	15	33	12	19	11			0	0	5	28	28	20	12	6		% of all dead trees		
%	0.0	0.0	0.0	0.3	0.4	0.7	0.7	1.0			0.0	0.0	0.5	0.8	1.9	0.7	1.1	0.6			0.0	0.0	0.9	4.7	4.9	3.7	2.2	1.1		% of remaining trees		






Figure S1. Site map and experimental design at the high elevation site (Sigira)




Figure S2. Site map and experimental design at the mid elevation site (Rubona)






Figure S3. Site map and experimental design at the low elevation site (Makera)








Figure S4. The relation between height and base diameter and the relative growth rate (RGR) for height (H-RGR) and base diameter (D-RGR) in relation to height and base diameter, respectively, for all 20 species measured tri-monthly over two years at all three sites. The grey marked height and base diameter intervals are used to calculate the standardised H-RGR and D-RGR, respectively. Negative height RGR of small trees are mainly due to soil erosion effects between the census while negative height and diameter RGR on larger trees mainly are due to broken stems causing shorter trees and/or for multistem trees changes of stem used for the measurements.




Figure S5 a-t. The development of the stem base diameter measured every third month over two years and the relative growth rate (RGR) between measurement intervals for 20 species grown at three sites along an elevation gradient (High, blue square/bar; Mid, orange circle/bar; Low, red triangle/bar). Each marker and bar show site level mean, including standard errors for bars. Full species names are given in Table 2.




Figure S6 a-t. The development of the stem height measured every third month over two years and the relative growth rate (RGR) between measurement intervals for 20 species grown at 3 sites along an elevation gradient (High, blue square/bar; Mid, orange circle/bar; Low, red triangle/bar). Each marker and bar show site level mean, including standard errors for bars. Full species names are given in Table 2.