

1 **Manuscript type:** *Ideas and perspective*

2

3 **Title:**

4 Ideas and perspectives. The fluctuating nature of oxygen shapes the ecology of aquatic habitats and their  
5 biogeochemical cycles: the aquatic oxyscape

6

7 **Authors**

8 Marco Fusi<sup>1,\*</sup>, Sylvain Rigaud<sup>2</sup>, Giovanna Guadagnini<sup>3</sup>, Alberto Barausse<sup>3</sup>, Ramona Marasco<sup>4</sup>, Daniele  
9 Daffonchio<sup>4</sup>, Julie Régis<sup>2</sup>, Louison Huchet<sup>2</sup>, Capucine Camin<sup>2</sup>, Laura Pettit<sup>1</sup>, Cristina Vina-Herbon<sup>1</sup>, Folco  
10 Giomi<sup>5,\*</sup>

11

12 <sup>1</sup>Joint Nature Conservation Committee, Monkstone House, City Road, Peterborough, PE1 1JY

13 <sup>2</sup>Univ. Nîmes, EA 7352 CHROME, Rue Du Dr Georges Salan, 30021 Nîmes, France

14 <sup>3</sup>Department of Biology, University of Padova, Padova, Italy

15 <sup>4</sup>Biological and Environmental Sciences and Engineering Division (BESE), Red Sea Research Center  
16 (RSRC), King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi  
17 Arabia

18 <sup>5</sup>Independent Researcher, Padua, Italy

19

20 \*equal contribution and co-corresponding authors:

21 Marco Fusi: marco.fusi@jncc.gov.uk

22 Folco Giomi: folcog@gmail.com

23

24 **Abstract**

25 Oxygen availability is a pivotal factor for ecosystem functioning and the resistance of organisms to the  
26 effect of climate change in aquatic habitats. Although extensive work has been made to assess the effect of  
27 oxygen on marine and freshwater biota, many studies did not capture the ecological importance of oxygen  
28 variations. Overlooking the fluctuating nature of oxygen may cause potential biases in the design and  
29 implementation of management policies of aquatic habitats. Conceptual perspectives on the dynamic nature  
30 of oxygen fluctuations have been raised in the scientific community to enhance the understanding of the  
31 effect of oxygen on the physiology and the ecology of aquatic species and the biogeochemical functioning  
32 of the ecosystems. A growing number of empirical works are outlining a novel conceptual framework that  
33 considers the magnitude of oxygen fluctuation as a key variable that explains adaptation to stress conditions.  
34 Oxygen in productive aquatic habitats shows large fluctuations at diel scale, exposing aquatic species from  
35 conditions of extreme supersaturation to anoxia. Recent research indicates that such fluctuation tunes the  
36 physiological plasticity of the animal in response to thermal stresses. In this contribution, we provide  
37 compelling evidence based on current research that the fluctuating oxygen landscape, here defined as  
38 “*oxyscape*”, has an important role in aquatic animal physiology and adaptation and the ecosystem  
39 biogeochemistry. We propose that the oxyscape should be considered in the modelling and managing  
40 policies of aquatic ecosystems.

41

42 **Keyword**

43 Biogeochemical cycles; Climate Change; Hyperoxia; Hypoxia; Marine Ecosystem Assessment; Microbial  
44 processes; Oxygen Fluctuations; Supersaturation

45

## 46 **1. Introduction to environmental dissolved oxygen variability and its ecological relevance**

47 The actual assembly of organism communities in a given habitat is determined by the specific environmental  
48 conditions that select specific sets of species. Climate anomalies and changes raise concerns on ecosystem  
49 stability and habitat preservation, with consequences ranging from species substitution to species extinction  
50 (Antão et al., 2020). To model and predict such effects on community composition and stability, it is  
51 necessary to learn about the limits of the environmental conditions that challenge the different species  
52 (Bennett et al., 2021). However, in many cases, such conditions are resumed into averaged parameters that  
53 do not capture the whole range to which organisms are exposed to (Bernhardt et al., 2020).

54 In aquatic environments, chemical and physical parameters greatly vary at different time and spatial scales,  
55 following cyclical fluctuation or stochastic variations (Shaw et al., 2013; Kerrison et al., 2011; Vargas et al.,  
56 2017, 2022; Manzello, 2010; Challener et al., 2016; Duarte et al., 2013). The lack of details on such variations  
57 in time and space challenges our understanding of how the species adapt their physiology, limiting our  
58 estimation of their resilience to ongoing environmental changes (Booth et al., 2023a). The perception of  
59 such knowledge gap is progressively moving the research interest on species' eco-physiological response to  
60 chemical and physical parameters, such as temperature, salinity or pH, and the associated biogeochemical  
61 cycles, toward more accurate assessments of their natural variability (Antão et al., 2020; Bernhardt et al.,  
62 2020). Increasing efforts are now devoted to upgrade eco-physiological approaches to integrate the relevant  
63 environmental background (Giomi et al., 2019; Marasco et al., 2023; Bitter et al., 2018, 2021; Pörtner et al.,  
64 2023).

65 Oxygen variability exerts a deep effect on aquatic life yet remains underestimated. Most studies dealing with  
66 oxygen variation have addressed the decline of mean oxygen availability in oceans (Breitburg et al., 2018)  
67 and the occurrence and repercussions of events of environmental hypoxia (Diaz and Rosenberg, 2008;  
68 Bickler and Buck, 2007). How short-term (i.e., daily) oxygen variations shape aquatic life and  
69 biogeochemical processes remains largely overlooked. These variations are generally of greater amplitudes  
70 and more regular than seasonal ones (Limburg et al., 2020; Bates et al., 2018). We refer to deviation from  
71 the saturation of oxygen concentrations as the ratio between the actual concentration and the saturation  
72 concentration of oxygen that would be observed in water under equilibrium conditions between air and  
73 water. The fine measure of the range of variation of aquatic oxygen and its effect on organisms and

74 biogeochemical cycles has the potential to elucidate unpredicted mechanisms of resilience and tolerance to  
75 ongoing climate change.

76

## 77 **2. Oxygen trends and stochastic variations**

78 A gradual decline of the total oxygen content of about 2% since the second half of the 20<sup>th</sup> century has  
79 been recently reported, with a further reduction of up to 7% predicted for the following century (Schmidtke  
80 et al., 2017; Breitburg et al., 2018). In parallel, open-ocean oxygen minimum zones (OMZs) have largely  
81 expanded (Stramma et al., 2010) together with the increasing occurrence of severe hypoxia events whose  
82 predictability has decreased as their occurrence spreads across all oceans (Diaz and Rosenberg, 2008). The  
83 main cause of deoxygenation is attributed to global warming and is exacerbated by the discharge of nutrients  
84 and pollutants (Ito et al., 2016). Topography of water basins also affects the incidence of hypoxia events  
85 influencing the rate of mixing of oxygenated layers and the resident time of water bodies (Breitburg and  
86 Grégoire, 2018; Laffoley and Baxter, 2019). For example, in the Baltic Sea, the geomorphology of the  
87 coastline can explain up to 80% of the hypoxic phenomena (Virtanen et al., 2019).

88 The gradual decrease of total oxygen in oceans has been reported in several studies that analyse long time  
89 trends (Keeling et al., 2010; Schmidtke et al., 2017; Breitburg et al., 2018) and from reports at the global  
90 geographical scale (Laffoley and Baxter, 2016). However, unexpected oxygen profiles have also been  
91 reported in several specific habitats and at different time scales suggesting that different abiotic and biotic  
92 drivers modulate oxygen dynamics (Craig and Hayward, 1983). For instance, the alternation of oxygen  
93 biological production and consumption determines massive fluctuation of its availability in highly  
94 productive aquatic habitats, such as mangrove forests, salt marshes, coral reefs (Giomi et al., 2019; Fusi et  
95 al., 2021; Booth et al., 2021), kelp forests (Krause-Jensen et al., 2016), plankton blooms in the open oceans  
96 (Riser and Johnson, 2008; Benoiston et al., 2017), freshwater lakes (Andersen et al., 2017) or even in  
97 agricultural drainage channels (Booth et al., 2023). The seasonal variation of diel oxygen fluctuations is  
98 increasingly reported at all latitudes and habitats and occur in a vast range of water bodies independently  
99 of the scale, the hydrological features, and the local biotic components (Figure 1). For example, fluctuations  
100 of 150% in spring and 30% in winter occur in the Venice lagoon (Figure 1, Location 5), and even more  
101 pronounced fluctuations occur in mangroves, corals and seagrasses (Figure 1, Locations 10-12; Giomi et

102 al., 2019). The magnitude of oxygen fluctuations is site-specific and depends, among other factors, on the  
103 solar radiation and water temperature and their effect on primary producers' photosynthetic and respiration  
104 activity, the cycles of water column stratification and mixing, and the nutrient loads. In highly productive  
105 marine environments, increased dissolved oxygen concentration may occur in spring due to the  
106 intensification of photosynthetic activity, followed by a decrease of available oxygen during the warmest  
107 months and a progressive recovery in autumn and winter (Cowan et al., 1996; Bartoli et al., 2001; Kim et  
108 al., 2019). Spatial differences in oxygen saturation occur between surface and bottom water masses because  
109 of isolation driven by water stratification, the decrease of light penetration, and the coupling with increased  
110 oxygen demand close to the sediments at the bottom of the water column (Figure 2). In summer, enhanced  
111 benthic respiration is determined by higher water temperature, which decreases oxygen solubility and  
112 enhances oxygen demand. Lower vertical mixing, higher water residence time, and higher turbidity due to  
113 higher concentration of organic matter, further decrease oxygen concentrations down to hypoxia in the  
114 deep water layers (Figure 2; Talke et al., 2009; Schmidt et al., 2019). Conversely, dissolved oxygen increases  
115 in the euphotic part of the water column because of the enrichment of photosynthetic communities (Spietz  
116 et al., 2015).

117

### 118 **3. The oxygen variation in the day-life of aquatic organisms**

119 Marked oxygen variations occur at a diel scale, driven by temperature and light diel cycles and by the balance  
120 between photosynthesis and respiration (Winter et al., 2019). The amplitude of diel oxygen variation can  
121 exceed the average seasonal variation, thus being variable with higher explanatory power for the physiology  
122 of aquatic life (Giomi et al., 2019). While water temperature is only linked to meteorological changes, the  
123 oxygen concentration in productive aquatic environments is also regulated by the biota component  
124 (Chapman, 2021). Community composition of primary producers (Power and Cardinale, 2009), rainfall  
125 level (Mallin et al., 1993), nutrients' runoff (Kinney and Roman, 1998), biotic networks (Graham et al.,  
126 2018; Breen and Mann, 1976), and bacterial community dynamics (Guo et al., 2022), govern, at a small  
127 scale, high spatial and temporal change in the concentrations of dissolved oxygen. The interaction of all  
128 these factors strongly affects dissolved oxygen fluctuations beyond the levels that can be predicted from  
129 just the meteorological trends.

130 Oxygen availability plays an important role in the physiology of aquatic species (Pörtner, 2010; Hochachka,  
131 1991; Stillman and Somero, 2000). In the design of experiments on the physiology of aquatic animals,  
132 oxygen is frequently treated as a fixed factor and maintained around the water saturation. However,  
133 organisms inhabiting aquatic environments experience ample diel oxygen fluctuations and are adapted to  
134 endure or respond to the experienced variation (Blewett et al., 2022; Morash et al., 2018). Oxygen may be  
135 systematically depleted at night and fully recovered during the day, changes that certainly have consequences  
136 on the tolerance to acute stressors such as hypoxia and warming. Dependent on the intensity of the  
137 photosynthetic activity, dissolved oxygen concentrations can reach levels far above the physical saturation  
138 (Giomi et al., 2019; McArley et al., 2020). Under such hyperoxic conditions, aquatic organisms experience  
139 oxygen availability well beyond their physiological needs, but the effects on their physiology and life history  
140 are largely unknown.

141 Hyperoxia benefits the metabolic performance of fishes and invertebrates, especially under acute warming  
142 that may be very intense in tidal ponds or shallow coastal habitats (Fusi et al., 2021; Booth et al., 2021;  
143 Giomi et al., 2019; McArley et al., 2020; Booth et al., 2023a). It extends the upper thermal tolerance of  
144 aquatic animals during acute events of daily warming and their survival to anomalous heat waves (McArley  
145 et al., 2022). Within the current trend of climate change, the beneficial effect of hyperoxia against extreme  
146 heating events can have underestimated positive consequences on the survival of the animals. For instance,  
147 an important consequence of exposure to the hyperoxic conditions experienced during the daytime under  
148 high heating regimes is the enhanced capability to extract oxygen from the poorly oxygenated water during  
149 the night and endure hypoxic or even anoxic periods (Giomi et al., 2019; Booth et al., 2021).

150 Oxygen fluctuations are also appreciated as a factor contributing to the structuring of organismal networks  
151 and in the synchronization of life history dynamics such as spawning or larval recruitment (Garzke et al.,  
152 2019; Viaroli and Christian, 2004).

153

#### 154 **4. The role of oxygen fluctuation in coastal biogeochemical cycles**

##### 155 ***4.1 Biogeochemical cycles in the pelagic compartment in relation to oxygen fluctuation at daily scale***

156 In productive coastal ecosystems the biogeochemical cycles of key chemical elements are strongly  
157 interconnected with oxygen dynamics. Nitrogen and phosphorus are typically key elements which limit

158 primary production in shallow coastal-marine ecosystems, whose emissions from river basins (or other  
159 sources, such as previously enriched sediments) can lead to eutrophication with detrimental effects on  
160 biodiversity and human welfare (Palmeri et al., 2013). The biogeochemical cycles of nitrogen and  
161 phosphorus are composed of multiple physical, chemical and biological processes, of which some are  
162 accelerated by oxygen availability, such as nitrification and mineralization, while some others are enhanced  
163 by its absence, such as denitrification and P release from sediments. The regular daily alternation of  
164 contrasting conditions of below- and above-saturation dissolved oxygen concentrations in the water  
165 column, particularly in the euphotic zone, promotes changes in the relative abundance of different nutrient  
166 forms (Figure 3). This becomes clear when looking, for example, at the nitrogen cycle in the water column  
167 and its complex relationship with oxygen availability: the presence of dissolved oxygen speeds up the  
168 mineralization of organic nitrogen to ammonium as well as nitrification, i.e., the microbial oxidation of  
169 ammonium into nitrites and then nitrates. The absence of oxygen makes denitrification possible, i.e., the  
170 reduction of nitrates into inert gaseous nitrogen is promoted by facultative aerobic bacteria which, in the  
171 absence of anoxia, will prefer oxygen to nitrates as an electron acceptor (Palmeri et al., 2013). Ammonium,  
172 nitrites and nitrates are important nitrogen forms which are bioavailable to primary producers, and  
173 oscillating oxygen conditions in the water column at the diel scales can promote their removal via the  
174 nitrification-denitrification chain. High oxygen concentrations are also associated with high photosynthetic  
175 activity (e.g., planktonic) and, therefore, with the fast removal of dissolved inorganic nutrients from water  
176 through uptake by primary producers (Caron, 1994). The picture is made more complex by the typically  
177 non-linear dependence of biogeochemical reactions on oxygen concentration (e.g., nitrification and  
178 denitrification) and on temperature (in the case of most biogeochemical processes), which can both change  
179 strongly throughout the day, especially in shallow productive ecosystems, altering reaction rates at sub-daily  
180 scales in a way which cannot be appreciated if only the daily mean in dissolved oxygen is considered, but  
181 which needs to be quantified if we are to mechanistically predict future biogeochemical cycling under  
182 climate change (Caballero-Alfonso et al., 2015). A further source of complexity is given by the feedbacks  
183 between abiotic and biotic components of pelagic ecosystems. The growth of primary producers, both  
184 microscopic and macroscopic, is on the one hand affected by the presence of bioavailable nutrient forms,  
185 and on the other hand their biomass can reduce light availability by increasing shading or turbidity, limiting

186 light penetration and, so, oxygen production and warming in deeper waters. These examples highlight the  
187 importance of higher frequency measurements of oxygen fluctuations, nutrient concentrations, and  
188 microbial community abundance, composition and activity in the water column at sub-daily scales, to get a  
189 better, more quantitative grasp of biogeochemical cycling in coastal water bodies (Meire et al., 2013). A  
190 general framework to assess the short-term (hourly) oxygen fluctuations in relation to the movement and  
191 related physical properties of water masses, i.e., of processes such as stratification, residence time,  
192 thermohaline flows, in addition to other processes (photosynthesis, respiration, solubility changes, etc.) is  
193 important to future oxygen modelling.

194 Of course, the intertwined oxygen and nutrient dynamics in the water column cannot be understood, at  
195 least in relatively shallow aquatic ecosystems, without looking also at the interactions between the pelagic  
196 and the benthic compartment. This takes place through vertical transport of matter (settling, resuspension,  
197 convection, upwelling currents, etc.) and is also affected by stratification, gradient-driven diffusive fluxes,  
198 and organism movements and vertical migrations.

199

#### 200 ***4.2 Biogeochemical cycles in the benthic compartment in relation to oxygen fluctuation at daily scale***

201 When temperature and oxygen concentration in the water column fluctuate at short daily time scale, yet  
202 with very large amplitude (from anoxia/hypoxia to hyperoxia), the impact on biogeochemical cycles and  
203 chemical transfers is mostly unknown. Based on the trends observed at seasonal “steady-state” conditions,  
204 a conceptual model on the location of the major benthic processes in the sediment and the resulting fluxes  
205 at the sediment-water interface within the diurnal temporal scale can be drawn (Figure 4).

206 As oxygen penetration depth and concentration in sediment is known to rapidly evolve in response to the  
207 oxygen concentration in the overlying water (Glud, 2008), the importance of aerobic process in surface  
208 sediment is also expected to fluctuate at the diurnal timescale, with higher contribution during the day than  
209 during the night. The fast kinetics of the reductive dissolution/oxidative precipitation of Mn and Fe oxy-  
210 hydroxides and the redox cycle of S chemical species (i.e.,  $\text{H}_2\text{S}/\text{SO}_4^{2-}$ ) suggests that those chemical species  
211 should evolve with a similar trend over daily timescales, with a reasonable short term (minutes-hours) delay  
212 (Rigaud et al., 2018). Consequently, we expect a vertical fluctuation of those biogeochemical redox  
213 processes in the sediment and related fluxes under diel oxygen fluctuation (Figure 4). The low oxygen



214 concentration at night should induce a lower oxygen penetration depth and an upward distribution of  $\text{NO}_3^-$   
215 , Mn and Fe oxyhydroxides and  $\text{SO}_4^{2-}$  reductions. If the oxygen concentration is low enough, the reductive  
216 dissolution of Mn and Fe oxyhydroxides may directly occur at the sediment-water interface, inducing their  
217 release to the water column with other chemical species associated with these phases (i.e.,  $\text{PO}_4^{3-}$  and most  
218 trace elements). In the specific case of anoxia, the sulfate-reduction process may occur at the sediment  
219 surface, releasing  $\text{H}_2\text{S}$  to the water column. It is expected that in such a condition, the release of dissolved  
220 metals (Fe, Mn and trace elements), which also present a rapid kinetic for metal sulfide formation in the  
221 presence of S(-II), can be reduced and likely reverted (Figure 4). In contrast, the increase of oxygen  
222 concentration during the day promotes oxygen penetration in the sediment and the oxidation of reduced  
223 species accumulated during the night, such as dissolved Mn, Fe,  $\text{NH}_4^+$  and eventually  $\text{H}_2\text{S}$ , preventing their  
224 release from the sediment. The reconstitution of the Mn/Fe oxyhydroxide reservoir in surface sediment  
225 favours the trapping of  $\text{PO}_4^{3-}$  and trace elements associated with Fe and Mn cycles. The oxidation of  $\text{NH}_4^+$   
226 induces the formation of  $\text{NO}_2^-$  and  $\text{NO}_3^-$  that may be released from the sediment.

227 In shallow coastal areas, the light radiation may also reach the sediment surface allowing the photosynthetic  
228 activity of the microphytobenthos to occur in surface sediment (Figure 4). The local source of oxygen at  
229 the sediment surface occurs because photosynthesis strongly enhances the biogeochemical processes  
230 described above (Denis et al., 2012; Rigaud et al., 2018). The oxygen released in the water column enhances  
231 the water hyperoxia and consumes nutrients and  $\text{CO}_2$  during photosynthesis. This creates a very peculiar  
232 situation that modifies the direction and intensity of chemical fluxes at the sediment-water interface and  
233 the chemical composition of water and surface sediment in coastal areas at the diurnal scale. For those rapid  
234 redox-sensitive chemical species, processes and fluxes are thus expected to evolve, between such extreme  
235 night/day fluctuations, with a transient response related to the reaction kinetics and transport within the  
236 sediment surface.

237 For OM mineralization processes, the effects of short-term oxygen fluctuation are more challenging to  
238 predict. It will also be dependent on the capacity of heterotrophs to react/adapt to the changing oxygen  
239 concentrations and eventually shift between different metabolic processes (for microorganisms where  
240 variables metabolic pathways coexist) or to activate/cease the process in relation to physicochemical  
241 conditions and oxidant availability (for microorganisms with specialized metabolisms). Consequently, the

242 dynamic response of the microbial community to short-term oxygen and chemical fluctuations needs to be  
243 investigated in more detail in relation to microbial communities' resilience and adaptation capacity (see  
244 section 5).

245 In productive coastal areas, the benthic organisms, including macro and meiofaunal species, adapt to large  
246 and rapid daily oxygen variation by modifying their behaviour. The bioturbating activity may be favoured  
247 during high oxygen concentration and lowered during conditions of low oxygen availability. We expect that,  
248 during the night oxygen deficiency, the benthic organism activity is reduced, inducing a decrease in the  
249 sediment reworking intensity. This is accompanied by a reduction of the biologically-mediated solute fluxes  
250 at the sediment-water interface and oxygen penetration depth in the sediment. In contrast, during oxygen  
251 supersaturation occurring in the day, bioturbation is intensified and may induce a short-term change in the  
252 contribution of biologically-mediated reactions and transfers. Here also, a delay between the oxygen  
253 fluctuation in the water column and the biogeochemical response of the sediment, related to the biological  
254 activity, would depend on site characteristics (oxygen concentration and range of fluctuation, bioturbating  
255 species). For instance, in areas where anoxia events are recurrent, the absence of macrofauna prevents  
256 bioturbation and its influence on the benthic biogeochemistry (Nilsson and Rosenberg, 1997).

257 In response to large amplitude and diel oxygen fluctuations, the sediment' biogeochemical cycles are also  
258 expected to fluctuate (Figure 4). However, as the resulting benthic biogeochemistry cycles are associated  
259 with the dynamics of three interdependent compartments (i.e., geochemical composition, microbial  
260 communities, bioturbating-faunal communities), each presenting its kinetics and responses to oxygen  
261 fluctuations, we expect the sediment to be permanently maintained under unsteady-state conditions. This  
262 makes it difficult to quantitatively predict the resulting OM mineralization rates, the proportion of involved  
263 processes, and the chemical transfers at the sediment-water interface. The recent development of tools  
264 which can measure the chemical composition and sediment-water fluxes over the short-term scale (i.e.,  
265 microsensors, eddy covariance technic, gradient-flux method), in addition to benthic microbial and  
266 macrofaunal activity (e.g., improved sediment profile imaging), may be beneficial to produce experimental  
267 data that will help to fill these gaps in understanding. At the same time, specific efforts should be focused  
268 on assessing the isolated response to those three different compartments. Those data should then be  
269 implemented to appropriately calibrate non-steady-state coupled pelagic-benthic biogeochemical models,

270 which can predict the resulting biogeochemical functioning of productive coastal ecosystems and, thus, to  
271 predict their fate under climatic changes.

272

## 273 **5. The interrelationship of oxygen fluctuation and aquatic microbial communities**

274 Oxygen is a two-faced element that acts as the terminal electron acceptor in aerobic respiration (by far the  
275 most efficient energy metabolism) and as an element of toxicity because the reduction of O<sub>2</sub> molecules  
276 partly results in reactive oxygen species. For this reason, oxygen remains a strong evolutionary force  
277 dominating functional interactions and the spatial structure of many microbial communities (Fenchel and  
278 Finlay, 2008). Oxygen availability determines microbial metabolism, and many studies have been carried  
279 out to elucidate the different kinds of communities and metabolisms occurring under anaerobic or aerobic  
280 conditions (Sandrin et al., 2009). Marine microorganisms in productive coastal environments also  
281 experience large oxygen diel fluctuations, especially at the boundary layer between sediment/primary  
282 producers and water (Pacherres et al., 2022).

283 Assessing the full environmental oxygen variability in aquatic habitats using high-resolution temporal and  
284 spatial scales relevant to microorganisms can reveal in detail the complexity of the patterns of community  
285 dynamics and diversity in such oxygen-variable environments (Berg et al., 2022). For example, oxygen  
286 largely varies during the day in tropical clear shallow waters because of the photosynthetic activity of  
287 seagrasses, corals and mangroves (Giomi et al., 2019). Therefore, the microbial communities in these  
288 ecosystems can experience environmental conditions ranging from hyperoxia to anoxia (see the oxygen  
289 profile in Figure 2). These fluctuations imply that the microbial communities have to shift their composition  
290 and function according to the oxygen availability (Fenchel and Finlay, 2008; Booth et al., 2023b, 2019).  
291 Microorganisms generally have a short generation time (and high turnover) that favour timely adaptation  
292 to new conditions in a changing environment (Steiner et al., 2019). During low oxygen availability at night-  
293 time, bacterial communities will shift their composition and function toward anaerobic function (Figure 5).  
294 End products of anaerobic metabolisms such as ammonium, sulphur and methane are consumed when  
295 oxygen becomes available again during the daylight and the recovery of photosynthesis, re-establishing  
296 aerobic respiration as the central metabolism of the microbial communities (Fenchel and Finlay, 2008). For  
297 example, in the Bohai Sea (China), under oxygen concentration lower than 4.2 mg/L the microbial

298 communities were dominated by bacteria of the *Anaerolineaceae* (Guo et al., 2022). In Chilean coast  
299 ecosystems, the composition of the bacterial communities is regulated by the seasonality of the upwelling  
300 waters, with the partitioning of community composition driven by dissolved oxygen. *Bacteroidetes*, SAR11,  
301 SAR86 and *Alphaproteobacteria* dominated in waters containing dissolved oxygen concentrations higher than  
302 70  $\mu\text{M}$ . In contrast, taxonomic groups such as Arctic96BD-19, SUP05, SAR324 and *Desulfobacterales* were  
303 observed at dissolved oxygen concentrations below 70  $\mu\text{M}$  to undetectable levels (Aldunate et al., 2018).  
304 The continuous variation of oxygen concentration enhances the diel cycling of nutrients in productive  
305 coastal environments, highlighting the importance of assessing the ecologically relevant oxygen fluctuations  
306 to determine the effective microbial functionality of the marine coastal ecosystem (Trowbridge et al., 2017).  
307 Incorporating methods to assess the ecologically relevant variability of oxygen availability, intensity and  
308 synchronicity in future experimental designs will generate a more accurate prediction of the response of  
309 bacterial communities and therefore their potential for biogeochemical cycling to climate change, better  
310 informing the development of management strategies to mitigate detrimental stressor impact on ecosystems  
311 effectively. The current challenge in microbial ecology is to understand the widespread temporal and spatial  
312 environmental oxygen variability and switch from the “mean conditions paradigm” often used for  
313 investigations of microbial ecological processes to a dynamic model able to capture the microbial changing  
314 and functional potential under oxygen fluctuating conditions (Fusi et al., 2022).

315

## 316 **6. Implications for marine assessment and management**

317 Oxygen production is considered an ecosystem service (<https://cices.eu/>) because of the benefits humans  
318 receive, but, to date, there is limited inclusion of the impacts of deoxygenation or oxygen fluctuation in  
319 marine environmental policies. There are only few areas whose oxygen dynamics has been extensively  
320 monitored like in North America (i.e., Chesapeake bay, Gulf of Mexico, Long Island Sound) and included  
321 in management plans for environmental protections (e.g.,  
322 <https://coastalscience.noaa.gov/project/operational-gulf-of-mexico-hypoxia-monitoring/>). However,  
323 globally, there has been a lack of oxygen dynamics’ integration into the computation of biodiversity  
324 indicators that assess aquatic communities (Breitburg et al., 2019; Chen et al., 2022), and in particular the  
325 oxygen dynamic at a daily scale remains largely neglected is marine assessment and management. Policies

326 on pollution or nutrient control had a positive effect on the oxygen level in ocean water, however, there is  
327 still a clear missing link on specific policies to monitor and manage daily oxygen variability. While attention  
328 has been given to ocean warming and acidification, daily oxygen dynamics have been overlooked in  
329 assessing marine habitat or species' sensitivity to climate changes and anthropogenic disturbances.  
330 Including ecologically relevant variations of oxygen availability into aquatic biodiversity indicators, in  
331 addition to continuous and high temporal resolution oxygen and chemical concentration measurements in  
332 the sediment and water column, offers a great opportunity to refine and produce more robust predictors  
333 that will be able to disentangle better the response of aquatic ecosystems to climate change and  
334 anthropogenic disturbances (Dafforn et al., 2012).

335 Current hypoxic thresholds for aquatic ecosystem management are generally calculated on averages and  
336 minima oxygen concentrations in water and they risk overlooking the effect of the oxygen fluctuation which  
337 are more relevant for communities' physiology (Tomasetti and Gobler, 2020). Enhancing monitoring  
338 programs by capturing the daily oxygen fluctuation will allow detection of when night-time low oxygen  
339 content is followed by a daytime recovery period, often reaching oxygen supersaturation. This alternation  
340 allows the aquatic communities to sustain their homeostasis in a fluctuating environment. Therefore, proper  
341 monitoring can inform a correct management to conserve, protect, and restore coastal water mosaic  
342 patterns of primary producers to ensure the ecological relevant fluctuations of dissolved oxygen.

343 In this contribution, based on recent research, we provide evidence that the fluctuating oxygen creates a  
344 spatial and temporal heterogeneous aquatic oxygen landscape that we have defined here as “oxyscape”. The  
345 oxyscape—having an important role for aquatic animal physiology and adaptation and the ecosystem  
346 biogeochemistry—should be considered in the modelling and managing policies of aquatic ecosystems to  
347 capture the ecologically relevant oxygen fluctuation. Understanding the oxyscape can help to reconsider  
348 the dynamics of many productive aquatic ecosystems and we, therefore, call on ecologists to rethink their  
349 models and experiments in this light. This would increase understanding of variations and resilience of  
350 communities to changes in environmental conditions at the scales of space and time relevant to individual  
351 components.

352

353 **Author contributions**

354 MF and FG conceived the study and wrote sections 1 to 3. AB, GG, SR, FG wrote section 4.1 and 4.2 and  
355 provided data for the Mediterranean sites. MF, RM and DD wrote section 5 and provided data for the Red  
356 Sea. MF, LP and CVH wrote section 6. All the authors discussed and reviewed the final version.

357

### 358 **Competing interests**

359 The contact author has declared that none of the authors have competing interests.

360

### 361 **Financial support**

362 D.D. acknowledge the financial support of King Abdullah University of Science and Technology (KAUST)  
363 through the Circular Carbon Economy initiative (grant no. REI/1/4483-01-01). D.D also acknowledge the  
364 Competitive Research Grant (CRG-7-3739), “The role of the bacterial symbiome at the gill-water (air)  
365 interface in the evolution toward territorialisation (Microlanding),” 1 April 2019 to 31 March 2022. We are  
366 very grateful to the LabEx DRIIHM, French Programme “Investissements d’Avenir” (ANR-11-LABX-  
367 0010), which is managed by the French ANR, under the Human- Environment Observatory  
368 “Mediterranean coast” (OHM Littoral Méditerranéen) for its funding that allowed to collect data from the  
369 Prevost lagoon, the Excellence Initiative of Aix-Marseille University - A\*MIDEX, a French  
370 “Investissements d’Avenir” programme (reference number ANR-11-IDEX-0001- 02), the PACA region  
371 and the CNRS-INSU-EC2CO program (PREDHYP- O2 program) for its funding that allowed to collect  
372 data from the Berre lagoon, and the Occitanie region and the University of Nîmes for the funding of PhD  
373 Thesis of Julie Régis. We also thank all the members who allowed us to collect data from the coastal  
374 Mediterranean lagoons in France (syndicat mixte GIPREB, Mairie de Chateauneuf-les-Martigues,  
375 Compagnons de Maguelone). The PREDHYPO program has received funding from Excellence Initiative  
376 of Aix-Marseille University - A\*MIDEX, a French “Investissements d’Avenir” programme (reference  
377 number ANR-11-IDEX-0001- 02), the PACA region and the CNRS-INSU-EC2CO program (PREDHYP-  
378 O2 program).

379 **References**

- 380 Aldunate, M., De la Iglesia, R., Bertagnolli, A. D., and Ulloa, O.: Oxygen modulates bacterial community  
381 composition in the coastal upwelling waters off central Chile, *Deep. Res. Part II Top. Stud. Oceanogr.*, 156,  
382 68–79, <https://doi.org/10.1016/j.dsr2.2018.02.001>, 2018.
- 383 Andersen, M. R., Kragh, T., and Sand-Jensen, K.: Extreme diel dissolved oxygen and carbon cycles in  
384 shallow vegetated lakes., *Proceedings. Biol. Sci.*, 284, 20171427, <https://doi.org/10.1098/rspb.2017.1427>,  
385 2017.
- 386 Antão, L. H., Bates, A. E., Blowes, S. A., Waldock, C., Supp, S. R., Magurran, A. E., Dornelas, M., and  
387 Schipper, A. M.: Temperature-related biodiversity change across temperate marine and terrestrial systems,  
388 *Nat. Ecol. Evol.*, 4, 927–933, <https://doi.org/10.1038/s41559-020-1185-7>, 2020.
- 389 Bartoli, M., Castaldelli, G., Nizzoli, D., Gatti, L. G., and Viaroli, P.: Benthic Fluxes of Oxygen, Ammonium  
390 and Nitrate and Coupled-uncoupled Denitrification Rates within Communities of Three Different Primary  
391 Producer Growth Forms, *Mediterr. Ecosyst.*, 225–233, [https://doi.org/10.1007/978-88-470-2105-1\\_29](https://doi.org/10.1007/978-88-470-2105-1_29),  
392 2001.
- 393 Bates, A. E., Helmuth, B., Burrows, M. T., Duncan, M. I., Garrabou, J., Guy-Haim, T., Lima, F., Queiros,  
394 A. M., Seabra, R., Marsh, R., Belmaker, J., Bensoussan, N., Dong, Y., Mazaris, A. D., Smale, D., Wahl, M.,  
395 and Rilov, G.: Biologists ignore ocean weather at their peril, *Nature*, 560, 299–301, 2018.
- 396 Bennett, J. M., Sunday, J., Calosi, P., Villalobos, F., Martínez, B., Molina-Venegas, R., Araújo, M. B., Algar,  
397 A. C., Clusella-Trullas, S., Hawkins, B. A., Keith, S. A., Kühn, I., Rahbek, C., Rodríguez, L., Singer, A.,  
398 Morales-Castilla, I., and Olalla-Tárraga, M. Á.: The evolution of critical thermal limits of life on Earth, *Nat.*  
399 *Commun.*, 12, 1–9, <https://doi.org/10.1038/s41467-021-21263-8>, 2021.
- 400 Benoiston, A. S., Ibarbalz, F. M., Bittner, L., Guidi, L., Jahn, O., Dutkiewicz, S., and Bowler, C.: The  
401 evolution of diatoms and their biogeochemical functions, *Philos. Trans. R. Soc. B Biol. Sci.*, 372,  
402 <https://doi.org/10.1098/rstb.2016.0397>, 2017.
- 403 Berg, P., Huettel, M., Glud, R. N., Reimers, C. E., and Attard, K. M.: Aquatic Eddy Covariance: The Method  
404 and Its Contributions to Defining Oxygen and Carbon Fluxes in Marine Environments, *Ann. Rev. Mar.*  
405 *Sci.*, 14, 431–455, <https://doi.org/10.1146/annurev-marine-042121-012329>, 2022.
- 406 Bernhardt, J. R., O'Connor, M. I., Sunday, J. M., and Gonzalez, A.: Life in fluctuating environments, *Philos.*

407 Trans. R. Soc. Lond. B. Biol. Sci., 375, 20190454, <https://doi.org/10.1098/rstb.2019.0454>, 2020.

408 Bickler, P. E. and Buck, L. T.: Hypoxia tolerance in reptiles, amphibians, and fishes: Life with variable  
409 oxygen availability, *Annu. Rev. Physiol.*, 69, 145–170,  
410 <https://doi.org/10.1146/annurev.physiol.69.031905.162529>, 2007.

411 Bitter, M. C., Wong, J. M., Dam, H. G., Donelan, S. C., Kenkel, C. D., Komoroske, L. M., Nickols, K. J.,  
412 Rivest, E. B., Salinas, S., Burgess, S. C., Morash, A. J., Neufeld, C., MacCormack, T. J., and Currie, S.: The  
413 importance of incorporating natural thermal variation when evaluating physiological performance in wild  
414 species, *J. Exp. Biol.*, 221, <https://doi.org/10.1242/JEB.164673>, 2018.

415 Bitter, M. C., Wong, J. M., Dam, H. G., Donelan, S. C., Kenkel, C. D., Komoroske, L. M., Nickols, K. J.,  
416 Rivest, E. B., Salinas, S., Burgess, S. C., and Lotterhos, K. E.: Fluctuating selection and global change: A  
417 synthesis and review on disentangling the roles of climate amplitude, predictability and novelty, *Proc. R.*  
418 *Soc. B Biol. Sci.*, 288, <https://doi.org/10.1098/rspb.2021.0727>, 2021.

419 Blewett, T. A., Binning, S. A., Weinrauch, A. M., Ivy, C. M., Rossi, G. S., Borowiec, B. G., Lau, G. Y.,  
420 Overduin, S. L., Aragao, I., and Norin, T.: Physiological and behavioural strategies of aquatic animals living  
421 in fluctuating environments, *J. Exp. Biol.*, 225, jeb242503, <https://doi.org/10.1242/jeb.242503>, 2022.

422 Booth, J., Fusi, M., Giomi, F., Chapman, E., Diele, K., and McQuaid, C.: Diel oxygen fluctuation drives  
423 the thermal response and metabolic performance of coastal marine ectotherms, *Proc. R. Soc. B Biol. Sci.*,  
424 288, 20211141, 2021.

425 Booth, J. M., Fusi, M., Marasco, R., Mbobo, T., and Daffonchio, D.: Fiddler crab bioturbation determines  
426 consistent changes in bacterial communities across contrasting environmental conditions, *Sci. Rep.*, 9,  
427 <https://doi.org/10.1038/s41598-019-40315-0>, 2019.

428 Booth, J. M., Giomi, F., Daffonchio, D., Mcquaid, C. D., and Fusi, M.: Disturbance of primary producer  
429 communities disrupts the thermal limits of the associated aquatic fauna, *Sci. Total Environ.*, 872, 162135,  
430 <https://doi.org/10.1016/j.scitotenv.2023.162135>, 2023a.

431 Booth, J. M., Fusi, M., Marasco, R., and Daffonchio, D.: The microbial landscape in bioturbated mangrove  
432 sediment: A resource for promoting nature-based solutions for mangroves, *Microb. Biotechnol.*, in press,  
433 <https://doi.org/10.1111/1751-7915.14273>, 2023b.

434 Breen, P. A. and Mann, K. H.: Changing lobster abundance and the destruction of kelp beds by sea urchins,



435 Mar. Biol., 34, 137–142, <https://doi.org/10.1007/BF00390755>, 1976.

436 Breitburg, D. and Grégoire, M.: The ocean is losing its breath, *Unesco*, 11, 1–43, 2018.

437 Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert,  
438 D., Gutiérrez, D., Isensee, K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C.,  
439 Rabalais, N. N., Roman, M. R., Rose, K. A., Seibel, B. A., Telszewski, M., Yasuhara, M., and Zhang, J.:  
440 Declining oxygen in the global ocean and coastal waters, *Science* (80-. ), 359, eaam7240,  
441 <https://doi.org/10.1126/science.aam7240>, 2018.

442 Breitburg, D. L., Conley, D., Isensee, K., Levin, L. A., Limburg, K. E., and Williamson, P.: What can we  
443 do? Adaptation and solutions to declining ocean oxygen, *Ocean deoxygenation everyone's Probl. Causes,*  
444 *impacts, consequences Solut.*, 545–562, 2019.

445 Caballero-Alfonso, A. M., Carstensen, J., and Conley, D. J.: Biogeochemical and environmental drivers of  
446 coastal hypoxia, *J. Mar. Syst.*, 141, 190–199, <https://doi.org/10.1016/j.jmarsys.2014.04.008>, 2015.

447 Caron, D. A.: Inorganic nutrients, bacteria, and the microbial loop, *Microb. Ecol. Ecol.*, 28, 295–298,  
448 <https://doi.org/10.1007/BF00166820>, 1994.

449 Challenger, R. C., Robbins, L. L., and McClintock, J. B.: Variability of the carbonate chemistry in a shallow,  
450 seagrass-dominated ecosystem: Implications for ocean acidification experiments, *Mar. Freshw. Res.*, 67,  
451 163–172, <https://doi.org/10.1071/MF14219>, 2016.

452 Chapman, D.: *Water Quality Assessments - A Guide to Use of Biota, Sediments and Water in*  
453 *Environmental Monitoring - Second Edition*, E&FN Spon, London, 2021.

454 Chen, H., Costanza, R., and Kubiszewski, I.: Legitimacy and limitations of valuing the oxygen production  
455 of ecosystems, *Ecosyst. Serv.*, 58, 101485, <https://doi.org/10.1016/j.ecoser.2022.101485>, 2022.

456 Cowan, J. L. W., Pennock, J. R., and Boynton, W. R.: Seasonal and interannual patterns of sediment-water  
457 nutrient and oxygen fluxes in Mobile Bay, Alabama (USA): Regulating factors and ecological significance,  
458 *Mar. Ecol. Prog. Ser.*, 141, 229–245, <https://doi.org/10.3354/meps141229>, 1996.

459 Craig, H. and Hayward, T.: Oxygen Supersaturation in the Ocean: Biological Versus Physical Contributors,  
460 735, 1983–1986, 1983.

461 Dafforn, K. a., Simpson, S. L., Kelaher, B. P., Clark, G. F., Komyakova, V., Wong, C. K. C., and Johnston,  
462 E. L.: The challenge of choosing environmental indicators of anthropogenic impacts in estuaries, *Environ.*

463 Pollut., 163, 207–217, <https://doi.org/10.1016/j.envpol.2011.12.029>, 2012.

464 Denis, L., Gevaert, F., and Spilmont, N.: Microphytobenthic production estimated by in situ oxygen  
465 microprofiling: Short-term dynamics and carbon budget implications, *J. Soils Sediments*, 12, 1517–1529,  
466 <https://doi.org/10.1007/s11368-012-0588-8>, 2012.

467 Diaz, R. J. and Rosenberg, R.: Spreading dead zones and consequences for marine ecosystems, *Science* (80-  
468 .), 321, 926–929, <https://doi.org/10.1126/science.1156401>, 2008.

469 Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., and Marbà, N.: The role of coastal plant  
470 communities for climate change mitigation and adaptation, *Nat. Clim. Chang.*, 3, 961–968,  
471 <https://doi.org/10.1038/nclimate1970>, 2013.

472 Fenchel, T. and Finlay, B.: Oxygen and the spatial structure of microbial communities, *Biol. Rev.*, 83, 553–  
473 569, <https://doi.org/10.1111/j.1469-185X.2008.00054.x>, 2008.

474 Fusi, M., Daffonchio, D., Booth, J., and Giomi, F.: Dissolved Oxygen in Heterogeneous Environments  
475 Dictates the Metabolic Rate and Thermal Sensitivity of a Tropical Aquatic Crab, *Front. Mar. Sci.*, 8, 1–9,  
476 <https://doi.org/10.3389/fmars.2021.767471>, 2021.

477 Fusi, M., Marasco, R., Ramond, J., Barausse, A., and Baldanzi, S.: Editorial: Fluctuating Habitats :  
478 Ecological Relevance of Environmental Variability and Predictability on Species , Communities , and  
479 Ecosystems, *Front. Ecol. Evol.*, 10, 1–4, <https://doi.org/10.3389/fevo.2022.907622>, 2022.

480 Garzke, J., Connor, S. J., Sommer, U., and O’connor, M. I.: Trophic interactions modify the temperature  
481 dependence of community biomass and ecosystem function, *PLoS Biol.*, 17, 1–27,  
482 <https://doi.org/10.1371/journal.pbio.2006806>, 2019.

483 Giomi, F., Barausse, A., Duarte, C. M., Booth, J., Agusti, S., Saderne, V., Anton, A., Daffonchio, D., and  
484 Fusi, M.: Oxygen supersaturation protects coastal marine fauna from ocean warming, *Sci. Adv.*, 5, 1–8,  
485 <https://doi.org/10.1126/sciadv.aax1814>, 2019.

486 Glud, R. N.: Oxygen dynamics of marine sediments, *Mar. Biol. Res.*, 4, 243–289,  
487 <https://doi.org/10.1080/17451000801888726>, 2008.

488 Graham, N. A. J., Wilson, S. K., Carr, P., Hoey, A. S., Jennings, S., and MacNeil, M. A.: Seabirds enhance  
489 coral reef productivity and functioning in the absence of invasive rats, *Nature*, 559, 250–253,  
490 <https://doi.org/10.1038/s41586-018-0202-3>, 2018.

491 Guo, X., Song, G., Li, Y., Zhao, L., and Wang, J.: Switch of Bacteria Community Under Oxygen Depletion  
492 in Sediment of Bohai Sea, *Front. Mar. Sci.*, 9, 1–13, <https://doi.org/10.3389/fmars.2022.833513>, 2022.

493 Hochachka, P. W.: Temperature: the ectothermy option, in: *Phylogenetic and biochemical perspectives:*  
494 *Biochemistry and molecular biology of fishes*, 1, vol. 1, 313–322, [https://doi.org/10.1016/B978-0-444-](https://doi.org/10.1016/B978-0-444-89124-2.50016-6)  
495 [89124-2.50016-6](https://doi.org/10.1016/B978-0-444-89124-2.50016-6), 1991.

496 Ito, T., Nenes, A., Johnson, M. S., Meskhidze, N., and Deutsch, C.: Acceleration of oxygen decline in the  
497 tropical Pacific over the past decades by aerosol pollutants, *Nat. Geosci.*, 9, 443–447,  
498 <https://doi.org/10.1038/ngeo2717>, 2016.

499 Keeling, R. F., Körtzinger, A., and Gruber, N.: Ocean deoxygenation in a warming world, *Ann. Rev. Mar.*  
500 *Sci.*, 2, 199–229, <https://doi.org/10.1146/annurev.marine.010908.163855>, 2010.

501 Kerrison, P., Hall-Spencer, J. M., Suggett, D. J., Hepburn, L. J., and Steinke, M.: Assessment of pH  
502 variability at a coastal CO<sub>2</sub> vent for ocean acidification studies, *Estuar. Coast. Shelf Sci.*, 94, 129–137,  
503 <https://doi.org/10.1016/j.ecss.2011.05.025>, 2011.

504 Kim, H., Takayama, K., Hirose, N., Onitsuka, G., Yoshida, T., and Yanagi, T.: Biological modulation in the  
505 seasonal variation of dissolved oxygen concentration in the upper Japan Sea, *J. Oceanogr.*, 75, 257–271,  
506 <https://doi.org/10.1007/s10872-018-0497-6>, 2019.

507 Kinney, E. H. and Roman, C. T.: Response of primary producers to nutrient enrichment in a shallow  
508 estuary, *Mar. Ecol. Prog. Ser.*, 163, 89–98, <https://doi.org/10.3354/meps163089>, 1998.

509 Krause-Jensen, D., Marba, N., Sanz-Martin, M., Hendriks, I. E., Thyrring, J., Carstensen, J., Sejr, M. K.,  
510 and Duarte, C. M.: Long photoperiods sustain high pH in Arctic kelp forests, *Sci. Adv.*, 2, e1501938–  
511 [e1501938](https://doi.org/10.1126/sciadv.1501938), <https://doi.org/10.1126/sciadv.1501938>, 2016.

512 Laffoley, D. and Baxter, J. M.: *Explaining Ocean Warming: Causes, scale, effects and consequences*, 458  
513 pp., <https://doi.org/http://dx.doi.org/10.2305/IUCN.CH.2016.08.en>, 2016.

514 Laffoley, D. D. A. and Baxter, J. M.: *Ocean Deoxygenation: Everyone’s Problem-Causes, Impacts,*  
515 *Consequences and Solutions*, 2019.

516 Limburg, K. E., Breitburg, D., Swaney, D. P., and Jacinto, G.: Ocean Deoxygenation: A Primer, *One Earth*,  
517 2, 24–29, <https://doi.org/10.1016/j.oneear.2020.01.001>, 2020.

518 Mallin, M. A., Paerl, H. W., Rudek, J., and Bates, P. W.: Regulation of estuarine primary production by

519 watershed rainfall and river flow, *Mar. Ecol. Prog. Ser.*, 93, 199–203,  
520 <https://doi.org/10.3354/meps093199>, 1993.

521 Manzello, D. P.: Ocean acidification hot spots: Spatiotemporal dynamics of the seawater CO<sub>2</sub> system of  
522 eastern Pacific coral reefs, *Limnol. Oceanogr.*, 55, 239–248, <https://doi.org/10.4319/lo.2010.55.1.0239>,  
523 2010.

524 Marasco, R., Fusi, M., Coscolín, C., Barozzi, A., Almendral, D., Bargiela, R., Gohlke, C., Christopher, P.,  
525 Dittrich, J., Gohlke, H., Matesanz, R., Sanchez-carrillo, S., Mapelli, F., Chernikova, T. N., Golyshin, P. N.,  
526 Ferrer, M., and Daffonchio, D.: Enzyme adaptation to habitat thermal legacy microbiomes, *Nat. Commun.*,  
527 in press, <https://doi.org/10.1038/s41467-023-36610-0>, 2023.

528 McArley, T. J., Sandblom, E., and Herbert, N. A.: Fish and hyperoxia—From cardiorespiratory and  
529 biochemical adjustments to aquaculture and ecophysiology implications, *Fish Fish.*, 1–32,  
530 <https://doi.org/10.1111/faf.12522>, 2020.

531 McArley, T. J., Morgenroth, D., Zena, L. A., Ekström, A. T., and Sandblom, E.: Experimental hyperoxia  
532 (O<sub>2</sub> supersaturation) reveals a gill diffusion limitation of maximum aerobic performance in fish, *Biol. Lett.*,  
533 18, 20220401, <https://doi.org/10.1098/rsbl.2022.0401>, 2022.

534 Meire, L., Soetaert, K. E. R., and Meysman, F. J. R.: Impact of global change on coastal oxygen dynamics  
535 and risk of hypoxia, *Biogeosciences*, 10, 2633–2653, <https://doi.org/10.5194/bg-10-2633-2013>, 2013.

536 Morash, A. J., Neufeld, C., MacCormack, T. J., and Currie, S.: The importance of incorporating natural  
537 thermal variation when evaluating physiological performance in wild species, *J. Exp. Biol.*, 221,  
538 <https://doi.org/10.1242/JEB.164673>, 2018.

539 Nilsson, H. C. and Rosenberg, R.: Benthic habitat quality assessment of an oxygen stressed fjord by surface  
540 and sediment profile images, *J. Mar. Syst.*, 11, 249–264, [https://doi.org/10.1016/S0924-7963\(96\)00111-X](https://doi.org/10.1016/S0924-7963(96)00111-X),  
541 1997.

542 Pacherres, C. O., Ahmerkamp, S., Koren, K., Richter, C., and Holtappels, M.: Ciliary flows in corals  
543 ventilate target areas of high photosynthetic oxygen production, *Curr. Biol.*, 32, 4150-4158.e3,  
544 <https://doi.org/10.1016/j.cub.2022.07.071>, 2022.

545 Palmeri, L., Barausse, A., and Jorgensen, S. E.: *Ecological processes handbook*, edited by: Palmeri, L.,  
546 Barausse, A., and Jorgensen, S. E., CRC Press, Boca Raton FL, 375 pp., 2013.

547 Pörtner, H.-O.: Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-  
548 related stressor effects in marine ecosystems., *J. Exp. Biol.*, 213, 881–893, 2010.

549 Pörtner, H.-O., Scholes, R. J., Arneith, A., Barnes, D. K. A., Burrows, M. T., Diamond, S. E., Duarte, C.  
550 M., Kiessling, W., Leadley, P., Managi, S., McElwee, P., Midgley, G., Ngo, H. T., Obura, D., Pascual, U.,  
551 Sankaran, M., Shin, Y. J., and Val, A. L.: Overcoming the coupled climate and biodiversity crises and their  
552 societal impacts, *Science* (80-. ), 380, eabl4881, <https://doi.org/10.1126/science.abl4881>, 2023.

553 Power, L. D. and Cardinale, B. J.: Species richness enhances both algal biomass and rates of oxygen  
554 production in aquatic microcosms, *Oikos*, 118, 1703–1711, [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0706.2009.17585.x)  
555 [0706.2009.17585.x](https://doi.org/10.1111/j.1600-0706.2009.17585.x), 2009.

556 Rigaud, S., Deflandre, B., Maire, O., Bernard, G., Duchêne, J. C., Poirier, D., and Anschutz, P.: Transient  
557 biogeochemistry in intertidal sediments: New insights from tidal pools in *Zostera noltei* meadows of  
558 Arcachon Bay (France), *Mar. Chem.*, 200, 1–13, <https://doi.org/10.1016/j.marchem.2018.02.002>, 2018.

559 Riser, S. C. and Johnson, K. S.: Net production of oxygen in the subtropical ocean, *Nature*, 451, 323–325,  
560 <https://doi.org/10.1038/nature06441>, 2008.

561 Sandrin, T. R., Dowd, S. E., Herman, D. C., and Maier, R. M.: Aquatic environments., in: *Environment*  
562 *Microbiology*, Elsevier, 103–122, 2009.

563 Schmidt, S., Diallo, I. I., Derriennic, H., Fallou, H., and Lepage, M.: Exploring the susceptibility of turbid  
564 estuaries to hypoxia as a prerequisite to designing a pertinent monitoring strategy of dissolved oxygen,  
565 *Front. Mar. Sci.*, 6, 1–8, <https://doi.org/10.3389/fmars.2019.00352>, 2019.

566 Schmidtko, S., Stramma, L., and Visbeck, M.: Decline in global oceanic oxygen content during the past five  
567 decades, *Nature*, 542, 335–339, <https://doi.org/10.1038/nature21399>, 2017.

568 Shaw, E. C., Munday, P. L., and McNeil, B. I.: The role of CO<sub>2</sub> variability and exposure time for biological  
569 impacts of ocean acidification, *Geophys. Res. Lett.*, 40, 4685–4688, <https://doi.org/10.1002/grl.50883>,  
570 2013.

571 Spietz, R. L., Williams, C. M., Rocap, G., and Horner-Devine, M. C.: A dissolved oxygen threshold for  
572 shifts in bacterial community structure in a seasonally hypoxic estuary, *PLoS One*, 10, 1–18,  
573 <https://doi.org/10.1371/journal.pone.0135731>, 2015.

574 Steiner, P. A., De Corte, D., Geijo, J., Mena, C., Yokokawa, T., Rattei, T., Herndl, G. J., and Sintes, E.:

575 Highly variable mRNA half-life time within marine bacterial taxa and functional genes, *Environ. Microbiol.*,  
576 21, 3873–3884, <https://doi.org/10.1111/1462-2920.14737>, 2019.

577 Stillman, J. H. and Somero, G. N.: A comparative analysis of the upper thermal tolerance limits of eastern  
578 Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical zonation, acclimation, and  
579 phylogeny., *Physiol. Biochem. Zool.*, 73, 200–208, <https://doi.org/10.1086/316738>, 2000.

580 Stramma, L., Schmidtko, S., Levin, L. A., and Johnson, G. C.: Ocean oxygen minima expansions and their  
581 biological impacts, *Deep. Res. Part I Oceanogr. Res. Pap.*, 57, 587–595,  
582 <https://doi.org/10.1016/j.dsr.2010.01.005>, 2010.

583 Talke, S. A., De Swart, H. E., and De Jonge, V. N.: An idealized model and systematic process study of  
584 oxygen depletion in highly turbid estuaries, *Estuaries and Coasts*, 32, 602–620,  
585 <https://doi.org/10.1007/s12237-009-9171-y>, 2009.

586 Tomasetti, S. J. and Gobler, C. J.: Dissolved oxygen and pH criteria leave fisheries at risk, *Science (80-. )*,  
587 368, 372–373, <https://doi.org/10.1126/science.aba4896>, 2020.

588 Trowbridge, C. D., Davenport, J., Cottrell, D. M., Harman, L., Plowman, C. Q., Little, C., and McAllen, R.:  
589 Extreme oxygen dynamics in shallow water of a fully marine Irish sea lough, *Reg. Stud. Mar. Sci.*, 11, 9–16,  
590 <https://doi.org/10.1016/j.rsma.2017.01.008>, 2017.

591 Vargas, C. A., Lagos, N. A., Lardies, M. A., Duarte, C., Manríquez, P. H., Aguilera, V. M., Broitman, B.,  
592 Widdicombe, S., and Dupont, S.: Species-specific responses to ocean acidification should account for local  
593 adaptation and adaptive plasticity, *Nat. Ecol. Evol.*, 1, 1–7, <https://doi.org/10.1038/s41559-017-0084>,  
594 2017.

595 Vargas, C. A., Cuevas, L. A., Broitman, B. R., San Martin, V. A., Lagos, N. A., Gaitán-Espitia, J. D., and  
596 Dupont, S.: Upper environmental pCO<sub>2</sub> drives sensitivity to ocean acidification in marine invertebrates,  
597 *Nat. Clim. Chang.*, 12, 200–207, <https://doi.org/10.1038/s41558-021-01269-2>, 2022.

598 Viaroli, P. and Christian, R. R.: Description of trophic status, hyperautotrophy and dystrophy of a coastal  
599 lagoon through a potential oxygen production and consumption index - TOSI: Trophic Oxygen Status  
600 Index, *Ecol. Indic.*, 3, 237–250, <https://doi.org/10.1016/j.ecolind.2003.11.001>, 2004.

601 Virtanen, E. A., Norkko, A., Nyström Sandman, A., and Viitasalo, M.: Identifying areas prone to coastal  
602 hypoxia - The role of topography, *Biogeosciences*, 16, 3183–3195, <https://doi.org/10.5194/bg-16-3183->

603 2019, 2019.


604 Winter, G., Hetzel, Y., Huang, P., Hipsey, M. R., Mulligan, R. P., and Hansen, J.: Coastal processes, extreme  
605 events and forecasting, in: *Marine Extreme*, Routledge, 17, 2019.

606

607 **Figure captions**

608

609 **Figure 1.** Range of oxygen saturations in several productive aquatic habitats. The dashed line represents  
 610 100% oxygen saturation. The violin plots report the densities of the frequencies of oxygen saturation  
 611 measured at each ecosystem across different season (based on data availability). Data source is specified for  
 612 each environment in the Supplementary Dataset for Figure 1 and 2.




613

614




615 **Figure 2.** Fluctuation of oxygen saturation measured at the bottom (4 m) and surface (30 cm) oxygen  
616 saturation of the Rove canal in France that highlight the contrast between the two depths of eutrophicated  
617 coastal environment from the winter to summer. The analysis of these case studies revealed different power  
618 to predict oxygen variation.



619

620


621 **Figure 3.** Nutrient concentration, dissolved oxygen and water temperature measurements made in the  
 622 water of a salt marsh of the Northern Venice Lagoon (Italy) on August 13th, 2015. A) Concentration of  
 623 different nitrogen forms in the marsh water (notice the two different vertical axes) over time in the central  
 624 part of the day. Measurements are made in the water entering the marsh (flood phase, left of the vertical  
 625 gray line which indicates the tidal peak) and quitting from the salt marsh (ebb phase, right of the vertical  
 626 gray line). After some time spent in the marsh, which is flooded by tide twice per day, water becomes  
 627 depleted in inorganic nutrients, presumably due to the uptake by primary producers, exchanges with the  
 628 sediments, and (in the case of ammonium) nitrification. Concurrently, water is enriched with dissolved  
 629 organic nitrogen related to the intense biological activity within the marsh. B) Dissolved oxygen and  
 630 temperature of the water entering the marsh (flood phase, left of the vertical gray line which indicates the  
 631 tidal peak) and quitting from the salt marsh (ebb phase, right of the vertical gray line) over time in the  
 632 central part of the day. While in the marsh, which is a biologically productive habitat compared to the  
 633 surrounding waters, the water becomes enriched with dissolved oxygen paralleled by an increase in water  
 634 temperature.



635

636


637 **Figure 4.** Scheme of the benthic biogeochemical cycle in relation to diel oxygen fluctuation. (A) The relative  
 638 evolution of the bottom water chemical composition, (B) evolution of the location of the most important  
 639 biogeochemical processes in the benthic compartment. (C) Relative influence of external drivers on benthic  
 640 processes. (D) Relative evolution of resulting flux direction and intensity. Note that the graphs are not  
 641 drawn to scale. Although concentrations of  $\text{PO}_4^{3-}$  and  $\text{NH}_4^+$  and of Mn and Fe may behave differently,  
 642 they are represented as similar in this figure for simplification.



643

644

645 **Figure 5.** Daily bacterial community cycle scheme in relation to diel oxygen fluctuation. Relative shift of  
 646 microbial community controlled by the oxygen availability that control the ratio between aerobic and  
 647 anaerobic taxa and their relative function. Oxygen fluctuation is important to determine the diel cycling of  
 648 the bacterial communities that in turn affect the biogeochemistry and the overall functioning of the aquatic  
 649 ecosystems. Note that the graphs are not drawn to scale.



650

651