

Sap flow and leaf gas exchange response to drought and heatwave in urban green spaces in a Nordic city

Joyson Ahongshangbam^{1,4}, Liisa Kulmala^{2,3}, Jesse Soininen¹, Yasmin Frühauf¹, Esko Karvinen², Yann Salmon^{1,3}, Anna Lintunen^{1,3}, Anni Karvonen¹, and Leena Järvi^{1,4}

¹Institute for Atmospheric and Earth System Research (INAR)/Physics, Faculty of Science, University of Helsinki, Helsinki, Finland

²Finnish Meteorological Institute, Helsinki, Finland

³Institute for Atmospheric and Earth System Research/Forest Sciences, Viikki Plant Science Centre (ViPS), Faculty of Agriculture and Forestry, University of Helsinki, Helsinki, Finland

⁴Helsinki Institute of Sustainability Science, University of Helsinki, Finland

Correspondence: Joyson Ahongshangbam (joyson.ahongshangbam@helsinki.fi)

Abstract. ~~Urban vegetation plays a role in offsetting urban CO₂ emissions, mitigating heat through tree transpiration and shading, and acting as deposition surface for pollutants.~~ ~~Urban vegetation plays an important role in offsetting urban CO₂ emissions and mitigating heat through tree transpiration and shading.~~ With frequent heatwave events and the accompanying drought, the functioning of urban trees is severely affected in terms of photosynthesis and transpiration rate. The detailed response is however still unknown despite tree functioning having crucial effects on the ecosystem services they provide. We conducted sap flux density (J_s) and leaf gas exchange measurements of three trees per species (*Tilia cordata*, *Tilia × europaea*, *Betula pendula*, *Malus spp.*) located at four types of urban green areas (Park, Street, Forest, Orchard) in Helsinki, Finland, over two contrasting summers 2020 and 2021. Summer 2021 had a strong heatwave and drought, whereas summer 2020 was more typical for Helsinki. In this study, our aim was to understand the responses of urban tree transpiration and leaf gas exchange to heatwave and drought and examine the main environmental drivers controlling the transpiration rate during these periods in urban green areas. We observed varying responses of ~~sap flux density~~ ~~tree water use~~ during the heatwave period at the four urban sites. J_s was found to be 35-67% higher during the heatwave as compared to the non-heatwave period at the Park, Forest, and Orchard sites but ~~but similar J_s was observed at the Street site during the heatwave as compared to the non-heatwave period~~ ~~no significant difference was found at the Street site.~~ Our results showed that J_s was higher (31-63%) at all sites during drought as compared to non-dry periods. The higher J_s during the heatwave and dry periods were mainly driven by the high atmospheric demand for evapotranspiration represented by the vapor pressure deficit (VPD), suggesting that the trees were not experiencing severe enough heat or drought stress that stomatal control would have decreased transpiration. Accordingly, photosynthetic potential (A_{max}), stomatal conductance (g_s), and transpiration (E) at the leaf level did not change at the four sites during heatwave and drought periods. Only g_s was significantly reduced during the drought period at the Park site. VPD explained 55-69% variations in the daily mean J_s during heatwave and drought periods at all sites while at the Forest site, the increase of J_s with the rise in VPD saturates after certain level of VPD, which might be evident due to low soil water availability at the Forest site during these hot and dry periods. ~~except at the Forest site where the saturation of J_s at high VPD was evident due~~

to low soil water availability. Overall, the heat and drought conditions were untypically harsh for the region but not excessive enough to restrict stomatal control and the increased transpiration indicating that ecosystem services such as cooling was not at risk.

Keywords: climate extreme, drought, sap flux, transpiration, urban trees

1 Introduction

Urbanization is increasing rapidly and transforming the natural environment, land cover, and ecological functions. Urbanization enhances CO₂ emissions and the local urban heat island (UHI) effect, leading to harsher conditions in cities and a decrease in human thermal comfort when compared to more natural surroundings (Oke et al., 1989; Roth et al., 1989). These challenges highlight the need for adequate urban planning in the long-term development of urban areas. Urban green areas play a vital role in compensating urban CO₂ emissions and mitigating the UHI effect as they have the potential for carbon sequestration and storage, and regulation of water for cooling the surrounding and surface energy balance (Lindén et al., 2016; Bowler et al., 2010). Urban trees also provide other ecosystem services such as cooling effect through shading, pollutant deposition infiltration, aesthetics and recreation, buffer for noise and wind, and soil conservation (Brack, 2002; Jo, 2002; Jim and Chen, 2009; Pataki et al., 2009; Har). Several studies have highlighted the CO₂ sequestration potential and annual carbon storage in urban vegetation (Nowak and Crane, 2002; Davies et al., 2011; Muñoz-Vallés et al., 2013; Nowak et al., 2013), and addressed urban green areas as a way of mitigating the cities' global GHG emissions in cities (Dhakal, 2010; Paloheimo and Salmi, 2013; Pataki et al., 2021). The role of urban trees in mitigating UHI through shading by tree canopies and cooling effect by transpiration have been reported in many urban studies (Rahman et al., 2019; Pataki et al., 2011). Trees provide cooling effect through two main primary mechanisms. Firstly, urban trees reduce surface temperature by shading, which results in less absorption and storage of incoming short-wave radiation by surfaces, thereby lowering the local air temperature. Secondly, through the process of transpiration where water taken up by roots is released through leaf stomates. The energy consumed to evaporate the water released from leaf stomata provides cooling effect on the leaf surface and lower nearby air temperature by advection. In urban condition, trees are subjected to human disturbances such as construction activities, dense building and vandalism (Czaja et al., 2020) and climate change impacts including extreme weather such as heatwaves and drought. All these affect the potential of trees to mitigate and adapt to climate change, and thus it is important to understand the response and functions of urban trees during extreme climate events.

In general, environmental conditions for urban trees are often more extreme than those in a natural forest stand, having e.g. higher air temperature, lower air humidity, and more limited soil water and nutrient availability (Nielsen et al., 2007). All these differences in local microclimate, growing conditions, species type, disturbances and management activities affected the functioning of urban trees. The rise in temperature during summer heatwaves affects leaf temperature potentially leading to leaf damages (Kunert et al., 2022; Atkin and Tjoelker, 2003; Ghannoum and Way, 2011), and vapor-pressure deficit (VPD) affecting transpiration and photosynthesis through stomatal control (Lloyd and Farquhar, 2008). In urban areas, tree reactions to heatwaves are rarely studied, but trees in natural forests can adapt to the rising temperature by enhancing growth

and utilized water more efficiently, provided enough moisture in the soil (Winbourne et al., 2020). The foliar temperatures of urban trees are often higher, which limits photosynthesis and transpirations through enzyme activity and stomatal regulation (Bussotti et al., 2014), hence such acclimation may not be apparent in urban settings. The response of urban trees to climate change has also been found to differ from that of forest trees and moreover, urban trees have been found to be more susceptible to multiple stresses enhanced by climate change (Bussotti et al., 2014). Droughts have occurred more frequently in recent times causing severe symptoms for trees even in areas that are considered to be rather moist high latitude areas (Hartmann et al., 2022). Drought usually co-occurs with high air temperature during summer severely affecting the functioning of trees and potentially resulting in leaf damage, reduced carbon assimilation and transpiration through lowered stomatal conductance (Bussotti et al., 2014; Winbourne et al., 2020). Tree usually response to drought in two ways by either avoiding the large decrease in water potential and relative water content or tolerate by maintaining the physiological process during drought (Villar-Salvador et al., 2004; De Micco and Aronne, 2012). However, the effect of an urban environment on the tree's response to stress such as drought and heatwaves are not known, particularly in the boreal urban environment. A few studies have investigated the potential cooling effect of urban trees during heat and drought (Gillner et al., 2015b), and the impact of drought on urban tree functions (Rötzer et al., 2021). The urban tree functions were found to be affected by species type, growing conditions, local climate and water availability, particularly during heatwave and drought periods. Nonetheless, urban tree responses to extreme heat and drought are poorly addressed and, with complex urban stressors and heterogeneity, need further study. Especially information on the impact of heat and drought on urban trees in high-latitude cities is lacking. Global warming is more prominent at high latitudes making urban ecosystems in high latitudes particularly vulnerable. Thus, it is important to understand the responses of trees to heat and drought in various urban environments in the boreal region.

In this study, we measured the functions of a set of typical urban tree species, tree, particularly their transpiration and leaf gas exchanges, during heatwave and drought periods in a boreal urban environment in Helsinki, Finland. In particular, our focus was on the response of the variability of sap flow rate patterns during heat and dry period tree water use patterns in different urban green areas, and on the environmental controls of sap flow rate tree water use. We addressed the following research questions:

1. How does heatwave affect transpiration and leaf gas exchange rates of urban trees?
2. How does drought affect transpiration and leaf gas exchange rates of urban trees?
3. What are the main environmental drivers affecting transpiration rates during heatwave and drought in urban green areas?
How do environmental drivers affect transpiration rates during heatwave and drought periods in urban green areas?

We hypothesize that H1) even though a heatwave increases sap flux density due to increased VPD in the short term, it decreases the photosynthetic potential due to lower stomatal conductance and also decreasing the role of VPD as a driver, heatwave increases sap flux density due to increased VPD decrease the photosynthetic potential due to lower stomatal conductance, H2) drought decreases both sap flux density transpiration and the photosynthetic potential. H3) the role of VPD as a driver decreases during heatwave and drought periods, again due to lower stomatal conductance To answer these questions and test the hypotheses, we conducted continuous sap flux and manual leaf gas exchange measurements at four urban green areas (Park,

90 Street, Forest and Orchard) in Helsinki during the summers of 2020 and 2021. ~~The summer of 2021 was hot and dry as the mean air temperature in July 2021 was 21.6°C, being 21% higher than in July 2020 (16.7°C) and 19% higher than the average mean temperature in July (18.1 °C) during a climatic reference period (1991 to 2020). The total precipitation for the months of June and July 2021 (86 mm) was 51% lower than during June and July in 2020 (177 mm) and 27% lower than the average total precipitation in June and July during the climatic reference period (117 mm).~~

95 2 Methods

2.1 Sites description

The study was conducted in the vicinity of the University of Helsinki Kumpula campus, located 4 km northeast of the Helsinki city centre. The Kumpula area is characterized by heterogeneous land-use cover (Figure 1a), particularly by diverse urban vegetation. Within the study area, four sites were selected to be studied: a park with sparse trees ('Park'), a [single line roadside trees roadside plantation](#) ('Street'), an urban forest ('Forest'), and an apple orchard ('Orchard'). These sites are located close to a micrometeorological eddy covariance station (FI-Kmp, 60°12'11.3"N 24°57'40.4"E) which is also an Associated Ecosystem Station of ICOS (Integrated Carbon Observation system), and part of the SMEAR (Station for Measuring Ecosystem Atmosphere Relations) III station (Vesala et al., 2008; Järvi et al., 2009). Overall, Helsinki is a humid continental region (according to Köppen climate classification), with annual precipitation of 652 mm yr⁻¹ and annual mean temperature of 6.5 °C during the 105 30-year climatic reference period 1991-2020 (FMI, 2021). [The summer of 2021 was hot and dry as the mean air temperature in July 2021 was 21.6°C, being 21% higher than in July 2020 \(16.7°C\) and 19% higher than the average mean temperature in July \(18.1 °C\) during a climatic reference period \(1991 to 2020\). The total precipitation for the months of June and July 2021 \(86 mm\) was 51% lower than during June and July in 2020 \(177 mm\) and 27% lower than the average total precipitation in June and July during the climatic reference period \(117 mm\).](#)

110 The urban park (Park; Figure 1b) is located in the Kumpula botanical garden, South-west of the FI-Kmp measurement tower. The site is characterized by a mixture of *Tilia species* trees, and a ground layer of short vegetation comprised mainly of lawn species, clovers (*Trifolium repens*) and mosses. The ground vegetation was mowed using an automatic mowing device leaving the clippings on-site and irrigation was activated on dry and warm days within the wider park area. However, the mowing and irrigation were restricted in-between the study trees (0.25 ha area) during the measurement period but the tree roots reached 115 the irrigated area.

The roadside plantation (Street; Figure 1 c) is located on the Hermannin rantatie road, 0.8 km east of the FI-Kmp measurement tower. It consists of a row of *Tilia × europaea*, which is a hybrid of *T. cordata* and *T. platyphyllos*. The species is the most commonly planted urban tree in Helsinki and Nordic countries in general, comprising 44% of Helsinki's Street trees (Sjöman et al., 2012). The street trees grow over a soil patch spreading 60 m long and 2.7 m wide, with an average tree spacing of 8.2 m. 120 Normally, the trees are regularly trimmed and maintained by the city gardening company but they were non-managed during our study.

Urban forest (Forest; Figure 1d) is a small forest patch (25x30 m²) located between the FI-Kmp measurement tower in the north and the Kumpula garden in the south. This site is dominated by mature *Betula pendula* trees. This site was the only non-managed study site and was regenerated naturally. Other deciduous trees such as *Betula pubescens*, *Alnus glutinosa*, *Acer platanoids*, and *Ulmus glabra* were also found in this urban forest as a species mixture. The ground layer was sparse consisting mainly of *Aegopodium podagraria* and bare spots.

The apple orchard (Orchard; Figure 1e) is located in the Kumpula school garden 0.9 km west of the FI-Kmp measurement tower. The site is characterized by scattered apple trees (20 trees per 30 m x 30 m area) planted over a managed lawn. There was no irrigation. The lawn in our measurement area was manually mowed a few times during the summer.

More detailed descriptions of all four sites are given in Table 1 and of soil properties; [particularly about soil water retention properties](#) in Appendix A1. All sites were equipped with continuous measurements of sap flux and meteorological variables from June 2020 to September 2021, except the Orchard site, where data was recorded only from June-September 2021. The continuous measurements were accompanied by manual measurements of leaf gas exchange during the summer months (June-August).

Table 1. The four study sites, the dominating tree species, their mean diameter at breast height (DBH), sapwood area, height, years since plantation (age) and soil particle type.

	Park	Street	Forest	Orchard
Latitude	60°12'08.4"N	60°11'51.6"N	60°12'07.7"N	60°12'30.17"N
Longitude	24°57'21.4"E	24°58'13.2"E	24°57'33.0"E	24°56'57.77"E
Tree species	<i>Tilia cordata</i>	<i>Tilia × europaea</i>	<i>Betula pendula</i>	<i>Malus spp.</i>
Drought strategies	Drought tolerant	Drought tolerant	Drought tolerant	Drought tolerant
DBH (cm)	26.3	19.5	23.6	30
Sapwood area (cm²)	433.8	271.9	349.7	397.0
Height (m)	12.5	10	22	6.5
Age (years)	26	34	35	72 (approx.)
Soil type	Sand moraine	Fine sand moraine	Sand moraine	Sand clay

2.2 Sap flux measurements

Sap flux measurements were conducted using the Thermal Dissipation Probe (TDP; Granier, 1985). TDP sensors consist of two thermocouple needles (20-30 mm long) where one needle acts as a heating probe and the other as a reference probe. The thermocouples measure the temperature difference between the heated probe and the reference probe, which is used to calculate the sap flux density and also the whole-tree water use when scaled up. In our study, we selected three sample trees at each site (a total of 12 trees) based on high sun exposure and dominant position at the site. In each tree, a TDP sensor was inserted into the stem xylem at a height of 1.3 m, where a vertical distance of 10 cm was kept between the heated and the reference probe. In the




Figure 1. (a) Surface cover of the study area in Helsinki (StromJan, 2020) with the location of the monitoring sites, and surface covers at the (b) Park, (c) Street, (d) Forest, and (e) Orchard measurement sites.

Street and Forest sites, the sensors were installed higher, at 2 m, in order to avoid damage or disturbance [from pedestrians](#). The sensors were installed on the northern side of the stem. To protect and minimize the effect of the thermal gradient, the sensors were insulated with reflective aluminum foil after installation. The upper probe was heated and the temperature difference (dT) was recorded every 1 min using a datalogger (Datataker DT80M). The sap flux density (J_s , $\text{g cm}^{-2}\text{h}^{-1}$) was calculated at 1 min interval based on Granier's equation (eqn 1):

$$J_s = 42.84 * ((dT_{max} - dT)/dT)^{1.231}, \quad (1)$$

where dT_{max} is the maximum dT where zero J_s was observed. Zero flux condition was based on Lu et al. (2004) where it was derived as the average of local daily maximum dT of seven consecutive nights. Processing of raw sap flux data was conducted in R (RStudio Team, 2020). Further, daily tree water use was calculated by multiplying the daily sap flux density with the sapwood area. For *Tilia cordata*, *Tilia × europaea* and *Betula pendula*, the sapwood area was derived from the literature using species-specific diameter relationship (Gebauer et al., 2008; Hernandez-Santana et al., 2015) and for Orchard, we calculated

the sapwood area by coring the stem of the apple trees. For further analysis, the sap flux density data were selected only for the growing season (June to September) for both 2020 and 2021. To compare the sap flux density of the four sites, only sunny days data were considered. Sunny days were selected based on the criteria where the daily total R_g was greater than 200 $W m^{-2}$, there was no precipitation, and the mean daytime vapor pressure deficit (VPD) was greater than 0.33 kPa (Riikonen et al., 2016). Sap flux data were averaged to half-hourly, daily and monthly values for each site. In addition, we normalized the sap flux density using VPD by simply dividing the halfhourly J_s by the corresponding halfhourly VPD data. This is done in order to assess the effect of VPD and also to examine the dependency of J_s on other environmental variables during these heatwave and drought periods.~~In addition, we normalized the sap flux density by dividing J_s with VPD to examine the dependency of J_s on other environmental variables.~~

2.3 Leaf gas exchange

Leaf gas exchange was measured using a portable gas exchange system (Walz GFS-3000, Heinz Walz, GmbH, Germany) with a standard measuring head (8 cm^2 cuvette, 2x4 cm). At each site, leaf gas exchange was measured during the summers of 2020 and 2021 with approximately four-week-interval from the same trees as the sap flux was measured. However, at the Park site, one different tree was selected for leaf gas exchange measurements to replace one of the sap flow-equipped tree as it was difficult to reach the canopy with a manlift. The measurements were made on the southern or southwestern side of each tree and conducted mostly during local morning time (8 AM-12 PM). The measurements were performed on a healthy single leaf. At the Park site, measurements were made at three heights of the canopy (top, middle, and bottom) whereas, at the Street and Forest sites, only two heights at the top and bottom were monitored. At the Orchard site, only one measurement was made in the middle of the canopy of each tree.

During each measurement, the CO_2 level was set to ambient conditions i.e. 415 ppm but the temperature was not set to any value and was rather following the ambient conditions. In case the assimilation at the first 10 min was very low (under 1.5 $\mu mol m^{-2} s^{-1}$), a different leaf from the branch was selected and the same measurement steps were repeated. The steps involved setting photosynthetically active radiation (PAR) at 1200 $\mu mol m^{-2} s^{-1}$ for 12 minutes and then increasing to a level of 1500 $\mu mol m^{-2} s^{-1}$. After reaching the maximum level, PAR gradually decreased down to $<1 \mu mol m^{-2} s^{-1}$ over a period of 43 minutes. Altogether, once set, 15 different PAR intensities were included. A simple light response curve was fitted to the net CO_2 exchange ($NE(PAR)$, $\mu mol m^{-2} s^{-1}$), as follows:

$$NE(PAR) = (A_{max} * PAR) / (\beta + PAR) - R, \quad (2)$$

where R is the plant respiration i.e. NE measured at $PAR = 0$ ($\mu mol m^{-2} s^{-1}$), A_{max} is the maximum rate of photosynthesis ($\mu mol m^{-2} s^{-1}$) and β ($m^{-2} s^{-1}$) is the half-saturation constant describing the light intensity where photosynthesis is half of the rate of A_{max} .

From the above fitting, only A_{max} is considered in our analysis. Other variables of leaf gas exchange, namely stomatal conductance (g_s , $mmol m^{-2} s^{-1}$) and transpiration (E , $mmol m^{-2} s^{-1}$) were also recorded during the measurements. Maximum stomatal conductance and transpiration were calculated based on momentary g_s and E at $PAR = 1100 \mu mol m^{-2} s^{-1}$.

During the manual measurement campaigns in the summer of 2021, three leaf samples per site were also collected monthly in order to measure their relative water content (RWC). The samples were collected during the late afternoon (4 PM) and the fresh weight (FW) was measured. After that, the leaf samples were soaked overnight and turgid weight (TW) was measured. Later on, the samples were oven dried at 60 °C for 24h and the dry weight (DW) was measured. The RWC was calculated based on the equation below:

$$\text{RWC}(\%) = ((\text{FW} - \text{DW}) / (\text{TW} - \text{DW})) * 100, \quad (3)$$

2.4 Meteorological and soil data

At all four sites, meteorological variables, including air temperature (Air T, °C) and relative humidity (RH, %) were measured at a height of 1.5-1.8 m with a weather sensor (HMP110, Vaisala, Vantaa, Finland, at Park, Street and Forest sites; HC2A, Rotronic, Bassersdorf, Germany, at Orchard site). Soil sensors (Hydra-probe 2 SDI-12, Stevens, Oregon, USA, except ML3 ThetaProbe, Delta-T, Cambridge, UK, sensors in the Orchard) were installed at 10 and 30 cm depth to measure soil temperature (Soil T, °C) and soil moisture (SM m³ m⁻³). Data were recorded continuously at 1 min intervals and then converted into half-hourly averaged data. Furthermore, the vapor pressure deficit (VPD, kPa) was calculated using Air T and RH based on saturated vapor pressure. Photosynthetically active radiation (PAR, W m⁻²) and precipitation data were collected from the SMEAR III station FI-Kmp measurement tower and roof of a nearby building, respectively (Vesala et al., 2008; Järvi et al., 2009).

2.5 Data and statistical analysis

According to Fischer and Schär (2010), the heatwave is defined as a spell of at least six consecutive days with maximum temperatures exceeding the local 90th percentile of the control period. In our study, heatwave (Appendix A1) was defined as the period of consecutive of at least 6 days when the local daily maximum air temperature of the year was exceeding the daily maximum air temperature of the control period (1991-2020). Here, we have considered the control period as an average of daily maximum temperature for the last 30 years. Accordingly, our study period was categorized into heatwave (21.9 °C; 17 June 2021 to 18 July 2021), pre-heatwave (16.8 °C; 1 June 2021 to 16 June 2021), post-heatwave (16.8 °C; 19 July 2021 to 31 August 2021) and no heatwave (17.5 °C; 1 July 2020 to 31 July 2020) periods. Accordingly, our study period was categorized into heatwave (17 June 2021 to 18 July 2021), pre-heatwave (1 June 2021 to 16 June 2021), post-heatwave (19 July 2021 to 31 August 2021) and no heatwave (1 July 2020 to 31 July 2020) periods. To determine the drought period, monthly Standardised Precipitation-Evapotranspiration Index (SPEI, (Vicente-Serrano et al., 2010) was calculated and found out that June (SPEI = -0.7) and July (SPEI = -0.3) 2021 had mild drought conditions. According to the particle size distribution, all other sites were determined as sandy loams except for the Orchard which was clay. According to Hagemann and Stacke (2015), the wilting point and field capacity of sandy loam is 10 % and 22.9 %, respectively and of clay is 25 % and 38.4 %, respectively. We calculated the relative extractable soil water (REW) from soil moisture data, field capacity and wilting point of the site according to Granier et al. (1999). We considered days with precipitation less than 1 mm and mean REW in the depth of 10cm less than 0.45 as a dry period for all sites. As a result, the dry period was (22 June 2021 to 27 July 2021) and wet period (28

July 2021 to 31 August 2021). To determine the drought period, Standardised Precipitation-Evapotranspiration Index (SPEI, (Vicente-Serrano et al., 2010) was calculated. Based on the index, June (SPEI = -0.7) and July (SPEI = -0.3) 2021 had mild drought conditions. We separated the measurement period further into dry (22 June 2021 to 27 July 2021) and wet periods (28 July 2021 to 31 August 2021) based on soil moisture ($0.17 \text{ m}^3 \text{ m}^{-3}$) and daily precipitation (dry period has precipitation less than 1 mm).

To test the hypotheses, Kruskal Wallis test followed by Dunn's posthoc test was performed to examine differences in J_s between the sites and different climatic periods (dry/wet/heatwave/post-heatwave/no heatwave). First, polynomial regression with 2nd order degree was fitted between daytime mean J_s and daytime daily VPD. Second, multiple linear regression analysis was performed between the J_s and different meteorological variables such as VPD, PAR, soil moisture and soil temperature at 30 cm to determine the effect of the meteorological variables and the relative importance of these meteorological variables in controlling the daily J_s was assessed using Student's t-test (t) between the individual meteorological and daily J_s . Data processing (post-processing of sap flux, meteorological and gas exchange data, statistical analysis) and visualization were conducted in Python (Python version 3).

3 Results

3.1 Weather conditions

The summer of 2021 (particularly July) was warm and dry, as compared to the summer of 2020, and the climatic reference period (1991 - 2020). The mean air temperature in July 2021 was 21.6°C , being 21% higher than in July 2020 (16.7°C) and 19% higher than the average July during the climatic reference period (18.1°C) (Figure 2a). At the four urban sites, high air temperature ($20.2\text{-}21.6^\circ\text{C}$), high soil temperature ($15.7\text{-}18.5^\circ\text{C}$), high VPD (0.9 - 1.1 kPa), and low soil moisture ($0.1\text{-}0.4 \text{ m}^3 \text{ m}^{-3}$) were observed during July 2021 (Figure 3 for Park site, Appendices A2, A3 and A4 for other sites). The total precipitation for the months of June and July in 2021 (86 mm) was 51% lower than during June and July 2020 (177 mm) and 27% lower than on average for the climatic reference period (117 mm) (Figure 2b).

The climatic conditions varied across the four sites during summer 2021. The mean of the meteorological variables (air temperature, VPD, soil temperature and soil moisture) during the summer months (June, July, August) of 2021 were considered for comparison. Higher (8%) mean air temperature was observed at both Orchard (19.2°C) and Street sites (19.3°C) as compared to Forest (17.9°C) and Park (17.8°C) sites (Table 2). Similar mean VPD was found at the Orchard (0.78 kPa), Park (0.75 kPa) and Forest sites (0.75 kPa), but 11% higher mean VPD at the Street site (0.84 kPa). Also, the mean soil temperature at the Street site (19.9°C) was 24-34% higher than at the Park (16.1°C), Orchard (15.2°C) and Forest sites (14.8°C). Mean soil moisture content varied largely as the Orchard site had the highest ($0.37 \text{ m}^3 \text{ m}^{-3}$) and the Forest site the lowest ($0.09 \text{ m}^3 \text{ m}^{-3}$) whereas Park and Street sites had 0.13 and $0.22 \text{ m}^3 \text{ m}^{-3}$ respectively. Soil moisture conditions were recovered after the rainfall at all the sites except the Forest site, where soil moisture remained low throughout the late summer after the hot and dry July 2021 (Appendix A3).




Figure 2. (a) Monthly mean air temperature and (b) monthly total precipitation for the year 2020 and 2021, and the climatic reference period of 30 years (1991-2020).

250 3.2 Variability in [sap flow rate](#) tree water-use in 2021

Based on sunny days, the mean [sap flow rate](#) daily water-use of the trees in the Park, Street, Forest and Orchard were $0.32 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$, $0.42 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$, $0.20 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$ and $0.46 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$, respectively and differed significantly between the sites (Figure 4a) ($P < 0.05$). The mean daytime sap flux density (J_s) for the summer period (June-August) was highest at the Orchard site with $20.6 \pm 0.3 \text{ g cm}^{-2} \text{ h}^{-1}$, lowest with $8.1 \pm 0.1 \text{ g cm}^{-2} \text{ h}^{-1}$ at the Forest site, and $14.4 \pm 0.2 \text{ g cm}^{-2} \text{ h}^{-1}$ and $17.7 \pm 0.3 \text{ g cm}^{-2} \text{ h}^{-1}$ at the Park and Street sites, respectively. The monthly mean J_s differed significantly at the four sites ($P < 0.05$) (Figure 4b).




Figure 3. Meteorological and soil data from 2021 showing hourly a) air temperature (Air T), b) water vapor deficit (VPD), c) soil temperature (Soil T) and d) soil moisture measured at the Park site, and e) daily mean Photosynthetically active radiation (PAR) and daily sum precipitation data measured at the SMEARIII station. The orange shading indicates the heatwave period during the summer of 2021 and the black vertical line indicates the onset of the wet period. The yellow markers in panel (c) denote the dates of manual leaf gas measurements.

3.3 The effect of heatwave on sap flux density

During heatwave period, the mean [sap flow rate](#)[daily water use](#) of the trees at the Park, Street, Forest and Orchard sites were $0.38 \pm 0.02 \text{ kg cm}^{-2} \text{ day}^{-1}$, $0.42 \pm 0.02 \text{ kg cm}^{-2} \text{ day}^{-1}$, $0.24 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$ and $0.52 \pm 0.02 \text{ kg cm}^{-2} \text{ day}^{-1}$,

Table 2. Monthly mean air temperature (Air T, °C), [mean](#) vapor pressure deficit (VPD, kPa), [mean](#) soil temperature (Soil T, °C) and [mean](#) soil moisture [content](#) ($\text{m}^3 \text{m}^{-3}$) for the four study sites in 2021.

		Air T	VPD	Soil T	Soil moisture
Park	June	18.4	0.84	14.4	0.16
	July	20.2	0.96	17.5	0.09
	August	14.8	0.45	14.7	0.15
	September	8.7	0.40	11.0	0.19
Street	June	19.8	0.93	18.5	0.23
	July	21.6	1.07	22.8	0.25
	August	16.1	0.51	18.5	0.25
	September	10.0	0.40	12.7	0.29
Forest	June	18.0	0.77	13.1	0.16
	July	20.7	1.01	16.9	0.07
	August	15.2	0.48	14.5	0.06
	September	8.7	0.34	11.2	0.06
Orchard	June	21.1	0.81	15.5	0.44
	July	21.1	1.05	15.7	0.28
	August	15.7	0.49	14.6	0.40
	September	9.4	0.35	12.9	0.49

260 respectively. At the Park site, the mean J_s during heatwave period was 59% higher than during no heatwave period and 39%
higher than during post-heatwave period but there was no significant difference with the pre-heatwave period. At the Street site,
there was no significant difference in the mean J_s between heatwave, no heatwave, pre-heatwave and post-heatwave periods.
At the Forest site, the mean J_s during heatwave period was 13% higher than during pre-heatwave and 67% higher than during
post-heatwave periods. At the Orchard site, the mean J_s during heatwave period was 35% higher than during post-heatwave
265 period but there was no significant difference with pre-heatwave period (Figure 5a, $P < 0.05$). At the Forest and Orchard sites,
data from no heatwave period were not available.

When normalized with VPD, there were no significant differences in the J_s between heatwave, no heatwave and pre-
heatwave and post-heatwave periods in the Park, Forest or Orchard (Figure 5b). At the Street site, the normalized J_s during
heatwave period was 33% lower than during no heatwave and 7% lower than during pre-heatwave but it did not differ from
270 that of post-heatwave period.

3.4 The effect of drought on sap flux density

During dry period, the mean [sap flow ratedaily-water-use](#) at the Park, Street, Forest and Orchard site was $0.36 \pm 0.02 \text{ kg cm}^{-2}$
 day^{-1} , $0.42 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$, $0.20 \pm 0.01 \text{ kg cm}^{-2} \text{ day}^{-1}$ and $0.50 \pm 0.02 \text{ kg cm}^{-2} \text{ day}^{-1}$, respectively. The mean J_s




Figure 4. (a) Daily water use (DWU) of the trees and (b) monthly mean sap flux density (J_s) at the four urban vegetation sites: Park (*Tilia cordata*), Street (*Tilia × europaea*), Forest (*Betula pendula*) and Orchard (*Malus spp.*). The letters indicate the significant differences between the (a) sites and (b) monthly mean J_s at each site ($P < 0.05$). [Note: there was no data available for August month at the Street site due to technical and sensor problems.](#)

was significantly higher during dry period than during wet period at all sites (Figure 6a, $P < 0.05$). At the Park, Street, Forest and Orchard sites, the mean J_s was 66%, 31%, 43% and 53%, respectively, higher during dry period than wet period.

The normalized J_s was significantly lower during dry than wet period at all sites, with 16% lower at the Park site, 48% lower at the Street site, 28% lower at the Forest site and 26% lower at the Orchard site (Figure 6b).

[We separated the period into three periods: only heat, only dry and both heat-dry by intersection of the heatwave and dry days which were defined earlier \(as describe in section 2.5\) and analysed the effect of only heat, only dry and both heat-dry on sap flux density and found out that the effect of heat was relatively higher than the effect of dry and both heat and dry at Park, Forest and Orchard while at the Street site, the effect of heat, dry and both are similar \(Appendix A6\).](#)

3.5 Leaf gas exchange during the heatwave and drought periods

We compared leaf gas exchange variables, namely A_{max} , g_s and E , between the different heatwave periods and between dry and wet periods (Table 3). No significant differences in these three variables were found at the Park, Forest and Orchard sites between the different heatwave periods; however, at the Street site, A_{max} and E were significantly ($P < 0.05$) higher during

Table 3. Average of leaf gas exchange variables: Maximum assimilation (A_{max} , $\mu\text{mol m}^{-2} \text{s}^{-1}$), Stomatal Conductance (g_s , $\text{mmol m}^{-2} \text{s}^{-1}$), Transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$) during heatwave, no heatwave, pre-heatwave and post-heatwave periods and during dry and wet periods at the four sites. The letters indicate the significant difference between the various heatwave periods or drought periods. [No data available at the Street site during pre-heatwave and at the Orchard site during no heatwave period.](#)

Site	Type / Period	A_{max}	g_s	E
Park	heatwave	13.6 ± 1.3	112.1 ± 14.1	1.5 ± 0.1
	pre-heatwave	13.6 ± 1.3	123.5 ± 0	1.9 ± 0
	post-heatwave	15.3 ± 1.1	141.3 ± 11	1.1 ± 0.1
	no heatwave	17.2 ± 1.4	147.8 ± 15.5	1.5 ± 0.1
	dry	15.0 ± 0.8	114.5 ± 7.7^a	1.5 ± 0.1
	wet	14.3 ± 2.0	163.3 ± 17.5^b	1.7 ± 0.2
Street	heatwave	10.9 ± 1.0^a	91.4 ± 7.3	1.2 ± 0.1^a
	pre-heatwave	-	-	-
	post-heatwave	6.6 ± 0.8^b	64.2 ± 12.0	0.8 ± 0.1^b
	no heatwave	8.7 ± 1.2^a	73.0 ± 11.1	0.8 ± 0.1^a
	dry	10.9 ± 1.0^a	91.4 ± 7.3	1.2 ± 0.1
	wet	6.6 ± 0.8^b	64.2 ± 12.0	0.8 ± 0.1
Forest	heatwave	16.4 ± 1.0	105.7 ± 17.8	1.3 ± 0.2
	pre-heatwave	16.3 ± 0	128.4 ± 0	1.5 ± 0
	post-heatwave	10.7 ± 2.9	104.2 ± 31.2	1.1 ± 0.3
	no heatwave	17.4 ± 2.3	133.5 ± 12.1	1.4 ± 0.1
	dry	16.4 ± 1.0	105.7 ± 17.8	1.3 ± 0.2
	wet	10.7 ± 2.9	104.2 ± 31.2	1.1 ± 0.3
Orchard	heatwave	14.1 ± 0.7	126.8 ± 5.4	0.7 ± 0.2
	pre-heatwave	13.3 ± 0	125.3 ± 0	2.0 ± 0
	post-heatwave	13.6 ± 1.0	157.3 ± 12.9	1.8 ± 0.1
	no heatwave	-	-	-
	dry	13.1 ± 1.4	135.4 ± 13.6	1.6 ± 0.1
	wet	14.0 ± 0.9	170.9 ± 20.4	1.9 ± 0.2




Figure 5. The sap flux densities at the four urban vegetation sites during different periods regarding a heatwave. a) mean J_s of the whole day (b) mean normalized J_s by VPD on sunny days. The letters indicate the significant differences between the different periods within each site ($P < 0.05$).

heatwave than post-heatwave periods but there was no significant difference as compared to no heatwave period. Also, g_s showed no difference between the different heatwave periods at the Street site.

Comparing leaf gas exchange variables between dry and wet periods, we found that at the Street site, A_{max} was significantly higher ($P < 0.05$) during dry period than wet period but no significant differences in g_s and E between dry and wet periods were found. Also, at the Park site, g_s was found to be significantly lower ($P < 0.05$) during dry period than wet period. At the Forest and Orchard sites, no significant differences were found in A_{max} , g_s and E between dry and wet periods.

The monthly relative leaf water content (RWC) as a proxy of leaf water potential showed that RWC was found to be lower (4-35 %) during July as compared to June and August at the Forest and Orchard sites; however, RWC was found to be higher (5-8 %) during July than the other summer months at Park and Street sites (Appendix A5).




Figure 6. Sap flux density (J_s) during dry and wet periods at the four study sites. Based on sunny days, (a) data represented mean J_s of the whole day and (b) normalized J_s by VPD. The letters indicate the significant differences J_s between the dry and wet at each site ($P < 0.05$)

295 3.6 Environmental control on sap flux density

We tested the relationship between the daily daytime mean VPD and J_s using 2nd order polynomial regression (Figure 7, Table 4). VPD explained the variation in J_s less during the heatwave than during post-heatwave and pre-heatwave except for Forest where the VPD was not a significant driver at all during pre-heatwave. During wet period, VPD explained a higher share of the variation of J_s except for Street where the wet data coverage is very low (Fig. 7).

300 Multiple linear regression between J_s and a higher number of environmental variables (VPD, PAR, Soil T and SM) showed that PAR and VPD were the significant drivers for J_s at the Park, Street and Orchard site but PAR was the only significant driver at the Forest site (Table 5). In addition, soil moisture was a significant variable in certain circumstances at certain sites such as in Park during heatwave and dry periods and in Forest, during post-heatwave, dry and wet periods i.e. in the latter part of the growing season in general.




Figure 7. Observed (dots) and modeled (lines) relationship between the daytime mean daily VPD and J_s during the heatwave periods and both dry and wet periods. Panel (a),(b),(c),(d) show the relationship between VPD and J_s at the Park, Street, Forest and Orchard site during the various heatwave periods. Panel (e),(f),(g),(h) show the relationship between VPD and J_s at the Park, Street, Forest and Orchard site during the dry and wet periods. The model is 2nd order polynomial fit (see Table 4).

Table 4. Relationship between vapor pressure deficit and daily mean sap flux density using 2nd order polynomial fit (see Fig. 7). The values are showing the squared coefficient of correlation (adj R^2) between the two variables. Relationships statistically significant at 0.05, 0.01 and 0.001 levels are marked with *, ** and *** respectively and non-significant as ns.

	no heatwave	pre-heatwave	heatwave	post-heatwave	dry	wet
Park	0.53***	0.73***	0.57***	0.90***	0.60***	0.85***
Street	0.80***	0.62***	0.55***	0.75***	0.58***	0.22 ^{ns}
Forest	-	0.16 ^{ns}	0.43***	0.44***	0.25**	0.48***
Orchard	-	0.77***	0.69***	0.91***	0.67***	0.90***

305 4 Discussion

In this study, we assessed the response of urban tree water use and leaf gas exchange to heat and drought during two contrasting summers 2020 and 2021, the latter being hot and dry, across four urban green areas in Helsinki. The results showed that tree water use, measured with sap flux density increased during the heatwave and the dry period but carbon assimilation, stomatal conductance and transpiration at leaf level did not generally change during these various periods. The increase in sap flux density during hot and dry periods varied across the urban sites that had different tree species and growing conditions. VPD

310

Table 5. Multiple linear regression between daily mean sap flux density and various environmental drivers (VPD, PAR, Soil temperature and Soil moisture) as independent variables. t stat value indicates the relative importance of the variables in controlling the daily sap flux variations. Relationships statistically significant at 0.05, 0.01 and 0.001 levels are marked with *, ** and *** respectively and non-significant as ns.

Site		All data	no heatwave	pre-heatwave	heatwave	post-heatwave	dry period	wet period
Park	R ²	0.63	0.59	0.75	0.74	0.93	0.74	0.92
	Intercept	5 (1.4) ^{ns}	-1.6 (-0.2) ^{ns}	-16.6 (-0.9) ^{ns}	-25.15 (-1.6) ^{ns}	4.1 (1.1) ^{ns}	-13.9(-1.1) ^{ns}	3.0 (0.7) ^{ns}
	VPD	9 (7.6)***	6.2 (3.1)***	9.1 (3.5)***	10.4 (5.1)***	9.3 (8.0)***	9.5 (5.2)***	11.0 (5.7) ^{ns}
	PAR	0.01 (3.5)***	0.01 (1.14) ^{ns}	0.01 (1.4) ^{ns}	0.03 (3.1)***	0.02 (5.4)***	0.02 (3.6)***	0.01 (4.1)**
	Soil T	-0.1 (-0.3) ^{ns}	0.5 (1.1) ^{ns}	1 (1.4) ^{ns}	0.8 (1.32) ^{ns}	-0.1 (-0.6) ^{ns}	0.4 (0.7) ^{ns}	0.03 (0.1) ^{ns}
	SM	-9.4 (-1.0) ^{ns}	-10.5 (-0.4) ^{ns}	45.6 (1.0) ^{ns}	79.6 (2.3)***	-8.0 (-1.0) ^{ns}	48.1 (2.1)***	-15.9 (-2.2) ^{ns}
Street	R ²	0.73	0.84	0.65	0.71	0.75	0.71	-
	Intercept	5.9 (2.27)*	6.2 (0.3) ^{ns}	7.0 (1.0) ^{ns}	10.6 (1.1) ^{ns}	40.7 (1.3) ^{ns}	22.5 (2.0) ^{ns}	-
	VPD	9.3 (7.8)**	17.5 (6.4)***	7.6 (2.5)*	4.6 (2.6)*	6.1 (1.4) ^{ns}	5.9 (3.4)***	-
	PAR	0.03 (7.1)**	0.02 (2.5)*	0.02 (1.7) ^{ns}	0.03 (4.4)***	0.01 (1.0) ^{ns}	0.03 (4.0)***	-
	Soil T	-0.3 (-3.3)**	-0.5 (-0.9) ^{ns}	0.07 (0.2) ^{ns}	-0.1 (-0.5) ^{ns}	-1.1 (-1.2) ^{ns}	-0.7 (-1.6) ^{ns}	-
	SM	12.6 (2.4)*	12.0 (0.2) ^{ns}	-9.7 (-0.7) ^{ns}	-13.9 (-1.0) ^{ns}	-34.5(-0.5) ^{ns}	-8.3 (-0.8) ^{ns}	-
Forest	R ²	0.56	-	0.93	0.67	0.68	0.63	0.72
	Intercept	2.4 (1.4) ^{ns}	-	91.0 (2.4) ^{ns}	-2.3 (-0.1) ^{ns}	-16.2(-1.8) ^{ns}	-17.7(-2.1)*	-17.3(-1.5) ^{ns}
	VPD	2.1 (2.0) ^{ns}	-	2.9 (1.5) ^{ns}	3.5 (1.7) ^{ns}	-0.3 (-0.3) ^{ns}	2.9(1.9) ^{ns}	0.1(0.03) ^{ns}
	PAR	0.02 (5.1)***	-	0.01 (1.3) ^{ns}	0.02 (3.4)***	0.02(4.99)***	0.02(3.6)***	0.02(3.5)***
	Soil T	-0.1 (-1.1) ^{ns}	-	-1.0 (-0.9) ^{ns}	-0.6 (-0.8) ^{ns}	-0.8 (-4.4)***	0.01(0.03) ^{ns}	-0.8(-3.4)**
	SM	-1.4 (-0.2) ^{ns}	-	-345.0 (-3.1)*	137.7 (1.1) ^{ns}	515.0(3.6)***	222.1(5.0)***	515.2(2.8)**
Orchard	R ²	0.89	-	0.86	0.77	0.93	0.74	0.92
	Intercept	9.8 (1.7) ^{ns}	-	-88.2 (-2.3) ^{ns}	57.9 (0.8) ^{ns}	27.0 (2.2)*	-31.2(-1.4) ^{ns}	3.1 (0.8) ^{ns}
	VPD	9.2 (8.0)***	-	5.9 (1.9) ^{ns}	5.4 (2.1) ^{ns}	7.8 (5.4)***	7.1 (3.4)**	11 (5.8)***
	PAR	0.04 (8.2)***	-	0.03 (2.9)*	0.05 (4.5)***	0.04 (7.6)***	0.05 (4.9)***	0.02 (4.07)***
	Soil T	-0.5 (-1.5) ^{ns}	-	0.6 (0.5) ^{ns}	-2.8 (-0.7) ^{ns}	-1.6 (-2.1)*	2.1 (1.6) ^{ns}	0.03 (0.1) ^{ns}
	SM	1.5 (0.4) ^{ns}	-	160.3 (3.2)*	-32.3 (-1.5) ^{ns}	-2.9 (-0.8) ^{ns}	-1.6 (-0.2) ^{ns}	-15.9(-2.2)***

Values indicate coefficient (*t*)^{sig}

was the main driver of tree water use during the heatwave and dry. ~~However, the VPD explained less of the variation in the transpiration during the heatwave and dry period as compared to other periods; giving support to our hypothesis H3.~~ Overall, it seems that the hot and dry conditions were not severe enough to trigger notable physiological adaptation as the urban trees in our study continued to function typically during the summer of 2021. Even though the air temperature was notably higher and precipitation notably lower than during previous summers, we did not get full support for the hypotheses H1 and H2 and conclude that the severe weather events did not alter the stomatal action and therefore the observed photosynthetic potential. However, we found some interesting insights that we will discuss further in this chapter.

4.1 Site variability

In our study, we observed that the four urban vegetation sites in Helsinki exhibit variable microclimatic conditions where air temperature under the canopy, soil temperature and soil moisture content were different. The highest air and soil temperatures were measured at the Street site, where impervious surfaces increase the temperatures due to heat storage. The high air temperature at the Orchard site is due to exposure of the site to direct sunlight throughout the day heating the garden. Also, the high soil moisture at the Orchard site is mainly due to the difference in soil type where the sand clay in Orchard has higher water holding capacity than sand moraine soil type at the other three sites. ~~At the Street site, VPD was clearly higher as compared to the other three sites and this is likely due to the larger cover of impervious surface at the Street site where the air temperature is marginally increased. VPD was similar at all other sites except the Street site, where it was clearly higher. We speculate that the highest deficit at the Street site is again due to the large fraction of impervious surfaces (Whitlow et al., 1992; Kjelgren and Montague, 1998).~~ Similar variability of meteorological conditions between different urban forests were found in Los Angeles metropolitan city; where the urban forest located near the city were warmer with high VPD and lower photosynthetically active radiation as compared to the urban forests located closer to the coast (Pataki et al., 2011). Among 10 different tree species in the city of Basel (Switzerland), the tree crown temperature was lower in the park than the street; reporting that it is species-specific for the cooling effect of urban trees (Leuzinger et al., 2010). The difference in microclimatic conditions has been observed to vary depending on the type of vegetation, the composition of the species, the amount of green cover and the impervious surface in urban vegetation (Perini and Magliocco, 2014; Kjelgren and Clark, 1992). ~~We observed that sap flow rate tree water use varied significantly across the four urban sites; Orchard and Street sites had the highest sap flow rate tree water use during the summer of 2021 and it was the lowest at the Forest site. The higher sap flow rate water use at the Orchard and Street sites was mainly due to site conditions such as high soil moisture content and tree characteristics such as bigger stem size. Water availability at the Street and Orchard ($0.23-0.49 \text{ m}^3 \text{ m}^{-3}$) was higher as compared to the Park and Forest ($0.06-0.19 \text{ m}^3 \text{ m}^{-3}$).~~ Many studies have shown high variability in transpiration tree water use across different urban green areas (Pataki et al., 2011; McCarthy and Pataki, 2010; Sushko et al., 2021). For example in the high-latitude city of Gothenburg (Sweden), *T. europaea* had two times higher daytime transpiration rates in a park compared to a street site (Konarska et al., 2016). ~~Although, the assymmetric measurement setup in our study limits us to compare the sites statistically but we observed that Street sites had the highest sap flow rate during the summer of 2021 and the lowest at the Forest site and we speculate that the differences might be related to growing conditions such as high soil moisture content and~~

345 ~~tree species. The found differences in sap flow rate tree water use between the sites in our study can also be related to different~~
~~tree species growing at the sites.~~ The sap flow rate of *Tilia cordata* at the Park was lower than that of *Malus spp* at the Orchard
and *Tilia × europaea* at the Street but higher than that of *Betula pendula* at the Forest site. The low ~~sap flow rate tree water use~~
of *Betula pendula* at the Forest site could be reduced due to the rather strong stomatal control typical for *Betula pendula*, i.e.
it closes the stomata easily during dry conditions (Zapater et al., 2013), whereas *Tilia cordata* found at the Park site has less
350 sensitive stomatal control (Leuschner et al., 2019), i.e. it keeps stomata open even in mild drought as it can tolerate drought
better than *Betula Pendula*. ~~Other studies in the streets of Munich and Helsinki have reported variability of transpiration rates,~~
~~mainly due to the differences tree species. In Munich, the transpiration of Tilia cordata Mill. were three times higher than~~
~~water use of Robinia pseudoacacia L. tree in the (Rahman et al., 2019) and in the street trees of Helsinki, Alnus glutinosa have~~
~~four times higher tree water use than Tilia × vulgaris (Riikonen et al., 2016)~~ Differences in water use between different urban
355 ~~tree species have also been reported before. Transpiration rate tree water use of Tilia cordata Mill. were reported to be three~~
~~times higher than water use of Robinia pseudoacacia L. tree in the streets of Munich, Germany (Rahman et al., 2019) and~~
~~water use of Tilia × vulgaris were found to be 1/4 lower than water use of Alnus glutinosa in the streets of Helsinki, Finland~~
~~(Riikonen et al., 2016). In general, the daily tree water use found in this latter study was lower (in the years 2008-2011) as~~
~~compared to the urban trees we found at the four studied sites in Helsinki. Overall, the transpiration rate tree water use across~~
360 ~~different urban sites varies not only due to site and climatic conditions but also due to the planted tree species.~~

4.2 Transpiration rate and leaf gas exchange during drought and heatwave

We observed varying responses of ~~sap flux density tree water use~~ during heatwave and drought periods at the studied sites. At
the Park, Forest and Orchard sites, J_s increased by 35-67% during the heatwave compared to periods of no heatwave and/or
post-heatwave, whereas the heatwave did not affect J_s in the Street site. Pre-heatwave period did not differ from the heatwave
365 period in terms of J_s in the Park and Orchard sites, but there was a small increase (13%) at the Forest site during the heatwave
period. During the dry period, J_s was significantly higher than during wet period at all sites. The leaf gas exchanges such as
 A_{max} , g_s and E did not change or reduce during the heatwave and dry period; thus, indicating no changes in the photosynthetic
potential during these periods. Hence, we conclude that the weather was yet not severe enough to support the hypotheses H1
and H2 in our study.

370 VPD is the driving force for transpiration, so an increase in VPD leads to an increase in transpiration ~~and tree water use~~
unless stomata in the leaves close to limit transpiration. ~~The ratio between J_s and VPD was significantly reduced during all~~
~~periods and at all sites, which indicates the substantial role of VPD; however, the relative importance of VPD over daily sap~~
~~flow variation differed at the four sites and during different periods of heatwave and drought. The response of sap flow with~~
~~VPD was less sensitive during heatwave and drought periods as compared to other heatwave periods and wet period (Table~~
375 ~~4). The ratio between J_s and VPD significantly decreased due to drought and the relative importance of VPD explaining the~~
~~observed variation in sap flow was lower in dry and heatwave conditions compared to other periods; giving support to our~~
~~hypothesis H3.~~ In theory, these results could indicate, that the trees limited their water transport via stomatal control in harsh
conditions. However, that was not clearly captured by the leaf-level measurements which on the other hand, may not represent

the conditions over the different periods as well as these automatic measurements. Moreover, the leaf-level measurements were
380 based on single-day measurements, which might not fully cover the heatwave or drought periods in our study. Together these
results indicate that the observed increase in transpiration at the studied sites was caused by an increase in the driving force
for transpiration, VPD. ~~Similar responses were also observed in the city of Dresden, Germany where the transpiration rate
and maximum stomatal conductance of *Tilia cordata* and *Corylus colurna* were largest during dry summer months in 2013
(Gillner et al., 2015a).~~

385 Stomatal control limits plant transpiration during drought. For example, Rötzer et al. (2021) found a substantial reduction
of 63 % in the transpiration rate in urban *Tilia cordata* and *Robinia pseudoacacia* trees in the city of Würzburg, Germany
during the European drought in 2018. During the dry period of this study, soil moisture decreased by 18-62% at the four sites
as compared to wet period. During heatwave period, soil moisture decreased by 30-58% as compared with the pre-heatwave
period at all sites except for the Street site. The availability of soil moisture at the Street site allowed an increase in A_{max}
390 and E during the heatwave period. However, the observed reductions in soil moisture did not seem to be enough to cause
strong stomatal regulation of transpiration (i.e. no change in g_s except for the Park site and no reduction in J_s). Interestingly, it
was previously reported that the non-stomatal origin limitation was responsible for reductions of photosynthesis in temperate
forest during European drought 2018 (Gourlez de la Motte et al., 2020) and these non-stomatal effects might play some role
in our urban site but yet to study further, and it may be also due to tree species behaviours (isohydril or anisohydril) towards
395 the drought response. Previous studies have also reported that transpiration during the local extreme high temperature was
maintained when there was sufficient water availability in the soil in different urban green sites in Los Angeles Metropolitan,
US (Pataki et al., 2011). Similarly, previous studies have shown that VPD is a significant driver for sap flow in *Tilia × vulgaris*
street trees in Helsinki (Riikonen et al., 2016), and VPD and solar radiation for daytime transpiration rates in seven different
tree species, including *Betula pendula*, in Gothenburg, Sweden (Konarska et al., 2016). Similarly to our results regarding the
400 Park site, Konarska et al. (2016) also found that the maximum stomatal conductance was reduced by 50% in the studied species
in Gothenburg even though transpiration rate remained high during dry conditions compared to wet conditions.

During the heatwave and the dry period, VPD explained most of the variation in the daily mean J_s at the Park (57-60%),
Street (55-58%) and Orchard sites (62-69%). Similarly, previous studies in high latitude cities have reported that VPD correlates
well with J_s in street trees (adj. $R^2 = 0.74$) in Helsinki, Finland (Riikonen et al., 2016), and in urban trees ($R^2=0.44-0.75$) in
405 Gothenburg, Sweden (Konarska et al., 2016). Also in Boston, Massachusetts, VPD has been shown to correlate with J_s (R^2
= 0.63, (Winbourne et al., 2020)). However, VPD did not explain daily variation in J_s at the forest site of our study. Also,
 J_s saturated after reaching certain VPD levels, especially at Forest and Orchard sites, and the saturation took place already in
rather low VPD levels in the case of the Forest site compared to the Orchard site. The difference in saturation levels may be
species-specific or caused by differences in the availability of soil moisture as the Orchard site had higher soil moisture than
410 the Forest site during the heatwave and dry period. *Tilia* at the Park and Street sites seemed to be more linearly correlated with
VPD, ~~i.e. suggesting that~~ transpiration continues to increase with increasing VPD during the stressful periods ~~at these sites.~~
With sufficient water supply by irrigation, less saturation with VPD was observed in urban trees previously (Winbourne et al.,
2020; Marchin et al., 2022). Similarly, the non-saturation of J_s with high VPD observed at the Park site may be due to irrigation

during the dry period. Several previous studies in urban trees have shown that the relationship between VPD and transpiration
415 are species-specific and it is more typical that VPD increases linearly in diffuse-porous trees but saturates in ring-porous trees
(Bush et al., 2008; Rahman et al., 2019). All the species studied here are diffuse-porous.

In addition to VPD, J_s was also explained by other environmental variables. The relative importance of soil moisture, soil
temperature and solar radiation in explaining J_s differed significantly between the four studied urban sites. In addition to VPD,
PAR was among the main environmental drivers of J_s at Park, Orchard and Street sites, whereas PAR alone explained J_s at
420 the Forest site (Table 5). Soil temperature was significantly related with J_s only at the Street site. When the different climatic
periods were analyzed separately, soil conditions (moisture and/or temperature) affected J_s during heatwave (t stat = 2.3,
P <0.05) and dry periods (t stat = 2.1, P <0.05) at the Park site, but not at the Street, Forest and Orchard sites. This can
be explained by the irrigation provided at the Park site during the dry period. Overall, the environmental variables that best
explained urban tree transpiration during hot and dry conditions were VPD and PAR, and the relative importance of these two
425 varied depending on the tree species, growing conditions and irrigation practices.

The high J_s during the heatwave in the studied green areas suggests transpirational cooling. Trees at the Orchard and Street
sites had the highest transpiration rates and trees in the Forest had the lowest transpiration rates during the dry and heatwave
periods. These differences were mainly due to tree species, their drought strategies (Gillner et al., 2017) and growing conditions
of the sites, particularly soil moisture availability. Lower transpiration in *Betula pendula* at the Forest site indicates that *Betula*
430 *pendula* trees growing in an urban forest do not cool the environment as much as *Tilia* or *Malus*. Several previous studies have
reported that the transpirational cooling effect of urban trees during hot and dry days increases or sustains the transpiration
rates in order to prevent excessive heat accumulation (Gillner et al., 2015b; Duarte et al., 2016; Drake et al., 2018; Urban et al.,
2017; Ibsen et al., 2021). Keeping the stomata open in hot and dry conditions cools down the internal leaf temperatures enabling
maintaining photosynthesis (De Kauwe et al., 2019; Urban et al., 2017; Drake et al., 2018). The response depends on the species
435 tolerance to drought, water use efficiency, microclimatic conditions and site heterogeneity (Bussotti et al., 2014; Winbourne
et al., 2020; Rennenberg et al., 2006). Especially *Tilia cordata* is known for its anisohydric behavior (i.e. stomatal control is
not strong) during heat and drought and an associated increase in transpiration rates causing a cooling effect in different urban
conditions (Moser et al., 2017). However, also *Betula pendula*, which typically shows isohydric behavior (i.e. strong stomatal
control), increased J_s during the heatwave and dry periods in the present study. Contrasting and species-specific response of
440 trees to heat and drought have been observed in several urban trees (Gillner et al., 2017; Osone et al., 2014), however here, a
rather constant pattern of heat and drought responses was observed between the studied species.

In our study, we have observed these responses of urban tree transpiration and leaf gas exchanges pattern during heatwave
and dry period; however, there were challenges and limitation to evaluate about detail comparison of tree species response
at the four urban sites as the tree species are different in different site conditions. Also, the limited measurements of leaf
445 gas exchanges could not be addressed more about water use efficiency during these local extreme period. Further study with
complex model capturing the effect of site conditions and tree species behaviour separately would be useful in addressing the
main factor affecting the different responses of urban vegetation during the heatwave and dry period.

5 Conclusions

We conclude that the heat and drought that occurred in Helsinki in 2021 were still not extreme enough to damage or dampen the gas exchange functioning of urban trees. Against our hypotheses, photosynthetic potential did not reduce due to lowered stomatal conductance during heatwave and drought conditions as the transpiration and photosynthetic potential during these periods stayed high suggesting stable ecosystem services such as cooling and also carbon sequestration during rough conditions. However, the significant role of VPD during heatwave and drought periods was well supported in our study but its overall significance decreased during drought periods. The observed responses of varying tree transpiration during heat and drought across the four urban green areas are mainly due to VPD and further investigations would be needed to differentiate the role of other factors such as growing conditions (soil water availability), tree species and tree size with proper sampling strategies and complex models. ~~The observed responses of tree transpiration and leaf gas exchanges during heat and drought across the four urban green areas are mainly due to tree species type and VPD. Also, site-to-site variability in urban tree water use was also largely explained by differences in tree size, varying growing conditions, and soil moisture availability, apart from tree species type and VPD.~~ As urbanization and the occurrence of climate extreme events are rising, particularly in high-latitude regions, the role of urban green areas in mitigating climate change and cooling local microclimate is significant in cities. Further studies of the cooling potential of urban trees will provide a better understanding and support the mitigation and city planning in the future.

Data availability. Datasets of sap flux density, meteorological and leaf gas exchange measurements at the four urban sites in Helsinki is stored in <https://doi.org/10.5281/zenodo.7525319>

Appendix A

Table A1. The soil properties of the four urban vegetation sites. Soil sample analyzed from the top 30 cm of soil.

	Park	Street	Forest	Orchard
Soil type	Sand moraine	Fine sand moraine	Sand moraine	Sandy clay
Bulk density (kg/l)	1.15	1.07	1.14	1.02
Main particle size distribution	66% sand 21% silt 8% clay	48% sand, 26% silt, 11% clay	71% sand 15% silt 11% clay	27% sand, 31% silt, 42% clay
Carbon content (%)	3.7	3.3	3.9	3.9
Nitrogen content (%)	0.252	0.168	0.329	0.32
C:N ratio	14.8	21.8	11.9	12.3
pH	5.6	7.2	6.5	5.9
<u>Soil porosity</u>	<u>41.6 %</u>	<u>41.6 %</u>	<u>41.6 %</u>	<u>46.1 %</u>
<u>Field capacity</u>	<u>22.9 %</u>	<u>22.9 %</u>	<u>22.9 %</u>	<u>38.4 %</u>
<u>Wilting point</u>	<u>10 %</u>	<u>10 %</u>	<u>10 %</u>	<u>13.4 %</u>
<u>Available water capacity</u>	<u>12.9 %</u>	<u>12.9 %</u>	<u>12.9 %</u>	<u>13.4 %</u>

*Available water capacity = Field capacity - Wilting point




Figure A1. Heatwave detection using the daily maximum air temperature summer 2021 and the control period (1991-2020). The red color indicates the period where the daily maximum air temperature in the summer of 2021 exceeded the control period.




Figure A2. Meteorological condition at the Street site showing hourly a) air temperature (Air T), b) water vapor deficit (VPD), c) soil temperature (Soil T) and d) soil moisture measured and e) daily mean photosynthetically active radiation (PAR) and daily sum precipitation data measured at the SMEARIII station. The yellow markers in panel (c) denote the dates of manual leaf gas measurements.




Figure A3. Meteorological condition at the Forest site showing hourly a) air temperature (Air T), b) water vapor deficit (VPD), c) soil temperature (Soil T) and d) soil moisture measured and e) daily mean photosynthetically active radiation (PAR) and daily sum precipitation data measured at the SMEARIII station. The yellow markers in panel (c) denote the dates of manual leaf gas measurements.




Figure A4. Meteorological condition at the Orchard site showing hourly a) air temperature (Air T), b) water vapor deficit (VPD), c) soil temperature (Soil T) and d) soil moisture measured and e) daily mean photosynthetically active radiation (PAR) and daily sum precipitation data measured at the SMEARIII station. The yellow markers in panel (c) denote the dates of manual leaf gas measurements.




Figure A5. Monthly values of relative water content (RWC %) at the four urban sites.




Figure A6. Pattern of sap flux density during heat and dry period at the four urban sites.

Author contributions. Conceptualization: JA, LK, LJ; Data collection: JA, JS, YF, AK, EK, EV; Formal analysis: JA, JS, AK; Funding acquisition: LK, LJ; Supervision: LK, LJ; Visualization: JA, JS; Writing—original draft preparation: JA; Writing—review and editing: JA, LK, LJ, YS, AL. All authors have read and agreed to the article.

470 *Competing interests.* The authors declare that there is no conflict of interest.

Acknowledgements. We thank the Academy of Finland (CarboCity project, decision numbers: 321527, 325549, 337549, and 337552), the Academy of Finland ACCC Flagship (decision numbers: 337549, 337552) and the Strategic Research Council working under the Academy of Finland (CO-CARBON project, decision numbers: 335201 and 335204). This project has also received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement 101037319 (PAUL project). We would like to thank Eki, Jarkko

475 Mäntylä, Elisa Vainio and Teemu Paljakka for their technical support during the field measurements.

References

- Atkin, O. K. and Tjoelker, M. G.: Thermal acclimation and the dynamic response of plant respiration to temperature, *Trends in Plant Science*, 8, 343–351, [https://doi.org/10.1016/S1360-1385\(03\)00136-5](https://doi.org/10.1016/S1360-1385(03)00136-5), 2003.
- Bowler, D. E., Buyung-Ali, L., Knight, T. M., and Pullin, A. S.: Urban greening to cool towns and cities: A systematic review of the empirical
480 evidence, *Landscape and Urban Planning*, 97, 147–155, <https://doi.org/10.1016/j.landurbplan.2010.05.006>, 2010.
- Brack, C. L.: Pollution mitigation and carbon sequestration by an urban forest, *Environmental Pollution*, 116, S195–S200, [https://doi.org/10.1016/S0269-7491\(01\)00251-2](https://doi.org/10.1016/S0269-7491(01)00251-2), 2002.
- Bush, S. E., Pataki, D. E., Hultine, K. R., West, A. G., Sperry, J. S., and Ehleringer, J. R.: Wood anatomy constrains stomatal responses to atmospheric vapor pressure deficit in irrigated, urban trees, *Oecologia*, 156, 13–20, <https://doi.org/10.1007/s00442-008-0966-5>, 2008.
- 485 Bussotti, F., Pollastrini, M., Killi, D., Ferrini, F., and Fini, A.: Ecophysiology of urban trees in a perspective of climate change., *Agrochimica*, 58, 247–268, <https://www.cabdirect.org/cabdirect/abstract/20153144491>, publisher: Università degli Studi di Pisa, 2014.
- Czaja, M., Kołton, A., and Muras, P.: The Complex Issue of Urban Trees—Stress Factor Accumulation and Ecological Service Possibilities, *Forests*, 11, 932, <https://doi.org/10.3390/f11090932>, number: 9 Publisher: Multidisciplinary Digital Publishing Institute, 2020.
- Davies, Z. G., Edmondson, J. L., Heinemeyer, A., Leake, J. R., and Gaston, K. J.: Mapping an urban ecosystem service: quantifying
490 above-ground carbon storage at a city-wide scale, *Journal of Applied Ecology*, 48, 1125–1134, <https://doi.org/10.1111/j.1365-2664.2011.02021.x>,
_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2011.02021.x>, 2011.
- De Kauwe, M. G., Medlyn, B. E., Pitman, A. J., Drake, J. E., Ukkola, A., Griebel, A., Pendall, E., Prober, S., and Roderick, M.: Examining the evidence for decoupling between photosynthesis and transpiration during heat extremes, *Biogeosciences*, 16, 903–916, <https://doi.org/10.5194/bg-16-903-2019>, publisher: Copernicus GmbH, 2019.
- 495 De Micco, V. and Aronne, G.: Morpho-Anatomical Traits for Plant Adaptation to Drought, in: *Plant Responses to Drought Stress: From Morphological to Molecular Features*, edited by Aroca, R., pp. 37–61, Springer, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-32653-0_2, 2012.
- Dhakar, S.: GHG emissions from urbanization and opportunities for urban carbon mitigation, *Current Opinion in Environmental Sustainability*, 2, 277–283, <https://doi.org/10.1016/j.cosust.2010.05.007>, 2010.
- 500 Drake, J., Tjoelker, M., Varhammar, A., Medlyn, B., Reich, P., Leigh, A., Pfautsch, S., Blackman, C., López, R., Aspinwall, M., Crous, K., Duursma, R., Kumarathunge, D., De Kauwe, M., Jiang, M., Nicotra, A., Tissue, D., Choat, B., Atkin, O., and Barton, C.: Trees Tolerate an Extreme Heatwave via Sustained Transpirational Cooling and Increased Leaf Thermal Tolerance, *Global Change Biology*, 24, <https://doi.org/10.1111/gcb.14037>, 2018.
- Duarte, A. G., Katata, G., Hoshika, Y., Hossain, M., Kreuzwieser, J., Arneth, A., and Ruehr, N. K.: Immediate and potential long-term effects
505 of consecutive heat waves on the photosynthetic performance and water balance in Douglas-fir, *Journal of Plant Physiology*, 205, 57–66, <https://doi.org/10.1016/j.jplph.2016.08.012>, 2016.
- Fischer, E. M. and Schär, C.: Consistent geographical patterns of changes in high-impact European heatwaves, *Nature Geoscience*, 3, 398–403, <https://doi.org/10.1038/ngeo866>, number: 6 Publisher: Nature Publishing Group, 2010.
- FMI: The Finnish Meteorological Institute, <https://en.ilmatiiteenlaitos.fi/open-data>, 2021.
- 510 Gebauer, T., Horna, V., and Leuschner, C.: Variability in radial sap flux density patterns and sapwood area among seven co-occurring temperate broad-leaved tree species, *Tree Physiology*, 28, 1821–1830, <https://doi.org/10.1093/treephys/28.12.1821>, 2008.

- Ghannoum, O. and Way, D. A.: On the role of ecological adaptation and geographic distribution in the response of trees to climate change, *Tree Physiology*, 31, 1273–1276, <https://doi.org/10.1093/treephys/tpr115>, 2011.
- Gillner, S., Korn, S., and Roloff, A.: Leaf-Gas Exchange of Five Tree Species at Urban Street Sites, *Arboriculture & Urban Forestry*, 41, 113–124, 2015a.
- Gillner, S., Vogt, J., Tharang, A., Dettmann, S., and Roloff, A.: Role of street trees in mitigating effects of heat and drought at highly sealed urban sites, *Landscape and Urban Planning*, 143, 33–42, <https://doi.org/10.1016/j.landurbplan.2015.06.005>, 2015b.
- Gillner, S., Korn, S., Hofmann, M., and Roloff, A.: Contrasting strategies for tree species to cope with heat and dry conditions at urban sites, *Urban Ecosystems*, 20, <https://doi.org/10.1007/s11252-016-0636-z>, 2017.
- Gourlez de la Motte, L., Beauclaire, Q., Heinesch, B., Cuntz, M., Foltýnová, L., Šigut, L., Kowalska, N., Manca, G., Ballarin, I. G., Vincke, C., Roland, M., Ibrom, A., Lousteau, D., Siebicke, L., Neiryink, J., and Longdoz, B.: Non-stomatal processes reduce gross primary productivity in temperate forest ecosystems during severe edaphic drought, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375, 20190527, <https://doi.org/10.1098/rstb.2019.0527>, publisher: Royal Society, 2020.
- Granier, A.: Une nouvelle méthode pour la mesure du flux de sève brute dans le tronc des arbres, *Annales des Sciences Forestières*, 42, 193–200, <https://doi.org/10.1051/forest:19850204>, publisher: EDP Sciences, 1985.
- Granier, A., Bréda, N., Biron, P., and Villette, S.: A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands, *Ecological Modelling*, 116, 269–283, [https://doi.org/10.1016/S0304-3800\(98\)00205-1](https://doi.org/10.1016/S0304-3800(98)00205-1), 1999.
- Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Climate Dynamics*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- Hardiman, B. S., Wang, J. A., Hutyra, L. R., Gately, C. K., Getson, J. M., and Friedl, M. A.: Accounting for urban biogenic fluxes in regional carbon budgets, *Science of The Total Environment*, 592, 366–372, <https://doi.org/10.1016/j.scitotenv.2017.03.028>, 2017.
- Hartmann, H., Bastos, A., Das, A. J., Esquivel-Muelbert, A., Hammond, W. M., Martínez-Vilalta, J., McDowell, N. G., Powers, J. S., Pugh, T. A., Ruthrof, K. X., and Allen, C. D.: Climate Change Risks to Global Forest Health: Emergence of Unexpected Events of Elevated Tree Mortality Worldwide, *Annual Review of Plant Biology*, 73, 673–702, <https://doi.org/10.1146/annurev-arplant-102820-012804>, [_eprint: https://doi.org/10.1146/annurev-arplant-102820-012804](https://doi.org/10.1146/annurev-arplant-102820-012804), 2022.
- Hernandez-Santana, V., Hernandez-Hernandez, A., Vadeboncoeur, M. A., and Asbjornsen, H.: Scaling from single-point sap velocity measurements to stand transpiration in a multispecies deciduous forest: uncertainty sources, stand structure effect, and future scenarios, *Canadian Journal of Forest Research*, 45, 1489–1497, <https://doi.org/10.1139/cjfr-2015-0009>, publisher: NRC Research Press, 2015.
- Ibsen, P. C., Borowy, D., Dell, T., Greydanus, H., Gupta, N., Hondula, D. M., Meixner, T., Santelmann, M. V., Shiflett, S. A., Sukop, M. C., Swan, C. M., Talal, M. L., Valencia, M., Wright, M. K., and Jenerette, G. D.: Greater aridity increases the magnitude of urban nighttime vegetation-derived air cooling, *Environmental Research Letters*, 16, 034011, <https://doi.org/10.1088/1748-9326/abdf8a>, publisher: IOP Publishing, 2021.
- Jim, C. Y. and Chen, W. Y.: Ecosystem services and valuation of urban forests in China, *Cities*, 26, 187–194, <https://doi.org/10.1016/j.cities.2009.03.003>, 2009.
- Jo, H.-k.: Impacts of urban greenspace on offsetting carbon emissions for middle Korea, *Journal of Environmental Management*, 64, 115–126, <https://doi.org/10.1006/jema.2001.0491>, 2002.
- Järvi, L., Hannuniemi, H., Hussein, T., Junninen, H., Aalto, P. P., Hillamo, R., Mäkelä, T., Keronen, P., Siivola, E., Vesala, T., and Kulmala, M.: The urban measurement station SMEAR III: Continuous monitoring of air pollution and surface–atmosphere interactions in Helsinki,

- Finland, <https://helda.helsinki.fi/handle/10138/233627>, accepted: 2018-03-20T15:08:03Z Publisher: Boreal Environment Research Publishing Board, 2009.
- 550 Kjelgren, R. and Clark, J.: Microclimates and tree growth in three urban spaces, *Journal of Environmental Horticulture*, 10, 139–145, https://digitalcommons.usu.edu/psc_facpub/658, 1992.
- Kjelgren, R. and Montague, T.: Urban tree transpiration over turf and asphalt surfaces, *Atmospheric Environment*, 32, 35–41, [https://doi.org/10.1016/S1352-2310\(97\)00177-5](https://doi.org/10.1016/S1352-2310(97)00177-5), aDS Bibcode: 1998AtmEn..32...35K, 1998.
- 555 Konarska, J., Uddling, J., Holmer, B., Lutz, M., Lindberg, F., Pleijel, H., and Thorsson, S.: Transpiration of urban trees and its cooling effect in a high latitude city, *International Journal of Biometeorology*, 60, 159–172, <https://doi.org/10.1007/s00484-015-1014-x>, 2016.
- Kunert, N., Hajek, P., Hietz, P., Morris, H., Rosner, S., and Tholen, D.: Summer temperatures reach the thermal tolerance threshold of photosynthetic decline in temperate conifers, *Plant Biology*, n/a, <https://doi.org/10.1111/plb.13349>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/plb.13349>, 2022.
- 560 Leuschner, C., Wedde, P., and Lübbe, T.: The relation between pressure–volume curve traits and stomatal regulation of water potential in five temperate broadleaf tree species, *Annals of Forest Science*, 76, 1–14, <https://doi.org/10.1007/s13595-019-0838-7>, number: 2 Publisher: BioMed Central, 2019.
- Leuzinger, S., Vogt, R., and Körner, C.: Tree surface temperature in an urban environment, *Agricultural and Forest Meteorology*, 150, 56–62, <https://doi.org/10.1016/j.agrformet.2009.08.006>, 2010.
- 565 Lindén, J., Fonti, P., and Esper, J.: Temporal variations in microclimate cooling induced by urban trees in Mainz, Germany, *Urban Forestry & Urban Greening*, 20, 198–209, <https://doi.org/10.1016/j.ufug.2016.09.001>, 2016.
- Lloyd, J. and Farquhar, G. D.: Effects of rising temperatures and [CO₂] on the physiology of tropical forest trees, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363, 1811–1817, <https://doi.org/10.1098/rstb.2007.0032>, 2008.
- Lu, P., Urban, L., and Zhao, P.: Granier’s thermal dissipation probe (TDP) method for measuring sap flow in trees: theory and practice, <https://publications.csiro.au/rpr/pub?list=BRO&pid=procite:ac6fc6af-67dc-4b32-ac82-5d3c9678573a>, 2004.
- 570 Marchin, R. M., Backes, D., Ossola, A., Leishman, M. R., Tjoelker, M. G., and Ellsworth, D. S.: Extreme heat increases stomatal conductance and drought-induced mortality risk in vulnerable plant species, *Global change biology*, 28, 1133–1146, publisher: Wiley Online Library, 2022.
- McCarthy, H. R. and Pataki, D. E.: Drivers of variability in water use of native and non-native urban trees in the greater Los Angeles area, *Urban Ecosystems*, 13, 393–414, <https://doi.org/10.1007/s11252-010-0127-6>, 2010.
- 575 Moser, A., Rahman, M. A., Pretzsch, H., Pauleit, S., and Rötzer, T.: Inter- and intraannual growth patterns of urban small-leaved lime (*Tilia cordata* mill.) at two public squares with contrasting microclimatic conditions, *International Journal of Biometeorology*, 61, 1095–1107, <https://doi.org/10.1007/s00484-016-1290-0>, 2017.
- Muñoz-Vallés, S., Cambrollé, J., Figueroa-Luque, E., Luque, T., Niell, F. X., and Figueroa, M. E.: An approach to the evaluation and management of natural carbon sinks: From plant species to urban green systems, *Urban Forestry & Urban Greening*, 12, 450–453, <https://doi.org/10.1016/j.ufug.2013.06.007>, 2013.
- 580 Nielsen, C., Bühler, O., and Kristoffersen, P.: Soil Water Dynamics and Growth of Street and Park Trees, *Arboriculture & Urban Forestry*, 33, 231–245, <https://doi.org/10.48044/jauf.2007.027>, 2007.
- Nowak, D. J. and Crane, D. E.: Carbon storage and sequestration by urban trees in the USA, *Environmental Pollution*, 116, 381–389, [https://doi.org/10.1016/S0269-7491\(01\)00214-7](https://doi.org/10.1016/S0269-7491(01)00214-7), 2002.
- 585

- Nowak, D. J., Greenfield, E. J., Hoehn, R. E., and Lapoint, E.: Carbon storage and sequestration by trees in urban and community areas of the United States, *Environmental Pollution*, 178, 229–236, <https://doi.org/10.1016/j.envpol.2013.03.019>, 2013.
- Oke, T. R., Crowther, J. M., McNaughton, K. G., Monteith, J. L., Gardiner, B., Jarvis, P. G., Monteith, J. L., Shuttleworth, W. J., and Unsworth, M. H.: The micrometeorology of the urban forest, *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 324, 335–349, <https://doi.org/10.1098/rstb.1989.0051>, publisher: Royal Society, 1989.
- Osone, Y., Kawarasaki, S., Ishida, A., Kikuchi, S., Shimizu, A., Yazaki, K., Aikawa, S.-I., Yamaguchi, M., Izuta, T., and Matsumoto, G. I.: Responses of gas-exchange rates and water relations to annual fluctuations of weather in three species of urban street trees, *Tree Physiology*, 34, 1056–1068, <https://doi.org/10.1093/treephys/tpu086>, 2014.
- Paloheimo, E. and Salmi, O.: Evaluating the carbon emissions of the low carbon city: A novel approach for consumer based allocation, *Cities*, 30, 233–239, <https://doi.org/10.1016/j.cities.2012.04.003>, 2013.
- Pataki, D. E., Emmi, P. C., Forster, C. B., Mills, J. I., Pardyjak, E. R., Peterson, T. R., Thompson, J. D., and Dudley-Murphy, E.: An integrated approach to improving fossil fuel emissions scenarios with urban ecosystem studies, *Ecological Complexity*, 6, 1–14, <https://doi.org/10.1016/j.ecocom.2008.09.003>, 2009.
- Pataki, D. E., McCarthy, H. R., Litvak, E., and Pincetl, S.: Transpiration of urban forests in the Los Angeles metropolitan area, *Ecological Applications*, 21, 661–677, <https://doi.org/10.1890/09-1717.1>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1890/09-1717.1>, 2011.
- Pataki, D. E., Alberti, M., Cadenasso, M. L., Felson, A. J., McDonnell, M. J., Pincetl, S., Pouyat, R. V., Setälä, H., and Whitlow, T. H.: The Benefits and Limits of Urban Tree Planting for Environmental and Human Health, *Frontiers in Ecology and Evolution*, 9, <https://www.frontiersin.org/article/10.3389/fevo.2021.603757>, 2021.
- Perini, K. and Magliocco, A.: Effects of vegetation, urban density, building height, and atmospheric conditions on local temperatures and thermal comfort, *Urban Forestry & Urban Greening*, 13, 495–506, <https://doi.org/10.1016/j.ufug.2014.03.003>, 2014.
- Rahman, M. A., Moser, A., Rötzer, T., and Pauleit, S.: Comparing the transpirational and shading effects of two contrasting urban tree species, *Urban Ecosystems*, 22, 683–697, <https://doi.org/10.1007/s11252-019-00853-x>, 2019.
- Rennenberg, H., Loreto, F., Polle, A., Brilli, F., Fares, S., Beniwal, R. S., and Gessler, A.: Physiological responses of forest trees to heat and drought, *Plant Biology (Stuttgart, Germany)*, 8, 556–571, <https://doi.org/10.1055/s-2006-924084>, 2006.
- Riikonen, A., Järvi, L., and Nikinmaa, E.: Environmental and crown related factors affecting street tree transpiration in Helsinki, Finland, *Urban Ecosystems*, 19, 1693–1715, <https://doi.org/10.1007/s11252-016-0561-1>, 2016.
- Roth, M., Oke, T. R., and Emery, W. J.: Satellite-derived urban heat islands from three coastal cities and the utilization of such data in urban climatology, *International Journal of Remote Sensing*, 10, 1699–1720, <https://doi.org/10.1080/01431168908904002>, publisher: Taylor & Francis eprint: <https://doi.org/10.1080/01431168908904002>, 1989.
- RStudio Team: RStudio Integrated Development Environment for R, RStudio, PBC., Boston, MA, <httpwww.rstudio.com>, 2020.
- Rötzer, T., Moser-Reischl, A., Rahman, M. A., Hartmann, C., Paeth, H., Pauleit, S., and Pretzsch, H.: Urban tree growth and ecosystem services under extreme drought, *Agricultural and Forest Meteorology*, 308–309, 108 532, <https://doi.org/10.1016/j.agrformet.2021.108532>, 2021.
- Sjöman, H., Östberg, J., and Bühler, O.: Diversity and distribution of the urban tree population in ten major Nordic cities, *Urban Forestry & Urban Greening*, 11, 31–39, <https://doi.org/10.1016/j.ufug.2011.09.004>, 2012.
- StromJan: StromJan/Raster4H: Final version, <https://doi.org/10.5281/zenodo.4005833>, 2020.

- Sushko, S., Yaroslavtsev, A., Tsuwareva, N., and Valentini, R.: Capacity of *Quercus robur* L. and *Tilia cordata* Mill. trees in providing urban ecosystem services in boreal climate, pp. EGU21–7125, <https://doi.org/10.5194/egusphere-egu21-7125>, conference Name: EGU General Assembly Conference Abstracts ADS Bibcode: 2021EGUGA..23.7125S, 2021.
- 625 Urban, J., Ingwers, M. W., McGuire, M. A., and Teskey, R. O.: Increase in leaf temperature opens stomata and decouples net photosynthesis from stomatal conductance in *Pinus taeda* and *Populus deltoides* x *nigra*, *Journal of Experimental Botany*, 68, 1757–1767, <https://doi.org/10.1093/jxb/erx052>, 2017.
- Vesala, T., Järvi, L., Launiainen, S., Sogachev, A., Rannik, , Mammarella, I., Ivola, E. S., Keronen, P., Rinne, J., Riikonen, A., and Nikinmaa, E.: Surface–atmosphere interactions over complex urban terrain in Helsinki, Finland, *Tellus B: Chemical and Physical Meteorology*, 60, 630 188–199, <https://doi.org/10.1111/j.1600-0889.2007.00312.x>, 2008.
- Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718, <https://doi.org/10.1175/2009JCLI2909.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2010.
- Villar-Salvador, P., Planelles, R., Oliet, J., Peñuelas-Rubira, J. L., Jacobs, D. F., and González, M.: Drought tolerance and transplanting performance of holm oak (*Quercus ilex*) seedlings after drought hardening in the nursery, *Tree Physiology*, 24, 1147–1155, <https://doi.org/10.1093/treephys/24.10.1147>, 2004.
- 635 Whitlow, T. H., Bassuk, N. L., and Reichert, D. L.: A 3-Year Study of Water Relations of Urban Street Trees, *Journal of Applied Ecology*, 29, 436–450, <https://doi.org/10.2307/2404512>, publisher: [British Ecological Society, Wiley], 1992.
- Winbourne, J. B., Jones, T. S., Garvey, S. M., Harrison, J. L., Wang, L., Li, D., Templer, P. H., and Hutya, L. R.: Tree Transpiration and Urban Temperatures: Current Understanding, Implications, and Future Research Directions, *BioScience*, 70, 576–588, <https://doi.org/10.1093/biosci/biaa055>, 2020.
- 640 Zapater, M., Bréda, N., Bonal, D., Pardonnet, S., and Granier, A.: Differential response to soil drought among co-occurring broad-leaved tree species growing in a 15- to 25-year-old mixed stand, *Annals of Forest Science*, 70, 31–39, <https://doi.org/10.1007/s13595-012-0233-0>, number: 1 Publisher: BioMed Central, 2013.