

# 1 **High-resolution spatial patterns and drivers of terrestrial** 2 **ecosystem carbon dioxide, methane, and nitrous oxide fluxes** 3 **in the tundra**

4 Anna-Maria Virkkala<sup>1,2</sup>, Pekka Niittynen<sup>3</sup>, Julia Kemppinen<sup>4</sup>, Maija E. Marushchak<sup>5</sup>, Carolina Voigt<sup>5</sup>,  
5 Geert Hensgens<sup>6</sup>, Johanna Kerttula<sup>5</sup>, Konsta Happonen<sup>7</sup>, Vilna Tyystjärvi<sup>8</sup>, Christina Biasi<sup>5,11</sup>, Jenni  
6 Hultman<sup>9,10</sup>, Janne Rinne<sup>9</sup>, Miska Luoto<sup>2</sup>

7 <sup>1</sup>Woodwell Climate Research Center, Falmouth, 149 Woods Hole Road, MA, USA

8 <sup>2</sup>University of Helsinki, Department of Geosciences and Geography, Gustaf Hällströmin katu 2, 00014

9 University of Helsinki, Finland

10 <sup>3</sup>University of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35 FI-40014,

11 Jyväskylä, Finland

12 <sup>4</sup>Geography Research Unit, University of Oulu, P.O. Box 8000 FI-90014, Oulu, Finland

13 <sup>5</sup>University of Eastern Finland, Department of Environmental and Biological Sciences, P.O. Box 1627 FI-  
14 70211, Kuopio, Finland

15 <sup>6</sup>Vrije Universiteit Amsterdam, Department of Earth and Climate, De Boelelaan 1085, 1081 HV, Amsterdam,  
16 the Netherlands

17 <sup>7</sup>Youth Research Society, Kumpulantie 3 A, 00520 Helsinki, Finland

18 <sup>8</sup>Finnish Meteorological Institute, Climate System Research Unit, Erik Palménin aukio 1, FI-00560 Helsinki,  
19 Finland

20 <sup>9</sup>Natural Resources Institute Finland, Latokartanonkaari 9, 00790 Helsinki, Finland

21 <sup>10</sup>University of Helsinki, Department of Microbiology, Viikinkaari 9, 00014 University of Helsinki, Finland

22 <sup>11</sup>Department of Ecology, University of Innsbruck, Sternwartstrasse 15, 6020, Innsbruck, Austria

23 *Correspondence to:* Anna-Maria Virkkala, avirkkala@woodwellclimate.org

24

25 **Abstract.** Arctic terrestrial greenhouse gas (GHG) fluxes of carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous  
26 oxide (N<sub>2</sub>O) play an important role in the global GHG budget. However, these GHG fluxes are rarely studied  
27 simultaneously, and our understanding of the conditions controlling them across spatial gradients is limited.  
28 Here, we explore the magnitudes and drivers of GHG fluxes across fine-scale terrestrial gradients during the  
29 peak growing season (July) in sub-Arctic Finland. We measured chamber-derived GHG fluxes and soil  
30 temperature, soil moisture, soil organic carbon and nitrogen stocks, soil pH, soil carbon-to-nitrogen (C/N) ratio,  
31 soil dissolved organic carbon content, vascular plant biomass, and vegetation type from 101 plots scattered  
32 across a heterogeneous tundra landscape (5 km<sup>2</sup>). We used these field data together with high-resolution remote  
33 sensing data to develop machine learning models for predicting (i.e., upscaling) daytime GHG fluxes across the  
34 landscape at 2-m resolution. Our results show that this region was on average a daytime net GHG sink during  
35 the growing season. Although our results suggest that this sink was driven by CO<sub>2</sub> uptake, it also revealed small  
36 but widespread CH<sub>4</sub> uptake in upland vegetation types, almost surpassing the high wetland CH<sub>4</sub> emissions at the  
37 landscape scale. Average N<sub>2</sub>O fluxes were negligible. CO<sub>2</sub> fluxes were controlled primarily by annual average  
38 soil temperature and biomass (both increase net sink) and vegetation type, CH<sub>4</sub> fluxes by soil moisture  
39 (increases net emissions) and vegetation type, and N<sub>2</sub>O fluxes by soil C/N (lower C/N increases net source).  
40 These results demonstrate the potential of high spatial resolution modeling of GHG fluxes in the Arctic. They  
41 also reveal the dominant role of CO<sub>2</sub> fluxes across the tundra landscape, but suggest that CH<sub>4</sub> uptake in dry  
42 upland soils might play a significant role in the regional GHG budget.

## 43 **1 Introduction**

44 Over the past millennia, Arctic soils in the treeless tundra biome have played an important role in the global  
45 climate system by accumulating large amounts of carbon (C) and nitrogen (N), thus cooling the climate  
46 (Hugelius et al., 2014, 2020; Strauss et al., 2017). However, the ongoing climate warming is changing the C and  
47 N cycles, leading to potentially increased net greenhouse gas (GHG) emissions from Arctic ecosystems to the  
48 atmosphere (Belshe et al., 2013; McGuire et al., 2012; Masyagina and Menyailo, 2020). Yet, even the current  
49 GHG balance of Arctic ecosystems is insufficiently understood due to severe gaps in flux measurement  
50 networks and poorly performing coarse-resolution models (Virkkala et al., 2021; Treat et al., 2018c). Thus, the  
51 contribution of Arctic regions to the global climate feedback remains uncertain.

52 One of the main uncertainties in understanding the Arctic GHG balance is related to the inadequately quantified  
53 magnitudes of all three main GHG fluxes - carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) -  
54 which show pronounced spatial variability across the diverse terrestrial environmental gradients in tundra  
55 (Virkkala et al., 2018; Pallandt et al., 2021; Voigt et al., 2020). In most tundra ecosystems, CO<sub>2</sub> fluxes are the  
56 largest flux driving the GHG balance due to the strong growing season photosynthetic activity and relatively  
57 high non-growing season respiratory CO<sub>2</sub> losses (Natali et al., 2019; Euskirchen et al., 2012; Heiskanen et al.,  
58 2021). However, growing evidence points to the importance of CH<sub>4</sub> and N<sub>2</sub>O fluxes, which are more potent  
59 GHGs than CO<sub>2</sub> (Voigt et al., 2017b). All three gasses have distinct spatiotemporal dynamics (Emmerton et al.,  
60 2014; Bruhwiler et al., 2021). However, only a few studies have simultaneously considered the contributions of  
61 all three main GHG fluxes to the tundra GHG balance (Voigt et al., 2017b; Kelsey et al., 2016; Brummell et al.,  
62 2012; Wagner et al., 2019).

63 The largest fine-scale differences in Arctic GHG fluxes occur in ecosystems with spatially varying soil moisture  
64 conditions (McGuire et al., 2012). Broadly speaking, the Arctic can be divided into wetlands and drier uplands  
65 (i.e., shrublands, grasslands, and barren lands; see e.g. Treat et al., 2018a; Virkkala et al., 2021). Wetlands cover  
66 between 5 and 25 % of the Arctic (Olefeldt et al., 2021; Kåresdotter et al., 2021; Reynolds et al., 2019). They  
67 are hotspots for soil C and N stocks and have the potential for high CH<sub>4</sub> emissions (Euskirchen et al., 2014;  
68 Hugelius et al., 2020); therefore they have been intensively studied (Rinne et al., 2018; Peltola et al., 2019;  
69 Turetsky et al., 2014). However, uplands cover the largest part of the Arctic (75 to 95 %) and can have  
70 significant variability in environmental conditions and GHG fluxes. These uplands have been relatively well  
71 studied for CO<sub>2</sub> fluxes (Williams et al., 2006; Cahoon et al., 2012a). Upland CH<sub>4</sub> and N<sub>2</sub>O fluxes, on the other  
72 hand, remain less well understood in terms of their magnitudes and drivers (Virkkala et al., 2018; Voigt et al.,  
73 2020). There are still likely some GHG flux hotspots to be discovered and coldspots to be verified, particularly  
74 in the upland tundra ecosystems.

75 The Arctic tundra is characterised by fine-scale environmental heterogeneity even within upland and wetland  
76 tundra environments. Thus, local-scale study settings that cover the main spatial environmental gradients are  
77 highly important (Treat et al., 2018c; Davidson et al., 2017). Such fine-scale variabilities are often measured  
78 with chambers, but most chamber-based study designs are limited to relatively small environmental gradients  
79 focusing on only a handful of different land cover types and environmental variables, leaving large gaps in our  
80 understanding of GHG flux hotspots (Virkkala et al. 2018). In this study, using an extensive spatial study design

81 with chamber GHG flux measurements from 101 plots, we aim to understand the magnitudes and environmental  
82 drivers of Arctic terrestrial CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes in a heterogeneous tundra landscape dominated by upland  
83 heaths. By combining in-situ measurements and remote sensing data, we investigate the fine-scale (2 m) spatial  
84 heterogeneity of GHG fluxes across the landscape, and estimate the contribution of the three gases to the total  
85 landscape-scale GHG flux.


86

## 87 **2 Materials and Methods**

### 88 **2.1 Study area**

89 The field measurements were collected during 2016-2018 in a sub-Arctic tundra environment in Kilpisjärvi  
90 (Gilbjesjávri in Northern Sámi language), northwestern Finland (69.06 N, 20.81 E). The study area is located on  
91 an elevational gradient between two fells, Saana (Sána; 1029 m.a.s.l) and Korkea-Jehkats (Jiehkkaš; 960 m.a.s.l),  
92 and the valley in between (~600 m.a.s.l.). The study area is above a mountain birch (*Betula pubescens* ssp.  
93 *czerepanovii*) forest and is dominated by dwarf-shrub evergreen and deciduous heaths. Dominant vascular plant  
94 species are, e.g., *Empetrum nigrum* ssp. *hermaphroditum*, *Betula nana*, *Vaccinium myrtillus*, *Vaccinium vitis-*  
95 *idaea*, and *Phyllodoce caerulea*. Vegetation in the wetlands is dominated by species common to fen wetlands,  
96 such as *Eriophorum* sp. or *Carex* sp. Mesic meadows are rich in forbs and grasses whereas barren heaths  
97 accommodate mostly lichens (e.g. *Cladonia* spp.) and mat-forming cushion plants (e.g. *Diapensia lapponica*) with  
98 scattered patches of *E. nigrum* and *B. nana*. Soils in the area are shallow (mean organic layer depth 6.6 cm, mean  
99 mineral layer depth 13.0 cm), and permafrost is absent from soils but can be found in the bedrock above 800 m  
100 a.s.l. (King and Seppälä, 1987). The environment is relatively undisturbed but experiences reindeer (*Rangifer*  
101 *tarandus tarandus*) grazing. The mean annual temperature in Saana fell (1002 m.a.s.l.) is -3.1 °C and the annual  
102 precipitation in Kilpisjärvi village ca. 5 km from the study site (480 m.a.s.l.) is 518 mm in 1991-2018 (Finnish  
103 Meteorological Institute, 2019a, b).

104 Our study design covered an area of ca. 3 x 1.5 km and consisted of 101 plots with GHG flux measurements and  
105 their supporting environmental data (Fig. 1). To produce continuous maps of soil temperature, moisture,  
106 vegetation type, biomass, soil C/N, and soil organic carbon stock, we utilized an extended dataset where some of  
107 the variables were measured from 50 plots while others from more than 6000 plots (Table S1). We selected the  
108 plots based on a combination of stratified sampling and systematic grid approaches, and the plots contain a variety  
109 of environmental gradients and habitats as well as the transition zones between them (Kemppinen et al., 2021).  
110 We recorded the locations of the plots using a hand-held Global Navigation Satellite System receiver with an  
111 accuracy of up to ≤6 cm under optimal conditions (GeoExplorer GeoXH 6000 Series; Trimble Inc., Sunnyvale,  
112 CA, USA).



113  
 114 **Figure 1:** The distribution of the main vegetation types across the Arctic tundra (Dinerstein et al., 2017; European  
 115 Space Agency, 2017) and the location of our study area (a), the distribution of plots (b) and environmental  
 116 conditions derived from statistical upscaling of in-situ measurements (see Sect. 2.4.2 Machine learning models)  
 117 across the study area (c). Soil moisture and temperature represent mean daytime (8 am to 8 pm) conditions from  
 118 the 1st of July to the 2nd of August and annual soil temperature is an average within the entire year (July 2017-  
 119 June 2018). Other variables represent growing season conditions and are considered static in this study. The aerial  
 120 image is produced by the National Land Survey of Finland (accessed in 2016).

## 121 2.2 Data

122 We measured GHG fluxes from 101 plots during the peak growing season (from now on, growing season).  
123 Environmental conditions explaining these GHG fluxes were measured at each plot. Most environmental variables  
124 had near complete spatial coverage; missing data were filled using the environmental predictions (see Sect. 2.4.2  
125 Machine learning models, Table S1). We used additional in-situ environmental data to upscale and visualize  
126 environmental conditions across the entire landscape (see Sect. 2.4.2 Machine learning models and Fig. 2):  
127 continuous soil moisture loggers (50 plots), continuous soil temperature loggers (250), soil samples for carbon  
128 and nitrogen stock and C/N estimation (168), and vegetation classification data (6524). The full set of variables  
129 at a plot consisted of the plot for GHG flux measurements, and of a nearby complementing plot (max. 2 m distance)  
130 where we monitored soil moisture and temperature continuously and did a vegetation classification. The additional  
131 plot was separated from the GHG plot to avoid disturbance of the continuous recordings. The additional plot was  
132 carefully situated to similar vegetation and microtopographic conditions as the GHG plot. Soil samples were  
133 collected as close as possible to the GHG plot.

### 134 2.2.1 Chamber measurements

135 We measured GHG exchange using a static, non-steady state non-flow-through system (Livingston and  
136 Hutchingson, 1995) composed of an acrylic chamber (20 cm diameter, 25 cm height). The chamber was placed  
137 on top of a collar and ventilated before each measurement. Prior to the measurements, we installed steel collars,  
138 which were 21 cm in diameter and 6 - 7 cm in height. Each collar was visited once during the growing-season,  
139 and measurements were conducted between 10 am and 5 pm. Although we did not have any temporal replicates,  
140 the spatial variation in our plots covered most of the temperature variation during the growing season. For more  
141 details, see Sect. S1.

142 For CO<sub>2</sub> flux measurements, transparent and opaque chamber measurements were conducted during 1<sup>st</sup> of July  
143 and 27<sup>th</sup> of July, 2018. The chamber included a small fan, a CO<sub>2</sub> probe GMP343 and an air humidity and  
144 temperature probe HMP75 (Vaisala, Finland). In the chamber, CO<sub>2</sub> concentration, air temperature and relative  
145 air humidity were recorded at 5-s intervals for 90 s. Photosynthetically active radiation was logged manually  
146 outside the chamber at 10-s intervals during the same period using a MQ-200 quantum sensor with a hand-held  
147 meter (Apogee Instruments, Inc, USA). MQ-200 measures photosynthetic photon flux density (PPFD) at a  
148 spectral range from 410 to 655 nm in  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . For more details of the equipment, see Happonen et al.  
149 (2022).

150 We progressively decreased the light intensity of net ecosystem exchange (NEE) measurements from ambient  
151 conditions to ca. 80 %, 50 % and 30 % PPFD by shading the chamber with layers of white mosquito net  
152 (replicate measurements per collar = 5 - 9). Ecosystem respiration (ER) was measured in dark conditions (0  
153 PPFD), which were obtained by covering the chamber with a space blanket (replicates = 2 - 3). Before flux  
154 calculations, we discarded the first 0 - 5 s as well as the last 5 s of the measurements to remove potentially  
155 disturbed observations. Fluxes were calculated from the concentration change within the chamber headspace  
156 over time using linear regression (for performance statistics see Sect. S2).

157 We standardized NEE, GPP, and ER measurements conducted at different light and temperature conditions to  
158 allow across-plot comparison of the fluxes. We fitted light-response curves using a non-linear hierarchical  
159 bayesian model with the plot as a random effect (Sect. S5). We used the Michaelis-Menten equation to model  
160 instantaneous NEE with plot-specific ER, maximum photosynthetic rate ( $GPP_{max}$ ) and the half-saturation  
161 constant (K) as parameters using the same formula as in (Williams et al., 2006; Cahoon et al., 2012b). ER also  
162 had an exponential air temperature (T) response (for more details, see (Happonen et al., 2022)). We used this  
163 model to predict NEE at dark (0 PPFD, i.e. ER) and average light (600 PPFD) conditions, and an air temperature  
164 of 20 °C at each plot. 20 °C was chosen as it represents a typical air temperature inside the chamber during flux  
165 measurements, and 600 PPFD because it is widely used in tundra literature (Dagg and Lafleur, 2011; Shaver et  
166 al., 2007). We then subtracted ER from the NEE normalized to average light conditions to arrive at an estimate  
167 of normalized gross primary productivity (GPP). Negative values in NEE indicate a net sink of CO<sub>2</sub> from the  
168 atmosphere to the ecosystems. GPP and ER are given as positive values.

169 We measured CH<sub>4</sub> and N<sub>2</sub>O fluxes with an opaque chamber (0 PPFD). Measurements were conducted during  
170 the 2<sup>nd</sup> of July and 2<sup>nd</sup> of August, 2018. Five gas samples were taken within a 50-min enclosure time and  
171 transferred into 12-mL vials (Labco Exetainer, Labco Ltd.). The vials were pre-evacuated in the laboratory and  
172 filled with 25 mL of the sample in the field. Gas samples were analyzed at the University of Eastern Finland  
173 with a gas chromatograph (Agilent 7890B; Agilent Technologies, Santa Clara, CA, USA), equipped with an  
174 autosampler (Gilson Inc., Middleton, WI, USA), with thermal conductivity detector (TCD) for CO<sub>2</sub>, flame  
175 ionization detector (FID) for CH<sub>4</sub> and an electron capture detector (ECD) for N<sub>2</sub>O. We calculated gas  
176 concentrations from GC peak areas relative to peak areas derived by analyzing gas standards (CO<sub>2</sub>: 7  
177 concentration levels ranging from 0-10000 ppm; CH<sub>4</sub>: 7 concentration levels ranging from 0-100 ppm; N<sub>2</sub>O: 5  
178 concentration levels ranging from 0-5000 ppb). Fluxes were calculated from the concentration change within the  
179 chamber headspace over time using linear regression. Quality control was based on visual inspection and  
180 RMSE. We also verified that the RMSE was less than 3 \* standard deviation of gas standards in a similar  
181 concentration range. Negative values in these fluxes represent net CH<sub>4</sub> and N<sub>2</sub>O sinks from the atmosphere to  
182 the ecosystems.

### 183 2.2.2 Soil temperature and moisture data

184 Soil moisture and soil temperature were measured simultaneously during the flux measurements. We measured  
185 soil moisture with a time-domain reflectometry sensor (FieldScout TDR 300; Spectrum Technologies Inc.,  
186 Plainfield, IL, USA; 0 to 7.5 cm depth). Soil temperature measurements conducted at the same time as CO<sub>2</sub> flux  
187 measurements were taken with a thermometer (TD 11 thermometer; VWR International bvba; Leuven,  
188 Germany; 6.0 to 7.5 cm depth). Soil temperature measurements (TM-80N measure and ATT-50 sensor)  
189 conducted at the same time as CH<sub>4</sub> and N<sub>2</sub>O flux measurements were taken with a thermometer in the uppermost  
190 10 cm. We refer to these variables as soil moisture and soil temperature throughout the text.

191 Temperature loggers (Thermochron iButton DS1921G and DS1922L, San Jose, CA, USA and TMS-4; TOMST  
192 s.r.o., Prague, Czech Republic) monitored temperatures at 7.5 cm and 6.5 cm (iButton and TMS-4, respectively)  
193 belowground at 0.25–4 h intervals (Sect. S3). We calculated a variable describing average soil temperature

194 conditions during the previous 12 months by averaging the iButton measurements from the study design  
195 (n=138) from July, 2017 to June 2018. We refer to this variable as annual soil temperature. In addition to  
196 temperature, the TMS-4 loggers also monitored soil moisture (raw time-domain transmission data between 1  
197 and 4095) to a depth of c. 14 cm (Wild et al., 2019). The raw time-domain transmission data was transformed  
198 into volumetric water content (VWC%) (Tyystjärvi et al., 2022).

199 These continuous soil moisture and temperature data were used to upscale soil microclimatic conditions at 2-  
200 hour timesteps during daytime (8 am to 8 pm) and from the 1st of July to the 2nd of August (see section Models  
201 used to predict environmental conditions). This period was chosen because the GHG fluxes were measured  
202 during this period and we did not want to extrapolate outside our main measurement campaign. Moreover, this  
203 period represents the peak growing season of this region.

### 204 2.2.3 Vegetation data

205 We took images from CH<sub>4</sub> and N<sub>2</sub>O collars on the measurement day and used them to classify the dominant  
206 vegetation to five distinct classes, following the Circumpolar Arctic Vegetation Map physiognomic  
207 classification system (Walker et al. 2005) with minor modifications (Fig. 1). We used the following classes:  
208 barren (< 10 % vegetation cover), evergreen shrub, deciduous shrub, meadow (graminoids and forbs), and  
209 wetlands. The sample sizes were not even between vegetation types, rather they roughly represent the spatial  
210 coverage of each vegetation type (8 observations of barren, 38 of evergreen shrub, 14 of deciduous shrub, 26 of  
211 meadow, and 15 of wetland). We utilized a larger dataset of 6524 vegetation descriptions in total estimated in  
212 the field and from aerial imagery from the study design to create the vegetation type map (for more details, see  
213 Sect. S4.1). We collected biomass samples from above-ground vascular plants using the clip-harvest method  
214 during late peak season, between 17<sup>th</sup> of July and 10<sup>th</sup> of August. Samples were collected within the chamber  
215 collars and were oven-dried at 70 °C for 48 h and weighed after drying. We refer to this variable as biomass (g  
216 dry-weight m<sup>-2</sup>).

### 217 2.2.4 Soil sampling and analyses

218 We measured the thickness of the organic and mineral soil layers using a metal probe reaching up to 80 cm  
219 depth. We collected soil samples (ca. 1 dl) from the organic and mineral layers using metal soil core cylinders (4  
220 - 6 cm Ø, 5 - 7 cm height) during August in 2016-2018. The organic samples were collected from the top soil,  
221 and mineral samples directly below the organic layer which was on average 6.6 cm deep. Large roots were  
222 excluded from the samples. The soil samples were freeze-dried and analysed in the Laboratory of Geosciences  
223 and Geography and Laboratory of Forest Sciences (University of Helsinki). Bulk density (kg m<sup>-3</sup>) was estimated  
224 by dividing the dry weight by the sample volume. Soil organic layer pH was analyzed following ISO standard  
225 10390. Total carbon and nitrogen content (C%, N%) analyses were done using Vario Elementar Micro cube and  
226 Vario Elementar Max -analyzer (Elementar Analysensysteme GmbH, Germany). Prior to CN% analysis,  
227 mineral samples were sieved through a 2 mm plastic sieve. Organic samples were homogenized by hammering  
228 the material into smaller pieces.

229 Soils in this landscape are acidic and likely have a minimal amount of carbonates; consequently, we assumed  
230 C% to equal organic C%. Soil organic carbon and nitrogen stocks were calculated for the entire soil horizon up  
231 to 80 cm (in 95 % of plots soil depth was less than that). Some plots lacked CN% data (30 % of the plots), and  
232 therefore, we used soil organic matter content estimated with the loss-on-ignition method according to SFS 3008  
233 (1990) to derive C%. We utilized a similar stock calculation framework using the bulk density, layer depth, and  
234 C% and N% data as in Kemppinen et al. (2021) except we used average bulk density and mineral C% estimates  
235 in each vegetation type in case that information was missing in stock calculation.

236 Soil samples for dissolved organic carbon concentration analyses in dry soil were collected between the 5th and  
237 14th of July 2018. After the collection, samples were stored at 4 °C and then dried at 60 °C for at least 5 days.  
238 Extraction of dissolved organic carbon was done using pure water extractions with 0.5 to 3 grams of dried soil  
239 added to 40 ml of water following the WEOC protocol from (Hensgens et al., 2021). Extracts were immediately  
240 filtered (0.7µm) using glass fibre filters, diluted, acidified to remove inorganic carbon, and measured on a  
241 Shimadzu TOC V-CPN analyzer set on the Nonpurgeable Organic Carbon mode. We refer to this variable as  
242 dissolved organic carbon.

#### 243 **2.2.5 Remotely sensed data**


244 Remotely sensed optical and light detection and ranging-based (LiDAR) data describing topographic,  
245 vegetation, snow, and surficial deposit conditions was used for upscaling the in-situ measured environmental  
246 variables (Fig. 2, Sect. S4 and Fig. S1).

#### 247 **2.3 Statistical analyses**

248 We investigated the dependencies of GPP, ER, NEE, CH<sub>4</sub> flux, and N<sub>2</sub>O flux on environmental variables using  
249 statistical analyses which included analysis of variance (ANOVA), and machine learning modeling and  
250 prediction. We developed machine learning models, in which we 1) upscaled environmental data (annual soil  
251 temperature, soil temperature, soil moisture, soil C/N, soil organic carbon stock, vegetation type, biomass) using  
252 remotely sensed variables as predictors; 2) modeled GHG fluxes using the environmental data as predictors, and  
253 3) upscaled GHG fluxes using the upscaled environmental data maps at a 2-meter spatial resolution across the  
254 landscape (Fig. 2). This two-step upscaling approach enabled us to focus on the relationships between GHG  
255 fluxes with their physical and ecological, in-situ measured environmental controls instead of the remotely  
256 sensed data that are proxies by nature. We ran all analysis in the R statistical programming environment (R Core  
257 Team 2020; version 4.0.3).

258





259

260 **Figure 2:** The upscaling framework used in this study. We first linked GHG fluxes to the in-situ environmental  
 261 drivers using machine learning models. Then we trained three machine learning models to upscale environmental  
 262 conditions across the landscape using remote sensing data. Then we used the GHG flux models and environmental  
 263 predictions to upscale GHG fluxes across the landscape throughout the entire growing season.

### 264 2.3.1 Analysis of variance (ANOVA)

265 We used one-way ANOVAs to test for vegetation type differences in environmental conditions, GHG fluxes,  
 266 and tested significance using multiple comparisons with a Tukey's honest significant difference test ( $p < 0.05$ ).  
 267  $\text{CH}_4$  flux, soil moisture, soil organic carbon and nitrogen stock, and biomass were not normally distributed, thus  
 268 we used Kruskal-Wallis test instead of ANOVA.

### 269 2.3.2 Machine learning models

270 We modeled our response variables using three machine-learning methods (generalized boosted regression  
 271 models, GBM; random forest, RF, and support vector machine regression, SVM), all of which have been widely  
 272 used in flux upscaling studies (see e.g. (Natali et al., 2019; Peltola et al., 2019; Tramontana et al., 2016). These  
 273 three approaches are non-parametric and can handle linear and non-linear relationships and different data  
 274 distributions. We chose RFs and GBMs because they utilize several decision trees in an ensemble model  
 275 framework and thus avoid overfitting, have high accuracy, are highly adaptable, and are not significantly  
 276 impacted by outliers. We chose SVMs because they are good at generalizing the relationships in the data. Based  
 277 on these models, we visualized the partial dependence plots characterizing the relationships between the  
 278 response and predictor variables while accounting for the average effect of the other predictors in the model  
 279 using the pdp package (Greenwell, 2017). Further, we calculated variable importance using the vip package  
 280 (Greenwell et al., 2020). Variable importance scores were estimated by randomly permuting the values of the  
 281 predictor in the training data and exploring how this influenced model performance based on the adjusted  $R^2$   
 282 values, with the idea that random permutation would decrease model performance (Breiman, 2001). We used  
 283 100 simulations to calculate 100 importance scores which were averaged. A standard deviation across these

284 scores was used as an uncertainty estimate, together with the differences in average importance across models.  
285 For more details, see Sect. S5.

286 We used ten topography, snow, vegetation, and surficial deposits variables to construct landscape-wide  
287 predictors matching the in-situ environmental conditions that we used to model the GHG flux values. These  
288 variables were the following: elevation, topographic wetness index, topographic position index at 5 and 30 m  
289 radii, aspect, slope, potential incoming solar radiation, normalized difference vegetation index, snow cover  
290 duration, and surface deposits. Soil organic carbon stocks, dissolved organic carbon, soil C/N, biomass, and  
291 annual soil temperature models were calibrated only once and a single prediction was made to the landscape.  
292 Soil temperatures and moisture vary throughout the growing season, thus, we calibrated each model at each time  
293 step and created 231 predictions over the study period (every 2 hours between 8 am and 8 pm from July 1st until  
294 August 2nd). For each variable, an ensemble prediction was produced by calculating a median prediction over  
295 the three predictions from the different modeling methods.

296 We examined the relationship between the five primary response variables (GPP, ER, NEE, CH<sub>4</sub> flux, N<sub>2</sub>O flux)  
297 and environmental predictors that describe (i) soil resources and conditions (soil moisture, soil C/N, soil pH)  
298 which are relevant to, for example, the growth of organisms (Nobrega and Grogan, 2008; Happonen et al.,  
299 2022); (ii) soil C and N stocks and dissolved organic carbon which are one of the main sources for the GHG  
300 emissions (Bradley-Cook and Virginia, 2018); (iii) soil temperatures which regulate enzymatic processes (St  
301 Pierre et al., 2019; Mauritz et al., 2017); and (iv) biomass and vegetation type which describe resource-use  
302 strategies, carbon inputs to soils and plant photosynthetic capacity, and integrate multiple environmental  
303 properties into one variable (Magnani et al., 2022). We excluded soil pH and soil nitrogen stock from modeling  
304 analyses due to high correlations (Pearson's  $r > 0.7$ ) with soil moisture and soil organic carbon stock,  
305 respectively. Further, dissolved organic carbon was excluded due to its low importance in all the models. We  
306 did not use air temperature as a predictor as we already controlled for it in CO<sub>2</sub> fluxes in the light-response  
307 model, and we assumed that soil microbes regulating CH<sub>4</sub> and N<sub>2</sub>O cycling are most importantly driven by soil  
308 temperatures (Kuhn et al., 2021). The final predictors for our models were soil moisture, soil temperature,  
309 annual soil temperature, soil organic carbon stock, soil C/N, biomass, and vegetation type. The machine learning  
310 parameters tuned for each model can be found from Sect. S5.

311 We used the machine learning models to predict GHG fluxes across the landscape for each 2-hour time step  
312 from July 1st until August 2nd. Similar to the environmental predictions, an ensemble prediction was produced  
313 by calculating a median prediction over the three predictions from the different modeling methods. As our focus  
314 was on understanding the spatial patterns in the mean growing season fluxes, we averaged GHG flux predictions  
315 over the study period. However, a visualization of the predicted mean daily patterns in soil moisture and  
316 temperatures, and the consequent GHG fluxes is provided in the supplementary material (Fig. S2).  
317 To compare the magnitude of all three important GHGs, namely CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O, we calculated the radiative  
318 forcing strength of the three GHGs over a 100-year period from our measurements and ensemble predictions.  
319 We used the Global Warming Potential (GWP; 27 for CH<sub>4</sub> and 273 for N<sub>2</sub>O (IPCC 2021)) and sustained GWP  
320 (45 for CH<sub>4</sub> and 270 for N<sub>2</sub>O (Neubauer 2015)), which are, to our knowledge, the best and most widely used  
321 approaches that exist to compare flux magnitudes. We acknowledge that these approaches are designed to

322 quantify an effect of a change in emission to the radiative forcing, and are thus not fully suitable to be used to  
323 quantify the climatic effect of natural continuous fluxes in our study (Mathijssen et al., 2022; Frohling et al.,  
324 2006).

325

326 For all of our models, we used a leave-one-plot-out cross validation scheme in which each plot was iteratively  
327 left out from the data set, and the remaining data were used to predict fluxes for the excluded plot to assess the  
328 predictive performance of the models (Bodesheim et al., 2018). Estimates of bias were calculated as an average  
329 of the absolute error (MAE) between prediction and actual observation. Coefficient of determination ( $R^2$ ) was  
330 used to determine the strength of the linear relationship between the observed and predicted fluxes.

331 The root mean squared error (RMSE) was used to estimate the model error. The same evaluation metrics were  
332 also calculated based on the prediction to the full model training data to represent model fit (Virkkala et al.  
333 2021); see table S3 which presents these for the individual models. Uncertainty in GHG flux predictions was  
334 derived by bootstrapping (fractional resampling with replacement based on vegetation type classes). We subset  
335 the model training data into 30 different data sets, all of which had the same number of observations as the  
336 original data itself. These 30 data sets were then used to produce 30 individual predictions for a subset of the  
337 times with all three machine learning models and their ensemble for each response variable (Sect. S5). The  
338 uncertainty estimates represent how different distributions of the input data as well as model parameters  
339 influence the upscaled flux maps.

340


## 341 **3 Results**

### 342 **3.1 Environmental conditions and GHG fluxes across vegetation types**

343 We observed large variability in GHG fluxes and environmental conditions within and across vegetation types  
344 (Fig. 3, Table S2). The variability within the vegetation types differed depending on the flux and environmental  
345 variable considered (e.g., meadow class had large variability in GPP and evergreen shrub class in soil C/N).  
346 Frequently, wetlands differed clearly from the other vegetation types. While wetlands had high  $\text{CH}_4$  emissions,  
347 all the other vegetation types with significantly lower soil moisture showed  $\text{CH}_4$  uptake. Meadows were a  
348 significantly larger net  $\text{CO}_2$  sink than evergreen shrub sites, while other vegetation types had intermediate NEE  
349 values. The  $\text{N}_2\text{O}$  fluxes were low from all vegetation types, and varied between small sinks and small sources.

350

351




352  
353  
354  
355  
356  
357  
358

**Figure 3:** The vegetation types considered in this study (a), the distribution of GHG fluxes (b), and environmental conditions (c) across the vegetation types. Lines represent Tukey's test results (\* =  $p \leq 0.05$ , \*\* =  $p \leq 0.01$ , \*\*\* =  $p \leq 0.001$ ). The box corresponds to the 25th and 75th percentiles, and the line within the box represents the median. The lines denote the 1.5 IQR of the lower and higher quartile, where IQR is the inter-quartile range, or distance between the first and third quartiles.

### 359 3.2 The performance of environmental and greenhouse gas flux models

360 The predictive performance of the ensemble environmental variable models was rather high but varied  
 361 depending on the variable ( $R^2$ : 0.09-0.71; Fig. S3). The predictive performance of the GHG models was for  
 362 most variables lower ( $R^2$ : 0.00-0.52), with  $\text{N}_2\text{O}$  flux models being close to random and GPP models performing  
 363 the best (Fig. 4). Model fit was significantly higher than predictive performance for all the fluxes (Table S3).  
 364 The scatterplots of observed and cross-validation-based predicted GHG fluxes suggest that the highest flux  
 365 estimates are often predicted most poorly, but the mean fluxes in each vegetation type were predicted  
 366 accurately, as expected.

367  
368



369


370 **Figure 4:** The correlation between observed and predicted values based on the ensemble model predictions (i.e.,  
 371 median of the three machine learning model outputs). Model predictive performance is described with mean  
 372 absolute error (MAE), R<sup>2</sup> (Rsquared), and RMSE (root mean square error).

### 373 3.3 Drivers of greenhouse gas fluxes

374 The most important controlling variables and the response shapes differed depending on the GHG flux (Fig. 5,  
 375 Fig. 6 and Fig. S4), and sometimes also depending on the machine learning model type applied. CO<sub>2</sub> fluxes  
 376 were driven by annual average soil temperature, biomass, and vegetation type. In addition, soil organic carbon  
 377 stocks were an important predictor for ER. Soil moisture and vegetation type were the most important predictors  
 378 for CH<sub>4</sub> fluxes, and soil C/N and soil moisture for N<sub>2</sub>O fluxes. In general, warmer and wetter conditions  
 379 increased net emissions of CH<sub>4</sub> and N<sub>2</sub>O and net uptake of CO<sub>2</sub>. Some fluxes were further positively correlated  
 380 with soil organic carbon stocks (ER, CH<sub>4</sub> flux) and negatively with soil C/N (GPP, ER, N<sub>2</sub>O). The importance  
 381 for variables explaining the N<sub>2</sub>O flux is low because the model predictive performance is close to random.


382

383



384  
 385  
 386  
 387

**Figure 5:** The variable importance of the environmental variables used to predict GHG fluxes. The models were generalized boosted regression models (GBM), random forest (RF), and support vector machine regression (SVM).



388

389

390

391


**Figure 6:** Partial dependence plots showing the relationships between GHG fluxes and environmental conditions across the three models (generalized boosted regression models, GBM; random forest, RF; and support vector machine regression, SVM). The y-axis of the plot (yhat) represents the marginal effect of the

392 predictor on the response and should not be directly compared with observed or predicted values, rather the  
393 shape and direction of the response instead. RFs and GBMs are based on decision trees, where trees are split  
394 based on a certain threshold in the data, which can be seen as thresholds in the partial dependence plots as well.  
395 SVMs map the data into a high-dimensional space where a hyperplane is fit to separate them, creating smoother  
396 response shapes. Partial dependence plots for GPP and ER are found in Fig. S4.


### 397 **3.4 Spatial patterns and contributions in greenhouse gas flux predictions**

398 The model predictions show large spatial variability in GHG fluxes across the landscape (Fig. 7, Fig. S5). Net  
399 CO<sub>2</sub> uptake as well as GPP and ER were highest in the warm and productive meadow locations of the valley  
400 whereas CH<sub>4</sub> and N<sub>2</sub>O fluxes were highest in the eastern parts of the landscape that is dominated by wetlands.  
401 The prediction suggests small but widespread net CH<sub>4</sub> uptake across the entire upland region. CO<sub>2</sub> was the most  
402 important flux contributing to the net GHG sink (Fig. 8). Mean fluxes calculated based on the upscaled flux  
403 maps differ from the in-situ based ones, particularly for wetland CH<sub>4</sub> emissions (Fig. 8, Fig. S6).





405 **Figure 7:** Ensemble predictions of growing season GHG fluxes, averaged over the 1st of July to the 2nd of  
 406 August (only daytime variability between 8 am and 8 pm considered) and photographs summarizing the main  
 407 sink-source patterns in the landscape. Note that the southwestern corner of the study design has mountain birch  
 408 forest for which we did not have any data; we did not have measurements from the northeastern corner either.  
 409  
 410  
 411



412  
 413  
 414 **Figure 8:** Growing season mean and percentile (0.025 and 0.975) GHG fluxes in CO<sub>2</sub> equivalents based on in-  
 415 situ data and upscaled flux predictions, averaged across the entire study period (only daytime variability  
 416 between 8 am and 8 pm considered) and across vegetation types. Note that the scale for the x axis is different for  
 417 each gas species, and that the uncertainties in in-situ versus predicted mean fluxes cannot be directly compared  
 418 with each other. The uncertainty in in-situ wetland CH<sub>4</sub> continues up to 6.7 but was cropped for visualization  
 419 purposes. The same graph using the sustained GWP approach can be found in the Supplement Fig. 6 and  
 420 demonstrates the potentially larger role of CH<sub>4</sub> fluxes over a 100-year horizon in this landscape.

## 421 4 Discussion

### 422 4.1 CO<sub>2</sub> fluxes driven by both biotic and abiotic variables

423 Our results show the importance of several environmental variables for CO<sub>2</sub> fluxes, demonstrating the strong  
 424 dependence of GPP and ER on a wide range of soil microclimatological, hydrological, soil biogeochemical, and  
 425 ecological processes (Sørensen et al., 2019; Dagg and Lafleur, 2011; Nobrega and Grogan, 2008; Cahoon et al.,  
 426 2016). Overall, the relationships with environmental conditions and GPP and ER were rather similar. Biomass  
 427 was a more important predictor than vegetation type for all the CO<sub>2</sub> fluxes, indicating that the quantity of plant  
 428 material producing and emitting carbon was potentially more important than the different types of plants  
 429 associated with CO<sub>2</sub> cycling in this study setting (Happonen et al., 2022). The high importance of plant-related  
 430 variables (e.g., leaf area index) as drivers of spatial variability in CO<sub>2</sub> fluxes has been previously found in other  
 431 tundra landscapes (Marushchak et al. 2013 and references therein).

432  
 433 Our models also show that annual soil temperatures have a different and stronger relationship with CO<sub>2</sub> fluxes  
 434 than instantaneous growing season soil temperatures, and these two soil temperature variables are indeed not  
 435 strongly correlated with each other in this study design. This is because annual soil temperatures are driven by

436 winter soil temperatures which increase with thicker snow cover that is found particularly in the valley and in  
437 microtopographic depressions, which are colder in the summer. Moreover, annual soil temperatures integrate  
438 many other environmental conditions from the entire year: they reflect growing season length and temperature  
439 conditions, regulate C and N availability, and control vegetation and microbial community composition and  
440 functioning over long time scales. These conditions have been shown to be important drivers of CO<sub>2</sub> fluxes  
441 across a range of Arctic sites (Zona et al., 2022; Lund et al., 2010). Similar to these previous studies, we  
442 observed that plots with warmer annual soil conditions have larger growing season GPP and ER fluxes and  
443 stronger net uptake. Our results also show other logical relationships between environmental conditions and CO<sub>2</sub>  
444 fluxes. For example, GPP and ER increased with soil moisture (Nobrega and Grogan, 2008). However, at  
445 around 50-60 % VWC this relationship plateaued and turned negative. This was likely due to the lack of oxygen  
446 for plant roots restricting growth of non-aerenchymous plants. Furthermore, the lack of oxygen allows only  
447 anoxic metabolic pathways for microbes, such as CH<sub>4</sub> production in methanogenesis, where CO<sub>2</sub> production is  
448 low (Bridgham et al., 2013). Further, soil organic carbon stock was an important predictor for ER, but not so  
449 much for GPP. This was likely related to the higher soil carbon contents boosting decomposition (Schlesinger  
450 and Andrews, 2000).

#### 451 **4.2 Small but consistent net CH<sub>4</sub> uptake mostly driven by soil moisture**

452 Net CH<sub>4</sub> flux was strongly controlled by soil moisture due to its effect on regulating the anoxic and oxic soil  
453 conditions, and therefore CH<sub>4</sub> production (methanogenesis) and CH<sub>4</sub> consumption (CH<sub>4</sub> oxidation, or  
454 methanotrophy) (Kelsey et al., 2016; Christensen et al., 1996; Treat et al., 2018b). Our results demonstrate that  
455 the rate of CH<sub>4</sub> emissions increases sharply in water-logged soil conditions, i.e. at soil moisture levels of > 60  
456 VWC% (Vainio et al. 2021). In drier conditions (VWC < 60%), soils contain more oxygen, which prevents CH<sub>4</sub>  
457 production and increases net CH<sub>4</sub> uptake. This result supports findings from recent studies that show that drier  
458 upland tundra areas can be habitats for methane oxidizing bacteria which can use CH<sub>4</sub> from the atmosphere as  
459 their main energy source, transforming these environments to net CH<sub>4</sub> sinks (Christiansen et al. 2015; Juncher  
460 Jørgensen et al. 2015; Lau et al. 2015; Emmerton et al. 2014; Wagner et al. 2019; St Pierre et al. 2019; Voigt et  
461 al. 2023). Given the large area of the Arctic, even minor fluxes such as those observed here for CH<sub>4</sub> uptake can  
462 be of global importance. This CH<sub>4</sub> uptake can strengthen the GHG sink of the Arctic and prevent CH<sub>4</sub> from  
463 entering the atmosphere.

464  
465 Our results show that net CH<sub>4</sub> uptake increases not only in drier conditions but also in soils with low C/N and  
466 soil organic carbon stocks. This is likely due to microbes needing and getting C and energy from the atmosphere  
467 due to limited soil C supply (Lau et al., 2015; Juutinen et al., 2022), and the capability of methanotrophs to  
468 effectively compete against classical heterotrophs dependent on larger organic macromolecules in these  
469 environments. The models did not clearly identify a particular vegetation type controlling net CH<sub>4</sub> uptake,  
470 however some individual models demonstrated deciduous shrubs and meadows to be more closely related to net  
471 CH<sub>4</sub> uptake (Larmola et al., 2010). Overall, our results indicate that net CH<sub>4</sub> uptake potential is present in any  
472 kind of upland tundra vegetation type (Fig. S7) as long as the abiotic conditions for microbes responsible for  
473 atmospheric CH<sub>4</sub> consumption are favourable.

474 Methane fluxes had a rather uniform distribution across the mineral upland regions (i.e., small but consistent net  
475 uptake). High CH<sub>4</sub> emissions were located in wetland regions dominated by high soil organic carbon stocks and  
476 moisture levels. Our observations demonstrated similar, or even higher net CH<sub>4</sub> uptake than previous studies.  
477 For example, dry tundra was CH<sub>4</sub> neutral in a recent Arctic-Boreal CH<sub>4</sub> flux synthesis (mean=3.83, median= -  
478 0.01 mg CH<sub>4</sub> m<sup>-2</sup> d<sup>-1</sup>; primarily based on growing season daytime fluxes; (Kuhn et al., 2021) whereas our study  
479 showed higher uptake rates for the non-wetland plots (mean=-2.05, median=-1.81 mg CH<sub>4</sub> m<sup>-2</sup> d<sup>-1</sup>). However,  
480 studies focusing on individual sites have recorded similar CH<sub>4</sub> flux magnitudes as observed here (Emmertson et  
481 al., 2014; Lau et al., 2015), but to the best of our knowledge, such extensive spatial patterns in CH<sub>4</sub> flux uptake  
482 using fine spatial resolution models as presented here have not been published so far.

### 483 **4.3 N<sub>2</sub>O fluxes remain neglectable and unpredictable**

484 We observed moderate, and to a large extent unpredictable variability in N<sub>2</sub>O fluxes in this landscape. The  
485 differences in average fluxes between the vegetation types were small. Based on our observations, most  
486 vegetation types were on average N<sub>2</sub>O sinks or neutral but deciduous and evergreen shrubs and meadows had  
487 some variability from moderate N<sub>2</sub>O sinks (up to -200 μg N m<sup>-2</sup> d<sup>-1</sup>) to moderate N<sub>2</sub>O sources (up to 400 μg N  
488 m<sup>-2</sup> d<sup>-1</sup>). Overall, our average N<sub>2</sub>O fluxes were close to zero and thus low in the light of the recent review (Voigt  
489 et al., 2020), which demonstrated that vegetated soils in permafrost regions are often small but evident sources  
490 of N<sub>2</sub>O during the growing season (~30 μg N m<sup>-2</sup> d<sup>-1</sup>), and that barren or sparsely vegetated soils serve as  
491 substantial sources of N<sub>2</sub>O (~455 μg N m<sup>-2</sup> d<sup>-1</sup>). The relatively small N<sub>2</sub>O fluxes observed here can be explained  
492 by the nitrogen-limited nature of the studied soils and the strong competition between plants and microbes for  
493 nutrients: with shallow soils and low stocks of soil organic nitrogen, nitrogen release in labile forms by  
494 mineralization remains low (Voigt et al., 2020). Most of the data in the synthesis came from ecosystems that are  
495 not as much nitrogen-limited as our site (e.g., peatlands, grasslands).

496  
497 We were unable to explain the patterns in N<sub>2</sub>O fluxes with the predictors used here. This was likely related to  
498 the relatively low variability in N<sub>2</sub>O fluxes in most of the plots in general, and the complexity of the soil  
499 microbial N cycle, where N<sub>2</sub>O is produced (nitrification, denitrification, DNRA) and consumed (denitrification)  
500 by multiple co-occurring processes, differently regulated by environmental variables (Butterbach-Bahl et al.,  
501 2013). Nevertheless, the most important driver of N<sub>2</sub>O flux was soil C/N, and the models suggested that lower  
502 C/N ratios were linked to higher net N<sub>2</sub>O emissions. This is expected as the excess soil N in soils with low C:N  
503 ratio allows more rapid N mineralization, nitrification and denitrification as compared to microbial  
504 immobilisation which accelerates N<sub>2</sub>O emissions (Klemetsson et al., 2005; Liimatainen et al., 2018). Further,  
505 N<sub>2</sub>O emissions were highest in the wetlands, similar to (Ma et al., 2007) who explained this by high ammonia or  
506 nitrate levels boosting N<sub>2</sub>O production. The uppermost soil layers were also likely not fully saturated by water at  
507 the time of the wetland measurements, which can induce higher N<sub>2</sub>O emissions in oxic but still moist conditions,  
508 which allow aerobic nitrification and anaerobic denitrification to co-occur (Voigt et al., 2020; Takakai et al.,  
509 2008). In contrast to C fluxes, vegetation type did not play an important role for N<sub>2</sub>O fluxes. This might be  
510 related to our study having no measurements in the previously observed, clear N<sub>2</sub>O flux hot spots located in  
511 barren permafrost peatlands, such as peat plateaus or palsas, with thick organic layers and high inorganic N  
512 content (Repo et al., 2009; Voigt et al., 2017a).

#### 513 **4.4 The sub-Arctic tundra landscape is a strong growing season GHG sink**

514 Our results demonstrate a high level of spatial heterogeneity in the growing season GHG fluxes across the  
515 landscape, with areas acting as both net CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O sinks and sources in some parts of it. Areas acting  
516 as GHG sinks covered most of the landscape (CO<sub>2</sub>: 91 %, CH<sub>4</sub>: 87 %, N<sub>2</sub>O: 73 %; 62 % of the area was a sink  
517 for all the three GHGs). We observed clear differences in flux magnitudes driven by key environmental  
518 conditions. Moist, and carbon and nitrogen-rich meadows and deciduous shrub heaths were strong GHG sinks.  
519 Wet sedge-dominated fens were GHG sinks with CH<sub>4</sub> emissions being compensated by net CO<sub>2</sub> uptake. Barren  
520 lands and evergreen shrubs were more resource-limited and closer to GHG neutral. These results are interesting  
521 in the light of the shrubification patterns observed across the entire Arctic (Myers-Smith et al., 2011; Parker et  
522 al., 2015; Vowles and Björk, 2018), and indicate that deciduous or evergreen shrub expansion may increase or  
523 decrease the growing season GHG sink. If shrubs expand to meadows, the GHG sink may decrease, whereas if  
524 they invade barren areas, the GHG sink may increase. However, our results did not quantify this change over  
525 time, or cover the entire year to confirm the net annual effect.

526

527 Our results indicate that this heterogeneous Arctic landscape was a cumulative net GHG sink during the  
528 measurement period during daytime (8 am to 8 pm) in July 2018. The July budget for CO<sub>2</sub> was -4.7 g C m<sup>-2</sup>  
529 month<sup>-1</sup>, for CH<sub>4</sub> 0.73 mg C m<sup>-2</sup> month<sup>-1</sup> and for N<sub>2</sub>O -10.0 µg N m<sup>-2</sup> month<sup>-1</sup>. The CO<sub>2</sub> sink is relatively small,  
530 likely due to the high cover of patchy and sparsely vegetated areas that were often CO<sub>2</sub> sources. This small sink  
531 value is an overestimation of the sink activity considering the whole course of the day as we did not have  
532 measurements from the night time and did thus not upscale fluxes in night-time conditions when ecosystems are  
533 net CO<sub>2</sub> sources due to the lack of light required for photosynthesis. It also overestimates the importance of CO<sub>2</sub>  
534 as a radiative forcing agent, since ecosystem CO<sub>2</sub> production during autumn and winter contributes substantially  
535 to the annual C balance (Celis et al., 2017; Commane et al., 2017), thereby reducing the CO<sub>2</sub> sink strength on an  
536 annual basis. Further, CH<sub>4</sub> uptake might continue even in rather cold conditions as long as soils remain dry and  
537 unfrozen (Emmertson et al., 2014). Nevertheless, our results demonstrate that net CO<sub>2</sub> uptake plays the most  
538 important role for the net growing season GHG budget. CH<sub>4</sub> emissions from wetlands are almost balanced by  
539 the net CH<sub>4</sub> uptake of other ecosystems. The role of N<sub>2</sub>O fluxes for the net GHG budget across the entire  
540 landscape is negligible for the growing season.

#### 541 **4.5 Methodological considerations in GHG flux modeling**

542 Our study creates new understanding about high-resolution upscaling of GHG fluxes by incorporating more  
543 chamber measurements, predictors, models, and environmental gradients compared to earlier efforts (Fox et al.,  
544 2008; Dinsmore et al., 2017; Räsänen et al., 2021; Juutinen et al., 2022; Vainio et al., 2021). For example, we  
545 included chamber measurements from 101 plots whereas earlier local-scale upscaling studies have usually had  
546 circa 30 plots. Further, we included eight different environmental predictors while other studies have often used  
547 only one or two, focusing on predictors describing vegetation type or soil moisture. Finally, we studied a tundra  
548 landscape that consists of almost all the main vegetation types of the entire Arctic, whereas earlier studies have  
549 investigated a narrower range of vegetation conditions, with a focus on wet ecosystems. However, at the same  
550 time, our models showed some signs of overfitting as demonstrated by the high model fit performance statistics  
551 and the mismatch between model fit and predictive performance statistics (Supplementary Text S5.3). This is a

552 common issue in upscaling (Kemppinen et al. 2018; Shi et al. 2022), and could indicate that the models have  
553 potentially learned to fit some noise or specific patterns unique to the training set instead of broadly  
554 generalizable relationships. Nevertheless, the relationships we observed were logical and comparable to those  
555 observed in other studies - both based on spatial and time series study designs (e.g., positive soil moisture-CH<sub>4</sub>  
556 flux or soil temperature-ER relationships (Euskirchen et al. 2014; Davidson et al. 2016; Zona et al. 2023)).  
557 Moreover, our study is based on a dataset focusing on spatial variation in GHG fluxes and correlations between  
558 variables. Therefore, the dataset should not directly be used to infer causal relationships or estimates of flux  
559 change over time (Damgaard 2019), and we advise caution when extrapolating these results to areas outside our  
560 study domain or different time periods.

561

562 Our study showed that using means of in-situ GHG fluxes in each vegetation class to derive a landscape-level  
563 GHG budget might produce significantly different results compared to the upscaled budget. This was apparent  
564 particularly for CH<sub>4</sub> fluxes, where the in-situ based average wetland CH<sub>4</sub> emission was more than two times  
565 larger CH<sub>4</sub> compared to the upscaled one. This mismatch is likely explained by the heterogeneity of  
566 environmental conditions and CH<sub>4</sub> fluxes within the wetland class that the chamber measurements alone could  
567 not cover (Fig. S7). A multivariate machine learning modeling approach with variables describing not only  
568 vegetation type but also soil moisture and other conditions were likely able to characterize the resulting CH<sub>4</sub>  
569 flux variability in a more representative way. For example, our soil moisture maps showed high variation in soil  
570 moisture between ca. 50 and 70 VWC% within the wetland areas, and high CH<sub>4</sub> emissions were observed only  
571 in areas with 60 VWC%. Overall, this result suggests that simple land cover-based upscaling efforts might lead  
572 to biased budget estimates, especially when spatial variability within land cover types is high, emphasizing the  
573 need for multivariate modeling in flux upscaling.

574

575 The performance of our models varied from good (GPP, CH<sub>4</sub> flux), moderate (ER and NEE) to low (N<sub>2</sub>O). CH<sub>4</sub>  
576 fluxes - both sources and sinks - were most accurately modeled, providing important support for future studies  
577 predicting not only the large CH<sub>4</sub> emissions but also the previously unquantified CH<sub>4</sub> uptake in Arctic  
578 landscapes. The lower predictive performance of the models for other GHG fluxes might be explained by the  
579 dynamic nature of fluxes not being represented in our spatial study design with no temporal chamber replicates  
580 in the plots, our models lacking important predictors, or our model structure not being ideal. The performance of  
581 the models could potentially be improved by describing plant functional composition using plant traits  
582 (Happonen et al., 2022), and including more detailed information about soil nutrients (e.g., soil nitrate or  
583 ammonium concentrations as soil C/N captures only very roughly how much N is available) or microbial  
584 communities (e.g., communities or genes associated with nitrification or methanogenesis or methanotrophy;  
585 (Pessi et al., 2022)).

586

587 Rainfall events are another source of uncertainty in our upscaling because they might increase soil moisture  
588 levels and activate processes related to methanogenesis, photosynthesis and respiration as well as nitrogen  
589 cycling. While our soil moisture predictions should capture these variations in soil wetness, we only made  
590 measurements once per plot under clear conditions and do not have information about how GHG fluxes might

591 respond to rainfall events. We might thus underestimate some of the instantaneous and longer-term changes in  
592 GHG fluxes during and after rain (see Text S1 and Fig. S10 for details).

593

594 We chose to use in-situ environmental data as predictors of GHG fluxes in our upscaling framework instead of  
595 linking remotely sensed variables with GHG fluxes directly. This was done to increase understanding about the  
596 mechanistic and ecological relationships but required us to first produce spatially continuous maps of  
597 environmental conditions, which might have added an additional layer of uncertainty into our framework.  
598 However, the most important environmental variables (i.e., soil moisture, temperature, biomass) had a high  
599 predictive performance. Nevertheless, future studies could explore the performance and information derived by  
600 upscaling GHG fluxes using high-resolution satellite or drone-derived remotely sensed indices directly (Siewert  
601 and Olofsson, 2020; Vainio et al., 2021; Berner et al., 2018).

602

603 Overall, the performance of our machine learning models predicting spatial variability in GHG fluxes was  
604 weaker than in other studies focusing on temporal variability (e.g., (López-Blanco et al., 2017; Celis et al.,  
605 2017), even though we had a comprehensive set of environmental measurements. Our results thus highlight the  
606 need for more focus on the spatial patterns in GHG fluxes. While the temporal variability is widely  
607 acknowledged as a source of uncertainty in GHG budget estimates (Baldocchi et al., 2018), the spatial  
608 variability may be just as important but remains insufficiently studied (Treat et al., 2018c). Study designs  
609 focusing on spatial variation in GHG fluxes using a combination of intensive measurement campaigns, remotely  
610 sensed datasets, and modeling approaches are informative although they do not produce direct information on  
611 the trends and drivers of GHG flux change following climate change. They provide new knowledge about the  
612 heterogeneity in GHG fluxes and their environmental drivers which is highly important for understanding flux  
613 magnitudes from local to global scales. Further, they can be used as a space-for-time substitution to understand  
614 ecosystem functions in locations that are assumed to be at different stages of development. Moreover, this  
615 knowledge is valuable for designing representative field studies in the future.

616

## 617 **5 Conclusions**

618 This study showed that predicting fluxes in heterogeneous tundra landscapes at high spatial resolutions is  
619 possible for CH<sub>4</sub>, GPP, and to some extent also NEE and ER fluxes but remains a challenge for N<sub>2</sub>O fluxes. This  
620 is a promising result for future high spatial resolution modeling studies that aim to understand the fine-scale  
621 biogeochemistry of the rapidly changing Arctic environments. Our study further demonstrates high spatial  
622 variability of GHG fluxes which is driven by a multitude of vegetation, soil microclimatological, hydrological,  
623 and biogeochemical conditions. The upscaling shows the importance of net CO<sub>2</sub> uptake for the peak growing  
624 season net GHG budget, and suggests that annual soil temperature and vegetation parameters are the most  
625 important drivers. Most importantly, it reveals small but widespread CH<sub>4</sub> uptake across the entire upland tundra  
626 in our domain that almost surpasses the high wetland CH<sub>4</sub> emissions. This provides more evidence to the  
627 relatively unquantified but important CH<sub>4</sub> sink in the Arctic GHG budget.

628

629 **Code/Data availability**

630 The field data, analysis codes and most of the results are available in a repository (Virkkala et al. 2023).  
631 Upscaling results for each individual timestep were not included in the repository due to their large size, but  
632 they can be acquired from the author upon request.

633 **Author contribution**

634 AMV and ML conceptualized the research with input from PN, JK, MEM, and CV. AMV, PN, JK, MEM, CV,  
635 GH, KH, VT, JH and ML contributed to data collection. AMV analyzed the data and wrote the manuscript draft.  
636 All the coauthors reviewed and edited the manuscript.

637 **Competing interests**

638 The authors declare no competing interests.

639 **Acknowledgements**

640 The authors would like to acknowledge the support by the research assistants during the data collection as well  
641 as Kilpisjärvi Biological Station. AMV was supported by The Finnish Cultural Foundation, Alfred Kordelin  
642 Foundation, Väisälä fund, and Jenny and Antti Wihuri Foundation, and the Gordon and Betty Moore foundation  
643 (grant #8414). AMV and ML acknowledge the Academy of Finland funding (grant #286950). AMV and GH  
644 acknowledge the Svenska Sällskapet för Antropologi och Geografi for funding. CV was supported by the  
645 Academy of Finland project MUFFIN (no. 332196). ML acknowledges Academy of Finland funding (grant  
646 #342890). CB acknowledges funding from Academy of Finland general research grant (project N-PERM,  
647 decision Nr. 341348). CB, CV, and MEM acknowledge Academy of Finland/Russian Foundation for Basic  
648 Research project NOCA (decision no. 314630). PN was funded by the Academy of Finland (project number  
649 347558). JK was funded by the Academy of Finland (project number 349606). JH was funded by the Academy  
650 of Finland (grant #308128). We acknowledge funding for fieldwork and equipment by the Nordenskiöld  
651 samfundet, Tiina and Antti Herlin foundation, and Maa- ja vesiteknikan tuki ry. JR was supported by the  
652 Greenhouse Gas Fluxes and Earth System Feedbacks (GreenFeedBack) project from the European Union's  
653 Horizon Europe Framework Programme for Research and Innovation (project no. 101056921), funded by the  
654 European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily  
655 reflect those of the European Union. Neither the European Union nor the granting authority can be held  
656 responsible for them.

657

658 **References**

659 European Space Agency (ESA), E. S.: Land Cover CCI Product User Guide Version 2 Tech. Rep, 2017.

660 Baldocchi, D., Chu, H., and Reichstein, M.: Inter-annual variability of net and gross ecosystem carbon fluxes: A  
661 review, *Agric. For. Meteorol.*, 249, 520–533, <https://doi.org/10.1016/j.agrformet.2017.05.015>, 2018.



- 662 Belshe, E. F., Schuur, E. A. G., and Bolker, B. M.: Tundra ecosystems observed to be CO<sub>2</sub> sources due to  
663 differential amplification of the carbon cycle, *Ecol. Lett.*, 16, 1307–1315, <https://doi.org/10.1111/ele.12164>,  
664 2013.
- 665 Berner, L. T., Jantz, P., Tape, K. D., and Goetz, S. J.: Tundra plant above-ground biomass and shrub dominance  
666 mapped across the North Slope of Alaska, *Environ. Res. Lett.*, 13, 035002, <https://doi.org/10.1088/1748-9326/aaaa9a>, 2018.
- 668 Bradley-Cook, J. I. and Virginia, R. A.: Landscape variation in soil carbon stocks and respiration in an Arctic  
669 tundra ecosystem, west Greenland, *Arct. Antarct. Alp. Res.*, 50, S100024,  
670 <https://doi.org/10.1080/15230430.2017.1420283>, 2018.
- 671 Breiman, L.: Random Forests, *Mach. Learn.*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- 672 Bridgman, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from wetlands:  
673 biogeochemical, microbial, and modeling perspectives from local to global scales, *Glob. Chang. Biol.*, 19,  
674 1325–1346, <https://doi.org/10.1111/gcb.12131>, 2013.
- 675 Bruhwiler, L., Parmentier, F.-J. W., Crill, P., Leonard, M., and Palmer, P. I.: The Arctic Carbon Cycle and Its  
676 Response to Changing Climate, *Current Climate Change Reports*, 7, 14–34, <https://doi.org/10.1007/s40641-020-00169-5>, 2021.
- 678 Brummell, M. E., Farrell, R. E., and Siciliano, S. D.: Greenhouse gas soil production and surface fluxes at a  
679 high arctic polar oasis, *Soil Biol. Biochem.*, 52, 1–12, <https://doi.org/10.1016/j.soilbio.2012.03.019>, 2012.
- 680 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide  
681 emissions from soils: how well do we understand the processes and their controls?, *Philos. Trans. R. Soc. Lond.*  
682 *B Biol. Sci.*, 368, 20130122, <https://doi.org/10.1098/rstb.2013.0122>, 2013.
- 683 Cahoon, S. M. P., Sullivan, P. F., Shaver, G. R., Welker, J. M., Post, E., and Holyoak, M.: Interactions among  
684 shrub cover and the soil microclimate may determine future Arctic carbon budgets, *Ecol. Lett.*, 15, 1415–1422,  
685 <https://doi.org/10.1111/j.1461-0248.2012.01865.x>, 2012a.
- 686 Cahoon, S. M. P., Sullivan, P. F., Post, E., and Welker, J. M.: Large herbivores limit CO<sub>2</sub> uptake and suppress  
687 carbon cycle responses to warming in West Greenland, *Glob. Chang. Biol.*, 18, 469–479,  
688 <https://doi.org/10.1111/j.1365-2486.2011.02528.x>, 2012b.
- 689 Cahoon, S. M. P., Sullivan, P. F., and Post, E.: Greater Abundance of *Betula nana* and Early Onset of the  
690 Growing Season Increase Ecosystem CO<sub>2</sub> Uptake in West Greenland, *Ecosystems*, 19, 1149–1163,  
691 <https://doi.org/10.1007/s10021-016-9997-7>, 2016.
- 692 Celis, G., Mauritz, M., Bracho, R., Salmon, V. G., Webb, E. E., Hutchings, J., Natali, S. M., Schädel, C.,  
693 Crummer, K. G., and Schuur, E. A. G.: Tundra is a consistent source of CO<sub>2</sub> at a site with progressive  
694 permafrost thaw during 6 years of chamber and eddy covariance measurements,  
695 <https://doi.org/10.1002/2016jg003671>, 2017.
- 696 Christensen, T. R., Prentice, I. C., Kaplan, J., Haxeltine, A., and Sitch, S.: Methane flux from northern wetlands  
697 and tundra. An ecosystem source modelling approach, *Tellus B Chem. Phys. Meteorol.*, 48, 652–661,  
698 <https://doi.org/10.1034/j.1600-0889.1996.t01-4-00004.x>, 1996.
- 699 Christiansen, J. R., Romero, A. J. B., Jørgensen, N. O. G., Glaring, M. A., Jørgensen, C. J., Berg, L. K., and  
700 Elberling, B.: Methane fluxes and the functional groups of methanotrophs and methanogens in a young Arctic  
701 landscape on Disko Island, West Greenland, *Biogeochemistry*, 122, 15–33, <https://doi.org/10.1007/s10533-014-0026-7>, 2015.
- 703 Commane, R., Lindaas, J., Benmergui, J., Luus, K. A., Chang, R. Y.-W., Daube, B. C., Euskirchen, E. S.,  
704 Henderson, J. M., Karion, A., Miller, J. B., Miller, S. M., Parazoo, N. C., Randerson, J. T., Sweeney, C., Tans,  
705 P., Thoning, K., Veraverbeke, S., Miller, C. E., and Wofsy, S. C.: Carbon dioxide sources from Alaska driven  
706 by increasing early winter respiration from Arctic tundra, *Proc. Natl. Acad. Sci. U. S. A.*, 114, 5361–5366,  
707 <https://doi.org/10.1073/pnas.1618567114>, 2017.
- 708 Dagg, J. and Lafleur, P.: Vegetation Community, Foliar Nitrogen, and Temperature Effects on Tundra CO<sub>2</sub>

- 709 Exchange across a Soil Moisture Gradient, *Arct. Antarct. Alp. Res.*, 43, 189–197, <https://doi.org/10.1657/1938-4246-43.2.189>, 2011.
- 710
- 711 Davidson, S. J., Santos, M. J., Sloan, V. L., Reuss-Schmidt, K., Phoenix, G. K., Oechel, W. C., and Zona, D.: Upscaling CH<sub>4</sub> Fluxes Using High-Resolution Imagery in Arctic Tundra Ecosystems, *Remote Sensing*, 9, 1227, <https://doi.org/10.3390/rs9121227>, 2017.
- 712
- 713
- 714 Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E. C., Jones, B., Barber, C. V., Hayes, R., Kormos, C., Martin, V., Crist, E., Sechrest, W., Price, L., Baillie, J. E. M., Weeden, D., Suckling, K., Davis, C., Sizer, N., Moore, R., Thau, D., Birch, T., Potapov, P., Turubanova, S., Tyukavina, A., de Souza, N., Pintea, L., Brito, J. C., Llewellyn, O. A., Miller, A. G., Patzelt, A., Ghazanfar, S. A., Timberlake, J., Klöser, H., Shennan-Farpón, Y., Kindt, R., Lillesø, J.-P. B., van Breugel, P., Graudal, L., Voge, M., Al-Shammari, K. F., and Saleem, M.: An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm, *Bioscience*, 67, 534–545, <https://doi.org/10.1093/biosci/bix014>, 2017.
- 715
- 716
- 717
- 718
- 719
- 720
- 721
- 722 Dinsmore, K. J., Drewer, J., Levy, P. E., George, C., Lohila, A., Aurela, M., and Skiba, U. M.: Growing season CH<sub>4</sub> and N<sub>2</sub>O fluxes from a subarctic landscape in northern Finland; from chamber to landscape scale, <https://doi.org/10.5194/bg-14-799-2017>, 2017.
- 723
- 724
- 725 Emmerton, C. A., St. Louis, V. L., Lehnerr, I., Humphreys, E. R., Rydz, E., and Kosolofski, H. R.: The net exchange of methane with high Arctic landscapes during the summer growing season, <https://doi.org/10.5194/bg-11-3095-2014>, 2014.
- 726
- 727
- 728 Euskirchen, E. S., Bret-Harte, M. S., Scott, G. J., Edgar, C., and Shaver, G. R.: Seasonal patterns of carbon dioxide and water fluxes in three representative tundra ecosystems in northern Alaska, *Ecosphere*, 3, art4, <https://doi.org/10.1890/es11-00202.1>, 2012.
- 729
- 730
- 731 Euskirchen, E. S., Edgar, C. W., Turetsky, M. R., Waldrop, M. P., and Harden, J. W.: Differential response of carbon fluxes to climate in three peatland ecosystems that vary in the presence and stability of permafrost, *J. Geophys. Res. Biogeosci.*, 119, 1576–1595, <https://doi.org/10.1002/2014jg002683>, 2014.
- 732
- 733
- 734 Fox, A. M., Huntley, B., Lloyd, C. R., Williams, M., and Baxter, R.: Net ecosystem exchange over heterogeneous Arctic tundra: Scaling between chamber and eddy covariance measurements, *Global Biogeochem. Cycles*, 22, 2008.
- 735
- 736
- 737 Frolking, S., Roulet, N., and Fuglestedt, J.: How northern peatlands influence the Earth's radiative budget: Sustained methane emission versus sustained carbon sequestration, *J. Geophys. Res.*, 111, <https://doi.org/10.1029/2005jg000091>, 2006.
- 738
- 739
- 740 Greenwell, B., Brandon, Greenwell, M., Bradley, and Boehmke, C.: Variable Importance Plots—An Introduction to the vip Package, <https://doi.org/10.32614/rj-2020-013>, 2020.
- 741
- 742 Greenwell, B. M.: pdp: An R Package for Constructing Partial Dependence Plots, *R J.*, 9, 421, 2017.
- 743
- 744
- 745 Happonen, K., Virkkala, A.-M., Kemppinen, J., Niittynen, P., and Luoto, M.: Relationships between above-ground plant traits and carbon cycling in tundra plant communities, *J. Ecol.*, 110, 700–716, 2022.
- 746
- 747
- 748
- 749 Hensgens, G., Laudon, H., Johnson, M. S., and Berggren, M.: The undetected loss of aged carbon from boreal mineral soils, *Sci. Rep.*, 11, 6202, <https://doi.org/10.1038/s41598-021-85506-w>, 2021.
- 750
- 751 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., and Others: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 2014.
- 752
- 753
- 754 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon
- 755

- 756 and nitrogen are vulnerable to permafrost thaw, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 20438–20446,  
757 <https://doi.org/10.1073/pnas.1916387117>, 2020.
- 758 Juncher Jørgensen, C., Lund Johansen, K. M., Westergaard-Nielsen, A., and Elberling, B.: Net regional methane  
759 sink in High Arctic soils of northeast Greenland, *Nat. Geosci.*, 8, 20–23, <https://doi.org/10.1038/ngeo2305>,  
760 2015.
- 761 Juutinen, S., Aurela, M., Tuovinen, J.-P., Ivakhov, V., Linkosalmi, M., Räsänen, A., Virtanen, T., Mikola, J.,  
762 Nyman, J., Vähä, E., Loskutova, M., Makshtas, A., and Laurila, T.: Variation in CO<sub>2</sub> and CH<sub>4</sub> fluxes among  
763 land cover types in heterogeneous Arctic tundra in northeastern Siberia, *Biogeosciences*, 19, 3151–3167,  
764 <https://doi.org/10.5194/bg-19-3151-2022>, 2022.
- 765 Kåresdotter, E., Destouni, G., Ghajarnia, N., Hugelius, G., and Kalantari, Z.: Mapping the vulnerability of arctic  
766 wetlands to global warming, *Earths Future*, 9, <https://doi.org/10.1029/2020ef001858>, 2021.
- 767 Kelsey, K. C., Leffler, A. J., Beard, K. H., Schmutz, J. A., Choi, R. T., and Welker, J. M.: Interactions among  
768 vegetation, climate, and herbivory control greenhouse gas fluxes in a subarctic coastal wetland, *Journal of*  
769 *Geophysical Research: Biogeosciences*, 121, 2960–2975, 2016.
- 770 Kempainen, J., Niittynen, P., Virkkala, A.-M., Happonen, K., Riihimäki, H., Aalto, J., and Luoto, M.: Dwarf  
771 Shrubs Impact Tundra Soils: Drier, Colder, and Less Organic Carbon, *Ecosystems*,  
772 <https://doi.org/10.1007/s10021-020-00589-2>, 2021.
- 773 King, L. and Seppälä, M.: Permafrost thickness and distribution in Finnish Lapland-Results of geoelectrical  
774 soundings, *Polarforschung*, 57, 127–147, 1987.
- 775 Klemetsson, L., Von Arnold, K., Weslien, P., and Gundersen, P.: Soil CN ratio as a scalar parameter to predict  
776 nitrous oxide emissions, *Glob. Chang. Biol.*, 11, 1142–1147, <https://doi.org/10.1111/j.1365-2486.2005.00973.x>,  
777 2005.
- 778 Kuhn, M. A., Varner, R. K., Bastviken, D., Crill, P., MacIntyre, S., Turetsky, M., Walter Anthony, K., McGuire,  
779 A. D., and Olefeldt, D.: BAWLD-CH 4: a comprehensive dataset of methane fluxes from boreal and arctic  
780 ecosystems, *Earth System Science Data*, 13, 5151–5189, 2021.
- 781 Larmola, T., Tuittila, E.-S., Tirola, M., Nykänen, H., Martikainen, P. J., Yrjälä, K., Tuomivirta, T., and Fritze,  
782 H.: The role of Sphagnum mosses in the methane cycling of a boreal mire, *Ecology*, 91, 2356–2365,  
783 <https://doi.org/10.1890/09-1343.1>, 2010.
- 784 Lau, M. C. Y., Stackhouse, B. T., Layton, A. C., Chauhan, A., Vishnivetskaya, T. A., Chourey, K., Ronholm, J.,  
785 Mykityczuk, N. C. S., Bennett, P. C., Lamarche-Gagnon, G., Burton, N., Pollard, W. H., Omelon, C. R.,  
786 Medvigy, D. M., Hettich, R. L., Pfiffner, S. M., Whyte, L. G., and Onstott, T. C.: An active atmospheric  
787 methane sink in high Arctic mineral cryosols, *ISME J.*, 9, 1880–1891, <https://doi.org/10.1038/ismej.2015.13>,  
788 2015.
- 789 Liimatainen, M., Voigt, C., Martikainen, P. J., Hytönen, J., Regina, K., Óskarsson, H., and Maljanen, M.:  
790 Factors controlling nitrous oxide emissions from managed northern peat soils with low carbon to nitrogen ratio,  
791 *Soil Biol. Biochem.*, 122, 186–195, <https://doi.org/10.1016/j.soilbio.2018.04.006>, 2018.
- 792 Livingston, G. P. and Hutchinson, G. L.: Enclosure-based measurement of trace gas exchange: applications  
793 and sources of error.[Bokförf.], 1995.
- 794 López-Blanco, E., Lund, M., Williams, M., Tamstorf, M. P., Westergaard-Nielsen, A., Exbrayat, J.-F., Hansen,  
795 B. U., and Christensen, T. R.: Exchange of CO<sub>2</sub> in Arctic tundra: impacts of meteorological variations and  
796 biological disturbance, *Biogeosciences*, 14, 4467, 2017.
- 797 Lund, M., Lafleur, P. M., Roulet, N. T., Lindroth, A., Christensen, T. R., Aurela, M., Chojnicki, B. H.,  
798 Flanagan, L. B., Humphreys, E. R., Laurila, T., Oechel, W. C., Olejnik, J., Rinne, J., Schubert, P., and Nilsson,  
799 M. B.: Variability in exchange of CO<sub>2</sub> across 12 northern peatland and tundra sites, *Glob. Chang. Biol.*, 16,  
800 2436–2448, <https://doi.org/10.1111/j.1365-2486.2009.02104.x>, 2010.
- 801 Magnani, M., Baneschi, I., Giamberini, M., Raco, B., and Provenzale, A.: Microscale drivers of summer CO<sub>2</sub>  
802 fluxes in the Svalbard High Arctic tundra, *Sci. Rep.*, 12, 763, <https://doi.org/10.1038/s41598-021-04728-0>,

- 803 2022.
- 804 Masyagina, O. V. and Menyailo, O. V.: The impact of permafrost on carbon dioxide and methane fluxes in  
805 Siberia: A meta-analysis, *Environ. Res.*, 182, 109096, <https://doi.org/10.1016/j.envres.2019.109096>, 2020.
- 806 Mathijssen, P. J. H., Tuovinen, J.-P., Lohila, A., Väiranta, M., and Tuittila, E.-S.: Identifying main uncertainties  
807 in estimating past and present radiative forcing of peatlands, *Glob. Chang. Biol.*, 28, 4069–4084,  
808 <https://doi.org/10.1111/gcb.16189>, 2022.
- 809 Mauritz, M., Bracho, R., Celis, G., Hutchings, J., Natali, S. M., Pegoraro, E., Salmon, V. G., Schädel, C., Webb,  
810 E. E., and Schuur, E. A. G.: Nonlinear CO<sub>2</sub> flux response to 7 years of experimentally induced permafrost thaw,  
811 *Glob. Chang. Biol.*, 23, 3646–3666, <https://doi.org/10.1111/gcb.13661>, 2017.
- 812 Ma, W. K., Schautz, A., Fishback, L.-A. E., Bedard-Haughn, A., Farrell, R. E., and Siciliano, S. D.: Assessing  
813 the potential of ammonia oxidizing bacteria to produce nitrous oxide in soils of a high arctic lowland ecosystem  
814 on Devon Island, Canada, *Soil Biol. Biochem.*, 39, 2001–2013, <https://doi.org/10.1016/j.soilbio.2007.03.001>,  
815 2007.
- 816 McGuire, A. D., Christensen, T. R., Hayes, D. J., Heroult, A., Euskirchen, E., Yi, Y., Kimball, J. S., Koven, C.,  
817 Lafleur, P., Miller, P. A., Oechel, W., Peylin, P., and Williams, M.: An assessment of the carbon balance of  
818 arctic tundra: comparisons among observations, process models, and atmospheric inversions, *Biogeosci.*  
819 *Discuss.*, 9, 4543, <https://doi.org/10.5194/bg-9-3185-2012>, 2012.
- 820 Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K. D., Macias-  
821 Fauria, M., Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P., Hermanutz, L., Trant, A., Collier, L. S.,  
822 Weijers, S., Rozema, J., Rayback, S. A., Schmidt, N. M., Schaepman-Strub, G., Wipf, S., Rixen, C., Ménard, C.  
823 B., Venn, S., Goetz, S., Andreu-Hayles, L., Elmendorf, S., Ravolainen, V., Welker, J., Grogan, P., Epstein, H.  
824 E., and Hik, D. S.: Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities,  
825 <https://doi.org/10.1088/1748-9326/6/4/045509>, 2011.
- 826 Natali, S. M., Watts, J. D., Rogers, B. M., Potter, S., Ludwig, S. M., Selbmann, A.-K., Sullivan, P. F., Abbott, B.  
827 W., Arndt, K. A., Birch, L., Björkman, M. P., Bloom, A. A., Celis, G., Christensen, T. R., Christiansen, C. T.,  
828 Commane, R., Cooper, E. J., Crill, P., Czimczik, C., Davydov, S., Du, J., Egan, J. E., Elberling, B., Euskirchen,  
829 E. S., Friborg, T., Genet, H., Göckede, M., Goodrich, J. P., Grogan, P., Helbig, M., Jafarov, E. E., Jastrow, J. D.,  
830 Kalhori, A. A. M., Kim, Y., Kimball, J. S., Kutzbach, L., Lara, M. J., Larsen, K. S., Lee, B.-Y., Liu, Z., Loranty,  
831 M. M., Lund, M., Lupascu, M., Madani, N., Malhotra, A., Matamala, R., McFarland, J., McGuire, A. D.,  
832 Michelsen, A., Minions, C., Oechel, W. C., Olefeldt, D., Parmentier, F.-J. W., Pirk, N., Poulter, B., Quinton, W.,  
833 Rezanezhad, F., Risk, D., Sachs, T., Schaefer, K., Schmidt, N. M., Schuur, E. A. G., Semenchuk, P. R., Shaver,  
834 G., Sonntag, O., Starr, G., Treat, C. C., Waldrop, M. P., Wang, Y., Welker, J., Wille, C., Xu, X., Zhang, Z.,  
835 Zhuang, Q., and Zona, D.: Large loss of CO<sub>2</sub> in winter observed across the northern permafrost region, *Nat.*  
836 *Clim. Chang.*, 9, 852–857, <https://doi.org/10.1038/s41558-019-0592-8>, 2019.
- 837 Nobrega, S. and Grogan, P.: Landscape and ecosystem-level controls on net carbon dioxide exchange along a  
838 natural moisture gradient in Canadian low arctic tundra, *Ecosystems*, 11, 377–396,  
839 <https://doi.org/10.1007/s10021-008-9128-1>, 2008.
- 840 Olefeldt, D., Hovemyr, M., Kuhn, M. A., Bastviken, D., Bohn, T. J., Connolly, J., Crill, P., Euskirchen, E. S.,  
841 Finkelstein, S. A., Genet, H., Grosse, G., Harris, L. I., Heffernan, L., Helbig, M., Hugelius, G., Hutchins, R.,  
842 Juutinen, S., Lara, M. J., Malhotra, A., Manies, K., McGuire, A. D., Natali, S. M., O'Donnell, J. A., Parmentier,  
843 F.-J. W., Räsänen, A., Schädel, C., Sonntag, O., Strack, M., Tank, S. E., Treat, C., Varner, R. K., Virtanen, T.,  
844 Warren, R. K., and Watts, J. D.: The boreal–Arctic Wetland and Lake Dataset (BAWLD), *Earth Syst. Sci. Data*,  
845 13, 5127–5149, <https://doi.org/10.5194/essd-13-5127-2021>, 2021.
- 846 Pallandt, M., Kumar, J., Mauritz, M., Schuur, E., Virkkala, A.-M., Celis, G., Hoffman, F., and Göckede, M.:  
847 Representativeness assessment of the pan-Arctic eddy-covariance site network, and optimized future  
848 enhancements, <https://doi.org/10.5194/bg-2021-133>, 2021.
- 849 Parker, T. C., Subke, J.-A., and Wookey, P. A.: Rapid carbon turnover beneath shrub and tree vegetation is  
850 associated with low soil carbon stocks at a subarctic treeline, *Glob. Chang. Biol.*, 21, 2070–2081,  
851 <https://doi.org/10.1111/gcb.12793>, 2015.
- 852 Peltola, O., Vesala, T., Gao, Y., Rätty, O., Alekseychik, P., Aurela, M., Chojnicki, B., Desai, A. R., Dolman, A.

- 853 J., Euskirchen, E. S., Friborg, T., Göckede, M., Helbig, M., Humphreys, E., Jackson, R. B., Jocher, G., Joos, F.,  
854 Klatt, J., Knox, S. H., Kowalska, N., Kutzbach, L., Lienert, S., Lohila, A., Mammarella, I., Nadeau, D. F.,  
855 Nilsson, M. B., Oechel, W. C., Peichl, M., Pypker, T., Quinton, W., Rinne, J., Sachs, T., Samson, M., Schmid,  
856 H. P., Sonntag, O., Wille, C., Zona, D., and Aalto, T.: Monthly gridded data product of northern wetland  
857 methane emissions based on upscaling eddy covariance observations, *Earth System Science Data*, 11, 1263–  
858 1289, <https://doi.org/10.5194/essd-11-1263-2019>, 2019.
- 859 Pessi, I. S., Viitamäki, S., Virkkala, A.-M., Eronen-Rasimus, E., Delmont, T. O., Marushchak, M. E., Luoto, M.,  
860 and Hultman, J.: In-depth characterization of denitrifier communities across different soil ecosystems in the  
861 tundra, *Environ Microbiome*, 17, 30, <https://doi.org/10.1186/s40793-022-00424-2>, 2022.
- 862 Räsänen, A., Manninen, T., Korhikoski, M., Lohila, A., and Virtanen, T.: Predicting catchment-scale methane  
863 fluxes with multi-source remote sensing, *Landsc. Ecol.*, 36, 1177–1195, [https://doi.org/10.1007/s10980-021-](https://doi.org/10.1007/s10980-021-01194-x)  
864 01194-x, 2021.
- 865 Reynolds, M. K., Walker, D. A., Balser, A., Bay, C., Campbell, M., Cherosov, M. M., Daniëls, F. J. A., Eidesen,  
866 P. B., Ermokhina, K. A., Frost, G. V., Jedrzejek, B., Jorgenson, M. T., Kennedy, B. E., Kholod, S. S.,  
867 Lavrinenko, I. A., Lavrinenko, O. V., Magnússon, B., Matveyeva, N. V., Metúsalemsson, S., Nilsen, L., Olthof,  
868 I., Pospelov, I. N., Pospelova, E. B., Pouliot, D., Razzhivin, V., Schaepman-Strub, G., Šibík, J., Telyatnikov, M.  
869 Y., and Troeva, E.: A raster version of the Circumpolar Arctic Vegetation Map (CAVM), *Remote Sens.*  
870 *Environ.*, 232, 111297, <https://doi.org/10.1016/j.rse.2019.111297>, 2019.
- 871 Repo, M. E., Susiluoto, S., Lind, S. E., Jokinen, S., Elsakov, V., Biasi, C., Virtanen, T., and Martikainen, P. J.:  
872 Large N<sub>2</sub>O emissions from cryoturbated peat soil in tundra, *Nat. Geosci.*, 2, 189–192,  
873 <https://doi.org/10.1038/ngeo434>, 2009.
- 874 Rinne, J., Tuittila, E.-S., Peltola, O., Li, X., Raivonen, M., Alekseychik, P., Haapanala, S., Pihlatie, M., Aurela,  
875 M., Mammarella, I., and Vesala, T.: Temporal Variation of Ecosystem Scale Methane Emission From a Boreal  
876 Fen in Relation to Temperature, Water Table Position, and Carbon Dioxide Fluxes, *Global Biogeochem. Cycles*,  
877 32, 1087–1106, <https://doi.org/10.1029/2017GB005747>, 2018.
- 878 Schlesinger, W. H. and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7–20,  
879 <https://doi.org/10.1023/A:1006247623877>, 2000.
- 880 Shaver, G. R., L. E. Street, Rastetter, E. B., M. T. Van Wijk, and Williams, M.: Functional Convergence in  
881 Regulation of Net CO<sub>2</sub> Flux in Heterogeneous Tundra Landscapes in Alaska and Sweden, *J. Ecol.*, 95, 802–817,  
882 2007.
- 883 Siewert, M. B. and Olofsson, J.: Scale-dependency of Arctic ecosystem properties revealed by UAV, *Environ.*  
884 *Res. Lett.*, 15, 094030, <https://doi.org/10.1088/1748-9326/aba20b>, 2020.
- 885 Sørensen, M. V., Graae, B. J., Classen, A., Enquist, B. J., and Strimbeck, R.: Drivers of C cycling in three  
886 arctic-alpine plant communities, *Arct. Antarct. Alp. Res.*, 51, 128–147,  
887 <https://doi.org/10.1080/15230430.2019.1592649>, 2019.
- 888 St Pierre, K. A., Danielsen, B. K., Hermesdorf, L., D’Imperio, L., Iversen, L. L., and Elberling, B.: Drivers of  
889 net methane uptake across Greenlandic dry heath tundra landscapes, *Soil Biol. Biochem.*, 138, 107605,  
890 <https://doi.org/10.1016/j.soilbio.2019.107605>, 2019.
- 891 Strauss, J., Schirrmeister, L., Grosse, G., Fortier, D., Hugelius, G., Knoblauch, C., Romanovsky, V., Schädel,  
892 C., Schneider von Deimling, T., Schuur, E. A. G., Shmelev, D., Ulrich, M., and Veremeeva, A.: Deep Yedoma  
893 permafrost: A synthesis of depositional characteristics and carbon vulnerability, *Earth-Sci. Rev.*, 172, 75–86,  
894 <https://doi.org/10.1016/j.earscirev.2017.07.007>, 2017.
- 895 Takakai, F., Desyatkin, A. R., Lopez, C. M. L., Fedorov, A. N., Desyatkin, R. V., and Hatano, R.: CH<sub>4</sub> and N<sub>2</sub>O  
896 emissions from a forest-alas ecosystem in the permafrost taiga forest region, eastern Siberia, Russia, *J. Geophys.*  
897 *Res.*, 113, <https://doi.org/10.1029/2007jg000521>, 2008.
- 898 Tramontana, G., Jung, M., Schwalm, C. R., Ichii, K., Camps-Valls, G., Ráduly, B., Reichstein, M., Arain, M.  
899 A., Cescatti, A., Kiely, G., and Others: Predicting carbon dioxide and energy fluxes across global FLUXNET  
900 sites with regression algorithms, *Biogeosciences*, 13, 4291–4313, 2016.

- 901 Treat, C. C., Anthony Bloom, A., and Marushchak, M. E.: Nongrowing season methane emissions—a significant  
902 component of annual emissions across northern ecosystems, <https://doi.org/10.1111/gcb.14137>, 2018a.
- 903 Treat, C. C., Bloom, A. A., and Marushchak, M. E.: Nongrowing season methane emissions—a significant  
904 component of annual emissions across northern ecosystems, *Glob. Chang. Biol.*, 24, 3331–3343, 2018b.
- 905 Treat, C. C., Marushchak, M. E., Voigt, C., Zhang, Y., Tan, Z., Zhuang, Q., Virtanen, T. A., Räsänen, A., Biasi,  
906 C., Hugelius, G., Kaverin, D., Miller, P. A., Stendel, M., Romanovsky, V., Rivkin, F., Martikainen, P. J., and  
907 Shurpali, N. J.: Tundra landscape heterogeneity, not interannual variability, controls the decadal regional carbon  
908 balance in the Western Russian Arctic, *Glob. Chang. Biol.*, 24, 5188–5204, <https://doi.org/10.1111/gcb.14421>,  
909 2018c.
- 910 Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., Minkinen, K., Moore, T.  
911 R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S.,  
912 Waddington, J. M., White, J. R., Wickland, K. P., and Wilmking, M.: A synthesis of methane emissions from 71  
913 northern, temperate, and subtropical wetlands, *Glob. Chang. Biol.*, 20, 2183–2197,  
914 <https://doi.org/10.1111/gcb.12580>, 2014.
- 915 Tyystjärvi, V., Kempainen, J., Luoto, M., Aalto, T., Markkanen, T., Launiainen, S., Kieloaho, A.-J., and Aalto,  
916 J.: Modelling spatio-temporal soil moisture dynamics in mountain tundra, *Hydrol. Process.*, 36,  
917 <https://doi.org/10.1002/hyp.14450>, 2022.
- 918 Vainio, E., Peltola, O., Kasurinen, V., Kieloaho, A.-J., Tuittila, E.-S., and Pihlatie, M.: Topography-based  
919 statistical modelling reveals high spatial variability and seasonal emission patches in forest floor methane flux,  
920 *Biogeosciences*, 18, 2003–2025, <https://doi.org/10.5194/bg-18-2003-2021>, 2021.
- 921 Virkkala, A.-M., Virtanen, T., Lehtonen, A., Rinne, J., and Luoto, M.: The current state of CO<sub>2</sub> flux chamber  
922 studies in the Arctic tundra: A review, *Progress in Physical Geography: Earth and Environment*, 42, 162–184,  
923 <https://doi.org/10.1177/0309133317745784>, 2018.
- 924 Virkkala, A.-M., Aalto, J., Rogers, B. M., Tagesson, T., Treat, C. C., Natali, S. M., Watts, J. D., Potter, S.,  
925 Lehtonen, A., Mauritz, M., Schuur, E. A. G., Kochendorfer, J., Zona, D., Oechel, W., Kobayashi, H.,  
926 Humphreys, E., Goeckede, M., Iwata, H., Lafleur, P. M., Euskirchen, E. S., Bokhorst, S., Marushchak, M.,  
927 Martikainen, P. J., Elberling, B., Voigt, C., Biasi, C., Sonnentag, O., Parmentier, F.-J. W., Ueyama, M., Celis,  
928 G., St Loius, V. L., Emmerton, C. A., Peichl, M., Chi, J., Järveoja, J., Nilsson, M. B., Oberbauer, S. F., Torn, M.  
929 S., Park, S.-J., Dolman, H., Mammarella, I., Chae, N., Poyatos, R., López-Blanco, E., Røjle Christensen, T.,  
930 Jung Kwon, M., Sachs, T., Holl, D., and Luoto, M.: Statistical upscaling of ecosystem CO<sub>2</sub> fluxes across the  
931 terrestrial tundra and boreal domain: regional patterns and uncertainties, *Glob. Chang. Biol.*,  
932 <https://doi.org/10.1111/gcb.15659>, 2021.
- 933 Voigt, C., Marushchak, M. E., Lamprecht, R. E., Jackowicz-Korczyński, M., Lindgren, A., Mastepanov, M.,  
934 Granlund, L., Christensen, T. R., Tahvanainen, T., Martikainen, P. J., and Biasi, C.: Increased nitrous oxide  
935 emissions from Arctic peatlands after permafrost thaw, *Proc. Natl. Acad. Sci. U. S. A.*, 114, 6238–6243,  
936 <https://doi.org/10.1073/pnas.1702902114>, 2017a.
- 937 Voigt, C., Lamprecht, R. E., Marushchak, M. E., Lind, S. E., Novakovskiy, A., Aurela, M., Martikainen, P. J.,  
938 and Biasi, C.: Warming of subarctic tundra increases emissions of all three important greenhouse gases—carbon  
939 dioxide, methane, and nitrous oxide, *Glob. Chang. Biol.*, 23, 3121–3138, 2017b.
- 940 Voigt, C., Marushchak, M. E., Abbott, B. W., Biasi, C., Elberling, B., Siciliano, S. D., Sonnentag, O., Stewart,  
941 K. J., Yang, Y., and Martikainen, P. J.: Nitrous oxide emissions from permafrost-affected soils, *Nature Reviews*  
942 *Earth & Environment*, 1, 420–434, <https://doi.org/10.1038/s43017-020-0063-9>, 2020.
- 943 Voigt, C., Virkkala, AM., Hould Gosselin, G. et al. Arctic soil methane sink increases with drier conditions and  
944 higher ecosystem respiration. *Nature Climate Change*. <https://doi.org/10.1038/s41558-023-01785-3>, 2023.
- 945 Vowles, T. and Björk, R. G.: Implications of evergreen shrub expansion in the Arctic, *J. Ecol.*, 107, 650–655,  
946 <https://doi.org/10.1111/1365-2745.13081>, 2018.
- 947 Wagner, I., Hung, J. K. Y., Neil, A., and Scott, N. A.: Net greenhouse gas fluxes from three High Arctic plant  
948 communities along a moisture gradient, *Arct. sci.*, 5, 185–201, <https://doi.org/10.1139/as-2018-0018>, 2019.

949 Wild, J., Kopecký, M., Macek, M., Šanda, M., Jankovec, J., and Haase, T.: Climate at ecologically relevant  
950 scales: A new temperature and soil moisture logger for long-term microclimate measurement, *Agric. For.*  
951 *Meteorol.*, 268, 40–47, <https://doi.org/10.1016/j.agrformet.2018.12.018>, 2019.

952 Williams, M., Street, L. E., van Wijk, M. T., and Shaver, G. R.: Identifying Differences in Carbon Exchange  
953 among Arctic Ecosystem Types, *Ecosystems*, 9, 288–304, <https://doi.org/10.1007/s10021-005-0146-y>, 2006.

954 Zona, D., Lafleur, P. M., Hufkens, K., Bailey, B., Gioli, B., Burba, G., Goodrich, J. P., Liljedahl, A. K.,  
955 Euskirchen, E. S., Watts, J. D., Farina, M., Kimball, J. S., Heimann, M., Göckede, M., Pallandt, M.,  
956 Christensen, T. R., Mastepanov, M., López-Blanco, E., Jackowicz-Korczynski, M., Dolman, A. J., Marchesini,  
957 L. B., Commane, R., Wofsy, S. C., Miller, C. E., Lipson, D. A., Hashemi, J., Arndt, K. A., Kutzbach, L., Holl,  
958 D., Boike, J., Wille, C., Sachs, T., Kalhori, A., Song, X., Xu, X., Humphreys, E. R., Koven, C. D., Sonntag,  
959 O., Meyer, G., Gosselin, G. H., Marsh, P., and Oechel, W. C.: Earlier snowmelt may lead to late season declines  
960 in plant productivity and carbon sequestration in Arctic tundra ecosystems, *Sci. Rep.*, 12, 3986,  
961 <https://doi.org/10.1038/s41598-022-07561-1>, 2022.

962 National Land Survey of Finland. 2019. Digital elevation model. National Land Survey of Finland.  
963 [https://www.maanmittauslaitos.fi/en/maps-and-spatial-data/expert-users/product-descriptions/elevation-model-](https://www.maanmittauslaitos.fi/en/maps-and-spatial-data/expert-users/product-descriptions/elevation-model-2-m)  
964 [2-m](https://www.maanmittauslaitos.fi/en/maps-and-spatial-data/expert-users/product-descriptions/elevation-model-2-m). Last accessed 19/04/2019

965  
966 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical  
967 Computing, Vienna, Austria. <https://www.R-project.org/>.

968 Virkkala, A.-M., Niittynen, P., Kemppinen, J., Marushchak, M.E, Voigt, C., Hensgens, G., Kerttula, J., Happonen,  
969 K., Tyystjärvi, V., Biasi, C., Hultman, J., Rinne, J., & Luoto, M. Data and code for "High-resolution spatial  
970 patterns and drivers of terrestrial ecosystem carbon dioxide, methane, and nitrous oxide fluxes in the tundra" [Data  
971 set]. Zenodo. <https://doi.org/10.5281/zenodo.7760577>, 2023.

972  
973  
974  
975  
976  
977  
978  
979  
980