



Bacterioplankton dark CO₂ fixation in oligotrophic waters

Afrah Allothman^{1*}, Daffne López-Sandoval^{1,2}, Carlos M. Duarte¹, Susana Agustí¹

¹Red Sea Research Centre, Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal, 23955, Saudi Arabia

5 ²Coastal and Marine Resources Core Lab (CMR), King Abdullah University of Science and Technology (KAUST), Thuwal, 23955, Saudi Arabia

*Corresponding author**: afrah.alothman@kaust.edu.sa

Abstract

10 Dark CO₂ fixation by bacteria is believed to be particularly important in oligotrophic ecosystems. However, only a few studies have characterized the role of bacterial dissolved inorganic carbon (DIC) fixation in global carbon dynamics. Therefore, this study quantified the primary production (PP), total bacteria dark CO₂ fixation (TB_{DIC} fixation), and heterotrophic bacterial production (HBP) in the warm and oligotrophic Red Sea using stable isotope labeling and cavity ring-down spectroscopy (¹³C-CRDS). Additionally, we assessed the contribution of bacterial
15 DIC fixation (TB_{DIC} %) relative to the total DIC fixation (Total_{DIC} fixation). Our study demonstrated that TB_{DIC} fixation increased the Total_{DIC} fixation from 2.03 to 60.45 μg C L⁻¹ d⁻¹ within the photic zone, contributing 13.18 % to 71.68 % with an average value of 33.95 ± 0.02 % of the photic layer Total_{DIC} fixation. The highest TB_{DIC} fixation values were measured at the surface and deep (400 m) water with an average value of 5.23 ± 0.45 μg C L⁻¹ d⁻¹, and 4.95 ± 1.33 μg C L⁻¹ d⁻¹, respectively. These findings suggest that the non-photosynthetic processes such as
20 anaplerotic DIC reactions and chemo-autotrophic CO₂ fixation extended to the entire oxygenated water column. On the other hand, the % of TB_{DIC} contribution to Total_{DIC} fixation increased as primary production decreased (R² = 0.45, p < 0.0001), suggesting the relevance of increased dark DIC fixation when photosynthetic production was low or absent, as observed in other systems. Therefore, when estimating the total carbon dioxide production in the ocean, dark DIC fixation must also be accounted as a crucial component of the carbon dioxide flux in addition to
25 photosynthesis.

1 Introduction

Bacteria are the central nodes of the microbial loop and play an essential role in the flux of organic carbon in marine ecosystems through different metabolic pathways (Azam et al., 1983). Most studies on the metabolism of marine
30 bacteria have focused on quantifying the uptake of organic compounds by heterotrophic bacteria and how it relates to bacterial growth and reproduction (Ducklow and Kirchman, 2000; Kirchman, 2000). However, heterotrophic marine bacteria can also metabolize CO₂ through anaplerotic carboxylation reactions, which form the basis of several metabolic pathways (Dijkhuizen and Harder, 1984). Such reactions are essential components of metabolic
35 pathways in bacteria that enable the synthesis of fatty acids, amino acids, vitamins, and nucleotides (Dijkhuizen and Harder, 1984; Erb, 2011), which also fuel microbial food webs.

Wood and Werkman (1936 first proposed that heterotrophic bacteria contribute to dark dissolved inorganic carbon (DIC) fixation, a discovery that was widely embraced by the scientific community. A decade later, when radioactive



40 isotope techniques emerged in the field, Steemann-Nielsen (Steemann-Nielsen, 1952) first reported on the possible
importance of dark DIC fixation to the total carbon flux in the ocean, suggesting that it could represent between 1 %
and 30 % of photosynthetic CO₂ fixation (Steemann-Nielsen, 1952; Nielsen, 1960). Subsequent quantifications in
the northern to southern Pacific and Atlantic oceans reported that dark DIC fixation accounted for approximately
>10 % of photosynthetic CO₂ fixation in temperate and equatorial areas, and between 10 % to 50 % of the light
45 DIC fixation in marine surface water, which has been directly associated with high bacterial activity (Prakash et al.,
1991, Li et al., 1993, Markager, 1998, Li and Dickie, 1991). A study that evaluated the role of Arctic bacterial dark
CO₂ incorporation suggested that the depletion and limitation of labile organic carbon compounds could enhance the
utilization of bicarbonate by chemoautotroph or heterotroph microorganisms to achieve metabolic balance (Alonso-
Sáez et al., 2010). Additionally, it was assumed that dark CO₂ utilization was mainly attributable to bacterial
50 metabolism, as it was highly positively correlated with heterotrophic bacterial production (Alonso-Sáez et al.,
2010). In general, anaplerotic CO₂ fixed by bacteria has been estimated to contribute up to 8 % of heterotroph
carbon biomass production in the ocean (Romanenko, 1964) and contributes significantly to the carbon flux
dynamics of many marine ecosystems (Alonso-Sáez et al., 2010, Yakimov et al., 2014, Zhou et al., 2017., Signori et
al., 2017, and Lliro's et al., 2011).

55

Dark CO₂ fixation by marine bacteria is thought to play an essential role under oligotrophic conditions, contributing
up to 30 % of bacterial production (González et al., 2008; Palovaara et al., 2014). Genetic studies have demonstrated
that increases in the abundance of associated anaplerotic enzyme transcripts coincided with a sudden increase in
bacterial metabolism, which could contribute significantly to the total DIC fixation rates in oligotrophic
60 environments (Baltar et al., 2016). Additionally, the recent discovery of light-driven CO₂ incorporation by
proteorhodopsin-containing flavobacterium *Polaribacter* sp. highlighted the significant role of anaplerotic
metabolism in heterotrophs (González et al., 2008). Therefore, whereas the total primary production of oceanic
ecosystems is typically attributed to photosynthesis, dark chemo-autotrophic and anaplerotic metabolism are also
important contributors (Baltar and Herndl, 2019).

65

Dark CO₂ fixation by bacteria is, therefore, likely to be highly relevant in the Red Sea, a predominantly oligotrophic,
landlocked system with no river inflow and with limited connection to the Indian Ocean (Edwards, 1987; Grasshoff,
1969). These features result in a limited nutrient input and an oligotrophication gradient from south to north (Wafar
et al., 2016). The Red Sea is also characterized by high surface temperatures ranging from 20 to 33.1 °C (Chaidez et
70 al., 2017, Shaltout, 2019) and warm deep-water temperatures of up to ~21.5 °C at depths below 300 m (Yao &
Hoteit, 2018).

Here we assess the contribution of dark CO₂ assimilation to the Red Sea bacterioplankton production using ¹³C
stable isotope as a tracer. In this study, both the dark bicarbonate synthesis by bacteria (TB_{DIC}) and light bicarbonate
synthesis by photosynthetic phytoplankton (PP) were quantified using ¹³C as a tracer in the Red Sea water column.

75 Additionally, ¹³C stable isotope fluxes were used to estimate bacterial production (BP) in the dark. Our study also



assessed the variations in dark CO₂ fixation rates at different depths through the water column in both open and coastal water bodies, and the relationship with water temperature. Moreover, our study estimated the contribution of dark bicarbonate synthesis (TB_{DIC} %) to total CO₂ fixation (Total_{DIC} fixation) by accounting for dark and light CO₂ fixation rates.

80

2 Methods

2.1 Sample collection and processing

85 A total of 59 water samples were collected over the course of four oceanographic cruises in the Eastern Red Sea, including the Center Competitive Found (CCF), Deep Cruise (DC), Deep Coral Survey (DCS), and Red Sea Decade Expedition (RSDE). Samples were also obtained from a time series fixed station (pelagic) and two other coastal sites (a lagoon and reef located in Abu Shusha) in the Central Red Sea of Saudi Arabia (Fig. 1). The cruises were conducted on board R/V *Thuwal* (DC, DCS), R/V *Al Azizi* (CCF), and *OceanXplorer* (RSDE) between August 2017
90 and June 2022. During the CCF oceanographic cruise, water samples were collected at five different depths ranging from the surface to the bottom layer of the photic zone. During the DC and DCS oceanographic cruises, time series and coastal station water samples we collected from surface water at a 3–5 m depth. During the RSDE cruise, water samples were collected from the surface at a 5 m depth, as well as from 400 m. Vertical environmental profiles of temperature, salinity, and photosynthetic active radiation were obtained for all of the studied stations as described by
95 López-Sandoval et al. (2021).






Figure 1: Stations sampled during the four oceanographic cruises [CCF (blue dots), DC (green dots), DCS (red dots), RSDE (black dots)] time series and coastal stations (orange dots) along the Eastern Red Sea conducted between 2017 and 2022.

At each station, water samples for isotope labeling analysis were collected in the morning using 12 L Niskin bottles with a rosette system or 10 L Niskin bottles deployed manually (López-Sandoval et al., 2021). For the deepest water (400 m), the water samples were collected using 1.5 L Niskin bottles attached to a remotely-operated underwater vehicle (ROV) and a submarine on board *OceanXplorer*. Water samples were collected directly from the Niskin bottles, prefiltered through 100 μ m mesh filters to remove larger zooplankton, and transferred into 10 L acid-washed carboy containers. The water samples were distributed into three transparent 2 L or 500 mL (^{13}C -PP; López-Sandoval, et al. 2019) polycarbonate (PC) bottles for light primary production measurements and another three dark 2 L or 500 mL PC bottles for measuring dark bacterial DIC uptake rate (^{13}C -TB_{DIC}). All water samples were enriched with ^{13}C -sodium bicarbonate solution (99.8 atom % 4 g/L of $\text{NaH}^{13}\text{CO}_3$; López-Sandoval et al., 2019) to a final carbon concentration of $\sim 153 \mu\text{mole } ^{13}\text{C L}^{-1}$ in each bottle. Additionally, during the DC and DCS oceanographic cruises, three dark PC bottles were enriched with ^{13}C -D-glucose substrate at a final concentration of 100 nM to measure bacterial production (^{13}C -BP) as described by Koshikawa et al. (1999). All PC bottles were incubated in tanks placed on the vessel's deck with a circulating seawater system to maintain surface water temperature and receive natural solar radiation. Moreover, a separate tank was attached to a chiller to mimic the water temperature at a 400 m depth. Additionally, coastal water samples were incubated in a similar outdoor setup at the Coastal & Marine Resources Core Lab (CMR) at King Abdullah University of Science and Technology (KAUST).

The bottles for PP were covered with neutral-density nets to reduce the light intensity according to the matching light received at the assigned depth. After 4–6 hours of incubation, the water samples collected during the CCF cruise and the time series station were filtered through pre-combusted Whatman GF/F filters (López-Sandoval et al., 2021). The rest of the samples were filtered through 25 mm diameter 3 μm Silver membranes (STERLITECH) to collect the size fraction above the picoplankton size. Afterward, the filtrate was collected into a 25 mm diameter (0.2 μm) Silver membrane filter (STERLIECH). The samples collected during the RSDE cruise were pre-filtered through 3 μm polycarbonate membrane filters before incubation. The natural isotopic composition of particulate organic carbon was measured in surface and deep seawater at each station. The collected filters were placed in small Petri dishes containing 150 μl (50 %) HCl to remove carbonate from the filters, allowed to dry for 12 hours, and stored at $-20 \text{ }^\circ\text{C}$ until required for downstream analyses.

130

2.1.1 Chlorophyll-*a* concentration and nutrients

Samples for chlorophyll-*a* (Chl-*a*) and nutrient analyses were collected at each depth within the photic layer during the CCF cruise and from surface water during the time series study (López-Sandoval et al., 2021). Samples for Chl-*a*



135 analysis were collected from all cruises and coastal stations. The water samples (200–500 ml) were filtered through
25 mm Whatman GF/F filters (0.7 μm) and then extracted in 90 % acetone in the dark as described by Prabowo and
Agusti (2019) and López-Sandoval et al. (2021). After 24 hours, the extracted pigment was measured using a Trilogy
Fluorometer equipped with a CHL-NA module (Turner Designs; San Jose, USA) calibrated with pure Chl-a (Prabowo
and Agusti, 2019). Water samples for inorganic nutrient concentration were collected and frozen until analyzed in the
140 laboratory. Nutrient concentrations were determined with a Segmented Flow Analyzer (SEAL AA3 Analytical Inc.;
WI, USA) following standard autoanalyzer methods (Hansen and Koroleff, 1999).

2.1.2 Heterotrophic bacteria abundance

The abundance of heterotrophic bacteria was quantified in each water sample. Briefly, 1.8 mL aliquots were
145 obtained from each sample, fixed with 25 % glutaraldehyde, flash-frozen in liquid nitrogen, and stored at $-80\text{ }^{\circ}\text{C}$ for
later analysis. The samples were stained with SYBR Green I intermediate solution (1:100) for the determination of
bacteria cell abundance by flow cytometry (Gasol and Moran, 2016) using either a FACSCanto II (Becton
Dickinson) or a Flow Cytometer Cube 8 (Sysmex).

2.1.3 Dissolved inorganic carbon (DIC)

150 The $\delta^{13}\text{C}$ of DIC in natural waters and after enrichment with $\text{NaH}^{13}\text{CO}_3^-$ was analyzed in seawater samples placed in
15 mL small glass tubes and treated with 0.05 % mercuric chloride (HgCl_2 -Sigma-Aldrich) to stop any biological
activity after sampling (Dickson et al., 2007). After fixation, all samples were kept in a dark and cool place until
analyzed in the laboratory. DIC measurements were conducted using an AutoMate Prep Device coupled with
Picarro's LIAISON interface and IsoCO2 WS-CRDS system (Santa Clara, California, USA).

155

2.1.4 Primary production (PP) and total dark bacteria DIC fixation (TB_{DIC})

The carbon content and the $\delta^{13}\text{C}$ values from PP and TB_{DIC} incubation filters were analyzed using a combustion
module (CM) attached to a cavity-ring down spectroscopy analyzer (CM-CRDS-G2201-I, Picarro), and each filter
160 was analyzed for 600 s. The combustion module converted the sample into the required gas (CO_2) by fast
combustion, after which the gas was transferred to the isotopic analyzer to measure the $^{13}\text{C}/^{12}\text{C}$ ratio ($\delta^{13}\text{C}$). Inside
the cavity, spectrum peaks were generated according to the measured wavelength absorbed by the gas of interest
($^{13}\text{CO}_2$ and $^{12}\text{CO}_2$), where each peak corresponded to the ^{13}C and ^{12}C concentrations (López-Sandoval et al., 2019).

Before analyzing the filters, CRDS-Picarro was calibrated using VPDB standards from the International Atomic
165 Energy Agency (IAEA) including IAEA-CH-6, C3, and 303B with $\delta^{13}\text{C}$ values of $-10.45\text{ }‰$, $-24.72\text{ }‰$, and $+450\text{ }‰$,
respectively. Additionally, Reston Stable Isotopic Laboratory standards (United States Geological Survey, UGS)
were also used, including USG62 ($-14.79\text{ }‰$), USG40 ($-26, 39\text{ }‰$), and USG41a ($+36.55\text{ }‰$, López-Sandoval et al.,
2019).



170 The ^{13}C and ^{12}C isotopic mass from the ^{13}C -enriched sample at the end of incubation time (h) was determined to
calculate PP ($\mu\text{g C}^{-1} \text{h}^{-1}$) using the ^{13}C -CRDS-PP method (López-Sandoval et al., 2019). Particularly, ^{13}C -PP was
calculated as the isotopic shift of particulate organic carbon (POC) from the samples incubated in the light ($\delta^{13}\text{C}_{\text{POC-Light}}$)
relative to the dark isotopic composition of the samples ($\delta^{13}\text{C}_{\text{POC-Dark}}$). Additionally, the isotopic shift of the
enriched DIC ($\delta^{13}\text{C}_{\text{DIC-Enriched}}$) relative to the natural DIC samples ($\delta^{13}\text{C}_{\text{DIC-Natural}}$) was also calculated. The
175 production was converted to carbon uptake rates considering the particulate organic carbon measured at the end of
incubation in the light-enriched samples per volume filtered in L ($\text{POC-}\mu\text{g C L}^{-1}$), and the carbon fixation rate was
calculated per time unit (hours of incubation) following Eq (1) below:

$$^{13}\text{C-PP} = ((\delta^{13}\text{C}_{\text{POC-Light}} - \delta^{13}\text{C}_{\text{POC-Dark}}) / (\delta^{13}\text{C}_{\text{DIC-Enriched}} - \delta^{13}\text{C}_{\text{DIC-Natural}})) \times \text{POC} / v/t$$

180

The TB_{DIC} fixation rate ($\mu\text{g C}^{-1} \text{h}^{-1}$) was measured using a similar equation as for the ^{13}C -PP method. However, the
uptake rate was calculated as the isotopic shift of particulate organic carbon from the samples incubated in the dark
($\delta^{13}\text{C}_{\text{POC-Dark}}$) relative to the natural isotopic composition of the samples ($\delta^{13}\text{C}_{\text{POC-Natural}}$). Additionally, we also
calculated the isotopic shift of the enriched DIC ($\delta^{13}\text{C}_{\text{DIC-Enriched}}$) relative to the natural DIC samples ($\delta^{13}\text{C}_{\text{DIC-Natural}}$).
185 The production rate was also converted to carbon uptake rates considering the particulate organic carbon measured
at the end of incubation in the dark-enriched samples and the volume filtered in L ($\text{POC-}\mu\text{g C L}^{-1}$). Additionally, the
carbon fixation rate was calculated per time unit (hours of incubation) following Eq (2) below:

$$^{13}\text{C-TB}_{\text{DIC}} = ((\delta^{13}\text{C}_{\text{POC-Dark}} - \delta^{13}\text{C}_{\text{POC-Natural}}) / (\delta^{13}\text{C}_{\text{DIC-Enriched}} - \delta^{13}\text{C}_{\text{DIC-Natural}})) \times \text{POC} / v/t$$

190

The $\text{Total}_{\text{DIC}}$ fixation was calculated as the sum of the ^{13}C -PP measured in the light and ^{13}C - TB_{DIC} measured in the
dark following Eq (3) below:

$$\text{Total}_{\text{DIC}} \text{ fixation} = ^{13}\text{C-PP}_{\text{Light}} + ^{13}\text{C-TB}_{\text{DIC-Dark}}$$

195

The contribution of the TB_{DIC} to the total carbon production was calculated relative to the $\text{Total}_{\text{DIC}}$ fixation following
Eq (4) below:

$$\text{TB}_{\text{DIC}} \% = ^{13}\text{C-TB}_{\text{DIC-Dark}} / \text{Total}_{\text{DIC}} \text{ fixation}$$

200

The data for PP and TB_{DIC} were converted to carbon uptake per day using a local photoperiod of 12 hours of
daytime for photosynthesis and 12 hours of nighttime for dark DIC fixation.

2.1.5 Bacterial production (BP)

205



Bacterial production (BP) was measured based on glucose uptake and was expressed in $\mu\text{g C L}^{-1} \text{h}^{-1}$ as described by Middelburg et al. (2000). ^{13}C incorporation was calculated as glucose uptake and defined as the difference between the fraction of ^{13}C in the natural isotopic composition sample (F_{Natural}) relative to the ^{13}C fraction in the enriched sample (F_{Enriched}) following Eq (5) below:

$$210 \quad \text{Glucose uptake rate (BP-}\mu\text{g C L}^{-1} \text{h}^{-1}) = [(F_{\text{Enriched}} - F_{\text{Natural}}) \times \text{POC}]/v/t$$

where $F = {}^{13}\text{C} / ({}^{13}\text{C} + {}^{12}\text{C})$; which is also expressed as $R/(R+1)$, where R is the carbon isotope ratio obtained from the measured $\delta {}^{13}\text{C}$ values and it is calculated following Eq (6, Middelburg et al., 2000) below:

$$R = (\delta {}^{13}\text{C}/1000 + 1) \times \text{V-PDB (Vienna Pee Dee Belemnite),}$$

215 where V-PDB = 0.0112372. Additionally, the uptake rate was calculated by considering the particulate organic carbon measured in the samples at the end of incubation per time unit (hours of incubation) and per volume filtered in L. The BP rates were reported on a per-day basis considering a 24 hours cycle.

2.1.6 Statistical analysis

220 All statistical analyses were conducted using the JMP PRO 16 software (JMP®, Version <16.1> SAS Institute Inc., Cary, NC, 1989–2019). A p -value ≤ 0.05 was considered statistically significant. The relationship and correlation between variables were explored using Spearman's nonparametric correlations and linear regression, and means were compared using one-way ANOVA and Student's t-test.

3 Results

225

The seawater temperature during the study period ranged from a minimum of 21.0 °C in deep waters (400 m depth) to a maximum of 32.2°C in surface waters, showing no latitudinal changes. Salinity, on the other hand, ranged from 38.21 to 40.80 (Table 1) and increased significantly with latitude ($\rho = 0.71$, $p < 0.0001$). Our results indicated that 230 silicate (SiO_2) levels were generally low, ranging from 0.44 to 1.65 μM , whereas phosphate (PO_4) ranged from undetectable to 0.31 μM . Both SiO_2 and PO_4 decreased with increasing latitude ($\rho = -0.42$, $p < 0.0007$, and $\rho = -0.51$, $p < 0.0001$, respectively). Nitrate (NO_3) (Table 1), however, did not exhibit any significant spatial variability throughout the study, with an average ($\pm\text{SE}$) low value of $0.57 \pm 0.05 \mu\text{M}$ (Mean \pm SE) from the surface to a 30 m depth, and a higher mean value of $2.44 \pm 0.55 \mu\text{M}$ below 30 m depth. Surface ($3.66 \pm 0.16 \text{ m}$) Chl-a averaged $0.22 \pm$ 235 $0.5 \text{ Chl-a } \mu\text{g L}^{-1}$ in the south, $0.38 \pm 0.05 \mu\text{g L}^{-1}$ in the central coastal stations, and $0.34 \pm 0.90 \mu\text{g L}^{-1}$ in the northern stations, and ranged from 0.04 to 0.81 $\mu\text{g L}^{-1}$, with the coastal stations, which exhibited the highest individual values (Table 1). However, during the winter in the northern stations, Chl-a concentration peaked and reached up to 0.69 $\mu\text{g L}^{-1}$, probably reflecting nutrient entrainment due to convective mixing. In addition, Chl-a concentration measured



240 during CCF cruise through different photic zone layers showed an increase toward the maximum chlorophyll a depth, located at, on average 61.66 m depth, where it reached up to $0.48 \pm 0.036 \mu\text{g L}^{-1}$.

Table 1: Value ranges (minimum-maximum) of the environmental and biological parameters obtained during the study along the Red Sea at different depths. The data in the table are Chlorophyll-a concentration (Chl-a), seawater Temperature (Temp), Salinity (Sal), Nutrient concentrations (SiO_2 , PO_4 , and NO_3), heterotrophic bacteria abundance (BACT), and Bacterial Production (BP). N/A: data not available.

Cruise	Date	Lat °N	Long °E	Depth m	Temp °C	Sal	Chl-a ($\mu\text{g L}^{-1}$)	SiO_2 μM	PO_4 μM	NO_3 μM	BACT cells mL^{-1}	BP $\mu\text{g C L}^{-1} \text{d}^{-1}$
CCF (Open water)	16/03/18–21/03/18	17.35–22.23	38.38–40.42	5–90	24.20–27.69	38.21–38.90	0.08–0.69	0.57–1.65	0.04–0.28	0.01–5.31	N/A	N/A
Deep Cruise (Open water)	04/04/19–09/04/19	18.67–24.46	37.01–40.22	5	24.00–27.00	38.30–40.00	0.14–0.19	0.59–1.03	0.06–0.20	0.14–0.60	$3.25\text{E}+05$ – $7.84\text{E}+05$	0.12–0.46
Deep Coral Survey (open water)	18/01/20–23/01/20	22.30–25.75	36.34–38.86	5	23.11–24.90	39.62–40.23	0.37–0.62	0.83–1.03	0.01–0.11	0.27–1.04	$4.38\text{E}+05$ – $6.57\text{E}+05$	0.04–0.08
Time series (coastal)	21/08/17–05/02/18	22.31	38.96	3	24.40–32.10	39.30–39.57	0.18–0.81	0.44–1.02	0.01–0.31	0.44–1.12	$2.00\text{E}+05$ – $4.13\text{E}+05$	
Reef (coastal)	12/11/19	22.32	39.02	3	29.90	39.78	0.40	N/A	N/A	N/A	N/A	0.69
Lagoon (coastal)	22/10/19	22.39	39.14	3	32.20	40.80	0.70	N/A	N/A	N/A	N/A	0.92
RSDE (open water)	07/02/22–05/06/22	19.44–24.15	36.59–39.44	5	25.13–28.87	38.45–40.16	0.07–0.49	N/A	N/A	N/A	$5.87\text{E}+04$ – $1.86\text{E}+05$	
				400	21.00–21.72	39.29–40.54	N/A	N/A	N/A	$1.90\text{E}+04$ – $3.48\text{E}+04$		

250 Heterotrophic bacteria cell abundance ranged from 1.90×10^4 cells mL^{-1} at 400 m depth to 7.84×10^5 cells mL^{-1} in surface waters, averaging $3.78 \pm 0.38 \times 10^5$ cells mL^{-1} in the photic layer (Table 1). There was no significant difference in bacterial abundance between open and coastal water (F ratio = 0.19, df = 1, p = 0.66). The bacterial production measured as the glucose uptake rate in the dark (HBP) varied from $0.04 \mu\text{g C L}^{-1} \text{d}^{-1}$ recorded in the northern stations to 0.69 and $0.92 \mu\text{g C L}^{-1} \text{d}^{-1}$ (Table 1) in the coastal reef and lagoon stations, respectively.

255 Additionally, HBP was significantly different between coastal and open waters (F ratio = 11.07, df = 1, p < 0.005), with higher rates observed in the coastal stations ($0.60 \pm 0.21 \mu\text{g C L}^{-1} \text{d}^{-1}$) compared to the open waters ($0.17 \pm 0.04 \mu\text{g C L}^{-1} \text{d}^{-1}$). HBP increased with increasing temperature ($R^2 = 0.71$, p < 0.0001).

260 Primary production rates across the study ranged from $0.84 \mu\text{g C L}^{-1} \text{d}^{-1}$ to $47.76 \mu\text{g C L}^{-1} \text{d}^{-1}$ (Table 2), and were significantly higher at the coastal stations ($17.77 \pm 3.60 \mu\text{g C L}^{-1} \text{d}^{-1}$) compared to open waters ($7.24 \pm 0.71 \mu\text{g C L}^{-1} \text{d}^{-1}$) (F ratio = 20.17, df = 1, p < 0.0001, Fig. 2A). PP also tended to increase with increasing temperature ($\rho = 0.35$, p < 0.001), but was independent of Chl-a concentration ($\rho = 0.12$, p = 0.52), and declined with increasing nitrate concentration ($\rho = -0.46$, p < 0.014, Fig. 4). Primary production throughout the photic zone (CCF open water)




265 decreased from $15 \mu\text{g C L}^{-1} \text{d}^{-1}$ at the surface to $<1 \mu\text{g C L}^{-1} \text{h}^{-1}$ at the base the photic layer (1 % PAR). In contrast, the concentration of particulate organic carbon across the photic layer changed gradually, from $95.13 \pm 10.23 \mu\text{g C L}^{-1}$ at the surface to $18.25 \pm 1.28 \mu\text{g C L}^{-1}$ at the base of the photic zone.

270 **Table 2:** Range (minimum-maximum) and mean \pm SE of dark DIC uptake rates by bacteria (TB_{DIC}), primary production (PP), the percentage contribution of TB_{DIC} ($\text{TB}_{\text{DIC}} \%$) relative to the total DIC fixation (Total $_{\text{DIC}}$ fixation), and PP obtained at different water depths in different cruises.

Cruise	Lat (°N)	Long (°E)	Depth (m)	TB_{DIC} ($\mu\text{g C L}^{-1} \text{d}^{-1}$)	PP ($\mu\text{g C L}^{-1} \text{d}^{-1}$)	PP %	$\text{TB}_{\text{DIC}} \%$	Mean Depth (m)	Mean PP ($\mu\text{g C L}^{-1} \text{d}^{-1}$)	Mean TB_{DIC} ($\mu\text{g C L}^{-1} \text{d}^{-1}$)	Mean $\text{TB}_{\text{DIC}} \%$	Mean PP %
CCF (open water)	17.35–22.23	38.38–40.42	5–90	1.07–2.69	9.84–15.36	28.22%–86.82%	13.18%–71.78%	3.66	11.15 ± 1.67	5.23 ± 0.54	36 ± 0.02	63 ± 0.02
Deep Cruise (open water)	18.67–24.46	37.01–40.22	5	2.05–12.45	2.69–16.71	45.48%–70.87%	29.13%–54.52%	12.00	12.71 ± 0.79	2.33 ± 0.11	15 ± 0.01	84 ± 0.01
Deep Coral Survey (open water)	22.30–25.75	36.34–38.86	5	2.69–10.61	3.16–14.68	34.49%–82.68%	17.32%–65.51%	28.00	10.33 ± 2.04	2.04 ± 0.25	17 ± 0.01	82 ± 0.01
Time series (coastal)	22.31	38.96	3	3.44–12.69	3.46–47.76	34.49%–84.37%	15.63%–65.51%	45.60	5.91 ± 0.68	1.89 ± 0.35	23 ± 0.02	76 ± 0.02
Reef (coastal)	22.32	39.02	3	3.44	4.31	55.61%	44.39%	61.66	2.12 ± 0.69	1.66 ± 0.50	45 ± 0.04	54 ± 0.04
Lagoon (coastal)	22.39	39.14	3	10.11	15.88	61.10%	38.90%	84.00	1.35 ± 0.45	1.42 ± 0.35	52 ± 0.01	47 ± 0.11
RSDE (open water)	19.44–24.15	36.59–39.44	5	1.89–5.65	2.15–6.21	48.13%–75.55%	24.45%–55.43%	400.00	N/A	4.95 ± 1.19	40 ± 0.07	59 ± 0.07
			400	1.37–6.33	N/A	N/A	N/A					

275




280 **Figure 2:** (A) Primary Production (PP) and (B) Total dark bacteria DIC fixation (TB_{DIC}) measured in open and coastal waters. Boxplots indicate the 95 % confidence intervals with ± 1 SD deviation. The central line in the box represents the median for each group of samples.



285 The ^{13}C -dark TB_{DIC} fixation rate varied from $1.07 \mu\text{g C L}^{-1} \text{d}^{-1}$ to a maximum of $12.69 \mu\text{g C L}^{-1} \text{d}^{-1}$ (Table 2) with no latitudinal pattern. We found significant differences in TB_{DIC} between coastal ($6.92 \pm 0.81 \mu\text{g C L}^{-1} \text{d}^{-1}$) and open water stations ($3.34 \pm 0.37 \mu\text{g C L}^{-1} \text{d}^{-1}$); (F ratio = 19.89, df = 1, $p < 0.0001$, Fig. 2B), but not ($p = 0.84$) between surface ($5.24 \pm 0.54 \mu\text{g C L}^{-1} \text{d}^{-1}$) and 400 m samples ($4.95 \pm 1.20 \mu\text{g C L}^{-1} \text{d}^{-1}$), which showed the highest values compared to other depths (Fig. 3). Additionally, TB_{DIC} exhibited a weak relationship with HBP, and bacterial abundance (Fig. 4), and on average, TB_{DIC} ($4.09 \pm 0.38 \mu\text{g C L}^{-1} \text{d}^{-1}$) exceeded the HBP rate ($0.26 \pm 0.06 \mu\text{g C L}^{-1} \text{d}^{-1}$) by over one order of magnitude. We found no significant correlation between TB_{DIC} and temperature, and nutrient concentration (Fig. 4). However, TB_{DIC} showed a high significant correlation with POC ($\rho = 0.80$, $p < 0.0001$), and a weak tendency to increase with increasing Chl-a ($\rho = 0.38$, $p < 0.01$, Fig. (4)).

290



295 **Figure 3:** Total bacteria DIC uptake rate ($\text{TB}_{\text{DIC}} \mu\text{g C L}^{-1} \text{d}^{-1}$) measured at the photic zone and in deep waters (400 m), where different letters indicate statistically significant differences (“A” is significantly higher than “B”). Each box plot indicates the 95 % confidence intervals with ± 1 SD deviation. The central line in the box represents the median for each group of samples.

300

The $\text{Total}_{\text{DIC}}$ fixation rate which is the uptake rate of both TB_{DIC} and PP fixation rate was ranged from 2.03 up to $60.45 \mu\text{g C L}^{-1} \text{d}^{-1}$. We found a significant and positive correlation between TB_{DIC} fixation and PP ($\rho = 0.53$, $p < 0.0001$, Fig. 4). On average, the contribution of TB_{DIC} % to the $\text{Total}_{\text{DIC}}$ daily fixation was $33.95 \pm 0.02 \%$, ranging from 13% to 72% across samples (Table 2). The TB_{DIC} % contribution was independent of the absolute TB_{DIC} fixation rate ($R^2 = 0.05$, $p = 0.09$), and was negatively correlated with PP ($R^2 = 0.45$, $p < 0.0001$, Fig. 5A). and did not

305



310

differ between open and coastal waters (F ratio = 0.008, df = 1, p = 0.92). The high TB_{DIC} at 400 m (1.37 to 6.33 $\mu\text{g C L}^{-1} \text{d}^{-1}$, Table 2), implied that the contribution of TB_{DIC} at 400 m was high (40 %) relative to surface PP (Fig. 5B and Table 2).




Figure 4: Scatterplot matrix plot, lower diagonal, and Spearman correlation coefficient (upper diagonal) among Total Bacterial Dark DIC uptake (TB_{DIC}), Total Bacterial Dark DIC uptake contribution ($TB_{DIC} \%$), Primary Production (PP), Particulate Organic Carbon (POC), Temperature (Temp), Bacterial Production (BP), Nitrate (NO_3), Chlorophyll -a concentration (Chl-a), and bacterial abundance. * $p < 0.01$, ** $p < 0.001$, *** $p < 0.0001$.

315






Figure 5: Figure (A) shows the significant probability of the relationship between the percentage contribution of TB_{DIC} ($TB_{DIC} \%$) and the primary productivity (PP), whereas (B) shows the total bacterial DIC uptake contribution ($TB_{DIC} \%$) to the total DIC fixation within the photic zone and the deep 400 m. The boxes with the same letters are not statistically significant, as determined by pairwise comparisons via Student's t-test.

4 Discussion

Although the contribution of heterotrophic bacteria to dark CO_2 fixation was discovered more than 80 years ago (Wood & Werkman, 1936), quantitative assessments of dark heterotrophic bacteria CO_2 fixation remain few (Braun et al., 2021), particularly relative to primary production rates. Our study successfully used the light and dark ^{13}C -bicarbonate additions coupled with CRDS-Picarro to accurately quantify the DIC incorporation by phytoplankton, heterotrophic bacteria, and chemo-autotrophs during daytime and nighttime, respectively. Decades after S. Nielsen (1952) introduced the ^{14}C method to measure phytoplankton primary production, the ^{13}C -PP method has recently garnered increasing attention as an alternative to the use of radioactive isotopes. Here, we extend the use of ^{13}C -bicarbonate additions to also resolve dark DIC uptake by bacteria, providing, to the best of our knowledge, the first application of this method coupled with CRDS-Picarro. Indeed, most previous studies measured dark DIC fixation using the ^{14}C -bicarbonate method (Baltar et al., 2010; Reinthaler et al., 2010; Yakimov et al., 2014; Zhou et al., 2017; Alonso-Sáez et al., 2010; Signori et al., 2017; Llirós et al., 2011) or genome and 16S ribosomal analysis (González et al., 2008; Yakimov et al., 2014), whereas very few studies have used ^{13}C -bicarbonate to measure dark DIC fixation (Roslev et al., 2004). One of the main advantages of using the stable isotope instead of radioactive isotopes additions is that it greatly reduces the health and safety issues associated with using radioactive products. Additionally, this method can be easily applied to characterize field samples (Middelburg et al., 2000; Boschker and Middelburg, 2002). Recent studies have demonstrated that the phytoplankton photosynthesis quantification results obtained with the ^{13}C -PP method coupled with CRDS-Picarro were similar to those achieved with the ^{14}C -method (López-Sandoval et al. 2018; López-Sandoval et al. 2019). Cavity ring down spectroscopy (CRDS), particularly with the Picarro analyzer, offers several advantages for stable isotope analysis applications. CRDS can detect trace amounts of isotopes in samples with very high sensitivity. This makes it particularly useful for analyzing low-concentration samples like oligotrophic waters. The high sensitivity of CRDS also allows for high precision and accuracy in isotopic measurements. (Berden et al., 2000; López-Sandoval et al., 2019).

Previous studies have confirmed that bacterial dark CO_2 uptake contributes significantly to the carbon flux dynamics of oligotrophic waters, shallow waters, coastal waters, and deep seas (Alonso-Sáez et al., 2010; Yakimov et al., 2014; Zhou et al., 2017., Signori et al., 2017, and LLiro's et al., 2011). Here, we confirmed the relevance of the dark CO_2 fixation processes in the oligotrophic Red Sea. Our estimates were within the range of reported dark DIC fixation rate, which range from $0.001 \mu g C L^{-1} d^{-1}$ in surface waters of the Subtropical North Atlantic and tropical estuarine systems (Reinthaler et al., 2010; Signori et al., 2017) to $206 \mu g C L^{-1} d^{-1}$ in an eutrophic Mediterranean coastal lagoon (LLiro's et al., 2011). In contrast, the highest values recorded in our study were found



355 in coastal waters and were similar to those reported from oligotrophic ocean waters (Alonso-Sáez et al., 2010;
LLiro's et al., 2011; Zhou et al., 2017). The deepest dark CO₂ fixation rate measurement yet reported was carried
out in the Mediterranean Sea at 4900 m, estimated at 0.096 ± 0.02 (Yakimov et al., 2014). We recorded a relatively
high TB_{DIC} uptake rate in Red Sea surface waters, toward the base of the photic zone but reaching high values at 400
m. Increases in dark DIC fixation rates at depth have been observed in the tropical South China Sea at depths
360 between 200 and 1500 m, with rates even exceeding those at the surface (Zhou et al., 2017). The high TB_{DIC} values
reported in the surface and deep water in our study suggest that dark DIC fixation not only contributes significantly
to the carbon fixation dynamics of surface water, where bacterial abundance is high, but also throughout the entire
(oxygenated) water column (Reinthal et al., 2010; Yakimov et al., 2014; Zhou et al., 2017). Therefore, given the
high TB_{DIC} values measured in the oligotrophic Red Sea ecosystem, our findings highlighted the importance of
365 accounting for dark DIC fixation in total carbon production estimations.

The Red Sea is characterized by a warm temperature throughout the water column (Shaltout, 2019), with average
temperature of 21.46 ± 0.23 °C at 400 m depth recorded in our study. Here, temperature was found to have a
positive correlation with PP but no correlation with the TB_{DIC} uptake rate, whereas Chl-a had no correlation with PP
as PP decreased gradually down the photic zone while Chl-a showed the maximum values in the bottom of the
370 photic zone. However, Chl-a showed a positive correlation with TB_{DIC}. Nitrate showed a negative correlation with
PP, suggesting nitrate depletion by the more productive phytoplankton communities, whereas TB_{DIC} was positively
correlated with PP. Studies conducted in the North Atlantic Ocean and tropical estuarine systems have reported a
weak relationship between dark CO₂ fixation and temperature, nutrients, and Chl-a (Reinthal et al., 2010; Signori
et al., 2017). In contrast, a study conducted in the South China Sea reported that dark CO₂ fixation rates increased in
375 with temperature and nutrient concentration (Zhou et al., 2017).

The significant negative relationship between TB_{DIC} % and PP confirmed the relevance of the dark CO₂ fixation
toward the most oligotrophic waters. Bacterial dark CO₂ fixation rate (TB_{DIC}) contributed significantly to the total
DIC uptake within the photic zone, where it reached up to 72% of the photosynthetic primary production and up to
52% of the surface PP in the deep water at 400 m. In the eastern North Atlantic Ocean, dark CO₂ fixation was
380 reported to support 72% of the prokaryotic carbon demand in the mesopelagic layers below 200 m (Baltar et al.,
2010). Baltar and Herndl (2019) analyzed data collected over the course of 30 years and found that the dark
CO₂ uptake rate contributed up to 22% of the total PP at the euphotic layer (0–150 m). Additionally, increasing
evidence has suggested that dark CO₂ uptake by heterotrophic bacteria contributes significantly to surface
CO₂ fixation, contributing up to 30% of the DIC uptake in some oligotrophic waters (González et al., 2008;
385 Palovaara et al., 2014). Similarly, in our study we recorded an average contribution of TB_{DIC} to total DIC uptake of
33.95 % within the photic zone and the deep 400 m water. The increase in the contribution of TB_{DIC} to the carbon
flux with decreasing PP in the Red Sea highlights the importance of dark DIC fixation as a key mechanism driving
plankton communities, particularly in highly oligotrophic environments with low (surface) or absent (deep water)
primary production. Additionally, the relevance of dark CO₂ incorporation is likely significant in oligotrophic and
390 nutrient-depleted environments where the availability of labile organic carbon is limited (González et al., 2008,



Alonso-Sáez et al., 2010). Overall, our results confirm the relevance of dark CO₂ fixation to the oligotrophic Red Sea.

Our findings indicate that TB_{DIC} exceeded HBP, as reported in previous studies (Zhou et al., 2017), confirming the important role of TB_{DIC} in fueling bacterial metabolism in oligotrophic waters, with low levels of labile organic matter for bacterial growth such as surface and deep oligotrophic waters. The high TB_{DIC} contribution to total DIC uptake can support the growth of the bacterial community, providing a path to support bacterial metabolism, respiration and carbon flux in the microbial loop (Zhou et al., 2017). The importance of dark CO₂ fixation processes seemed to increase with depth in the Red Sea, as the dark/light ratio of CO₂ fixation rate increased in deeper waters, reaching up to 1.13 ± 0.65 toward the base of the photic zone. These findings were consistent with those of Baltar and Herndl (2019), who reported that the dark/light ratio reached 1 at 120–160 m depths from data collected along the ALOHA and BATS; the longest oceanic time series of the Atlantic and Pacific Ocean, highlighting the urgent need to account for dark DIC fixation in future studies on total primary production. Considering the total net primary production in the ocean to be approximately 50 Pg C y⁻¹ as reported by Field et al. (1998) and based on the potential TB_{DIC} % contribution of 13.18% to 71.78% to the Total_{DIC} fixation reported in our study, we estimated that approximately 6.5 to as much as 35.5 Pg C y⁻¹ could be added to the global primary production estimation. This would be a considerable amount of carbon productivity by heterotrophs that is not being accounted for in current carbon flux and production estimations, which could represent a significant source of carbon in surface and deep waters (Baltar and Handle, 2019).

5 Conclusions

Using a stable isotope method, our study demonstrated the substantial contribution of dark CO₂ assimilation by heterotrophic bacteria in the oligotrophic Red Sea. The results presented herein represent a first attempt to estimate and confirm the role of dark heterotrophic bacteria CO₂ assimilation to the carbon flux dynamics along the Red Sea at different depths and in different water bodies. Even though temperature, a uniquely influential feature of the warm Red Sea, appeared to have a weak correlation with TB_{DIC}, our study confirmed that the importance of anaplerotic CO₂ incorporation and chemo-autotrophic process would be significant in environments with low or absent primary productivity. Moreover, due to the large fraction of Total_{DIC} fixation generated from the contribution of TB_{DIC} in the surface and the deep water, as reported in our study and other studies, it is essential to account for the contribution of heterotrophic dark CO₂ fixation to the total DIC fixation as a source of prokaryotic carbon demand.

Data availability. The data are presented in the manuscript; and can be requested from the corresponding author

Author contribution. Each of the authors made substantial contributions to the conception, design, and execution of this study. Prof. Susana Agustí was responsible for the development of the study design and goals, data analysis, critical revisions of the manuscript, and overall project coordination. Prof. Carlos Duarte contributed to the development of the study design, interpretation of the data, and critical revisions of the manuscript, in addition to his



coordination to conduct the project. In addition, Prof. Carlos provided the raw data run by CRDS-Picarro in his laboratory. Dr. Daffne López-Sandoval was involved in the data collection and analysis, and provided critical feedback on the manuscript. Afrah Alothman was responsible for data collection, data analysis and drafting the
430 manuscript. In addition, Afrah reviewed any critical revisions made by the co-authors. All authors have read and approved the final manuscript for submission.

Competing interests. The authors declare that they have no conflict of interest.

435 *Acknowledgment:* We would like to express our deepest gratitude to each and every one involved in this study, and the King Abdullah University of Science and Technology who have made this research possible. First and foremost, we would like to thank and show our grateful to all participants who generously shared their time and experiences to enclosed this research including all the crew members and the scientific leaders of the vessels and cruises involved
440 in this study. In addition, we would like to thank Coastal & Marine Resources Core Lab (CMR) team at King Abdullah University of Science and Technology (KAUST) who were behind the possibility of conducting coastal experiments outdoor set up. We also extend our appreciation to Mongi Ennasri who provided the support and assistance throughout the samples analysis in the CRDS-Picarro. Each and every one of these individuals and organizations support enabled us to collect and analyze the data, and to disseminate our findings.

445 *Financial support.* This work was supported by King Abdullah University of Science and Technology (KAUST).

References

- Alonso-Sáez, L., Galand, P. E., Casamayor, E. O., Pedros-Alio, C., & Bertilsson, S.: High bicarbonate assimilation in the dark by Arctic bacteria. *The ISME journal*, 4(12), 1581-1590, 2010.
- 450 Azam, F., Fenchel, T., J. G., Gray, J. S., Meyer-Reil, L. A., & Thingstad, F.: The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series*, 10, 257-263, 1983.
- Baltar, F., & Herndl, G. J.: Is dark carbon fixation relevant for oceanic primary production estimates. *Biogeosciences*, 16, 3793-3799, 2019.
- Baltar, F., Aristegui, J., Sintes, E., Gasol, J. M., Reinthaler, T., & Herndl, G. J.: Significance of non-sinking particulate
455 organic carbon and dark CO₂ fixation to heterotrophic carbon demand in the mesopelagic northeast Atlantic. *Geophysical Research Letters*, 37(9), 2010.
- Baltar, F., Lundin, D., Palovaara, J., Lekunberri, I., Reinthaler, T., Herndl, G. J., & Pinhassi, J.: Prokaryotic responses to ammonium and organic carbon reveal alternative CO₂ fixation pathways and importance of alkaline phosphatase in the mesopelagic North Atlantic. *Frontiers in Microbiology*, 7, 1670, 2016.
- 460 Berden, G., Peeters, R., & Meijer, G.: Cavity ring-down spectroscopy: Experimental schemes and applications. *International Reviews in Physical Chemistry*, 19(4), 565-607, 2000.



- Braun, A., Spona-Friedl, M., Avramov, M., Elsner, M., Baltar, F., Reinthaler, T., Herndl, G.J. and Griebler, C.: Reviews and syntheses: Heterotrophic fixation of inorganic carbon—significant but invisible flux in environmental carbon cycling. *Biogeosciences*, 18(12), 3689-3700, 2021.
- 465 Boschker, H. T. S., & Middelburg, J. J.: Stable isotopes and biomarkers in microbial ecology. *FEMS Microbiology Ecology*, 40(2), 85-95, 2002.
- Chaidez, V., Dreano, D., Agusti, S., Duarte, C. M., & Hoteit, I.: Decadal trends in Red Sea maximum surface temperature. *Scientific Reports*, 7(1), 1-8, 2017.
- Dickson, A. G., Sabine, C. L., & Christian, J. R.: Guide to best practices for ocean CO₂ measurements. *North Pacific Marine Science Organization*, 3, 191, 2007.
- 470 Dijkhuizen, L., & Harder, W.: Current views on the regulation of autotrophic carbon dioxide fixation via the Calvin cycle in bacteria. *Antonie van Leeuwenhoek*, 50(5-6), 473-487, 1984.
- Ducklow, H. Kirchman, D.: Bacterial production and biomass in the oceans. *Microbial Ecology of the Oceans*, 1, 85-120, 2000.
- 475 Edwards, F. J.: Climate and oceanography. *Red sea*, 1, 45-68, 1987.
- Erb, T. J.: Carboxylases in natural and synthetic microbial pathways. *Applied and Environmental Microbiology*, 77(24), 8466-8477, 2011.
- Gasol, J. M., & Morán, X. A. G.: Flow cytometric determination of microbial abundances and its use to obtain indices of community structure and relative activity. In *Hydrocarbon and Lipid Microbiology Protocols* Springer, Berlin, Heidelberg, 159-187, 2016.
- 480 González, J.M., Fernández-Gómez, B., Fernández-Guerra, A., Gómez-Consarnau, L., Sánchez, O., Coll-Lladó, M., Del Campo, J., Escudero, L., Rodríguez-Martínez, R., Alonso-Sáez, L. and Latasa, M.: Genome analysis of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria). *Proceedings of the National Academy of Sciences*, 105(25), 8724-8729, 2008.
- 485 Grasshoff, K.: Zur Chemie des Roten Meeres und des Inneren Golfs von Aden nach Beobachtungen von FS" Meteor" während der Indischen Ozean Expedition. *Borntraeger*, 1964/65, 1969.
- Hansen, H. P., & Koroleff, F.: Determination of Nutrients. *Methods of Seawater Analysis*, 159-228, 1999.
- Kirchman, D. L.: Uptake and regeneration of inorganic nutrients by marine heterotrophic bacteria. *Microbial Ecology of the Oceans*, 28, 255–271 2000.
- 490 Koshikawa, H., Harada, S., Watanabe, M., Kogure, K., Ioriya, T., Kohata, K., ... & Akehata, T.: Influence of plankton community structure on the contribution of bacterial production to metazooplankton in a coastal mesocosm. *Marine Ecology Progress Series*, 186, 31, 1999.
- Li, W. K. W., & Dickie, P. M.: Light and dark 14C uptake in dimly-lit oligotrophic waters: relation to bacterial activity. *Journal of plankton research*, 13(supp1), 29-44, 1991.
- 495 Li, W. K. W., Irwin, B. D., & Dickie, P. M.: Dark fixation of 14C: variations related to biomass and productivity of phytoplankton and bacteria. *Limnology and Oceanography*, 38(3), 483-494, 1993.



- Llirós, M., Alonso-Sáez, L., Gich, F., Plasencia, A., Auguet, O., Casamayor, E. O., & Borrego, C. M.: Active bacteria and archaea cells fixing bicarbonate in the dark along the water column of a stratified eutrophic lagoon. *FEMS Microbiology Ecology*, 77(2), 370-384, 2011.
- 500 López-Sandoval, D. C., Delgado-Huertas, A., Carrillo-de-Albornoz, P., Duarte, C. M., & Agustí, S.: Use of cavity ring-down spectrometry to quantify ^{13}C -primary productivity in oligotrophic waters. *Limnology and Oceanography: Methods*, 17(2), 137-144, 2019.
- López-Sandoval, D. C., Duarte, C. M., & Agustí, S.: Nutrient and temperature constraints on primary production and net phytoplankton growth in a tropical ecosystem. *Limnology and Oceanography*, 66(7), 2923-2935, 2021.
- 505 Markager, S.: Dark uptake of inorganic ^{14}C in oligotrophic oceanic waters. *Journal of Plankton Research*, 20(9), 1813-1836, 1998.
- Middelburg, J. J., Barranguet, C., Boschker, H. T., Herman, P. M., Moens, T., & Heip, C. H.: The fate of intertidal microphytobenthos carbon: An in situ ^{13}C -labeling study. *Limnology and Oceanography*, 45(6), 1224-1234, 2000
- Nielsen, E. S.: Dark fixation of CO_2 and measurements of organic productivity. With remarks on chemo-synthesis. *Physiologia Plantarum*, 13(2), 348-357, 1960.
- 510 Palovaara, J., Akram, N., Baltar, F., Bunse, C., Forsberg, J., Pedrós-Alió, C., ... & Pinhassi, J.: Stimulation of growth by proteorhodopsin phototrophy involves regulation of central metabolic pathways in marine planktonic bacteria. *Proceedings of the National Academy of Sciences*, 111(35), E3650-E3658, 2014.
- Prabowo, D.A. and Agustí, S.: Free-living dinoflagellates of the central Red Sea, Saudi Arabia: Variability, new records and potentially harmful species. *Marine Pollution Bulletin*, 141, pp.629-648, 2019.
- 515 Prakash, A., Sheldon, R. W., & Sutcliffe Jr, W. H.: Geographic variation of oceanic ^{14}C dark uptake. *Limnology and oceanography*, 36(1), 30-39, 1991.
- Reinthal, T., van Aken, H. M., & Herndl, G. J.: Major contribution of autotrophy to microbial carbon cycling in the deep North Atlantic's interior. *Deep Sea Research Part II: Topical Studies in Oceanography*, 57(16), 1572-1580, 2010.
- 520 Romanenko, V. I.: Heterotrophic assimilation of CO_2 by bacteria flora water. *Microbiology (Mikrobiologi)*, 33(4), 610-614, 1964.
- Roslev, P., Larsen, M. B., Jørgensen, D., & Hesselsoe, M.: Use of heterotrophic CO_2 assimilation as a measure of metabolic activity in planktonic and sessile bacteria. *Journal of Microbiological Methods*, 59(3), 381-393, 2004.
- Shaltout, M.: Recent Sea surface temperature trends and future scenarios for the Red Sea. *Oceanologia*, 61(4), 484-504, 2019.
- 525 Signori, C. N., Valentin, J. L., Pollery, R. C., & Enrich-Prast, A.: Temporal variability of dark carbon fixation and bacterial production and their relation with environmental factors in a tropical estuarine system. *Estuaries and Coasts*, 41(4), 1089-1101, 2018.
- Steemann-Nielsen, E.: On the determination of the activity for measuring primary production. *J. Cons. Int. Explor. Mer*, 18, 117-140, 1952.
- 530 Wafar, M., Qurban, M. A., Ashraf, M., Manikandan, K. P., Flandez, A. V., & Balala, A. C.: Patterns of distribution of inorganic nutrients in Red Sea and their implications to primary production. *Journal of Marine Systems*, 156, 86-98, 2016.



- 535 Wood, H. G., & Werkman, C. H.: The utilisation of CO₂ in the dissimilation of glycerol by the propionic acid bacteria. *Biochemical Journal*, 30(1), 48, 1936.
- Yakimov, M.M., La Cono, V., Smedile, F., Crisafi, F., Arcadi, E., Leonardi, M., Decembrini, F., Catalfamo, M., Bargiela, R., Ferrer, M. and Golyshin, P.N.: Heterotrophic bicarbonate assimilation is the main process of de novo organic carbon synthesis in hadal zone of the Hellenic Trench, the deepest part of Mediterranean Sea. *Environmental Microbiology Reports*, 6(6), 709-722, 2014.
- 540 Yao, F., & Hoteit, I.: Rapid red sea deep water renewals caused by volcanic eruptions and the North Atlantic oscillation. *Science advances*, 4(6), eaar5637, 2018.
- Zhou, W., Liao, J., Guo, Y., Yuan, X., Huang, H., Yuan, T., & Liu, S.: High dark carbon fixation in the tropical South China Sea. *Continental Shelf Research*, 146, 82-88, 2017.