

# Short-term variation of pH in seawaters around coastal areas of Japan: Characteristics and forcings

Tsuneo Ono<sup>1</sup>, Daisuke Muraoka<sup>2</sup>, Masahiro Hayashi<sup>3</sup>, Makiko Yorifuji<sup>3\*</sup>, Akihiro Dazai<sup>4</sup>, Shigeyuki Omoto<sup>5</sup>, Takehiro Tanaka<sup>6</sup>, Tomohiro Okamura<sup>7</sup>, Goh Onitsuka<sup>7</sup>, Kenji Sudo<sup>7</sup>, Masahiko Fujii<sup>8#</sup>, Ryuji Hamanoue<sup>8</sup>, and Masahide Wakita<sup>9</sup>

<sup>1</sup>Fisheries Resources Institute, Japan Fisheries Research and Education Agency, Yokohama, 236-8648, Japan

<sup>2</sup>Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Miyako, 027-0097, Japan

<sup>3</sup>Marine Ecology Research Institute, Kashiwazaki, 945-0017, Japan

<sup>4</sup>Center for Sustainable Society, Minami-Sanriku, 986-0775, Japan

10 <sup>5</sup>Eight-Japan Engineering Consultants Inc., Okayama, 700-8617, Japan

<sup>6</sup>Satoumi Research Institute, Okayama, 704-8194, Japan

<sup>7</sup>Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Hatsukaichi, 739-0452, Japan

<sup>8</sup>Faculty of Environmental Earth Science, Hokkaido University, Sapporo, 060-0810, Japan

<sup>9</sup>Mutsu Institute for Oceanography, Japan Agency for Marine-Earth Science and Technology, Mutsu, 035-0022, Japan

15 \*now at National Institute of Advanced Industrial Science and Technology, Tsukuba, 305-8567, Japan

#now at Otsuchi Coastal Research Center, Atmosphere and Ocean Research Institute, the University of Tokyo, Otsuchi, 028-1102, Japan

*Correspondence to:* Tsuneo Ono (tono@fra.affrc.go.jp)

**Abstract.** The pH of coastal seawater varies based on several local forcings, such as water circulation, terrestrial inputs, and biological processes, and these forcings are changing along with global climate change. Understanding the mechanism of pH variation in each coastal area is thus important for a realistic future projection that considers changes in these forcings. From 2020 to 2021, we performed parallel year-round observations of pH and related ocean parameters at five stations around the Japanese coast (Miyako Bay, Shizugawa Bay, Kashiwazaki Coast, Hinase Archipelago, and Ohno Strait) to understand the characteristics of short-term pH variations and their forcings. Annual variability (~1 standard deviation) of pH and aragonite saturation state ( $\Omega_{ar}$ ) were 0.05–0.09 and 0.25–0.29, respectively, for three areas with low anthropogenic pressures (Miyako Bay, Kashiwazaki Coast, and Shizugawa Bay), while it increased to 0.16–0.21 and 0.52–0.58, respectively, in two areas with medium anthropogenic pressures (Hinase Archipelago and Ohno Strait in Seto Inland Sea). Statistical assessment of temporal variability at various timescales revealed that most of the annual variabilities in both pH and  $\Omega_{ar}$  were derived by short-term variation at a timescale of < 10 days, rather than seasonal-scale variation. Our analyses further illustrated that most of the short-term pH variation was caused by biological processes, while both thermodynamic and biological processes equally contributed to the temporal variation in  $\Omega_{ar}$ . The observed results showed that short-term acidification with  $\Omega_{ar} < 1.5$  occurred occasionally in Miyako and Shizugawa Bays, while it occurred frequently in the Hinase Archipelago and Ohno Strait. Most of such short-term acidified events were related to short-term low-salinity events. Our analyses showed that the amplitude of short-term pH

variation was linearly correlated with that of short-term salinity variation, and its regression coefficient at the time of high  
35 freshwater input was positively correlated with the nutrient concentration of the main river that flows into the coastal area.

## 1 Introduction

The ocean is witnessing a reduction in its pH due to anthropogenic CO<sub>2</sub> sequestration both in open (e.g., Bates et al.,  
2014; Iida et al., 2021; Jiang et al., 2019; Lauvset et al., 2015; Takahashi et al., 2014) and coastal oceans (e.g., Carstensen and  
Duarte, 2019; Duarte et al., 2013; Hauri et al., 2013; Ishida et al., 2021; Ishizu et al., 2019; Yao et al., 2022). In the coastal  
40 ocean, pH shows short time variation caused by several processes such as water mass changes (e.g., Johnson et al., 2013; Ko  
et al., 2016; Wakita et al., 2021), coastal upwelling (e.g., Barton et al., 2012; Booth et al., 2012; Feely et al., 2008, 2016;  
Vargas et al., 2015), rivers inputs (e.g., Cai et al., 2017; Fujii et al., 2021; Gomez et al., 2021; Salisbury et al., 2008; Salisbury  
and Jönsson, 2018), terrestrial nutrients' inputs (e.g., Cai et al., 2011; Guo et al., 2021; Kessouri et al., 2021; Provoost et al.,  
2010; Sunda and Cai, 2012; Wallace et al., 2014), and various coastal biological processes (e.g., Delille et al., 2009; Lowe et  
45 al., 2019; Mongin et al., 2016; Ricart et al., 2021; Yamamoto-Kawai et al., 2021). The amplitude of short-term pH variation  
often exceeds that of the decadal-scale long-term pH trend, and hence, blurs the signal of anthropogenic CO<sub>2</sub>-induced  
acidification of coastal seawater (e.g., Borges and Gypens, 2010; Duarte et al., 2013, Johnson et al., 2013; Provoost et al.,  
2010; Salisbury and Jönsson, 2018). Short-term pH variations in coastal waters are important for local ecosystems as they are  
mostly caused by natural forcings that have been acting before the industrial period, and hence, the local ecosystem is expected  
50 to adapt to such short-term pH variations as long as they are natural in terms of timing and amplitude. For example, *Ostrea  
lurida*, a native oyster in Netarts Bay, Oregon, USA, has adjusted its spawning season before and after the summer upwelling  
season so that its larvae can avoid low-pH waters (Waldbusser et al., 2014). Several anthropogenic perturbations, such as  
changes in land use, sewerage treatment, vanishment of seagrass bed, and modification of coastal topography, can change  
these forcings, shifting the timing and/or amplitude of natural short-term pH variation (e.g., Hoshiba et al., 2021; Papalexioiu  
and Montanari, 2019). Understanding the present situation as well as the mechanism of short-term pH variation in coastal  
60 waters is thus critical for evaluating the risk of acidification in coastal areas.

Japan consists of 14,125 islands distributed in a wide latitudinal range from 20° N to 45° N in the western North  
Pacific, containing diverse coastal environments from coral reefs to seasonal floating sea ice. Japan is a highly developed  
country, and a significant portion of its coastal area has experienced various types of anthropogenic perturbations. The  
60 country's Ministry of the Environment (MOE) conducts regular pH monitoring at over 2,000 coastal stations around Japan  
from the early 1980s until the present, and the obtained data showed significant variability in the multi-decadal pH trend from  
−0.012 y<sup>−1</sup> to +0.009 y<sup>−1</sup> throughout the stations (Ishizu et al., 2019). The observed range of pH trend within the Japanese coast  
is equivalent to 85% of that observed in 83 coastal systems in the world (Carstensen and Duarte, 2019), and this result suggests  
that the Japanese coastal area can be considered as a “sample shelf” for coastal acidification studies. The MOE monitors pH

65 monthly and seasonally, and several of pH stations of MOE, especially those in northern areas, lack wintertime observations (Ishizu et al., 2019). We thus need additional pH observations with a higher time resolution to understand the characteristics of short-term pH variation at a timescale of < 30 days around the Japanese coast to assess its variability and mechanisms. A number of scientists have already started such pH monitoring in coastal stations in Japan (e.g., Christian and Ono, 2019; Fujii et al., 2021, 2023; Ishida et al., 2021; Wakita et al., 2021). However, most of these observations were started recently and the  
70 summarisation of observed results among these stations is yet to be made. In this study, we carried out the first synthesis effort of such high-resolution pH monitoring stations along coastal areas in Japan that are operated by different founders/programs, by summarising continuous monitoring data of pH observed during 2020–2021 at five stations around the Japanese coast. Here, we describe and discuss the amplitude of pH variation in each timescale, similarity, and dissimilarity in the characteristics of variation, and their forcings.

75

## 2 Observations and settings of study areas



**Figure 1: Map of the five study stations along the coast of Japan.**


farms in the coastal area (Kakehi et al., 2018). Additional nutrients to the bay are provided by three rivers: Hei, Tsugaruishi,

Hydrographic monitoring, including pH monitoring, was performed at five stations around the coast of Japan in the following stations, Miyako Bay, Shizugawa Bay, Kashiwazaki Coast, Hinase Archipelago, and Ohno Strait (Fig. 1), from 2020 to 2021. Miyako Bay and Kashiwazaki Coast were selected to represent coastal environment with relatively low anthropogenic nutrient loadings, while the other three stations were selected from major farming areas of pacific oyster. The detailed settings of the areas and observation procedures are described in the following sections.

### 2.1 Miyako Bay

Miyako Bay is located in the northern part of Honshu Island facing the western North Pacific, with a bay area of 24 km<sup>2</sup> and 4.8-km wide bay mouth (Fig. 2(a)). The outer bay area is usually occupied by temperate western North Pacific water, which brings enough nutrients in winter to support seaweed

95 and Tashiro (25.8, 5.16, and 3.78 m<sup>3</sup> s<sup>-1</sup>, respectively, for annual average flow rate; Okada et al., 2014), although the quantity of nutrients input by the rivers are limited to low levels (57, 11, and 8 tN y<sup>-1</sup> for Hei, Tsugaruishi, and Tashiro Rivers, respectively; Bernardo et al., 2023). With a population of 60,000 residents within the hinterland, Miyako Bay has maintained good water quality with 1–2 mg L<sup>-1</sup> of chemical oxygen demand (MOE, 2022). While kelp beds are well developed near the shoreline, wakame seaweed (*Undaria pinnatifida*) farmyards are developed in the middle of the bay.




**Figure 2: Map of Miyako Bay with the location of the station (solid circle). Overlaid map shows detailed structure of the station. Grey square and solid triangle represent locations of settling tank and water intake, respectively.**

The monitoring site was located in front of the Miyako Field Station of the Japan Fisheries Research and Education Agency, which is located north of the bay mouth (141° 58' 5" E and 39° 41' 28" N; Fig 2). This area frequently encounters severe winter storms, so we set the monitoring station in the settling tank (3.6 × 2.8 × 5.4 m) of the field station, in which coastal water is continuously pumped from the water intake located 200 m off the coastline and 1–3 m in depth (Fig. 2). Sensors for pH (Kimoto Electronic, SPS-14), dissolved oxygen (DO) (JFE Advantech, AROW2), and salinity/water temperature (JFE Advantech, ACTW) were then moored at a depth of 1 m in the settling tank. Mechanical precision of each sensor is ±0.003 pH units for pH, ±2% for DO, ±0.008 for salinity and ±0.01 °C for water temperature, respectively. The difference in pH between water intake and settling tank was measured for two weeks during the monitoring period, and it was confirmed that the difference in pH between the water intake and settling tank was < 0.006 pH units. The sampling frequency of each sensor was set to 1 h. All sensors were replaced every 2 months because of the limitation of batteries, and the DO sensor was calibrated by air-saturated pure water and sodium sulphite solution, while the pH sensor was calibrated against

tris(hydroxymethyl)aminomethane and 2-amino-2-methyl-1-propanol-buffered artificial seawaters provided by FUJIFILM Wako Pure Chemical Corporation (Cat. No. 017-28191 and 010-28181, respectively) at the beginning of each deployment. Along with these measurements, discrete water samples were taken from the tank at a depth of 1 m using a 1.5-L Niskin bottle every week. Subsamples for salinity, nutrients, dissolved inorganic carbon (DIC), and total alkalinity (TA) were then taken and stored. Salinity and nutrients were measured at the Yokohama Laboratory of the Japan Fisheries Research and Education Agency using a salinometer (Guildline Instruments 8400 B) and continuous flow analyser (Seal Analytical QuAAtro 39), respectively, while DIC and TA were measured at the Mutsu Institute for Oceanography of the Japan Agency for Marine-Earth

125

Science and Technology (MIO-JAMSTEC) using a carbon coulometer (UIC CM 5012 with Nippon ANS MODEL 3000A) and an open-cell titrator (Kimoto Electric ATT-15) calibrated against the seawater reference materials provided by KANSO Corporation (Wakita et al., 2021). Measurement precision was  $\pm 1 \mu\text{mol kg}^{-1}$  for both DIC and TA. The pH of the seawater at the time of each sampling ( $\text{pH}_{\text{discrete}}$ ) was calculated from DIC and TA using the CO2sys program v2.1 (Pierrot and Wallace, 2006), with the settings of Lueker et al. (2000) for the dissociation constant of carbonate, Dickson (1990) for the dissociation constant of bisulphate, and Lee et al. (2010) for the aqueous boron concentration. The drift of the pH sensor during deployment was corrected based on the difference between  $\text{pH}_{\text{discrete}}$  and the pH measured by the sensor. The relationship between the measured TA and salinity was evaluated as a linear function, and the time series of TA throughout the sensor deployment was calculated based on the salinity–TA relationship. The saturation states of calcite ( $\Omega_{\text{ca}}$ ) and aragonite ( $\Omega_{\text{ar}}$ ) were then calculated from water temperature, salinity, nutrient concentrations, DIC, and TA using the CO2sys program. This monitoring program, founded by the Study on Biological Effects of Acidification and Hypoxia (BEACH) of the Environment Research and Technology Development Fund of the Environmental Restoration and Conservation Agency, started on 1 July 2020 and ended on 21 September 2021.



**Figure 3: (a) Detailed map of Kashiwazaki Coast with the location of the station (solid circle). (b) Locations of water intake and settling tank in Kashiwazaki Observatory.**

## 2.2 Kashiwazaki Coast

Kashiwazaki City is located at the centre of Honshu Island facing the Sea of Japan. Its coastline has little flexure, with the periodic occurrence of shallow sandy beaches and small rocky reefs (Fig. 3(a)). While benthic biomass is quite low in sandy beach coastal areas, local *Sargassum* seaweed beds are formed off rocky reefs. The low-nutrient Tsushima Warm Current flows throughout the year off the narrow band of low-salinity coastal water (Niigata Prefectural Institute for Fisheries and Oceanography, 1998). Kashiwazaki City has a population of 79,000 with a cultivated land area of 4,890 ha (mainly rice paddy fields) and ~200 plants of manufacturing industries. Sewage waters from these civil activities are released into coastal areas via two small rivers, Sabaishi and Ukawa Rivers, which bring low salinity and nutrients. The quality of off-Kashiwazaki coastal waters has been well maintained, with a chemical oxygen demand of  $< 2 \text{ mg L}^{-1}$  (MOE, 2022). However, 30 years of monitoring by the Marine Ecology Research Institute shows that

the pH of off-Kashiwazaki coastal water has decreased at a rate of  $-0.003 \text{ y}^{-1}$  because of the increase in water temperature and concentration of atmospheric  $\text{CO}_2$  (Ishida et al., 2021).

Similar to the Miyako site, sensors for pH, DO, and salinity/water temperature were set within the water tank (3,000 L), to which coastal water was continuously pumped from the water intake set in front of the Marine Ecology Research Institute (138 °35' 24" E and 37 °25' 30" N, Fig. 3(b)). The settings of each sensor, water sampling, and measurements were the same as those of the Miyako site. This monitoring program, also founded by BEACH, started on 16 July 2020 and ended on 24 August 2021.

### 2.3 Shizugawa Bay



**Figure 4: Detailed map of Shizugawa Bay with the location of the station. Open circles represent stations installed by the Ocean Acidification Adaptation Project. Solid circle represents the location of station S3 used in this study.**

Shizugawa Bay is located  $\sim 100$  km south of Miyako Bay, with a bay area of  $46.8 \text{ km}^2$  and a  $6.6\text{-km}$  wide bay mouth (Fig. 4). The Oyashio-oriented cold-water and Kuroshio-oriented warm-water masses intermittently occupy the outer bay, and the former water brings high quantities of nutrients to this bay in winter, similar to that in the Miyako Bay. The large Kitakami River (water transport of  $390 \text{ m}^3 \text{ s}^{-1}$ ), flows out to the sea south of Shizugawa Bay, bringing nutrients of  $6 \times 10^3 \text{ tN y}^{-1}$  to the coastal area (Sugimura et al., 2015). In addition, 10 small rivers flow directly into the bay, and the total nutrient flux provided by these rivers was estimated at  $270 \text{ tN y}^{-1}$  (Yamamoto et al., 2018). Shizugawa Bay has been widely used for the aquaculture industry, especially for culturing Pacific oysters and silver salmon. Historically, the extent of aquaculture utilisation occasionally becomes too heavy, resulting in the emergence of low-oxygen deep waters in the bay caused by the degradation of organic materials derived from aquaculture. After the Great East Japan Earthquake in 2011, oyster farmers in Shizugawa Bay decided to reduce the number of their oyster rafts to one third of what they had before the earthquake to reduce the environmental impact of aquaculture on the bay ecosystem, which was on the way to recovery from the tsunami disaster. As a result, the DO of the bottom water in the bay showed a remarkable increase **by**  $> 1 \text{ mg L}^{-1}$  (Yamamoto et al., 2017).

Since 2020, hydrographic conditions, including pH, have been observed at four stations within Shizugawa Bay (Fig. 4) by the Ocean Acidification Adaptation Project (OAAP) founded by the Nippon Foundation (Fujii et al., 2023). Detailed settings of the observations are described in Fujii et al. (2023), and here, we reproduce its brief outline: Sensors for water temperature, salinity, DO, and pH were set at a depth of 1 m in each station to collect data at a temporal resolution of 1 h. Water samples for salinity, nutrients, DIC, and TA were taken from each station monthly, and the sampling interval was enhanced to every 15 days during summer. Salinity, nutrients, DIC, and TA were measured by the same methods as for the Miyako site, although measurements were made at the Kesen-numa Laboratory of Miyagi Prefectural Institute for Fisheries Sciences and MIO-JAMSTEC for nutrients and other properties, respectively. Similar to the Miyako site,  $\text{pH}_{\text{discrete}}$  was calculated from DIC and TA and used for the drift correction of the pH sensors. The time series of TA throughout the sensor deployment was also calculated based on the observed TA–salinity relationship. In this study, we used data from Station S3 (Fig.4) as a representative of Shizugawa Bay, as this station is located in oyster farming areas, the largest aquaculture industry, which is the largest source of organic carbon in this bay. The measurement of water temperature, salinity, and pH was started on 20 August 2020, while that of DO on 27 April 2021. All parameters are continuously measured till date; however, we use data from August 2020 to December 2021 to maintain synchronicity with data of other stations.

205

## 2.4 Hinase Archipelago




**Figure 5: Detailed map of Hinase Archipelago with the location of the station. Open circles represent stations installed by the Ocean Acidification Adaptation Project. Solid circle represents the location of station H2 used in this study.**

The Seto Inland Sea is the largest inner sea in Japan ( $\sim 2.32 \times 10^4$  km<sup>2</sup>) surrounded by the Honshu, Shikoku, and Kyushu Islands, and the Hinase Archipelago is located in the middle of the Seto Inland Sea. This area consists of four major waterways and many small canals divided by Honshu Island and eight other small islands with a water depth of  $\sim 6$  m (Fig. 5). The Seto Inland Sea is a moderately eutrophic area, receiving 53.0 km<sup>3</sup> y<sup>-1</sup> of freshwater and  $1.41 \times 10^5$  tN y<sup>-1</sup> of nitrate from the surrounding lands (Higashi et al., 2018; Nishijima, 2018). The water

exchange rate between the Seto Inland Sea and the outer North Pacific is also high (residence time of ~8 months; Yanagi and Ishii, 2004), and as a result, the nutrient concentration of surface water in the Seto Inland Sea is maintained at a moderate level (~0.4  $\mu\text{M}$  of DIN near the Hinase Archipelago; Tsukamoto and Yanagi, 1998). Historically, nutrient loading was the highest in the 1980s (~ $2.35 \times 10^5$  tN  $\text{y}^{-1}$ ; Nishijima, 2018). While the long-standing efforts of local communities succeeded in reducing it to the current level, significant quantities of organic nutrients still remained in the bottom sediment, creating local internal sources of nutrients in the Seto Inland Sea (e.g., Yamamoto et al., 2021). Although there is no large point source of nutrients, such as chemical plants, near the Hinase Archipelago, this area also receives nutrients from the Bizen city, that has a population of 36,000 residents. The Chikusa River, with an average flow rate of  $33.5 \text{ m}^3 \text{ s}^{-1}$ , flows in the eastern edge of this area, while several other small streams provide additional freshwater. The coastal area of the southern Hinase Archipelago is filled with eelgrass beds, while the northern coastal areas are partially filled with artificial shorelines, such as port facilities and breakwaters. Oyster aquaculture is widely used in open-water areas. The OAAP also launched four stations in this area (Fig. 5). Settings of sensors, water sampling, and its measurements were the same as those of Shizugawa Bay, with the exception that nutrient measurements were done by the Okayama Prefectural Institute of Fisheries Sciences (see Fujii et al., 2023 for details). In this study, we used data from Station H2 (Fig.5) to represent the Hinase Archipelago, as this station is located in the centre of oyster farming areas. Measurements of water temperature, salinity, and pH were started on 29 August 2020, while measurements of DO was started on 10 June 2021. All parameters are continuously measured till date; however, we use data from August 2020 to December 2021.

240

## 2.5 Ohno Strait



**Figure 6: Detailed map of Ohno Strait with the location of the station (solid circle).**

Ohno Strait is the small strait between Honshu and Miyajima Islands, which are located at the southern boundary of Hiroshima Bay in the western Seto Inland Sea (Fig. 6). As part of the Seto Inland Sea, the surface water surrounding Ohno Strait is moderately eutrophic (~2  $\mu\text{M}$  of DIN; Tsukamoto and Yanagi, 1998), although it is slightly lower than the Hinase Archipelago, reflecting an east-west gradient of surface nutrient concentration. The Ohta River, which flows through Hiroshima city with a population of 1.2 million, adds  $1.93 \times 10^3$  tN  $\text{y}^{-1}$  of nitrogen into Hiroshima Bay (Yamamoto et al., 2002). Part of the Ohta River water plume flows into the Ohno Strait (Abo and Onitsuka, 2019), providing nutrients to the surface layer of this strait. Industrial areas with a population of 115,000 are developed along the western coast of this strait,



255 providing another nutrient source to this area. The strait itself is lightly used for oyster farming, and the bottom sediment in the strait is mainly filled by silt and shell fragments.

The monitoring station was launched in the experimental raft off the Hatsukaichi Field Station of the Japan Fisheries Research and Education Agency, with a water depth of 3–6 m depending on the tide. A set of sensors (Kimoto Electronic SPS-14 for pH, JFE Advantech AROW2 for DO, and JFE Advantech ACTW for salinity/water temperature) were fixed from the raft at a  
260 depth of 1 m, and each parameter was measured at an interval of 1 h. The methods and frequencies for sensor calibration and discrete water sampling were the same as those of the Miyako site. This station was launched on 22 June 2021 by the OAAP and is continuing to operate till date; however, we used data from June 2021 to December 2021.

### 3 Results

#### 265 3.1 Temporal variation of water temperature, salinity, and DO

Figure 7 shows the time series of water temperature, salinity, and DO at the five stations. All stations showed similar seasonal variations in water temperature, with the highest and lowest temperatures in July and February, respectively (Fig. 7(a)). The seasonal amplitude was the **largest** in Hinase and the **smallest** in Miyako. At all stations, water temperature showed significant day-to-day variation with a timescale of < 10 days, and also diurnal variations. We discuss this further in Section  
270 4.1.

Time series of salinity in Shizugawa, Hinase, and Ohno show similar patterns of seasonal variation, low in summer/autumn and high in winter (Fig. 7(b)), suggesting that the main source of freshwater in these areas are rainfall events (during the rainy season in June/July and typhoon season in August–October). In contrast, in Kashiwazaki, salinity is high in summer and low in winter, suggesting that the main freshwater source in this area is snowfall in winter. The salinity of the  
275 Miyako site shows two low-salinity peaks, one in spring and the other in autumn, suggesting that this area is affected by both snowmelt waters and typhoon events (JFA, 2004). In Miyako, Shizugawa, Hinase, and Ohno, where the freshwater input is significant in summer/autumn, salinity frequently showed short-term drawdown that is synchronised with local rainfall events. The amplitude of sporadic salinity drawdown was extremely **large** in Hinase and Ohno, where the surface salinity temporally reached to less than 10 (Fig. 7(b)). Amplitude of seasonal annual variation in these two areas reaches over 25, which  
280 corresponds to the largest annual salinity amplitude among the worlds' 83 stations collected in Carstensen and Duarte (2019). We discuss the effect of these short-term low salinity events on pH in Section 4.3.

DO showed similar patterns of seasonal variation: low in summer and high in winter, although the durations were short in Hinase and Ohno (Fig. 7 (c)). This pattern suggests that the seasonal variation of DO is mainly driven by variation of oxygen solubility induced by water temperature rather than biological processes (see Section 4.2 for more details). This  
285 seasonal variation overlaps with day-to-day variation with a duration of ~10 days, and the amplitude of such day-to-day





Figure 7: Time series of (a) water temperature, (b) salinity, and (c) DO in the five stations. Legends of colour plots are the same for all panels as shown in Fig. 7(b).



290 **Figure 8:** (a) Time series of pH in each station. (b) An example of detailed pH time series. Data in Shizugawa Bay from 2 to 9 February 2021 are presented here. (c) Time series of  $pH_{min}$  in each station. Legends of colour plots are the same for all panels as shown in Fig. 8(c).

variation is especially significant in Hinase and Ohno in summer and autumn. Because of this short-term day-to-day variation, DO in Hinase and Ohno were occasionally below  $150 \mu\text{mol kg}^{-1}$  in summer and autumn, indicating that water conditions in these two areas occasionally become significantly undersaturated in DO, even in surface waters.

### 3.2 Temporal variation of pH

Figure 8(a) shows the time series of pH after drift correction at five stations (hereafter, all pH values are reported in total scale in this paper). Although the mechanistic precision of the pH sensor is  $\pm 0.003$  pH units, the uncertainty of the pH value measured by the glass electrode is mainly controlled by the precision of its drift correction. In the case of Miyako, where drift correction was performed weekly, we set two pH sensors in the same settling tank for two weeks to evaluate the reliability of pH data after the drift correction process. Values of pH obtained from the two sensors matched with that of 2 standard deviations (SD) of  $\pm 0.010$  pH units, and we used this value as the uncertainty of pH values obtained at Miyako, Kashiwazaki, and Ohno stations. In Shizugawa and Hinase, drift corrections were made at longer intervals, and Fujii et al. (2023) evaluated the uncertainty of pH at these two stations at  $\pm 0.015$  pH units. Figure 8(a) shows significant daily fluctuations far higher than these pH uncertainties. The diurnal amplitude was the **largest** in winter in Shizugawa Bay, where the difference in pH between daytime and night-time exceeded 0.8 pH units. **Large** diurnal variations in pH and/or  $p\text{CO}_2$  are often measured in shallow coastal ecosystems (e.g., Fujii et al., 2021; Ricart et al., 2021; Waldbusser et al., 2014). In most cases, **large** diurnal pH variation is observed in summer, and the **largest** diurnal amplitude of pH in winter has rarely been reported. Such patterns raise the possibility of biofouling in pH sensors (e.g., Venkatesan et al., 2017), although visual inspection at the time of sensor exchange did not show significant adherence of biomes during winter. However, the detailed pH variation in each day (Fig. 8(b)) showed that during the night-time it was relatively low and constant, even during significant diurnal variation. This is probably because the organic material produced by the fouling biomes during daytime had settled down from the sensors, and hence, the effect of the decomposition of organic materials during night-time remained low even in the **large** diurnal variation period. We do not further discuss diurnal variations of pH, as we do not have definite evidence for the effect of biofouling in winter in Shizugawa Bay. We alternatively calculated the daily minimum pH ( $\text{pH}_{\text{min}}$ ), that is usually observed at night, and analysed the day-to-day variation of  $\text{pH}_{\text{min}}$  at various timescales (Fig. 8(c)). Several rearing experiments have suggested that coastal organisms are affected by  $\text{pH}_{\text{min}}$  rather than the daily average pH (e.g., Onitsuka et al., 2018), therefore, the daily variation of  $\text{pH}_{\text{min}}$  was analysed instead of pH.

$\text{pH}_{\text{min}}$  showed a similar pattern of seasonal variations at all five stations, with high values in winter and low values in summer/autumn (Fig. 8(c)). Seasonal amplitude was the **smallest** in Miyako and the **largest** in Hinase and Ohno (Fig. 7(a)). In Hinase and Ohno, annual range of  $\text{pH}_{\text{min}}$  variation reached over 1.1 pH units, which was larger than that of 83 stations around the world listed in Carstensen and Duarte (2019) except for Baltic Sea. This seasonal variation overlapped with the short-term drawdown event of  $\text{pH}_{\text{min}}$ , which was frequently observed in summer/autumn. The timing of such events was


325 synchronised with that of short-term low-salinity events (Fig. 7(b)). This result indicates that either rainfall or increased river  
 flow causes several short-term processes in these coastal areas, causing short-term variation in pH.

### 3.3 Relationship between TA and salinity

Figure 9 shows the observed relationship between  
 330 TA and salinity based on discrete water samples obtained at  
 each station. TA in Miyako, Kashiwazaki, and Shizugawa  
 varied in a narrow range between 2,222 and 2,236  $\mu\text{mol}$   
 $\text{kg}^{-1}$  at a salinity of 33.5, which is approximately equal to  
 that in the surface waters of the western North Pacific in the  
 335 corresponding latitudinal range (Takatani et al., 2014). This  
 similarity of TA range indicates that coastal waters in these  
 three areas are occasionally not significantly different from  
 their open water sources when salinity is satisfactorily high  
 ( $\sim 33.5$ ). In contrast, in Hinase and Ohno, we obtained TA  
 340 of 2,222 and 2182  $\mu\text{mol kg}^{-1}$ , respectively, if we extended  
 their regression equation to a salinity of 34.0. These values  
 again fit with those of Kuroshio waters (Takatani et al.,  
 2014), reflecting that the ultimate source water for the Seto  
 Inland Sea is Kuroshio. It is noteworthy that the maximum  
 salinity values observed in Hinase and Ohno were 31.67 and  
 31.43, respectively (Fig. 7(b)). These results indicate that  
 345 the open waters of the Hinase and Ohno areas are already  
 diluted from their ultimate source (i.e., Kuroshio water),  
 and hence, have already witnessed several modifications  
 through coastal processes within the Seto Inland Sea.  
 350

The freshwater endmember in each regression line  
 represents the TA of the main freshwater source in each  
 area, although the absence of a low-salinity data point leads  
 to a high uncertainty in estimating the endmember. Table 1

355 describes the list of TA at the mouth of the major freshwater inflow in each coastal area. In most study areas, calculated  
 freshwater endmember of TA-S regressions agreed with the observed TA of one of freshwater source (Table 1), indicating that



**Figure 9: Salinity–TA relationship based on discrete water samples obtained from (a) Miyako, (b) Kashiwazaki, (c) Shizugawa, (d) Hinase, and (e) Ohno stations. Regression equations are provided below the X-axes.**

non-dilution/concentration changes of TA (Jiang et al. 2014) was relatively small in these areas, and by comparing these values with the freshwater endmember calculated from Fig. 9, we can speculate which river was the actual major freshwater source for each area. For the Miyako station, which is located at the northern edge of Miyako Bay, the Tashiro River becomes a major controller of salinity because of its proximity to the station. Two small rivers are neighbouring the Kashiwazaki site, and both rivers can act as salinity controllers for this site based on TA. Interestingly, all small rivers that flow into Shizugawa Bay had far higher TA than the calculated freshwater endmember, reflecting the existence of limestone areas in their basin. Rather, the TA of Kitakami River, despite flowing into a different bay south of Shizugawa, fitted with that of the freshwater endmember, suggesting that this river is the main source of freshwater for Station S3 because of its large flow rate. In the Hinase Archipelago, all surrounding rivers had similar TA, and hence, we cannot determine which rivers are the main freshwater sources for Station H2. The freshwater endmember calculated in Ohno Strait was significantly higher than TA of its neighbouring large river (Ohta River), suggesting an additional contribution from small rivers directly facing this strait.

Table 1. TA of freshwater sources in each coastal area, with calculated TA of freshwater endmember from the salinity–TA relationship (Fig. 9). Uncertainty terms of freshwater endmember was calculated as standard error of sal = 0 intercept in least squares fitting.

Area	Freshwater endmember of TA from Fig. 9	Freshwater source in each area	TA in each freshwater source
Miyako Bay	158±84 $\mu\text{mol kg}^{-1}$	Hei River	415 $\mu\text{mol kg}^{-1}$ <sup>a</sup>
		Tsugaruishi River	482 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
		Tashiro River	142 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
Kashiwazaki Coast	735±132 $\mu\text{mol kg}^{-1}$	Sabaishi River	845 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
		Ukawa River	655 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
Shizugawa Bay	280±172 $\mu\text{mol kg}^{-1}$	Kitakami River	351 $\mu\text{mol kg}^{-1}$ <sup>a</sup>
		Small rivers in Shizugawa Bay	725–1108 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
Hinase Archipelago	516±169 $\mu\text{mol kg}^{-1}$	Chikusa River	525 $\mu\text{mol kg}^{-1}$ <sup>a</sup>
		Small rivers in the Archipelago	495–640 $\mu\text{mol kg}^{-1}$ <sup>b</sup>
Ohno Strait	497±36 $\mu\text{mol kg}^{-1}$	Ohta River	291 $\mu\text{mol kg}^{-1}$ <sup>a</sup>

a: values referred from Kobayashi (1960)


b: measured in this study

### 3.4 Temporal variation of parameters derived from $\text{pH}_{\text{min}}$ and TA

Based on the regression equation obtained from Fig. 9, we calculated the time series of TA from the time series of salinity at each station (Fig. 10(a)). We then calculated the time series of DIC,  $p\text{CO}_2$ ,  $\Omega_{\text{ar}}$ , and  $\Omega_{\text{ca}}$  from  $\text{pH}_{\text{min}}$  and TA, and here, we show  
395 the results of  $p\text{CO}_2$  and  $\Omega_{\text{ar}}$  in Fig. 10(b) and Fig. 10(c). In this calculation we used time series of  $\text{pH}_{\text{min}}$  and TA estimated from salinity, and uncertainty of these parameters were in the order of  $\pm 0.010$  pH units and  $\pm 20 \mu\text{mol kg}^{-1}$  for  $\text{pH}_{\text{min}}$  and TA, respectively (see Sections 3.2 and 3.3 for details). Uncertainty of derived parameters by using these values were estimated by using CO2sys program as  $\pm 15 \mu\text{atm}$  and  $\pm 0.07$  for  $p\text{CO}_2$  and  $\Omega_{\text{ar}}$ , respectively.

Since TA was calculated as a linear equation of salinity, it was expected to have a resemblance between the TA and  
400 salinity patterns (Fig. 10(a)). Sporadic decreases in TA corresponding to low salinity events in Hinase and Ohno were noticeable. Water with low TA has low buffer capacity, and hence, it has particularly high risks of low pH and high  $p\text{CO}_2$  (e.g., Carstensen and Duarte, 2019; Salisbury, 2008). Figure 10(b) shows an appearance of such risk, extremely high  $p\text{CO}_2$  over  $1,500 \mu\text{atm}$  during the low salinity period in Hinase and Ohno. The annually averaged  $p\text{CO}_2$  values were 404, 450, 393, 589, and  $621 \mu\text{atm}$  for Miyako, Kashiwazaki, Shizugawa, Hinase, and Ohno, respectively, showing that surface waters are in  
405 equilibrium with or even higher than current atmospheric concentration of  $\text{CO}_2$  (416 ppm at 2021; WDCGG, 2023). Generally, estuary areas tend to become sources of atmospheric  $\text{CO}_2$ , while “open” coastal areas such as marginal seas, continental shelves, and large bays tend to become sinks of atmospheric  $\text{CO}_2$  (e.g., Borges et al., 2005; Kubo et al., 2017; Laruelle et al., 2010; Tokoro et al., 2020). Typically, terrestrial input of organic matter from rivers is mostly decomposed in estuaries, causing high  $p\text{CO}_2$  and low pH, as well as a high nutrient flux to the outer estuary. In an “open” coastal area, nutrients transported from the  
410 estuary activate high primary production, which causes low  $p\text{CO}_2$  and high pH in surface waters. The Seto Inland Sea itself is considered a sink of atmospheric  $\text{CO}_2$  (e.g., Tokoro et al., 2020), and hence, it can be treated as an open coastal area. Our results indicate that despite the high distance from the large river mouth, both Hinase Archipelago and Ohno Strait can be classified as “estuaries,” or at least as areas that receive a significant quantity of particulate organic matter from land.

The  $\Omega_{\text{ar}}$  was high in summer and low in winter in Miyako, Kashiwazaki, and Shizugawa (Fig. 10(c)), showing an  
415 opposite pattern to the seasonal variation of  $\text{pH}_{\text{min}}$ . This was due to the **large** seasonal variation in the solubility of aragonite induced by the seasonal change in water temperature (see Section 4.2). In contrast, in Hinase and Ohno, the amplitude of seasonal variation of pH was **large** enough to overcome seasonal variation of aragonite solubility, and as a result,  $\Omega_{\text{ar}}$  changed to show a seasonal maximum in winter and a seasonal minimum in summer. Short-term variation of  $\Omega_{\text{ar}}$  linked to that of salinity also overlapped with seasonal scale variation, and as a result, significantly low  $\Omega_{\text{ar}}$  conditions were frequently observed  
420 in Hinase and Ohno. Fujii et al. (2023) described that during summer–autumn in the Hinase Archipelago,  $\Omega_{\text{ar}}$  occasionally falls below the threshold level of the larvae of the Pacific oyster *Crassostrea gigas*, with the effect of ocean acidification (OA) becoming detectable in the rearing experiment ( $\sim \Omega_{\text{ar}} = 1.5$ ; Waldbusser et al., 2015). This study showed that the situation was almost the same in Ohno Strait (Fig. 10(c)). It should be highlighted that Fujii et al. (2023) also noted that actual damaged



425 **Figure 10: Time series of (a) TA, (b)  $p\text{CO}_2$ , and (c)  $\Omega_{\text{ar}}$  in the five stations. Legends of colour plots are the same for all panels as shown in Fig. 10(a). Solid line in Fig. 10(b) represents the  $p\text{CO}_2$  value in equilibrium with the present atmospheric  $\text{CO}_2$  concentration of 416 ppm. Dashed line in Fig. 10(c) represents the experimentally obtained threshold of the ocean acidification effect for the larvae of Pacific oyster *Crassostrea gigas* ( $\Omega_{\text{ar}} = 1.5$ , Waldbusser et al. [2015]).**



larvae were not detected based on microscopic inspection of Hinase ( $n = 1062$ ). The rearing experiment of Waldbusser et al. (2015) was conducted on the Oregon coast, where *C. gigas* is non-native. Therefore, it is not unrealistic that there is a difference in the tolerance for OA between the local population of Oregon and Seto Inland Sea, the latter being the native habitat of *C. gigas*. Kurihara et al. (2007) examined the effect of low pH in the Seto Inland Sea population of *C. gigas*, and found that the larvae of this population are affected by OA at a pH of 7.4 (National Bureau of Standards [NBS] scale). This threshold approximately corresponds to a pH of 7.27 in the total scale, and in this case, larvae of Pacific oysters are considered to be the same in most seasons both in Hinase and Ohno (Fig. 8(c)). In the experiment by Kurihara (2007), the rearing experiment was occupied only at the control (pH of 8.2 in the NBS scale) and acidified (pH of 7.4 in the NBS scale) conditions, and hence, true threshold for the Seto Inland Sea Pacific oyster population can exist between a pH of 7.27 and 8.07. We thus need further studies, including new rearing experiments, to determine why Pacific oyster larvae are still safe in the present Hinase Archipelago.

A low  $\Omega_{ar}$  level below 1.6 was also detected in Shizugawa Bay (Fujii et al., 2023; see also Table 2 and Fig. 10(c) in this study), and Miyako Bay (Table 2 and Fig. 10(c)), but its duration was only four days and one day, respectively, throughout the study period. In Kashiwazaki,  $\Omega_{ar}$  was above this level throughout the observation period. Other calcifiers are important for fisheries around the Japanese coast, such as Ezo abalone (*Haliotis discus hannai*) and short-spined sea urchin (*Strongylocentrotus intermedius*), have lower  $\Omega_{ar}$  thresholds than that of Pacific oyster, 1.1 (Onitsuka et al., 2018) for *H. discus hannai* and 1.12 (Zhan et al., 2016) for *S. intermedius*. Therefore, Shizugawa Bay, Miyako Bay, and the Kashiwazaki Coast have a low risk of OA, at least from the viewpoint of fisheries. We, however, should note that we have investigated only a few marine species so far, and there may be many unknown species that are vulnerable to low pH/low  $\Omega_{ar}$  condition. We must enhance our knowledge of biological responses according to species, especially for those with low economic importance, such that we can evaluate the total risk posed by OA to the coastal ecosystem.

450

## 4 Discussion

### 4.1 Quantification of short-term variability in properties of water

We focused on short-term variations of pH and related parameters with timescales shorter than one month, which cannot be detected by regular MOE monitoring based on water sampling. To assess this quantitatively, we calculated the SD of each parameter at different timescales (annual, monthly, and 10-days) as follows (see also Appendix A):

$SD_a$	annual standard deviation of each data
$SD_{a<m>}$	annual standard deviation of monthly average
$SD_m$	monthly standard deviation of each data
$avgSD_m^a$	annual average of $SD_m$

460	$\text{maxSD}_m^a$	annual maximum of $\text{SD}_m$
	$\text{SD}_{10}$	moving 10-day standard deviation of each data
	$\text{avgSD}_{10}^a$	annual average of $\text{SD}_{10}$
	$\text{maxSD}_{10}^a$	annual maximum of $\text{SD}_{10}$
	$\text{avgSD}_{10}^m$	monthly average of $\text{SD}_{10}$

465 The results are listed in Table 2. As shown in Figures 7 and 9, both the amplitude and frequency of short-term variation differed significantly among the seasons for almost all parameters. To determine seasonal differences in the extent of short-term variation,  $\text{avgSD}_{10}^m$  in each month was further listed in Table 3.

The  $\text{SD}_{a<m>}$  of water temperature were in the range of 3.9 - 7.2, and this was almost same with that of  $\text{SD}_a$  (Table 2). This implies that large-time scale variation over one month (ca., seasonal scale) was the main component of observed annual temporal variation (Fig. 7(a)). The  $\text{avgSD}_m^a$  of water temperature (0.9–1.5) were < 25% of the  $\text{SD}_a$  (3.9–7.2), and the  $\text{avgSD}_{10}^a$  (0.5–0.8) was approximately half of  $\text{avgSD}_m^a$  in each area (Table 2). In the case of salinity, on the other hand,  $\text{SD}_{a<m>}$  (0.34 - 3.91) was about 70 % of  $\text{SD}_a$  (0.63 - 5.34), indicating that variations with shorter time scale than one month contributes some parts of annual variation in the case of salinity. The  $\text{avgSD}_m^a$  of salinity (0.33–2.37) was approximately half of the  $\text{SD}_a$  (0.66–5.34) in each area, and  $\text{avgSD}_{10}^a$  (0.26–1.40) was approximately two thirds of  $\text{avgSD}_m^a$ . Seasonal variability of  $\text{avgSD}_{10}^m$  was also higher than that of water temperature (Table 3), and as a result,  $\text{avgSD}_{10}^m$  of salinity occasionally exceeded the  $\text{SD}_a$  in several months. Interestingly,  $\text{avgSD}_{10}^a$  and  $\text{avgSD}_m^a$  were almost the same in Miyako, Kashiwazaki, and Shizugawa (Table 2), and the  $\text{maxSD}_{10}^a$  for salinity was higher than the  $\text{maxSD}_m^a$ . Such a result can be attributed to the perturbation of salinity that mainly occurs within the timescale of 10-days, and monthly SD of salinity is essentially determined by the amplitude and frequency of 10-days salinity variation included within that month. Although the  $\text{avgSD}_{10}^a$  of salinity was lower than  $\text{avgSD}_m^a$  in Hinase and Ohno, the  $\text{maxSD}_{10}^a$  was still higher than the  $\text{maxSD}_m^a$ , indicating that the 10-days variation is also an influential component of monthly variation in these two areas.

The relative contribution of short-term variation to annual variability for other parameters varied between the above two extremes (water temperature and salinity). In the cases of DO and pH,  $\text{SD}_{a<m>}$  was on the same level with  $\text{SD}_a$  in Miyako, Kashiwazaki and Shizugawa, but it declined to about 80 % of  $\text{SD}_a$  in Hinase and Ohno. The  $\text{avgSD}_m^a$  was around 25 - 30% of the  $\text{SD}_a$  in Miyako and Kashiwazaki, but it increased to over 50% in Shizugawa, Hinase, and Ohno both for DO and pH. The  $\text{maxSD}_m^a$  exceeded  $\text{SD}_a$  in Hinase and Ohno both for DO and pH. Even the  $\text{maxSD}_{10}^a$  exceeded  $\text{SD}_a$  in Ohno in the case of DO, while it exceeds in Shizugawa, Hinase, and Ohno in the case of pH (Table 2). These results indicate that although the main driver of the annual variation for DO and pH were seasonal variation in Miyako and Kashiwazaki, shorter variation also contributed in Shizugawa, Hinase, and Ohno stations. In the cases of  $\Omega_{ar}$ ,  $\text{SD}_{a<m>}$  showed lower values than  $\text{SD}_a$  in all stations, indicating larger contribution of short-term variation compared to the case of DO and pH. The  $\text{avgSD}_m^a$  was approximately half of  $\text{SD}_a$  for all stations, while the  $\text{avgSD}_{10}^a$  was approximately one third of the  $\text{avgSD}_m^a$ . Both  $\text{maxSD}_m^a$  and  $\text{maxSD}_{10}^a$  exceeded  $\text{SD}_a$  in most areas. These results indicate that short-term variation in the 10-days scale is the significant driver of

Table 2. Statistical aspects of environmental parameters in each station. Annual values are calculated based on the last 1-year data to avoid biases that come from different time length.

Area	Miyako	Kashiwa-zaki	Shizu-gawa	Hinase	Ohno	
495	Water temp. [°C]					
500	SD <sub>a</sub>	4.9	7.2	5.6	6.6	3.9
	SD <sub>a&lt;m&gt;</sub>	4.8	7.0	5.6	6.6	3.9
	avgSD <sub>m</sub> <sup>a</sup>	0.9	1.2	1.3	1.5	1.3
	maxSD <sub>m</sub> <sup>a</sup>	2.1	2.2	2.2	2.7	2.1
	avgSD <sub>10</sub> <sup>a</sup>	0.5	0.6	0.6	0.6	0.8
505	maxSD <sub>10</sub> <sup>a</sup>	1.5	1.9	2.3	1.9	2.1
	Salinity					
	SD <sub>a</sub>	0.66	0.63	0.79	2.44	5.34
	SD <sub>a&lt;m&gt;</sub>	0.34	0.50	0.64	1.79	3.91
510	avgSD <sub>m</sub> <sup>a</sup>	0.33	0.37	0.43	1.28	2.37
	maxSD <sub>m</sub> <sup>a</sup>	<b>1.10</b>	<b>0.64</b>	<b>1.04</b>	<b>5.67</b>	<b>7.47</b>
	avgSD <sub>10</sub> <sup>a</sup>	0.26	0.25	0.37	0.81	1.40
	maxSD <sub>10</sub> <sup>a</sup>	<b>1.31</b>	<b>0.73</b>	<b>1.71</b>	<b>6.00</b>	<b>7.93</b>
515	Oxygen [μmol kg <sup>-1</sup> ]					
	SD <sub>a</sub>	60	51	26	53	44
	SD <sub>a&lt;m&gt;</sub>	60	50	26	40	37
	avgSD <sub>m</sub> <sup>a</sup>	17	13	13	36	35
	maxSD <sub>m</sub> <sup>a</sup>	28	20	22	<b>62</b>	<b>57</b>
520	avgSD <sub>10</sub> <sup>a</sup>	9	7	10	11	22
	maxSD <sub>10</sub> <sup>a</sup>	22	17	21	40	<b>48</b>
	pH <sub>min</sub>					
	SD <sub>a</sub>	0.05	0.09	0.09	0.21	0.16
525	SD <sub>a&lt;m&gt;</sub>	0.04	0.09	0.08	0.16	0.11
	avgSD <sub>m</sub> <sup>a</sup>	0.02	0.03	0.05	0.10	0.09
	maxSD <sub>m</sub> <sup>a</sup>	<b>0.06</b>	0.05	<b>0.09</b>	<b>0.23</b>	<b>0.24</b>
	avgSD <sub>10</sub> <sup>a</sup>	0.02	0.02	0.03	0.06	0.07
	maxSD <sub>10</sub> <sup>a</sup>	<b>0.06</b>	0.04	<b>0.10</b>	<b>0.26</b>	<b>0.31</b>
530	Ω <sub>ar</sub>					
	SD <sub>a</sub>	0.25	0.26	0.29	0.58	0.52
	SD <sub>a&lt;m&gt;</sub>	0.24	0.23	0.16	0.41	0.33
	avgSD <sub>m</sub> <sup>a</sup>	0.12	0.14	0.23	0.31	0.31
535	maxSD <sub>m</sub> <sup>a</sup>	0.21	0.22	<b>0.48</b>	<b>0.89</b>	<b>0.78</b>
	avgSD <sub>10</sub> <sup>a</sup>	0.09	0.11	0.15	0.22	0.25
	maxSD <sub>10</sub> <sup>a</sup>	<b>0.25</b>	0.21	<b>0.63</b>	<b>1.27</b>	<b>0.81</b>

Numbers are shown in bold when they exceed SD<sub>a</sub>.

Table 3. Monthly average of 10-day standard variation (avgSD<sub>10<sup>th</sup></sub>) for each parameter on each month in each station. Numbers shown by bold represent annual maximum and the second maximum

Area	Parameter	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sept.	Oct.	Nov.	Dec.	
Miyako	Water Temp. [°C]	0.26	0.17	0.15	0.28	0.64	<b>0.72</b>	<b>0.86</b>	0.58	0.70	0.53	0.35	0.45	
	Salinity	0.03	0.12	0.28	0.17	0.15	0.10	0.32	<b>0.71</b>	<b>0.78</b>	0.29	0.08	0.05	
	oxygen [ $\mu\text{mol kg}^{-1}$ ]	8.9	13.1	10.8	<b>14.4</b>	<b>13.5</b>	8.4	9.5	6.0	7.3	5.2	3.3	5.6	
	pH	0.01	0.02	<b>0.03</b>	<b>0.03</b>	<b>0.03</b>	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
	$\Omega_{\text{ar}}$	0.03	0.08	0.11	<b>0.14</b>	0.12	<b>0.14</b>	0.08	0.13	0.13	0.07	0.04	0.04	
Kashiwazaki	Water Temp. [°C]	0.70	<b>0.76</b>	0.34	0.34	0.51	0.55	0.61	0.57	0.58	0.74	0.59	<b>1.08</b>	
	Salinity	0.25	<b>0.41</b>	0.20	0.20	<b>0.38</b>	0.15	0.37	0.20	0.13	0.21	0.37	0.30	
	oxygen [ $\mu\text{mol kg}^{-1}$ ]	9.9	<b>11.7</b>	5.0	4.2	6.1	6.9	5.3	3.8	4.1	6.0	5.8	<b>10.8</b>	
	pH	0.02	0.02	0.02	0.02	0.02	<b>0.03</b>	0.02	0.02	0.02	0.02	0.02	<b>0.03</b>	
	$\Omega_{\text{ar}}$	0.08	<b>0.13</b>	0.09	0.08	0.10	0.08	<b>0.13</b>	0.12	0.10	0.12	0.11	<b>0.15</b>	
Shizugawa	Water Temp. [°C]	0.30	0.32	0.23	0.61	0.94	<b>1.15</b>	0.68	<b>1.13</b>	0.61	0.46	0.44	0.40	
	Salinity	0.04	0.13	0.32	0.58	0.56	0.32	<b>0.77</b>	0.61	<b>0.66</b>	0.23	0.18	0.09	
	oxygen [ $\mu\text{mol kg}^{-1}$ ]	--	--	--	--	<b>16.8</b>	10.1	9.0	<b>13.3</b>	7.4	7.0	5.6	9.1	
	pH	0.02	<b>0.04</b>	0.03	<b>0.04</b>	0.03	0.02	0.03	<b>0.05</b>	0.03	0.02	0.02	0.01	
	$\Omega_{\text{ar}}$	0.10	0.18	0.17	0.18	0.15	0.14	0.18	<b>0.27</b>	0.16	<b>0.20</b>	0.13	0.06	
Hinase	Water Temp. [°C]	<b>0.88</b>	--	0.51	0.41	0.60	0.34	0.71	<b>1.00</b>	0.49	0.71	0.65	0.72	
	Salinity	0.05	--	0.19	0.20	0.61	1.33	<b>2.41</b>	<b>2.17</b>	1.16	0.40	0.28	0.10	
	oxygen [ $\mu\text{mol kg}^{-1}$ ]	--	--	--	--	--	--	--	--	16.4	13.1	8.2	7.5	
	pH	0.01	--	0.06	--	0.02	0.09	<b>0.12</b>	<b>0.15</b>	0.08	0.04	0.03	0.02	
	$\Omega_{\text{ar}}$	0.07	--	0.19	--	0.11	<b>0.38</b>	<b>0.58</b>	0.34	0.20	0.14	0.10	0.09	
Ohno	Water Temp. [°C]	--	--	--	--	--	<b>1.26</b>	0.90	<b>1.13</b>	0.61	0.54	0.49	0.44	
	Salinity	--	--	--	--	--	0.68	<b>2.75</b>	<b>3.60</b>	2.03	0.44	0.22	0.10	
	oxygen [ $\mu\text{mol kg}^{-1}$ ]	--	--	--	--	--	17.9	29.6	<b>31.0</b>	29.3	<b>29.7</b>	9.5	4.6	
	pH	--	--	--	--	--	0.04	0.09	<b>0.14</b>	<b>0.10</b>	0.07	0.02	0.01	
	$\Omega_{\text{ar}}$	--	--	--	--	--	0.26	<b>0.46</b>	<b>0.47</b>	0.27	0.18	0.07	0.04	
Tokyo Bay*	Water Temp. [°C]	0.36	0.31	0.55	0.45	0.56	0.76	<b>0.84</b>	<b>0.96</b>	0.82	0.57	0.28	0.51	
	Salinity	0.34	0.35	0.79	0.72	0.67	1.35	<b>2.11</b>	<b>1.65</b>	1.41	0.86	0.53	0.46	
	pH	0.02	0.04	0.05	0.07	0.08	<b>0.15</b>	<b>0.15</b>	0.13	0.13	0.08	0.04	0.02	

\*See Section 4.3 for the detail of the data in Tokyo Bay.

annual temporal variation for  $\Omega_{ar}$ , which was similar to the case of salinity.

We further investigated the seasonal dependency of the 10-days SD in each area based on Table 3. The  $\max\text{SD}_{10^a}$  for DO,  $\text{pH}_{\min}$ , and  $\Omega_{ar}$  occurred at approximately the same time, and moreover, seasonal distribution of  $\text{avgSD}_{10^a}$  showed positive correlation with statistical significance ( $r^2 > 0.5$ ) among these three parameters. These results indicate that the 10-days variations of these three parameters were caused by the same processes (ca. biological activities).

However, the specific timing of the occurrence of the annual maximum for these three parameters differed among the areas. In Miyako, the annual maximum of  $\text{avgSD}_{10^m}$  for DO,  $\text{pH}_{\min}$ , and  $\Omega_{ar}$  occurred in spring, when biological activities are the highest. In Shizugawa, Hinase, and Ohno, the maximum  $\text{avgSD}_{10^m}$  for these parameters occurred during late summer and early autumn, which approximately overlapped with the timing of the occurrence of the annual maximum of  $\text{avgSD}_{10^m}$  of salinity (Table 3). In Kashiwazaki, the maximum  $\text{avgSD}_{10^m}$  for DO,  $\text{pH}_{\min}$ , and  $\Omega_{ar}$  occurred in winter, when strong perturbation of river flow induced by heavy snowfall caused an annual maximum  $\text{avgSD}_{10^m}$  for both salinity and water temperature. Overall, the investigated statistical aspects of short-term variations indicated that several physical processes related to the 10-days salinity variation had derivatively induced biological processes that caused short-term variations in DO,  $\text{pH}_{\min}$ , and  $\Omega_{ar}$  in these coastal areas.

Temporal variation of pH and the related parameters are mainly analysed with interannual, seasonal and diurnal time scales (e.g., Bauman and Smith 2018; Carstensen et al., 2019; Hassoun et al., 2017; Lowe et al., 2018; Provoost et al., 2010; Rosenau et al., 2021), but many studies have also highlighted existence of significant day-to-day scale pH variation in coastal areas, especially in estuaries (e.g., Bednarzek et al., 2022; Frieder et al., 2012; Fujii et al., 2021; Hoffman et al., 2012; Johnson et al., 2013). The present results illustrated that such day-to-day variation with the timescales from 10 to 30 days plays significant contribution to the annual variation of pH and the related parameters in the coastal areas in Japan. Indeed, amplitude of  $\text{SD}_a$  and  $\text{avgSD}_m^a$  of  $\text{pH}_{\min}$  observed in Hinase and Ohno was almost similar with those observed in US estuaries by former study (Bednarzek et al., 2022; Hoffman et al., 2011), indicating that such high contribution of short-term variation to total variability is common amongst the world's estuaries. In several cites off California coasts, it was reported that eventual coastal upwelling brings carbon and nutrient to the coast and cause multi-days scale pH variation (e.g., Frieder et al., 2012; Kessouri et al., 2021). In Japan coasts, on the other hand, the observed short-term variations in pH seemed to be connected strongly with that of salinity, suggesting that the change in riverine input plays significant rolls in the short-term pH variation.

#### 4.2 Classification of temporal variations into thermodynamic and non-thermodynamic components

To determine the processes that contribute to the short-term variation in biogeochemical properties, we divided the observed temporal variations of each parameter into thermodynamic and non-thermodynamic components using the following equation:

$$C_i = C_i(\text{eq}) + C_i(\text{diseq}) \quad (1)$$

575


where  $C_i$  represents the observed concentration of parameter  $i$ , while  $C_i(\text{eq})$  represents the estimated concentration of parameter  $i$  in equilibrium with the current atmosphere under the observed water temperature and salinity.  $C_i(\text{diseq})$  represents the difference between  $C_i$  and  $C_i(\text{eq})$ . This is the similar idea of AOU, while  $\text{DO}(\text{diseq})$  is equivalent to AOU multiplied by -1. For DO,  $\text{DO}(\text{eq})$  was calculated from water temperature and salinity based on the formulation of Weiss (1970) with a fixed atmospheric pressure of 1 atm. Both  $\text{pH}(\text{eq})$  and  $\Omega_{\text{ar}}(\text{eq})$  were calculated using the CO2sys program with the same set of constants as described in Section 2.1, using water temperature, salinity, estimated TA, and a fixed atmospheric  $\text{CO}_2$  mole fraction of 415 ppm.

Figure 11 shows the temporal variation of  $\text{DO}(\text{eq})$ ,  $\text{pH}(\text{eq})$ , and  $\Omega_{\text{ar}}(\text{eq})$ , while Fig. 12 shows  $\text{DO}(\text{diseq})$ ,  $\text{pH}_{\text{min}}(\text{diseq})$ , and  $\Omega_{\text{ar}}(\text{diseq})$ . We also calculated 1-year SD, monthly SD, and 10-days SD for these parameters, and the results are listed in Table 4. The annual temporal variation was approximately the same between  $\text{DO}(\text{eq})$  and  $\text{DO}(\text{diseq})$  (Table 4), but their origin was significantly different. While seasonal variation played a dominant role in the temporal variation of  $\text{DO}(\text{eq})$  (Fig. 11(a), Table 4), short-term variations played a significant role in that of  $\text{DO}(\text{diseq})$  (Fig. 12(a), Table 4). In the case of pH, the temporal variation of  $\text{pH}(\text{eq})$  resembles that of salinity (Figs. 7(b) and 11(b)), and the amplitude of  $\text{pH}(\text{eq})$  ( $\sim 0.25$ , Fig. 11(b)) was approximately one order smaller than that of  $\text{pH}(\text{diseq})$  ( $\sim 1.2$ , Fig. 12(b)). This result indicated that the thermodynamic process had a negligible contribution to the temporal variation of pH. Interestingly, the relative contributions of monthly SD and 10-days SD of  $\text{pH}(\text{diseq})$  to the 1-year SD showed a similar structure to that of  $\text{pH}_{\text{min}}$  (Table 4), suggesting that the short-term variation at the scale of 10-days was the main driver of annual temporal variation for  $\text{pH}_{\text{min}}(\text{diseq})$ , which is similar to the cases of  $\text{pH}_{\text{min}}$  and salinity. Temporal variation of  $\Omega_{\text{ar}}(\text{eq})$  had both seasonal- and short-term scale components (Table 4), and the pattern of seasonal variation resembled that of water temperature, while that of short-term variation resembled that of salinity (Fig. 11(c)).  $\Omega_{\text{ar}}(\text{diseq})$  showed a similar scale of 1-year variability to that of  $\Omega_{\text{ar}}(\text{eq})$  (Fig. 11(c) and Fig. 12(c), Table 4), similar to the case of DO. The temporal pattern of  $\Omega_{\text{ar}}(\text{diseq})$  was quite similar to that of  $\text{pH}_{\text{min}}(\text{diseq})$ , suggesting that short-term variation at the 10-days scale contributes significantly to the annual temporal variation of  $\Omega_{\text{ar}}(\text{diseq})$ , similar to the case of  $\text{pH}_{\text{min}}(\text{diseq})$ . Overall, the non-thermodynamic component contributed approximately half of the observed annual variation in DO and  $\Omega_{\text{ar}}$ , and in the case of pH, the non-thermodynamic component played a dominant role in the annual variation. For all these biogeochemical parameters, the short-term variation at the 10-days scale contributed a significant percentage of the temporal variation of the non-thermodynamic component.

To determine the specific process that drives the non-thermodynamic temporal variation of these properties, we examined the correlation between  $\text{pH}_{\text{min}}(\text{diseq})$  and  $\text{DO}(\text{diseq})$  (Fig. 13(a)). These two properties showed significant positive correlation, indicating that the non-thermodynamic component of temporal variations was driven mainly by biological processes. Several studies have also detected a positive relationship between variations in oxygen saturation and the metabolic component of pH in coastal waters, suggesting the primary influence of biological processes on pH dynamics (e.g., Baumann

605





655

**Figure 11:** Time series of (a) DO(eq), (b) pH(eq), and (c)  $\Omega_{ar}$ (eq) in the five stations. Legends of colour plots are the same for all panels as shown in Fig. 11(b).







Figure 12: Time series of (a)  $\text{DO}(\text{diseq})$ , (b)  $\text{pH}_{\min}(\text{diseq})$ , and (c)  $\Omega_{\text{ar}}(\text{diseq})$  in the five stations. Legends of colour plots are the same for all panels as shown in Fig. 12(a).

660



**Figure 13:** Plot of (a)  $\text{pH}_{\min}(\text{diseq})$  and (b)  $\text{DIC}_{\max}(\text{diseq})$  against  $\text{DO}(\text{diseq})$ , respectively. The purple dashed line in Fig. 13(b) represents the theoretical line when assuming that biological production / degradation of organic matter occur with the Redfield relationship ( $-\Delta \text{pH}_{\min}(\text{diseq}) / \Delta \text{DO}(\text{diseq}) = 106/138$ ). The red dashed line represents the regression line of Hinase data.

665 and Smith, 2018; Lowe et al., 2019). We further investigated the correlation between  $\text{DIC}_{\text{max}}(\text{diseq})$  and  $\text{DO}(\text{diseq})$  (Fig. 13(b));  
the former was calculated as  $\text{pH}_{\text{min}}(\text{diseq})$  and  $\Omega_{\text{ar}}(\text{diseq})$ . Most data obtained in Miyako and Ohno and approximately 50%  
of data obtained in Shizugawa approximately followed a linear regression line with the slope of  $-\Delta\text{DIC}_{\text{max}}(\text{diseq})/\Delta\text{DO}(\text{diseq})$   
= 0.77 (~106/138) crossing the origin, suggesting that the observed temporal variations of both  $\text{DIC}_{\text{max}}(\text{diseq})$  and  $\text{DO}(\text{diseq})$   
were induced by the production/decomposition of planktonic oceanic particles with C : O<sub>2</sub> ratio of 106 : 138 (Redfield et al.,  
670 1963) in these two areas. Such Redfield-type relationship between short-term variation of DO and that of DIC had also been  
observed in the coastal area of East China Sea (Guo et al., 2021). The other data obtained in Shizugawa, as well as most data  
obtained in Hinase and Kashiwazaki followed the regression lines with steeper slopes than that of open ocean stoichiometry.  
The respiratory quotients of estuarine ecosystems can occasionally be as high as 1.5 (Wang et al., 2018) or even 2.0 (Giblin et  
al., 1997), when affected by the anaerobic respiration process in estuarine sediments. The observed high -  
675  $\Delta\text{DIC}_{\text{max}}(\text{diseq})/\Delta\text{DO}(\text{diseq})$  ratio thus suggests that lateral propagation of biological processes in the neighbouring estuaries  
contributes significantly to the observed temporal variations of  $\text{DIC}_{\text{max}}(\text{diseq})$  and  $\text{DO}(\text{diseq})$  in these three areas. On the other  
hand, turnover time of oxygen in ocean mixed layer due to air-sea gas exchange (in the range from several days to two weeks;  
Izett et al., 2018; Qin et al., 2021) is about ten-times shorter than that of CO<sub>2</sub> (in the order of several months; e.g., Jones et al.,  
2014), reflecting their abundance in seawater. The  $\Delta\text{DO}(\text{diseq})$  that had once caused by biological processes, therefore, can  
680 recover to zero with shorter time scale than that of  $\Delta\text{DIC}_{\text{max}}(\text{diseq})$  through the air-sea gas exchange process. The observed  
difference in  $-\Delta\text{DIC}_{\text{max}}(\text{diseq})/\Delta\text{DO}(\text{diseq})$  ratio between Hinase and Ohno can also be explained by this mechanism, if large  
gas exchange process (ca., high wind speed and/or high wave height) have occurred in Hinase while it had not occurred in Ohno.  
However, monthly-averaged wind speed was always higher in Ohno than in Hinase in 2020 autumn (Japan Meteorological  
Agency weather record data base, <https://www.data.jma.go.jp/stats/etrn/index.php>), when low  $\Delta\text{DO}(\text{diseq})$  and high  
685  $\Delta\text{DIC}_{\text{max}}(\text{diseq})$  were mainly observed in this study (Figure 12). Similarly, there is no significant difference in wind speed  
between two groups of Shizugawa data (one following Hinase data and the another following Ohno data). We thus conclude  
that relatively faster recovery of  $\Delta\text{DO}(\text{diseq})$  compared to that of  $\Delta\text{DIC}_{\text{max}}(\text{diseq})$  through air-sea gas exchange is not the main  
cause of the observed high  $-\Delta\text{DIC}_{\text{max}}(\text{diseq})/\Delta\text{DO}(\text{diseq})$  ratio, although this process may partly contribute to create high ratio  
in these areas.

690

### 4.3 Controlling factor of the amplitude of short-term pH variation.

Our analyses in Sections 3.2 and 3.4 clearly shows that severe low pH/low  $\Omega_{\text{ar}}$  situations occur only at the short-term pH  
drawdown events that coincide with rainfall events in the coastal areas of Japan (Fig. 9(c) and 10(c)). As the amplitude of pH  
variation related to rainfall events differed among the five coastal areas, it is important to understand the controlling factor that  
695 determines the amplitude of short-term pH variation. To investigate this aspect, we analysed the relationship of statistical short-




Figure 14: Plots  $\text{avgSD}_{10}^m(\text{pH})$  against  $\text{avgSD}_{10}^m(\text{sal})$  for (a) Tokyo Bay, (b) Hinase and Ohno, and (c) Miyako, Kashiwazaki, and Shizugawa, respectively. The red dashed line represents regression line of Tokyo Bay, while the black dashed lines represent that of (b) Hinase and Ohno, and (c) Miyako, Kashiwazaki, and Shizugawa, respectively. Green and blue dashed lines represent regression lines of Hinase and Ohno, and Miyako, Kashiwazaki, and Shizugawa, respectively, with the data of months with the  $\text{avgSD}_{10}^m(\text{sal})$  higher than 1.0.

700




Figure 15: Plot of  $\Delta\text{avgSD}_{10}^m(\text{pH}) / \Delta\text{avgSD}_{10}^m(\text{sal})$  against  $\overline{\text{NO}_3}$  for each coastal area category. The blue and red dots represent slope calculated from all months from the months with  $\text{avgSD}_{10}^m(\text{sal})$  higher than 1.0, respectively. The dashed lines represent the regression lines.

705

term variabilities between salinity and  $\text{pH}_{\min}$ . Since short-term variation mainly occurred in the timescale of  $< 10$  days for both salinity and  $\text{pH}_{\min}$  (see discussion in Section 4.1), we plotted the  $\text{avgSD}_{10^m}$  of  $\text{pH}_{\min}$  against that of salinity (Fig. 14). In this analysis, we introduced continuous monitoring data of pH and salinity obtained in Tokyo Bay (off Kawasaki Artificial Island, <https://www.tbeic.go.jp/MonitoringPost/manual/aboutObservedPoint02.pdf>) to obtain information on highly eutrophic coastal areas. The details of the observation methods and the location settings of this station are described in the Appendix B.

In Tokyo Bay,  $\text{avgSD}_{10^m}$  of  $\text{pH}_{\min}$  (hereafter  $\text{avgSD}_{10^m}(\text{pH})$ ) linearly increased with an increase in  $\text{avgSD}_{10^m}$  of salinity ( $\text{avgSD}_{10^m}(\text{sal})$ , Fig. 14(a)), suggesting that freshwater transports the sources of biological processes (both organic carbon and dissolved nutrients) at a constant concentration regardless of the flow rate. On the other hand, in Hinase and Ohno,  $\text{avgSD}_{10^m}(\text{pH})$  increased with that of  $\text{avgSD}_{10^m}(\text{sal})$  at the same rate as observed in Tokyo Bay when  $\text{avgSD}_{10^m}(\text{sal})$  was low but  $\Delta\text{avgSD}_{10^m}(\text{pH}) / \Delta\text{avgSD}_{10^m}(\text{sal})$  changed to a low value when  $\text{avgSD}_{10^m}(\text{sal})$  exceeded 1.0 (Fig. 14(b)). Generally, suspended particle export of the river increases linearly with that of river discharge, while percentage of particle organic materials (POM) within suspended particles decreases along with the increase of suspended particle export (e.g., Bukaveckas 2022; Coynel et al., 2005; Point et al., 2007; Zhang et al., 2013). As this result, increasing rate of the POM transport with that of river discharge tends to become weak when river discharge becomes high enough (Coynel et al., 2005; Kim et al., 2020). If we assume that short term pH variation is mainly caused by the degradation of POM discharged from rivers into coastal areas and subsequential biological production by using nutrients released from the degraded POM (e.g., Kubo et al., 2017; Salisbury et al., 2008; Tokoro et al., 2020), we can expect that  $\Delta\text{avgSD}_{10^m}(\text{pH}) / \Delta\text{avgSD}_{10^m}(\text{sal})$  will also become small at the months with high river discharge (i.e., high  $\text{avgSD}_{10^m}(\text{sal})$ ). The observed result in this study also indicated that the concentration of biologically active materials transported by freshwater into the Seto Inland Sea was diluted at a high freshwater flow rate, while it was not diluted in the rivers flowing into Tokyo Bay, as the latter receives far higher anthropogenic nutrients' loadings from its drainage basin than those received by the Seto Inland Sea. In the coastal area receiving further low anthropogenic nutrients' loadings such as Miyako, Kashiwazaki, and Shizugawa,  $\Delta\text{avgSD}_{10^m}(\text{pH}) / \Delta\text{avgSD}_{10^m}(\text{sal})$  was much lower than that of Hinase and Ohno when  $\overline{\text{SD}}_{10}(\text{sal})$  exceeded 1.0 (Fig. 14(c)).

We determined the nutrients' concentrations of the main freshwater sources for each of the three coastal area categories (i.e., Tashiro, Sabaishi, and Kitakami Rivers for Miyako + Kashiwazaki + Shizugawa, Chikusa and Ohta Rivers for Hinase + Ohno, Ara, Tama, and Tsurumi Rivers for Tokyo Bay, respectively) from the MOE Public Water Quality Database, and calculated the weighted mean nitrate concentration ( $\overline{\text{NO}_3}$ ) of river waters that flow into each coastal area using water transport as the weight. We then found that  $\Delta\text{avgSD}_{10^m}(\text{pH}) / \Delta\text{avgSD}_{10^m}(\text{sal})$  observed at high freshwater input (that is,  $\text{avgSD}_{10^m}(\text{sal}) > 1.0$ ) showed a linear relationship with  $\overline{\text{NO}_3}$  in each coastal area category (Fig. 15). Even if we compare  $\Delta\text{avgSD}_{10^m}(\text{pH}) / \Delta\text{avgSD}_{10^m}(\text{sal})$  for all months, positive correlation against  $\overline{\text{NO}_3}$  was still significant. These results imply that the amplitude of short-term variation of  $\text{pH}_{\min}$  in coastal areas is principally determined by the quantity of nutrients transported by freshwater from the hinterland. We, however, need to note that the riverine nitrate concentration can be an implicit function of other controlling factors. For example, we recently observed that the quantity of suspended organic

particles in river waters, as well as accumulated organic materials in riverine and estuarine sediments, are linearly correlated with the riverine nitrate concentration (unpublished data). Hence, the  $\overline{\text{NO}_3}$  in Fig. 15 may actually be an indicator of the transport of these materials. We thus need further detailed observations, especially at the time of increase in water level of the river, to understand practical processes that cause short-term pH variation in coastal waters.

745

## 5 Conclusion

In this study, we synthesised data from continuous pH monitoring of five coastal areas in Japan from 2020 to 2021. Annual variability ( $\sim 1$  SD) of pH and  $\Omega_{\text{ar}}$  were 0.05–0.09 pH units and 0.25–0.29, respectively, for three areas with low anthropogenic nutrients' loadings (Miyako Bay, Kashiwazaki Coast, and Shizugawa Bay), while they increased to 0.16–0.21 pH units and 0.52–0.58, respectively, in two areas with medium anthropogenic nutrients' loadings (Hinase Archipelago and Ohno Strait in Seto Inland Sea). Statistical assessment of temporal variability at various timescales revealed that most of the annual variabilities in both pH and  $\Omega_{\text{ar}}$  were controlled by short-term variation with a timescale of  $< 10$  days, rather than the seasonal-scale variation. Our analyses further illustrated that most of the short-term pH variation (and hence, annual variation) was caused by biological processes, while both thermodynamic and biological processes equally contributed to the temporal variation in  $\Omega_{\text{ar}}$ . Biological alteration of pH mainly occurred in oceanic areas in Miyako Bay and Ohno Strait, whereas it occurred mainly in estuarine areas in Kashiwazaki Coast and Hinase Archipelago. In Shizugawa Bay, both oceanic and estuarine areas were responsible for the biological alteration of pH.


The observed results show that short-term acidification with  $\Omega_{\text{ar}}$  of  $< 1.5$  occurred occasionally in Miyako and Shizugawa Bays, while it occurred frequently in the Hinase Archipelago and Ohno Strait. Many such short-term acidification events were related to short-term low-salinity events. Our analyses showed that the amplitude of short-term pH variation was linearly correlated with that of short-term salinity variation, and its regression coefficient at the time of high freshwater input was positively correlated with the nutrient concentration of the main river that flows into the coastal area.

Fortunately, no marine organism that has been damaged by ocean acidification has been detected in coastal areas of Japan (Fujii et al., 2023). However, our study showed that  $\Omega_{\text{ar}}$  in Japanese coastal areas occasionally drops to a level that is potentially hazardous for marine organisms, such as Pacific oysters ( $< 1.5$ , Waldbusser et al., 2015) even in the present state. It is already known that the pH in Japanese coastal areas is decreasing at the same rate as that of the open ocean (Ishida et al., 2021; Ishizu et al., 2019), and hence, the extent, duration, and frequency of such short-term low pH and  $\Omega_{\text{ar}}$  situations will increase in coastal areas in Japan in the future (Fujii et al., 2023). Our study indicates that the amplitude of the short-term drawdown of pH related to low-salinity events will decrease if we can reduce the nutrient concentration of rivers. Similar effect of anthropogenic nutrient reduction to diminish short-term  $\Omega_{\text{ar}}$  drawdown in coastal waters are also reported by Kessouri et al. (2021). We should note, however, that a certain percentage of Japanese coastal areas are now suffering due to the problem of

770

low biological productivity derived from the decreased anthropogenic nutrient input (e.g., Yamamoto et al., 2021). We must consider the balance between the risk of ocean acidification and oligotrophication when we control anthropogenic loadings to coastal waters in the future. We should also note that not only the concentration of dissolved nutrients, but also other  
775 biologically active materials such as suspended organic materials in river waters and accumulated organic materials in riverine and estuarine sediments may be responsible for short-term pH variations at low salinity events (e.g., Carstensen and Duarte, 2019; Salisbury et al., 2008). Our analysis revealed that seawaters in both the Hinase Archipelago and Ohno Strait were oversaturated with CO<sub>2</sub> all through the years, and hence, it is considered that more organic matter than that biologically produced within these areas was decomposed. In such cases, not only the reduction of dissolved nutrients but also the reduction  
780 of particulate organic matter transported from rivers to coastal areas will contribute to the suppression of short-term pH drawdown. Many studies have already mentioned the contribution of seaweed/seagrass beds to the effective capture of suspended organic sediments in estuaries (e.g., Barcelona et al., 2021; Levia-Duenas et al., 2023; Potouroglou et al., 2017), and hence, the conservation and/or development of seaweed/seagrass beds in estuarine and coastal areas will contribute to the reduction of organic matter transport from rivers to coastal areas at times of high river flow. To specify effective measures  
785 against the current coastal acidification in Japan, further detailed observations, especially at the time of increase in water level of the river, are needed such that we can obtain a detailed understanding of practical processes that cause short-term pH variation in coastal waters.

## 6 Appendix A: Schematic explanation of $SD_a$ , $SD_{a<m>}$ , $SD_m$ , and $SD_{10}$



## 790 7 Appendix B: Observation methods and location settings of Tokyo Bay data

Detailed data for Tokyo Bay (Kawasaki Artificial Island, Section 4.3), referred to by us, are described in the home page of data holders: <https://www.tbeic.go.jp/MonitoringPost/manual/aboutObservedPoint02.pdf>. Here, we provide a brief summary.

795 Kawasaki Artificial Island (KAI) is located in the centre of Tokyo Bay, the bay with the highest population in its basin area (2.9 million) and highest nutrient loadings ( $210 \text{ tN d}^{-1}$ ) in Japan (Fig. A1). KAI is a huge concrete cylinder with a diameter of 200 m, in which an air vending system for the trans-bay freeway tunnel is stored. An autonomous ocean profiling system



was set off KAI, which included a YSI 6600V2-4M multiple ocean sensor for the measurement of water temperature, salinity, and pH. The mechanical measurement resolution of pH was  $\pm 0.01$  pH units. The sensors were cleaned and calibrated on a monthly basis. Although the pH sensor was calibrated against NBS buffers, we used pH data without conversion to a total scale, as we used these pH data only for the analysis of temporal variability. Vertical profiles were measured hourly, and the obtained data were distributed from the public database managed by the Tokyo Bay Environmental Information Center (<https://www.tbeic.go.jp/MonitoringPost/ViewGraph/ViewGraph?buoyId=02>). We downloaded KAI data pertaining to July 2020 to December 2021 from this database and used them for analysis.

## 8 Data Availability

Data from Tokyo Bay can be downloaded from the Tokyo Bay Environmental Information Center Public Database (<https://www.tbeic.go.jp/MonitoringPost/ViewGraph/ViewGraph?buoyId=02>). Other data used in this study are provided in the Supplementary Materials.

## 9 Author contribution

TO and DM performed field measurements and water sampling in Miyako Bay. MH and MY performed field measurements and water sampling on the Kashiwazaki Coast. AD performed field measurements and water sampling in Shizugawa Bay. SO, TT, MF, and RH performed field measurements and water samplings in the Hinase Archipelago. TO, GO, and KK performed field measurements and water sampling in Ohno Strait. TO undertook the measurements of discrete carbonate samples taken in Miyako Bay and Kashiwazaki Coast, while MW undertook the measurements of those taken in Shizugawa Bay, Hinase Archipelago, and Ohno Strait. Analyses of data obtained by these efforts were performed by TO.

## 10 Acknowledgements

We thank all fishers who helped us with their boats and offered their support for monitoring in Shizugawa Bay and the Hinase Archipelago. We also thank Dr. Yukihiro Nojiri, Dr. Hideaki Nakata, Dr. Osamu Matsuda, Dr. Tetsuo Yanagi, and the reviewers of this paper for their fruitful advice and discussions during our analyses. This study was funded both by the Ocean Acidification Adaptation Project (OAAP) of the Nippon Foundation and the Study of Biological Effects of Acidification and Hypoxia (BEACH) of the Environment Research and Technology Development Fund Grant Number JPMEERF20202007 of the Environmental Restoration and Conservation Agency of Japan.

## 11 References

- Abo, K. and Onitsuka, G.: Characteristics of currents induced by heavy rainfall in Hiroshima Bay in July 2018, Proceedings of the Japan Society of Civil Engineers, Ser. B, 75, 1051-1056, in Japanese, 2019.
- 825 Barcelona, A., Oldham, C., Colomer, J., Garcia-Orellana, J., and Serra, T.: Particle capture by seagrass canopies under an oscillatory flow, *Coastal Engineering*, 169, 103972, <https://doi.org/10.1016/j.coastaleng.2021.103972>, 2021.
- Barton, A., Hales, B., Waldbusser, G. G., Langdon, C. and Feely, R. A.: The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects, *Limnology and Oceanography*, 57, 698-710, <https://doi.org/10.4319/lo.2012.57.3.0698>, 2012.
- 830 Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J. and Santana-Casiano, J. M.: A time-series view of changing surface ocean chemistry due to ocean uptake of anthropogenic CO<sub>2</sub> and ocean acidification, *Oceanography*, 27, 126-141, <https://doi.org/10.5670/oceanog.2014.16>, 2014.
- Baumann, H. and Smith, E. M.: Quantifying metabolically driven pH and oxygen fluctuations in US nearshore habitats at diel to interannual timescales, *Estuaries and Coasts*, 41, 1102-1117, <https://doi.org/10.1007/s12237-017-0321-3>, 2018.
- 835 Bednaršek, N., Beck, M. W., Pelletier, G., Applebaum, S. L., Feely, R. A., Butler, R., Byrne, M., Peabody, B., Davis, J. and Štrus, J.: Natural analogues in pH variability and predictability across the Coastal Pacific estuaries: extrapolation of the increased oyster dissolution under increased pH amplitude and low predictability related to ocean acidification, *Environmental Science and Technology*, 56, 9015-9028, <https://doi.org/10.1021/acs.est.2c00010>, 2022.
- Bernardo, L. P. C., Fujii, M., and Ono, T.: Development of a high-resolution marine ecosystem model for predicting the combined impacts of ocean acidification and hypoxia, *Frontiers in Marine Science*, 10, 1174892, <https://doi.org/10.3389/fmars.2023.1174892>, 2023.
- Booth, J. A. T., McPhee-Shaw, E. E., Chua, P., Kingsley, E., Denny, M., Phillips, R., Bograd, S. J., Zeidberg, L. D. and Gilly, W. F.: Natural intrusions of hypoxic, low pH water into nearshore marine environments on the California coast, *Continental Shelf Research*, 45, 108-115, <https://doi.org/10.1016/j.csr.2012.06.009>, 2012.
- 845 Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO<sub>2</sub> in the coastal ocean: Diversity of ecosystems counts, *Geophysical Research Letters*, 32, L14601, <https://doi.org/10.1029/2005GL023053>, 2005.
- Borges, A. V. and Gypens, N.: Carbonate chemistry in the coastal zone responds more strongly to eutrophication than to ocean acidification, *Limnology and Oceanography*, 55, 346-353, 2010.
- Bukaveckas, P. A.: Carbon dynamics at the river–estuarine transition: a comparison among tributaries of Chesapeake Bay, *Biogeosciences*, 19, 4209-4226, <https://doi.org/10.5194/bg-19-4209-2022>, 2022.
- 850 Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., Chou, W.-C., Zhai, W., Hollibaugh, J. T., Wang, Y., Zhao, P., Guo, X., Gundersen, K., Dai, M., and Gong, G.-C.: Acidification of subsurface coastal waters enhanced by eutrophication, *Nature Geoscience*, 4, 1297, <https://doi.org/10.1038/ngeo1297>, 2011.
- Cai, W.-J., Huang, W.-J., Luther III, G. W., Pierrot, D., Li, M., Testa, J., Xue, M., Joesoef, A., Mann, R., Brodeur, J., Xu, Y.-

- 855 Y., Chen, B., Waldbusser, G. G., Cornwell, J. and Kemp, W. M.: Redox reactions and weak buffering capacity lead to acidification in the Chesapeake Bay, *Nature Communications*, 8, 369, <https://doi.org/10.1038/s41467-017-00417-7>, 2017.
- Carstensen, J. and Duarte, C. M.: Drivers of pH variability in coastal ecosystems, *Environmental Science and Technology*, 53, 4020-4029, <https://doi.org/10.1021/acs.est.8b03655>, 2019.
- Christian, J. and Ono, T.: Ocean acidification and deoxygenation in the North Pacific Ocean, North Pacific Marine Science Organization, Sidney, 116 pp., 2019.
- 860 Coynel, A., Seyler, P., Etcheber, H., Meybeck, M. and Orange, D.: Spatial and seasonal dynamics of total suspended sediment and organic carbon species in the Congo River, *Global Biogeochemical Cycles*, 19, GB4019, <https://doi.org/10.1029/2004GB002335>, 2005.
- Delille, B., Borges, A. V. and Delille, D.: Influence of giant kelp beds (*Macrocystis pyrifera*) on diel cycles of  $p\text{CO}_2$  and DIC  
865 in the Sub-Antarctic coastal area, *Estuarine, Coastal and Shelf Science*, 81, 114-122, <https://doi.org/10.1016/j.ecss.2008.10.004>, 2009.
- Dickson, A. G.: Standard potential of the reaction:  $\text{AgCl(s)} + 12\text{H}_2(\text{g}) = \text{Ag(s)} + \text{HCl(aq)}$ , and the standard acidity constant of the ion  $\text{HSO}_4^-$  in synthetic sea water from 273.15 to 318.15 K, *The Journal of Chemical Thermodynamics*, 22, 113-127, [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.
- 870 Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J. A., and McCulloch, M.: Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH, *Estuaries and Coasts*, 36, 221-236, <https://doi.org/10.1007/s12237-013-9594-3>, 2013.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D. and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 320, 1490e1492, <https://doi.org/10.1126/science.1155676>, 2008.
- 875 Feely, R. A., Alin, S. R., Carter, B., Bednarsek, N., Hales, B., Chan, F., Hill, T. M., Gaylord, B., Sanford, E., Byrne, R. H., Sabine, C. L., Greeley, D. and Juraneck, L.: Chemical and biological impacts of ocean acidification along the west coast of North America, *Estuarine, Coastal and Shelf Science*, 183, 260-270, <https://doi.org/10.1016/j.ecss.2016.08.043>, 2016.
- Frieder, C. A., Nam, S. H., Martz, T. R. and Levin, L. A.: High temporal and spatial variability of dissolved oxygen and pH in a nearshore California kelp forest, *Biogeoscience*, 9, 3917-3930, <https://doi.org/10.5194/bg-9-3917-2012>, 2012.
- 880 Fujii, M., Takao, S., Yamaka, T., Akamatsu, T., Fujita, Y., Wakita, M., Yamamoto, A., and Ono, T.: Continuous monitoring and future projection of ocean warming, acidification, and deoxygenation on the subarctic coast of Hokkaido, Japan, *Frontiers in Marine Science*, 8, 590020, <https://doi.org/10.3389/fmars.2021.590020>, 2021.
- Fujii, M., Hamanoue, R., Bernardo, L. P., Ono, T., Dazai, A., Oomoto, S., Katao, S., Wakita, M., and Tanaka, T.: Observed and projected impacts of coastal warming, acidification and deoxygenation on Pacific oyster (*Crassostrea gigas*) farming:  
885 Case studies in the Hinase Area, Okayama Prefecture and Shizugawa Bay, Miyagi Prefecture, Japan, *Biogeoscience*, in press, 2023.
- Giblin, A. E., Hopkinson, C. S. and Tucker, J.: Benthic metabolism and nutrient cycling in Boston Harbor, Mountains, *Estuaries*,

- 20, 346-364, <https://doi.org/10.2307/1352349>, 1997.
- 890 Gomez, F. A., Wanninkhof, R., Barbero, L. and Lee, S.-K.: Increasing river alkalinity slows ocean acidification in the northern Gulf of Mexico, *Geophysical Research Letters*, 48, e2021GL096521, <https://doi.org/10.1029/2021GL096521>, 2021.
- Guo, X., Yao, Z., Gao, Y., Luo, Y., Xu, Y., and Zhai, W.: Seasonal variability and future projection of ocean acidification on the East China Sea shelf off the Changjiang Estuary, *Frontiers in Marine Science*, 8, 770034, <https://doi.org/10.3389/fmars.2021.770034>, 2021.
- 895 Hauri, C., Gruber, N., Vogt, M., Doney, S. C., Feely, R. A., Lachkar, Z., Leinweber, A., McDonnell, A. M. P., Munnich, M. and Plattner, G.-K.: Spatiotemporal variability and long-term trends of ocean acidification in the California Current System, *Biogeosciences*, 10, 193-216, <https://doi.org/10.5194/bg-10-193-2013>, 2013.
- Higashi, H., Sato, Y., Yoshinari, H., Maki, H., Koshikawa, H., Kanaya, G. and Uchiyama, Y.: Freshwater discharge from small river basins and its impacts on reproductivity of coastal hydrodynamic simulation in Seto Inland Sea, *Proceedings of the Japan Society of Civil Engineers, Ser. B*, 74, 1135-11140, in Japanese, 2018.
- 900 Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan, A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., Crook, E. D., Kroeker, K. J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C. and Martz, T. R.: High-Frequency Dynamics of Ocean pH: A Multi-Ecosystem Comparison, *PLoS One*, 6, e28983, <https://doi.org/10.1371/journal.pone.0028983>, 2012.
- Hoshiba, Y., Hasumi, H., Itoh, S., Matsumura, Y. and Nakada, S.: Biogeochemical impacts of flooding discharge with high 905 suspended sediment on coastal seas: A modelling study for a microtidal open bay, *Scientific Reports*, 11, 21322, <https://doi.org/10.1038/s41598-021-00633-8>, 2021.
- Iida, Y., Takatani, Y., Kojima, A. and Ishii, M.: Global trends of ocean CO<sub>2</sub> sink and ocean acidification: an observation-based reconstruction of surface ocean inorganic carbon variables, *Journal of Oceanography*, 77, 323-358, <https://doi.org/10.1007/s10872-020-00571-5>, 2021.
- 910 Ishida, H., Isono, R. S., Kita, J. and Watanabe, Y. W.: Long-term ocean acidification trends in coastal waters around Japan, *Nature Scientific Reports*, 11, 5052, <https://doi.org/10.1038/s41598-021-84657-0>, 2021.
- Ishizu, M., Miyazawa, Y., Tsunoda, T., and Ono, T.: Long-term trends in pH in Japanese coastal seawater, *Biogeosciences*, 16, 4747-4763, <https://doi.org/10.5194/bg-16-4747-2019>, 2019.
- Izett, R. W., Manning, C. C., Hamme, R. C., and Tortell, P. D.: Refined estimates of net community production in the Subarctic 915 Northeast Pacific derived from  $\Delta O_2/Ar$  measurements with N<sub>2</sub>O-based corrections for vertical mixing, *Global Biogeochemical Cycles*, 32, 326-350, <https://doi.org/10.1002/2017GB005792>, 2018.
- Japan Fisheries Agency, Final report of the research for forest - river - ocean linkage for the creation of appropriate fisheries environment. 118pp., in Japanese, 2004.
- Jiang, Z.-P., Tyrrell, T., Hydes, D. J., Dai, M., and Hartman, S. E.: Variability of alkalinity and the alkalinity-salinity 920 relationship in the tropical and subtropical surface ocean, *Global Biogeochemical Cycles*, 28, 729-742,

- <https://doi.org/10.1002/2013GB004678>, 2014.
- Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S. K. and Olsen, A.: Surface ocean pH and buffer capacity: past, present and future, *Nature Scientific Report*, 9, 18624, <https://doi.org/10.1038/s41598-019-55039-4>, 2019.
- 925 Johnson, Z. I., Wheeler, B. J., Blinebry, S. K., Carlson, C. M., Ward, C. S. and Hunt, D. E.: Dramatic variability of the carbonate system at a temperate coastal ocean site (Beaufort, North Carolina, USA) is regulated by physical and biogeochemical processes on multiple timescales, *PLoS ONE*, 8, e85117, <https://doi.org/10.1371/journal.pone.0085117>, 2013.
- Jones, D. C., Ito, T., Takano, Y., and Hsu, W.-C.: Spatial and seasonal variability of the air-sea equilibration timescale of carbon dioxide, *Global Biogeochemical Cycles*, 28, 1163-1178, <https://doi.org/10.1002/2014GB004813>, 2014.
- 930 Kakehi, S., Naiki, K., Kodama, T., Wagawa, T., Kuroda, H., and Ito, S.-I.: Projections of nutrient supply to a wakame (*Undaria pinnatifida*) seaweed farm on the Sanriku Coast of Japan, *Fisheries Oceanography*, 27, 323-335, <https://doi.org/10.1111/fog.12255>, 2018.
- Kessouri, F., McWilliams, J. C., Bianchi, D., Sutula, M., Renault, L., Deutsch, C., Feely, R. A., McLaughlin, K., Ho, M., Howard, E. M., Bednars'ek, N., Damien, P., Molemaker, J. and Weisberg, S. B.: Coastal eutrophication drives acidification, oxygen loss, and ecosystem change in a major oceanic upwelling system, *Proceedings of the National Academy of Sciences*, 935 118, e2018856118, <https://doi.org/10.1073/pnas.2018856118>, 2021.
- Kim, J., Blair, N. E., Ward, A. S. and Goff, K.: Storm-induced dynamics of particulate organic carbon in Clear Creek, Iowa: An intensively managed landscape critical zone observatory story, *Frontiers in Water*, 2, 578261, <https://doi.org/10.3389/frwa.2020.578261>, 2020.
- 940 Ko, Y. H., Lee, K., Noh, J. H., Lee, C. M., Kleypas, J. A., Jeong, H. J. and Kim, K. Y.: Influence of ambient water intrusion on coral reef acidification in the Chuuk Lagoon, located in the coral-rich western Pacific Ocean, *Geophysical Research Letters*, 43, 3830-3838, <https://doi.org/10.1002/2016GL068234>, 2016.
- Kobayashi, J.: Study on average water quality of Japanese rivers and its features, *Studies in Agriculture*, 48, 63-104, in Japanese, 1960.
- 945 Kubo, A., Maeda, Y., and Kanda, J.: A significant net sink for CO<sub>2</sub> in Tokyo Bay, *Nature Scientific Reports*, 7, 44355, <https://doi.org/10.1038/srep44355>, 2017.
- Kurihara, H., Kato, S. and Ishimatsu, A.: Effects of increased seawater pCO<sub>2</sub> on early development of the oyster *Crassostrea gigas*, *Aquatic Biology*, 1, 91-98, 2007.
- 950 Laruelle, G. G., Dürr, H. H., Slomp, C. P. and Borges, A. V.: Evaluation of sinks and sources of CO<sub>2</sub> in the global coastal ocean using a spatially explicit typology of estuaries and continental shelves, *Geophysical Research Letters*, 37, L15607, <https://doi.org/10.1029/2010GL043691>, 2010.
- Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A. and Tjiputra, J.: Trends and drivers in global surface ocean pH over the past three decades, *Biogeoscience*, 12, 1285-1298, <https://doi.org/10.5194/bg-12-1285-2015>, 2015.
- Lee, K., Kim, T.-W., Bryne, R. H., Millero, F. J., Feely, R. A. and Liu, Y.-M.: The universal ratio of boron to chlorinity for

- the North Pacific and North Atlantic oceans, *Geochimica et Cosmochimica Acta*, 74, 1801-1811,  
955 <https://doi.org/10.1016/j.gca.2009.12.027>, 2010.
- Leiva-Dueñas, C., Graversen, A. E. L., Banta, G. T., Holmer, M., Masque, P., Stæhr, P. A. U. and Krause-Jensen, D.: Capturing  
of organic carbon and nitrogen in eelgrass sediments of southern Scandinavia, *Limnology and Oceanography*, online first,  
<https://doi.org/10.1002/lno.12299>, 2023.
- Lowe, A. T., Bos, J., Ruesink, J.: Ecosystem metabolism drives pH variability and modulates long-term ocean acidification in  
960 the Northeast Pacific coastal ocean, *Nature Scientific Reports*, 9, 963, <https://doi.org/10.1038/s41598-018-37764-4>, 2019.
- Lueker, T. J., Dickson, A. G. and Keeling, C. D.: Ocean  $p\text{CO}_2$  calculated from dissolved inorganic carbon, alkalinity, and  
equations for K1 and K2: validation based on laboratory measurements of  $\text{CO}_2$  in gas and seawater at equilibrium, *Marine  
Chemistry*, 70, 105-119, 2000. [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0)
- Ministry of Environment, Japan: Public Water Quality Database, available at [https://water-pub.env.go.jp/water-pub/mizu-](https://water-pub.env.go.jp/water-pub/mizu-site/mizu/kousui/dataMap.asp)  
965 [site/mizu/kousui/dataMap.asp](https://water-pub.env.go.jp/water-pub/mizu-site/mizu/kousui/dataMap.asp) (last access: October 29 2022), 2022.
- Mongin, M., Baird, M. E., Tilbrook, B., Matear, R. J., Lenton, A., Herzfeld, M., Wild-Allen, K., Skerratt, J., Margvelashvili,  
N., Robson, B. J., Duarte, C. M., Gustafsson, M. S. M., Ralph, P. J. and Steven, A. D. L.: The exposure of the Great Barrier  
Reef to ocean acidification, *Nature Communications*, 7, 10732, <https://doi.org/10.1038/ncomms10732>, 2016.
- Niigata Prefectural Institution of Fisheries Oceanography: 1997 Annual report of ocean monitoring for ocean climate forecast  
970 for fisheries, 106 pp., in Japanese, 1998.
- Nishijima, W.: Management of nutrient concentrations in the Seto Inland Sea, *Bulletin on Coastal Oceanography*, 56, 13-19,  
in Japanese, 2018.
- Okada, T., Maruya, Y., Nakayama, K. and Iseri, E.: A study on restoration of eelgrass (*Zostera marina*) damaged by tsunami  
in Miyako Bay, *Proceedings of the Japan Society of Civil Engineers, Ser. B*, 70, 1186-1190, in Japanese, 2014.
- 975 Onitsuka, T., Takami, H., Muraoka, D., Matsumoto, Y., Nakatsubo, A., Kimura, R., Ono, T., Nojiri, Y.: Effects of ocean  
acidification with  $p\text{CO}_2$  diurnal fluctuations on survival and larval shell formation of Ezo abalone, *Haliotis discus hannai*,  
*Marine Environmental Research* 134, 28-36, <https://doi.org/10.1016/j.marenvres.2017.12.015>, 2018.
- Papalexioiu, S. M. and Montanari, A.: Global and regional increase of precipitation extremes under global warming, *Water  
Resources Research*, 55, 4901-4914, <https://doi.org/10.1029/2018WR024067>, 2019.
- 980 Pierrot, D., Lewis, E., and Wallace, D.W.R.: MS Excel Program Developed for  $\text{CO}_2$  System Calculations. ORNL/CDIAC-  
105a. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak  
Ridge, Tennessee. [https://doi.org/10.3334/CDIAC/otg.CO2SYS\\_XLS\\_CDIAC105a](https://doi.org/10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a), 2006.
- Point, D., Bareille, G., Amouroux, D., Etcheber, H. and Donard, O. F. X.: Reactivity, interactions and transport of trace  
elements, organic carbon and particulate material in a mountain range river system (Adour River, France), *Journal of  
985 Environmental Monitoring*, 9, 157-167, <https://doi.org/10.1039/b616312b>, 2007.
- Potouroglou, M., Bull, J. C., Krauss, K. W., Kennedy, H. A., Fusi, M., Daffonchio, D., Mangora, M. M., Githaiga, M. N.,

- Diele, K., and Huxham, M.: Measuring the role of seagrasses in regulating sediment surface elevation, *Nature Scientific Reports*, 7, 11917, <https://doi.org/10.1038/s41598-017-12354-y>, 2017.
- 990 Provoost, P., Heuven, S. v., Soetaert, K., Laane, R. W. P. M. and Middelburg, J. J.: Seasonal and long-term changes in pH in the Dutch coastal zone, *Biogeosciences*, 7, 3869-3878, <https://doi.org/10.5194/bg-7-3869-2010>, 2010.
- Qin, C., Zhang, G., Zheng, W., Han, Y., and Liu, S.: High-resolution distributions of O<sub>2</sub>/Ar on the northern slope of the South China Sea and estimates of net community production, *Ocean Science*, 17, 249-264, <https://doi.org/10.5194/os-17-249-2021>, 2021.
- 995 Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of sea-water, *The Sea*, vol. 2, Hill, M. N. (ed), pp. 26-77, Interscience, New York, 1963.
- Ricart, A. M., Ward, M., Hill, T. M., Sanford, E., Kroeker, K. J., Takeshita, Y., Merolla, S., Shukla, P., Ninokawa, A. T., Elsmore, K. and Gaylord, B.: Coast-wide evidence of low pH amelioration by seagrass ecosystems, *Global Change Biology*, 27, 2580-2591, <https://doi.org/10.1111/gcb.15594>, 2021.
- 1000 Rosenau, N. A., Galavotti, H., Yates, K. K., Bohlen, C. C., Hunt, C. W., Liebman, M., Brown, C. A., Pacella, S. R., Largier, J. L., Nielsen, K. J., Hu, X., McCutcheon, M. R., Vasslides, J. M., Poach, M., Ford, T., Johnston, K. and Steele, A.: Integrating High-resolution coastal acidification monitoring data across seven United States Estuaries, *Frontiers in Marine Science*, 8, 679913, <https://doi.org/10.3389/fmars.2021.679913>, 2021.
- Salisbury, J., Green, M., Hunt, C., and Campbell, J.: Coastal acidification by rivers: A threat to shellfish?, *EOS*, 89, 513-528, 2008. <https://doi.org/10.1029/2008EO500001>
- 1005 Salisbury, J. E. and Jönsson, B. F.: Rapid warming and salinity changes in the Gulf of Maine alter surface ocean carbonate parameters and hide ocean acidification, *Biogeochemistry*, 141, 401-418, <https://doi.org/10.1007/s10533-018-0505-3>, 2018.
- Sugimura, A., Takeda, A., Nihei, Y. and Ohtsuki, K.: Evaluation of sediment, nutrient and organic matter transport from a wide range of land to coastal ocean, *Proceedings of the Japan Society of Civil Engineers, Ser. B*, 71, 1423-1428, in Japanese, 1010 2015.
- Sunda, W. G. and Cai, W.-J.: Eutrophication induced CO<sub>2</sub>-acidification of subsurface coastal waters: Interactive effects of temperature, salinity, and atmospheric *p*CO<sub>2</sub>, *Environmental Science and Technology*, 46, 10651-10659, <https://doi.org/10.1021/es300626f> 2012.
- 1015 Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., Sweeney, C. and Munro, D. R.: Climatological distributions of pH, *p*CO<sub>2</sub>, total CO<sub>2</sub>, alkalinity, and CaCO<sub>3</sub> saturation in the global surface ocean, and temporal changes at selected locations, *Marine Chemistry*, 164, 95-125, <https://doi.org/10.1016/j.marchem.2014.06.004>, 2014.
- Takatani, Y., Enyo, K., Iida, Y., Kojima, A., Nakano, T., Sasano, D., Kosugi, N., Midorikawa, T., Suzuki, T. and Ishii, M.: Relationships between total alkalinity in surface water and sea surface dynamic height in the Pacific Ocean, *Journal of*

- 1020 Geophysical Research, 119, 2806-2814, <https://doi.org/10.1002/2013JC009739>, 2014.
- Tokoro, T., Nakaoka, S.-i., Takao, S., Kuwae, T., Kubo, A., Endo, T. and Nojiri, Y.: Contribution of biological effects to carbonate system variations and the air–water CO<sub>2</sub> flux in urbanised bays in Japan, *Journal of Geophysical Research*, 126, e2020JC016974, <https://doi.org/10.1029/2020JC016974>, 2020.
- 1025 Tsukamoto, H. and Yanagi, T.: Temporal and Spatial variations in water temperature, salinity, pH, nutrient salts and chlorophyll-a in the Seto Inland Sea, *Umi no Kenkyu*, 7, 11-19, in Japanese, 1998.
- Vargas, C. A., Contreras, P. Y., Pérez, C. A., Sobarzo, M., Saldías, G. S. and Salisbury, J.: Influences of riverine and upwelling waters on the coastal carbonate system off Central Chile and their ocean acidification implications, *Journal of Geophysical Research*, 121, 1468-1483, <https://doi.org/10.1002/2015JG003213>, 2015.
- 1030 Venkatesan, R., Senthilkumar, P., Vedachalam, N., and Muruges, P.: Biofouling and its effects in sensor mounted moored observatory system in Northern Indian Ocean, *International Biodeterioration & Biodegradation*, 116, 198-204, <http://dx.doi.org/10.1016/j.ibiod.2016.10.034>, 2017
- 1035 Wakita, M., Sasaki, K., Nagano, A., Abe, H., Tanaka, T., Nagano, K., Sugie, K., Kaneko, H., Kimoto, K., Okunishi, T., Takada, M., Yoshino, J. and Watanabe, S.: Rapid reduction of pH and CaCO<sub>3</sub> saturation state in the Tsugaru Strait by the intensified Tsugaru Warm Current during 2012–2019, *Geophysical Research Letters*, 48, e2020GL091332, <https://doi.org/10.1029/2020GL091332>, 2021.
- Waldbusser, G. G. and Salisbury, J. E.: Ocean acidification in the coastal zone from an organism’s perspective: Multiple system parameters, frequency domains, and habitats, *Annual Review of Marine Science*, 6, 221-47, <https://doi.org/10.1146/annurev-marine-121211-172238>, 2014.
- 1040 Waldbusser, G. G., Hales, B., Langdon, C. J., Haley, B. A., Schrader, P., Brunner, E. L., Gray, M. W., Miller, C. A. and Gimenez, I.: Saturation-state sensitivity of marine bivalve larvae to ocean acidification, *Nature Climate Change*, 5, 273-280, <https://doi.org/10.1038/NCLIMATE2479>, 2015.
- Wallace, R. B., Baumann, H., Grear, J. S., Aller, R. C. and Gobler, C. J.: Coastal ocean acidification: The other eutrophication problem, *Estuarine, Coastal and Shelf Science*, 148, 1-13, <https://doi.org/10.1016/j.ecss.2014.05.027>, 2014.
- 1045 Wang, S. R., Iorio, D. D., Cai, W.-J. and Hopkinson, C. S.: Inorganic carbon and oxygen dynamics in a marsh-dominated estuary, *Limnology and Oceanography*, 63, 47-71, <https://doi.org/10.1002/lno.10614>, 2018.
- Weiss, R. F.: The solubility of nitrogen, oxygen and argon in water and seawater, *Deep-Sea Research*, 17, 721-735, 1970.
- World Data Centre for Greenhouse Gases: Current State of GHCs, [https://gaw.kishou.go.jp/publications/global\\_mean\\_mole\\_fractions](https://gaw.kishou.go.jp/publications/global_mean_mole_fractions), last cited 2023
- 1050 Yamamoto, H., Yoshiki, K., Komatsu, T., Sasa, S., Hayama, M., Murata, H. and Yanagi, T.: Numerical evaluation of environmental capacity of aquaculture in Shizugawa Bay - comparison before and after the Great East Japan Earthquake-, *Proceedings of the Japan Society of Civil Engineers, Ser. B*, 73, 1339-1344, 2017.
- Yamamoto, H., Yoshiki, K., Komatsu, T., Sasa, S. and Yanagi, T.: Numerical evaluation of carrying capacity of aquaculture



- in Shizugawa Bay using land-sea integrated ecosystem model, Proceedings of the Japan Society of Civil Engineers, Ser. B, 74, 1279-1284, in Japanese, 2018.
- 1055 Yamamoto, T., Ishida, M. and Seiki, T.: Long-term variation in phosphorus and nitrogen concentrations in the Ohta River water, Hiroshima, Japan as a major factor causing the change in phytoplankton species composition Bulletin of Japan Society of Fisheries Oceanography, 102-109, in Japanese, 2002.
- Yamamoto, T., Orimoto, K., Asaoka, S., Yamamoto, H. and Onodera, S.-i.: A Conflict between the legacy of eutrophication and cultural oligotrophication in Hiroshima Bay, *Oceans*, 2, 546-565, <https://doi.org/10.3390/oceans2030031>, 2021.
- 1060 Yamamoto-Kawai, M., Ito, S., Kurihara, H. and Kanda, J.: Ocean acidification state in the highly eutrophic Tokyo Bay, Japan: Controls on seasonal and interannual variability, *Frontiers in Marine Sciences*, 8, 642041, <https://doi.org/10.3389/fmars.2021.642041>, 2021.
- Yanagi, T. and Ishii, D.: Open ocean originated phosphorus and nitrogen in the Seto Inland Sea, Japan, *Journal of Oceanography*, 60, 1001-1005, 2004.
- 1065 Yao, H., Wang, J., Han, Y., Jiang, X. and Chen, J.: Decadal acidification in a subtropical coastal area under chronic eutrophication, *Environmental Pollution*, 293, 118487, <https://doi.org/10.1016/j.envpol.2021.118487>, 2022.
- Zhan, Y., Hu, W., Zhang, W., Liu, M., Duan, L., Huang, X., Chang, Y. and Li, C.: The impact of CO<sub>2</sub>-driven ocean acidification on early development and calcification in the sea urchin *Strongylocentrotus intermedius*, *Marine Pollution Bulletin*, 112, 291-302, <https://doi.org/10.1016/j.marpolbul.2016.08.003>, 2016.
- 1070 Zhang, L. J., Wang, L., Cai, W.-J., Liu, D. M. and Yu, Z. G.: Impact of human activities on organic carbon transport in the Yellow River, *Biogeosciences*, 10, 2513-2524, <https://doi.org/10.5194/bg-10-2513-2013>, 2013.