

Central Arctic Ocean surface-atmosphere exchange of CO₂ and CH₄ constrained by direct measurements

John Prytherch^{1,2,3}, Sonja Murto^{1,3}, Ian Brown⁴, Adam Ulfsbo⁵, Brett F. Thornton^{3,6}, Volker Brüchert^{3,6}, Michael Tjernström^{1,3}, Anna Lunde Hermansson⁷, Amanda T. Nylund⁷, Lina A. Holthusen^{8*}

¹Department of Meteorology, Stockholm University, Stockholm, 10691, Sweden

²Department of Earth Sciences, Uppsala University, Uppsala, 75236, Sweden

³Bolin Centre for Climate Research, Stockholm, 10691, Sweden

⁴Plymouth Marine Laboratory, Plymouth, PL1 3DH, United Kingdom

⁵Department of Marine Sciences, University of Gothenburg, Gothenburg, 40530, Sweden

⁶Department of Geological Sciences, Stockholm University, Stockholm, 10691, Sweden

⁷Department of Mechanics and Maritime Sciences, Chalmers University of Technology, Gothenburg, 41296, Sweden

⁸Chemical Oceanography Department, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, 24105, Germany

* now at: Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, 26129, Germany

Correspondence to: John Prytherch (john.prytherch@misu.su.se)

Abstract. The Central Arctic Ocean (CAO) plays an important role in the global carbon cycle, but the current and future exchange of the climate-forcing trace gases methane (CH₄) and carbon dioxide (CO₂) between the CAO and the atmosphere is highly uncertain. In particular, there are very few observations of near surface gas concentrations or direct air-sea CO₂ flux estimates, and no previously reported direct air-sea CH₄ flux estimates from the CAO. Furthermore, the effect of sea ice on the exchange is not well understood. We present direct measurements of the air-sea flux of CH₄ and CO₂, as well as air-snow fluxes of CO₂ in the summertime CAO North of 82.5 N from the Synoptic Arctic Survey (SAS) expedition carried out on the Swedish icebreaker *Oden* in 2021.

Measurements of air-sea CH₄ and CO₂ flux were made using floating chambers deployed in leads accessed from sea ice and from the side of *Oden*, and air-snow fluxes determined from chambers deployed on sea ice. Gas transfer velocities determined from fluxes and surface water dissolved gas concentrations exhibited a weaker wind speed dependence than existing parameterisations, with a median sea-ice lead gas transfer rate of 2.5 cm hr⁻¹, applicable over the observed 10-m wind speed range (1-11 m s⁻¹). Average observed air-sea CO₂ flux was -7.6 mmol m⁻² day⁻¹, and the average air-snow CO₂ flux -1.1 mmol m⁻² day⁻¹. Extrapolating these fluxes and the corresponding sea-ice concentrations gives an August and September flux for the CAO of -1.75 mmol m⁻² day⁻¹, within the range of previous indirect estimates.

Average observed air-sea CH₄ flux of 3.5 μmol m⁻² day⁻¹, accounting for sea ice concentration, equates to an August and September CAO flux of 0.35 μmol m⁻² day⁻¹, lower than previous estimates and implying that the CAO is a very small, << 1%, contributor to the Arctic flux of CH₄ to the atmosphere.

Deleted: ²... Ian Brown⁴³... Adam Ulfsbo⁴⁴... Brett F. Thornton^{32...65}... Volker Brüchert^{32...65}... Michael Tjernström^{1,32}... Anna Lunde Hermansson¹⁶... Amanda T. Nylund¹⁶... Lina A. Holthusen^{8*7} ... [1]

Formatted: Superscript

Deleted: ²

Deleted: ³

Deleted: ⁴...epartment of Marine Sciences, University of Gothenburg, Gothenburg, 405301319 ... [2]

Deleted: ⁵

Deleted: ⁶

Deleted: ⁷

Deleted: 's...(CAO) plays an important role in the global carbon cycle, but the current and future role in the...exchange of the climate-forcing trace gases methane (CH₄) and carbon dioxide (CO₂) between the CAO andwith...the atmosphere is highly uncertain. In particular, in particular ... [3]

Formatted: Subscript

Deleted: ...he effect of sea ice on the exchange is not well understood... There are no previously reported direct air-sea methane (CH₄) flux estimates from the CAO...e present direct measurements of the air-sea flux of CH₄...and carbon dioxide (...O₂)...as well as air-snow-ice ... [4]

Deleted: and air-snow fluxes determined from chambers deployed on sea ice. Gas transfer velocities determined from F...luxes and surface water dissolved gas concentrations from surface water were used to determine gas transfer velocities that ...xhibited a weaker wind speed dependence than existing parameterisations, with a median sea-ice lead gas transfer rate in sea-ice leads...f 2.5 ... [5]

Formatted: Font: Italic

Deleted: CO₂ ...ir-sea CO₂ flux wasere...-7.6 mmol m⁻² day⁻¹, and the average air-snow CO₂ flux -1.1 mmol m⁻² day⁻¹. Extrapolating these fluxes and the corresponding sea-...ce concentrations gives an August and September flux for the CAO of -1.75 mmol m⁻² day⁻¹ and a corresponding annual CAO uptake of 126.6 Tg yr⁻¹... within the range of previous indirect estimates... [6]

Formatted: Superscript

Deleted: or 9.2 Gg yr⁻¹, ...ower than previous estimates and implying that the CAO is a very small, << 1%, contributor to the Arctic flux of CH₄ to the atmosphere from the Arctic ... [7]

1 Introduction

The Arctic is on average warming up to 4 times faster than the global average rate (Rantanen et al., 2022), manifested in dramatic reductions in sea-ice extent (Onarheim et al., 2018) and thickness (Kwok, 2018). Reduced sea-ice cover in a warming Arctic is expected to have complex effects on air-sea gas exchanges (Parmentier et al., 2013), in particular for the exchange of the two gases whose rising atmospheric concentrations are principally responsible for the observed global warming, carbon dioxide (CO₂) and methane (CH₄). The parameterisation of gas transfer in the presence of sea ice is still under debate but is expected to have large impacts on polar carbon budgets: for example, Arctic Ocean CO₂ uptake estimates are highly dependent on the parameterisation choice, with resulting uptake differences of 50 Tg C yr⁻¹, ~30 % of the total (Yasanaka et al., 2018). The Arctic Ocean corresponds to 5-14% of the global ocean CO₂ sink from 3% of the global ocean area (Bates and Mathis, 2009; Yasanaka et al., 2016; Yasanaka et al., 2018).

The Central Arctic Ocean's (CAO) role in carbon cycling and exchange of climate-forcing trace gases with the atmosphere is uncertain due to several factors, including limited observations of dissolved gas concentrations and air-sea fluxes. The CAO is defined as the deep-water part of the Arctic Ocean, excluding the shallow shelf seas (Jakobsson, 2002). Thus defined, the CAO has an average depth of 2748 m and an area of 4.5 x 10⁶ km², which corresponds to about 47% of the surface area of the entire Arctic Ocean.

Sources of CH₄ from the Arctic Ocean are poorly constrained and spatially variable (Thornton et al., 2016a). Oceans have long been seen as generally a weak source of CH₄. The shelf seas of the Arctic and particularly the extensive shallow East Siberian Shelf are believed to be a large source, though are poorly constrained. Here, with potential sources of CH₄ from thawing subsea permafrost and riverine input (Weber et al., 2019; Manning et al., 2020), fluxes of 9-286 μmol m⁻² day⁻¹ were observed with direct measurement (Thornton et al., 2020), and 187-238 μmol m⁻² day⁻¹ were estimated from seawater concentrations (Thornton et al., 2016b). While the integrated magnitude of these Arctic marine methane emissions remains under debate, it appears to presently be on the scale of ~4-5 Tg CH₄ yr⁻¹ or 100 μmol m⁻² day⁻¹ (Thornton et al., 2016a; 2016b; 2020), ~1% of global CH₄ emissions (Saunio et al., 2020). There are previously no direct CH₄ flux measurements available from the CAO. The estimates that are available are derived from near-surface (typically ~ 10 m depth) concentrations (Manning et al., 2022; Damm et al., 2018; Fenwick et al., 2017; Lorenson et al., 2016). Seawater CH₄ measurements from the North American Arctic shelf and Canada Basin suggest small fluxes, 0.3 – 2.2 μmol m⁻² day⁻¹ (Manning et al., 2022; Fenwick et al., 2017). Excess CH₄ may be transported from shallow sea areas across the Arctic Ocean in the water column and also frozen in sea ice and in brines within the ice (Damm et al., 2018). Seawater CH₄ concentrations from the Beaufort Shelf and CAO were used to estimate an Arctic-wide CH₄ flux of 7.5 μmol m⁻² day⁻¹, and a potential flux, if the ice would disappear from currently ice-covered areas, of 18-63 μmol m⁻² day⁻¹ (Lorenson et al., 2016). Emissions of CH₄ derived from changing seawater concentrations following a winter storm ~150 km north of Svalbard were 19 μmol m⁻² day⁻¹ during and 5 μmol m⁻² day⁻¹ afterwards (Silyakova et al., 2022). Much higher fluxes, 125 μmol m⁻² day⁻¹, extending for at least 50 km over wintertime (November and April) sea-ice leads at latitudes up to 82°N, have been derived from aircraft-based atmospheric profile

Moved down [1]: The Central Arctic Ocean's (CAO) role in carbon cycling and exchange of climate-forcing trace gases with the atmosphere is uncertain due to several factors, including limited observations of dissolved gas concentrations and air-sea fluxes.

Moved down [2]: Throughout this manuscript we use the convention that positive fluxes are upwards, a release of gas from the surface to the atmosphere, and negative fluxes are a downwards flux, indicating an uptake of gas by the surface from the atmosphere.

Moved (insertion) [1]

Deleted: methane

Formatted: Subscript

185 measurements (Kort et al., 2012). These fluxes, observed over deep water, are ascribed to local CH₄ production associated with the mixed sea ice and lead environment. The large range of indirect estimates of CAO CH₄ emissions presents a challenge to modeling and highlights the need for direct measurements.

Deleted: The authors ascribe t

Formatted: Subscript

190 The CAO is generally undersaturated with respect to atmospheric CO₂. This undersaturation is due to a combination of the low temperatures and resulting lower gas saturation, dilution due to freshwater input, limited atmospheric equilibration due to the presence of sea ice, CaCO₃ dissolution (Fransson et al., 2017), vertical mixing and primary production, particularly in shelf seawater advected to central areas (Bates et al., 2006). Average Arctic Ocean uptake in summer months, with sea-ice concentration (SIC) of ~50% and a CO₂ partial pressure (pCO_{2W}) undersaturation of ~ 80 μ atm, is estimated at -4 mmol m⁻² day⁻¹ (Yasunaka et al., 2018). The majority of existing Arctic Ocean observations are from coastal regions, with average air-sea fluxes in the Eastern Arctic Ocean 0 –10 mmol m⁻² day⁻¹ (Manizza et al., 2019) and average air-sea fluxes in the Western Arctic Ocean somewhat higher, 0 –20 mmol m⁻² day⁻¹, primarily due to low SIC in the Chukchi Sea (Bates and Mathis, 2009; Ouyang et al., 2022). In the CAO, observations of pCO_{2W} and the air-sea flux are especially sparse. Yasunaka et al., (2018), using a self-organising map to extrapolate the sparse near-surface CO₂ observations, determine the mean flux for the CAO region to be below the uncertainty of their method (uncertainty 2.8 to 3.7 mmol m⁻² day⁻¹). Coupled ocean-biogeochemistry modelling gives an annual CAO flux of -2.2 ± 4.0 Tg CO₂ yr⁻¹ (Manizza et al., 2019). Earlier estimates from the Canada Basin suggested fluxes smaller than -3 mmol m⁻² day⁻¹ in periods with SIC near 100%, but an uptake of around -55 mmol m⁻² day⁻¹ in summer months, with lower SIC (Bates et al., 2006). High fluxes can occur when water undersaturated with CO₂, high winds and open water coincide, with CO₂ fluxes up to -86 mmol m⁻² day⁻¹ estimated in June close to the pack edge north of Svalbard (Fransson et al., 2017). Fluxes determined from direct, eddy covariance, measurements in areas adjacent to ice or from lead water surfaces are of the order of -10 mmol m⁻² day⁻¹, in both central and coastal areas of the Arctic Ocean (Prytherch et al., 2017; Dong et al., 2021; Prytherch and Yelland, 2021).

Deleted: For CO₂, t

205 The presence of sea ice makes the Arctic Ocean a highly heterogenous and dynamic environment over a wide range of spatial scales, complicating robust model representation of surface exchange processes. Wind speed is the primary forcing of gas exchange in the ice-free ocean (Wanninkhof et al., 2009). Sea ice limits air-sea gas exchange, but also contributes to its forcing through additional sea-ice dependent physical processes that impact interfacial mixing. This implies a non-linear relationship of gas transfer rate to sea-ice cover (Loose et al. 2014). The highest rates of gas exchange in pack ice regions occur through open water areas such as leads. Leads appear on scales from < 1 m to > 10 km, and can open, change size and close throughout all seasons (Marcq and Weiss, 2012). Sea-ice dependent processes impacting gas exchange from lead surfaces are: shear between floating ice and the underlying water (Lovely et al. 2015); upper ocean stability, stratification from freshwater during ice melt and convection-driven turbulent mixing from surface buoyancy during freeze periods (MacIntyre et al. 2010). Also, modifications to the wind stress input from ice-edge form drag, ice-wave interactions and short fetch conditions (Bigdeli et al. 2018). Direct ship-based eddy covariance measurements support an inverse linear scaling of exchange with SIC (Butterworth & Miller, 2016; Prytherch et al. 2017) while laboratory and indirect radon isotope measurements support an enhanced exchange (Fanning & Torres, 1991; Loose et al. 2011; 2017). Recent direct and indirect gas exchange observations

suggest that gas exchange will be reduced in the presence of sea ice (Rutgers van der Loeff et al. 2014; Prytherch and Yelland, 2021).

Strong, near-surface gradients in dissolved gas concentrations are prevalent in Arctic sea-ice regions (e.g. Miller et al., 2019; Ahmed et al., 2020; Dong et al., 2021). Unaccounted for, this stratification will bias flux estimates and parameterisations of gas transfer derived from sub-surface concentrations. Using eddy covariance (EC) flux measurements, Dong et al. (2021) showed that surface fCO_2 in the summertime Arctic marginal ice zone was $39 \mu\text{atm}$ lower than that measured from their ship's intake at 6 m depth. They determined that this was partly due to effects on CO_2 solubility from meltwater cooling and freshening, with approximately half of the reduction from other factors, presumably photosynthesis. Miller et al. (2019) report pCO_{2-w} differences between the surface and an intake at 7 m depth of between -180 and $+160 \mu\text{atm}$ during sampling in the Canadian Arctic Archipelago and Hudson Bay. The authors note that "The temperature differences between the underway system and the shallower samples were often contrary to the pCO_2 gradients..., indicating that pCO_2 was not simply controlled by surface heating and cooling". Due to surface longwave emission and the resulting cool skin effect (Woolf et al., 2016), temperature differences are always present between the sea surface and intake depths even when the upper ocean is well mixed.

Sea-ice can be porous due to brine channels within the ice and can exchange gases with the atmosphere (Delille et al. 2014). Ice-atmosphere fluxes of CO_2 are typically smaller than those through water surfaces, on the order of $1 \text{ mmol m}^{-2} \text{ day}^{-1}$, ~~but can be a significant contributor to regional fluxes in sea ice areas. Ice-atmosphere fluxes are largely dependent on~~ temperature (Delille et al., 2014) and snow cover thickness (Geilfus et al., 2012; Nomura et al., 2010). ~~Ice-atmosphere CH_4 fluxes are less well known.~~ Sea ice cores obtained on the Siberian shelf have been supersaturated with CH_4 , while cores from the CAO were close to equilibrium, but atmospheric fluxes were not determined (Damm et al., 2015). There are no reported measurements of ice-atmosphere CO_2 or CH_4 flux in the summertime CAO.

The high sensitivity and small footprint of the flux chamber technique makes it attractive for measurements in small water bodies such as leads. The technique has been frequently used in inland water bodies but is criticised because the chamber isolates the air-water interface from the wind and may modify the gas exchange process. While this isolation suggests that chambers reduce gas transfer, previous studies have often suggested an overestimation of k when determined from chambers, resulting from artefact turbulence at the water surface introduced by the chamber, particularly at very low wind speeds (Matthews et al., 2003; Vachon et al., 2010). The gas transfer of poorly soluble gases is controlled by mixing on the water side of the interfacial layer, and the mixing itself depends on forcings such as wind speed (Wanninkhof et al., 2009). If water is able to advect into the chamber rapidly and with minimal modification by the chamber, then the gas transfer within the chamber will be representative of the outside environment. Appropriate chamber designs with minimal penetration into the water layer, high surface to volume ratios, and short measuring intervals minimize such measurement bias.

To remedy some of the uncertainties reviewed above we here report direct measurements of the air-sea flux of CO_2 and CH_4 , gas transfer velocities determined from the air-sea flux measurements and surface water dissolved gas concentrations,

Moved (insertion) [3]

Deleted: and

Formatted: Subscript

and air-~~snow~~ fluxes of CO₂ during the period of rapid sea-ice melt in the summertime CAO. ~~We relate the gas transfer velocities to wind speed and lead width, discuss the measurement uncertainties and estimate regional CAO fluxes.~~

Deleted: ice

255 2. Methods

Measurements were obtained during the Synoptic Arctic Survey (SAS) expedition (Snoeijs-Leijonmalm et al., 2022) carried out on the Swedish icebreaker *Oden* in 2021. The science operations of the expedition began on August 1, when *Oden* reached the ice edge north of Svalbard (80.71°N, 11.20°E). *Oden* transited to the North Pole along 30°E, then towards the northern coast of Greenland along the Lomonosov Ridge, then East from the Morris Jesup Rise towards Svalbard, with relevant science
260 operations finishing on September 11 (Fig. 1, Fig. 2a).

Ice stations were carried out throughout the expedition during which *Oden* halted in the ice for periods ranging between several hours and 2 days in order to perform winch operations such as Conductivity Temperature Depth sensor and Niskin bottle casts and net deployments. During long-duration stations, sampling was carried out directly from the sea ice at locations within a radius of ~200 m of *Oden*, or to a radius of ~500m using *Oden*'s helicopter. During shorter stations where
265 on-ice work was not possible, measurements were performed from over the side of the ship (overside).

2.1 Gas flux measurements

Measurements of air-water CO₂ and CH₄ flux were made using floating chambers (e.g., Cole et al., 2010). Two types of floating chamber sampling were performed: ice-based sampling, with the chamber placed onto lead water or meltpond surfaces accessed from sea ice and allowed to float freely; and overside sampling during the shorter stations, with the chamber lowered
270 over *Oden*'s side to the water surface, resulting in sampling closely adjacent to the ship. Fluxes measured overside may be influenced by the presence of *Oden* impacting the gas exchange rate both through modification of the near-surface winds and through modification of ocean near-surface turbulence. ~~Measurements were made as far from any propellor movement or water flushing as possible but there may still have been some influence on the measurements.~~ In the presence of upper-ocean dissolved gas concentration gradients, mixing induced by *Oden* may modify the air-sea concentration difference that drives
275 the flux. As such, overside and ice-based flux and surface water gas concentration measurements are presented separately throughout this study.

Deleted: Furthermore, i

Two floating chamber flux systems were used during SAS, both comprising a gas analyser and air pump connected through tubing in a closed loop with the chamber. The first system used a Los Gatos Research (LGR) Greenhouse Gas Analyser cavity-enhanced laser spectrometer, measuring mixing ratios of CO₂, CH₄ and H₂O. The second system used a
280 Li-COR 7200RS non-dispersive infrared (NDIR) spectrometer, measuring CO₂ and H₂O mixing ratios. Surface fluxes of a gas species are determined from the gradient with time of the gas mixing ratio in the chamber, with an approximately linear gradient required for a successful flux measurement. The sampling time for each flux measurement was approximately 10 minutes, after which the chamber was manually raised from the surface and equilibrated with atmosphere, before being

Deleted:

Moved (insertion) [6]

replaced to begin the next sampling period. Each chamber deployment during SAS typically consisted of ~8 sampling periods (from 2 to 10). The mean of the fluxes measured within each deployment is used in the subsequent analysis, with the standard error of fluxes indicating the variability within each deployment. Fluxes were converted to $\text{mmol m}^{-2} \text{day}^{-1}$ using the chamber surface area, the volume of the total chamber system (chamber, tubing, and analyser measurement cell), chamber air pressure and the ideal gas law.

The same chambers were used with both analysers. The chambers are custom built, consisting of upturned polyethylene bowls, with foam floats attached and rubber stoppers in the roof through which the tubing is inserted (Supplemental Materials Fig. S1). The chambers are lightweight, low profile (volume 7489 ml, surface area 0.078 m^2), have small chamber wall intrusion depths ($< 3 \text{ cm}$) and are allowed to float freely. Wind is the dominant source of surface mixing energy in the open ocean and most large water bodies. Chamber flux measurement necessarily involves isolation of an area of water from the wind. The chamber design and sampling choices are made to minimise known measurement biases resulting from anchoring effects (Lorke et al., 2015), and from chamber size and shape and sampling duration (Matthews et al., 2003; Mannich et al., 2019). Similar chamber designs to that deployed here have been shown to determine gas exchange rates in agreement with those determined in streams and small lakes from tracer releases (Cole et al., 2010), surface dissipation measurements and IR imagery (Gálfalk et al., 2013). With appropriate flux chamber design (i.e., the relatively small and lightweight chambers used here) it is assumed that the wind-induced interfacial layer mixing, the controlling factor for air-sea gas transfer of poorly soluble species, is advected within the chamber with minimal modification of the mixing by the chamber.

Air-snow CO_2 flux measurements were made using a PP systems CPY-4 chamber connected to an EGM-4 NDIR analyser and mixed by a fan inside the chamber (e.g. Miller et al. 2015). The transparent chamber (volume 2344 ml, surface area 167 cm^2) was covered to minimise insolation effects, and deployed on undisturbed snow surfaces. The chamber collar was pressed 1 cm into snow to prevent air leaks. The sampling time was also 10 minutes here.

Measurements of surface atmosphere CO_2 and CH_4 flux were also made throughout the expedition by EC from a system installed on Oden's foremast (Prytherch et al., 2017). Post-cruise analysis determined that fluxes measured in sea-ice regions were below the limit of detection of this system. While the strong observed CO_2 undersaturation resulted in air-sea fluxes at the lead spatial scale that would be within the typical detectable range of EC systems, the high sea-ice concentration within the EC footprint (of approximately km^2 scale from the 20 m EC measurement height) throughout the expedition substantially reduced the signal size (e.g. Prytherch et al., 2017), and as such EC measurements are not used in the analysis here (see Supplementary Material).

Throughout this manuscript we use the convention that positive fluxes are upwards, i.e., indicating a release of gas from the surface to the atmosphere, and negative fluxes are a downwards flux, indicating uptake of gas by the surface from the atmosphere.

Deleted: in order to minimise biases on gas exchange rates from chamber effects (Cole et al., 2010; Lorke et al., 2015).

Moved up [6]: Surface fluxes of a gas species are determined from the gradient with time of the gas mixing ratio in the chamber, with an approximately linear gradient required for a successful flux measurement. The sampling time for each flux measurement was approximately 10 minutes, after which the chamber was manually raised from the surface and equilibrated with atmosphere, before being replaced to begin the next sampling period. Each chamber deployment during SAS typically consisted of ~8 sampling periods (from 2 to 10). The mean of the fluxes measured within each deployment is used in the subsequent analysis, with the standard error of fluxes indicating the variability within each deployment. Fluxes were converted to $\text{mmol m}^{-2} \text{day}^{-1}$ using the chamber surface area, the volume of the total chamber system (chamber, tubing, and analyser measurement cell), chamber air pressure and the ideal gas law.

Deleted: and air-ice

Deleted: both

Deleted: and on ice surfaces immediately following removal of covering snow...

Formatted: Superscript

Deleted: or ice

Deleted: eddy covariance (

Deleted:)

Deleted: of the EC system

Formatted: Subscript

Deleted: , due primarily to the

Deleted: they

Moved (insertion) [2]

Deleted: an

345 2.2 Water sampling

Onboard *Oden*, an underway seawater intake system continuously pumped water from a depth of approximately 8 m. Water temperature at 8 m depth was measured close to the underway line inlet with 2 hull contact sensors, and the temperature and salinity of the underway line water was measured using a Seabird SBE45 thermosalinograph (TSG) located in *Oden's* main lab. Downstream of the TSG on the underway line, a [Pro Oceanus](#) CO₂-Pro CV membrane equilibration sensor measures equilibrated CO₂ mixing ratio from which pCO_{2w} was calculated. This sensor was factory-calibrated prior to the expedition and performed an automated zeroing procedure every 6 hours. The manufacturer states the accuracy to be ± 3 ppm.

Surface water was sampled during the expedition to determine dissolved gas concentrations and carbonate system variables. Water samples were taken using bottles submerged at depths of 0-10 cm and using syringes at depths of 0-5 cm. During some stations a Ruttner sampler was used to obtain water samples at depths from 0.5 to 2.5 m. During underway chamber flux sampling, water samples were obtained using a lowered bucket, with the resulting water depth range for bottle and syringe samples corresponding approximately to the height of the bucket (0-30 cm). Temperature and practical salinity during each sampling were measured with a WTW 340i conductivity probe. [Discrete sampling at depth and from the underway line followed the washing and repeat overflow protocols described in Dickson et al. \(2007\). For surface water bottle sampling, the bottles were first washed with seawater then submerged by hand gradually while lying sideways to minimise air-water mixing as it was not possible to follow standard overfilling protocols \(Dickson et al., 2007\). Syringe sampling is not described in the carbon system sampling protocols, but follows Bastviken et al. \(2003\). The syringes were repeatedly washed with seawater prior to sampling.](#)

Samples of dissolved inorganic carbon (DIC) and total alkalinity (TA) were collected in Pyrex® borosilicate bottles (250 ml) according to Dickson et al. (2007), without being poisoned with mercury chloride (HgCl₂). Samples were stored short-term ([typically less than 12 hours](#), +4°C and dark) and thermostated to 25°C in a water bath prior to analysis. DIC was determined using a coulometric titration method based on Johnson et al. (1987) with a modified Single Operator Multiparameter Metabolic Analyzer (SOMMA) system (coulometer type UIC 5012) and the mean difference and standard deviation of duplicate sample analysis were $5.0 \pm 2.5 \mu\text{mol kg}^{-1}$ (n=12). TA was determined using a semi-open cell potentiometric (Orion ROSS 8102BN) titration (Metrohm Dosimat 665) method using a 5-point Gran evaluation (Haraldsson et al., 1997) and the mean difference and standard deviation of duplicate sample analysis were $3.4 \pm 2.5 \mu\text{mol kg}^{-1}$. The accuracy in DIC and TA was ensured by routine analysis of certified reference material (CRM Batch #181 and #191) obtained from A. G. Dickson of Scripps Institution of Oceanography (La Jolla, CA, USA). The pCO_{2w} was calculated from measured DIC and TA using the CO2SYSv1.1 MATLAB toolbox (Lewis and Wallace, 1998; van Heuven et al., 2011) and the dissociation constants of Lueker et al. (2000).

Samples for CH₄ analysis from surface leads, buckets, CTD and the pumped underway line were collected into 500 ml borosilicate bottles poisoned with 100 μl of saturated HgCl₂. Bottles were allowed to overfill with the bottle volume three times to remove gas bubbles, then transferred to the laboratory. Prior to analysis, samples were placed into a water bath at

Deleted:

Deleted: -

Surface buoyancy flux into lead waters was determined following MacIntyre et al. (2009) using the net radiative fluxes and turbulent heat fluxes made onboard *Oden* (Vüllers et al., 2021) and water surface temperature and salinity. Turbulent fluxes were gap-filled using the bulk estimates (Smith, 1988).

2.5 Gas transfer velocity calculation

The air-sea flux, F_X , of a poorly soluble gas species X , such as CO₂ and CH₄, can be represented as (e.g., Wanninkhof et al., 2009; Fairall et al., 2022):

$$F_X = kK_{0X}\Delta f_X, \quad (1)$$

where K_{0X} is the aqueous phase solubility of X (mol m⁻³ atm⁻¹) and Δf_X is the difference in the fugacity of X between the surface water and air. Fugacity is partial pressure corrected for non-ideality, this correction is very small (< 1%) for CO₂ and CH₄ in summertime Arctic conditions (McGillis et al., 2006), and partial pressures are used in all calculations reported here. The gas transfer velocity, k , represents the kinetic forcing of the flux, dependent on both molecular diffusivity, D , water viscosity, ν , and processes that impact transfer across the water surface interfacial layer. To account for the dependence of diffusivity on gas species, temperature and salinity, k is commonly normalised in terms of the non-dimensional Schmidt number Sc ($Sc = \nu/D$):

$$k_{660} = k, \quad (2)$$

here 660 is the Schmidt number of CO₂ in seawater at 20°C and the exponent n depends on the hydrodynamics of the interfacial layer, and is commonly chosen to be 0.5 for a wavy surface (e.g., Jähne et al., 1987).

We determine k_{660} from Eqs. (1) and (2) using measurements of F_X from the floating chamber, measured partial pressure differences, and solubilities and Sc calculated from the polynomial relationships summarised in Wanninkhof (2014), with the solubility relationships for CO₂ and CH₄ originally determined by Weiss (1974) and Wiesenburg and Guinasso (1979) respectively. Uncertainties in k_{660} are determined from combination of the flux and surface water partial pressure standard errors. Where the partial pressure comprised only one sample and so had no error value, the mean partial pressure standard error for that species is used instead.

3. Results

Unless otherwise stated, data is presented as means and standard deviations. The air-water fluxes and coincident supporting measurements are shown in Supplementary Material Table S1, and air-ice and air-snow fluxes are shown in Table S2. The uncertainties of the flux measurements shown in the Tables and Figures are the standard error of the samples within each

Deleted: .

Deleted: T

Deleted: directly from the flux measurement standard error.

Deleted: Parameterisation of k is commonly made in terms of wind speed, the dominant source of surface mixing energy in the open ocean. Chamber flux measurement necessarily involves isolation of an area of water from the wind. With appropriate flux chamber design (i.e., relatively small and lightweight chambers) it is assumed that the wind-induced interfacial layer mixing, the controlling factor for air-sea gas transfer of poorly soluble species, is advected within the chamber with minimal modification of the mixing by the chamber. This is further discussed in section 4.2.

Deleted: Stated flux uncertainties are the standard error of the flux measurements within each chamber deployment.

Deleted: s

Deleted: m

chamber deployment, and the uncertainties of the surface partial pressures in Table S1 and the Figures are the standard error of the samples comprising each measurement.

3.1 Meteorological and seawater conditions

480 Average wind speeds at 10 m, U_{10} , during SAS were $5.4 \pm 2.4 \text{ m s}^{-1}$ (Figs. 2b, 3a). The winds speeds corresponding to the air-water flux measurements were representative of the expedition as a whole with an average of $5.1 \pm 2.7 \text{ m s}^{-1}$. Wind speeds were generally moderate, with $\sim 80\%$ of the U_{10} measurements between 2 and 6.5 m s^{-1} (Fig. 3a). The highest winds occurred early in the expedition, on August 8 and 9, with 20-minute average U_{10} reaching 13.4 m s^{-1} . The wind distribution is similar to that observed on previous summertime CAO measurement campaigns, though without the passage of frontal systems and the associated higher winds that occurred during some campaigns (e.g., Tjernström et al., 2012; Vüllers et al., 2021). Sea ice concentration was generally high, with 90% of the expedition occurring in SIC $> 75\%$, and 60% of the expedition in SIC $> 95\%$ (Fig. 3b). The notable exception is a period from September 3 to September 7 when Oden was operating in a region of mixed sea ice and open water north of Greenland (Fig. 2c, Fig. 1).

490 Surface water temperature in leads sampled during flux measurements were $-1.4 \pm 0.5 \text{ }^\circ\text{C}$ (Fig. 4a), and the salinity of those samples showed high variation, ranging from 1.2 to 32.9, with an average 26.3 ± 8.5 (Fig. 4b). Surface water temperatures were generally at or very close to the seawater freezing point. Salinity lower than 22 occurred when ice movement caused a release of fresh meltpond water into leads, resulting in strong near-surface gradients. The temperature and salinity measured from the 8 m depth intake showed less variation, $-1.2 \pm 0.2 \text{ }^\circ\text{C}$ and 30.8 ± 1.2 , respectively. Surface temperatures determined by infrared sensors, including both ice and water surfaces, were $-2.0 \pm 1.7 \text{ }^\circ\text{C}$ over the course of the expedition. 495 Approximately 20% of the expedition occurred with surface temperatures below $-3 \text{ }^\circ\text{C}$, and 30% with surface temperatures above $-1 \text{ }^\circ\text{C}$ (Fig. 3c).

Throughout the expedition, intermittent periods with falling surface temperatures occurred, associated with clear sky conditions and surface longwave cooling (e.g., Prytherch and Yelland, 2021). From August 25 there was a regime shift, with protracted periods of surface (ice) temperature below the seawater freezing point, and deeper intermittent surface temperature drops, falling to around $-10 \text{ }^\circ\text{C}$, indicating the beginning of the autumn freeze up (Supplementary Material Section 2 and Fig. 500 S3).

Deleted: 2

3.2 CH₄ and CO₂ partial pressures

Atmospheric CH₄ partial pressure, pCH_4a , measured using the chamber LGR analyser during equilibration periods, increased from approximately $1.98 \text{ } \mu\text{atm}$ to $2.01 \text{ } \mu\text{atm}$ over the course of the expedition (mean and standard deviation $1.99 \pm 0.02 \text{ } \mu\text{atm}$; Fig. 4c). The surface seawater pCH_4w was determined as the average of two GC-based sampling methods (Section 2.2), with the two methods having a mean standard error of $2.82 \pm 1.54 \text{ } \mu\text{atm}$ with the syringe-based samples consistently higher (Supplemental Materials Section 1; Figure S2a). The averaged surface seawater pCH_4w measurements were slightly

oversaturated throughout the expedition, $5.37 \pm 3.13 \mu\text{atm}$ (Fig. 4c and Table S1). Oversight samples were slightly higher and more variable ($5.88 \pm 4.32 \mu\text{atm}$) than ice-based samples ($5.05 \pm 2.26 \mu\text{atm}$), with the highest oversight sample on August 29 at $16.0 \mu\text{atm}$. The pCH_4 sampled from 8 m depth was lower and less variable than the surface measurements, with an average of $2.70 \pm 0.68 \mu\text{atm}$. The 8 m pCH_4 was closer to equilibrium than the surface waters, but still slightly oversaturated, most notably from August 18 to September 04, where pCH_4 averaged $3.1 \pm 0.8 \mu\text{atm}$.

Near surface gradients of CH_4 saturation determined from Ruttner bottle sampling and on-ship analysis during the latter part of the expedition showed little variation with depth over the upper 2.5 m (Fig. 5). A pronounced freshening of the near-surface water was observed on September 3 and 8, and to a lesser extent on August 27. This is likely to be either a residual layer from ice melt or due to leakage of fresh meltpond water resulting from ice movement caused by *Oden*'s nearby passing.

Atmospheric CO_2 partial pressure, $p\text{CO}_2a$, from chamber equilibration periods was $402.4 \pm 3.0 \mu\text{atm}$ (Fig. 4d). The surface seawater $p\text{CO}_2w$ was determined as the average of GC-based syringe samples and bottle samples analysed for DIC and TA from which $p\text{CO}_2$ was calculated (Section 2.2). The two methods had a mean standard error of $25.5 \pm 22 \mu\text{atm}$ (Supplemental Materials Section 1; Figure S2b). The resulting averaged surface seawater $p\text{CO}_2w$ was always undersaturated, with a mean value of $227 \pm 71 \mu\text{atm}$ (Fig. 4d and Table S1). The $p\text{CO}_2w$ from oversight samples was higher and more variable ($288 \pm 69 \mu\text{atm}$) than that from the ice-based samples ($189 \pm 39 \mu\text{atm}$). The seawater $p\text{CO}_2w$ from 8 m depth was closer to equilibrium but still strongly undersaturated during most of the expedition, on average $315 \pm 28 \mu\text{atm}$. There was a strong gradient in $p\text{CO}_2w$ between the surface measurements and those from the 8 m intake from August 14 to the end of the expedition (Fig. 4d). Both depths were undersaturated throughout the expedition with the surface $p\text{CO}_2w$ on average $88 \mu\text{atm}$ lower than at 8 m depth. The period prior to August 13 had noticeably higher surface water $p\text{CO}_2w$ than later periods, close to or above the 8 m depth $p\text{CO}_2w$. The surface measurements in this early period were from both oversight and ice-based sampling, suggesting this relatively reduced or reversed $p\text{CO}_2w$ gradient was not primarily an artefact of *Oden*'s influence on the upper water column.

3.3 Lead air-sea fluxes

All measured air-sea CH_4 fluxes were positive (emission from the water to the atmosphere), in the direction of the air-sea concentration gradient, and the majority of the fluxes were less than $5 \mu\text{mol m}^{-2} \text{day}^{-1}$ (Fig. 6a and Table S1). The two highest fluxes, on August 11 and August 29, were coincident with the strongest air-sea $p\text{CH}_4$ differences (Fig. 4c). The average air-sea CH_4 flux was $3.5 \pm 4.4 \mu\text{mol m}^{-2} \text{day}^{-1}$. The oversight flux measurements were on average higher and more variable ($6.7 \pm 6.0 \mu\text{mol m}^{-2} \text{day}^{-1}$) than the ice-based measurements ($1.5 \pm 1.1 \mu\text{mol m}^{-2} \text{day}^{-1}$).

Measured air-sea CO_2 fluxes, using both the LGR- and Li-COR 7200-based systems, were all negative, showing uptake of CO_2 by seawater. The flux was in the direction of the air-sea concentration gradient, with a range of -1.5 to $-20.7 \text{ mmol m}^{-2} \text{day}^{-1}$ (Fig. 6a and Table S1). The average air-sea CO_2 flux was $-7.3 \pm 5.7 \text{ mmol m}^{-2} \text{day}^{-1}$. As for the CH_4 fluxes,

- Deleted: 6
- Deleted: 21
- Deleted: 71
- Deleted: 7.12
- Deleted: 5
- Deleted: 21
- Deleted: 6
- Deleted: 68
- Deleted: 96
- Deleted: 71
- Deleted: slightly oversaturated
- Deleted: ; however,
- Deleted: from
- Deleted: the 8 m pCH_4 was slightly oversaturated
- Deleted: .
- Formatted: Font: Not Italic
- Deleted: Measurements of CH_4 in ice cores were generally undersaturated, but water sampled immediately below the ice was often oversaturated, with an average saturation of 141% (range 67% to 210%) from the 9 coring locations sampled between August 27 and September 11....
- Formatted: Subscript
- Formatted: Font: Italic
- Formatted: Font: Italic, Subscript
- Deleted: from all sampling
- Deleted: n
- Deleted: average

the average magnitude of the overside CO₂ flux measurements was higher and more variable ($-12.7 \pm 7.3 \text{ mmol m}^{-2} \text{ day}^{-1}$) than the ice-based measurements ($-4.8 \pm 2.0 \text{ mmol m}^{-2} \text{ day}^{-1}$).

565 The largest CO₂ fluxes occurred in the early part of the expedition, prior to August 13, during overside sampling and in a period with notably smaller air-sea $p\text{CO}_2$ gradient than later. The large overside flux measurements on August 8 occurred during high winds, but other high overside measurements on August 10 and 11 occurred during light winds $< 4 \text{ m s}^{-1}$ (Fig. 2b). These measurements may be biased by the presence of *Oden*, or the interaction between *Oden* and the floating chamber, affecting the near-surface turbulence that drives the flux. These biases likely depend on the interaction of conditions such as the wind speed and direction relative to *Oden*, the location and topology of sea ice, and the position of the chamber relative to *Oden* and to the sea ice, all of which varied from one sampling to another. As such, all overside flux measurements are treated separately to the ice-based measurements.

575 Thin, grease or frazil ice was present at times during the expedition, especially during the latter part of the expedition, from August 26, coinciding with the end of the melt season and the onset of the freeze up. Ice-based sampling when grease ice was present resulted in lower fluxes on average of CH₄ ($0.9 \pm 0.6 \text{ } \mu\text{mol m}^{-2} \text{ day}^{-1}$), indicating that grease ice may impede gas exchange, presumably through the ice layer presenting both a physical barrier and reducing the action of wind on the surface water. However, the presence of grease ice did not change the average flux of CO₂, and the mean reduction of CH₄ flux is within the standard deviation. The presence of the chamber may modify the effect of the grease ice on gas exchange in an unquantified way, trapping ice inside or outside the chamber, and modifying the radiative balance at the surface, affecting ice formation.

Deleted: Grease ice impedes gas exchange, and i

Deleted: .

Deleted: For CO₂,

Deleted: .

3.4 Gas transfer velocities

Gas transfer velocities, k_{660} , determined directly from air-sea chamber flux measurements using Eqs. (1) and (2) were generally less than 10 cm hr^{-1} , with the notable exception of k_{660} determined from overside CO₂ flux measurements prior to August 13 when winds and fluxes were high (Fig. 6b). For all measurements, the average k_{660} was $6.7 \pm 9.7 \text{ cm hr}^{-1}$ (median 3.3 cm hr^{-1}). The average CH₄-flux derived k_{660} was $4.2 \pm 4.6 \text{ cm hr}^{-1}$ while the average CO₂-flux derived k_{660} was $8.0 \pm 11.4 \text{ cm hr}^{-1}$. The distribution of k_{660} is non-Gaussian and the median values from both gases combined, and for CH₄ and CO₂ separately, are similar: 3.3, 3.8 and 3.1 cm hr^{-1} respectively. The ice-based measurements were lower and had less scatter, the median k_{660} of both gases was 2.5 cm hr^{-1} (average $3.0 \pm 3.4 \text{ cm hr}^{-1}$). Many of the lower k_{660} measurements were made in the presence of grease or frazil ice, with the median k_{660} of grease-ice affected, ice-based samples of both gases being 1.1 cm hr^{-1} (average $1.4 \pm 0.9 \text{ cm hr}^{-1}$).

Deleted: c

Deleted: c

Deleted: quadratic

Deleted: 188

Deleted: 1.932

Deleted: 18

Deleted: This fit is very close to the parameterisation previously determined from eddy covariance measurements of CO₂ exchange in a large CAO lead (Prytherch and Yelland, 2021).

590 The wind speed dependence of the measurements, determined with non-linear least squares fit of the form $y = a \times x^b$, was approximately cubic, though the correlation with the data was weak ($a = 0.02$, $b = 2.9$, $r^2 = 0.31$; Fig. 7). The two measurements with greatest uncertainty (standard errors of 289 and 45 cm hr^{-1}) were excluded from the fit. The relatively few data points (36) and the weakness of the correlation means that caution is needed in interpretation or application of this relationship. At low wind speeds ($< 3 \text{ m s}^{-1}$), the measurements are higher than previously determined power law

610 parameterisations, suggesting that processes other than wind speed may be contributing to interfacial mixing. With the exception of one value, these low wind k_{660} are relatively small, similar to the low-wind values of previous parameterisations determined with non-zero intercepts (e.g., Wanninkhof et al., 2009).

The overside CO₂ flux derived k_{660} were higher and more scattered than the other measurements and may have been affected by their close proximity to *Oden's* hull and the additional turbulence that this may induce in the ocean. The wind speed dependence of the fit determined to only ice-based measurements (again excluding the measurement with highest standard error) was weaker and had a weaker correlation ($a = 0.999$, $b = 0.533$, $r^2 = 0.13$). The ice-based measurements would not have been affected by any such bias, and the wind speed dependence of ice-based CO₂- and CH₄-derived k_{660} was weaker, close to linear, and generally below previous open ocean and lead water k_{660} parameterisations. The contribution to the observed k_{660} wind speed dependence of forcings particular to the sea-ice environment are discussed in Section 4.2.

615 The measurements were also compared with the parametric Wave Age Gas Transfer (WAGT) model of gas exchange in sea ice regions (Bigdeli et al., 2018). The WAGT model was run in both SIC mode, with SIC set to 90%, and in fetch mode using the lead width measurements. The WAGT fetch mode has good agreement with the measurements at wind speeds below 6 m s⁻¹ whereas the SIC mode overpredicts. Both WAGT modes over predict at higher wind speeds.

3.5 Air-snow fluxes

625 Chamber measurements of air-snow CO₂ flux showed consistent small fluxes, on average, -1.1 ± 1.2 mmol m⁻² day⁻¹, with the largest magnitude flux -2.9 mmol m⁻² day⁻¹ (Fig. 6c and Table S2). No clear dependence of the atmospheric flux on surface temperature was found, with the largest magnitude fluxes occurring at temperatures close to and above freezing, but some relatively large fluxes (up to -2.8 mmol m⁻² day⁻¹) also being observed during the coldest measured surface conditions, with temperatures $< -2^\circ\text{C}$ (Fig. 8).

4. Discussion

630 4.1 Near-surface concentration gradients

Strong near-surface CO₂ gradients are often observed in Arctic waters. Lower CO₂ partial pressure at the surface relative to 8 m (Fig. 4d) may result from the input of unsaturated water from melting sea ice, from differences in biological activity, and from the increased solubility resulting from the generally lower temperature and fresher surface waters (Figs. 4a, b). Solubility differences between the surface and 8 m in the measurements reported here on average account for a $p\text{CO}_{2W}$ difference of 10 μatm , or 11% of the total average difference between the depths. The fresher surface layer observed on several days (August 27, September 3 and 8; Fig. 5) indicates a stratification that could act to suppress near-surface mixing, leading to lower surface gas concentrations and thus lower fluxes. In the coarse vertical resolution observations reported here, near-surface CH₄

Deleted: 23

Deleted: 706

Deleted: 07

Deleted: ,

Deleted: and chamber flux measurement biases,

Deleted: s

Deleted: /ice

Deleted: In contrast, measurements of air-ice CO₂ flux on ice surfaces following the removal of the snow cover were more variable (mean -0.5 ± 2.2 mmol m⁻² day⁻¹), with the largest magnitude flux of -6.3 mmol m⁻² day⁻¹ and positive, emission fluxes of up to 1.7 mmol m⁻² day⁻¹.

Deleted: for either the snow or ice surface

Deleted: air-snow

Deleted: and air-ice up to -1.5 mmol m⁻² day⁻¹

Moved (insertion) [5]

Deleted: The higher $p\text{CH}_{4W}$ at the surface relative to at 8 m depth (Fig. 4c), together with the higher $p\text{CH}_{4W}$ observed immediately under the ice indicates a near-surface source of CH₄. This supports the hypothesised role of sea ice in transporting CH₄ from productive shallow sea areas and/or providing an environment for the local generation of CH₄ (Damm et al., 2018).

660 saturations decrease in the presence of lower salinity, suggesting that in these cases the greater solubility in fresher water compensates for any stratification effect.

The results here further demonstrate the concentration gradients that can be present between the surface and the sampling depths used for the majority of the reported ocean 'surface' gas concentration measurement (e.g., Miller et al., 2019). Larger air-sea concentration differences will, all else being equal, cause a larger air-sea flux, and thus any gradient between the surface and the water sampling depth will bias both bulk-method fluxes (i.e., fluxes calculated using Eq. (1) and a parameterisation of k ; Wanninkhof et al., 2009) and gas transfer velocity measurements. As such gradients are common, and are not easy to determine or correct for, many parameterisations of gas transfer incorporate such biases.

If the surface partial pressure, T and S measurements from this study are used to determine bulk-method fluxes, the CO_2 fluxes are on average 77% (with the $k-U_{10}$ relationship of Ho et al., 2006) or 25% (with the $k-U_{10}$ relationship of Prytherch and Yelland, 2021) higher than the measured chamber fluxes (Table 1). In contrast, if the 8 m depth partial pressure, T and S measurements are used instead, the bulk fluxes are 1% (Ho et al., 2006) and 30% lower (Prytherch and Yelland, 2021) than the measured fluxes. For CH_4 the effects of the gradients are similar (Table 1), with bulk fluxes derived from surface measurements being larger than the chamber fluxes by 156% and 80% (for the Ho et al., 2006, and Prytherch and Yelland, 2021 $k-U_{10}$ relationships, respectively) and the bulk fluxes derived from 8 m measurements being smaller than the chamber flux measurements by 28% and 49% respectively.

675 4.2 Gas transfer in the presence of sea ice

Greater lead width and fetch enables greater wind-driven mixing, while measurements close to ice edges may also be affected by the interaction of wind and waves with the ice edge. The wind speed dependence of all (36) measurements (Fig. 7) is approximately cubic with a weak correlation (r^2 0.31). The k_{660} measurements, normalised by U_{10} , do not show a clear dependence on lead width as determined by laser ranging (Fig. 9). For overside measurements lead width was the distance from Oden's hull to the ice edge. In contrast, the U_{10} normalised ice-based measurements do have a dependence on lead width; the correlation of a non-linear least squares fit of the form $y = a \times x^b$ is weak ($a = 0.208$, $b = 0.34$, $r^2 = 0.27$) but stronger than the fit of these measurements to wind speed. Properly accounting for such fetch effects would require determination of lead dimensions, relative wind direction and ice freeboard, but the relatively small variation of the ice-based measurements with width suggests that simpler approaches to parameterising k in sea-ice regions, with either a wind speed dependence or a constant value, may be sufficient.

The buoyancy flux measurements enable identification of measurements made in conditions of surface cooling-induced convection, which can enhance gas exchange (McGillis et al., 2004), but no clear enhancement of gas transfer was apparent (Fig. 7). Many of the convective measurements were also in leads of < 10 m width (Figs. 7, 9) complicating determination of the impact of either convection or fetch on gas exchange. Many, but not all, of the lower gas exchange rates were measured when grease ice was present on the lead water, even at higher wind speeds (> 7 m s^{-1}) when low lead widths

Moved up [5]: Strong near-surface CO_2 gradients are often observed in Arctic waters. Using EC flux measurements, Dong et al. (2021) showed that surface $f\text{CO}_2$ in the summertime Arctic marginal ice zone was $39 \mu\text{atm}$ lower than that measured from their ship's intake at 6 m depth. They determined that this was partly due to effects on CO_2 solubility from meltwater cooling and freshening, with approximately half of the reduction from other factors, presumably photosynthesis. Miller et al. (2019) report $p\text{CO}_{2W}$ differences between the surface and an intake at 7 m depth of between -180 and $+160 \mu\text{atm}$ during sampling in the Canadian Arctic Archipelago and Hudson Bay. The authors note that "The temperature differences between the underway system and the shallower samples were often contrary to the $p\text{CO}_2$ gradients..., indicating that $p\text{CO}_2$ was not simply controlled by surface heating and cooling". Due to surface longwave emission and the resulting cool skin effect (Woolf et al., 2016), temperature differences are always present between the sea surface and intake depths even when the upper ocean is well mixed. Solubility differences between the surface and 8 m in the measurements reported here on average account for a $p\text{CO}_{2W}$ difference of $10 \mu\text{atm}$, or 11% of the total average difference between the depths. The majority of the observed undersaturation is therefore likely to be due to biological activity.

Moved up [3]: Using EC flux measurements, Dong et al. (2021) showed that surface $f\text{CO}_2$ in the summertime Arctic marginal ice zone was $39 \mu\text{atm}$ lower than that measured from their ship's intake at 6 m depth. They determined that this was partly due to effects on CO_2 solubility from meltwater cooling and freshening, with approximately half of the reduction from other factors, presumably photosynthesis. Miller et al. (2019) report $p\text{CO}_{2W}$ differences between the surface and an intake at 7 m depth of between -180 and $+160 \mu\text{atm}$ during sampling in the Canadian Arctic Archipelago and Hudson Bay. The authors note that "The temperature differences between the underway system and the shallower samples were often contrary to the $p\text{CO}_2$ gradients..., indicating that $p\text{CO}_2$ was not simply controlled by surface heating and cooling". Due to surface longwave emission and the resulting cool skin effect (Woolf et al.,

Deleted: Strong near-surface CO_2 gradients are often observed in observed in Arctic waters. Using EC flux measurements, Dong et al. (2021) showed that surface $f\text{CO}_2$ in the summertime Arctic ma (... [8]

Deleted: equation

Deleted: The high sensitivity and small footprint of the flux chamber technique makes it attractive for measurements in small water bodies such as leads. The technique has been frequently (... [9]

Deleted: close to

Deleted: quadratic

Deleted: and in close agreement with that from a large Arctic lead of ~ 100 - 400 m width (Prytherch and Yelland, 2021)

Deleted: ²

Deleted:

Deleted: 041

Deleted: 501

Deleted: 16

Deleted: are

Deleted: i

provided sheltered conditions (Fig. 7). Grease ice is expected to reduce gas transfer rates, and all high k_{660} measurements (> 5 cm hr^{-1}), both overside and ice-based, were obtained when grease ice was not present.

810 The WAGT) model was designed to use estimates of the area of open water from SIC products. Although different ice products have experienced significant improvements in increased resolution in recent years, they are still prone to missing critical small-scale features such as open or refrozen leads. Passive radiometer data also treats a ~20 cm thin ice cover as open water (e.g. Hoffman et al., 2019), with severe impacts on gas transfer estimates.

4.3 Regional fluxes

815 The fluxes and gas transfer measurements presented here provide direct measurement-based constraints for the role of the CAO in CO_2 and CH_4 atmospheric exchange. Air-sea chamber fluxes were made at latitudes from approximately 82.5°N to 90°N . These latitudes bound an area of $2.2 \times 10^6 \text{ km}^2$, approximately corresponding to the deep-water areas of the Amundsen basin. The IHO-defined central Arctic deep basin ($4.7 \times 10^6 \text{ km}^2$), and the bathymetrically-defined $4.5 \times 10^6 \text{ km}^2$ “Central Arctic Ocean Basin” delineated by Jakobsson (2002) and used here as the definition of the CAO are larger. In all definitions, 820 the CAO corresponds to the Arctic Ocean deep water > 2400 m depth, as a strong contrast to the shallow shelf seas (average depths ~ 50 -250 m) which surround the Arctic Ocean. The average SIC during this portion of the expedition was 90%, determined from AMSR-2 6.25 km^2 observations using the ASI algorithm (Sprenn et al., 2008).

The measured air-sea CO_2 fluxes are of similar magnitude to the EC measurements of Prytherch et al. (2017) (average $-7.8 \text{ mmol m}^{-2} \text{ day}^{-1}$), Prytherch and Yelland (2021) (0 to $-20 \text{ mmol m}^{-2} \text{ day}^{-1}$, average $-12.4 \text{ mmol m}^{-2} \text{ day}^{-1}$, measured in 825 higher winds), and of Dong et al. (2021), (0 to $-40 \text{ mmol m}^{-2} \text{ day}^{-1}$, measured in SIC $> 60\%$). Averaging observations regionally, the mean chamber-derived CO_2 air-sea fluxes of $-7.6 \text{ mmol m}^{-2} \text{ day}^{-1}$ correspond to SIC-accounted CAO fluxes of $-3.42 \times 10^9 \text{ mol day}^{-1}$, or $-150.5 \text{ Gg CO}_2 \text{ day}^{-1}$. The flux measurements reported here were made during the summer and the beginning of the autumn freeze up in conditions of persistent, strong $p\text{CO}_2$ undersaturation. The annual cycle of Arctic Ocean surface $p\text{CO}_2$ is poorly constrained, primarily due to the rarity of reported observations outside of summer months (e.g. Yasunaka et al., 2018; Else et al., 2012). As such, while we can estimate an annual CAO air-sea flux by averaging the fluxes and SIC (90%), 830 the year-round flux of $-55.0 \text{ Tg CO}_2 \text{ yr}^{-1}$ is a very uncertain estimate. Furthermore, while density stratification from sea-ice melt in summer months may suppress upper ocean mixing, leading to reduced air-sea concentration gradients and thus reduced summer fluxes relative to other seasons, this stratification was not evident in the measurements reported here. Thus, due to the higher SIC and lower primary productivity in winter months, our annual flux estimate is likely to represent a high estimate. If 835 the average bulk flux of $-8.4 \text{ mmol m}^{-2} \text{ day}^{-1}$, as determined from near-surface CO_2 concentrations and the k relationship of Prytherch and Yelland (2021), is used instead of the average observed flux, the CAO air-sea flux is slightly larger, $-3.78 \times 10^9 \text{ mol day}^{-1}$, or $-166.4 \text{ Gg CO}_2 \text{ day}^{-1}$.

The measured air-snow CO_2 fluxes ($-1.1 \pm 1.2 \text{ mmol m}^{-2} \text{ day}^{-1}$) are a similar magnitude to previously reported EC-determined air-snow fluxes from the summertime CAO (Prytherch and Yelland, 2021; $-0.7 \text{ mmol m}^{-2} \text{ day}^{-1}$, maximum -6.5 840 $\text{mmol m}^{-2} \text{ day}^{-1}$). They also span a similar range (-2.9 to $0.25 \text{ mmol m}^{-2} \text{ day}^{-1}$), including the observed positive emission fluxes,

Moved down [4]: 5. Regional fluxes and conclusions¶

Deleted: → While the observed gas transfer velocities had a similar wind speed dependence to that of Prytherch and Yelland (2021), the dependence of the ice-based measurements, free from any influence of *Oden* on the exchange rate, was much weaker. In pack ice regions where water surfaces are small, the results reported here show that gas exchange can be appropriately represented with a constant k_{660} , 2.5 cm hr^{-1} . In areas such as the marginal ice zone MIZ with mixed sea ice and water and larger fetches, gas exchange should be represented using a wind-speed dependent parameterisation such as Prytherch and Yelland (2021).¶

Deleted:

Deleted: Although the

Deleted: only

Deleted: , we can estimate an approximate annual CAO air-sea flux by averaging the fluxes and SIC (90%) year-round to yield $-55.0 \text{ Tg CO}_2 \text{ yr}^{-1}$.

Deleted: D

Deleted: ,

Deleted: . However,

Deleted: hat

Deleted: ,

Deleted: and

Deleted: limit due to the higher SIC and lower primary productivity in winter months

Deleted: ,

Deleted: ,

Deleted: annual

Deleted: -

Deleted: 60.1 Tg yr^{-1} .

Deleted: $\text{mmol m}^{-2} \text{ day}^{-1}$

Deleted: ,

Deleted: upwards

875 as previously reported air-snow chamber flux measurements (Geilfus et al., 2012, Nomura et al., 2010, 2013, 2018, Delille et al., 2014, Geilfus et al., 2015). However, none of these previously reported air-snow flux measurements are from the summertime or autumn CAO. Using the observed expedition SIC and the average flux through snow-covered sea ice of $-1.1 \text{ mmol m}^{-2} \text{ day}^{-1}$, yields a CAO regional air-snow flux of $-4.46 \times 10^9 \text{ mol day}^{-1}$, or $-196.1 \text{ Gg day}^{-1}$. Combining the air-sea and air-snow chamber flux estimates with constant SIC (90%), the average atmospheric CO_2 flux for the CAO is $-1.75 \text{ mmol m}^{-2} \text{ day}^{-1}$.

880 The average air-sea CH_4 flux of $3.5 \mu\text{mol m}^{-2} \text{ day}^{-1}$, accounting for SIC, equates to a CAO flux of $1.58 \times 10^6 \text{ mol day}^{-1}$, or 25.3 Mg day^{-1} . Similarly, the average bulk CH_4 flux of $5 \mu\text{mol m}^{-2} \text{ day}^{-1}$ derived from near-surface CH_4 concentrations and the k relationship of Prytherch and Yelland (2021) corresponds to 36.1 Mg day^{-1} for the CAO area. The annual cycle of Arctic Ocean surface CH_4 is much less well known than for CO_2 , in particular the origin of the dissolved CH_4 driving the observed fluxes. As such the extrapolation of our observations and bulk estimates to annual fluxes (9.2 Gg yr^{-1} and 13.2 Gg yr^{-1} respectively) is also highly uncertain. Both the directly observed and bulk flux-derived annual estimates are likely to be high estimates due to the higher SIC and minimal release of CH_4 from within sea ice during winter.

Stratification of near surface waters by freshening from melt water may reduce air-sea gas flux by both reducing the turbulent mixing below the interfacial layer, dampening the gas transfer velocity, and by impeding the replenishment of the equilibrating surface waters from the bulk water below, reducing the concentration difference that drives the exchange. 890 Suppression of turbulent mixing may contribute to the observed difference between chamber and bulk fluxes, however, the surface layer freshwater stratification apparent in surface layers did not coincide with CH_4 stratification (Fig. 5).

Both the direct chamber CH_4 fluxes and bulk CH_4 fluxes are of similar magnitude to previously reported flux estimates derived from seawater CH_4 measurements in summer and winter at more southerly Arctic marine locations (e.g. Lorenson et al., 2016; Manning et al., 2020, 2022; Fenwick et al., 2017; Silyakova et al., 2022), but are approximately 35 times smaller than aircraft-based observations of fluxes from wintertime leads over the western CAO's Canada Basin, south of 82°N (Kort et al., 2012). This suggests a strong seasonal dependence in CAO CH_4 flux, or that the aircraft-based measurements of large fluxes represent episodic rather than typical widespread emissions. If we assume that the fluxes reported by Kort et al. are more typical of the Canada Basin and distinct from our primarily Amundsen basin study area, this suggests a much higher net annual CH_4 flux from the CAO of up to $\sim 0.1 \text{ Tg yr}^{-1}$. The CAO has a mean depth $> 2700 \text{ m}$ (Jakobsson, 2002), meaning any 900 CH_4 emissions must be from near-surface production or long-range transport of dissolved CH_4 ; seafloor sources are too deep to significantly affect the surface $p\text{CH}_4$ in the CAO.

5. Conclusions

The direct flux and gas transfer measurements presented in this study offer insights into the role of the CAO in atmospheric exchange of climate forcing trace gases, informing future climate modelling and carbon budget analysis. They provide a measurement-based constraint on the magnitude of the CAO CO_2 flux, with the observed CAO CO_2 flux in agreement with 905

Deleted: ,

Deleted: and extrapolated year-round, $-126.6 \text{ Tg yr}^{-1}$

Deleted: The direct flux and gas transfer measurements presented in this study offer insights into the role of the CAO in Arctic CO_2 atmospheric exchange. They provide a measurement-based constraint on the magnitude of the CAO CO_2 flux, with the observed CAO CO_2 flux in agreement with the observation-based estimates of Bates et al. (2006) and Yasunaka et al. (2018), and an order of magnitude greater than the coupled ocean-biogeochemistry model estimate of Manizza et al., 2019.

Formatted: Subscript

Formatted: Subscript

Deleted: Extrapolating to annual fluxes gives

Deleted: as

Deleted: a

Deleted: t

Deleted: †

Deleted: The total CH_4 flux to the atmosphere from the Arctic Ocean is dominated by the shallow coastal shelf emissions, where surface $p\text{CH}_4$ can be greatly enhanced by bubbles from shallow gas seeps reaching the atmosphere. Additionally, direct terrestrial riverine inputs of CH_4 into the shelf seas produce much higher $p\text{CH}_4$ near the coastline (Weber et al., 2019; Manning et al., 2020). While the integrated magnitude of these Arctic marine methane emissions remains under debate, it appears to presently be on the scale of $\sim 4\text{-}5 \text{ Tg CH}_4 \text{ yr}^{-1}$ or $100 \mu\text{mol m}^{-2} \text{ day}^{-1}$ (Thornton et al., 2016a; 2016b; 2020), $\sim 1\%$ of global CH_4 emissions (Saunois et al., 2020). The direct measurement-based constraints determined here thus show that the CAO is a very small additional contributor to the Arctic Ocean CH_4 flux in summer and likely also year-round, informing future climate modelling and carbon budget analysis.

Moved (insertion) [4]

Deleted: Regional fluxes and c

940 the observation-based estimates of Bates et al. (2006) and Yasunaka et al., (2018), and an order of magnitude greater than the coupled ocean-biogeochemistry model estimate of Manizza et al., 2019. For CH₄, the direct measurement-based constraints determined here show that the CAO is a very small additional contributor to the Arctic Ocean CH₄ flux in summer, with the mean CAO CH₄ flux of 25.3 Mg day⁻¹ comprising approximately 1/500th of the estimate of integrated Arctic marine CH₄ emissions of 4-5 Tg yr⁻¹ (Thornton et al, 2016a; 2016b; 2020) if that estimate is assumed evenly distributed over the year. Arctic marine CH₄ emissions are dominated by the shallow coastal shelf emissions.

945 While the observed gas transfer velocities had a similar, if weakly correlated, wind-speed dependence to that of Prytherch and Yelland (2021), the wind-speed dependence of the ice-based measurements, free from any influence of *Oden* on the exchange rate, was much weaker. The measurements reported here, from pack ice regions with small water surfaces, could be appropriately represented with a constant k_{660} , 2.5 cm hr⁻¹ instead of a relationship directly or indirectly (i.e. wave or turbulence-based) dependent on wind speed. In areas such as the marginal ice zone MIZ with mixed sea ice and water and larger fetches, the direct and indirect influence of wind on interfacial mixing will be greater and gas exchange should be represented using a parameterisation incorporating wind-speed such as Prytherch and Yelland (2021).

950 Author contributions

JP designed the experiment, JP and SM carried out the flux measurements and water sampling. JP, IB, VB, AU, AH, AN and LH analysed the water samples. JP, BT and MT developed the ship-based instrumentation, and JP and VB developed the chamber flux systems. JP prepared the manuscript with contributions from all co-authors.

Acknowledgements

955 The Swedish Polar Research Secretariat (SPRS) provided access to the icebreaker (*I/B*) *Oden* and logistical support. We would like to thank Capt. Mattias Petersson and the crew of *Oden* for their invaluable support throughout the field campaign. Thanks also to the Chief Scientist Pauline Snoeijts Leijonmalm for the planning and coordination of SAS2021.

Deleted: c

Financial support

960 JP and SM were supported by the Knut and Alice Wallenberg Foundation (grant no. 2016-0024). AU was supported by the Swedish Research Council Formas (grant no. 2018-01398) and the Hasselblad Foundation (grant no. 2019-1218). LAH was supported by the Arctic Research Icebreaker Consortium (ARICE, EU, <https://arice-h2020.eu>) (grant no. 730 965). This paper contributes to the science plan of the Surface Ocean-Lower Atmosphere Study (SOLAS), which is partially supported by the U.S. National Science Foundation (Grant OCE-1840868) via the Scientific Committee on Oceanic Research (SCOR).

965 **References**

- [Ahmed, M. M. M., Else, B. G. T., Capelle, D., Miller, L. A., and Papakyriakou, T.: Underestimation of surface pCO₂ and air-sea CO₂ fluxes due to freshwater stratification in an Arctic shelf sea, Hudson Bay, *Elementa: Science of the Anthropocene*, 8, <https://doi.org/10.1525/elementa.084>, 2020.](https://doi.org/10.1525/elementa.084)
- 970 [Bastviken, D., Ejlertsson, J., Sundh, I., and Tranvik, L.: METHANE AS A SOURCE OF CARBON AND ENERGY FOR LAKE PELAGIC FOOD WEBS, *Ecology*, 84, 969–981, \[https://doi.org/10.1890/0012-9658\\(2003\\)084\\[0969:maasoc\\]2.0.co;2\]\(https://doi.org/10.1890/0012-9658\(2003\)084\[0969:maasoc\]2.0.co;2\), 2003.](https://doi.org/10.1890/0012-9658(2003)084[0969:maasoc]2.0.co;2)
- Bates, N. R. and Mathis, J. T.: The Arctic Ocean marine carbon cycle: evaluation of air-sea CO₂ exchanges, ocean acidification impacts and potential feedbacks, *Biogeosciences*, 6, 2433–2459, <https://doi.org/10.5194/bg-6-2433-2009>, 2009.
- 975 Bates, N. R., Moran, S. B., Hansell, D. A., and Mathis, J. T.: An increasing CO₂ sink in the Arctic Ocean due to sea-ice loss, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006gl027028>, 2006.
- Bigdeli, A., Hara, T., Loose, B., and Nguyen, A. T.: Wave Attenuation and Gas Exchange Velocity in Marginal Sea Ice Zone, *J. Geophys. Res. Oceans*, 123, 2293–2304, <https://doi.org/10.1002/2017jc013380>, 2018.
- 980 Butterworth, B. J. and Miller, S. D.: Air-sea exchange of carbon dioxide in the Southern Ocean and Antarctic marginal ice zone, *Geophys. Res. Lett.*, 43, 7223–7230, <https://doi.org/10.1002/2016gl069581>, 2016.
- 985 Cole, J. J., Bade, D. L., Bastviken, D., Pace, M. L., and Van de Bogert, M.: Multiple approaches to estimating air-water gas exchange in small lakes, *Limnol. Oceanogr. Methods*, 8, 285–293, <https://doi.org/10.4319/lom.2010.8.285>, 2010.
- Damm, E., Rudels, B., Schauer, U., Mau, S., and Dieckmann, G.: Methane excess in Arctic surface water- triggered by sea ice formation and melting, *Sci Rep*, 5, <https://doi.org/10.1038/srep16179>, 2015.
- 990 Damm, E., Bauch, D., Krumpfen, T., Rabe, B., Korhonen, M., Vinogradova, E., and Uhlig, C.: The Transpolar Drift conveys methane from the Siberian Shelf to the central Arctic Ocean, *Sci Rep*, 8, <https://doi.org/10.1038/s41598-018-22801-z>, 2018.
- Delille, B., Vancoppenolle, M., Geilfus, N.-X., Tilbrook, B., Lannuzel, D., Schoemann, V., Becquevort, S., Carnat, G., Delille, D., Lancelot, C., Chou, L., Dieckmann, G. S., and Tison, J.-L.: Southern Ocean CO₂ sink: The contribution of the sea ice, *J. Geophys. Res. Oceans*, 119, 6340–6355, <https://doi.org/10.1002/2014jc009941>, 2014.
- 995

- Dickson, A.G., Sabine, C.L. and Christian, J.R. (Eds.). Guide to Best Practices for Ocean CO₂ Measurements. PICES Special Publication 3, 191 pp. 2007.
- 1000
- Dong, Y., Yang, M., Bakker, D. C. E., Liss, P. S., Kitidis, V., Brown, I., Chierici, M., Fransson, A., and Bell, T. G.: Near-Surface Stratification Due to Ice Melt Biases Arctic Air-Sea CO₂ Flux Estimates, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021gl095266>, 2021.
- 1005 [Else, B. G. T., Papakyriakou, T. N., Galley, R. J., Mucci, A., Gosselin, M., Miller, L. A., Shadwick, E. H., and Thomas, H.: Annual cycles of pCO₂sw in the southeastern Beaufort Sea: New understandings of air-sea CO₂ exchange in arctic polynya regions, *J. Geophys. Res.*, 117, <https://doi.org/10.1029/2011jc007346>, 2012.](#)
- Fairall, C. W., Yang, M., Brumer, S. E., Blomquist, B. W., Edson, J. B., Zappa, C. J., Bariteau, L., Pezoa, S., Bell, T. G., and Saltzman, E. S.: Air-Sea Trace Gas Fluxes: Direct and Indirect Measurements, *Front. Mar. Sci.*, 9, <https://doi.org/10.3389/fmars.2022.826606>, 2022.
- 1010
- Fanning, K. A. and Torres, L. M.: 222Rn and 226Ra: indicators of sea-ice effects on air-sea gas exchange, *Polar Research*, 10, 51–58, <https://doi.org/10.3402/polar.v10i1.6727>, 1991.
- 1015
- Fenwick, L., Capelle, D., Damm, E., Zimmermann, S., Williams, W. J., Vagle, S., and Tortell, P. D.: Methane and nitrous oxide distributions across the North American Arctic Ocean during summer, 2015, *J. Geophys. Res. Oceans*, 122, 390–412, <https://doi.org/10.1002/2016jc012493>, 2017.
- 1020
- Fransson, A., Chierici, M., Skjelvan, I., Olsen, A., Assmy, P., Peterson, A. K., Spreen, G., and Ward, B.: Effects of sea-ice and biogeochemical processes and storms on under-ice water f CO₂ during the winter-spring transition in the high AO: Implications for sea-air CO₂ fluxes, *J. Geophys. Res. Oceans*, 122, 5566–5587, <https://doi.org/10.1002/2016jc012478>, 2017.
- 1025
- Geilfus, N.-X., Carnat, G., Papakyriakou, T., Tison, J.-L., Else, B., Thomas, H., Shadwick, E., and Delille, B.: Dynamics of pCO₂ and related air-ice CO₂ fluxes in the Arctic coastal zone (Amundsen Gulf, Beaufort Sea), *J. Geophys. Res.*, 117, n/a-n/a, <https://doi.org/10.1029/2011jc007118>, 2012.
- 1030
- Geilfus, N.-X., Galley, R. J., Crabeck, O., Papakyriakou, T., Landy, J., Tison, J.-L., and Rysgaard, S.: Inorganic carbon dynamics of melt-pond-covered first-year sea ice in the Canadian Arctic, *Biogeosciences*, 12, 2047–2061, <https://doi.org/10.5194/bg-12-2047-2015>, 2015.

- Gålfalk, M., Bastviken, D., Fredriksson, S., and Arneborg, L.: Determination of the piston velocity for water-air interfaces using flux chambers, acoustic Doppler velocimetry, and IR imaging of the water surface, *J. Geophys. Res. Biogeosci.*, 118, 770–782, <https://doi.org/10.1002/jgrg.20064>, 2013.
- 1035 Haraldsson, C., Anderson, L. G., Hassellöv, M., Hulth, S., and Olsson, K.: Rapid, high-precision potentiometric titration of alkalinity in ocean and sediment pore waters, *Deep Sea Research Part I: Oceanographic Research Papers*, 44, 2031–2044, [https://doi.org/10.1016/s0967-0637\(97\)00088-5](https://doi.org/10.1016/s0967-0637(97)00088-5), 1997.
- 1040 Ho, D. T., Law, C. S., Smith, M. J., Schlosser, P., Harvey, M., and Hill, P.: Measurements of air-sea gas exchange at high wind speeds in the Southern Ocean: Implications for global parameterizations, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006gl026817>, 2006.
- 1045 [Hoffman, J., Ackerman, S., Liu, Y., and Key, J.: The Detection and Characterization of Arctic Sea Ice Leads with Satellite Imagers, *Remote Sensing*, 11, 521, <https://doi.org/10.3390/rs11050521>, 2019.](#)
- Van Heuven, S., Pierrot, D., Rae, J. W. B., Lewis, E., and Wallace, D. W. R.: MATLAB Program Developed for CO2 System Calculations. ORNL/CDIAC-105b., https://doi.org/10.3334/CDIAC/OTG.CO2SYS_MATLAB_V1.1, 2011.
- 1050 Jähne, B., Münnich, K. O., Börsinger, R., Dutzi, A., Huber, W., and Libner, P.: On the parameters influencing air-water gas exchange, *J. Geophys. Res.*, 92, 1937, <https://doi.org/10.1029/jc092ic02p01937>, 1987.
- Jakobsson, M.: Hypsometry and volume of the Arctic Ocean and its constituent seas, *Geochem.-Geophys.-Geosyst.*, 3, 1–18, <https://doi.org/10.1029/2001gc000302>, 2002.
- 1055 Johnson, K. M., Sieburth, J. M., Williams, P. J. leB., and Brändström, L.: Coulometric total carbon dioxide analysis for marine studies: Automation and calibration, *Marine Chemistry*, 21, 117–133, [https://doi.org/10.1016/0304-4203\(87\)90033-8](https://doi.org/10.1016/0304-4203(87)90033-8), 1987.
- 1060 Kort, E. A., Wofsy, S. C., Daube, B. C., Diao, M., Elkins, J. W., Gao, R. S., Hints, E. J., Hurst, D. F., Jimenez, R., Moore, F. L., Spackman, J. R., and Zondlo, M. A.: Atmospheric observations of Arctic Ocean methane emissions up to 82° north, *Nature Geosci.*, 5, 318–321, <https://doi.org/10.1038/ngeo1452>, 2012.
- Kwok, R.: Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018), *Environ. Res. Lett.*, 13, 105005, <https://doi.org/10.1088/1748-9326/aae3ec>, 2018.
- 1065

- Lewis, E., and D. W. R. Wallace. Program Developed for CO₂ System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. 1988.
- Loose, B., Schlosser, P., Perovich, D., Ringelberg, D., Ho, D. T., Takahashi, T., Richter-Menge, J., Reynolds, C. M., McGillis, W. R., and Tison, J.-L.: Gas diffusion through columnar laboratory sea ice: implications for mixed-layer ventilation of CO₂ in the seasonal ice zone, *Tellus B: Chemical and Physical Meteorology*, 63, 23, <https://doi.org/10.1111/j.1600-0889.2010.00506.x>, 2011.
- Loose, B., McGillis, W. R., Perovich, D., Zappa, C. J., and Schlosser, P.: A parameter model of gas exchange for the seasonal sea ice zone, *Ocean Sci.*, 10, 17–28, <https://doi.org/10.5194/os-10-17-2014>, 2014.
- Loose, B., Kelly, R. P., Bigdeli, A., Williams, W., Krishfield, R., Rutgers van der Loeff, M., and Moran, S. B.: How well does wind speed predict air-sea gas transfer in the sea ice zone? A synthesis of radon deficit profiles in the upper water column of Arctic Ocean, *JGR Oceans*, 122, 3696–3714, <https://doi.org/10.1002/2016jc012460>, 2017.
- Lovely, A., Loose, B., Schlosser, P., McGillis, W., Zappa, C., Perovich, D., Brown, S., Morell, T., Hsueh, D., and Friedrich, R.: The Gas Transfer through Polar Sea ice experiment: Insights into the rates and pathways that determine geochemical fluxes, *J. Geophys. Res. Oceans*, 120, 8177–8194, <https://doi.org/10.1002/2014jc010607>, 2015.
- Lorenson, T. D., Greinert, J., and Coffin, R. B.: Dissolved methane in the Beaufort Sea and the Arctic Ocean, 1992-2009; sources and atmospheric flux, *Limnol. Oceanogr.*, 61, S300–S323, <https://doi.org/10.1002/lno.10457>, 2016.
- Lorke, A., Bodmer, P., Noss, C., Alshboul, Z., Koschorreck, M., Somlai-Haase, C., Bastviken, D., Flury, S., McGinnis, D. F., Maeck, A., Müller, D., and Premke, K.: Technical note: drifting versus anchored flux chambers for measuring greenhouse gas emissions from running waters, *Biogeosciences*, 12, 7013–7024, <https://doi.org/10.5194/bg-12-7013-2015>, 2015.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium, *Marine Chemistry*, 70, 105–119, [https://doi.org/10.1016/s0304-4203\(00\)00022-0](https://doi.org/10.1016/s0304-4203(00)00022-0), 2000.
- MacIntyre, S., Fram, J. P., Kushner, P. J., Bettez, N. D., O'Brien, W. J., Hobbie, J. E., and Kling, G. W.: Climate-related variations in mixing dynamics in an Alaskan arctic lake, *Limnol. Oceanogr.*, 54, 2401–2417, https://doi.org/10.4319/lo.2009.54.6_part_2.2401, 2009.

1100 MacIntyre, S., Jonsson, A., Jansson, M., Aberg, J., Turney, D. E., and Miller, S. D.: Buoyancy flux, turbulence, and the gas transfer coefficient in a stratified lake, *Geophys. Res. Lett.*, *37*, n/a-n/a, <https://doi.org/10.1029/2010gl044164>, 2010.

Manizza, M., Menemenlis, D., Zhang, H., and Miller, C. E.: Modeling the Recent Changes in the Arctic Ocean CO₂ Sink (2006–2013), *Global Biogeochem. Cycles*, *33*, 420–438, <https://doi.org/10.1029/2018gb006070>, 2019.

1105

[Mannich, M., Fernandes, C. V. S., and Bleninger, T. B.: Uncertainty analysis of gas flux measurements at air–water interface using floating chambers, *Ecohydrology & Hydrobiology*, *19*, 475–486, <https://doi.org/10.1016/j.ecohyd.2017.09.002>, 2019.](https://doi.org/10.1016/j.ecohyd.2017.09.002)

1110 Manning, C. C. M., Zheng, Z., Fenwick, L., McCulloch, R. D., Damm, E., Izett, R. W., Williams, W. J., Zimmermann, S., Vagle, S., and Tortell, P. D.: Interannual Variability in Methane and Nitrous Oxide Concentrations and Sea–Air Fluxes Across the North American Arctic Ocean (2015–2019), *Global Biogeochemical Cycles*, *36*, <https://doi.org/10.1029/2021gb007185>, 2022.

1115 Manning, C. C., Preston, V. L., Jones, S. F., Michel, A. P. M., Nicholson, D. P., Duke, P. J., Ahmed, M. M. M., Manganini, K., Else, B. G. T., and Tortell, P. D.: River Inflow Dominates Methane Emissions in an Arctic Coastal System, *Geophys. Res. Lett.*, *47*, <https://doi.org/10.1029/2020gl087669>, 2020.

1120 Matthews, C. J. D., St.Louis, V. L., and Hesslein, R. H.: Comparison of Three Techniques Used To Measure Diffusive Gas Exchange from Sheltered Aquatic Surfaces, *Environ. Sci. Technol.*, *37*, 772–780, <https://doi.org/10.1021/es0205838>, 2003.

Marq, S. and Weiss, J.: Influence of sea ice lead-width distribution on turbulent heat transfer between the ocean and the atmosphere, *The Cryosphere*, *6*, 143–156, <https://doi.org/10.5194/tc-6-143-2012>, 2012.

1125 McGillis, W. R. and Wanninkhof, R.: Aqueous CO₂ gradients for air–sea flux estimates, *Marine Chemistry*, *98*, 100–108, <https://doi.org/10.1016/j.marchem.2005.09.003>, 2006.

1130 Miller, L. A., Fripiat, F., Else, B. G. T., Bowman, J. S., Brown, K. A., Collins, R. E., Ewert, M., Fransson, A., Gosselin, M., Lannuzel, D., Meiners, K. M., Michel, C., Nishioka, J., Nomura, D., Papadimitriou, S., Russell, L. M., Sørensen, L. L., Thomas, D. N., Tison, J.-L., van Leeuwe, M. A., Vancoppenolle, M., Wolff, E. W., and Zhou, J.: Methods for biogeochemical studies of sea ice: The state of the art, caveats, and recommendations, edited by: Deming, J. W. and Ackley, S. F., *Elementa: Science of the Anthropocene*, *3*, <https://doi.org/10.12952/journal.elementa.000038>, 2015.

- 1135 Miller, L. A., Burgers, T. M., Burt, W. J., Granskog, M. A., and Papakyriakou, T. N.: Air-Sea CO₂ Flux Estimates in Stratified Arctic Coastal Waters: How Wrong Can We Be?, *Geophys. Res. Lett.*, 46, 235–243, <https://doi.org/10.1029/2018gl080099>, 2019.
- Mortin, J., Graversen, R. G., and Svensson, G.: Evaluation of pan-Arctic melt-freeze onset in CMIP5 climate models and reanalyses using surface observations, *Clim Dyn*, 42, 2239–2257, <https://doi.org/10.1007/s00382-013-1811-z>, 2013.
- 1140 Nomura, D., Yoshikawa-Inoue, H., Toyota, T., and Shirasawa, K.: Effects of snow, snowmelting and refreezing processes on air–sea-ice CO₂ flux, *J. Glaciol.*, 56, 262–270, <https://doi.org/10.3189/002214310791968548>, 2010.
- Nomura, D., Granskog, M. A., Assmy, P., Simizu, D., and Hashida, G.: Arctic and Antarctic sea ice acts as a sink for atmospheric CO₂ during periods of snowmelt and surface flooding, *J. Geophys. Res. Oceans*, 118, 6511–6524, <https://doi.org/10.1002/2013jc009048>, 2013.
- 1145 <https://doi.org/10.1002/2013jc009048>, 2013.
- Nomura, D., Granskog, M. A., Fransson, A., Chierici, M., Silyakova, A., Ohshima, K. I., Cohen, L., Delille, B., Hudson, S. R., and Dieckmann, G. S.: CO₂ flux over young and snow-covered Arctic pack ice in winter and spring, *Biogeosciences*, 15, 3331–3343, <https://doi.org/10.5194/bg-15-3331-2018>, 2018.
- 1150 <https://doi.org/10.5194/bg-15-3331-2018>, 2018.
- Onarheim, I. H., Eldevik, T., Smedsrud, L. H., and Stroeve, J. C.: Seasonal and Regional Manifestation of Arctic Sea Ice Loss, *J. Climate*, 31, 4917–4932, <https://doi.org/10.1175/jcli-d-17-0427.1>, 2018.
- Ouyang, Z., Li, Y., Qi, D., Zhong, W., Murata, A., Nishino, S., Wu, Y., Jin, M., Kirchman, D., Chen, L., and Cai, W.: The Changing CO₂ Sink in the Western Arctic Ocean From 1994 to 2019, *Global Biogeochemical Cycles*, 36, <https://doi.org/10.1029/2021gb007032>, 2022.
- 1155 <https://doi.org/10.1029/2021gb007032>, 2022.
- Parmentier, F.-J. W., Christensen, T. R., Sørensen, L. L., Rysgaard, S., McGuire, A. D., Miller, P. A., and Walker, D. A.: The impact of lower sea-ice extent on Arctic greenhouse-gas exchange, *Nature Clim Change*, 3, 195–202, <https://doi.org/10.1038/nclimate1784>, 2013.
- 1160 <https://doi.org/10.1038/nclimate1784>, 2013.
- Prytherch, J. and Yelland, M. J.: Wind, Convection and Fetch Dependence of Gas Transfer Velocity in an Arctic Sea-Ice Lead Determined From Eddy Covariance CO₂ Flux Measurements, *Global Biogeochem Cycles*, 35, <https://doi.org/10.1029/2020gb006633>, 2021.
- 1165 <https://doi.org/10.1029/2020gb006633>, 2021.

- Prytherch, J., Brooks, I. M., Crill, P. M., Thornton, B. F., Salisbury, D. J., Tjernström, M., Anderson, L. G., Geibel, M. C., and Humborg, C.: Direct determination of the air-sea CO₂ gas transfer velocity in Arctic sea ice regions, *Geophys. Res. Lett.*, **44**, 3770–3778, <https://doi.org/10.1002/2017gl073593>, 2017.
- 1170 Rantanen, M., Karpechko, A. Yu., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun Earth Environ*, **3**, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Rigor, I. G., Colony, R. L., and Martin, S.: Variations in Surface Air Temperature Observations in the Arctic, 1979–97, *J. Climate*, **13**, 896–914, [https://doi.org/10.1175/1520-0442\(2000\)013<0896:visato>2.0.co;2](https://doi.org/10.1175/1520-0442(2000)013<0896:visato>2.0.co;2), 2000.
- 1175 Rutgers van der Loeff, M. M., Cassar, N., Nicolaus, M., Rabe, B., and Stimac, I.: The influence of sea ice cover on air-sea gas exchange estimated with radon-222 profiles, *J. Geophys. Res. Oceans*, **119**, 2735–2751, <https://doi.org/10.1002/2013jc009321>, 2014.
- 1180 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F.,
- 1190 Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, **12**, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- Silyakova, A., Nomura, D., Kotovitch, M., Fransson, A., Delille, B., Chierici, M., and Granskog, M. A.: Methane release from open leads and new ice following an Arctic winter storm event, *Polar Science*, **33**, 100874, <https://doi.org/10.1016/j.polar.2022.100874>, 2022.
- Smith, S. D.: Coefficients for sea surface wind stress, heat flux, and wind profiles as a function of wind speed and temperature, *J. Geophys. Res.*, **93**, 15467, <https://doi.org/10.1029/jc093ic12p15467>, 1988.

1200

Snoeijs-Leijonmalm and the the SAS-Oden 2021 Scientific Party: Expedition Report SWEDARCTIC: Synoptic Arctic Survey 2021 with icebreaker Oden [urn:nbn:se:polar:diva-8882](https://nbn-resolving.org/urn:nbn:se:polar:diva-8882), 2022.

1205 Spreen, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2005jc003384>, 2008.

Thornton, B. F., Wik, M., and Crill, P. M.: Double-counting challenges the accuracy of high-latitude methane inventories, *Geophys. Res. Lett.*, 43, <https://doi.org/10.1002/2016gl071772>, 2016a.

1210 Thornton, B. F., Geibel, M. C., Crill, P. M., Humborg, C., and Mörth, C.-M.: Methane fluxes from the sea to the atmosphere across the Siberian shelf seas, *Geophys. Res. Lett.*, 43, 5869–5877, <https://doi.org/10.1002/2016gl068977>, 2016b.

1215 Thornton, B. F., Prytherch, J., Andersson, K., Brooks, I. M., Salisbury, D., Tjernström, M., and Crill, P. M.: Shipborne eddy covariance observations of methane fluxes constrain Arctic sea emissions, *Sci. Adv.*, 6, <https://doi.org/10.1126/sciadv.aay7934>, 2020.

Tjernström, M., Birch, C. E., Brooks, I. M., Shupe, M. D., Persson, P. O. G., Sedlar, J., Mauritsen, T., Leck, C., Paatero, J., Szczerzak, M., and Wheeler, C. R.: Meteorological conditions in the central Arctic summer during the Arctic Summer Cloud Ocean Study (ASCOS), *Atmos. Chem. Phys.*, 12, 6863–6889, <https://doi.org/10.5194/acp-12-6863-2012>, 2012.

1220

Upstill-Goddard, R. C., Rees, A. P., and Owens, N. J. P.: Simultaneous high-precision measurements of methane and nitrous oxide in water and seawater by single phase equilibration gas chromatography, *Deep Sea Research Part I: Oceanographic Research Papers*, 43, 1669–1682, [https://doi.org/10.1016/s0967-0637\(96\)00074-x](https://doi.org/10.1016/s0967-0637(96)00074-x), 1996.


1225 Vachon, D., Prairie, Y. T., and Cole, J. J.: The relationship between near-surface turbulence and gas transfer velocity in freshwater systems and its implications for floating chamber measurements of gas exchange, *Limnol. Oceanogr.*, 55, 1723–1732, <https://doi.org/10.4319/lo.2010.55.4.1723>, 2010.

1230 Vüllers, J., Achtert, P., Brooks, I. M., Tjernström, M., Prytherch, J., Burzik, A., and Neely III, R.: Meteorological and cloud conditions during the Arctic Ocean 2018 expedition, *Atmos. Chem. Phys.*, 21, 289–314, <https://doi.org/10.5194/acp-21-289-2021>, 2021.

- 1235 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr. Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- 1240 Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C., and McGillis, W. R.: Advances in Quantifying Air-Sea Gas Exchange and Environmental Forcing, *Annu. Rev. Mar. Sci.*, 1, 213–244, <https://doi.org/10.1146/annurev.marine.010908.163742>, 2009.
- Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nat Commun*, 10, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- [Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, 2, 203–215, \[https://doi.org/10.1016/0304-4203\\(74\\)90015-2\]\(https://doi.org/10.1016/0304-4203\(74\)90015-2\), 1974.](https://doi.org/10.1016/0304-4203(74)90015-2)
- 1245 Wiesenburg, D. A. and Guinasso, N. L., Jr.: Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and seawater, *J. Chem. Eng. Data*, 24, 356–360, <https://doi.org/10.1021/je60083a006>, 1979.
- 1250 Woolf, D. K., Land, P. E., Shutler, J. D., Goddijn-Murphy, L. M., and Donlon, C. J.: On the calculation of air-sea fluxes of CO₂ in the presence of temperature and salinity gradients, *J. Geophys. Res. Oceans*, 121, 1229–1248, <https://doi.org/10.1002/2015jc011427>, 2016.
- [Yang, M., Prytherch, J., Kozlova, E., Yelland, M. J., Parenkat Mony, D., and Bell, T. G.: Comparison of two closed-path cavity-based spectrometers for measuring air–water CO₂ and CH₄ fluxes by eddy covariance, *Atmos. Meas. Tech.*, 9, 5509–5522, <https://doi.org/10.5194/amt-9-5509-2016>, 2016.](https://doi.org/10.5194/amt-9-5509-2016)
- 1255 Yasunaka, S., Murata, A., Watanabe, E., Chierici, M., Fransson, A., van Heuven, S., Hoppema, M., Ishii, M., Johannessen, T., Kosugi, N., Lauvset, S. K., Mathis, J. T., Nishino, S., Omar, A. M., Olsen, A., Sasano, D., Takahashi, T., and Wanninkhof, R.: Mapping of the air–sea CO₂ flux in the Arctic Ocean and its adjacent seas: Basin-wide distribution and seasonal to interannual variability, *Polar Science*, 10, 323–334, <https://doi.org/10.1016/j.polar.2016.03.006>, 2016.
- 1260 Yasunaka, S., Siswanto, E., Olsen, A., Hoppema, M., Watanabe, E., Fransson, A., Chierici, M., Murata, A., Lauvset, S. K., Wanninkhof, R., Takahashi, T., Kosugi, N., Omar, A. M., van Heuven, S., and Mathis, J. T.: Arctic Ocean CO₂ uptake: an improved multiyear estimate of the air–sea CO₂ flux incorporating chlorophyll a concentrations, *Biogeosciences*, 15, 1643–1661, <https://doi.org/10.5194/bg-15-1643-2018>, 2018.
- 1265

Species, (unit)	Measured	Bulk, with surface pX , T , S		Bulk, with 8 m pX , T , S	
		Ho et al., 2006	P&Y, 2021	Ho et al., 2006	P&Y, 2021
CH ₄ ($\mu\text{mol m}^{-2} \text{ day}^{-1}$)	3.5 ± 4.4	9.0 ± 8.2	6.4 ± 5.8	2.5 ± 3.7	1.8 ± 2.6
CO ₂ ($\text{mmol m}^{-2} \text{ day}^{-1}$)	-7.3 ± 5.7	-12.9 ± 11.6	-9.1 ± 8.2	-7.3 ± 8.6	-5.1 ± 6.0

Table 1. Air-sea CH₄ and CO₂ Fluxes (mean \pm standard deviation) determined from chamber measurements and from bulk methods using the specified gas transfer relationship and partial pressure, temperature and salinity measurement location.



Deleted:

0 20
AMSR2 €

Deleted: s

Deleted: and s

1270

Figure 1. Map of the expedition route (red line) with chamber flux sampling locations (yellow circles) and numbers corresponding to Table S1. Sea-ice concentration is shown as determined for 2021-9-1 by AMSR-2 satellite observations and the ASI 6.25km² product.



Figure 2. Time series of: a) *Oden's* latitude; b) wind speeds as measured on *Oden*; c) SIC determined from the ASI 6.25km² product interpolated to *Oden's* position. Grey circles in each plot show averaged values corresponding to the times of flux chamber measurement.

1280




Figure 3. Occurrence probability (blue) and cumulative probability (red) during the expedition of a) wind speeds adjusted to 10 m height, b) SIC determined from the ASI 6.25km² product interpolated to *Oden's* position, c) surface ice or water temperature as measured with IR sensors onboard *Oden*.

1285








Figure 4. Time series of: a) seawater temperature at 8m depth and at the surface during chamber flux sampling, the corresponding freezing point determined from salinity measurement, and surface temperature (ice or water) measured from IR sensors onboard *Oden*; b) Salinity measured from the 8 m intake and at the surface during chamber flux sampling; c) partial pressures of CH₄ in water at specified depths and sampling locations and in the near-surface atmosphere; d) as per c for CO₂.

1290




1295 **Figure 5.** Near-surface profiles of CH₄ saturation (red) and salinity (blue) determined from ice-based Ruttner bottle sampling on the indicated day. An estimate of ice floe thickness is shown (thick black line), determined from auger drilling close to the sampling site.





1300 **Figure 6.** Time series of: a) air-sea fluxes of CH₄ (filled symbols) and CO₂ (non-filled symbols) measured with chamber flux systems at the specified sampling locations; b) $k_{\pm se}$ of CH₄ (filled symbols) and CO₂ (non-filled symbols) determined from air-sea chamber flux measurements using equations 1 and 2; c) snow-air CO₂ fluxes measured with the EGM-4 chamber flux system. Error bars are the standard error of the flux samples in each deployment. Outer black circles in a and b indicate measurements made in the presence of grease ice.

1305



Deleted:

Deleted: ice-air and



Deleted:

1310 Figure 7. Wind speed dependence of k_{660} determined from air-sea chamber flux measurements using Eqs. (1) and (2), and non-linear
 1315 least squares fits to the measurements ($y = a \times x^b$). Outer circles around the markers indicate measurements made in the presence
 of grease ice. Black crosses indicate measurements made with lead widths < 10 m. Diamonds indicate measurements made when
 surface buoyancy flux was negative (surface cooling). Also shown are three wind speed-based parameterisations of k_{660} . And the
parametric WAGT model in both SIC mode (with SIC = 90%) and fetch mode using lead width.

Deleted: three





Figure 8. Surface temperature dependence of ice-air and snow-air CO₂ fluxes measured with the EGM-4 chamber flux system.





Figure 9. Dependence on lead width of water-air chamber flux-derived k_{660} normalised by U_{10} . A non-linear least squares fit to the ice-based measurements ($y = a \times x^b$) is shown (blue line). Outer circles around the markers indicate measurements made in the presence of grease ice. Diamonds indicate measurements made when surface buoyancy flux was negative (surface cooling).



Deleted:

Deleted: 2

1325

1330

Page 1: [1] Deleted John Prytherch 24/08/2023 11:15:00

Page 1: [2] Deleted John Prytherch 24/08/2023 11:15:00

Page 1: [2] Deleted John Prytherch 24/08/2023 11:15:00

Page 1: [3] Deleted John Prytherch 27/09/2023 12:47:00

▼
▲
Page 1: [3] Deleted John Prytherch 27/09/2023 12:47:00

▼
▲
Page 1: [3] Deleted John Prytherch 27/09/2023 12:47:00

▼
▲
Page 1: [3] Deleted John Prytherch 27/09/2023 12:47:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [4] Deleted John Prytherch 26/09/2023 10:36:00

▼
▲
Page 1: [5] Deleted John Prytherch 26/09/2023 10:40:00

▼
▲
Page 1: [5] Deleted John Prytherch 26/09/2023 10:40:00

▼
▲
Page 1: [5] Deleted John Prytherch 26/09/2023 10:40:00

▼
▲
Page 1: [5] Deleted John Prytherch 26/09/2023 10:40:00

▼
▲
Page 1: [5] Deleted John Prytherch 26/09/2023 10:40:00

▼ ▲
Page 1: [6] Deleted John Prytherch 26/09/2023 11:28:00

▼ ▲
Page 1: [6] Deleted John Prytherch 26/09/2023 11:28:00

▼ ▲
Page 1: [6] Deleted John Prytherch 26/09/2023 11:28:00

▼ ▲
Page 1: [6] Deleted John Prytherch 26/09/2023 11:28:00

▼ ▲
Page 1: [6] Deleted John Prytherch 26/09/2023 11:28:00

▼ ▲
Page 1: [7] Deleted John Prytherch 26/09/2023 11:29:00

▼ ▲
Page 1: [7] Deleted John Prytherch 26/09/2023 11:29:00

▼ ▲
Page 14: [8] Deleted John Prytherch 27/09/2023 10:11:00

▼ ▲
Page 14: [9] Deleted John Prytherch 26/09/2023 13:45:00