



Biomineralization of Fe- and Mn-rich silicates by cyanobacteria under oxic and alkaline conditions

Karim Benzerara^{1*}, Agnes Elmaleh¹, Maria Ciobanu², Alexis De Wever¹, Paola Bertolino², Miguel Iniesto², Didier Jézéquel³, Purificación López-García², Nicolas Menguy¹, Elodie Muller¹, Fériel Skouri-Panet¹, Sufal Swaraj⁴, Rosaluz Tavera⁵, Christophe Thomazo^{6,7}, David Moreira²

¹Sorbonne Université, Muséum National d'Histoire Naturelle, UMR CNRS 7590. Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie (IMPMC), Paris, France

²Unité d'Ecologie Systématique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, Orsay, France

10 ³IPGP, CNRS UMR 7154, Université de Paris & UMR CARRTEL, INRAE-USMB, France

⁴Synchrotron SOLEIL, L'Orme des Merisiers, Saint-Aubin-BP 48, 91192 Gif-sur-Yvette, cedex, France

⁵Departamento de Ecología y Recursos Naturales, Universidad Nacional Autónoma de México, DF México, México

⁶Biogéosciences, CNRS UMR 6282, Université de Bourgogne Franche-Comté, France

⁷Institut Universitaire de France, Paris, France

15

*Correspondence to: Karim Benzerara (karim.benzerara@sorbonne-universite.fr)

20

Keywords: Biomineralization; Iron; Manganese; Cyanobacteria

Abstract. Iron and manganese are poorly soluble elements in oxic and alkaline solutions, whereas they are much more soluble under anoxic conditions. As a result, the formation of authigenic mineral phases rich in Fe and/or Mn has traditionally been viewed as diagnostic of global or local anoxic conditions. Here we reveal that some specific picocyanobacteria can biomineralize abundant, authigenic Fe(III)- and Mn(IV)-rich amorphous silicates under oxic conditions in an alkaline lake in Mexico. The resulting biominerals cluster as small globules arranged as rings around the division septum of cyanobacterial cells. These rings are enveloped within an organic, likely polysaccharidic envelope, and are preserved upon sedimentation. Based on their 16S rDNA sequence, these cyanobacteria were affiliated to the Synechococcales order. The high Fe and Mn enrichment of the silicates questions the systematic inference of anoxic conditions based on their detection. Moreover, this process scavenges iron from the water column, an overlooked biological contribution to the Fe cycle. Finally, it reveals a new case of controlled biomineralization of silicates by bacteria.

25
30



1 Introduction

35 Iron is the fourth most abundant element in the Earth's crust (Taylor, 1964). It is essential to a broad diversity of organisms,
particularly primary producers (Liu et al., 2021). However, Fe is poorly soluble in oxic, neutral to alkaline waters and
sometimes becomes a limiting nutrient for the biosphere (Tagliabue et al., 2017). Overall, its bioavailability affects the
efficiency of atmospheric carbon uptake in aqueous environments (Aumont and Bopp, 2006). Therefore, efforts have been
invested to model the Fe cycle in these environments (Boyd and Ellwood, 2010). Fe enters the cycle through different sources
40 such as aeolian dust, detrital particles, release from anoxic sediments, or hydrothermal vents (Jickells et al., 2005; Elrod et al.,
2004; Tagliabue et al., 2010). Biological activity sets a “ferrous wheel” by mediating the dissolution of particulate Fe and
degrading biogenic Fe-organic complexes on the one hand, and through Fe cell uptake and complexation by organic ligands
on the other hand. Adsorption on various settling mineral particles can scavenge iron out of this wheel. Overall, the balance
between removal and release controls the bioavailability of Fe. In any case, since dissolved Fe concentrations remain low
45 under these conditions, no significant authigenesis of Fe-rich mineral phases is expected in an oxic water column. By contrast,
soluble reduced Fe(II) was much more abundant in ancient aqueous environments before atmosphere oxygenation, 2.4 billion
years ago (Canfield, 2005). At that time, authigenesis of Fe-rich mineral phases was usual, either by precipitation of Fe(II)
phases and/or local oxidation of Fe(II) in Fe(III), resulting in the massive deposition of sedimentary Fe-rich formations such
as banded iron formations (BIFs) (Canfield, 2005). The nature of the authigenic minerals feeding these formations has been
50 widely discussed. Among the possible candidates, Fe(II)-rich silicates such as greenalite have been suggested (Rasmussen et
al., 2019). Like iron, reduced Mn(II) is highly soluble, while oxidized Mn (III, IV) precipitates as insoluble oxides under oxic
conditions (Johnson, 2019). Consequently, the finding of Fe- and/or Mn-rich authigenic minerals has often been considered as
an indication of past anoxic conditions (Rasmussen et al., 2019; Tosca et al., 2016).

Here, we examine the particulate fraction in the water column of a Mexican crater lake. We evidence a biological process that
55 forms Fe- and Mn-rich silicates under oxic and alkaline conditions.

2 Materials and Methods

2.1 Field campaign and sample collection

60 Cyanobacteria forming (Fe,Mn)-rich silicate rings were detected in all field campaigns to Lake La Preciosa at all depths (0 to
35 m), in May 2014, May 2016, March 2018 and May 2019. Temperature, dissolved oxygen concentration, turbidity and pH
were measured in situ over a depth profile using a YSI 6600 multiparameter probe and an optode O₂ SDOT nke in May 2016.
The detection limit for dissolved O₂ was ca 0.1 % (0.3 ± 0.2 μM). The oxygen sensor was calibrated against water-vapor-



saturated ambient air (100 % O₂ local saturation) and the “zero” O₂ was checked using a sodium sulphite solution at 10 wt%.
65 The pH sensor was calibrated using 3 buffers (4/7/10). Lake water was sampled with a Niskin bottle in the center of the lake at different depths from the surface down to 35 m, the maximum depth of Lake La Preciosa. Part of the water was filtered through 0.2 µm polyethersulfone (PES) filters on the same day and kept in sterile tubes for chemical analyses. Part of filtered solutions were acidified with nitric acid (2%) for major cation analyses. The other part was used for measurement of anion and orthosilicic acid (H₄SiO₄) concentrations, without pre-acidification. Lastly, microbial cells were concentrated 5000 times from
70 5 L unfiltered water samples using 0.2-µm pore-size Cell-Trap units (MEM-TEQ Ventures Ltd, Wigan, UK).

2.2 Fluorescence-activated cell sorting (FACS) and cell micromanipulation

Cyanobacterial cells were sorted with a BD FACSAria III Cell Sorter (Becton Dickinson and Company, San Jose, California, USA) equipped with one gas-state 633 nm laser (>18 mW, elliptical shape) and three solid state lasers at 488 nm (>20 mW,
75 elliptical shape), 405 nm (>50 mW, elliptical shape), and 375 nm (>7mW, top hat shape). The 488 nm laser was used for the analysis of forward scatter (FSC, 488/10, 1.0 ND filter), side scatter (SSC, 488/10), phycoerythrin (PE, 585/42, 556LP) and chlorophyll/phycoyanin (PerCP-Cy5-5, 695/40, 655LP). Light was detected by Hamamatsu R3896 photomultiplier tubes in C10562 sockets (Hamamatsu, 211 Hamamatsu City, Japan). The applied voltages were: FSC (220 V), SSC (324 V), PE (703 V) and PerCP_Cy5-5 (577 V). The trigger was set on the FSC with a threshold of 2000. The fluidic system was run at 45 psi
80 (3.102 bar) with an 85-µm nozzle. Samples were sorted at a speed of 6000-6500 events s⁻¹, with a flow rate of 1, corresponding to 0.2 psi over the sheath pressure. The sheath fluid consisted of sterile 1X PBS buffer (K₂HPO₄ 0.11%, KH₂PO₄ 0.03%, NaCl 0.8%). In addition to the excitation lasers, an additional BD Accudrop laser of 660 nm was used for Drop Delay setup. Lake La Preciosa cell suspensions were filtered on a 35 µm mesh just before sorting to avoid clogs. We used 1 µm UV Fluoresbrite Microspheres 24062 (Polysciences, Warrington, USA) for calibration of the log range and a pure culture of
85 *Synechococcus rubescens* as a positive control for phycoerythrin and chlorophyll/phycoyanin fluorescence (Wood et al., 1985). Twenty-five thousand events of one distinct phycoerythrin/chlorophyll-positive population (Pop1, Supplementary Fig. 6a-b) were sorted directly on a 13 mm, 0.4 µm pore size polycarbonate filters (MerckMillipore, Germany) using the most accurate sorting mode “Single-cell purity”. The filters were prepared and observed by SEM as described below. For DNA extraction, 20 cells were sorted under the same conditions, directly into a 0.2 mL microtube (Eppendorf) and stored at -20°C.
90 Cytometric data were processed using the BD FACSDIVA V9.0.1 software (Becton, Dickinson and Company, San Jose, California, USA).

Besides, we also isolated cyanobacterial cells by micromanipulation from samples collected in March 2018 using an Eppendorf PatchMan NP2 micromanipulator equipped with 6 µm-diameter microcapillaries (Eppendorf) mounted on a Leica DIII3000 B inverted microscope. Individual cells were rinsed twice with sterile 10 mM Tris pH 8.0 buffer. Several sets of 5 to 10 cells
95 were deposited in a volume of 0.4 µL of the same buffer in 0.2 mL tubes (Eppendorf) and stored frozen at -20°C until further processing.



2.3 16S rRNA gene sequencing and phylogenetic analysis

DNA was extracted from the FACS-sorted and the micromanipulated cells with the PicoPure DNA extraction kit (Applied Biosystems). In the case of the micromanipulated cells, whole genome amplification (WGA) was carried out on the PicoPure-extracted DNA using Multiple Displacement Amplification (MDA) with the REPLI-g WGA kit (Qiagen). 16S rRNA genes were amplified by PCR from the PicoPure-purified and WGA-amplified DNAs using the cyanobacterial-specific primers CYA-106F (5'-CGG ACG GGT GAG TAA CGC GTG A) and CYA-1380R (5'-ACG ACT TCG GGC GTG ACC). PCR amplifications were carried out in 25 µl reaction volume with GoTaq polymerase reaction mix (Promega, Lyon, France). PCR reactions consisted of an initial denaturing period (95°C for 3 min) followed by 35 cycles of denaturing (93°C for 45 s), annealing (55°C for 45 s), extension (72°C for 2 min), and a final extension period (72°C for 5 min). Because of the possible presence of more than one species in the samples, we constructed 16S rRNA gene clone libraries with the PCR amplicons using the Topo TA cloning system (Invitrogen) following the instructions provided by the manufacturer. After plating, positive transformants were screened by PCR amplification using the M13R and T7 flanking vector primers. Amplicons were sequenced by Sanger sequencing (Genewiz, Essex, UK).

All forward and reverse Sanger sequences were quality-controlled and merged using Codon Code Aligner (<http://www.codoncode.com/aligner/>). Then, we utilised Mafft (Katoh and Standley, 2013) to produce multiple sequence alignments including our sequences and the closest blast (Altschul et al., 1997) hits identified in GenBank. Gaps and ambiguously aligned regions were removed with TrimAl (Capella-Gutiérrez et al., 2009). A maximum likelihood phylogenetic tree was constructed with IQ-TREE (Nguyen et al., 2015) using the general time reversible (GTR) model and branch support estimated by ultrafast bootstrapping.

2.4 Sediment collection

A sediment core measuring 10 cm in length was collected from the bottom of Lake La Preciosa at 35-m depth during the field campaign in May 2016, using a gravitational Uwitec corer with a diameter of 90 mm. The sediment core was transferred into a glove-bag and placed under anoxic conditions (N₂ atmosphere) immediately after collection. It was then processed and sliced on the field into cm-scale fractions along the core's vertical axis. Oxygen levels in the glove-bag were monitored with an Oxi 340i WTW oxygen meter and were always below the detection limit of 0.1 mg/L. Back in France, sediments were vacuum-dried in a Jacomex™ glovebox (< 1 ppm O₂) before being analyzed by SEM.

125



2.5 Solution chemistry

Orthosilicic acid (H_4SiO_4) concentrations were determined by continuous flow colorimetric analyses (QuAAtro Axflow) at the Institut de Physique du Globe de Paris (IPGP, Paris, France) using non-acidified, 0.2- μm -filtered La Preciosa solutions. Anion concentrations were measured using ion chromatography (ICS1100 Dionex Thermofisher) using non-acidified, 0.2- μm -
130 filtered La Preciosa solutions. Concentrations of major cations were determined by inductively coupled plasma - optical emission spectrometry (ICP-OES, iCAP6200 Thermofisher), using acidified, 0.2- μm -filtered La Preciosa solutions. The uncertainty in the concentration measurements of orthosilicic acid, anions and cations was lower than 5%. Activities of anions, cations and orthosilicic acid as well as saturation indices of the surface water solutions of the lakes were calculated using the Visual MINTEQ software.

135

2.6 Confocal laser scanning microscopy (CLSM) and correlative with scanning electron microscopy (SEM)

For standard CLSM analyses, cells concentrated using Cell-Trap units were deposited on a glass slide, covered by a coverslip and sealed with nail polish. Samples were observed by CLSM using a Zeiss LSM 710. Excitation was performed at 405 and 488 nm and emission spectra were measured for each pixel of the images over the 405-720 wavelength range using a 34-
140 channel Quasar T-PMT detector. Data were processed using the Zeiss Zen software. For correlative CLSM-SEM, cells were first washed with milliQ water and then deposited on a coverslip and left to dry. The coverslip was mounted onto a KorrMik Life Sciences sample holder and the correlative Shuttle and Find software implemented in ZEN 2012 was used. CLSM analyses were conducted first. The same areas were relocated in the SEM with the Shuttle and Find software.

145 2.7 Scanning electron microscopy (SEM)

Two types of sample preparation were conducted for scanning electron microscopy (SEM) analyses: (i) in one case, cells were sorted by FACS directly on a 13 mm, 0.4 μm pore size polycarbonate membrane (MerckMillipore, Germany), washed with 1 mL sterile milli-Q water and dried at room temperature; (ii) in another case, cell suspensions were vitrified by plunge and freeze in liquid ethane on a gold planchet before being freeze-dried in a Leica EM ACE600 apparatus. Observations were
150 performed using a Zeiss ultra 55 field emission gun SEM. The acquisition of high spatial resolution images was performed at an accelerating voltage of 1 kV and a working distance of ~ 3 mm and a 20 nm aperture using an annular in-column InLens detector for detecting secondary electrons. For the search of silicate rings in samples, backscattered electron (BSE) images were acquired using an angle selective backscattered (AsB) detector at an accelerating voltage of 10 kV and a working distance of ~ 7.5 mm and a 60 nm aperture at high current. The elemental composition of mineral phases was determined under the
155 same conditions as BSE imaging by energy dispersive x-ray spectrometry (EDXS) using an EDS QUANTAX detector (Bruker). EDXS data were analyzed using the ESPRIT software package (Bruker).



2.8 Transmission electron microscopy (TEM), scanning transmission electron microscopy (STEM), and energy filtered-TEM (ETFEM)

160 Cells concentrated from the water column using Cell Trap units were washed before being deposited on a formvar-coated TEM grid and air-dried. Silicate rings were more fastidious to find in sediments because of the high contrast of all mineral particles composing them. Therefore, we enriched samples in rings by a weak acid leaching (step addition of acetic acid keeping pH at ~6). These samples were also deposited on a formvar-coated TEM grid for further analyses.

Observations of unfixed samples were carried out using a Jeol 2100F TEM microscope operating at 200 kV, equipped with a Schottky emitter, a STEM device, which allows Z-contrast imaging in the high-angular annular dark-field (HAADF) mode, a Jeol Si(Li) x-ray detector and a GIF 2001 Gatan energy filter. Semi-quantitative analyses of EDXS spectra were performed using the JEOL Analysis Station software. This was based on the use of K-factors. After subtracting out the background noise in the EDXS spectrum, the software performed a Gaussian fit of selected elemental peaks and calculated the area under the peaks. From this, the atomic percentage of each selected element was assessed. ETFEM elemental mapping of C, Fe and Mn

165 was performed using the 3-window technique (Hofer et al., 1997). This technique requires three energy-filtered images: two positioned before the ionisation edge (pre-edge images), which serve to calculate the background, and one positioned just after the edge (post-edge image). Calculated background image was subtracted from the post-edge image to give an elemental map, in which changes in background shape were considered. Maps were calculated for the C K-edge and Fe and Mn L_{2,3} edges using a 20-eV-wide (for C) or 40-eV-wide (for Fe, Mn) energy window for pre-edge and post-edge. Zero-loss images were

175 obtained by selecting elastically scattered electrons only, using a 10-eV-wide energy window.

2.9 Scanning transmission x-ray microscopy (STXM)

Cells suspended in the water column and sediment samples were prepared in the same way as for TEM and deposited on TEM grids. Scanning transmission x-ray microscopy analyses were performed on the HERMES beamline at the SOLEIL synchrotron (St. Aubin, France). This microscopy uses monochromated x-rays in the soft x-ray energy domain (200-2000 eV) which are focused onto a ~25 nm spot by a Fresnel zone plate. This approach provides images with a ~25 nm spatial resolution and spatially-resolved speciation information based on x-ray absorption near edge structure (XANES) spectroscopy. An image is obtained by positioning the sample at the focal point of the lens and raster scanning it in x and y, while recording the intensity of the transmitted x-rays. Some cells with rings were pre-located by short TEM imaging conducted prior to STXM

185 measurements in order to facilitate the analyses. However, some areas of the TEM grids remained pristine with no prior TEM analyses. A stack of STXM images of the areas of interest was acquired at a sequence of photon energies at the C K-edge first. The spectral resolution was 0.12 eV in the 282–291.5 eV energy range, where most of the narrowest XANES peaks were present. Then, stacks were acquired at the Mn L_{2,3}-edges (spectral resolution of 0.1 eV in the 642-650 and 653.75-662 eV



ranges) before Fe L_{2,3}-edges (spectral resolution of 0.15 eV in the 709-721 eV range). Areas free of particles were used to
190 measure the incident flux (I₀). Images were converted from transmitted intensity units to optical density units (OD) following
the formula $OD = -\log(I/I_0)$. XANES spectra were extracted and mapped using the aXis2000 software (McMaster University,
<http://unicorn.mcmaster.ca/axis/aXis2000-IDLVM.html>).

3 Results

195 3.1 A new biomineralization process of amorphous (Fe,Mn)-rich silicates

We sampled water at several depths in the Mexican crater Lake La Preciosa, up to ~40 m below water level and examined the
plankton (>0.2 μm). Among the broad morphological diversity of microorganisms observed by light microscopy, one frequent
morphotype appeared as doublets of coccoid cells measuring ~1 μm in diameter each and forming loose aggregates of up to
~50 cells (Supplementary Fig. 1). Their abundance was estimated to be about $2.3 (\pm 0.3) \times 10^5$ cells/mL. An opaque band
200 systematically separated each doublet of dividing cells. This trait morphologically resembles that of previously described
members of the *Cyanocadena* genus (Hindák, 1982). With electron microscopy, these bands appeared as bright rings with a
diameter of 1 μm (Fig. 1-2; Supplementary Fig. 2). They localized at the division septum of cells, and were sometimes observed
in planar view when detached from dividing cells (Fig. 1 b-d & 2a). Overall, these rings appear as traces of septation events.
In some cases, we observed single rings orientated perpendicularly in-between 2 pairs of dividing cells (Supplementary Fig.
205 1). This pattern suggests that cells divided along 2 consecutive perpendicular planes. By contrast, in rare cases, we observed a
series of aligned cells with rings perpendicular to the alignment, suggesting a single division direction only.

At higher magnification, the rings appeared as aggregates of globules measuring ~50-100 nm in diameter (Fig. 2b;
Supplementary Fig. 2) and presenting a nanoporous, sometimes fibrous texture (Supplementary Fig. 2T, V). The rings mostly
contained Fe and Si, with some variable amounts of Mg, Ca, K and Mn as shown by energy dispersive x-ray spectrometry
210 (EDXS) (Fig. 2 g-l & Supplementary Fig. 2). They systematically had a low Al content (1.02 ± 1.38 mol%, 25 measurements;
Table SI-1). The $(Fe+Mn+Ca+Mg)/(Si+Al)$ ratio was estimated at 1.5 ± 0.55 on average and varies between 0.8 up to 2 (25
measurements). Electron diffraction revealed that the grains comprising the rings were amorphous (Fig. 2c). No Fe- and/or
Mn-rich intracellular deposits were detected within the cells. The redox state of Mn and Fe in the rings was determined by x-



ray absorption near edge structure (XANES) spectroscopy using scanning transmission x-ray microscopy (STXM). Analyses
215 at the Fe and Mn L_{2,3} edges of pristine rings unirradiated by TEM beforehand showed that Fe and Mn composing the rings
were oxidized, *i.e.*, in the Fe(III) and Mn(IV) redox states (Supplementary Fig. 3).

3.2 Bacteria biomineralizing (Fe, Mn)-rich silicates are cyanobacteria phylogenetically close to *Cyanobium*

The coccoid cells forming the (Fe,Mn)-rich silicate rings showed strong autofluorescence by confocal laser scanning
220 microscopy (CLSM), with two maximum emission peaks at ~585 and 670 nm (Supplementary Fig. 4). Correlation of CLSM
with SEM confirmed that these autofluorescent cells were the ones forming (Fe,Mn)-rich silicate rings (Supplementary Fig.
5). Based on Wood *et al.* (Wood *et al.*, 1985) and a direct comparison with the autofluorescence of reference phycoerythrin-
containing *Synechococcus rubescens* cells, these two peaks were interpreted as resulting from the fluorescence of
phycoerythrin and chlorophyll/phycoyanin, respectively. This indicated that cells forming (Fe,Mn)-rich silicate rings are
225 cyanobacteria.

These phycoerythrin-containing cyanobacteria were efficiently enriched by fluorescence-activated cell sorting (FACS) based
on the autofluorescence properties of their particular pigments (Supplementary Fig. 6a-b). DNA was extracted from the
enriched cell fractions and, after PCR amplification, their 16S rRNA genes were sequenced. In parallel, micromanipulated
cells were also used for 16S rRNA gene sequencing after whole genome amplification (WGA). Maximum likelihood
230 phylogenetic tree reconstruction confirmed that these cells were cyanobacteria. More specifically, their sequences branched
within a group containing sequences classified as *Synechococcus* and *Cyanobium* (Fig. 3). The small differences between La
Preciosa sequences (<1%), suggested a population of very closely related cyanobacterial strains producing (Fe, Mn)-silicate
rings in this lake.

These 16S rRNA gene sequences were used to screen a large dataset of 16S rRNA gene metabarcoding sequences of plankton
235 from 11 Mexican lakes, including La Preciosa (Zeyen *et al.*, 2021; Iniesto *et al.*, 2022). We detected sequences closely related
(>99% identity) to those of the silicate-ring-producing cyanobacteria in nine of these eleven lakes: La Preciosa, Alchichica,
Atexcac, Aljojuca, Alberca de los Espinos, Pátzcuaro, Quechulac, Tecuitlapa and Yuriria (Supplementary Table 2). These
sequences were particularly abundant (representing >1% of all bacterial metabarcoding reads) in La Preciosa, Alchichica,



Atexcac, Aljojuca, and Alberca de los Espinos, sometimes reaching frequencies up to 15%. Interestingly, these lakes were
240 characterized by relatively high salinity and alkalinity values (Zeyen et al., 2021), whereas those where the sequences were
completely absent had the lowest values for these parameters.

3.3 (Fe, Mn)-rich silicates form under oxygenated conditions and within a polysaccharidic extracellular compartment

Although we have not experimentally determined the speciation of Fe in La Preciosa water column, we can infer that the
245 concentration of dissolved free Fe is likely below 1 nM (Liu and Millero, 2002) based on the high pH (between 8.8 and 9.0)
and oxic conditions (>25% of local O₂ water saturation), in particular in the upper part of the water column (Supplementary
Fig. 7). The rest of Fe in the water column of Lake La Preciosa may be (i) colloidal, inorganic, or organic (complexed by
dissolved organic molecules) Fe, in the <0.2 μm dissolved fraction, or (ii) particulate Fe complexed by microbial cells and/or
adsorbed/co-precipitated in inorganic particles. Measured dissolved Fe concentrations (*i.e.* free and colloidal Fe) were always
250 below the detection limit of ICP-AES (0.02 nM), except at a depth of 5 m in 2019, when Fe and Mn were anomalously high,
reaching 6 and 1 μM respectively. Here, Fe and Mn were most likely under a colloidal form, suggesting that the size of the
colloidal pool varies over time.

Dissolved Si concentrations were measured and found to be relatively high in Lake La Preciosa (5x10⁻⁴ M). Moreover, as
observed in several other alkaline Mexican lakes (Zeyen et al., 2021), Lake La Preciosa solutions were oversaturated with
255 amorphous sepiolite (Wollast et al., 1968), a hydrated Mg-silicate phase, at all depths and all times (Supplementary Fig. 7).

Lastly, chemical conditions prevailing locally where silicates form may be different from the ones in the lake. Indeed, we
found evidence of an extracellular compartment enclosing the rings. First, SEM observations at low-electron voltage of
lyophilized samples showed that silicate rings were embedded in a fibrillar matrix, with a mesh texture reminiscent of the
extracellular polymer substances (EPS) excreted by some bacteria (Fig. 1e-f). Second, energy-filtered TEM (EFTEM) imaging
260 showed that the globular grains were contained within a carbon-rich envelope (Supplementary Fig. 8). Finally, the detection
of a carboxylic-rich polymer using XANES spectroscopy at the C K-edge confirmed the polysaccharidic composition of this
carbonaceous envelope (Supplementary Fig. 8).



3.4 Biomineralized (Fe,Mn)-silicates are preserved upon sedimentation

265 The sediments were mostly composed of aragonite, anorthite and diatomaceous amorphous silica (Supplementary Fig. 9).
Moreover, in the bulk Fourier transform infrared spectroscopy spectra of the sediments, bands at 473, 540, 1026, 3625 and
3680 cm^{-1} could be affiliated to clay-like phases, a proportion of which possibly corresponded to the rings. Although they
could not be unambiguously recognized by bulk analyses, (Fe,Mn)-rich rings with well preserved, sometimes partly broken
morphology, were abundant enough so that they could be observed by electron microscopy in sediment samples with no prior
270 chemical treatment (Fig. 4 and Supplementary Fig. 10-12). TEM-EDXS showed that their chemical composition was similar
to that of the water-column rings with relatively high Mn (Mn/Si~0.04 atom/atom) and Fe (Fe/Si~0.36 atom/atom) contents.
The surface appearance of some rings suggested that they were contained within an organic envelope (Fig. 4a and
Supplementary Fig. 10 a), which was further supported by STXM analyses at the C K-edge on acetic-acid-leached samples
(Supplementary Fig. 4). Here, the spectral signature of carbon was different from that measured on water-column rings.
275 However, this may have resulted from artifactual damages induced by the acid-leaching of the samples, TEM pre-screening
and/or transformation of the organics upon sedimentation. Bulk chemical analyses of the sediments showed that Fe represented
up to 2% of the total sediment mass (including ignition loss). This provides some maximum estimate of the Fe-containing
silicate ring abundance.

4 Discussion

280

4.1 A new controlled biomineralization process concentrating Fe and Mn under oxic conditions

Our findings show that some picocyanobacteria biomineralize Fe- and Mn-rich silicates in oxic and alkaline water. Several
studies have documented the extracellular neof ormation of diverse clay-like phases by bacteria (Konhauser and Urrutia, 1999;
del Buey et al., 2021). They do so either by binding anions (e.g., H_3SiO_4^-) to positively-charged molecular sites on the cell
285 surface or extra-polymeric substances (EPS), and by ion-bridging with cations bound to negatively charged functional groups
and/or by increasing the local pH due to their metabolic activity (Zeyen et al., 2015). Here, while these mechanisms may also
occur owing to the evidenced EPS and the possible pH increase due to oxygenic photosynthesis, the control on
biomineralization by this specific bacterium appears much tighter. First, this biomineralization process is not seen in other
cyanobacteria populating Lake La Preciosa and therefore is likely mediated by some biological mechanisms specific to these
290 cyanobacteria. Moreover, (i) the biominerals have a specific ring morphology, with a definite diameter of ~1 μm ; (ii) they are
associated with the cell division plane, (iii) they are enclosed within an extracellular EPS compartment that might be involved
in mineral precipitation and control the unusual textural organization of (Fe, Mn)-silicates as a ring, and (iv) they are rich in
Fe and Mn, whereas the concentration of free Fe under these environmental conditions does not allow abiotic authigenesis of
Fe-rich silicates. This shows that these cyanobacteria manage to efficiently concentrate Fe and Mn locally from the water



295 column, thus likely involving some energy cost. Fe complexed by organic molecules is usually the most bioavailable form for
cyanobacteria (Beghoura et al., 2019). Alternatively, Fe contained in colloids and particles may be released into the water
column through biologically and photochemically-mediated dissolution (Baker and Croot, 2010). The uptake of Fe from one
or several of these sources might be achieved based on processes similar to those evidenced in many other cyanobacteria, such
as reductive uptake (Kranzler et al., 2014) and/or siderophore-mediated Fe(III) uptake (Swanner et al., 2015) but possibly with
300 high efficiency by these cyanobacteria. Mn is usually taken up in its reduced Mn(II) state through the same transporters as Fe
(Qiu et al., 2021). Since no intracellular precipitate rich in Mn and/or Fe was detected in the cyanobacteria, Fe(II) and Mn(II)
appear to be rapidly released after uptake at the division septum where they precipitate to form rings.
The chemical composition of the biomineralized phase (Al-free and a (Fe+Mn+Ca+Mg)/(Si+Al) ratio between 0.8 and 2) is
reminiscent of the composition of various phyllosilicate phases such as smectites and/or serpentines [greenalite or cronstedtite:
305 Fe/Si=1.5 to 4]. Alternatively, the lowest measured (Fe+Mg+Mn)/(Si+Al) values may correspond to Fe-bearing talcs
consistently with phases detected in La Preciosa sediments and Mexican microbialites (Zeyen et al., 2019). The initial phase
may consist of mostly Fe(II)- and Mn(II)-silicates that subsequently oxidized, since decades of experimental syntheses have
consistently shown that the synthesis of Fe(III)-smectites was achieved by starting from an Fe(II)-silicate gel (Baron et al.,
2016). Alternatively, amorphous (Fe, Mn)-silicate may form by the release of Fe(II) and Mn(II) at the septum of the bacterial
310 cells, which would spontaneously oxidize in the orthosilicate-rich water of Lake La Preciosa. This scenario is consistent with
the formation of Fe-Si complexes with an Fe/Si ratio between 1 and 2 observed in several experimental studies (Pokrovski et
al., 2003; Doelsch et al., 2001). Here, the high Si content of the rings may just be related to the Si-rich environmental conditions
and not the result of bacterial Si cycling.

315 4.2 Biological and environmental significance of this biomineralization process

The intensity and environmental distribution of this biomineralization process remain unknown. Several groups of
picocyanobacteria described using classical approaches seem to form Fe-rich precipitates as well, with diverse textural patterns
(Hindák, 2002). For example, *Cyanodictyon imperfectum*, detected in several lakes worldwide, forms rings composed of iron
320 oxides at its division septum (Cronberg and Weibull, 1981; Economou-Amilli and Spartinou, 1991). *Cyanocatena planctonica*
has also been described to form rings composed of iron oxides at the division septum (Hindák, 1982). All these cells divide
along one division plane only, a taxonomic feature different from most Lake La Preciosa cyanobacteria. Hindák (Hindák,
1982) described another cyanobacterium, *Cyanogranis ferruginea*, which divides along 2 planes and forms iron oxide rings
but not at the cell septa, in contrast with the cells in Lake La Preciosa. No Mn or Si enrichment was reported by any of these
325 studies but mineralogical analyses were too limited to be conclusive. One difficulty in inferring phylogenetic affiliations is
that sequence data for all these cyanobacterial morphotypes is not available. Although the morphological features of
picocyanobacteria alone cannot be used to reliably discriminate between different taxa (Callieri et al., 2012), the morphotype



observed in La Preciosa resemble *Cyanocatena* cells (Hindák, 1982), at least regarding their biomineralization pattern. Therefore, we propose to ascribe our new 16S rDNA sequence to this genus, as *Cyanocatena* sp. The taxonomic affiliation of this sequence allows to start investigating the potential prevalence of this unusual biomineralization process in different lakes by the analysis of massive metabarcoding datasets

Although we ignore the molecular mechanisms of this biomineralization process, several potential functions can be speculated for this trait: (i) High amounts of Fe and Mn are needed by cyanobacteria compared with other microorganisms (Raven et al., 1999; Nelson and Junge, 2015) but, at the same time, elevated intracellular concentrations of these elements may enhance oxidative stress, therefore requiring some appropriate homeostasis (Liu et al., 2021; Kranzler et al., 2013). Different systems are involved in bacterial intracellular storage of Fe, including different types of ferritin family proteins (Keren et al., 2004). The sequestration of high amounts of Fe and Mn by extracellular amorphous phases may contribute to homeostasis and provide a larger but less toxic Fe and Mn reserve (Brown et al., 2005; Cosmidis and Benzerara, 2022). (ii) The oxidation of Fe and Mn upon extracellular release may also generate protons which could be used to generate a proton gradient and gain energy, a mechanism suggested for Fe-oxidizing bacteria (Chan et al., 2004). (iii) Last, the sequestration of high amounts of Fe and/or Mn in these biominerals may be a way to divert these elements from other organisms and therefore serve competition purposes, as suggested for the production of siderophores by some cyanobacteria in the case of iron stress (Wilhelm, 1995).

5 Conclusions

Whatever its mechanism and function, the biomineralization of silicates appears as a yet overlooked Fe-scavenging process, which, together with the already known Fe-scavenging formation of lithogenic inorganic particles, removes Fe out of the water column. Cyanobacteria have been considered actors in the formation of some iron deposits by triggering the oxidation of dissolved Fe(II) from anoxic fluids and therefore inducing the precipitation of Fe-rich mineral phases (Crowe et al., 2008; Emerson and Moyer, 2002). The sheaths of some cyanobacteria can also serve as precipitated iron repository in iron-depositing hot springs (Brown et al., 2005). Here, we evidence an additional mechanism by which cyanobacteria may contribute to the formation of Fe-rich phases, involving Fe scavenging and directed nucleation and growth of Fe-precipitates within an organic matrix. The resulting biominerals show diagnostic features, which may help search for them in modern and ancient samples, following the example of the recent discovery of purported cyanobacteria microfossils with intracellular Fe-silicate nanocrystals in the 1.88 Ga Gunflint Iron Formation (Lepot et al., 2017). The embedding of Fe-rich silicate rings within EPS in La Preciosa may enhance their preservation and, concomitantly, the formation of these organominerals may favour the preservation of organic matter upon time, as shown by Keil et al. (Keil et al., 1994). Consistently, at least part of these rings withstand dissolution and remain preserved in Lake La Preciosa sediments, having therefore some potential for fossilization



over longer time periods. Overall, the existence of such a biomineralization process questions the systematic inference of
360 anoxic conditions based on the detection of such Fe- and Mn-rich phases in the sedimentary record.

Data availability

Author contributions

365 KB, AE, MC, PLG, RT, CT, DM designed the study. KB, AdW, PB, MI, DJ, PLG, RT, CT, DM collected the samples on the
field. KB, AE, MC, AdW, PB, MI, DJ, PLG, NM, EM, FSP, SS, RT, DM carried out the measurements. KB, AE, MC, ADW,
MI, NM, EM, DM analyzed the data. All authors wrote the manuscript.

Competing Interests

The authors declare that they have no conflict of interest.

370

Acknowledgements

This study was supported by the INSU programme Intervrvie, the ERC grants Calcyan (PI: K. Benzerara, Grant Agreement no.
307110) and Plast-Evol (PI: D. Moreira, Grant Agreement no. 787904), and the Agence Nationale de la Recherche (PI: P.
López-García, ANR Microbialites, grant number ANR-18-CE02-0013-02). We thank Lucie Franco and Michael Bourges for
375 helping with FACS experiments; Mélanie Poinot, Jena Johnson, Daniel Nothaft for help in the field; Franck Bourdelle for
help on STXM experiments; Mickael Trichet for help with lyophilisation. The authors acknowledge SOLEIL for the provision
of beamtime.

References

- 380 Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J.: Gapped BLAST and PSI-
BLAST: a new generation of protein database search programs, *Nucleic Acids Res*, 25, 3389–3402,
<https://doi.org/10.1093/nar/25.17.3389>, 1997.
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochemical Cycles*, 20,
<https://doi.org/10.1029/2005GB002591>, 2006.
- 385 Baker, A. R. and Croot, P. L.: Atmospheric and marine controls on aerosol iron solubility in seawater, *Marine Chemistry*, 120,
4–13, <https://doi.org/10.1016/j.marchem.2008.09.003>, 2010.



- Baron, F., Petit, S., Tertre, E., and Decarreau, A.: Influence of aqueous Si and Fe speciation on tetrahedral Fe(III) substitutions in nontronites: A clay synthesis approach, *Clays Clay Miner.*, 64, 230–244, <https://doi.org/10.1346/CCMN.2016.0640309>, 2016.
- 390 Beghoura, H., Gorgues, T., Aumont, O., Planquette, H. f., Tagliabue, A., and Auger, P.-A.: Impact of inorganic particles of sedimentary origin on global dissolved iron and phytoplankton distribution, *Journal of Geophysical Research: Oceans*, 124, 8626–8646, <https://doi.org/10.1029/2019JC015119>, 2019.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nature Geosci.*, 3, 675–682, <https://doi.org/10.1038/ngeo964>, 2010.
- 395 Brown, I. I., Mummey, D., and Cooksey, K. E.: A novel cyanobacterium exhibiting an elevated tolerance for iron, *FEMS Microbiol Ecol.*, 52, 307–314, <https://doi.org/10.1016/j.femsec.2004.11.020>, 2005.
- del Buey, P., Sanz-Montero, M. E., Braissant, O., Cabestrero, O., and Visscher, P. T.: The role of microbial extracellular polymeric substances on formation of sulfate minerals and fibrous Mg-clays, *Chem. Geol.*, 581, 120403, <https://doi.org/10.1016/j.chemgeo.2021.120403>, 2021.
- 400 Callieri, C., Cronberg, G., and Stockner, J. G.: Freshwater picocyanobacteria: single cells, microcolonies and colonial forms, in: *Ecology of Cyanobacteria II: Their Diversity in Space and Time*, edited by: Whitton, B. A., Springer Netherlands, Dordrecht, 229–269, https://doi.org/10.1007/978-94-007-3855-3_8, 2012.
- Canfield, D. E.: The early history of atmospheric oxygen: Homage to Robert M. Garrels, *Annual Review of Earth and Planetary Sciences*, 33, 1–36, <https://doi.org/10.1146/annurev.earth.33.092203.122711>, 2005.
- 405 Capella-Gutiérrez, S., Silla-Martínez, J. M., and Gabaldón, T.: trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses, *Bioinformatics*, 25, 1972–1973, <https://doi.org/10.1093/bioinformatics/btp348>, 2009.
- Chan, C. S., De Stasio, G., Welch, S. A., Girasole, M., Frazer, B. H., Nesterova, M. V., Fakra, S., and Banfield, J. F.: Microbial polysaccharides template assembly of nanocrystal fibers, *Science*, 303, 1656–1658, <https://doi.org/10.1126/science.1092098>, 2004.
- 410 Cosmidis, J. and Benzerara, K.: Why do microbes make minerals?, *Comptes Rendus. Géoscience*, 354, 1–39, <https://doi.org/10.5802/crgeos.107>, 2022.
- Cronberg, G. and Weibull, C.: *Cyanodictyon imperfectum*, a new chroococcal blue-green alga from Lake Trummen, Sweden, *Algological Studies/Archiv für Hydrobiologie, Supplement Volumes*, 101–110, https://doi.org/10.1127/algol_stud/27/1981/101, 1981.
- 415 Crowe, S. A., O’Neill, A. H., Katsev, S., Hehanussa, P., Haffner, G. D., Sundby, B., Mucci, A., and Fowle, D. A.: The biogeochemistry of tropical lakes: A case study from Lake Matano, Indonesia, *Limnology and Oceanography*, 53, 319–331, <https://doi.org/10.4319/lo.2008.53.1.0319>, 2008.
- Doelsch, E., Stone, W. E. E., Petit, S., Masion, A., Rose, J., Bottero, J.-Y., and Nahon, D.: Speciation and crystal chemistry of Fe(III) chloride hydrolyzed in the presence of SiO₄ ligands. 2. Characterization of Si–Fe aggregates by FTIR and ²⁹Si solid-state NMR, *Langmuir*, 17, 1399–1405, <https://doi.org/10.1021/la0013188>, 2001.
- 420



- Economou-Amilli, A. and Spartinou, M.: The diversity of *Cyanodictyon imperfectum* (Chroococcales, Cyanophyceae) in Lake Amvrakia, Greece, *Algological Studies/Archiv für Hydrobiologie, Supplement Volumes*, 105–114, 1991.
- Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: A missing source for global budgets, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020216>, 2004.
- 425 Emerson, D. and Moyer, C. L.: Neutrophilic Fe-oxidizing bacteria are abundant at the Loihi Seamount hydrothermal vents and play a major role in Fe oxide deposition, *Appl Environ Microbiol*, 68, 3085–3093, <https://doi.org/10.1128/AEM.68.6.3085-3093.2002>, 2002.
- Hindák, F.: On some planktonic coccoid blue-green algae characteristic by Fe-precipitates, *Algological Studies/Archiv für Hydrobiologie, Supplement Volumes*, 241–258, 1982.
- 430 Hindák, F.: Four new chroococcalean species (Cyanophyta/Cyanobacteria) from Western Slovakia, *Biologia - Section Botany*, 57, 415–422, 2002.
- Hofer, F., Grogger, W., Kothleitner, G., and Warbichler, P.: Quantitative analysis of EFTEM elemental distribution images, *Ultramicroscopy*, 67, 83–103, [https://doi.org/10.1016/S0304-3991\(96\)00106-4](https://doi.org/10.1016/S0304-3991(96)00106-4), 1997.
- Iniesto, M., Moreira, D., Benzerara, K., Reboul, G., Bertolino, P., Tavera, R., and López-García, P.: Planktonic microbial communities from microbialite-bearing lakes sampled along a salinity-alkalinity gradient, *Limnology and Oceanography*, 435 67, 2718–2733, <https://doi.org/10.1002/lno.12233>, 2022.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Science*, 308, 67–71, 440 <https://doi.org/10.1126/science.1105959>, 2005.
- Johnson, J. E.: From minerals to metabolisms: Evidence for life before oxygen from the geological record, *Free Radic Biol Med*, 140, 126–137, <https://doi.org/10.1016/j.freeradbiomed.2019.01.047>, 2019.
- Katoh, K. and Standley, D. M.: MAFFT multiple sequence alignment software version 7: improvements in performance and usability, *Mol Biol Evol*, 30, 772–780, <https://doi.org/10.1093/molbev/mst010>, 2013.
- 445 Keil, R. G., Montluçon, D. B., Prahl, F. G., and Hedges, J. I.: Sorptive preservation of labile organic matter in marine sediments, *Nature*, 370, 549–552, <https://doi.org/10.1038/370549a0>, 1994.
- Keren, N., Aurora, R., and Pakrasi, H. B.: Critical roles of bacterioferritins in iron storage and proliferation of cyanobacteria, *Plant Physiology*, 135, 1666–1673, <https://doi.org/10.1104/pp.104.042770>, 2004.
- Konhauser, K. O. and Urrutia, M. M.: Bacterial clay authigenesis: a common biogeochemical process, *Chemical Geology*, 450 161, 399–413, [https://doi.org/10.1016/S0009-2541\(99\)00118-7](https://doi.org/10.1016/S0009-2541(99)00118-7), 1999.
- Kranzler, C., Rudolf, M., Keren, N., and Schleiff, E.: Chapter Three - Iron in Cyanobacteria, in: *Advances in Botanical Research*, vol. 65, edited by: Chauvat, F. and Cassier-Chauvat, C., Academic Press, 57–105, <https://doi.org/10.1016/B978-0-12-394313-2.00003-2>, 2013.



- Kranzler, C., Lis, H., Finkel, O. M., Schmetterer, G., Shaked, Y., and Keren, N.: Coordinated transporter activity shapes high-affinity iron acquisition in cyanobacteria, *ISME J*, 8, 409–417, <https://doi.org/10.1038/ismej.2013.161>, 2014.
- 455 Lepot, K., Addad, A., Knoll, A. H., Wang, J., Troadec, D., Beche, A., and Javaux, E. J.: Iron minerals within specific microfossil morphospecies of the 1.88 Ga Gunflint Formation, *Nat. Commun.*, 8, 14890, <https://doi.org/10.1038/ncomms14890>, 2017.
- Liu, L.-M., Li, D.-L., Deng, B., Wang, X.-W., and Jiang, H.-B.: Special roles for efflux systems in iron homeostasis of non-siderophore-producing cyanobacteria, *Environ Microbiol*, 24, 551–565, <https://doi.org/10.1111/1462-2920.15506>, 2021.
- 460 Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Marine Chemistry*, 77, 43–54, [https://doi.org/10.1016/S0304-4203\(01\)00074-3](https://doi.org/10.1016/S0304-4203(01)00074-3), 2002.
- Nelson, N. and Junge, W.: Structure and energy transfer in photosystems of oxygenic photosynthesis, *Annual Review of Biochemistry*, 84, 659–683, <https://doi.org/10.1146/annurev-biochem-092914-041942>, 2015.
- 465 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q.: IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies, *Mol Biol Evol*, 32, 268–274, <https://doi.org/10.1093/molbev/msu300>, 2015.
- Pokrovski, G. S., Schott, J., Farges, F., and Hazemann, J.-L.: Iron (III)-silica interactions in aqueous solution: insights from X-ray absorption fine structure spectroscopy, *Geochimica et Cosmochimica Acta*, 67, 3559–3573, [https://doi.org/10.1016/S0016-7037\(03\)00160-1](https://doi.org/10.1016/S0016-7037(03)00160-1), 2003.
- 470 Qiu, G.-W., Jiang, H.-B., Lis, H., Li, Z.-K., Deng, B., Shang, J.-L., Sun, C.-Y., Keren, N., and Qiu, B.-S.: A unique porin mediates iron-selective transport through cyanobacterial outer membranes, *Environmental Microbiology*, 23, 376–390, <https://doi.org/10.1111/1462-2920.15324>, 2021.
- Rasmussen, B., Muhling, J. R., Tosca, N. J., and Tsikos, H.: Evidence for anoxic shallow oceans at 2.45 Ga: Implications for the rise of oxygenic photosynthesis, *Geology*, 47, 622–626, <https://doi.org/10.1130/G46162.1>, 2019.
- 475 Raven, J. A., Evans, M. C. W., and Korb, R. E.: The role of trace metals in photosynthetic electron transport in O₂-evolving organisms, *Photosynthesis Research*, 60, 111–150, <https://doi.org/10.1023/A:1006282714942>, 1999.
- Swanner, E. D., Wu, W., Hao, L., Wüstner, M. L., Obst, M., Moran, D. M., McIlvin, M. R., Saito, M. A., and Kappler, A.: Physiology, Fe(II) oxidation, and Fe mineral formation by a marine planktonic cyanobacterium grown under ferruginous conditions, *Frontiers in Earth Science*, 3, 2015.
- 480 Tagliabue, A., Bopp, L., Dutay, J.-C., Bowie, A. R., Chever, F., Jean-Baptiste, P., Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G., Aumont, O., Gehlen, M., and Jeandel, C.: Hydrothermal contribution to the oceanic dissolved iron inventory, *Nature Geosci*, 3, 252–256, <https://doi.org/10.1038/ngeo818>, 2010.
- Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron in ocean biogeochemistry, *Nature*, 543, 51–59, <https://doi.org/10.1038/nature21058>, 2017.
- 485 Taylor, S. R.: Abundance of chemical elements in the continental crust: a new table, *Geochimica et Cosmochimica Acta*, 28, 1273–1285, [https://doi.org/10.1016/0016-7037\(64\)90129-2](https://doi.org/10.1016/0016-7037(64)90129-2), 1964.



- Tosca, N. J., Guggenheim, S., and Pufahl, P. K.: An authigenic origin for Precambrian greenalite: Implications for iron formation and the chemistry of ancient seawater, *GSA Bulletin*, 128, 511–530, <https://doi.org/10.1130/B31339.1>, 2016.
- 490 Wilhelm, S. W.: Ecology of iron-limited cyanobacteria: a review of physiological responses and implications for aquatic systems, *Aquatic Microbial Ecology*, 09, 295–303, <https://doi.org/10.3354/ame009295>, 1995.
- Wollast, R., Mackenzie, F. T., and Bricker, O. P.: Experimental precipitation and genesis of sepiolite at earth-surface conditions, *American Mineralogist*, 53, 1645–1662, 1968.
- Wood, A. M., Horan, P. K., Muirhead, K., Phinney, D. A., Yentsch, C. M., and Waterbury, J. B.: Discrimination between types of pigments in marine *Synechococcus* spp. by scanning spectroscopy, epifluorescence microscopy, and flow cytometry¹, *Limnology and Oceanography*, 30, 1303–1315, <https://doi.org/10.4319/lo.1985.30.6.1303>, 1985.
- 495 Zeyen, N., Benzerara, K., Li, J., Groleau, A., Balan, E., Robert, J.-L., Esteve, I., Tavera, R., Moreira, D., and Lopez-Garcia, P.: Formation of low-T hydrated silicates in modern microbialites from Mexico and implications for microbial fossilization, *Frontiers in Earth Science*, 3, 1–23, <https://doi.org/10.3389/feart.2015.00064>, 2015.
- Zeyen, N., Benzerara, K., Menguy, N., Brest, J., Templeton, A. S., Webb, S. M., Gérard, E., Moreira, D., López-García, P., 500 Tavera, R., and Morin, G.: Fe-bearing phases in modern lacustrine microbialites from Mexico, *Geochimica et Cosmochimica Acta*, 253, 201–230, <https://doi.org/10.1016/j.gca.2019.03.021>, 2019.
- Zeyen, N., Benzerara, K., Beyssac, O., Daval, D., Muller, E., Thomazo, C., Tavera, R., López-García, P., Moreira, D., and Duprat, E.: Integrative analysis of the mineralogical and chemical composition of modern microbialites from ten Mexican lakes: What do we learn about their formation?, *Geochimica et Cosmochimica Acta*, 305, 148–184, 505 <https://doi.org/10.1016/j.gca.2021.04.030>, 2021.

Figures

510

515

520




Figure 1. Scanning electron microscopy imaging of (Fe,Mn)-silicate rings formed by dividing cells. (a-b): SEM images obtained with the secondary electron detector of an air-dried sample collected in 2018 (acceleration voltage: 5 keV; working



555 distance: 7 mm). (c-d): SEM images obtained with the secondary electron detector of air-dried samples collected in 2019 (acceleration voltage: 2 keV; working distance: 3 mm). (e-f): SEM images obtained with the InLens detector of freeze-dried samples collected in 2018. Arrows in (e) show some rings. Arrow in (f) shows the mesh texture of the polysaccharidic envelope of the rings.






Figure 2. Transmission electron microscopy analyses of bacterial (Fe,Mn)-silicate rings. (a) Scanning transmission electron microscopy image in high-angular annular dark field (STEM-HAADF) mode of dividing cells collected in 2019. Rings which are composed of high atomic number elements appear in bright. (b) STEM-HAADF image at higher magnification of a ring collected in 2018, showing that it is composed of globules, which themselves show a fibrous texture. (c) Selected area electron diffraction pattern of a ring, characteristic of an amorphous material. (d) Zero-loss energy-filtered transmission electron microscopy (EFTEM) image of a ring collected in 2018. (e) EFTEM image at the Fe $L_{2,3}$ edges. (f) EFTEM image at the C K-edge. (g) STEM-HAADF image of cells with rings chemically mapped using energy dispersive x-ray spectrometry (EDXS). (h) Chemical map of carbon based on the C $K\alpha$ emission line. (i) Chemical map of iron based on the Fe $K\alpha$ emission line. (j) Chemical map of manganese based on the Mn $K\alpha$ emission line. (k) Chemical map of silicon based on the Si $K\alpha$ emission line. (l) EDXS spectrum of the rings observed in the chemical maps.

605

610

615

620

625




Figure 3. Maximum likelihood phylogenetic tree of cyanobacterial 16S rRNA gene sequences showing the position of the cells carrying (Fe,Mn)-silicate rings. Numbers on branches are bootstrap proportions.




Figure 4. SEM, TEM and STXM analyses of cyanobacterial silicate rings in Lake La Preciosa sediments. (a-d): SEM image in the secondary electron detection mode and EDXS chemical maps of Fe, Si and Mg of a ring in pristine (non-acid-leached)



sediments collected in 2016 at a 3 cm depth in the sediment core. Note that the surface of the rings seems to be covered by a
685 smooth and folded layer in SE mode, suggesting that they are covered by organics. (e) Bright-field STEM image of a ring in
acid-leached sediments. (f-h) Corresponding STEM-EDXS chemical maps of Si, Mn and Fe. (i) STEM-EDXS spectrum
measured on the ring. (j) Carbon map obtained by subtracting an OD-converted STXM image aquired at 282 eV from an OD-
converted image at 288.5 eV. (k-n) Spectral analysis at the C K-edge of the image shown in (j). (k-m): maps of three carbon-
containing compounds; (k): carbon associated with rings; (l) carbon typically in a carbonate functional group; (m) carbon in a
690 carbonate functional group but with a different crystallographic orientation. (n) Corresponding spectra at the C K-edge.
Vertical dashed lines are at 284.8 eV (attributed to aromatic functional groups), 288.5 eV (attributed to carboxylics functional
groups), 290.3 eV (attributed to carbonate groups). The ring analyzed by STXM was recovered from an acid-leached sediment
sample at 1 cm depth in the sediment core (see Fig. SI-11 for TEM analyses).